Title:

Beyond static structures: quantitative dynamics in the eukaryotic RNA exosome complex

Authors:

Jobst Liebau[1], Daniela Lazzaretti[1], Anna Bichler[1], Michael Pilsl[2] and Remco Sprangers[1]*

Affiliations:

[1]Institute of Biophysics and Physical Biochemistry, Regensburg Center for Biochemistry, University of Regensburg, 93053 Regensburg, Germany.

[2]Structural Biochemistry Group, Regensburg Center for Biochemistry, University of Regensburg, 93053 Regensburg, Germany.

Corresponding Author: Remco Sprangers, E-mail: remco.sprangers@ur.de, phone: +49-941-943-7751, ORCID: 0000-0001-7323-6047

One sentence summary:

NMR complements static structures of large molecular machines with quantitative insights into functionally important motions.

Keywords:

protein dynamics, RNA exosome, NMR spectroscopy, methyl-TROSY, ^{19}F NMR

Abstract

Molecular machines play pivotal roles in all biological processes. Most structural methods, however, are unable to directly probe molecular motions. We demonstrate that dedicated NMR experiments can provide quantitative insights into functionally important dynamic regions in very large asymmetric protein complexes. We establish this for the 410 kDa eukaryotic RNA exosome complex that contains ten distinct protein chains. NMR data reveal site-specific interactions and conformational changes in regions that are invisible in static cryo-EM and crystal structures. In particular, we identify a flexible plug region that blocks an aberrant route of the RNA to the active site. Our work thus demonstrates that a combination of state-of-the-art structural biology methods can provide quantitative insights into molecular machines that go well beyond static images.

Main Text:

Protein dynamics are tightly coupled with function (*1–3*). Nuclear magnetic resonance (NMR) methods are particularly well suited to study dynamic processes in solution, at quasi atomic resolution and on a wide range of timescales (*4*). Recent advances in sample preparation combined with NMR pulse-sequence and hardware design (*5–11*) have made complexes over 100 kDa accessible to detailed solution NMR studies. This thus opens ample opportunities where NMR methods can complement static structural information obtained by e.g. single particle cryo-electron microscopy (cryo-EM) or *in silico* tools (*12, 13*).

Here, we study the eukaryotic RNA exosome, an essential 3′–5′ ribonuclease complex (**Fig. 1A**). In the cytoplasm, the exosome is involved in the canonical turnover of mRNA and in mRNA quality control; in the nucleus the complex degrades and processes a wide variety of RNA substrates (*14–16*). The exosome is a modular molecular machine that consists of an inert, nonameric core (Exo9; 300 kDa). This core contains an essential central channel that is formed by six distinct RNase PH-like subunits (Rrp41, Rrp45, Rrp43, Rrp46, Mtr3, Rrp42) and an substrate entrance pore that is formed by three cap subunits (Csl4, Rrp4, Rrp40) that contain RNA binding domains (*17*). Rrp41 and Rrp45 recruit the catalytic subunit Rrp44 (Dis3 in humans) (*18*) to assemble the catalytically active decameric complex (Exo10; 410 kDa). Within Rrp44, the RNB domain harbors processive exonucleolytic activity, while the PIN domain can hydrolyse RNA in an endonucleolytic manner (*19–21*). During catalysis, the 3′ end of a single-stranded RNA substrate is recruited by the cap subunits, threaded through the channel and is finally presented to Rrp44 (**Fig. 1A**). Several compartment-specific co-factors associate with the complex to convey substrate specificity (*20, 22*). Additionally, the exosome can recruit Rrp6, a second catalytic subunit that harbors distributive exonucleolytic activity (*22, 23*). Mutations in the exosome complex have been linked to multiple human diseases, underscoring its central functional importance (*24*). In the past, static structures of the human (*20, 25, 26*) and yeast (*23, 27–33*) exosome complexes have been reported that reveal its subunit organization and RNA interactions.

Static structures of the ctExo9 complex

Here, we complement this structural information and determined the structure of Exo9 from the eukaryotic thermophile *Chaetomium thermophilum* (ctExo9) by X-ray crystallography to 3.8 Å resolution (**fig. S1B, C, table S1A**) and by cryo-EM to 3.2 Å resolution (**Fig. 1B**, **fig. S1A, C, fig. S2,**

table S1B). The general architecture of ctExo9 is identical to that of yeast and human exosome

complexes (*20, 23, 25–32*). Despite this wealth of structural information, several disordered regions are

invisible in these structures. In particular, an extended loopregion in Rrp42 (Rrp42-EL) and the entry

loop in Rrp41 are largely unresolved, while the shorter exit loop of Rrp41 is only partially visible (**Fig.**

1C, fig. S1A, D, E). A priori, these "invisible" regions cannot be considered functionally unimportant,

as disordered regions are often directly involved in biological function (*3, 34, 35*).

NMR assignments in the exosome complex

To obtain insights complementary to the static structures, we turned to methyl-based NMR

spectroscopic methods (*10*). Such approaches have been successfully applied to large, highly

symmetric protein assemblies with molecular weights of up to 1 MDa (*36–38*) and to single-chain

proteins of up to 100 kDa (*39*). In that light, the eukaryotic exosome complex is significantly more

challenging to study, as it contains ten distinct protein chains with a total molecular weight of almost

half a megadalton. To render the exosome complex visible to NMR spectroscopy, we employed a

labeling scheme, in which one subunit at a time was labeled with NMR-active Ile-δ1[^{13}CH$_3$] and Met-

ϵ1[^{13}CH$_3$] methyl groups in an otherwise fully deuterated background ('IM-labeling') (**fig. S3**). Methyl

resonance assignments were obtained by exploiting a divide-and-conquer strategy, where we first

assigned resonances in the monomeric subunits Csl4, Rrp41 and Rrp45 (**Fig. 1D**, **fig. S4**). These

assignments were then transferred to the Exo9 and Exo10 complexes, assisted by numerous point

mutants (**table S6C**, exemplified in **fig. S5**). The assignment completion of the Ile-δ1 resonances was

close to 90% (**table S2**) providing a set of NMR probes that can report on interactions and dynamics

and that are well distributed over the complex.

Interactions between Exo9 and Rrp44

Based on chemical shift perturbations (CSPs) site-specific insights into intermolecular interactions can

be obtained. Chemical shifts of a number of resonances in the ring subunits Rrp41 and Rrp45 differ

significantly between the Exo9 and Exo10 complexes, whereas resonances in Csl4 were unaffected by

the addition of Rrp44 (**Fig. 1D, fig. S6**). These observations are in agreement with existing structural

information for yeast and human exosomes that show that Rrp44 is recruited to the Exo9 complex by

Rrp41 and Rrp45 (*20, 25, 28–31*). Our data thus reveal that interactions between ctExo9 and ctRrp44

are conserved and that methyl-TROSY methods can be exploited to identify interaction interfaces in
large asymmetric eukaryotic assemblies.

The methyl-TROSY methods that we deployed are "blind" to regions that are devoid of Ile or Met
residues. To also investigate such regions, we turned to ^{19}F NMR methods that were recently shown to
be excellent tools to study interactions and dynamics on a broad range of timescales (*40–43*), even for
larger complexes (*44–47*). First, we employed amber codon suppression to introduce a 4-
trifluoromethyl-L-phenylalanine (tfmF) into Rrp41 at position D113 (Rrp41^{D113tfmF}). Based on our
structures, this position is located next to a partially structured loop ('exit loop') that faces the Rrp44
interaction interface of Exo9 and that lines the exit site of the RNA channel (**fig. S1A, D**). Upon
addition of Rrp44 the resonance of Rrp41^{D113tfmF} shifts, demonstrating its spatial proximity to Rrp44
(**fig. S7A**). Second, we introduced a tfmF label at position Q86 in Rrp41 (Rrp41^{Q86tfmF}) that is located in
an extended loop close to the cap subunits and not visible in any of the structures ('entry loop', **fig.
S1A, D**). This resonance is not affected by the addition of Rrp44, in agreement with a remote location
of the entry loop from Rrp44 (**fig. S7B**). To probe if the "invisible" entry loop approaches the entry site
of the RNA channel, we assembled an exosome complex in which the cap subunit Csl4 was labeled
with a paramagnetic 2,2,6,6-tetramethylpiperidine-1-oxyl (TEMPO) spin-label at position E130
(Csl4$^{C122S, E130C-TEMPO}$, **fig. S1C**). Rrp41^{Q86tfmF} proved to be too remote to be affected by the Csl4 spin-
label. However, Rrp41^{G71tfmF}, for which the flourine label is located in the center of the entry loop,
displayed fluorine paramagnetic relaxation enhancements (PREs, Γ) that are a direct reporter of the
distance between the Csl4 spin-label and Rrp41^{G71tfmF}. The spin-label gives rise to enhanced R_1 and R_2
relaxation rates, establishing that the Rrp41 entry loop is located at the entry site of the RNA channel
(**fig. S8, table S3A**). To probe if the entry (Rrp41^{Q86tfmF}) and exit (Rrp41^{D113tfmF}) loops undergo motions
on the micro- to millisecond timescale we measured fluorine CPMG relaxation dispersion experiments
(**fig. S9**) that reveal no signs of chemical exchange, indicating that both loops move on a fast ($\ll$ms)
timescale in solution.

RNA threads through the exosome channel
To investigate interactions between the exosome and RNA in solution, we first assessed methyl CSPs in
the subunit-specific IM-labeled Exo9 and Exo10 complexes. These data reveal that Csl4, Rrp41 and
Rrp45 all interact with the substrate (**Fig. 2A**). In Csl4, CSPs are most pronounced in the S1 domain

(residues 98-178) indicating direct interactions with linear RNA (**Fig. 2C, D**). This is expected since S1

domains have been implicated in RNA binding (*48, 49*). In Rrp41 and Rrp45, resonances of residues

that line the channel are affected by RNA (**Fig. 2A, C**), consistent with RNA being threaded through

the channel (**Fig. 2D**). ^{19}F NMR data confirm the involvement of both the Rrp41 entry and exit loop in

RNA interactions (**Fig. 2B, fig. S8C**). Moreover, for the entry loop, we observe a reduction of the PRE

effect that is caused by Csl4$^{C122S, E130C-TEMPO}$ upon addition of RNA (**fig. S8A, B**). This indicates that the

dynamic entry loop is displaced away from Csl4 when RNA enters the exosome barrel (**fig. S8D**).

Based on previous RNA interaction studies and alignments of the ctRrp41 and ctRrp45 sequences with

corresponding archaeal, yeast and human sequences (**fig. S10**) we establish that the RNA coordination

via positively charged residues inside the channel is conserved among those species (*18, 50*).

RNA displaces a flexible loop at the channel exit

Next, we investigated the dynamics and function of an extended loop region in the Rrp42 subunit,

Rrp42-EL (**fig. S11A**) that is unresolved in both the X-ray and cryo-EM structure (**Fig. 1C**). To obtain

insights into the location of Rrp42-EL in the exosome complex, we engineered a double mutant,

Rrp42$^{C59S, A106C}$, in which the single wild-type Cys residue is replaced by a Ser residue (C59S) and a new

Cys residue is incorporated into Rrp42-EL (A106C). Subsequently, we attached a TEMPO spin-label to

Rrp42-EL (Rrp42$^{C59S, A106C-TEMPO}$) and reconstituted this subunit together with IM-labeled Csl4, Rrp41 or

Rrp45 into Exo9 and Exo10 complexes. Csl4 resonances are not affected by the spin-label (**Fig. 3A, E**),

indicating that Rrp42-EL does not approach the cap subunit Csl4. In contrast, a number of Rrp41 and

Rrp45 resonances display substantial PRE effects ($I_{para}/I_{dia} < 1$) in Exo9 and Exo10 (**Fig. 4B**, **C, F, G,**

fig. S12). The affected residues face the exit site of the Exo9 channel, and PREs are stronger in the

Exo10 complex than in the Exo9 complex indicating that the conformation of Rrp42-EL is affected by

Rrp44. In the presence of RNA substrate, PRE effects in Rrp45 are obliterated (**Fig. 4D, fig. S12**),

indicating that RNA displaces the loop away from Rrp45. Based on that, we conclude that Rrp42-EL

adopts two conformations: one, in which it is proximal to the channel and which is stabilized by Rrp44

('closed') and another, in which it is distant from the channel and which is preferred in the presence of

RNA ('open').

Rrp42-EL dynamics are modulated by Rrp44 and RNA

To obtain direct insights into the dynamics of Rrp42-EL, we labeled A106C with 3-Bromo-1,1,1-

trifluoro-acetone (BTFA) to form Rrp42[C59S, A106C-TFA] (**fig. S13**), which retains ribonucleolytic activity of

the Exo10 complex (**fig. S14**). Within the Exo9 complex, Rrp42[C59S, A106C-TFA] displays one fluorine

resonance (**Fig. 4A**); however, CPMG relaxation dispersion (**Fig. 4B, fig. S15B**) and chemical

exchange saturation transfer (CEST) (**fig. S15A**) measurements reveal the presence of a second, minor

conformation. Addition of single-stranded RNA results in a minor shift of the fluorine resonance

frequency (**Fig. 4A**) and restricts the motions of the loop considerably as evidenced by an attenuated

relaxation dispersion profile and a symmetric CEST dip (**Fig. 4B, fig. S15C, D**). To obtain additional

information on the localization and dynamics of this loop, we attached a spin-label to position A106C

(Rrp42[C59S, A106C-TEMPO]) and determined its PRE effect on Rrp41[D113tfmF] (**Fig. 4C, table S3B**). In the

absence of RNA, sizable PRE effects are visible for R_1 and R_2 relaxation rates, indicating that Rrp42-

EL and the Rrp41 exit loop come within less than ~10 Å from each other. This is in agreement with the

complementary methyl-TROSY data (**Fig. 3B**), where spin-labeled Rrp42-EL caused PRE effects close

to the exit loop of Rrp41 (e.g. I98 and I116). Upon addition of substrate, the fluorine PRE effects are

abolished (**Fig. 4C, H**), which implies that the Rrp41 exit loop and Rrp42-EL move apart.

We next turned to the Exo10 complex, in which the fluorine label in Rrp42-EL displays a second

downfield-shifted resonance implying the formation of a second, long-lived conformation (**Fig. 4D**).

This second conformation is induced by the C-terminal RNB-S1 domains of Rrp44, as Exo10

complexes that only contain the Rrp44 PIN domain or the PIN domain plus the two cold-shock (CS)

domains fail to stabilize the second conformation (**fig. S16**). Structurally, a direct interaction between

Rrp42-EL and the Rrp44-RNB-S1 domains is unlikely based on known structures of the human and *S.*

cerevisiae Exo10 complexes (**fig. S17**). It is, however, plausible that the CS domains are brought into

close spatial proximity of Rrp42-EL when Rrp44-RNB-S1 interacts with Exo9.

The dynamics of Rrp42-EL in the Exo10 complex can be probed using CEST (**Fig. 4E, fig. S15E**) and

longitudinal exchange (EXSY) (**fig. S15F**) experiments. Since EXSY experiments exclusively probe

motions on the slow NMR timescale, we conclude that the dynamics of Rrp42-EL are significantly

slowed down by Rrp44. [19]F PRE effects show that Rrp42-EL in the Exo10 complex is still in close

proximity to the Rrp41 exit loop (**Fig. 4F**), which corroborates the methyl-TROSY data (**Fig. 3B**).

Upon addition of RNA to Exo10, the downshifted [19]F resonance and PRE effects disappear and there

are no indications for motions in the CEST profiles (**Fig. 4D-F, J, fig. S15G, H**) implying that Rrp42-

EL is in the open state. Rrp42-EL likely adopts multiple conformations in the open state as evidenced

by a weak CPMG profile of Exo10 Rrp42$^{C59S,\ A106C\text{-}TFA}$ in the presence of RNA (**fig. S15H**) and a

shoulder of the open-state resonance in the absence of RNA (**Fig. 4D, fig. S15F, inset**).

Quantitative assessment of the dynamics of Rrp42-EL

Interestingly, we observe that upon addition of Rrp44 to the apo Exo9 complex Γ_2 rates are enhanced

(spectra in **Fig. 4C, F**) while Γ_1 rates are diminished (inversion recovery plots in **Fig 4C, F, table S3B**).

This apparent contradiction can be explained by the differential dependence of Γ_1 and Γ_2 on fast

timescale motions. As noted before (*51*), decreased order parameters (S^2) and increased internal

motions (τ_i) can result in enhanced Γ_1 rates, whereas Γ_2 rates are largely unaffected by motions that are

faster than the rotational correlation time (**Eq. S9, fig. S19**). Our data thus imply that Rrp42-EL is more

rigid in the closed conformation, that is more prominently populated in the Exo10 complex, than in the

ensemble of open conformations, which is mainly populated in the Exo9 complex.

To quantitatively assess the dynamics of the open-closed equilibrium and to obtain insights into order

parameters of Rrp42-EL in the two states, we globally fitted a two-site exchange model to the dynamics

experiments (CEST, RD, EXSY) of apo Exo9 and Exo10, and to ^{19}F PRE experiments in the Exo9 and

Exo10 complexes in the absence and presence of RNA (**fig. S15**, **fig. S18**, **fig. S20, table S4**). We

assumed that the chemical shifts, local correlation times and order parameters of the open and closed

states are the same in the Exo9 and Exo10 complexes. The analysis revealed that in Exo9, Rrp42-EL

adopts the closed conformation to 5 ±1 % and that the open to closed transition takes place at a rate

($k_{ex}=k_{open\rightarrow closed}+k_{closed\rightarrow open}$) of 5800 ±1400 s^{-1} (**Fig. 4G**). In the Exo10 complex the population of the

closed conformation is significantly higher (26 ±1 %), whereas the exchange rate is reduced to 35 ±35

s^{-1} (**Fig. 4I**). The order parameter (S^2) of the open conformation (~0.1) is significantly lower than of the

closed conformation (~0.7), revealing that the loop in the open state is highly flexible, whereas it is

stably fixed in the closed state. In the presence of RNA, the open conformation in the Exo9 and Exo10

complex is occupied to 100% (**Fig. 4C, F, H, J**), indicating that Rrp42-EL is fully displaced by

substrate RNA.

Rrp42-EL blocks an aberrant RNA access path

The observation that Rrp42-EL can dynamically interact with the channel exit of the exosome raises the

question of whether this is functionally relevant. To address this, we studied the activity of the Exo10

complex in the presence of full-length Rrp42 and with a version in which Rrp42-EL was truncated

($Rrp42^{\Delta93-125}$). The RNA degradation rate in Exo10 is unaffected by the truncation of Rrp42 (**fig. S21**),

which means that the loop displacement by substrate RNA comes at a low energetic cost. In the

canonical substrate route, RNA threads through the Exo9 channel; it has, however, been shown that

RNA can employ alternative paths to the Rrp44 active site that bypass the Exo9 channel (*33*, *52*). To

assess if RNA can access Rrp44 via a direct path, we introduced an extension into a channel-lining loop

of Rrp45, termed Rrp45-L, that has previously been shown to block the exosome channel in *S.*

cerevisiae (*17*, *52*) (**fig. S22**). We observe that Rrp45-L reduces the activity of the exosome to ~4% of

wild-type activity (**fig. S21D**), confirming that the "through-channel" path is the major route RNA

employs. Next, we combined channel-blocked Exo10 Rrp45-L with the $Rrp42^{\Delta93-125}$ mutant, in which

Rrp42-EL is truncated. Interestingly, we find that the activity in this exosome complex is partially

recovered (**Fig. 5A, fig. S21**). Based on that we conclude that Rrp42-EL functions as a barrier that

blocks an aberrant direct access path to the Rrp44 active site (**Fig. 5B**). Rrp42 in *C. thermophilum* thus

contains a previously unidentified flexible "one-way-plug" that readily allows for passage of substrate

RNA via the through-channel route but that prevents access to the Rrp44 active site via an aberrant

direct path. The latter would result in the potentially detrimental ability of the exosome to degrade

substrates that are not selected for processing or degradation by accessory factors, which interact with

the cap subunits of the exosome complex.

Conclusions

The "resolution revolution" in cryo-electron microscopy (*53*) and the remarkable performance of

structure prediction algorithms have substantially increased insights into the relationship between

protein structure and function. However, those methods provide static structural snapshots and often

lack information on loop regions, protein dynamics and transient interactions, all of which may be

crucial for protein function. Here, we demonstrate that dedicated NMR methods, such as CEST, EXSY,

relaxation dispersion, PRE experiments and a combination of methyl-TROSY and ^{19}F NMR, can

complement static structural information, even in large, fully asymmetric eukaryotic molecular

machines. In particular, we show that quantitative and functionally important information can be

obtained on regions that are "invisible" in structures derived from cryo-EM and X-ray crystallography.

Since large asymmetric or transiently formed complexes play a key role in virtually all aspects of

molecular biology, we envision that the strategies to study large complexes by NMR we laid out here

will be of future importance to gain a deeper understanding of how protein structure, interactions and

dynamics relate to function.

 Figures

Figure 1: Structure of ctExo9 and assignment of NMR spectra. (**A**) Schematic depiction of Exo10.

Throughout the manuscript the Csl4, Rrp41, Rrp42 and Rrp45 subunits are colored blue, red, green and

orange, respectively. The individual Rrp44 domains (PIN; pilT N-terminal, CS: cold-shock, RNB: RNA

binding and S1) are labeled and the exonucleolytic site is highlighted with a star. The path of the RNA

towards the active site is indicated in purple. (**B**) Side view (top) and top view (bottom) of the ctExo9

cryo-EM density map. (**C**) Zoom around the boxed region in panel **B**, top, that displays the cryo-EM

density around the invisible section of the extended loop in Rrp42. (**D**) Ile-δ1 region of methyl-TROSY

spectra for Csl4, Rrp41 and Rrp45 in the monomeric form (column 1) and when reconstituted into

Exo9 (column 2) or Exo10 (column 3). Resonance assignments are indicated.

Figure 2: RNA interaction in the exosome. (**A**) Ile-δ1 region of methyl-TROSY spectra in the absence (black) and presence of RNA for Csl4 (blue), Rrp41 (red) and Rrp45 (orange) reconstituted into Exo9 (top) and Exo10 (bottom). (**B**) 1D ^{19}F spectrum of Rrp41^{D113tfmF} (top; exit loop) and Rrp41^{Q86tfmF} (bottom; entry loop) reconstituted into Exo10 with (red) and without (black) RNA. The dashed line indicates the center of the resonance for Exo10. (**C**) RNA-induced CSPs of Ile-δ1 for Exo9 plotted onto the cryo-EM structure with the same coloring scheme as in panel **A**. Rrp42 is in green. For clarity only the outline of the subunit that is NMR active is shown. (**D**) Schematic depiction of Exo10 showing a tentative RNA path through the exosome channel. The coloring scheme is as in panel **A**. Rrp42 is omitted for clarity.

Figure 3: Localization of Rrp42-EL. PRE effects of Rrp42$^{C59S,\ A106C\text{-}TEMPO}$ on Ile-δ1 of (**A**) Csl4 in Exo9, (**B**) Rrp41 in Exo9 (red) and Exo10 (light red), (**C**) Rrp45 in Exo9 (orange) and Exo10 (light orange) without RNA and (**D**) Rrp45 in Exo9 (orange) and Exo10 (light orange) with RNA. n.d.: value not determined due to signal overlap. Ile-δ1 PREs of (**E**) Csl4 (in Exo9), (**F**) Rrp41 (in Exo10) and (**G**) Rrp45 (in Exo10) plotted onto the cryo-EM structure. Csl4 is shown in blue, Rrp41 in red, Rrp45 in orange and Rrp42 in green. Note, that Rrp42-EL is not visible in the structure (see Fig. 1C). For clarity only the outline of the subunit that is NMR active is shown.liebau_manuscript.odt

Figure 4: Dynamics of Rrp42-EL. (**A**) ^{19}F spectra of Rrp42$^{C59S, A106C-TFA}$ in Exo9 without (top) and with (bottom) RNA and (**B**) corresponding RD profiles. (**C**) Inversion recovery experiments (curves) and ^{19}F 1D spectra (insets) for Rrp41^{D113tfmF} in Exo9 Rrp42$^{C59S, A106C-TEMPO}$ without (left) and with (right) RNA for paramagnetic (in color) and diamagnetic (in black) samples. (**D**) ^{19}F spectra of Rrp42$^{C59S, A106C-TFA}$ in Exo10 without (top) and with (bottom) RNA and (**E**) corresponding CEST profiles, acquired for $B_1 = 25$ Hz. (**F**) Inversion recovery experiments (curves) and ^{19}F 1D spectra (insets) for Rrp41^{D113tfmF} in Exo10 Rrp42$^{C59S, A106C-TEMPO}$ without (left) and with (right) RNA for paramagnetic (in color) and diamagnetic (in black) samples. (**G-J**) Schematic depiction of Rrp42-EL dynamics in Exo9 (**G**) without and (**H**) with RNA and in Exo10 (**I**) without and (**J**) with RNA. Rrp41 is in red, Rrp45 in orange, Csl4 in blue and Rrp42 in green. The green loop depicts Rrp42-EL, the green dot A106C$^{TFA/TEMPO}$ and the red dot Rrp41^{D113tfmF}. Fits for panels **C** and **F** were obtained using **Eq. S1**. Fits for panels **B** and **E** were obtained using the model described in materials and methods.

292 **Figure 5: Rrp42-EL blocks an aberrant RNA access path.** RNA activity assays for channel-blocked

293 Exo10 Rrp45-L (gray crosses, solid line) and Exo10 Rrp45-L Rrp42$^{\Delta 93\text{-}125}$ (yellow circles, dashed line).

294 The lines are global linear fits to the linear activity regime. (**B**) Rrp42-EL allows on-path RNA to

295 access Rrp44 for degradation (left) but blocks a direct access path towards Rrp44 (right).

References

1. K. Henzler-Wildman, D. Kern, Dynamic personalities of proteins. *Nature* **450**, 964–972 (2007).

2. A. Sekhar, L. E. Kay, NMR paves the way for atomic level descriptions of sparsely populated, transiently formed biomolecular conformers. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 12867–12874 (2013).

3. M. Corbella, G. P. Pinto, S. C. L. Kamerlin, Loop dynamics and the evolution of enzyme activity. *Nat. Rev. Chem.* **7**, 536–547 (2023).

4. I. R. Kleckner, M. P. Foster, An introduction to NMR-based approaches for measuring protein dynamics. *Biochim. Biophys. Acta* **1814**, 942–968 (2011).

5. K. Pervushin, R. Riek, G. Wider, K. Wüthrich, Transverse relaxation-optimized spectroscopy (TROSY) for NMR studies of aromatic spin systems in ^{13}C-labeled proteins. *J. Am. Chem. Soc.* **120**, 6394–6400 (1998).

6. V. Tugarinov, P. M. Hwang, J. E. Ollerenshaw, L. E. Kay, Cross-correlated relaxation enhanced ^{1}H^{13}C NMR spectroscopy of methyl groups in very high molecular weight proteins and protein complexes. *J. Am. Chem. Soc.* **125**, 10420–10428 (2003).

7. P. Schanda, B. Brutscher, Very fast two-dimensional NMR spectroscopy for real-time investigation of dynamic events in proteins on the time scale of seconds. *J. Am. Chem. Soc.* **127**, 8014–8015 (2005).

8. P. Schanda, H. Van Melckebeke, B. Brutscher, Speeding up three-dimensional protein NMR experiments to a few minutes. *J. Am. Chem. Soc.* **128**, 9042–9043 (2006).

9. L. E. Kay, Solution NMR spectroscopy of supra-molecular systems, why bother? A methyl-TROSY view. *J. Magn. Reson.* **210**, 159–170 (2011).

10. S. Schütz, R. Sprangers, Methyl TROSY spectroscopy: A versatile NMR approach to study challenging biological systems. *Prog. Nucl. Mag. Res. Sp.* **116**, 56–84 (2020).

11. P. Rößler, D. Mathieu, A. D. Gossert, Enabling NMR studies of high molecular weight systems without the need for deuteration: The XL-ALSOFAST experiment with delayed decoupling. *Angew. Chem. Int. Edit.* **59**, 19329–19337 (2020).

12. Y. Cheng, Single-particle cryo-EM---How did it get here and where will it go. *Science* **361**, 876–880 (2018).

13. J. Jumper, R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates, A. Žídek, A. Potapenko, A. Bridgland, C. Meyer, S. A. A. Kohl, A. J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K. Kavukcuoglu, P. Kohli, D. Hassabis, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589 (2021).

14. C. Kilchert, S. Wittmann, L. Vasiljeva, The regulation and functions of the nuclear RNA exosome complex. *Nat. Rev. Mol. Cell Biol.* **17**, 227–239 (2016).

15. A. Łabno, R. Tomecki, A. Dziembowski, Cytoplasmic RNA decay pathways - Enzymes and mechanisms. *BBA-Mol. Cell Res.* **1863**, 3125–3147 (2016).

16. D. Schaeffer, A. Clark, A. A. Klauer, B. Tsanova, A. van Hoof, "Functions of the Cytoplasmic Exosome" in *RNA Exosome*, T. H. Jensen, Ed. (Springer US, New York, NY, 2010)*Advances in Experimental Medicine and Biology*, pp. 79–90.

17. E. V. Wasmuth, C. D. Lima, Exo- and endoribonucleolytic activities of yeast cytoplasmic and nuclear RNA exosomes are dependent on the noncatalytic core and central channel. *Mol. Cell* **48**, 133–144 (2012).

18. F. Bonneau, J. Basquin, J. Ebert, E. Lorentzen, E. Conti, The yeast exosome functions as a macromolecular cage to channel RNA substrates for degradation. *Cell* **139**, 547–559 (2009).

19. K. Januszyk, C. D. Lima, The eukaryotic RNA exosome. *Curr. Opin. Struc. Biol.* **24**, 132–140 (2014).

20. E.-M. Weick, M. R. Puno, K. Januszyk, J. C. Zinder, M. A. DiMattia, C. D. Lima, Helicase-dependent RNA decay illuminated by a cryo-EM structure of a human nuclear RNA exosome-Mtr4 complex. *Cell* **173**, 1663-1677.e21 (2018).

21. J. C. Zinder, C. D. Lima, Targeting RNA for processing or destruction by the eukaryotic RNA exosome and its cofactors. *Genes Dev* **31**, 88–100 (2017).

22. M. Schmid, T. H. Jensen, "The Nuclear RNA Exosome and Its Cofactors" in *The Biology of MRNA: Structure and Function*, M. Oeffinger, D. Zenklusen, Eds. (Springer International Publishing, Cham, 2019)*Advances in Experimental Medicine and Biology*, pp. 113–132.

23. E. V. Wasmuth, K. Januszyk, C. D. Lima, Structure of an Rrp6–RNA exosome complex bound to poly(A) RNA. *Nature* **511**, 435–439 (2014).

24. D. J. Morton, E. G. Kuiper, S. K. Jones, S. W. Leung, A. H. Corbett, M. B. Fasken, The RNA exosome and RNA exosome-linked disease. *RNA* **24**, 127–142 (2018).

25. P. Gerlach, J. M. Schuller, F. Bonneau, J. Basquin, P. Reichelt, S. Falk, E. Conti, Distinct and evolutionary conserved structural features of the human nuclear exosome complex. *Elife* **7**, e38686 (2018).

26. Q. Liu, J. C. Greimann, C. D. Lima, Reconstitution, activities, and structure of the eukaryotic RNA exosome. *Cell* **127**, 1223–1237 (2006).

27. S. Falk, F. Bonneau, J. Ebert, A. Kögel, E. Conti, Mpp6 incorporation in the nuclear exosome contributes to RNA channeling through the Mtr4 helicase. *Cell Reports* **20**, 2279–2286 (2017).

28. E. Kowalinski, A. Kögel, J. Ebert, P. Reichelt, E. Stegmann, B. Habermann, E. Conti, Structure of a cytoplasmic 11-subunit RNA exosome complex. *Mol. Cell* **63**, 125–134 (2016).

29. J.-J. Liu, C.-Y. Niu, Y. Wu, D. Tan, Y. Wang, M.-D. Ye, Y. Liu, W. Zhao, K. Zhou, Q.-S. Liu, J. Dai, X. Yang, M.-Q. Dong, N. Huang, H.-W. Wang, CryoEM structure of yeast cytoplasmic exosome complex. *Cell Res* **26**, 822–837 (2016).

30. D. L. Makino, M. Baumgärtner, E. Conti, Crystal structure of an RNA-bound 11-subunit eukaryotic exosome complex. *Nature* **495**, 70–75 (2013).

31. J. M. Schuller, S. Falk, L. Fromm, E. Hurt, E. Conti, Structure of the nuclear exosome captured on a maturing preribosome. *Science* **360**, 219–222 (2018).

32. A. Keidel, A. Kögel, P. Reichelt, E. Kowalinski, I. B. Schäfer, E. Conti, Concerted structural rearrangements enable RNA channeling into the cytoplasmic Ski238-Ski7-exosome assembly. *Mol. Cell*, doi: 10.1016/j.molcel.2023.09.037 (2023).

33. D. L. Makino, B. Schuch, E. Stegmann, M. Baumgärtner, C. Basquin, E. Conti, RNA degradation paths in a 12-subunit nuclear exosome complex. *Nature* **524**, 54–58 (2015).

34. V. Csizmok, A. V. Follis, R. W. Kriwacki, J. D. Forman-Kay, Dynamic protein interaction networks and new structural paradigms in signaling. *Chem. Rev.* **116**, 6424–6462 (2016).

35. A. R. Camacho-Zarco, V. Schnapka, S. Guseva, A. Abyzov, W. Adamski, S. Milles, M. R. Jensen, L. Zidek, N. Salvi, M. Blackledge, NMR provides unique insight into the functional dynamics and interactions of intrinsically disordered proteins. *Chem. Rev.* **122**, 9331–9356 (2022).

36. G. Mas, J.-Y. Guan, E. Crublet, E. C. Debled, C. Moriscot, P. Gans, G. Schoehn, P. Macek, P. Schanda, J. Boisbouvier, Structural investigation of a chaperonin in action reveals how nucleotide binding regulates the functional cycle. *Sci. Adv.* **4**, eaau4196 (2018).

37. T.L, Religa, R. Sprangers, L. E. Kay, Dynamic regulation of archaeal proteasome gate opening as studied by TROSY NMR. *Science* **328**, 5974 (2010).

38. R. Sprangers, L. E. Kay, Quantitative dynamics and binding studies of the 20S proteasome by NMR. *Nature* **445**, 618–622 (2007).

39. J. H. Overbeck, D. Stelzig, A.-L. Fuchs, J. P. Wurm, R. Sprangers, Observation of conformational changes that underlie the catalytic cycle of Xrn2. *Nat. Chem. Biol.* **18**, 1152–1160 (2022).

40. J. Liebau, M. Tersa, B. Trastoy, J. Patrick, A. Rodrigo-Unzueta, F. Corzana, T. Sparrman, M. E. Guerin, L. Mäler, Unveiling the activation dynamics of a fold-switch bacterial glycosyltransferase

by ^{19}F NMR. *J. Biol. Chem.* **295**, 9868–9878 (2020).

41. J. H. Overbeck, W. Kremer, R. Sprangers, A suite of ^{19}F based relaxation dispersion experiments to assess biomolecular motions. *Journal of biomolecular NMR* **74**, 753–766 (2020).

42. J. H. Overbeck, J. Vögele, F. Nussbaumer, E. Duchardt-Ferner, C. Kreutz, J. Wöhnert, R. Sprangers, Multi-site conformational exchange in the synthetic neomycin-sensing riboswitch studied by ^{19}F NMR. *Angew. Chem. Int. Ed.* **62**, e202218064 (2023).

43. C. Krempl, R. Sprangers, Assessing the applicability of ^{19}F labeled tryptophan residues to quantify protein dynamics. *J. Biomol. NMR* **77**, 55–67 (2023).

44. Y. Huang, X. Wang, G. Lv, A. M. Razavi, G. H. M. Huysmans, H. Weinstein, C. Bracken, D. Eliezer, O. Boudker, Use of paramagnetic ^{19}F NMR to monitor domain movement in a glutamate transporter homolog. *Nat. Chem. Biol.* **16**, 1006–1012 (2020).

45. L.-P. Picard, R. S. Prosser, Advances in the study of GPCRs by ^{19}F NMR. *Curr. Opin. Struct. Biol.* **69**, 169–176 (2021).

46. X. Wang, D. Liu, L. Shen, F. Li, Y. Li, L. Yang, T. Xu, H. Tao, D. Yao, L. Wu, K. Hirata, L. M. Bohn, A. Makriyannis, X. Liu, T. Hua, Z.-J. Liu, J. Wang, A genetically encoded F-19 NMR probe reveals the allosteric modulation mechanism of cannabinoid receptor 1. *J. Am. Chem. Soc.* **143**, 16320–16325 (2021).

47. L. Ye, X. Wang, A. McFarland, J. J. Madsen, ^{19}F NMR: A promising tool for dynamic conformational studies of G protein-coupled receptors. *Structure* **30**, 1372–1384 (2022).

48. M. Bycroft, T. J. P. Hubbard, M. Proctor, S. M. V. Freund, A. G. Murzin, The solution structure of the S1 RNA binding domain: a member of an ancient nucleic acid-binding fold. *Cell* **88**, 235–242 (1997).

49. B. M. Lunde, C. Moore, G. Varani, RNA-binding proteins: modular design for efficient function. *Nat. Rev. Mol. Cell Biol.* **8**, 479–490 (2007).

50. E. Lorentzen, A. Dziembowski, D. Lindner, B. Seraphin, E. Conti, RNA channelling by the archaeal exosome. *EMBO Rep* **8**, 470–476 (2007).

51. G. M. Clore, J. Iwahara, Theory, practice and applications of paramagnetic relaxation enhancement for the characterization of transient low-population states of biological macromolecules and their complexes. *Chem. Rev.* **109**, 4108–4139 (2009).

52. J. Han, A. van Hoof, The RNA exosome channeling and direct access conformations have distinct in vivo functions. *Cell Rep* **16**, 3348–3358 (2016).

53. W. Kühlbrandt, The resolution revolution. *Science* **343**, 1443–1444 (2014).

54. D. G. Gibson, L. Young, R.-Y. Chuang, J. C. Venter, C. A. Hutchison, H. O. Smith, Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods* **6**, 343–345 (2009).

55. S. J. Miyake-Stoner, C. A. Refakis, J. T. Hammill, H. Lusic, J. L. Hazen, A. Deiters, R. A. Mehl, Generating permissive site-specific unnatural aminoacyl-tRNA synthetases. *Biochemistry* **49**, 1667–1677 (2010).

56. E. Gasteiger, C. Hoogland, A. Gattiker, S. Duvaud, M. R. Wilkins, R. D. Appel, A. Bairoch, "Protein Identification and Analysis Tools on the ExPASy Server" in *The Proteomics Protocols Handbook*, J. M. Walker, Ed. (Humana Press, Totowa, NJ, 2005)*Springer Protocols Handbooks*, pp. 571–607.

57. W. Kabsch, XDS. *Acta Crystallogr. D* **66**, 125–132 (2010).

58. N. Eswar, B. Webb, M. A. Marti-Renom, M. s. Madhusudhan, D. Eramian, M. Shen, U. Pieper, A. Sali, Comparative Protein Structure Modeling Using Modeller. *Current Protocols in Bioinformatics* **15**, 5.6.1-5.6.30 (2006).

59. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta Crystallogr. D* **66**, 486–501 (2010).

60. D. Liebschner, P. V. Afonine, M. L. Baker, G. Bunkóczi, V. B. Chen, T. I. Croll, B. Hintze, L.-W. Hung, S. Jain, A. J. McCoy, N. W. Moriarty, R. D. Oeffner, B. K. Poon, M. G. Prisant, R. J. Read,

J. S. Richardson, D. C. Richardson, M. D. Sammito, O. V. Sobolev, D. H. Stockwell, T. C. Terwilliger, A. G. Urzhumtsev, L. L. Videau, C. J. Williams, P. D. Adams, Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in Phenix. *Acta Crystallogr. D* **75**, 861–877 (2019).

61. T. I. Croll, ISOLDE: a physically realistic environment for model building into low-resolution electron-density maps. *Acta Crystallogr. D* **74**, 519–530 (2018).

62. M. Schorb, I. Haberbosch, W. J. H. Hagen, Y. Schwab, D. N. Mastronarde, Software tools for automated transmission electron microscopy. *Nat. Methods* **16**, 471–477 (2019).

63. S. H. W. Scheres, RELION: Implementation of a Bayesian approach to cryo-EM structure determination. *J. Struct. Biol.* **180**, 519–530 (2012).

64. T. Bepler, A. Morin, M. Rapp, J. Brasch, L. Shapiro, A. J. Noble, B. Berger, Positive-unlabeled convolutional neural networks for particle picking in cryo-electron micrographs. *Nat. Methods* **16**, 1153–1160 (2019).

65. J. Guillerez, P. J. Lopez, F. Proux, H. Launay, M. Dreyfus, A mutation in T7 RNA polymerase that facilitates promoter clearance. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 5958–5963 (2005).

66. P. Schanda, E. Kupce, B. Brutscher, SOFAST-HMQC experiments for recording two-dimensional heteronuclear correlation spectra of proteins within a few seconds. *J. Biomol. NMR* **33**, 199–211 (2005).

67. F. Delaglio, S. Grzesiek, G. W. Vuister, G. Zhu, J. Pfeifer, A. Bax, NMRPipe: a multidimensional spectral processing system based on UNIX pipes. *J. Biomol. NMR* **6**, 277–293 (1995).

68. R. L. J. Keller, *The Computer Aided Resonance Assignment Tutorial* (CANTINA Verlag, Goldau, Switzerland, 2004; cara.nmr.ch).

69. A. J. Baldwin, An exact solution for $R_{2,eff}$ in CPMG experiments in the case of two site chemical exchange. *J. Magn. Reson.* **244**, 114–124 (2014).

70. P. Vallurupalli, A. Sekhar, T. Yuwen, L. E. Kay, Probing conformational dynamics in biomolecules via chemical exchange saturation transfer: a primer. *J. Biomol. NMR* **67**, 243–271 (2017).

71. J.-P. Demers, A. Mittermaier, Binding mechanism of an SH3 domain studied by NMR and ITC. *J. Am. Chem. Soc.* **131**, 4355–4367 (2009).

72. N. Bloembergen, L. O. Morgan, Proton relaxation times in paramagnetic solutions. Effects of electron spin relaxation. *J. Chem. Phys.* **34**, 842–850 (1961).

73. I. Solomon, Relaxation processes in a system of two spins. *Phys. Rev.* **99**, 559–565 (1955).

74. F. Madeira, M. Pearce, A. R. N. Tivey, P. Basutkar, J. Lee, O. Edbali, N. Madhusoodanan, A. Kolesnikov, R. Lopez, Search and sequence analysis tools services from EMBL-EBI in 2022. *Nucleic Acids Res* **50**, W276-279 (2022).

75. Y. Z. Tan, P. R. Baldwin, J. H. Davis, J. R. Williamson, C. S. Potter, B. Carragher, D. Lyumkis, Addressing preferred specimen orientation in single-particle cryo-EM through tilting. *Nat. Methods* **14**, 793–796 (2017).

Acknowledgments

We thank Iris Holdermann (MPI Tübingen) for support in the crystallization of the Exo9 complex, Janina Petters for help with the cloning, Nadine Stefan and Johanna Stöfl for technical assistance, Jan Overbeck and Philip Wurm for support in conducting NMR experiments and David Stelzig for assistance with the RNA activity assays. All present and past group members are acknowledged for critically discussing the project.

Funding:

This project is funded from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 89550 (to JL), by the German Research Foundation (Deutsche Forschungsgemeinschaft) under grant agreement No. SP 1324/3-1 and by European Research Council under the European Union's Seventh Framework Programme (FP7/2007–2013), ERC grant agreement No. 616052 (to RS).

Authors contributions:

Conceptualization: RS, JL, DL

Methodology: RS, JL, DL

Investigation: JL, DL, AB, MP

Formal Analysis: RS, JL, DL

Visualization: JL, DL

Funding acquisition: RS, JL

Project administration: RS

Supervision: RS

Writing – original draft: RS, JL, DL with consent of all the authors

Writing – review & editing: RS, JL, DL with consent of all the authors

Competing interests:

The authors declare no competing interests.

Data and materials availability:

All data are available in the manuscript or the supplementary materials. Atomic coordinates have been deposited in the Protein Data Bank (PDB) with accession code 8PEL (crystal structure) and 8R1O (cryo-EM structure). Cryo-EM maps have been deposited in the Electron Microscopy Data Bank (EMDB) with accession code EMD-18825.

Supplementary Materials

Materials and Methods

Figs. S1 to S22

Tables S1 to S9

References (54–75)