Title

HemK2 functions for sufficient protein synthesis and RNA stability through

eRF1 methylation during Drosophila oogenesis

Authors

Fengmei Xu[1], Ritsuko Suyama[1], Toshifumi Inada[2], Shinichi Kawaguchi[1]*, and

Toshie Kai[1]*

*Corresponding author

E-mail: kawaguchi.shinichi.fbs@osaka-u.ac.jp (KS)

E-mail: kai.toshie.fbs@osaka-u.ac.jp (KT)

Affiliations

[1] Graduate School of Frontier Biosciences, Osaka University, Osaka

565-0871, Japan

[2] Division of RNA and Gene regulation, Institute of Medical Science, The

University of Tokyo, Tokyo 108-8639, Japan.

Abstract

HemK2 is a well-conserved methyltransferase from yeast to human, but genuine substrates remain controversial, and its biological significance in the higher organism remains elusive. In this study, we report HemK2 functions for the methylation of eukaryotic Release Factor 1 (eRF1), which is required for female germline development in *Drosophila melanogaster.* Germline-knockdown of *hemk2* (*hemk2*-GLKD) causes apoptosis of germline cells, and severe downregulation of eRF1 methylation and protein synthesis. Overexpression of methylation-deficient eRF1 phenocopies the defects observed in *hemk2*-GLKD, suggesting that eRF1 is the primary substrate for methylation by HemK2. We show that *hemk2*-GLKD also results in significant downregulation of mRNAs in germline cells, and the defects in egg formation and protein synthesis are partially rescued by blocking the no-go decay pathway. In addition, *hemk2* knockdown increases disome formation, suggesting that perturbation of eRF1 methylation results in ribosome stalling, which in turn triggers ribosome quality control to degrade actively translated mRNAs. We propose that eRF1 methylation by HemK2 is required for the efficient protein production and mRNA stability, to ensure the formation of the high-quality eggs.

Introduction

Post-translational modifications (PTMs), such as methylation phosphorylation, glycosylation, acetylation, and ubiquitination, are chemical alterations occurring after protein synthesis. PTMs significantly impact protein functions, interactions, stability, and enzymatic activity (Ramazi and Zahiri, 2021; Ryšlavá et al., 2013). Among those over 400 types of different PTMs, methylation is one of the most widespread modifications. Methylation reaction is catalyzed by S-adenosylmethionine-dependent methyltransferases (MTases), transferring methyl groups to various biomolecules, not only to proteins but also to DNA and RNA at their cytosines and adenines (Chen et al., 2013; Dai et al., 2021; Ping et al., 2014; Wang et al., 2016). In *Drosophila* genome, 127 MTase genes have been annotated. These MTases were further categorized based on their substrate specificities. Lysine MTases and arginine MTases form large group in protein MTases. Previous studies on histone methylation have highlighted the significance of lysine/arginine modifications in transcription regulation (Gates et al., 2017; Zhao and Garcia, 2015). Besides lysine and arginine residues, however, other amino acid residues such as histidine (Clarke, 2013), aspartic acid (Sprung et al., 2008), asparagine (Klotz et al., 1990), and glutamine (Heurgué-Hamard et al., 2002) are also known to undergo a methylation.

HemK2 is a member of unclassified protein MTases because the different classes of substrates are known to this MTase. HemK2 has been previously

reported to catalyze methylation of histone H4 lysine 12 (H4K12), DNA

N6-adenine (6mA) and eukaryotic translation release factor 1 (eRF1).

HemK2-mediated histone H4K12 methylation regulates the proliferation of

prostate tumor cells (Metzger et al., 2019), while reduced genomic DNA 6mA

levels are associated with tumorigenesis in cancer patients (Xiao et al., 2018).

However, the enzymatic activity of HemK2 towards genomic DNA 6mA

remains controversial (Kweon et al., 2019; Ratel et al., 2006; Woodcock et al.,

2019). On the other hand, the role of HemK2 as a MTase for eRF1 has been

well studied in yeast and mammalian culture cells. HemK2 (N6AMT1 in

humans; YDR140w or Mtq2p in yeast) methylates the glutamine residue in the

GGQ motif of eRF1 (Figaro et al., 2008; Heurgué-Hamard et al., 2006; Liu et

al., 2010). Depletion of HemK2 in yeast (Mtq2p) resulted in sensitivity to

translation fidelity of antibiotics (Polevoda et al., 2006). Mouse HemK2

(PRED28) deficiency led to early embryonic lethality and impaired post

implantation development (Liu et al., 2010). However, whether these potential

substrates are methylated by HemK2 in the physiological conditions in

multicellular animals remains unclear.

Drosophila oogenesis is an ideal model for studying genetic regulation of

egg production. The female contains a pair of ovaries, each of which consist of

18–20 ovarioles containing the germarium at the anterior tip followed by

progressively developing egg chambers (Avilés-Pagán and Orr-Weaver, 2018;

Gleason et al., 2018; Rastegari et al., 2020). Each egg chamber contains one

oocyte and 15 nurse cells which support oocyte growth and is surrounded by a

monolayer of somatic follicle cells. Successful accomplishment of processes

such as stem cell maintenance, their asymmetric division and differentiation,

meiosis, and more importantly, the production of mRNAs and proteins to be

maternally deposited to eggs, is crucial for proper oogenesis. *Drosophila*

oogenesis is an energetically consuming process involving cellular

proliferation and growth that requires diverse protein synthesis (Lasko, 2012).

The role of HemK2 as a MTase has been well studied in yeast and

mammalian culture cells. However, its function in animals, especially in the

reproductive system where active gene expression and protein medication

take place, remains unclear. Here, we characterized HemK2's role in eRF1

methylation during *Drosophila* oogenesis. HemK2 was essential for egg

production; upon knocking down of *hemk2* expression, oogenesis stopped in

the mid stage and germline cells were degenerated by apoptosis. Perturbation

of *hemk2* expression caused severe reduction of eRF1 methylation but not

DNA 6mA and H4K12 methylation. Furthermore, ineffective termination

resulting from unmethylated eRF1 also decreased mRNA abundance through

the No-go-decay pathway. Taken together, these results suggest an essential

role of *hemk2*-mediated eRF1 methylation in ensuring efficient protein

production and mRNA stability, thereby regulating gene expression during

Drosophila oogenesis.

Results

Germline knockdown of *hemk2* exhibited female sterility marked by the developmental arrest in the mid-oogenesis

To explore function of a well-conserved methyltransferase, HemK2, encoded by *CG9960* in *Drosophila melanogaster*, we knocked down its expression by driving shRNA targeting its 3'UTR (BL#40837) with a germline driver, *NGT40; NosGal4-VP16* (Grieder et al., 2000). The oocyte starts accumulating proteins and RNAs provided by 15 nurse cells through the cytoplasmic channels (Figure 1A, B). The females subjected to germline-knockdown of *hemk2* (*hemk2*-GLKD), which resulted in ~65% of reduction of its mRNA level (Figure 1C, S1A), exhibited female sterility. *hemk2*-GLKD females contained small ovaries and did not lay any eggs at all (Figure 1B, D), suggesting the developmental arrest of egg chambers at stages 7/8 prior to the vitellogenic stages. In contrast, *hemk2*-GLKD males with ~75% reduction in the mRNA level did not show any significant defects in fertility (Figure S1F, G).

hemk2 expression is under di-cistronic control, being co-expressed as a single transcript along with the other open reading frame, *snapin*, in the upstream of the *hemk2* coding frame (Figure S1A) (Wall et al., 2005). In order to verify if the defects in *hemk2*-GLKD were caused by reduction of *hemk2* expression, each single transgene of N-terminal FLAG-tagged HemK2 or

Snapin devoid of shRNA targeting sequence was expressed in germline upon

the knockdown of the di-cistronic transcript (Figure 1E; Figure S1E). The

number of laid eggs and hatching rate in the females expressing *hemk2* in

germline were comparable with those of the control (Figure 1D), indicating that

the defect was rescued. In contrast, the *snapin* expression did not recover

those defects (Figure S1D). These results indicate the female sterility was

caused by the deterioration of *hemk2* function upon the knocking down of the

dicistronic transcript, and *hemk2* had essential roles in germline cells during

oogenesis.

To investigate *hemk2*'s function in somatic cells, *hemk2* was knocked

down in ovarian somatic cells including follicle cells by a somatic driver, *traffic*

jam-Gal4 (tj-Gal4). Upon *hemk2* somatic-knockdown (*hemk2*-STKD), the

females became sterile, and the ovaries were rudimentary (Figure S1B, C).

hemk2-STKD ovaries often contained the fused egg chambers and

disorganized follicle cells (Figure S1C). The fertility and developmental defects

were almost completely rescued by the expression of *hemk2* transgene by

tj-Gal4 (Figure S1B, C), suggesting that *hemk2* has a predominant role not

only in germline but also in somatic cells.

hemk2-GLKD ovaries showed pleiotropic phenotypes: hyperplasia of the

posterior follicle cells, failure in chromosome dispersion, and massive

apoptosis of germline cells (Figure 2). While all the egg chambers in the

control females were encapsulated by a monolayer of follicle cells, those in

most of *hemk2*-GLKD ovarioles contained multiple layered follicle cells at the

posterior region (Figure 2A; 94 %). Moreover, cells positive for

phospho-histone H3 (pH3), a marker of mitosis, were often enriched within the

multi-layered follicle cells of *hemk2*-GLKD during stage 7/8 (Figure 2A),

indicating their hyperplasia at posterior side of egg chambers.

Secondly, *hemk2*-GLKD nurse cells showed a chromosome dispersion

defect. The wildtype nurse cells in early oogenesis form polytene

chromosomes condensing into a characteristic '5-blob' conformation and

become dispersed completely by stage 6 (Dej and Spradling, 1999). However,

almost all ovarioles of *hemk2*-GLKD harbored nurse cell nuclei containing

chromosomes remained in a blob-like configuration even beyond stage 6

(Figure 2B, Figure S2A; 99%). In order to investigate the possible causes that

may results in this chromosomal defect, we examined the expression level of

the genes which were known to function for chromosomal dispersion

(Goodrich et al., 2004; Klusza et al., 2013). Among those genes, qRT-PCR

analysis revealed that the mRNA level of *otu* was most drastically reduced

upon *hemk2*-GLKD (Figure 2C), suggesting a potential correlation between *otu*

expression and the observed defect. The expression of N-terminal

FLAG-tagged Otu104 isoform in *hemk2*-GLKD ovaries significantly restored

the number of ovarioles containing dispersed chromosomes (Figure S2A).

These results suggest that *hemk2* functions for the chromosome dispersion

during oogenesis, at least in part, by ensuring the expression of *otu*.

Thirdly, *hemK2*-GLKD resulted in massive degeneration of egg chambers

in mid-oogenesis where the germline cell death led to degeneration of egg

chambers depending on physiological conditions and environmental cues as a

developmental checkpoint (Drummond-Barbosa and Spradling, 2001; McCall,

2004). Immunostaining of *hemk2*-GLKD with antibody against p53, an inducer

of apoptosis, revealed increased accumulation of p53 signals in the nuclei of

nurse cell in the majority of *hemk2*-GLKD ovarioles (Figure 2D; 93%),

suggesting that *hemk2*-GLKD resulted in germline cell death in mid-oogenesis.

Double-knockdown of *p53* with *hemk2* significantly restored the progression of

oogenesis in the *hemk2*-GLKD ovaries (Figure S2B). In addition, the

expression of p35, an inhibitor of executioner caspases DrICE and Dcp-1

(Fuchs and Steller, 2015; Werz et al., 2005) also suppressed the cell death in

hemk2-GLKD ovaries (Figure S2B). Although the females with p53-knocking

down or the ectopic expression of p35 in *hemk2*-GLKD females did not fully

restore the egg laying as many as the control, the hatching rates of these eggs

were comparable to that of the control (Figure S2C). Altogether, we conclude

that p53-dependent apoptosis takes place during the mid-oogenesis in

hemk2-GLKD although the trigger mechanism remains elucidate. However,

the chromosomes remained undispersed in *hemk2*-GLKD ovaries restored by

the repression of cell death (Figure S2B), suggesting that apoptosis and

chromosomal dispersion defects involve distinct pathways triggered by *hemK2*

downregulation.

193   *hemK2* **functions for eRF1 methylation and the catalytic activity is**

194   **essential for oogenesis**

195   Previous studies reported HemK2 functioned as methyltransferase for

196   three substrates, DNA N6-adenine (6mA) in human culture cells (Xiao et al.,

197   2018), histone H4 lysine 12 (H4K12) in A549 lung adenocarcinoma cells

198   (Metzger et al., 2019) and a glutamine residue in GGQ motif of eRF1 in yeast

199   (Heurgué-Hamard et al., 2005) and human (Figaro et al., 2008). However, its

200   genuine substrate and function in animals remains elusive. Firstly, we

201   examined 6mA modification of genomic DNA upon *hemk2*-GLKD by

202   immunostaining with an anti-6mA antibody. The pattern and the signal

203   intensity of 6mA staining remained unaffected in the germline of *hemk2*-GLKD

204   ovaries (Figure S3A), demonstrating that HemK2 was unlikely to function as

205   methyltransferase for genomic N6-adenine in *Drosophila* germline. In addition,

206   the amount of methylated H4K12 was comparable in *hemk2*-GLKD and the

207   control ovaries (Figure S3B), excluding H4K12 as a bona fide substrate in

208   *Drosophila* germline. Lastly, we examined methylation level of endogenous

209   eRF1 in *hemk2*-GLKD ovaries. Because antibody against methyl-eRF1 did not

210   work for Western blotting, we conducted eRF1 immunoprecipitation followed

211   by immunoblotting with an antibody against N5-methyl-glutamine (Methyl-Q).

212   The methylation level of eRF1 in *hemk2*-GLKD ovaries was considerably

213   reduced to 35% of that of the control, which was possibly due to the unaffected

214   somatic contribution (Figure 3A). Indeed, immunostaining of *hemk2*-GLKD

ovary with anti-methyl-eRF1 antibody revealed a dramatic reduction of eRF1

methylation in germline cells but not in the somatic follicle cells, while the

expression level of endogenous eRF1 was comparable in both cell types

(Figure 3B). Consistently, immunoprecipitation of germline-expressed

FLAG-tagged eRF1 had a scanty of methylation in *hemk2*-GLKD ovaries

compared to that of the control (Figure 3C), indicating that *Drosophila* HemK2

mediated methylation of eRF1 at glutamine residue(s). It has been reported

that a glutamine residue of a GGQ tripeptide motif in eRF1, which functions for

peptide release, is a well-conserved methylation site by HemK2 in other

organisms (Dinçbas-Renqvist et al., 2000; Figaro et al., 2008; Gao et al., 2020;

Kusevic et al., 2016; Svidritskiy and Korostelev, 2018). We also examined if

GGQ tripeptide motif of eRF1 was the methylation target site in S2 cells by

mutating the corresponding Gln at 185^{th} position of *Drosophila* eRF1 to Ala

(eRF1 Q185A) (Figure 3D). In agreement with those reported in other systems,

methylation of eRF1 Q185A was hardly detected in S2 cells even without

HemK2 knockdown, suggesting that this mutation perturbed the methylation of

eRF1 (Figure 3D). Taken together, these results suggest that Q185 is a

predominant methylation site in eRF1 by HemK2.

The rescue experiments described above confirmed that HemK2 had an

essential function *in vivo* for the egg development (Figure 1 and Figure S1).

We then examined the importance of amino acid residues required for

HemK2's catalytic activity during oogenesis in vivo. The highly conserved

NPPY motif of HemK2 is known as a binding pocket of the substrate and

essential for the methylation activity of HemK2 in human cell lines (Figure 4A)

(Gao et al., 2020; Metzger et al., 2019). Previous study also showed mutations

in the relatively-close residues to the substrates, Asn122 and Tyr125, in NPPY

motif of human HemK2 disrupted its catalytic activity for methylation of eRF1 *in*

vitro (Figure 4B) (Metzger et al., 2019). We mutated the corresponding

residues of *Drosophila* HemK2, 116th Asn and 119th Tyr, to alanine (N116A

and Y119A, respectively), and expressed them in germline of *hemk2*-GLKD

ovaries (Figure 4C; Figure S4A). While HemK2 Y119A did not rescue the

defects of *hemK2*-GLKD, such as degeneration of egg chambers, egg laying

and hatching rate, HemK2 N116A was able to recover them comparably to

those in the control (Figure 4C, D). These results suggest that Tyr119, but not

Asn116, was essential for HemK2 function in *Drosophila* germline cells.

Next, we investigated the catalytic activity of HemK2 for methylation of Gln

in the GGQ motif of eRF1 in S2 cells. Each FLAG-tagged HemK2 variant and

Myc-tagged eRF1 were co-transfected into S2 cells upon knockdown of

endogenous *hemk2* (Figure 4E). The immunoprecipitation of Myc-eRF1

followed by immunoblotting with anti-Methyl-Q antibody revealed that

methylation of eRF1 was significantly reduced upon *hemk2* knockdown, and

such a reduction was successfully rescued by the expression of wildtype

HemK2 as well as N116A mutant, but not Y119A (Figure 4E). These results

indicate that 119[th] Tyr but not 116[th] Asn of *Drosophila* HemK2 is indispensable

to mediate the methylation of eRF1 at 185th Gln.

Several studies previously reported that HemK1, a paralog of HemK2

containing a conserved NPPY motif and 42% homology to HemK2, catalyzed

the methylation of Gln in GGQ motif of mitochondrial release factors in human

cells (Fang et al., 2022; Ishizawa et al., 2008). To address the specificity of

cytosolic eRF1 methylation by HemK2, we examined eRF1 methylation in S2

cells by expression of HemK1 upon *hemk2* knockdown. The reduction of eRF1

methylation by *hemk2* knockdown was successfully rescued by the expression

of HemK2 but not by that of HemK1, indicating that eRF1 methylation is

uniquely catalyzed by HemK2 (Figure S4B).

eRF1 methylation ensures proper progression of oogenesis and efficient

translation

Next, we examined the necessity of eRF1 methylation during oogenesis

by knockdown of eRF1's expression in germline cells. Upon *eRF1*-GLKD,

ovaries became rudimentary and produced no eggs, showing a severer defect

than that of *hemk2*-GLKD ovaries (Figure 5A, B). Immunostaining of

eRF1-GLKD ovaries with a germline marker, a DEAD box RNA helicase, Vas,

further revealed agametic germaria containing very few germline cells (Figure

5C). Those defects by eRF1-GLKD were successfully rescued with the

expression of FLAG-tagged wildtype eRF1 in germline cells but not with that of

the methylation-deficient Q185A mutant (Figure 5B, C). These results suggest

that the 185[th] Gln position, and potentially its methylation, plays a vital role in

the survival of germline cells and progression of oogenesis. The milder

phenotype observed in *hemk2*-GLKD might have been resulted from the

incomplete perturbation of eRF1 methylation, possibly due to residual *hemk2*

expression escaped from knockdown or any other enzyme(s) catalyzing eRF1

methylation.

In addition, the expression of FLAG-eRF1 Q185A in the germline cells of

the wildtype ovaries resulted in failure in egg production (Figure 5D). Even in

the presence of wildtype eRF1, the expression of eRF1 Q185A caused severe

developmental arrest at mid-oogenesis as well as failure in chromosome

dispersion and massive apoptosis, resembling the defect of *hemk2*-GLKD

(Figure 5D; Figure S5A, B). More than 80% of ovarioles expressing eRF1

Q185A contained p53-positive egg chambers (Figure S5C). Taken together,

these results suggest that the expression of eRF1 Q185A phenocopies

hemk2-GLKD; the loss of eRF1 methylation is the main reason causing defects in

hemk2-GLKD.

It has been reported that the methylation of a Gln residue in eRF1 GGQ

motif leads to the release of the nascent polypeptide and assures efficient

translational termination (Dinçbas-Renqvist et al., 2000; Figaro et al., 2008;

Heurgué-Hamard et al., 2002; Svidritskiy and Korostelev, 2018), and

perturbation of proper termination could potentially result in deficient

translational activity by inhibiting ribosome recycling. To assess the translation

activity in *hemk2*-GLKD ovaries, we performed an HPG

(L-homopropargylglycine) incorporation assay ex vivo with the dissected

ovaries. HPG is a methionine analog containing an alkyne moiety and can be

ligated to the fluorescent dye through a 'Click reaction' in the presence of

copper (II) sulfate (Shen et al., 2021). The fluorescent HPG signal representing

the newly synthesized proteins within 30 min in cytoplasm of germline cells

was remarkably reduced in *hemk2*-GLKD ovaries (Figure 5E), suggesting that

eRF1 methylation was required for efficient translation. Consistent with the

capability of eRF1 methylation in S2 cells and of rescuing the defects in

hemk2-GLKD ovaries (Figure 4C, D, E), the reduction of translation efficiency

by *hemk2*-GLKD was fully rescued by the expression of the wildtype *hemk2* as

well as N116A mutant but not the Y119A mutant (Figure 5E). In addition,

similar to that in *hemK2*-GLKD, the expression of methylation-deficient eRF1

Q185A in germline cells abolished the translation activity (Figure S5D).

Altogether, these results suggest that eRF1 methylation by HemK2 is required

for the efficient translation activity in *Drosophila* ovary.

hemk2-GLKD induced mRNA destabilization via No-Go Decay pathway

To understand the molecular defects that caused by *hemk2*-GLKD and

malfunction of eRF1, we performed the ovarian mRNA-seq analysis by next

generation sequencing. The late egg chambers in the control ovaries were cut

out to minimize the developmental difference between *hemk2*-GLKD and

control ovaries. Consistent with the RT-PCR results described above (Figure

1C and Figure 2C), the reductions of *hemk2* (log2FC=-1.27) and *otu*

(log2FC=-1.45) were severely downregulated in the mRNA-seq (Table S4). In

addition, 478 upregulated and 325 downregulated genes were detected

(log2FC < -1 or >1; q-val < 0.01). GO term analysis (Thomas et al., 2022) of

these differentially-expressed genes revealed that *hemk2*-GLKD resulted in

severe downregulation of many genes involved in the female reproductive

processes, such as female gamete generation, oogenesis and germ cell

development (Figure 6A). In contrast, genes involved in processes unrelated

to oogenesis were relatively upregulated (Figure 6B). To further examine the

impact of gene expression upon germline-knockdown of *hemk2*, we took

advantage of previous study reporting gene expression in different ovarian cell

types based on the single cell mRNA analysis; two germline and somatic cell

types were identified by the gene expression patterns. Among those, we

focused on genes expressed in 1 germline and 5 somatic cell types, the

represented cell types in early to mid-oogenesis to see the changes of mRNA

abundance. Of note, genes enriched in Germline 2 cell type (N=154), which

are expressed from region 2 onward, were most downregulated (average

log2FC = -0.55) in *hemk2*-GLKD versus control (Figure 6C) (Jevitt et al., 2020).

In contrast, the expression of genes enriched in 5 different somatic-cell types

did not show any significant change (Figure 6D; Figure S6A-E). These results

suggest that perturbation of eRF1 function in germline cells by

hemk2-knockdown specially downregulates mRNAs.

Previous studies have demonstrated that methylation of eRF1 at GGQ

motif enhances the release of nascent peptide and increases the rate of

translation termination by 10 to 100 times (Pierson et al., 2016). Thus, we

hypothesized that perturbation of the methylation on eRF1 reduced the release

rate of the nascent peptides, leading to the ribosome stalling at the stop

codons on the actively translated mRNAs. Such ribosome stalling is known to

trigger Ribosome-associated Quality Control (RQC), eliminating mRNAs being

actively translated with bound ribosomes in a process called no-go decay

(NGD) (Brandman and Hegde, 2016; Doma and Parker, 2006; Ikeuchi et al.,

2019; Simms et al., 2017; Tomomatsu et al., 2023). To address this possibility,

first, we investigated the ribosome stalling in *hemk2*-GLKD. We conducted

sucrose gradient sedimentation assay using *hemk2*-GLKD ovarian lysate to

examine whether ribosomes are collided by inefficient translation termination.

Ribosomes in the ovarian lysate were fixed with a translation inhibitor, emetine,

incubated with RNase to cleave the unprotected region of mRNA, and

analyzed through the sucrose gradient. The monosome was a major

population and a disome caused by a ribosome collision was hardly detected

in the control (Figure 6E). Upon *hemk2*-GLKD, however, the ratio of disome

significantly increased (Figure 6E, h*emk2*-GLKD), suggesting that the

inefficient release of the nascent polypeptide caused by unmethylated eRF1

resulted in the ribosome collision.

Next, we examined whether NGD contributed to severe defects observed

in *hemk2*-GLKD. To address this possibility, NGD pathway was simultaneously

knocked down together with *hemk2* in germline cells and the defects were

investigated (Figure 7 and Figure S7A). In the double-germline knockdown

(DGKD) ovaries of *hemk2* and *ski8*, a component of the Ski complex which

threads mRNAs into the exosome and promotes 3' to 5' degradation

(Anderson and Parker, 1998), the developmental defects in *hemk2*-GLKD

were partially rescued (Figure 7A). *ski8-hemk2* DGKD females contained later

stages of egg chambers and laid the comparable number of eggs, albeit the

hatching rate remained 40% of the control (Figure 7A, B). Because the Ski

complex is rather a component involved in general RNA degradation, we

further examined other factors which specifically function for NGD, such as

pelota (*pelo*), a critical effector of NGD pathway which functions for

dissociation of stalled ribosomes and RNA degradation (Becker et al., 2011;

Jamar et al., 2017; Kobayashi et al., 2010; Tsuboi et al., 2012). Intriguingly,

similar to *ski8-hemk2* DGKD, *pelo-hemk2* DGKD female also possessed later

egg chambers and laid the comparable numbers of eggs with 60% hatching

rate (Figure7A, B). We also examined another potential NGD component, an

endonuclease called *nonu-1*/*cue2* reported in *C. elegans* and *S. cerevisiae*

which cleaves mRNAs following the ribosome stalling during translation

(D'Orazio et al., 2019; Glover et al., 2020) (Figure S7B). Unlike *ski8* and *pelo*,

however, double-germline knockdown of *CG7139*, the *Drosophila* homolog of

nonu-1/*cue2*, and *hemk2* did not show any restoration of defects caused

(Figure 7A, B; Figure S7A). This suggests that in *Drosophila*, *CG7139* is not

essential in NGD pathway, or it may function redundantly with other

endonucleases during translation surveillance induced by *hemk2*-GLKD. In

addition, double-germline knockdown of *hemk2* and *upf3*, a factor involved in

nonsense-mediated decay (NMD) (Tan et al., 2022), did not rescue the defects

in *hemk2*-GLKD (Figure 7A, B), proving a specific action of NGD triggering

mRNA degradation. We confirmed the effective knocking down of each

examined component in ovaries by qRT-PCR (Figure S7A; ranging from 10 to

30 % compared to that of the control); considering the contribution of

non-knockdown somatic cell fractions to entire ovarian RNAs, the expression

of each factor was significantly reduced. Consistent with their capability to

rescue in the developmental morphology of *hemk2*-GLKD, the HPG assay

revealed that knockdown of *ski8* or *pelo*, but not *CG7139* or *upf3*, in

hemk2-GLKD ovaries partly rescued the reduction of translational activity

(Figure 7C, D). All together, these data suggest that the perturbation of eRF1

methylation in *hemk2*-GLKD ovaries leads to poor translation termination and

ribosome stalling, which triggers mRNA degradation specifically through the

NGD pathway.

Consistent with the observed restoration of oogenesis defects in

hemk2-GLKD ovaries, the expression levels of those downregulated

protein-coding RNAs were recovered by the additional germline knockdown of

ski8 or *pelo* (Figure S6F, G), further confirming that NGD was specifically

activated for the mRNA degradation upon *hemk2*-GLKD. Double-germline

knockdown of *ski8* or *pelo and hemk2* also restored the overall protein

synthesis, although the ribosome stalling was observed to similar extent of that

in the *hemk2*-single knockdown (Figure 6E). These results imply that

translation re-initiation could take place to synthesize proteins to some extent

when mRNAs are not cleared by NGD, partly restoring egg maturation (Figure

7A, B). Overall, these findings indicate that perturbation of eRF1 methylation in

hemk2-GLKD leads to inefficient release of synthesized peptides, triggering

RQC and activation of the NGD pathway, possibly by the ribosome stalling.

Discussion

The knockdown of *hemk2* resulted in female sterility, with a significant

disruption in oogenesis development observed in both ovarian germline and

somatic cells, leading to a developmental arrest in the mid-stage (Figure 1;

Figure S1). Curiously, *hemk2*-GLKD ovaries exhibited the morphological

defects, hyperplasia of the posterior follicle cells and failure in chromosomal

dispersion (Figure 2A, B). Previous studies reported that Delta ligand

expressed in germline cells Notch-Delta signaling triggers, inducing a

transition from mitotic cell cycle to endocycle in follicle cells (Deng et al., 2001;

Schaeffer et al., 2004; St. Johnston, 2001). The observed hyperplasia of the

posterior follicle cells in *hemk2*-GLKD ovaries may be attributed to the

downregulation of Delta ligand in germline cells, which in turn results in the

prolonged mitotic division of follicle cells even beyond stage 6.

In stage 6 and older egg chambers, almost all *hemk2*-GLKD ovarioles

exhibited the presence of undispersed chromosomes within nurse cell nuclei

(Figure 2B; Figure S2A). Some of genes involved in RNA regulation, such *otu*,

hfp, and *smn* were significantly downregulated in *hemk2*-GLKD ovaries (King

et al., 1981; Lee et al., 2009; Van Buskirk and Schüpbach, 2002). Similar to

that in *hemK2*-GLKD, mutations in these genes also produce defects in nurse

cell chromosome dispersal (Goodrich et al., 2004; Klusza and Deng, 2010; Lee

et al., 2009; Steinhauer and Kalfayan, 1992). Among those, one of two

isoforms coded by *otu* locus, Otu104, is known to function for the chromosome

dispersion of nurse cells (Goodrich et al., 2004; Klusza and Deng, 2010; Van

Buskirk and Schüpbach, 2002). Consistently, the expression of Otu104 slightly

rescued the chromosome dispersion defect observed in *hemk2*-GLKD (Figure

S2A). However, egg chambers remained small (Figure S2A), indicating the

impact of *hemk2*-GLKD on egg formation was not restored by Otu104

expression. Conversely, the repression of cell death in *hemk2*-GLKD ovaries

significantly rescued the egg formation in *hemk2*-GLKD ovaries, yet the nurse

cell chromosomes remained undispersed (Figure S2B). Collectively, our

results suggest that germline cell death in *hemk2*-GLKD is not attributed to the

chromosomal dispersion defect but possibly rather to ribosome stalling

resulting from the perturbation of eRF1 methylation (Wu et al., 2020). It has

been reported that the ribosome collision triggers two distinct stress response

pathways; transient collision evokes GCN2-mediated pathway for cell survival,

and severer collision activates MAPKK inducing apoptosis in human cell lines

(Wu et al., 2020). Our observation suggests that insufficient methylation by

hemk2 knockdown results in ribosome stalling and collision, thereby inducing

apoptosis in vivo, the translationally-active *Drosophila* ovaries.

Three potential substrates, DNA 6mA, histone H4K12 and eRF1, have

been previously identified for HemK2 in yeast and mammalian culture cells

(Figaro et al., 2008; Metzger et al., 2019; Xiao et al., 2018). Our study revealed

that *hemk2* knockdown did not affect methylation patterns of DNA 6mA or

histone H4K12 (Figure S3A, B), but caused a significant decrease in eRF1

methylation levels (Figure 3A-C), demonstrating the substrate specificity of

Drosophila HemK2 for eRF1. The 119[th] Tyr but not 116[th] Asn of *Drosophila*

HemK2 was indispensable and critical for its function in oogenesis and eRF1

methylation (Figure 4C-E). These results were discrepant from the fact that

both of the corresponding residues in human HemK2 were essential for eRF1

methylation *in vitro* (Metzger et al., 2019). Recent structural studies for human

HemK2, however, suggested that the side chain of Asn122 was not involved in

binding to methylated glutamine in the substrate biding pocket of HemK2 (Gao

et al., 2020), which is consistent with the dispensability of Asn116 for HemK2

methyltransferase activity.

The examination of murine HemK2's substrate specificity through a

peptide substrate array revealed a potential consensus motif:

G-Q-[SRYKLGAMTC]-[ARFGLWYCSQKH]-[LARCQFYT]-R, where [ ] denotes

any one residue among those listed inside (Kusevic et al., 2016). By using this

consensus motif, we identified several possible substrates of HemK2 among

Drosophila melanogaster proteins, such as CG17841, Spargel, Chd1, Ythdc1,

and eRF1. These potential substrates may be methylated by HemK2 in

particular conditions. However, the overexpression of the methylation-deficient

eRF1 Q185A mutant recapitulated the phenotypic characteristics observed in

hemk2-GLKD (Figure 5D), suggesting that eRF1 serves as the primary

substrate for *Drosophila* HemK2 in germline cells.

Previous studies have reported that the methylation of the conserved

GGQ motif in release factors significantly enhances the release of nascent

peptide and substantially increased the termination rate of protein synthesis by

a magnitude of 10 to 100 times (Pierson et al., 2016). In agreement with this,

the translation efficiency was remarkably reduced upon perturbation of eRF1

methylation by *hemk2*-GLKD (Figure 5E). Such a reduction in protein

synthesis as well as oogenesis defect and down regulation of mRNAs were

significantly restored by the blockade of a general RNA degradation or NGD

pathway, but not by NMD (Figure 7A-D). These results suggest that the

residual activity of the ribosome along with unmethylated eRF1 can manage to

synthesize the minimal amounts of polypeptides when mRNAs are provided

for the translation.

Our study provides evidence that HemK2 acts as a genuine protein

methyltransferase in *Drosophila* and plays a crucial role in modulating the

activity of eRF1 by glutamine methylation, which is indispensable for proper

germline development. Germline nurse cells produce large amounts of

proteins and mRNAs to transfer them to oocyte. Thus, it is important to

regulate the costly protein synthesis. However, the regulation mechanism of

eRF1 methylation by HemK2 remains to be fully elucidated and requires

further investigation. This mechanism may function to maximize egg formation

under favorable physiological conditions but suppress it under unfavorable

ones.

Materials and Methods

Fly stocks and cultures

All stocks and crosses were raised at room temperature (RT) or 25°C on standard food (5%(w/v) dry yeast, 5%(w/v) corn flower, 2%(w/v) rice bran, 10%(w/v) glucose, 0.7%(w/v) agar, 0.2%(v/v) propionic acid, 0.05%(v/v) p-hydroxy butyl benzoic acid). For knockdown experiments, the following TRiP lines obtained from Bloomington Drosophila Stock Center (IN, USA) were used in this study: *hemk2* (HMS02003, #40837), *eRF1* (HMS05812, #67900), *upf3* (HMJ22158, #58181), *cue2* (HMS02721, #44007), *ski8* (HMC04664, #57377), *p53* (HMS02286, #41720). attP40 (#25709) and attP2 (#25710) were used for injection. The following drivers were used to express transgenes or shRNAs: *NGT40* and *nosGal4-VP16* (Grieder et al., 2000), *Matα-tub-GAL4* (#7063), and *traffic jam-Gal4* (DGRC# 104055) (Hayashi et al., 2002). UASp-P35 transgenic line was a kind gift from Dr. Bergmann (Werz et al., 2005). Either the *y w* strain or the respective heterozygote was used as a control in each experiment.

Generation of transgenic fly lines

The generated transgenic lines were as follows: FLAG-HemK2 wildtype at P2, FLAG-HemK2 N116A at P2, FLAG-HemK2 Y119A at P2, FLAG-snapin at P2, FLAG-eRF1 wildtype at P2, FLAG-eRF1 Q185A at P2, FLAG-*otu*104 at P2. In each case, the coding sequence or its mutant form was amplified by PCR using *y w* genomic DNA (gDNA) or the cDNAs obtained by reverse

transcription from ovarian RNA of *y w* flies as template, and triple repeats of

the FLAG tag were added to the N-terminal region. Purified PCR fragments

were cloned into the *Xba*I-linearized pUASp-K10-attB vector (Koch et al., 2009)

by In-Fusion reaction (#639648, Takara Bio, Japan). The positive plasmids

were injected into embryo of nos-phiC31;;P{CaryP} attP2 (#25710) by using

the standard protocol.

hemk2 TRiP at P2, and *pelo* TRiP at P2 and P40 were generated in this

study. TRiP constructs were generated by introducing the annealed oligos

(TRiP-Fw/Rv) into *Nhe*I- and *Eco*RI-linearized pWalium22 vector (Perkins et

al., 2015) by DNA ligation. The UASp and Walium22 plasmids were then

injected into embryo of nos-phiC31; P{CaryP}attP40 (#25709) or

nos-phiC31;;P{CaryP} attP2 (#25710). The oligos used for annealing are listed

in Table S1.

Fluorescent immunostaining

Immunostaining was performed as previously reported (Lim et al., 2022).

Ovaries were dissected and teased apart slightly in phosphate-buffered saline

(PBS), followed by fixation with 5.3% (v/v) paraformaldehyde (PFA, Electron

Microscopy Sciences, United States) in PBS for 10 min at RT. Fixed ovaries

were washed for 30 min with several changes of PBX (PBS containing 0.2%

(v/v) Triton X-100), and incubated with PBX containing 4% (w/v) bovine serum

albumin (BSA) for at least 30 min of blocking. The incubation with primary

antibodies in PBX containing 0.4% (w/v) BSA was performed overnight at 4°C,

and ovaries were washed for 1 h with several changes of PBX. The secondary

antibody incubation was then conducted for >2 h at room temperature, and

ovaries were washed for 1 h with PBX and stained with 1 μg/ml of

4′6-diamidine-2′-phenylindole (DAPI) (SIGMA, United States) for 10 min in

PBX. After rinsing with PBS, the ovaries were mounted with Fluoro-KEEPER

Antifade (Nacalai Tesque, Japan) onto a slide.

The following primary antibodies were used in the indicated dilution;

mouse anti-FLAG (SIGMA [M2], Cat.# F1804, 1:200), guinea pig anti-Vas

(1:2,000) (Patil and Kai, 2010), rabbit anti-pH3 (Millipore, Cat.# 06-570, 1:250),

mouse anti-p53 (DSHB [25F4], 1:100), rabbit anti-Fibrillarin (Abcam, Cat.#

ab5821, 1:200), rabbit anti-DNA 6mA (pAbmAbs, Cat.# 202003, 1:1,000),

rabbit anti-eRF1 (SIGMA, Cat.# E8156, 1:200), rabbit anti-methyl-eRF1 (1:100)

(Lacoux et al., 2020). The following secondary antibodies were used in this

study; Alexa Fluor 488-, 555- conjugated goat antibodies against rabbit,

mouse, and guinea pig IgG (Invitrogen, Cat.#: A11029, A11034, A11073,

A21424, A21429, A21435) were used at 1:200 in PBX containing 0.4% (w/v)

BSA. Images of the ovaries were taken by Zeiss LSM 900 confocal

microscope and processed using Zen (Zeiss) and Fiji (Schindelin et al., 2012)

software.

Protein synthesis assay

The Click-iT® HPG Alexa Fluor 488 Protein Synthesis Assay kit (Molecular

probes, United States) was used to monitor newly synthesized proteins by

quantification of fluorescent HPG signals. The freshly dissected ovaries were

treated with 1 mM HPG in PBS for 30 min and fixed in 5.3% PFA in PBS for 10

min. Subsequently, the ovaries were permeabilized with PBX for 30 min by

washing three times (3% BSA/PBX for the first time, PBX only for the second

and third time). They were then incubated with a freshly prepared Click-iT®

reaction cocktail for 30 min. After washing with Click-iT® reaction rinse buffer,

the ovaries were incubated with HCS NuclearMask™ Blue Stain working

solution (1:2,000 in PBX) for 30 min according to manufacturer's protocol. The

samples were further immunostained with antibodies, and the images were

taken as described above.

Quantification analysis of HPG signal for translation activity

Each image was processed with Fiji software. The fluorescence intensity

of HPG signals in the cytosolic (non-nucleus) regions of nurse cells was

measured. Each small square (in identical size) was calculated after

background (blank area without any imaging) subtraction, and 2-5 squares per

egg chamber (stage 3-8) were analyzed. The mean value of all squares per

image was quantified as translation activity and 5 images were quantified per

genotype.

Immunoprecipitation and Western blotting

One hundred ovaries were dissected in cold PBS and homogenized with a

pestle in PBST (PBS with 0.1%(v/v) Tween-20) supplemented with proteinase

inhibitor cocktail (PIC, #11873580001, Roche, Switzerland). After

centrifugation at 15,000 xg for 10 min at 4°C, supernatant was collected, 10 µl

of Protein G magnetic beads (TAMAGAWA SEIKI, #TAS8848N1173) was

incubated with FLAG antibody (SIGMA [M2], Cat.# F1804) or (5 µl beads for

eRF1 case) for more than 1 h at 4°C with gentle rotation before adding lysate.

Subsequently the antibody-conjugated beads were washed twice with cold

PBST and incubated with lysate at 4°C for 3-4 h. Proteins were rinsed with

cold PBST three times and eluted from beads in 50 µl of sample buffer at 95°C

for 5 min.

Protein lysates or immunoprecipitates were separated by SDS-PAGE and

transferred to a 0.45 µm Clear Trans® PVDF membrane (Wako, Japan). The

membrane was blocked in PBST with 3% (w/v) skim milk (Nacalai Tesque,

Japan) and incubated with the following primary antibodies diluted in the signal

enhancer reagent HIKARI (Nacalai Tesque): rabbit anti-eRF1 (SIGMA, Cat.#

E8156, 1:1,000), rabbit anti-Methyl-Q (Millipore, Cat.# ABS2185, 1:1000),

mouse anti-FLAG (SIGMA [M2], Cat.# F1804, 1:1,000), mouse anti-c-Myc

(Wako [9E10], Cat.# 017-21871, 1:1,000), mouse anti-histone H4 (MAB

Institute, Cat.# 388-09171, 1:2000), rabbit anti-H4K12met (PTM BIO, Cat.#

PTM-685RM, 1:2,000), mouse anti-Tubulin (Santa Cruz [DM1A], Cat.#

sc-32293, 1:3,000), rabbit anti-HA (Cell signaling Technology [C29F4], Cat.#

3724, 1:1,000).

HRP-conjugated secondary antibodies used in this study were as follows:

goat anti-rabbit IgG, or anti-mouse IgG (1:3,000; #170-6515, #170-6516,

Bio-Rad, CA, United States). To avoid signals from IgG heavy chain

overlapping with eRF1, VeriBlot-HRP (Abcam, #ab131366, 1:1,000) was used

in Western blot analysis for immunoprecipitation of endogenous eRF1 or

FLAG-eRF1. Chemi-Lumi One (Nacalai Tesque, #07880-70) was used to

obtain the chemiluminescent signals, which were then imaged by ChemiDoc™

Touch MP (Bio-Rad). Band intensities were quantified from independent

experiments by Image Lab software (Bio-Rad).

Quantitative RT-PCR

Total RNAs were extracted from ovaries using TRIzol LS (Invitrogen) in

strict accordance with the manufacturer's instructions. Following this, the

purified RNA (1 µg) was subjected to DNase I treatment (Invitrogen) to

eliminate any genomic contamination, which was subsequently inactivated

using EDTA. cDNAs were synthesized by reverse transcription with

Superscript III system (Invitrogen) following the manufacturer's protocol. The

qPCR experiment was carried out using KAPA SYBR Fast qPCR Master Mix

(KAPA biosystems) in the StepOnePlus Real-Time PCR system (Applied

Biosystems). The relative expression levels were normalized to those of *rp49*

with three biological replicates and technical duplicates. The primer sequences

for detecting the expression levels of genes are shown in Table S2.

S2 cell experiments

Drosophila S2 cells were cultured at 28°C in Schneider medium

containing 10% (v/v) Fetal Bovine Serum (FBS), supplemented with penicillin

and streptomycin. The endogenous *hemk2* was knocked down by dsRNA

targeting a 186bp region starting from ATG codon of *snapin* gene (Figure S1).

The dsRNA was generated by *in vitro* AmpliScribe T7 High Yield transcription

kit (Lucigen, United States). The primers used for PCR are listed in Table S3.

Plasmids used for transfection were as follows: Myc-HemK2 wildtype,

Myc-HemK2 N116A, Myc-HemK2 Y119A, FLAG-eRF1 wildtype, FLAG-eRF1

Q185A, Myc-HemK1. The constructions were generated using the Gateway

system (Life technologies) and transfected by HilyMax (Dojindo Molecular

Technologies, Inc, Japan) following the manufacturer's instructions. The

immunoprecipitation and immunoblotting were conducted as described above.

Fertility test

To test female fertility, *y w* males were mated with three eclosed females

of each genotype for few days in vials with normal food, and then transferred to

apple juice plates with a bit of wet yeast paste. The total number of eggs laid

by mated females was counted, and the unhatched eggs were quantified to

determine the hatching rates after 24 hours of egg laying. To analyze the

fertility of *hemk2*-GLKD males, three *y w* virgin females were mated with

hemk2-GLKD males for longer time, and the following statistics were recorded.

Each experiment was conducted in three biological replicates.

mRNA-seq analyses

Total RNAs of each genotype in two biological replicates were extracted

from dissected ovaries using TRIzol LS (Invitrogen) in accordance with the

manufacturer's protocol. Because *hemk2* knockdown ovaries did not contain

later stages, regions later than mid oogenesis were eliminated by cutting the

ovaries for RNA preparation to minimize the developmental stage difference.

Poly(A)-tailed RNAs were isolated and concentrated using oligo-dT beads in

NEBNext® Poly(A) mRNA Magnetic Isolation Module (New England Biolabs,

USA), subsequently subjected to fragmentation and reverse transcription.

After ligation of adapters, PCR amplification was performed to generate the

libraries (NEBNext® UltraTM II Directional RNA Library Prep Kit, New England

Biolabs, USA). The cDNA library preparation and next-generation sequencing

were performed by Rhelixa Inc (Japan). Around 13 million paired sequences

were obtained for each sample. Reads with low quality and adapter sequences

were removed by fastp program (Chen et al., 2018). Subsequently, reads were

mapped on the *Drosophila* melanogaster genome reference (dm6) by STAR

program (Dobin et al., 2013). The counts of the mapped reads for genes were

analyzed by featureCounts (Liao et al., 2014). The statistical analysis for the

differential expression genes were performed using DESeq2 (Love et al.,

2014). The upregulated genes (log2FoldChnage > 1, p-adj < 0.01)) and

downregulated genes (log2FoldChange < -1, p-adj < 0.01) were selected and

subjected to the GO term overrepresentation analysis by PANTHER (Thomas

et al., 2022).

Monosome/Disome profiling

Ovaries were dissected in polysome lysis buffer (20 mM Tris-Cl, pH 7.5, 5

mM $MgCl_2$, 100 mM NaCl) supplemented with 0.1% NP-40, 200 µg/ml emetine

(E2375, SIGMA), and PIC. The dissected ovaries were snap frozen in liquid

nitrogen and stored at -80ºC with minimum volume of the buffer. Around 30

pairs of ovaries were homogenized with plastic pestle in 500 µl of polysome

lysis buffer supplemented with 0.1% CHAPS (Nacalai Tesque, Japan), 200

µg/ml emetine, and PIC. The lysate was centrifuged at 15,000 xg for 5 min at

4ºC. Supernatant was further centrifuged at 15,000 xg for 15 min at 4ºC and

250 µl of the supernatant was collected. The RNA concentration in the

supernatant was determined by Qubit RNA HS assay kit (Q32852, Invitrogen).

The RNA concentration in the supernatant was adjusted to 0.05 µg/µl by

adding polysome lysis buffer supplemented with 0.1% CHAPS, 200 µg/ml

emetine, and PIC. The resulted lysate was incubated with RNase A (1 µg/ml)

for 10 min at 25ºC to cleave the unprotected region of RNA. 200 µl of the

RNase A-treated sample was loaded on the top of the sucrose gradient

(10%-50% sucrose in polysome lysis buffer) in SW41Ti and centrifuged at

30,000 xg for 3 h at 4ºC. After centrifugation, the gradients were fractionated

from the top using piston gradient fractionator (BioComp, Canada).

Acknowledgments

We thank Dr Valérie Heurgué-Hamard (IBPC, Paris) for a generous gift of antibody against methyl-eRF1. We acknowledge Bloomington Drosophila Stock Centre and Kyoto Stock Center for the fly stocks. Assistance with sucrose gradient analysis was provided by the Center for Medical Research and Education, Graduate School of Medicine, Osaka University. We appreciate the insightful discussion and suggestions from all the members of KT's laboratory.

Author Contributions

All authors contributed to the experiment design. XF, SR, KS, and KT performed all experiments. XF, SR and KS performed the computational analyses. XF, SR, KS, and KT wrote the paper. KS, IT, and KT supervised the project.

Funding Statement

This work was supported by TAKEDA Bioscience Research Grant (J191503009) for KT, Grant-in-Aid for Scientific Research on Innovative Areas (20H05390) for KT, Grant-in-Aid for Transformative Research Areas (A) (21H05275) for KT, and Osaka University Institute for Datability Science "Transdisciplinary Research Project" (Na22990007) for KT and KS.

Data And Material Availability

NGS data sets have been deposited to the DNA Data Bank of Japan (DDBJ).

BioProject Accession: PRJDB17326. All fly strains generated for this study are

available upon request.

Conflict Of Interest Statement

The authors declare that the research was conducted in the absence of any

commercial or financial relationships that could be construed as a potential

conflict of interest.

References

Anderson, J.S., Parker, R.P., 1998. The 3' to 5' degradation of yeast mRNAs is a general mechanism for mRNA turnover that requires the SKI2 DEVH box protein and 3' to 5' exonucleases of the exosome complex. EMBO J. 17, 1497–1506. https://doi.org/10.1093/emboj/17.5.1497

Avilés-Pagán, E.E., Orr-Weaver, T.L., 2018. Activating embryonic development in Drosophila. Semin. Cell Dev. Biol. 84, 100–110. https://doi.org/10.1016/j.semcdb.2018.02.019

Becker, T., Armache, J.-P., Jarasch, A., Anger, A.M., Villa, E., Sieber, H., Motaal, B.A., Mielke, T., Berninghausen, O., Beckmann, R., 2011. Structure of the no-go mRNA decay complex Dom34–Hbs1 bound to a stalled 80S ribosome. Nat. Struct. Mol. Biol. 18, 715–720. https://doi.org/10.1038/nsmb.2057

Brandman, O., Hegde, R.S., 2016. Ribosome-associated protein quality control. Nat. Struct. Mol. Biol. 23, 7–15. https://doi.org/10.1038/nsmb.3147

Chen, C.-C., Wang, K.-Y., Shen, C.-K.J., 2013. DNA 5-methylcytosine demethylation activities of the mammalian DNA methyltransferases. J. Biol. Chem. 288, 9084–9091. https://doi.org/10.1074/jbc.M112.445585

Chen, S., Zhou, Y., Chen, Y., Gu, J., 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics 34, i884–i890. https://doi.org/10.1093/bioinformatics/bty560

Clarke, S.G., 2013. Protein methylation at the surface and buried deep: thinking outside the histone box. Trends Biochem. Sci. 38, 243–252. https://doi.org/10.1016/j.tibs.2013.02.004

Dai, X., Ren, T., Zhang, Y., Nan, N., 2021. Methylation multiplicity and its

clinical values in cancer. Expert Rev. Mol. Med. 23, e2.

https://doi.org/10.1017/erm.2021.4

Dej, K.J., Spradling, A.C., 1999. The endocycle controls nurse cell polytene

chromosome structure during *Drosophila* oogenesis. Development 126,

293–303. https://doi.org/10.1242/dev.126.2.293

Deng, W.-M., Althauser, C., Ruohola-Baker, H., 2001. Notch-Delta signaling

induces a transition from mitotic cell cycle to endocycle in *Drosophila*

follicle cells. Development 128, 4737–4746.

https://doi.org/10.1242/dev.128.23.4737

Dinçbas-Renqvist, V., Engström, Å., Mora, L., Heurgué-Hamard, V.,

Buckingham, R., Ehrenberg, M., 2000. A post-translational modification

in the GGQ motif of RF2 from Escherichia coli stimulates termination of

translation. EMBO J. 19, 6900–6907.

https://doi.org/10.1093/emboj/19.24.6900

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut,

P., Chaisson, M., Gingeras, T.R., 2013. STAR: ultrafast universal

RNA-seq aligner. Bioinformatics 29, 15–21.

https://doi.org/10.1093/bioinformatics/bts635

Doma, M.K., Parker, R., 2006. Endonucleolytic cleavage of eukaryotic mRNAs

with stalls in translation elongation. Nature 440, 561–564.

https://doi.org/10.1038/nature04530

D'Orazio, K.N., Wu, C.C.-C., Sinha, N., Loll-Krippleber, R., Brown, G.W.,

Green, R., 2019. The endonuclease Cue2 cleaves mRNAs at stalled

ribosomes during No Go Decay. eLife 8, e49117.

https://doi.org/10.7554/eLife.49117

Drummond-Barbosa, D., Spradling, A.C., 2001. Stem Cells and Their Progeny

Respond to Nutritional Changes during Drosophila Oogenesis. Dev.

Biol. 231, 265–278. https://doi.org/10.1006/dbio.2000.0135

Fang, Q., Kimura, Y., Shimazu, T., Suzuki, T., Yamada, A., Dohmae, N.,

Iwasaki, S., Shinkai, Y., 2022. Mammalian HEMK1 methylates

glutamine residue of the GGQ motif of mitochondrial release factors. Sci.

Rep. 12, 4104. https://doi.org/10.1038/s41598-022-08061-y

Figaro, S., Scrima, N., Buckingham, R.H., Heurgué-Hamard, V., 2008. HemK2

protein, encoded on human chromosome 21, methylates translation

termination factor eRF1. FEBS Lett. 582, 2352–2356.

https://doi.org/10.1016/j.febslet.2008.05.045

Fuchs, Y., Steller, H., 2015. Live to die another way: modes of programmed

cell death and the signals emanating from dying cells. Nat. Rev. Mol.

Cell Biol. 16, 329–344. https://doi.org/10.1038/nrm3999

Gao, J., Wang, B., Yu, H., Wu, G., Wan, C., Liu, W., Liao, S., Cheng, L., Zhu,

Z., 2020. Structural insight into HEMK2–TRMT112-mediated glutamine

methylation. Biochem. J. 477, 3833–3838.

https://doi.org/10.1042/BCJ20200594

Gates, L.A., Foulds, C.E., O'Malley, B.W., 2017. Histone Marks in the 'Driver's

Seat': Functional Roles in Steering the Transcription Cycle. Trends

Biochem. Sci. 42, 977–989. https://doi.org/10.1016/j.tibs.2017.10.004

Gleason, R.J., Anand, A., Kai, T., Chen, X., 2018. Protecting and Diversifying

the Germline. Genetics 208, 435–471.

https://doi.org/10.1534/genetics.117.300208

Glover, M.L., Burroughs, A.Max., Monem, P.C., Egelhofer, T.A., Pule, M.N.,

Aravind, L., Arribere, J.A., 2020. NONU-1 Encodes a Conserved

Endonuclease Required for mRNA Translation Surveillance. Cell Rep.

30, 4321-4331.e4. https://doi.org/10.1016/j.celrep.2020.03.023

Goodrich, J.S., Clouse, K.N., Schüpbach, T., 2004. Hrb27C, Sqd and Otu

cooperatively regulate *gurken* RNA localization and mediate nurse cell

chromosome dispersion in *Drosophila* oogenesis. Development 131,

1949–1958. https://doi.org/10.1242/dev.01078

Grieder, N.C., Cuevas, M.D., Spradling, A.C., 2000. The fusome organizes the

microtubule network during oocyte differentiation in *Drosophila*.

Development 127, 4253–4264. https://doi.org/10.1242/dev.127.19.4253

Hayashi, S., Ito, K., Sado, Y., Taniguchi, M., Akimoto, A., Takeuchi, H., Aigaki,

T., Matsuzaki, F., Nakagoshi, H., Tanimura, T., Ueda, R., Uemura, T.,

Yoshihara, M., Goto, S., 2002. GETDB, a database compiling

expression patterns and molecular locations of a collection of gal4

enhancer traps. genesis 34, 58–61. https://doi.org/10.1002/gene.10137

Heurgué-Hamard, V., Champ, S., Engström, A., Ehrenberg, M., Buckingham,

R.H., 2002. The hemK gene in Escherichia coli encodes the

N(5)-glutamine methyltransferase that modifies peptide release factors.

EMBO J. 21, 769–778. https://doi.org/10.1093/emboj/21.4.769

Heurgué-Hamard, V., Champ, S., Mora, L., Merkoulova-Rainon, T., Kisselev,

L.L., Buckingham, R.H., 2005. The Glutamine Residue of the

Conserved GGQ Motif in Saccharomyces cerevisiae Release Factor

eRF1 Is Methylated by the Product of the YDR140w Gene *. J. Biol.

Chem. 280, 2439–2445. https://doi.org/10.1074/jbc.M407252200

Heurgué-Hamard, V., Graille, M., Scrima, N., Ulryck, N., Champ, S., van

Tilbeurgh, H., Buckingham, R.H., 2006. The zinc finger protein Ynr046w

is plurifunctional and a component of the eRF1 methyltransferase in

yeast. J. Biol. Chem. 281, 36140–36148.

https://doi.org/10.1074/jbc.M608571200

Ikeuchi, K., Izawa, T., Inada, T., 2019. Recent Progress on the Molecular

Mechanism of Quality Controls Induced by Ribosome Stalling. Front.

Genet. 9, 743. https://doi.org/10.3389/fgene.2018.00743

Ishizawa, T., Nozaki, Y., Ueda, T., Takeuchi, N., 2008. The human

mitochondrial translation release factor HMRF1L is methylated in the

GGQ motif by the methyltransferase HMPrmC. Biochem. Biophys. Res.

Commun. 373, 99–103. https://doi.org/10.1016/j.bbrc.2008.05.176

Jamar, N.H., Kritsiligkou, P., Grant, C.M., 2017. The non-stop decay mRNA

surveillance pathway is required for oxidative stress tolerance. Nucleic

Acids Res. 45, 6881–6893. https://doi.org/10.1093/nar/gkx306

Jevitt, A., Chatterjee, D., Xie, G., Wang, X.-F., Otwell, T., Huang, Y.-C., Deng,

W.-M., 2020. A single-cell atlas of adult Drosophila ovary identifies

transcriptional programs and somatic cell lineage regulating oogenesis.

PLOS Biol. 18, e3000538. https://doi.org/10.1371/journal.pbio.3000538

King, R.C., Riley, S.F., Cassidy, J.D., White, P.E., Paik, Y.K., 1981. Giant

Polytene Chromosomes from the Ovaries of a Drosophila Mutant.

Science 212, 441–443. https://doi.org/10.1126/science.6782674

Klotz, A.V., Thomas, B.A., Glazer, A.N., Blacher, R.W., 1990. Detection of

methylated asparagine and glutamine residues in polypeptides. Anal.

Biochem. 186, 95–100. https://doi.org/10.1016/0003-2697(90)90579-X

Klusza, S., Deng, W.-M., 2010. poly is required for nurse-cell chromosome

dispersal and oocyte polarity in Drosophila. Fly (Austin) 4, 128–136.

Klusza, S., Novak, A., Figueroa, S., Palmer, W., Deng, W.-M., 2013. Prp22

and Spliceosome Components Regulate Chromatin Dynamics in

Germ-Line Polyploid Cells. PLoS ONE 8, e79048.

https://doi.org/10.1371/journal.pone.0079048

Kobayashi, K., Kikuno, I., Kuroha, K., Saito, K., Ito, K., Ishitani, R., Inada, T.,

Nureki, O., 2010. Structural basis for mRNA surveillance by archaeal

Pelota and GTP-bound EF1α complex. Proc. Natl. Acad. Sci. U. S. A.

107, 17575–17579. https://doi.org/10.1073/pnas.1009598107

Koch, R., Ledermann, R., Urwyler, O., Heller, M., Suter, B., 2009. Systematic

Functional Analysis of Bicaudal-D Serine Phosphorylation and

Intragenic Suppression of a Female Sterile Allele of BicD. PLoS ONE 4,

e4552. https://doi.org/10.1371/journal.pone.0004552

Kusevic, D., Kudithipudi, S., Jeltsch, A., 2016. Substrate Specificity of the

HEMK2 Protein Glutamine Methyltransferase and Identification of Novel

Substrates. J. Biol. Chem. 291, 6124–6133.

https://doi.org/10.1074/jbc.M115.711952

Kweon, S.-M., Chen, Y., Moon, E., Kvederaviciutė, K., Klimasauskas, S.,

Feldman, D.E., 2019. An Adversarial DNA N6-Methyladenine-Sensor

Network Preserves Polycomb Silencing. Mol. Cell 74, 1138-1147.e6.

https://doi.org/10.1016/j.molcel.2019.03.018

Lacoux, C., Wacheul, L., Saraf, K., Pythoud, N., Huvelle, E., Figaro, S., Graille,

M., Carapito, C., Lafontaine, D.L.J., Heurgué-Hamard, V., 2020. The

catalytic activity of the translation termination factor methyltransferase

Mtq2-Trm112 complex is required for large ribosomal subunit

biogenesis. Nucleic Acids Res. 48, 12310–12325.

https://doi.org/10.1093/nar/gkaa972

Lasko, P., 2012. mRNA Localization and Translational Control in Drosophila

Oogenesis. Cold Spring Harb. Perspect. Biol. 4, a012294.

https://doi.org/10.1101/cshperspect.a012294

Lee, L., Davies, S.E., Liu, J.-L., 2009. The spinal muscular atrophy protein

SMN affects Drosophila germline nuclear organization through the U

body–P body pathway. Dev. Biol., Special Section: Evolution of

Developmental Regulatory Systems 332, 142–155.

https://doi.org/10.1016/j.ydbio.2009.05.553

Liao, Y., Smyth, G.K., Shi, W., 2014. featureCounts: an efficient general

purpose program for assigning sequence reads to genomic features.

Bioinformatics 30, 923–930.

https://doi.org/10.1093/bioinformatics/btt656

Lim, L.-X., Isshiki, W., Iki, T., Kawaguchi, S., Kai, T., 2022. The Tudor

Domain-Containing Protein, Kotsubu (CG9925), Localizes to the Nuage

and Functions in piRNA Biogenesis in D. melanogaster. Front. Mol.

Biosci. 9, 818302. https://doi.org/10.3389/fmolb.2022.818302

Liu, P., Nie, S., Li, B., Yang, Z.-Q., Xu, Z.-M., Fei, J., Lin, C., Zeng, R., Xu,

G.-L., 2010. Deficiency in a Glutamine-Specific Methyltransferase for

Release Factor Causes Mouse Embryonic Lethality. Mol. Cell. Biol. 30,

4245–4253. https://doi.org/10.1128/MCB.00218-10

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change

and dispersion for RNA-seq data with DESeq2. Genome Biol. 15, 550.

https://doi.org/10.1186/s13059-014-0550-8

McCall, K., 2004. Eggs over easy: cell death in the Drosophila ovary. Dev. Biol.

274, 3–14. https://doi.org/10.1016/j.ydbio.2004.07.017

Metzger, E., Wang, S., Urban, S., Willmann, D., Schmidt, A., Offermann, A.,

Allen, A., Sum, M., Obier, N., Cottard, F., Ulferts, S., Preca, B.-T.,

Hermann, B., Maurer, J., Greschik, H., Hornung, V., Einsle, O., Perner,

S., Imhof, A., Jung, M., Schüle, R., 2019. KMT9 monomethylates

histone H4 lysine 12 and controls proliferation of prostate cancer cells.

Nat. Struct. Mol. Biol. 26, 361–371.

https://doi.org/10.1038/s41594-019-0219-9

Patil, V.S., Kai, T., 2010. Repression of Retroelements in Drosophila Germline

via piRNA Pathway by the Tudor Domain Protein Tejas. Curr. Biol. 20,

724–730. https://doi.org/10.1016/j.cub.2010.02.046

Perkins, L.A., Holderbaum, L., Tao, R., Hu, Y., Sopko, R., McCall, K.,

Yang-Zhou, D., Flockhart, I., Binari, R., Shim, H.-S., Miller, A., Housden,

A., Foos, M., Randkelv, S., Kelley, C., Namgyal, P., Villalta, C., Liu,

L.-P., Jiang, X., Huan-Huan, Q., Wang, X., Fujiyama, A., Toyoda, A.,

Ayers, K., Blum, A., Czech, B., Neumuller, R., Yan, D., Cavallaro, A.,

Hibbard, K., Hall, D., Cooley, L., Hannon, G.J., Lehmann, R., Parks, A.,

Mohr, S.E., Ueda, R., Kondo, S., Ni, J.-Q., Perrimon, N., 2015. The

Transgenic RNAi Project at Harvard Medical School: Resources and

Validation. Genetics 201, 843–852.

https://doi.org/10.1534/genetics.115.180208

Pierson, W.E., Hoffer, E.D., Keedy, H.E., Simms, C.L., Dunham, C.M., Zaher,

H.S., 2016. Uniformity of Peptide Release Is Maintained by Methylation

of Release Factors. Cell Rep. 17, 11–18.

https://doi.org/10.1016/j.celrep.2016.08.085

Ping, X.-L., Sun, B.-F., Wang, L., Xiao, W., Yang, X., Wang, W.-J., Adhikari, S.,

Shi, Y., Lv, Y., Chen, Y.-S., Zhao, X., Li, A., Yang, Y., Dahal, U., Lou,

X.-M., Liu, X., Huang, J., Yuan, W.-P., Zhu, X.-F., Cheng, T., Zhao,

Y.-L., Wang, X., Danielsen, J.M.R., Liu, F., Yang, Y.-G., 2014.

Mammalian WTAP is a regulatory subunit of the RNA

N6-methyladenosine methyltransferase. Cell Res. 24, 177–189.

https://doi.org/10.1038/cr.2014.3

Polevoda, B., Span, L., Sherman, F., 2006. The Yeast Translation Release

Factors Mrf1p and Sup45p (eRF1) Are Methylated, Respectively, by the

Methyltransferases Mtq1p and Mtq2p*. J. Biol. Chem. 281, 2562–2571.

https://doi.org/10.1074/jbc.M507651200

Ramazi, S., Zahiri, J., 2021. Post-translational modifications in proteins:

resources, tools and prediction methods. Database 2021, baab012.

https://doi.org/10.1093/database/baab012

Rastegari, E., Kajal, K., Tan, B.-S., Huang, F., Chen, R.-H., Hsieh, T.-S., Hsu,

H.-J., 2020. WD40 protein Wuho controls germline homeostasis via

TRIM-NHL tumor suppressor Mei-p26 in *Drosophila*. Development 147,

dev182063. https://doi.org/10.1242/dev.182063

Ratel, D., Ravanat, J.-L., Charles, M.-P., Platet, N., Breuillaud, L., Lunardi, J.,

Berger, F., Wion, D., 2006. Undetectable levels of N6-methyl adenine in

mouse DNA: Cloning and analysis of PRED28, a gene coding for a

putative mammalian DNA adenine methyltransferase. FEBS Lett. 580,

3179–3184. https://doi.org/10.1016/j.febslet.2006.04.074

Ryšlavá, H., Doubnerová, V., Kavan, D., Vaněk, O., 2013. Effect of

posttranslational modifications on enzyme function and assembly. J.

Proteomics 92, 80–109. https://doi.org/10.1016/j.jprot.2013.03.025

Schaeffer, V., Althauser, C., Shcherbata, H.R., Deng, W.-M., Ruohola-Baker,

H., 2004. Notch-Dependent Fizzy-Related/Hec1/Cdh1 Expression Is

Required for the Mitotic-to-Endocycle Transition in Drosophila Follicle

Cells. Curr. Biol. 14, 630–636.

https://doi.org/10.1016/j.cub.2004.03.040

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M.,

Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B.,

Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K., Tomancak, P.,

Cardona, A., 2012. Fiji - an Open Source platform for biological image

analysis. Nat. Methods 9, 10.1038/nmeth.2019.

https://doi.org/10.1038/nmeth.2019

Shen, Y., Liu, W., Zuo, J., Han, J., Zhang, Z.C., 2021. Protocol for visualizing

newly synthesized proteins in primary mouse hepatocytes. STAR

Protoc. 2, 100616. https://doi.org/10.1016/j.xpro.2021.100616

Simms, C.L., Yan, L.L., Zaher, H.S., 2017. Ribosome collision is critical for

quality control during no-go decay. Mol. Cell 68, 361-373.e5.

https://doi.org/10.1016/j.molcel.2017.08.019

Sprung, R., Chen, Y., Zhang, K., Cheng, D., Zhang, T., Peng, J., Zhao, Y.,

2008. Identification and validation of eukaryotic aspartate and

glutamate methylation in proteins. J. Proteome Res. 7, 1001–1006.

https://doi.org/10.1021/pr0705338

St. Johnston, H.L.-S. and D., 2001. Delta signaling from the germ line controls

986 the proliferation and differentiation of the somatic follicle cells during

987 Drosophila oogenesis. Genes Dev. 15, 1393–1405.

988 https://doi.org/10.1101/gad.200901

989 Steinhauer, W.R., Kalfayan, L.J., 1992. A specific ovarian tumor protein

990 isoform is required for efficient differentiation of germ cells in Drosophila

991 oogenesis. Genes Dev. 6, 233–243. https://doi.org/10.1101/gad.6.2.233

992 Svidritskiy, E., Korostelev, A.A., 2018. Conformational control of translation

993 termination on the 70S ribosome. Struct. Lond. Engl. 1993 26,

994 821-828.e3. https://doi.org/10.1016/j.str.2018.04.001

995 Tan, K., Stupack, D.G., Wilkinson, M.F., 2022. Nonsense-mediated RNA

996 decay: an emerging modulator of malignancy. Nat. Rev. Cancer 22,

997 437–451. https://doi.org/10.1038/s41568-022-00481-2

998 Thomas, P.D., Ebert, D., Muruganujan, A., Mushayahama, T., Albou, L.-P., Mi,

999 H., 2022. PANTHER: Making genome-scale phylogenetics accessible

1000 to all. Protein Sci. 31, 8–22. https://doi.org/10.1002/pro.4218

1001 Tomomatsu S., Watanabe A., Tesina P., Hashimoto S., Ikeuchi K., Li S.,

1002 Matsuo Y., Beckmann R., Inada T., 2023. Two modes of

1003 Cue2-mediated mRNA cleavage with distinct substrate recognition

1004 initiate no-go decay. Nucleic Acids Res. 51, 253–270.

1005 https://doi.org/10.1093/nar/gkac1172

1006 Tsuboi, T., Kuroha, K., Kudo, K., Makino, S., Inoue, E., Kashima, I., Inada, T.,

1007 2012. Dom34:Hbs1 Plays a General Role in Quality-Control Systems by

1008 Dissociation of a Stalled Ribosome at the 3′ End of Aberrant mRNA. Mol.

1009 Cell 46, 518–529. https://doi.org/10.1016/j.molcel.2012.03.013

1010 Van Buskirk, C., Schüpbach, T., 2002. half pint Regulates Alternative Splice

Site Selection in Drosophila. Dev. Cell 2, 343–353.

https://doi.org/10.1016/S1534-5807(02)00128-4

Wall, A.A., Phillips, A.M., Kelly, L.E., 2005. Effective Translation of the Second

Cistron in Two Drosophila Dicistronic Transcripts Is Determined by the

Absence of In-frame AUG Codons in the First Cistron*. J. Biol. Chem.

280, 27670–27678. https://doi.org/10.1074/jbc.M500255200

Wang, X., Feng, J., Xue, Y., Guan, Z., Zhang, D., Liu, Z., Gong, Z., Wang, Q.,

Huang, J., Tang, C., Zou, T., Yin, P., 2016. Structural basis of

N6-adenosine methylation by the METTL3–METTL14 complex. Nature

534, 575–578. https://doi.org/10.1038/nature18298

Werz, C., Lee, T.V., Lee, P.L., Lackey, M., Bolduc, C., Stein, D.S., Bergmann,

A., 2005. Mis-specified cells die by an active gene-directed process,

and inhibition of this death results in cell fate transformation in

Drosophila. Development 132, 5343–5352.

https://doi.org/10.1242/dev.02150

Woodcock, C.B., Yu, D., Zhang, X., Cheng, X., 2019. Human

HemK2/KMT9/N6AMT1 is an active protein methyltransferase, but does

not act on DNA in vitro, in the presence of Trm112. Cell Discov. 5, 50.

https://doi.org/10.1038/s41421-019-0119-5

Wu, C.C.-C., Peterson, A., Zinshteyn, B., Regot, S., Green, R., 2020.

Ribosome Collisions Trigger General Stress Responses to Regulate

Cell Fate. Cell 182, 404-416.e14.

https://doi.org/10.1016/j.cell.2020.06.006

Xiao, C.-L., Zhu, S., He, M., Chen, D., Zhang, Q., Chen, Y., Yu, G., Liu, J., Xie,

S.-Q., Luo, F., Liang, Z., Wang, D.-P., Bo, X.-C., Gu, X.-F., Wang, K.,

Yan, G.-R., 2018. N6-Methyladenine DNA Modification in the Human

Genome. Mol. Cell 71, 306-318.e7.

https://doi.org/10.1016/j.molcel.2018.06.015

Zhao, Y., Garcia, B.A., 2015. Comprehensive Catalog of Currently

Documented Histone Modifications. Cold Spring Harb. Perspect. Biol. 7,

a025064. https://doi.org/10.1101/cshperspect.a025064

1042

1043

Legends

Figure 1. HemK2 is essential for female germline development and functions in mid-oogenesis.

(A) Schematic representation of *Drosophila* oogenesis. The ovariole consists of the germarium, situated at the anterior tip of the ovariole, followed by a series of sequential stages of egg chamber development. The 16-cell germline cyst originates from the germarium and is encapsulated by a monolayer of somatic follicle cells to generate an egg chamber. One of the 16-cells is identified as the oocyte, which will eventually develop into a mature egg. The developmental checkpoint before the vitellogenesis phases, known as mid-oogenesis, is characterized at stage 7/8. **(B)** Morphological images of control and HemK2 germline knockdown (GLKD) ovaries. *hemk2*-GLKD causes female sterility marked by developmental arrest of egg chambers at stages 7-8 and disruption of vitellogenesis. scale bar is 250 µm. **(C)** qRT-PCR validation of knockdown efficiency in *hemk2*-GLKD ovaries. All values are normalized to *rp49* with *tubulin* as a control. Error bars indicate standard deviation (n=3). **(D)** Egg laying and hatching analysis. Sterility in *hemk2*-GLKD was rescued by wild-type FLAG-tagged HemK2 transgene expression driven by germline driver *NGT40* and *nosGal4-VP16*. Eggs are produced by 3 females every day (n=3). Error bars indicate standard deviation. **(E)** Staining of FLAG-tagged HemK2 (Green) and DAPI (Blue) for DNA in ovaries with each indicated genotype as in (D). Arrows indicate the degenerated egg chambers

1066   in HemK2-GLKD and later stage egg chambers in FLAG-HemK2 transgene

1067   expression. All scale bars are 50 µm. **(F)** Western blots detecting FLAG

1068   showing the germline expression of FLAG-HemK2 wildtype in ovary lysate.

1069   Tubulin was used as the loading control.

1070

Figure 2. HemK2 functions for multiple morphological aspects of

***Drosophila* germline.**

(A) Immunostaining of pH3 (Red) for mitosis and DAPI (Blue) for DNA in

ovaries (left panel). The hyperplasia of the posterior follicle cells in

hemk2-GLKD is denoted with dotted rectangle (upper right panel). All scale

bars are 20 µm. % of ovarioles with single or multi-layered follicle cells is

quantified (right panel, n=100). **(B)** Immunostaining of Fibrillarin (Green) for

nucleolus and DAPI (Blue) for DNA in stage7/8 egg chambers. Inset shows a

magnified image of nurse cell nucleus with dispersed or undispersed

chromosomes. All scale bars are 10 µm. **(C)** qRT-PCR analysis shows

expression level of each chromosome dispersion associated genes. Error bars

indicate standard deviation (n=3). **(D)** Immunostaining of *p53* (Green) for

apoptosis and DAPI (Blue) for DNA in ovaries (left panel). Arrows indicate the

accumulation of *p53* signals in the nuclei of nurse cells. All scale bars are 20

µm. % of ovarioles with or without *p53* signals is quantified (right panel,

n=100).

1087

Figure 3. The methylation of eRF1 was reduced by *hemk2*-GLKD

significantly.

(A) Immunoprecipitation of endogenous eRF1 in ovaries. Western blots

detecting N5-methyl-glutamine (Methyl-Q: bottom panel) showed the reduction

of eRF1 methylation by *hemk2*-GLKD. Heterozygous ovaries were used as the

control. Methyl-Q band intensity normalized to eRF1 was quantified (right

panel). Error bars indicate standard deviation (n=3), *p*-value is shown

(unpaired *t*-test). **(B)** Immunostaining of methyl-eRF1 (Green) and eRF1 (Red)

in *hemk2*-GLKD ovaries. Allows show the cytosol of the germline cells. All

scale bars are 20 μm. **(C)** Immunoprecipitation of FLAG-eRF1 expressed in

germline cells in ovaries. *Matα-tub-Gal4* was used to drive the germline

expression of FLAG-eRF1. Western blots detecting Methyl-Q showed the

significant reduction of FLAG-eRF1 methylation by *hemk2*-GLKD. Asterisks

indicate the nonspecific band. Quantification of Methyl-Q band intensity

normalized to FLAG-eRF1 (right panel) was performed as in (A). Error bars

indicate standard deviation (n=3), *p*-value is shown (unpaired *t*-test). **(D)**

Immunoprecipitation of FLAG-eRF1 Q185A in S2 cells. Western blots

detecting Methyl-Q showing a scanty of methylation in both conditions.

Asterisks indicate the nonspecific band.

1107

1108

Figure 4. eRF1 methylation is required for *hemk2* function.

(A) A schematic diagram showing MTase domain of *Drosophila* HemK2.
Alignment shows that the NPPY motif in the substrate binding pocket is highly
conserved among different species and HemK2 mutants were generated with
Asn116 to Ala (N116A) and Tyr119 to Ala (Y119A), respectively. **(B)** The
crystal structure of the human HemK2-TRMT112 bound with methyl-glutamine
(Methyl-Q) and S-adenosylhomocysteine (SAH) (Gao et al., 2020). Only active
site around Methyl-Q is shown. HemK2 and Methyl-Q are shown in gray and
green, respectively. N122 and Y125 are correspond to N116 and Y119 in
Drosophila HemK2, respectively. Hydrogen bonds around Methyl-Q product
are also shown. **(C)** Immunostaining of FLAG-tagged HemK2 variants (Green)
and DAPI (Blue) for DNA in ovaries with each indicated genotype. The defects
in *hemk2*-GLKD were rescued by transgene expression of the N116A, but not
Y119A HemK2 mutant. All scale bars are 50 μm. **(D)** Egg laying and hatching
analysis of each indicated genotype as in (C). Eggs are produced by 3 females
every day (n=3). Error bars indicate standard deviation. **(E)**
Immunoprecipitation of FLAG-eRF1 in S2 cells. Myc-HemK2 mutants and
FLAG-eRF1 were co-transfected with or without HemK2 knockdown (KD).
Western blots detecting Methyl-Q showing the reduction of FLAG-eRF1
methylation in *hemk2*-KD was rescued by expression of HemK2 wildtype and
N116A, but not Y119A HemK2 mutant. Quantification of Methyl-Q band
intensity normalized to FLAG-eRF1 was performed (bottom panel).

1131

Figure 5. eRF1 methylation ensures proper progression of oogenesis

and was needed for efficient translation.

(A) Morphological images of control and *eRF1*-GLKD ovaries. Scale bar is 250

µm. **(B)** Egg laying and hatching analysis of each indicated genotype. Eggs

are produced by 3 females every day (n=3). Error bars indicate standard

deviation. **(C)** Immunostaining of FLAG-tagged eRF1 WT/Q185A (Green), Vas

(Red) for germline cells and DAPI (Blue) for DNA in ovaries with each

indicated genotype as in (B). The defects in *eRF1*-GLKD were rescued by

transgene expression of wildtype eRF1 but not Q185A eRF1 mutant. All scale

bars are 20 µm. **(D)** Immunostaining of FLAG-tagged eRF1 Q185A (Red) and

DAPI (Blue) for DNA in ovaries. The overexpression of FLAG-eRF1 Q185A in

wildtype phenocopied *hemk2*-GLKD. Arrows indicate the dying egg chambers.

All scale bars are 50 µm. **(E)** Homopropargylglycine (HPG) protein synthesis

assay in ovaried with each indicated genotype (Green in left panels). FLAG

(Red) showing transgene expression was immunostained in parallel. All scale

bars are 20 µm. Quantification of HPG signal intensity in each genotype of the

left panels is shown in the right panel. Representative images were measured

(n=5). Error bars indicate standard deviation.

1150

Figure 6. HemK2 knockdown resulted the mRNA reduction in germline

cells and ribosome stalling.

(A) GO analysis of the downregulated genes in *hemk2*-GLKD ovary.

Overrepresentation of the GO biological process term was tested by

PANTHER (version 18.0). The fold enrichments were plotted and colored

according to their FDR values. **(B)** GO terms of the upregulated genes in

hemk2-GLKD ovary analyzed as in (A). **(C)** Volcano plot of the

germline-enriched genes in *hemk2*-GLKD ovaries. (orange: p-value < 0.05 and

log2FC < -1 or > 1). Cell type-enriched genes are defined according to the

single cell mRNA-seq analysis of *Drosophila* ovaries (Jevitt et al., 2020). **(D)**

Log2-fold changes of cell-type enriched coding genes in *hemk2*-GLKD ovaries

compared with the control (average +/- standard deviation). **(E)** Profiles of

ribosomes in ovarian lysates of the control (blue), *hemk2*-GLKD (red), and

double germline knockdown of *hemk2* and *ski8* (green). The mRNA-bound

monosomes and disomes were separated by the sucrose gradient

sedimentation. The elution profiles were recorded by the absorbance at 254

nm and normalized by the monosome peaks in each sample. The disome

peaks are shown in the inset.

1169

Figure 7. Defects during oogenesis in *hemk2*-GLKD were rescued by

blocking of the no-go decay (NGD) pathway.

(A) Immunostaining of DAPI (Blue) for DNA in ovaries with each indicated

genotype. The defects in *hemk2*-GLKD were rescued by double knockdown of

ski8 or *pelo*, but not *CG7139* or *upf3*. All scale bars are 50μm. **(B)** Egg laying

and hatching analysis of each indicated genotype as in (A). Eggs are produced

by 3 females every day (n=3). Error bars indicate standard deviation. **(C)** HPG

protein synthesis assay in ovaried with each indicated genotype as in (A). All

scale bars are 20 μm. **(D)** % of ovarioles labeled with HPG positive is

quantified (left panel, n=100). Quantification of HPG signal intensity in each

genotype in (C) is shown in the right panel. Representative images were

measured (n=5). Error bars indicate standard deviation.

1182

Supplementary Legends

Figure S1. HemK2 plays a predominant role not only in germline but also in somatic cells.

(A) A schematic drawing of *Drosophila* HemK2/Snapin locus. The primers of Snapin dsRNA and HemK2 qPCR are illustrated (bottom). **(B)** Egg laying and hatching analysis. Sterility in HemK2 somatic knockdown (STKD) was rescued by wild-type FLAG-tagged HemK2 transgene expression driven by somatic driver *traffic jam-Gal4*. Eggs are produced by 3 females every day (n=3). Error bars indicate standard deviation. **(C)** Immunostaining of FLAG-tagged HemK2 (Green) and DAPI (Blue) for DNA in ovaries with each indicated genotype as in (B). Arrows indicate the degenerated egg chambers in HemK2-STKD and later stage egg chambers in FLAG-HemK2 transgene expression. All scale bars are 50 μm. **(D)** Egg laying and hatching analysis. Sterility in *hemk2/snapin*-GLKD was not rescued by Snapin transgene expression driven by germline driver *NGT40 and nosGal4-VP16*. Eggs are produced by 3 females every day (n=3). Error bars indicate standard deviation. **(E)** Immunostaining of FLAG-tagged Snapin (Green) and DAPI (Blue) for DNA in ovaries with each indicated genotype as in (D). All scale bars are 50 μm. **(F)** qRT-PCR validation of knockdown efficiency in *hemk2*-GLKD testes. All values are normalized to *rp49* with *tubulin* as a control. Error bars indicate standard deviation (n=3). **(G)** Egg laying and hatching analysis in testes. Eggs are produced by 3 *yw* females every day mated with the indicated males (n=3). Error bars indicate

1205   standard deviation.

1206

Figure S2. The defects in *hemk2*-GLKD were rescued by the repression

of cell death partially.

(A) Immunostaining of FLAG-tagged otu104 (Red), Fibrillarin (Green) for

nucleolus and DAPI (Blue) for DNA in ovaries with each indicated genotype

(left panel). The chromosomal dispersion defect in *hemk2*-GLKD was partially

rescued by transgene expression of *otu*104 driven by germline driver *NGT40*

and nosGal4-VP16. All scale bars are 20 μm. % of ovarioles with dispersed or

undispersed chromosomes is quantified (right panel, n=100, stage 6 and older

egg chambers). **(B)** Immunostaining of Fibrillarin (Green) for nucleolus and

DAPI (Blue) for DNA in egg chambers with each indicated genotype. The

stages of egg chambers are determined and labeled (bottom). **(C)** Egg laying

and hatching analysis of each indicated genotype as in (B). Eggs are produced

by 3 females every day (n=3). Error bars indicate standard deviation.

1220

Figure S3. DNA 6mA and histone H4K12 remained unaffected by

***hemk2*-GLKD.**

(A) Immunostaining of Vas (Red) for germline cells, DNA N6-adenine (6mA)

(Green) and DAPI (Blue) for DNA in ovaries. DNA 6mA was unaffected by

hemk2-GLKD. All scale bars are 20 μm. **(B)** Immunoblotting of histone H4 and

H4K12met in ovary lysate. Tubulin was used as the loading control.

1227

Figure S4. eRF1 methylation is uniquely catalyzed by HemK2.

(A) Western blots detecting FLAG showing the germline expression of

FLAG-HemK2 N116A and Y119A mutants in ovary lysate. Tubulin was used

as the loading control. **(B)** Immunoprecipitation of FLAG-eRF1 in S2 cells.

Myc-HemK1/HemK2 and FLAG-eRF1 were co-transfected with or without

hemk2 KD. Western blots detecting Methyl-Q showing the reduction of

FLAG-eRF1 methylation in *hemk2*-KD was rescued by expression of HemK2

but not HemK1. Quantification of Methyl-Q band intensity normalized to

FLAG-eRF1was performed (bottom panel).

1237

Figure S5. The overexpression of FLAG-eRF1 Q185A in wildtype

phenocopied *hemk2*-GLKD.

(A) Western blots detecting eRF1 showing the expression of endogenous

eRF1 and FLAG-eRF1 Q185A in ovary lysate. *Matα-tub-Gal4* was used to

drive the germline expression of FLAG-eRF1 Q185A. Tubulin was used as the

loading control. **(B)** Egg laying and hatching analysis of each indicated

genotype. Eggs are produced by 3 females every day (n=3). Error bars

indicate standard deviation. **(C)** Immunostaining of p53 (Green) for apoptosis

and DAPI (Blue) for DNA in ovaries (left panel). The overexpression of

FLAG-eRF1 Q185A in wildtype phenocopied *hemk2*-GLKD in cell death.

Arrows indicate the accumulation of *p53* signals in the nuclei of nurse cells. All

scale bars are 20 μm. % of ovarioles with or without p53 signals is quantified

(right panel, n=100). **(D)** HPG protein synthesis assay in ovaried (left panels).

FLAG (Red) showing transgene expression was immunostained in parallel. All

scale bars are 20 μm. Quantification of HPG signal intensity in each genotype

of the left panels is shown in the right panel. Representative images were

measured (n=5), *p*-values are shown (unpaired *t*-test). Error bars indicate

standard deviation.

1256

Figure S6. *hemk2* knockdown resulted the mRNA reduction in germline

cells, but not in somatic cells.

(A-E) Volcano plot of the cell type-enriched genes in *hemk2*-GLKD ovaries.

(orange: p-value < 0.05 and log2FC < -1 or > 1). Cell type-enriched genes are

defined according to the single cell mRNA-seq analysis of *Drosophila* ovaries

(Jevitt et al., 2020). Somatic germarium (A), Polar cells (B), Stalk cells (C),

pre-follicle cells (D), and mitotic follicle cells (E). **(F, G)** Log2-fold changes of

cell-type enriched coding genes in *hemk2-ski8*-DGKD (F) and in

hemk2-pelo-DGKD (G) ovaries compared with the control (average +/-

standard deviation).

1267

1268 **Figure S7. Validations of the knockdown efficiency for each mRNA decay**

1269 **factor.** (A) qRT-PCR was performed to validate the knockdown of *hemk2*, *ski8*,

1270 *pelo*, *cg7139*, and *upf3*. Error bars indicate standard deviation. (B) *CG7139* is

1271 a *Drosophila* homolog of *nonu-1*/*cue2*. Smr nuclease and AAA_33 kinase

1272 domains showed the 30% and 20% identity, respectively.

Figure 1

A *Drosophila* oogenesis

previtellogenic stages   vitellogenesis

- nurse cell
- follicle cell
- oocyte
- nucleus

Germarium   St3   St5   St6   St7   St9   St12   St14

B   Germline knockdown (GLKD)

Control   *hemk2*-GLKD

250 µm

C

Relative mRNA level

Tub   hemk2

Control   *hemk2*-GLKD

D

# of eggs / day

Control, *hemk2*-GLKD, *hemk2*-GLKD; FLAG-HemK2, FLAG-HemK2 OE

% of egg hatching

N/A

Control, *hemk2*-GLKD, *hemk2*-GLKD; FLAG-HemK2, FLAG-HemK2 OE

F

| *hemk2*-GLKD: | - | + | + | - |
| FLAG-HemK2: | - | - | WT | WT |

FLAG

Tub

E   Control   *hemk2*-GLKD   *hemk2*-GLKD; FLAG-HemK2   FLAG-HemK2 OE

FLAG

DAPI FLAG

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7