Disrupting Homologous Recombination or Single-Strand Annealing Significantly Hinders CRISPR-Cas12a-Assisted Nonhomologous End-Joining Gene Editing Efficiency in *Mycobacterium abscessus*

Sanshan Zeng,[a,b,c,d] Yanan Ju,[a,b,c,e] Md Shah Alam,[a,b,c,d] Ziwen Lu,[a,b,c,d] H.M. Adnan Hameed,[a,b,c] Lijie Li,[a,b,c,e] Xirong Tian,[a,b,c,d] Cuiting Fang,[a,b,c,d] Xiange Fang,[a,b,c,d] Jie Ding,[a,b,c,f] Xinyue Wang,[g,h] Jinxing Hu,[h] Shuai Wang,[a,b,c,h]* Tianyu Zhang [a,b,c,d,e,f,g] *

[a] State Key Laboratory of Respiratory Disease, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou, China

[b] Guangdong-Hong Kong-Macao Joint Laboratory of Respiratory Infectious Diseases, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou, China

[c] China-New Zealand Joint Laboratory on Biomedicine and Health, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou, China

[d] University of Chinese Academy of Sciences, Beijing, China

[e] School of Basic Medical Sciences, Division of Life Science and Medicine, University of Science and Technology of China, Hefei, China

[f] Institute of Physical Science and Information Technology, Anhui University, Hefei, China

[g] Guangzhou National Laboratory, Guangzhou, China

[h] State Key Laboratory of Respiratory Disease, Guangzhou Chest Hospital, Guangzhou, China

* Corresponding authors: Shuai Wang, E.mail: wshuai1@163.com, Tel: +86-20-32015266; Tianyu Zhang, E.mail: zhang_tianyu@gibh.ac.cn, Tel: +86-20-32015270.

Running title: Influence of HR and SSA on *M. abscessus* Gene Editing Efficiency

ABSTRACT *Mycobacterium abscessus*, a fast-growing, non-tuberculous mycobacterium resistant to most antimicrobial drugs, poses a significant public health challenge because of its ability to cause many types of serious infections in humans. While genetic manipulation tools for *M. abscessus* are still being developed, the clustered regularly interspaced short palindromic repeats (CRISPR)-CRISPR-associated protein (Cas) systems have shown promise in generating highly specific double-strand breaks (DSBs) in its genome. These DSBs can be repaired by the error-prone nonhomologous end joining (NHEJ) mechanism, facilitating targeted gene editing. Here, our study marks a pioneering application of the CRISPR-NHEJ strategy in *M. abscessus*. Moreover, our research uncovered an unexpected finding: contrary to previous observations in *Mycobacterium tuberculosis* and other species, the inhibition of RecA or disruption of key genes in the homologous recombination or single-strand annealing pathways resulted in a significant decrease in NHEJ repair efficiency in *M. abscessus*. This discovery challenges the established perspectives and offers new insights into the interaction among the three DSB repair pathways in Mycobacterium species.

IMPORTANCE There are still very few genetic manipulation tools available for *Mycobacterium abscessus*.. Here we report the successful application of CRISPR-Cas12a-assisted nonhomologous end joining (NHEJ) in efficient gene editing in *M. abscessus*. Contrary to previous research suggesting that homologous recombination (HR) inhibition may enhance such editing efficiency in other Mycobacterium species, our results showed that disruption of either HR or single-strand annealing (SSA) DNA double-strand breaks (DSBs) repair pathways not only failed to enhance but also significantly reduced the gene editing efficiency in *M. abscessus*. This suggests that NHEJ repair in *M. abscessus* may require components from both HR and SSA pathways, highlighting a complex interaction among the DSB repair pathways in *M. abscessus.*

KEYWORDS *Mycobacterium abscessus*, CRISPR-Cas12a, DSB repair, NHEJ

The prevalence of lung infections caused by non-tuberculous mycobacteria (NTM) has been rising annually, posing an escalating threat to public health. Among these NTM, *Mycobacterium abscessus* is one of the primary causative agents, second only to *Mycobacterium avium* complex in both the United States and the humid regions of southern coastal China (1-3). Besides contagious pulmonary infections, *M. abscessus* can infect the skin, soft tissues, central nervous system, eyes, and various other parts of the human body (4). The intrinsic resistance of *M. abscessus* to a broad spectrum of antibiotics not only complicates the treatment, but often requires the use of multidrug regimens and prolonged therapies (5). The threat posed by *M. abscessus* necessitates the urgent development of genetic tools for drug target identification, resistance mechanism elucidation, and vaccine research.

Various attempts have been made at genetic manipulation in Mycobacterium species, such as the expression of *gp60* and *gp61* from the mycobacteriophage Che9c to enhance recombination in *Mycobacterium smegmatis, Mycobacterium tuberculosis* (6), and *M. abscessus* (7). CRISPR-assisted recombineering has further enabled the effectual construction of marker-free recombinants in Mycobacterium spp. After successful implication in *M. smegmatis* (8), this technique has recently been applied successfully for the first time by us in *M. abscessus* (9). Additionally, nonhomologous end joining (NHEJ), another DNA double-strand break (DSB) repair mechanism, presents the potential for genetic manipulation together with CRISPR, owing to its template-independent and error-prone repair characteristics (10, 11). The combined use of CRISPR and NHEJ for genetic manipulation has been proven effective in *M. smegmatis* and *M. tuberculosis*, with inhibition of the host's homologous recombination (HR) system by overexpressing RecX or disrupting *recA* gene (12). RecX, known to antagonize RecA, impedes the HR process by negatively regulating the array of biochemical reactions facilitated by RecA (13). However, attempts to apply these strategies to *M. abscessus* have not been reported so far. Here, we report that gene editing by the CRISPR-NHEJ strategy was successful and efficient in *M. abscessus*. Interestingly, on the contrary to previous findings, gene editing efficiency significantly decreased in *M. abscessus* once RecX was overexpressed or key genes in HR or single-strand annealing (SSA) were knocked-out.

We tried to apply the CRISPR-NHEJ strategy to *M. abscessus,* and used *M.*
smegmatis as a control. The pNHEJ-Cpf1 plasmid containing *Mycobacterium*
marinum NHEJ elements (*ligD*, *ku* and *nrgA*) as well as *cpf1* (*cas12a*) or the
pNHEJX-Cpf1 plasmid additionally expressing *recX^{Msm}* (*recX* gene of *M. smegmatis*,
Fig. S1) were transduced into *M. abscessus* GZ002 (14, 15) and *M. smegmatis* MC^2
155 by electroporation, respectively. Transformants were selected on
kanamycin-containing 7H10 plates. Strains carrying the corresponding plasmids were
cultured in 3 mL of 7H9 medium supplemented with 10% OADC, 0.2% glycerol,
0.05% Tween 80, and 100 μg/mL kanamycin for *M. abscessus* or 50 μg/mL for *M.*
smegmatis. Then 1 mL of the starter culture was inoculated into a 125 mL shake flask
containing 40 mL of 7H9 complete broth, and incubated at 37°C until OD$_{600}$=0.8-1.0.
After that, the bacteria were harvested for preparation of competent cells as
previously described (16). About 500 ng of the plasmid expressing crRNA (primers
for constructing crRNA are listed in Table S1) was mixed with 100 μL of competent
cells, transferred to a 2 mm electroporation cuvette, and incubated on ice for 10 min.
Then the cuvette was placed into an electroporation chamber for electroporation at a
voltage of 2,500 V, a capacitance of 25 μF, and a resistance of 1,000 Ω. After
electroporation, 2 mL of 7H9 medium was added to the electroporation cuvette to
resuspend the bacteria, which was then transferred to a 50 mL tube and incubated at
30°C for 4 to 5 hours. Then, cultures were plated separately on 7H11 agar
supplemented with the appropriate antibiotics and 200 ng/mL anhydrotetracycline
(ATc) for *M. abscessus* or 50 ng/mL for *M. smegmatis*. The editing efficiency was
determined by the ratio of colonies with altered target sequences to the total number
of randomly picked colonies. For each assessment, at least 20 colonies were randomly
selected for PCR (primers are shown in Table S1) and sequencing analysis. The
colonies that either lacked the target bands in PCR or showed sequencing results
deviating from the wild-type sequence were identified as colonies that had undergone
gene editing.

Here, we observed a surprising phenomenon. In the *M. abscessus* transformed
with pNHEJ-Cpf1 group, the gene editing efficiency of the randomly selected
nonessential gene *MAB_3513c* was 63.35 ± 4.738 % (Fig. 1A). However, the gene
editing efficiency in the *M. abscessus* transformed with pNHEJX-Cpf1 group was
only 18.35 ± 2.333 %. The editing efficiency in the pNHEJ-Cpf1 group was

approximately threefold higher than that in the strain with additional $recX^{Msm}$ expression (Fig. 1A, $P = 0.0068$), which is completely opposite to a previous research in *M. smegmatis* showing that it is necessary either to introduce additional $recX^{Msm}$ gene or to knock-out the *recA* gene to suppress HR to edit gene efficiently when using the CRISPR-NHEJ strategy (12). Consequently, researchers hypothesized that in *M. smegmatis*, HR and NHEJ are competing pathways (12). To verify this hypothesis, we repeated their study by editing the *MSMEG_1946* gene in *M. smegmatis* to compare the editing efficiency with or without expressing additional $recX^{Msm}$ gene. Although the gene editing efficiency remained high at $85 \pm 7.071\%$ in *M. smegmatis* containing pNHEJX-Cpf1 with overexpressed $recX^{Msm}$, in the group containing pNHEJ-Cpf1 without additional $recX^{Msm}$, the gene editing efficiency was still $78.35 \pm 11.31\%$. We did not observe an obvious decrease in editing efficiency (Fig. 1A, $P = 0.5650$). These results were clearly different from the earlier report (12).

Unlike HR, NHEJ repair was characterized by deletions of base pairs of random lengths (Fig. S2). To further validate the broad applicability of this strategy, we tried to edit other *M. abscessus* genes only using pNHEJ-Cpf1 without additional expression of $recX^{Msm}$. To date, we have successfully knocked-out over 30 genes (partially listed in Table S2) in *M. abscessus* using this method. A limitation of this study is that we did not edit more genes to compare the efficiencies with and without additional expression of the $recX^{Msm}$ widely. However, the overall efficiency range of editing different genes by different researchers using this method in *M. abscessus* was 30% - 90% which was generally sufficient for a common gene editing. This strategy has significantly improved our efficiency in acquiring mutants.

The surprising results spurred our further investigation. On the one hand, we tried the gene editing in the HR-disrupted *M. abscessus* strain as reported in *M. smegmatis* (11). On the other hand, though it was reported that the inhibition of SSA can enhance HR (17), but no study has been reported on whether disruption of SSA would affect the NHEJ-dependent gene editing in Mycobacteria, so, we tried to investigate weither blocking SSA or disrupting HR and SSA both would impact the NHEJ-dependent gene editing efficiency in *M. abscessus*.

We constructed selection-marker free, in-frame deletion knock-out strains of *recA*, *recC* or *recO* gene in *M. abscessus* (unpublished) using the CRISPR-assisted

recombineering method as described in our previous study (9). The RecA plays a key

role in HR, while the RecBCD complex composed of RecC plays an important role in

SSA independently of the RecA-dominated HR. In addition, it was shown that

deletion of the *recO* gene would result in defective repair of both HR and SSA (18-20).

Next, in these HR and SSA repair-deficient strains, we repeated the gene editing

experiments performed in the wild-type strain as described above to observe the

potential impact of HR (*recA*) or SSA (*recC*) or both (*recO*) repair defects on the gene

editing efficiency.

Our results showed that the gene editing efficiencies were 3.33 ± 2.877 %, 17.5 ±

3.536 % and 7.5 ± 3.536 % in the *recA*, *recC* or *recO* knock-out strains, respectively

(Fig. 1B). As expected, the gene editing efficiency in the *recA* gene knock-out strain

was much lower than that in the wild-type strain (P = 0.0004) and even significantly

lower than that in the wild-type strain overexpressing *recX^{Msm}* gene (P = 0.009, Fig.

1A and Fig. 1B). Similarly, the gene editing efficiencies in the *recC* or *recO* gene

knock-out strains were significantly lower than that in the wild-type strain (P =

0.0082 and P = 0.0056, respectively, Fig. 1B). All these results supported that

disruption of the HR or SSA repairing pathway would decrease the

CRISPR-Cas12a-assisted NHEJ gene editing efficiency.

Interestingly, the ranking of editing efficiencies in the knock-out strains was *recA*

(HR) ≤ *recO* (HR&SSA) ≤ *recC* (SSA). The significant difference in editing

efficiencies between two of the three knock-out strains was only observed between

recA and *recC* (P = 0.0156, Fig. 1B). This indicated that the gene editing efficiency

was influenced differently by the different genes in the HR or SSA DSB repair

pathways, and suggested that those in the HR pathway or gene playing a key role in

the two pathways might have a stronger influence than that only affects SSA. Even

though *recO* may influence both HR and SSA, it only affects a branch pathway of HR

unlike RecA which plays a key role in HR (21). However, the influence of each gene

in either HR or SSA pathways on the gene editing efficiency needs further

investigation.

Previous studies largely indicated that the three DSB repair mechanisms might

be competitive with each other for gene editing efficiency (12, 17), which contrasts

sharply with our findings that they cooperate for the completion of gene editing. Here

we report for the first time the successful application of the NHEJ-CRISPR strategy for the efficient gene editing but without inhibiting the HR pathway in *M. abscessus*, and also reveal a potential synergistic interaction between HR or SSA and NHEJ within this bacterium. For *M. smegmatis*, a study showing that disruption of HR would not enhance the DSB repair efficiency through the NHEJ pathway supports what we found here that disruption of HR did not affect the CRISPR-Cas12a-assisted NHEJ gene editing efficiency (Fig. 1A) (22). Our novel finding offers a new perspective for further exploration of the interrelationships between DSB repair mechanisms in mycobacteria and paves the way for future enhancement of specific repair pathway efficiencies by manipulating these relationships. More refined mechanism research of the relationship among HR, SSA, and NHEJ and their impact on the CRISPR-Cas12a-assisted NHEJ gene editing efficiency requires further investigation.

ACKNOWLEDGMENTS

This study was supported by the National Key R&D Program of China (2021YFA1300900), the National Natural Science Foundation of China (21920102003), the State Key Lab of Respiratory Disease, Guangzhou Institute of Respiratory Diseases, First Affiliated Hospital of Guangzhou Medical University (SKLRD-Z-202301, SKLRD-Z-202412), and Guangzhou Science and Technology (2024A04J4273), the Chinese Academy of Sciences, President's International Fellowship Initiative (2023VBC0015). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. All authors read and approved the final version of the manuscript.

We also extend our sincere appreciation to Professor Yichen Sun from the Chinese Academy of Medical Sciences and Peking Union Medical College for generously providing the pJV53-Cpf1 and pCR-ZEO plasmids for our study.

REFERENCES

1. Victoria L, Gupta A, Gómez JL, Robledo J. 2021. *Mycobacterium abscessus* complex: A Review of Recent Developments in an Emerging Pathogen. Front Cell Infect Microbiol 11:659997.

2. Floto RA, Olivier KN, Saiman L, Daley CL, Herrmann J-L, Nick JA, Noone PG, Bilton D, Corris P, Gibson RL, Hempstead SE, Koetz K, Sabadosa KA, Sermet-Gaudelus I, Smyth AR, Van Ingen J, Wallace RJ, Winthrop KL, Marshall BC, Haworth CS. 2016. US Cystic Fibrosis Foundation and European Cystic Fibrosis Society consensus recommendations for the management of non-tuberculous mycobacteria in individuals with cystic fibrosis. Thorax 71:88-90.

3. Liu C-F, Song Y-M, He W-C, Liu D-X, He P, Bao J-J, Wang X-Y, Li Y-M, Zhao Y-L. 2021. Nontuberculous mycobacteria in China: incidence and antimicrobial resistance spectrum from a nationwide survey. Infect Dis Poverty 10:59.

4. Griffith DE, Aksamit T, Brown-Elliott BA, Catanzaro A, Daley C, Gordin F, Holland SM, Horsburgh R, Huitt G, Iademarco MF, Iseman M, Olivier K, Ruoss S, von Reyn CF, Wallace RJ, Winthrop K. 2007. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. Am J Respir Crit Care Med 175:367-416.

5. Nessar R, Cambau E, Reyrat JM, Murray A, Gicquel B. 2012. *Mycobacterium abscessus*: a new antibiotic nightmare. J Antimicrob Chemother 67:810-818.

6. van Kessel JC, Hatfull GF. 2007. Recombineering in *Mycobacterium tuberculosis*. Nat Methods 4:147-152.

7. Wang S, Cai X, Yu W, Zeng S, Zhang J, Guo L, Gao Y, Lu Z, Hameed HMA, Fang C, Tian X, Yusuf B, Chhotaray C, Alam MDS, Zhang B, Ge H, Maslov DA, Cook GM, Peng J, Lin Y, Zhong N, Zhang G, Zhang T. 2022. Arabinosyltransferase C Mediates Multiple Drugs Intrinsic Resistance by Altering Cell Envelope Permeability in *Mycobacterium abscessus*. Microbiol Spectr 10:e0276321.

8. Yan M-Y, Yan H-Q, Ren G-X, Zhao J-P, Guo X-P, Sun Y-C. 2017. CRISPR-Cas12a-Assisted Recombineering in Bacteria. Appl Environ Microbiol 83:e00947-17.

9. Wang S, Zhang J, Hameed HMA, Ding J, Guan P, Fang X, Peng J, Su B, Ma S, Tan Y, M Cook G, Zhang G, Lin Y, Zhong N, Hu J, Liu J, Zhang T. 2023. Amino acid 17 in QRDR of Gyrase A plays a key role in fluoroquinolones susceptibility in mycobacteria. Microbiol Spectr 11:e0280923.

10. Singh A. 2017. Guardians of the mycobacterial genome: A review on DNA repair systems in *Mycobacterium tuberculosis*. Microbiology (Reading) 163:1740-1758.

11. Glickman MS. 2014. Double-Strand DNA Break Repair in Mycobacteria. Microbiol Spectr 2.

12. Yan M-Y, Li S-S, Ding X-Y, Guo X-P, Jin Q, Sun Y-C. 2020. A CRISPR-Assisted Nonhomologous End-Joining Strategy for Efficient Genome Editing in *Mycobacterium tuberculosis*. mBio 11:e0280923.

13. Prasad D, Muniyappa K. 2020. The extended N-terminus of Mycobacterium smegmatis RecX potentiates its ability to antagonize RecA functions. Biochim Biophys Acta Proteins Proteom 1868:140468.

257   14.   Guo J, Wang C, Han Y, Liu Z, Wu T, Liu Y, Liu Y, Tan Y, Cai X, Cao Y, Wang B, Zhang B,
258   Liu C, Tan S, Zhang T. 2016. Identification of Lysine Acetylation in *Mycobacterium abscessus*
259   Using LC-MS/MS after Immunoprecipitation. J Proteome Res 15:2567-78.

260   15.   Chhotaray C, Wang S, Tan Y, Ali A, Shehroz M, Fang C, Liu Y, Lu Z, Cai X, Hameed HMA,
261   Islam MM, Surineni G, Tan S, Liu J, Zhang T. 2020. Comparative Analysis of Whole-Genome
262   and Methylome Profiles of a Smooth and a Rough *Mycobacterium abscessus* Clinical Strain.
263   G3 (Bethesda) 10:13-22.

264   16.   Goude R, Roberts DM, Parish T. 2015. Electroporation of mycobacteria. Methods Mol Biol
265   1285:117-130.

266   17.   Tufariello JM, Malek AA, Vilchèze C, Cole LE, Ratner HK, González PA, Jain P, Hatfull GF,
267   Larsen MH, Jacobs WR. 2014. Enhanced Specialized Transduction Using Recombineering in
268   *Mycobacterium tuberculosis*. mBio 5:e01179-14.

269   18.   Cox MM. 1999. Recombinational DNA repair in bacteria and the RecA protein. Prog Nucleic
270   Acid Res Mol Biol 63:311-366.

271   19.   Gorna E, Alina, Bowater P, Richard, Dziadek J. 2010. DNA repair systems and the
272   pathogenesis of *Mycobacterium tuberculosis*: varying activities at different stages of infection.
273   Clin Sci 119:187-202.

274   20.   Gupta R, Barkan D, Redelman-Sidi G, Shuman S, Glickman MS. 2011. Mycobacteria exploit
275   three genetically distinct DNA double-strand break repair pathways. Mol Microbiol
276   79:316-330.

277   21.   Gupta R, Ryzhikov M, Koroleva O, Unciuleac M, Shuman S, Korolev S, Glickman MS. 2013.
278   A dual role for mycobacterial RecO in RecA-dependent homologous recombination and
279   RecA-independent single-strand annealing. Nucleic Acids Res 41:2284-95.

280   22.   Gupta R, Barkan D, Redelman-Sidi G, Shuman S, Glickman MS. 2011. Mycobacteria exploit
281   three genetically distinct DNA double-strand break repair pathways. Mol Microbiol
282   79:316-30.

FIG 1 CRISPR-Cas12a-NHEJ-assisted gene editing in *M. abscessus* and *M. smegmatis*. (A) Overexpression of the *recX^{Msm}* gene has different effects on the editing efficiency of *M. smegmatis* and *M. abscessus*. The plasmid expressing crRNA targeting *MSMEG_1946* or *MAB_3513c* was introduced into *M. smegmatis* or *M. abscessus* strains harboring pNHEJ-Cpf1 or pNHEJX-Cpf1, respectively. pN: pNHEJ-Cpf1, pNX: pNHEJX-Cpf1. (B) CRISPR-Cas12a-NHEJ gene editing efficiency in *M. abscessus* and its derivatives. Knocking out the *recA* (disruption of HR), *recC* (disruption of SSA), *recO* (disruption of HR and SSA) genes all resulted in a significant decrease in gene editing efficiency in *M. abscessus*. Editing efficiency was calculated by measuring the proportion of colonies with mutations at the target site among the total randomly selected clones and survival rate was defined as the ratio of the number of CFUs obtained from plates with ATc to the number of CFUs obtained from plates without ATc. ns, not significant. *, $p < 0.05$. **, $p < 0.01$. ***, $p < 0.001$.