Plasmodium RON11 triggers biogenesis of the merozoite rhoptry pair and is essential for erythrocyte invasion

David Anaguano[1,2], Opeoluwa Adewale-Fasoro[3,4], Grace S. Vick[2,5], Sean Yanik[3,4], James Blauwkamp[6], Manuel A. Fierro[1,2,#], Sabrina Absalon[6], Prakash Srinivasan[3,4] and Vasant Muralidharan[1,2,*]

[1]Department of Cellular Biology, [2]Center for Tropical and Emerging Global Diseases, University of Georgia, Athens, GA.

[3]Department of Molecular Microbiology and Immunology, and Johns Hopkins Malaria Institute, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD

[4]The Johns Hopkins Malaria Research Institute, Baltimore, MD, 21205, USA.

[5]Department of Infectious Diseases, College of Veterinary Medicine, University of Georgia, Athens, GA.

[6]Department of Pharmacology and Toxicology, Indiana University School of Medicine, Indianapolis IN.

[#]Current address: Department of Biomedical Sciences, College of Veterinary Medicine, Iowa State University, Ames IA

*Corresponding author: vasant@uga.edu

ABSTRACT

Malaria is a global and deadly human disease caused by the apicomplexan parasites of the genus *Plasmodium*. Parasite proliferation within human red blood cells (RBC) is associated with the clinical manifestations of the disease. This asexual expansion within human RBCs, begins with the invasion of RBCs by *P. falciparum*, which is mediated by the secretion of effectors from two specialized club-shaped secretory organelles in merozoite-stage parasites known as rhoptries. We investigated the function of the Rhoptry Neck Protein 11 (RON11), which contains seven transmembrane domains and calcium-binding EF-hand domains. We generated conditional mutants of the *P. falciparum* RON11. Knockdown of RON11 inhibits parasite growth by preventing merozoite invasion. The loss of RON11 did not lead to any defects in processing of rhoptry proteins but instead led to a decrease in the amount of rhoptry proteins. We utilized ultrastructure expansion microscopy (U-ExM) to determine the effect of RON11 knockdown on rhoptry biogenesis. Surprisingly, in the absence of RON11, fully developed merozoites had only one rhoptry each. The single rhoptry in RON11 deficient merozoites were morphologically typical with a bulb and a neck oriented into the apical polar ring. Moreover, rhoptry proteins are trafficked accurately to the single rhoptry in RON11 deficient parasites. These data show that in the absence of RON11, the first rhoptry is generated during schizogony but upon the start of cytokinesis, the second rhoptry never forms. Interestingly, these single-rhoptry merozoites were able to attach to host RBCs but are unable to invade RBCs. Instead, RON11 deficient merozoites continue to engage with RBC for prolonged periods eventually resulting in echinocytosis, a result of secreting the contents from the single rhoptry into the RBC. Together, our data show that RON11 triggers the *de novo* biogenesis of the second rhoptry and functions in RBC invasion.

INTRODUCTION

Malaria remains a devastating disease that affected approximately 249 million people worldwide in 2022, with the African region accounting for over 95% of the cases (World Health Organization 2023). The main causative agent of malaria is the unicellular protozoan parasite *Plasmodium falciparum*, which belongs to the apicomplexan phylum, along with other medically important organisms such as *Toxoplasma gondii* and *Cryptosporidium parvum*. Clinical symptoms of malaria in humans are associated with the continuous cycle of invasion of red blood cells (RBCs), development and replication within RBCs, and egress from RBCs to release new merozoites that perpetuate the cycle. Invasion is a process that has been targeted by multiple vaccination and antibody treatment strategies (Beeson et al. 2016; González-Sanz, Berzosa, and Norman 2023), which underscores the importance of understanding this process for the development of clinical interventions.

Invasion is a rapid and complex process involving multiple steps. Newly egressed merozoites initiate an early attachment, leading to RBC deformation and merozoite reorientation (Groomes, Kanjee, and Duraisingh 2022; Hart et al. 2023; Weiss et al. 2015). Once its apical end is reoriented towards the RBC, the merozoite appears to form a pore in the RBC membrane, secretes proteins from rhoptries into the RBC, which together with micronemal proteins on the merozoite membrane, form an irreversible tight junction at the interface merozoite-RBC (Cova, Lamarque, and Lebrun 2022; Geoghegan et al. 2021; Riglar et al. 2011; Vulliez-Le Normand et al. 2012; Weiss et al. 2015). Subsequently, the merozoite utilizes its actinomyosin motor to push through the tight junction and internalize within the newly formed parasitophorous vacuole (PV) (Blake, Haase, and Baum 2020; Frénal et al. 2017; Geoghegan et al. 2021; Perrin et al. 2018; Yahata et al. 2021). Within the PV, merozoites develop into the ring stage, progressing into the metabolically active forms, trophozoites, and finally into the replicative form, schizonts, which make 16-32 daughter merozoites. The parasite then initiates a cascade of events leading to the rupture of the RBC and PV membranes, resulting in the egress of merozoites (Dvorin and Goldberg 2022).

Rhoptries are specialized club-shaped secretory organelles in apicomplexans known to
be essential for parasite proliferation. Even though rhoptries have conserved roles
among apicomplexans, their numbers vary between species, for instance
Cryptosporidium spp. have a single rhoptry, *Plasmodium spp.* have two in the asexual
stages, and *Toxoplasma spp.* have eight to twelve rhoptries (Cova, Lamarque, and
Lebrun 2022; Mageswaran et al. 2021; Martinez et al. 2022; Segev-Zarko et al. 2022).
In *P. falciparum*, rhoptries are located at the apical end of the merozoite and secrete
proteins that facilitate different steps during invasion. Rhoptries are internally
compartmentalized into two regions: the neck and bulb, storing specific groups of
proteins that are secreted at each step of invasion (Counihan et al. 2013; Cova,
Lamarque, and Lebrun 2022; Liffner et al. 2021). Rhoptry neck proteins are assumed to
be released first, primarily mediating attachment to RBCs and the formation of the tight
junction through secretion of proteins from the RON complex (Srinivasan et al. 2011;
Weiss et al. 2015). In contrast, rhoptry bulb proteins are secreted after tight junction
formation and are associated with proper development within the PV, such as RAP1
and RAP2, which form the RAP complex (Ghosh et al. 2017; Riglar et al. 2011). During
invasion, rhoptries fuse their necks and bulbs to form a single rhoptry (Hanssen et al.
2013). After invasion is completed, this single rhoptry is assumed to be disassembled,
as rhoptries are absent during early ring stages. During schizogony, rhoptries are then
formed *de novo* by the fusion of Golgi-derived vesicles (Bannister et al. 2000; Krai,
Dalal, and Klemba 2014).

Rhoptry biogenesis begins early during schizogony, where the rhoptries remain
associated with the cytoplasmic extensions of the outer centriolar plaque (CP) branches
in a 1:1 ratio with each new rhoptry forming *de novo* through each remaining mitotic
event, until the last event where this ratio is broken and each outer CP branch is
observed with 2 rhoptries (Bannister et al. 2000; Liffner, Cepeda Diaz, Blauwkamp,
Anaguano, Frolich, et al. 2023; Rudlaff et al. 2020). The final rhoptry pairs undergo *de*
novo biogenesis instead of dividing from a common rhoptry precursor (Liffner, Cepeda
Diaz, Blauwkamp, Anaguano, Frolich, et al. 2023), as was previously assumed.
Rhoptries maintain a bulb shape throughout most of their biogenesis process, adopting
their final shape only after merozoite segmentation, with the formation of the neck in

each rhoptry (Bannister et al. 2000; Liffner, Cepeda Diaz, Blauwkamp, Anaguano,

Frolich, et al. 2023). However, the molecular factors regulating rhoptry biogenesis, their

structure, and internal compartmentalization are unknown.

Rhoptry neck protein 11, or RON11 (PF3D7_1463900), is a seven transmembrane

domain protein with a pair of calcium-binding EF-hand domains at its C-terminus that is

conserved in Apicomplexans. Initial attempts to disrupt *P. berghei* RON11 were

unsuccessful (Lasonder et al. 2008; van Ooij et al. 2008), while a recent knockdown

showed a potential role in sporozoite invasion and gliding motility (Bantuchai et al.

2019). In contrast, RON11 disruption in *T. gondii* showed only a minor fitness defect,

suggesting a non-essential role (Wang et al. 2016). However, nothing is known about

the role of RON11 in *P. falciparum*. In this study, we employed the tetR-DOZI

conditional knockdown system (Rajaram, Liu, and Prigge 2020) to investigate the

function of RON11 in the *P. falciparum* intraerythrocytic asexual life cycle. Our findings

revealed that RON11 is essential for the intraerythrocytic growth of *P. falciparum.*

Parasites lacking RON11 are unable to invade RBCs. Anti-RON11 antibodies are able

to block merozoite invasion into RBCs suggesting that RON11 plays an essential role

during invasion. Surprisingly, RON11 knockdown results in merozoites that have a

single rhoptry. These single rhoptry merozoites are unable to complete invasion, even

though they attach and secrete the contents from the single rhoptry into RBCs.

Collectively, our data show that RON11 plays an essential role in regulating the *de novo*

formation of the second rhoptry during merozoite segmentation and required for

merozoite invasion.

RESULTS

RON11 is essential for *P. falciparum* intraerythrocytic life cycle

To investigate the role of RON11 in *P. falciparum*, we generated RON11 conditional knockdown parasite lines employing the tetR-DOZI aptamer system, termed as RON11[apt] (Rajaram, Liu, and Prigge 2020). Using CRISPR/Cas9 technology, we integrated the tetR-DOZI system into the endogenous loci of RON11, leading to the expression of a hemagglutinin (HA)-tagged protein under the regulation of anhydrotetracycline (aTc) (Fig. 1A). PCR analysis of genomic DNA from RON11[apt] parasites showed correct integration at the endogenous loci (Fig. 1B). To test the efficiency of the knockdown system, we evaluated protein expression both in the presence (+RON11) and absence (-RON11) of aTc via western blotting. Lysates from synchronized RON11[apt] ring stage parasites cultured without aTc for 44 hours post-invasion (hpi) (until schizont stage) exhibited around 97% of reduction in protein expression (Fig. 1C), confirming the system's suitability for studying RON11 function. Parasite growth was measured using flow cytometry (Fig. 1D). These data show that RON11 is essential for parasite expansion, as -RON11 parasites failed to progress past their first life cycle (Fig. 1D).

Furthermore, we assessed the subcellular localization of RON11, which was previously observed in rhoptries of *P. berghei* schizonts (Bantuchai et al. 2019) (Fig. 1E). RON11 was localized with two well-known markers of the rhoptry neck (RON4) and rhoptry bulb (RAP1) (Fig. 1E). Our data showed the presence of RON11 in the rhoptry neck, as suggested by its colocalization with RON4 (Fig. 1E, top panels, and 1F). RON11 did not colocalize with RAP1, suggesting that RON11 is confined to the neck region (Fig. 1E, bottom panels, and 1F). Additionally, we observed RON11 at the parasite periphery of newly invaded ring-stage parasites (Supp. Fig. 1), suggesting that RON11 might localize to the parasitophorous vacuole (PV) after invasion, similar to RAP1 (Anaguano et al. 2023; Liffner, Cepeda Diaz, Blauwkamp, Anaguano, Frolich, et al. 2023; Riglar et al. 2011). Taken together, these data show RON11 localizes to the rhoptry neck in schizonts, and in early-ring-stage parasites, it localizes at the parasite periphery.

Figure 1. RON11 is essential for intraerythrocytic growth. (A) Schematic showing the integration of the repair plasmid used to introduce the tetR-DOZI system in the genomic loci of RON11. Cas9 introduces a double-stranded break at the C-terminus of the RON11 locus. The repair plasmid provides homology regions for double-crossover homologous recombination, introducing the system and providing a new 3'UTR to the gene. **(B)** PCR confirming integration at the RON11 locus. Amplicons were amplified from genomic DNA isolated from mutant and the parental line (NF54[attB]) (Nkrumah et al. 2006). Primers were designed to amplify two regions: between the C-terminus and the 3'UTR of RON11, and between the C-terminus of RON11 and the new aptamer tandem. **(C)** Western blot of lysates from RON11[apt] schizont grown in the presence and absence of 0.5 µM aTc. Lysates were probed with antibodies against HA (RON11) and EF1α (loading control). The protein marker sizes are shown on the left. (n = 4 biological replicates). **(D)** Growth of RON11[apt] parasites over 3 intraerythrocytic life cycles in the presence and absence of 0.5 µM aTc. Synchronous parasites were collected every 48 hours, stained with the DNA dye Hoechst 33342, and measured via flow cytometry. One representative data set of three biological replicates shown. (n = 3 technical replicates; error bars = standard deviation (SD)). **(E)** IFAs showing the localization of RON11 in RON11[apt] schizont with respect to two rhoptry markers: RON4 (neck) and RAP1 (bulb). Synchronous parasites were fixed with acetone and stained with specific antibodies. Images from left to right are phase-contrast, DAPI (nucleus, cyan), anti-HA (magenta), anti-RON4 or RAP1 (yellow),

and fluorescence merge. Z stack images were deconvolved and projected as a combined single image.
Representative images of 3 biological replicates. **(F)** Quantification of the colocalization of RON11[apt] with
RON4 and RAP1 as determined by the Pearson's correlation coefficient (PCC). (n = 3 biological
replicates, 4 schizonts per replicate. Error bars = standard error of the mean (SEM); ****p < 0.0001 by
unpaired two-tailed t-test).

RON11 is required for merozoite invasion

As RON11 is known to be highly expressed in schizont stages (Bozdech et al. 2003;
Llinás et al. 2006), and we showed its knockdown inhibits parasite growth (Fig. 1D).
Therefore, we wanted to determine if RON11 is necessary for schizont development,
egress or merozoite invasion. To define the specific stage of the intraerythrocytic
lifecycle that requires the function of RON11, we monitored the progression of
synchronized RON11[apt] parasites through the asexual life cycle, both in the presence
and absence of aTc. Samples were collected at all the stages, starting with tightly
synchronized schizonts (46-48 hpi), observed using Hema3-stained thin blood smears
and light microscopy. Our findings show that, both +RON11 and -RON11 parasites
developed normally from rings through schizonts, and after completion of schizogony,
daughter merozoites egressed at 48 hpi. (Fig. 2A). However, upon RON11 knockdown,
the egressed merozoites did not produce rings, instead, they accumulated around the
RBCs (Fig. 2A). To quantify our microscopy-based observations, we used flow
cytometry to measure parasite transition from schizonts into rings. We confirmed that
even though schizonts egressed in the absence of RON11, rings were not detected at
any time point post-egress (Fig. 2B). Together these findings show that the knockdown
of RON11 prevents merozoite invasion, while having no effect on intraerythrocytic
development or on the egress of daughter merozoites.

We hypothesized that RON11 may have a specific function in merozoite invasion. To
independently test this hypothesis, we generated antibodies against the N-terminus of
RON11 (aa32-442) (Supp. fig. 2A-B). As RON11 contains seven predicted
transmembrane domains, prediction of protein orientation on the rhoptry membrane
suggests that the N-terminus may face the rhoptry lumen. We hypothesized that this

region may become accessible to antibodies during merozoite invasion when rhoptry
contents are secreted. Therefore, if RON11 plays a role in merozoite invasion, then anti-
RON11 antibodies should block parasite invasion into the host RBC. Western blot
analysis detected a single high molecular weight protein of the expected size in schizont
lysate (Supp. fig. 2D), showing that anti-RON11 antibodies were specific for RON11.
We also confirmed the specificity of the anti-RON11 antibodies using
immunofluorescence microscopy (Supp. Fig. 2E). The anti-RON11 antibody staining
was colocalized with the anti-HA staining in the RON11[apt] (Supp. fig. 2E). We then
evaluated the ability of anti-RON11 antibodies to block merozoite invasion and therefore
reduce parasite multiplication. In these neutralization assays, parasites were incubated
with anti-RON11 antibodies and their ability to proliferate was measured. *P. falciparum*
parasites (3D7 strain) when incubated with various concentrations of total anti-RON11
IgG. Parasite growth was measured by flow cytometry (Fig. 2C). Antibodies against
RON11 exhibited concentration-dependent growth inhibitory activity, with around 80%
growth inhibition at a concentration of 15 mg/mL (Fig. 2C). This suggests that the N-
terminus of RON11 is accessible to antibodies during merozoite invasion and may
interfere with its function in RBC invasion. Together with our RON11 conditional
knockdown data, these findings support a putative role for RON11 in merozoite
invasion.

Figure 2. RON11 is required for merozoite invasion. (A) Representative Hema-3-stained blood-smears
showing the development of RON11^{apt} parasites in the presence and absence of 0.5 µM aTc.
Synchronous parasites were smeared, stained, and imaged at different time points by light microscopy.
(B) Tracking of the development of synchronous RON11^{apt} parasites over 8 hours post-egress in the
presence and absence of 0.5 µM aTc. Schizonts were tightly synchronized using 2 µM Compound 1.
Synchronized parasites were collected every 2 hours, stained with Hoechst 33342, and measured via
flow cytometry to distinguish between rings and schizonts. One representative data set of three biological
replicates shown. (n = 3 technical replicates; error bars = SD). **(C)** *In vitro* neutralization assay was
performed against 3D7 parasites using total IgG purified from rats immunized with rMBP-RON11 (N-
term). Potential influence of anti-MBP antibodies was evaluated by pre-incubating IgG with rMBP(3µM).
Parasites were stained with SYBR green and parasitemia was measured by flow cytometry. Growth
inhibitory activity is represented as the percentage (%) of inhibition relative to the "no-IgG" control. (n = 2-
4 biological replicates in duplicate; error bars = SEM).

RON11 is not required for merozoite attachment or eventual rhoptry discharge

Merozoite invasion is a complex multistep process, which involves the participation of multiple regulatory factors. In general terms, invasion can be divided into several discrete steps: attachment, apical end reorientation, RBC deformation, and pore formation, rhoptry secretion and tight junction formation, internalization and PVM sealing, and echinocytosis (Cowman et al. 2017; Groomes, Kanjee, and Duraisingh 2022). To dissect the specific role of RON11 during merozoite invasion, we investigated the following specific events during invasion: attachment, deformation, rhoptry secretion and echinocytosis.

First, we used live-cell imaging to examine the merozoite-RBC interactions in recently egressed RON11[apt] merozoites in the presence and absence of RON11 (Fig. 3A, Supp. Fig. 3, Movie S1, and Movie S2). As shown previously (Fig. 2A and 2B), RON11 knockdown completely inhibited invasion in merozoites that induced deformation (n = 0/53), in contrast to the +RON11 merozoites in which 77.2% (n = 27/35) of merozoites invaded successfully (Fig. 3A, B). These invasion defects in -RON11 merozoites were observable even though these merozoites were capable of generating strong deformations of the RBC membrane (Fig. 3B, C and Supp. Fig 3D) similar to +RON11 merozoites (p = 0.048). The -RON11 merozoites were capable of inducing echinocytosis (n = 33/44) in the attached RBCs (Fig 3D and Movie S2). We also noticed that -RON11 merozoites continue to engage with the RBCs (deformation) for longer periods of time and remain attached to RBCs, eventually leading to echinocytosis (Fig. 3C, D and Supp. Fig 3C). However, no significant difference between +RON11 and -RON11 merozoites was observed in the time that it took the merozoites from their initial contact with RBCs to the beginning of echinocytosis (Fig. 3D). Altogether, these data suggest that RON11 is not required for initial attachment, reorientation, and deformation of RBCs, but is essential for internalization into the host cell. The longer deformation eventually leads to echinocytosis, which is thought to be indicative of secretion of contents from rhoptries into the RBC. Our data suggests that while -RON11 parasites

are capable of inducing echinocytosis though the timing of echinocytosis is prolonged in
the absence of RON11 (Fig. 3D, E and Supp. Fig. 3B, C).

To confirm our observation that there was no defect in the attachment of -RON11
parasites to host RBCs, we used a cytochalasin-D-based strategy to observe
merozoite-RBC interactions (Riglar et al. 2011). Cytochalasin D is an actin
polymerization inhibitor that disrupts the activity of the merozoite actomyosin motor,
preventing internalization but not affecting the previous steps on merozoite invasion,
such as attachment and rhoptry secretion (Boyle et al. 2010; Geoghegan et al. 2021;
Riglar et al. 2011). Samples for analysis were collected from Compound 1 treated
mature schizonts that were washed and allowed to invade fresh RBCs in the presence
of cytochalasin D (Supp. fig. 3E).

We used light microscopy to quantify +RON11 and -RON11 merozoites attached to
RBCs. In merozoites incubated in cytochalasin D, no significant difference in the
numbers of attached +RON11 or -RON11 merozoites was observed (Supp. fig. 3F).
When merozoites were allowed to invade without cytochalasin D, we found more
attached -RON11 merozoites, compared to the cytochalasin-D-treated +RON11
merozoites (Supp. fig. 3G). As expected, the number of attached merozoites dropped
significantly when cytochalasin D was removed from +RON11 merozoites, as they were
able to invade RBCs and form rings (Supp. fig. 3G). Together, these data demonstrated
that RON11 is not required for merozoite attachment to RBCs.

Our live imaging data suggests that echinocytosis, albeit delayed, occurs in -RON11
parasites (Fig. 3A, D, E). To confirm that -RON11 parasites are able to secrete rhoptry
contents into the host RBC, we used IFAs to analyze the secretion of rhoptry proteins
from cytochalasin-D-treated merozoites into RBCs in both +RON11 and -RON11
parasites. We selected RAP1 as a rhoptry secretion marker because it has been
previously shown to be secreted into RBCs cytoplasm in the presence of cytochalasin D
(Riglar et al. 2011). We found that -RON11 merozoites were able to secrete RAP1 into
RBCs similarly to +RON11 merozoites (Fig. 3F). Comparing the number of attached
merozoites that secreted RAP1 into RBCs versus the ones who did not, no significant
differences were observed between +RON11 and -RON11 parasites (Fig. 3G). These

results, together with our observation of prolonged deformation prior to echinocytosis by

live video microscopy, suggest that RON11 deficient merozoites are able to eventually

secrete rhoptry contents into the RBC.

Figure 3. RON11 is not required for merozoite attachment nor rhoptry secretion. (A) Representative time-course images from Movie S1 (+RON11) and Movie S2 (-RON11) showing RON11[apt] merozoites (arrowhead) interacting with RBCs in the presence and absence of 0.5 µM aTc. Images show the different stages of merozoite invasion post RBC initial contact. One representative data set of three biological replicates shown. **(B)** Quantification of merozoites that induced deformation in RBCs that completed invasion or remained attached. (n = 3 biological replicates, 35 merozoites for +RON11 and 53 for -RON11). **(C)** Duration of merozoite-induced RBC deformation quantified using live video microscopy. (n = 3 biological replicates, 14 merozoites for +RON11 and 12 for -RON11; error bars = SEM; **p < 0.01 by unpaired two-tailed t-test). **(D)** Duration between merozoite first contact with RBCs and echinocytosis quantified using live video microscopy. (n = 3 biological replicates, 35 merozoites for +RON11 and 53 for -RON11; error bars = SEM; ns = non-significant by unpaired two-tailed t-test). **(E)** Duration of echinocytosis quantified using live video microscopy. (n = 3 biological replicates, 23 merozoites for +RON11 and 29 for -RON11; error bars = SEM; ***p < 0.001 by unpaired two-tailed t-test). **(F)** Representative IFAs showing RON11[apt] attached-merozoites secreting RAP1 into RBCs in the presence or absence of aTc, after incubation with cytochalasin D. Parasites were fixed with PFA and stained with specific antibodies. Images from left to right are phase-contrast, DAPI (nucleus, blue), anti-HA (green), anti-RAP1 (red), and fluorescence merge. Z stack images were deconvolved and projected as a combined single image. Representative images of 3 biological replicates. **(G)** Quantification of RON11[apt] attached-merozoites secreting RAP1 into RBCs. Slides were blinded and secretion events were scored and represented as the percentage of events per 50 attached merozoites (n = 3 biological replicates, 50 events per replicate. Error bars = SEM; ns = non-significant by unpaired two-tailed t-test).

RON11 knockdown results in merozoites with one rhoptry

Given previous observations that other rhoptry membrane-associated proteins, such as RAMA, play a role in rhoptry formation and transport of rhoptry proteins (Sherling et al. 2019), we sought to determine whether RON11 has a similar function in rhoptry biogenesis. Therefore, to determine if RON11 functions in rhoptry biogenesis, we employed ultrastructural expansion microscopy (UExM) in combination with NHS-ester staining, which allows us to observe rhoptry structures in high resolution (Liffner, Cepeda Diaz, Blauwkamp, Anaguano, Frölich, et al. 2023). We analyzed tightly synchronized schizonts that were cultured in the presence or absence of aTc. ML10 is a highly specific inhibitor of the cGMP-dependent protein kinase (PKG) that arrests

parasites prior to egress (Ressurreição et al. 2020), allowing for a tighter
synchronization.

We observed fully formed rhoptries exhibiting their distinctive structures: a slender and
elongated neck at the apex, connected to an apical ring, and a distal rotund bulb (Fig.
4A, top panels), as previously reported (Liffner, Cepeda Diaz, Blauwkamp, Anaguano,
Frolich, et al. 2023). Strikingly, -RON11 merozoites had only one rhoptry (Fig. 4A and
4B). These single rhoptries displayed no discernible structural alterations, as the neck,
and the bulb, were still present in the proper orientation positioned at the apical end of
the merozoite, as defined by the observed apical ring (Fig. 4A, bottom panels). In rare
cases, we observed -RON11 schizonts with several rhoptries amassed in regions that
appeared to be located outside of merozoites (Supp. fig. 4A). Basal complexes were
visible at their maximum contraction at the basal end of the analyzed merozoites
(Liffner, Cepeda Diaz, Blauwkamp, Anaguano, Frolich, et al. 2023), suggesting that
RON11 knockdown had no impact on merozoite segmentation, and confirming that the
analyzed merozoites were fully developed (Fig. 4A and Supp. fig. 4A). To validate this
finding, we quantified the number of merozoites per schizont and found that the
absence of RON11 had no effect on the total number of merozoites, even though the
number of rhoptries was reduced by half (Supp. fig. 4B, C). To our knowledge, this
marks the first observation of such a unique phenotype, single rhoptry merozoites (or
changes in rhoptry numbers), in apicomplexan parasites.

Figure 4. RON11 knockdown generates merozoites with single rhoptries. (A) Representative images of RON11[apt] schizonts showing the structure of rhoptries within fully developed merozoites in the presence and absence of 0.5 µM aTc. ML10-arrested schizonts were expanded by U-ExM, fixed with PFA, and stained with NHS-Ester (greyscale), anti-HA (magenta), and the DNA dye SYTOX (cyan). Selected Z stack images were projected as a combined single image. Number on image = Z-axis thickness of projection in µm. **(B)** Quantification of merozoites with single, dual or multiple rhoptries in the presence and absence of 0.5 µM aTc. (n = 4 biological replicates, 28 schizonts for +RON11 and 36 for -RON11; error bars = SEM; ****$p < 0.0001$ by unpaired two-tailed t-test)

To determine if RON11 knockdown disrupts the localization of rhoptry proteins, we analyzed the localization of other subcellular markers in segmented schizonts using UExM. We confirmed that RON11 localizes to the neck of fully formed rhoptries, along with its colocalization with RON4 (Fig. 5A). These UExM data show that the localization of RON4 and RAP1 remained unaltered in the single rhoptries in -RON11 parasites (Fig. 5A). Similarly, RON4 and RAP1 retained their localization in rhoptries of free -RON11 merozoites (Supp. fig. 5).

Since -RON11 parasites had only one rhoptry, this suggests that there is a defect in the biogenesis of the second rhoptry. This raised the question, does single rhoptry in -RON11 schizonts have the same amount of rhoptry proteins as the two rhoptries in +RON11 parasites? To answer this question, we assessed the expression of RON4 and RAP1 in fully matured +RON11 and -RON11 schizonts (48 hpi) using western blotting. The total amount of RON4 and RAP1 was reduced by approximately half in -RON11 parasites (Fig. 5B and 5C). This finding could be attributed to the reduction of the number of rhoptries by half, aligning with recent findings suggesting that rhoptry pairs form *de novo* during schizogony rather than through duplication from a single mother rhoptry (Liffner, Cepeda Diaz, Blauwkamp, Anaguano, Frolich, et al. 2023).

Our data show that the single rhoptry -RON11 merozoites are unable to invade host RBCs. The biogenesis as well as the rhoptry biogenesis defect could be due to defective maturation of other rhoptry proteins by the rhoptry-localized aspartic protease, Plasmepsin IX (PMIX) (Favuzza et al. 2020; Nasamu et al. 2017; Pino et al. 2017) . PMIX processes several rhoptry proteins, including invasion ligands (Favuzza et al. 2020; Nasamu et al. 2017; Pino et al. 2017). Therefore, we tested if RAP1 processing by PMIX is inhibited in -RON11 parasites (Fig. 5D). We assessed the cleavage of RAP1 by PMIX into 67 kDa (p67) and 82 kDa (p82) forms in lysates collected from mature +RON11 and -RON11 schizonts (48hpi) using western blotting (Fig. 5D). The ratio between the cleaved forms of RAP1 (p67:p82) did not differ between +RON11 and -RON11 parasites. These data suggest that PMIX activity was intact in the single rhoptry -RON11 merozoites.

To rule out a generalized defect in the secretory pathway caused by the loss of RON11,
we evaluated the expression of markers of other subcellular compartments, such as the
micronemes (AMA1) and merozoite surface (MSP1). We used IFAs and western
blotting on fully mature E64-treated schizonts cultured in the absence of RON11. We
observed no effect on protein content or secretion of MSP1 to the merozoite surface
under RON11 knockdown (Supp. fig. 6A and 6B). Similarly, AMA1 retained its
micronemal and secreted locations in the absence of RON11 (Supp. fig. 6C). These
data show that the loss of RON11 did not lead to a generalized defect in protein
secretion.

Figure 5. RON11 knockdown does not affect the localization and processing of neck and bulb rhoptry proteins. (A) Representative images of RON11[apt] parasite showing the localization of RON11 with respect to RON4 and RAP1 in the presence and absence of aTc. E64-arrested parasites were expanded by U-ExM, fixed with PFA, and stained with NHS-Ester (greyscale), anti-HA (magenta), and RON4 or RAP1 (yellow). Selected Z stack images were projected as a combined single image. Number on image = Z-axis thickness of projection in μm. **(B)** Western blot of parasite lysates isolated from E64-

arrested RON11[apt] parasites in the presence and absence of aTc. Samples were probed with antibodies

against RON4, RAP1 and EF1α (loading control). The protein marker sizes are shown on the left.

Representative blot of 4 biological replicates shown. **(C)** Quantification of the relative level of RON4 and

RAP1 total protein content in E64-arrested RON11[apt] parasites in the presence and absence of aTc. Band

intensities were normalized to the loading control, EF1α, and are presented as normalized arbitrary units

(A.U.) (n = 4 biological replicates; error bars = SEM; **p < 0.01 by unpaired two-tailed *t*-test). **(D)**

Quantification of RAP1 processing in E64-arrested RON11[apt] parasites in the presence and absence of

aTc. Band intensities were normalized to the ratio of processed RAP1 (p67/p82) (n = 3 biological

replicates; error bars = SEM; ns = non-significant by unpaired two-tailed *t*-test).

RON11 triggers the biogenesis of the last rhoptry pair

Rhoptries are first observed early during schizogony (Mahajan et al. 2008; Liffner,

Cepeda Diaz, Blauwkamp, Anaguano, Frolich, et al. 2023) and our data shows that -

RON11 merozoites have a single rhoptry (Fig. 4). When does rhoptry biogenesis fail

during schizogony and how does this generate single rhoptry merozoites? Because

schizogony is known to last approximately 15 hours (Voß et al. 2023), we designed a

rescue assay to define the window of time within schizogony where restoring RON11

might restore parasite growth. Synchronized ring-stage RON11[apt] parasites were grown

with or without aTc and when these parasites reached schizogony (44 to 50 hpi), aTc

was added back to the parasites at different time points (Fig. 6). The ability of these

schizonts to form rings was measured using flow cytometry. Supplementation with aTc

as late as 46 hpi restored the ability of -RON11 schizonts to generate ring stage

parasites (Fig. 6). However, aTc supplementation after 46 hpi was unable to rescue

growth (Fig. 6). These findings suggested that RON11 has an essential role during the

final hours of schizont development (Fig. 6).

Figure 6. RON11 knockdown phenotype is rescued by addition of aTc during the last 2 hours of the asexual life cycle. Growth of RON11[apt] parasites in the presence and absence of aTc. Tightly synchronous parasites were grown for 44 hpi and aTc was added back every two hours. Samples were collected at 24, 44 and 56 hpi, fixed in PFA, stained with the DNA dye Hoechst 33342, and measured via flow cytometry to differentiate between rings and schizonts. (n = 3 biological replicates; error bars = SEM; *ns = non-significant; ***p < 0.001; ****p < 0.0001 by one-way ANOVA compared to + RON11).

As our data shows that RON11 acts during the last hours of schizogony (Fig. 6), we hypothesized it might have a role in regulating the formation of the last rhoptry pair during late schizogony. To test our hypothesis, we employed UExM in combination with NHS-ester to follow the development of rhoptry biogenesis during schizogony in the presence and absence of RON11. It has been previously shown that during their biogenesis, rhoptries associate with the branches of the outer CPs in a ratio of one rhoptry per branch during the early mitotic events of schizogony (Fig. 7A) (Bannister et al. 2000; Liffner, Cepeda Diaz, Blauwkamp, Anaguano, Frolich, et al. 2023; Rudlaff et al. 2020). Once merozoite segmentation starts, each outer CP branch accommodates two rhoptries, which results in the final rhoptry pair within each merozoite (Fig. 7A) (Liffner, Cepeda Diaz, Blauwkamp, Anaguano, Frolich, et al. 2023). By analyzing at 44-48 hour synchronized schizonts, we found that the formation and segregation of rhoptries during the initial mitotic events was not inhibited in -RON11 parasites (Fig. 7B and 7C). Surprisingly, we discovered that the loss of RON11 impaired the *de novo* formation of the second rhoptry after the final mitotic event (Fig. 7D). Initiation of merozoite segmentation on those events was confirmed by the presence of basal

468   complex rings in association with the CP/rhoptry-complex densities (Fig. 7). Altogether

469   these data confirm our thesis that RON11 regulates the *de novo* biogenesis of the

470   second rhoptry after the last mitotic event during schizogony (Fig. 7).

Figure 7. RON11 is required for the *de novo* biogenesis of the final rhoptry pair during merozoite segmentation. (A) Schematic of the current model for *de novo* formation of rhoptries during schizogony. **(B, C and D)** Representative images of different steps of rhoptry biogenesis in RON11[apt] late schizonts in the presence and absence of aTc: **(B)** Two rhoptries associated to each branch of a CP in the presence and absence of aTc, **(C)** Single rhoptries being segregated with each branch of a dividing CP during mitosis, and **(D)** Final rhoptry pairs being segregated with a CP branch during the final mitotic event. Basal complexes are demarcated by a discontinuous line. Late-schizont parasites were expanded by U-ExM, fixed with PFA, and stained with NHS-Ester (greyscale), anti-HA (magenta), and anti-RAP1 (yellow). Selected Z stack images were projected as a combined single image. Number on image = Z-axis thickness of projection in µm.

DISCUSSION

Similar to most rhoptry proteins studied to date in the asexual RBC stages (Liffner et al. 2021), we have demonstrated that RON11 is indispensable for parasite development, particularly merozoite invasion of RBCs. This aligns with previous failed attempts to generate RON11 knockouts in *P. berghei* parasites which supported essentiality of RON11 in blood stages (Lasonder et al. 2008; van Ooij et al. 2008). We used a combination of live cell video microscopy cytochalasin D- based attachment assays to show that initial engagement with RBC occurs normally in absence of RON11 but merozoites fail to enter into RBCs. While the actomyosin motor and its components are well-established as being involved in merozoite internalization (Blake, Haase, and Baum 2020; Perrin et al. 2018), no rhoptry protein has been associated with a role in this final stage of invasion. Interestingly, we observed by live imaging that -RON11 merozoites that are unable to complete invasion, are still able to induce strong deformations on RBCs. Since gliding motility is required for deformation (Yahata et al. 2021; Hart et al. 2023), this suggests that RON11 may not be required for motility of *P. falciparum* merozoites, unlike previously observed for *P. berghei* sporozoites (Bantuchai et al. 2019). RON11 deficient merozoites engage and deform host cells for significantly longer before the RBCs undergo echinocytosis, a process that is thought to follow rhoptry secretion (Weiss et al. 2015). Therefore, RON11 may not be essential for rhoptry discharge but may play a role in the efficiency or timing of rhoptry secretion. In

agreement with this observation, -RON11 merozoites in the presence of cytochalasin D
proceeded to attach and secrete RAP1 into RBCs.

Following the kinetics of invasion by live cell microscopy also allows us to surmise the
step of invasion where RON11 functions. Previous studies using this approach revealed
key receptor-ligand interactions as molecular determinants of specific steps in
merozoite invasion (Weiss et al. 2015). As -RON11 merozoites are able to attach,
reorient and deform RBCs, it rules out a role for RON11 in these early steps. Following
deformation, the RH5-basigin interaction leads to an opening at the interface of the
merozoite and RBC initiating the secretion of rhoptry neck/bulb proteins. Blocking this
interaction leads to normal deformation but prevents rhoptry secretion and no
echinocytosis (Weiss et al. 2015). Finally, the AMA1-RON2 interaction mediates tight
junction formation that provides an anchor and allows the actomyosin motor to drive the
merozoite inside the RBC (Srinivasan et al. 2011). Peptide inhibitors of this interaction
lead to strong deformation and echinocytosis, a result of rhoptry secretion (Weiss et al.
2015). Interestingly, RON11 knockdown resulted in merozoites unable to internalize but
stay attached to the RBC and induce echinocytosis. This suggests that RON11 perhaps
functions downstream of the RH5-basigin interaction. Furthermore, deformation times
were significantly longer with -RON11 merozoites compared to +RON11 merozoites
(average of 21 sec and 6.5 sec respectively). Rhoptry secretion coincides with
internalization and immediately follows the deformation step. The longer duration of
deformation may therefore reflect a temporal defect in rhoptry secretion in the absence
of RON11. These data are consistent with IFA data that show that RON11 is not
required for the secretion of the rhoptry bulb protein, RAP1. However, since IFA uses
fixed cells, we cannot determine the kinetics of rhoptry secretion. Thus, it is possible
that RON11 may play a role in ensuring timely secretion of rhoptry contents after the
formation of the tight junction.

Calcium signaling plays a critical role during merozoite invasion. Following RBC
deformation and immediately before internalization a prominent calcium signal is
observed at the apical end of the invading merozoite, suggesting a potential role for
calcium sensing protein(s) at this step of invasion (Weiss et al. 2015). This signal occurs

downstream of RH5-basigin interaction and is not inhibited by AMA1-RON2 blocking peptide agonists (Weiss et al. 2015). It is possible that RON11 through its C-terminal EF hand motifs, which can bind calcium, may play a role in regulating the kinetics of rhoptry secretion and/or subsequent steps in invasion.

Rhoptries are critical secretory organelles conserved among apicomplexans, with differences in numbers depending on the species as well as between merozoite and sporozoite. During *P. falciparum* intraerythrocytic development, rhoptries are initially observed forming *de novo* during early schizogony (Liffner, Cepeda Diaz, Blauwkamp, Anaguano, Frolich, et al. 2023). Then, rhoptries replicate throughout multiple mitotic events until each merozoite contains two rhoptries by the end of the cycle. In contrast, *T. gondii* can have as many as 8-12 rhoptries, but only two are known to release their contents during invasion (Mageswaran et al. 2021). Rhoptry biogenesis remains a poorly understood process, primarily due to the small size of *P. falciparum*'s rhoptries, which, coupled with the limited resolution offered by conventional microscopy, has made their study challenging. Recent advances in microscopy techniques, such as UExM, have opened new possibilities for exploring these micron-sized secretory organelles (Liffner and Absalon 2023). Using UExM, we observed that RON11 is required for the proper biogenesis of rhoptry pairs, as its absence led to merozoites with single rhoptries, an unprecedented phenotype that has not been observed in any other apicomplexan system. Only three parasite proteins, Sortilin, RAMA and CERLI2, have been shown to have a role in the biogenesis of rhoptries in *P. falciparum* (Hallée et al. 2018; Liffner et al. 2022; Sherling et al. 2019). None of them share any structural similarities, localize to the same region within rhoptries, or generate a knockdown phenotype analogous to RON11. For instance, RAMA and CERLI2 loss causes morphological alterations in the rhoptry necks (Liffner et al. 2022; Sherling et al. 2019), while RON11's single rhoptries did not display any apparent structural change at the resolution of UExM imaging. Our data strongly suggests that RON11 is required for *de novo* rhoptry biogenesis at the end of schizogony, when daughter merozoites are being formed via cytokinesis. This thesis is supported by our initial observations by UExM that rhoptries are formed and segregated with the CPs without issues in -RON11 parasites during the early asynchronous mitotic events of schizogony. These observations align

with our findings that -RON11 parasite growth is rescued by reversing the knockdown
as late as 46 hours post invasion, in the 48-hour asexual life cycle.

In contrast, we observed that the *de novo* formation of the initial rhoptry was not
disrupted in -RON11 parasites during schizogony. However, during cytokinesis, we
noticed that one rhoptry was being distributed along with the dividing CPs into a nascent
-RON11 merozoite, instead of two rhoptries as observed in +RON11 parasites. These
observations are in agreement with our findings of a roughly 50% reduction in the
number of rhoptries per schizont. Interestingly, we observed that RON11 knockdown
also led to about a 50% reduction in the quantity of other rhoptry proteins, which is
consistent with the finding that only half the amount of rhoptries are formed. This is a
somewhat surprising result as prior work on other *P. falciparum* organelles suggest that
a defect in organelle biogenesis does not lead to a loss in expression of organellar
proteins. For example, upon loss of the parasite plastid known as the apicoplast,
proteins targeted to the apicoplast are synthesized and appear to be stuck in secretory
vesicles (Florentin et al. 2017, 2020; Yeh and DeRisi 2011). Our data suggest that
RON11 may function in a pathway that regulates synthesis of rhoptry proteins, either via
transcription or translation. These data suggest that a signaling pathway may exist that
ensures synthesis of rhoptry proteins only when rhoptry biogenesis occurs. Future
transcriptomic and proteomic studies should resolve the role of RON11 in this *de novo*
biogenesis pathway. Since RON11 has a pair of calcium binding EF-hand domains, it
raises the possibility that a calcium-dependent pathway may function in biogenesis of
rhoptries. During the final hours of schizogony, calcium-dependent pathways also
trigger egress of the merozoites from the host RBC (Dvorin and Goldberg 2022). This
study raises the possibility that a similar calcium signal may also kickstart, via RON11,
the *de novo* biogenesis of the second rhoptry to ensure that the egressing merozoites
have two rhoptries each for invading the next host RBC. It is unlikely that these are the
same calcium signals because blocking the egress-specific calcium signal using PKG
inhibitors does not prevent rhoptry biogenesis.

Since -RON11 merozoites with a single, functional rhoptry attach and secrete their
contents into the host RBC but fail to internalize into the host cell, this raises the

question: do *P. falciparum* merozoites need two rhoptries to invade host RBCs? One
possibility is that since the single rhoptry -RON11 merozoites have half as much rhoptry
contents compared to the +RON11 merozoites, the failure to internalize may be due to
insufficient amounts of invasion ligands such as RON2. It is possible that while RAP1 is
secreted from the single rhoptry in -RON11 parasites, secretion of other rhoptry neck
proteins is disrupted or the timing of rhoptry secretion is disrupted in the absence of
RON11. Distinguishing between these models will require developing tools to observe
rhoptry secretion in live parasites. Another possibility is that RON11 has another
biogenesis-independent function in merozoite invasion. Other rhoptry proteins have
been shown to have multiple independent functions (Counihan et al. 2017; Ito,
Schureck, and Desai 2017; Schureck et al. 2021; Sherling et al. 2019). Our data
suggests that RON11 may function in merozoite invasion because rhoptry proteins do
localize correctly in -RON11 parasites, they are processed accurately by PMIX, and
anti-RON11 antibodies inhibit intraerythrocytic growth of *P. falciparum.*

Together, these data suggest a model where RON11 has two functions in the final
hours of the intraerythrocytic life cycle of *P. falciparum.* During segmentation of the
schizont, RON11 is required to trigger the *de novo* biogenesis of the final rhoptry pair,
which ensures each merozoite ends up with two rhoptries. After the merozoite makes
contact with a new host RBC, RON11 is required for internalization into the new host
cell. Calcium signals play a critical role in both merozoite egress as well as invasion. It
remains to be seen if the calcium binding EF-hand domains of RON11 play a role in
either its function in rhoptry biogenesis or its function in merozoite invasion. The
discovery of the biological function of RON11 now provides an avenue to investigate the
molecular mechanisms driving rhoptry biogenesis and how it is linked to merozoite
invasion.

MATERIALS AND METHODS

Construction of RON11 plasmids

Genomic DNA was isolated from *P. falciparum* NF54^{attB} cultures using the QIAamp DNA blood kit (Qiagen). PCR products were inserted into the respective plasmids using the NEBuilder HiFi DNA Assembly system (NEB). All constructs used in this study were confirmed by sequencing. All primers used in this study are in Table 3.1.

For generation of the plasmid pKD-RON11-Apt, sequences of approximately 500 bp of homology to the RON11 (Pf3D7_1463900) C-terminus and 3'UTR were amplified using primer pairs P3-P4 and P5-P6, respectively (Table 3.1). Amplicons were then inserted into pKD (Rajaram, Liu, and Prigge 2020) digested by with AatII and AscI. For expression of a RON11 gRNA, oligo P7 was inserted into cut pUF1-Cas9 (Table 3.1).

Parasite culture and transfections

Plasmodium parasites were cultured in RPMI 1640 medium (NF54^{attB}, RON11^{apt}) supplemented with AlbuMAX I (Gibco), and transfected as described earlier (Kudyba et al. 2018).

For generation of RON11^{Apt} parasites, a mix of two plasmids (50 μg each) were transfected into NF54^{attB} parasites in duplicate. The plasmid mix contained the plasmid pUF1-Cas9-SBP1gRNA, which contains the DHOD resistance gene, and the marker-free plasmid pTOPO-SBP1-TbID. Drug pressure was applied 48 h after transfection, using 1μM DSM1 (Ganesan et al. 2011) and selecting for Cas9 expression. After parasites grew back from transfection, integration was confirmed by PCR, and then cloned using limiting dilution. After clonal selection, cultures were transferred to biotin-free medium without DSM1.

For generation of RON11^{apt} parasites, the pKD-RON11-Apt plasmid (20 μg) and the respective pUF1-Cas9 plasmid (50 μg) were transfected into NF54^{attB} parasites in duplicate. Before transfection pKD plasmids were digested overnight with EcoRV (NEB). The enzyme was then subjected to heat inactivation for 20 min at 65 °C and

then mixed with the pUF1-Cas9 plasmid. Transfected parasites were grown in 0.5 μM

anhydrous tetracycline (aTc) (Cayman Chemical). Drug pressure was applied 48 h after

transfection, using blasticidin (BSD) (Gibco) at a concentration of 2.5 μg/mL, selecting

for pKD-RON11-Apt expression. After parasites grew back from transfection, integration

was confirmed by PCR, and then cloned using limiting dilution. Clones were maintained

in mediums containing 0.5 μM aTc and 2.5 μg/mL BSD.

Growth assays

For all assays, aliquots of parasite cultures were incubated in 8 μM Hoechst 33342

(ThermoFisher Scientific) for 20 min at room temperature and then fluorescence was

measured using a CytoFlex S (Beckman Coulter) flow-cytometer. Flow cytometry data

were analyzed using FlowJo software (Tree Star, Inc.) and plotted using Prism

(GraphPad Software Inc.).

For the growth assay, synchronous ring-stage parasites were washed five times with

RPMI 1640 medium and split into two cultures, one resuspended in medium containing

0.5 μM aTc and 2.5 μg/mL BSD (+RON11), and the other one in medium containing

only 2.5 μg/mL BSD (-RON11). Cultures were then transferred to a 96-well plate at

0.2% parasitemia and grown for 6 days. Parasitemia was monitored every 48 h.

For the invasion vs egress assay, synchronous ring-stage parasites were washed and

split into two cultures with and without aTc, as previously described. Parasites were

then allowed to develop into mature schizonts to be isolated using a Percoll gradient

(Genesee Scientific). Enriched parasites were then incubated for 4 hours at 37 °C in

pre-warmed RPMI media supplemented with the PKG inhibitor, ML10 compound (25

nM) (obtained from S. Osborne, BEI resources) (Ressurreição et al. 2020). After

incubation, parasites were washed twice with pre-warmed RPMI media and transferred

immediately to pre-warmed RBCs at 2% hematocrit. Parasitemia was monitored every

two hours, for eight hours.

For the rescue assay, parasites were synchronized to a 1-hour window. Synchronized

ring-stage parasites were washed and split into six cultures in a 6-well plate (+ RON11,

- RON11, 44h, 46h, 48h and 50h). Culture "+RON11" was grown in the presence of aTc,
while the others were grown in absence of aTc. Parasites were allowed to develop into
schizonts and aTc was supplemented back into cultures at their corresponding time
point, except for the "-RON11" culture. Parasitemia was monitored at 44 and 56 hours
post infection (hpi).

Western blotting

Parasites were synchronized, washed, and split into two cultures with and without aTc,
as previously described. Synchronous parasites were then allowed to develop into
mature schizonts to be then incubated with E64 for 4 hours at 37 °C. After treatment,
culture pellets were treated with ice-cold 0.04% saponin in 1x PBS to isolate parasites
from host cells. The parasite pellets were subsequently solubilized in protein loading
dye with Beta-mercaptoethanol (LI-COR Biosciences) and used for SDS-PAGE.

Primary antibodies used in this study included mouse-anti-HA (6E2; Cell Signaling
Technology; 1:2000), rabbit-anti-PfEF1α (from Daniel Goldberg, 1:2000), mouse-anti-
RAP1 (2.29; from Jana McBride via the European Malaria Reagent Repository; 1:500)
(Hall et al. 1983), mouse-anti-RON4 (10H11; from Alan Cowman; 1:500) (Richard et al.
2010), mouse-anti-MSP1 (12.4; from Jana McBride via the European Malaria Reagent
Repository; 1:500) (McBride, Newbold, and Anand 1985). Secondary antibodies used
were IRDye 680 CW goat-anti-rabbit IgG and IRDye 800CW goat-anti-mouse IgG (Li-
COR Biosciences; 1:20 000). Membranes were imaged using the Odyssey Clx Li-COR
infrared imaging system (Li-COR Biosciences). Images were processed and analyzed
using ImageStudio (Li-COR Biosciences).

Microscopy and image analysis

For all microscopy assays, parasites were synchronized, washed, and split into two
cultures with and without aTc, as previously described.

For IFAs, cells were fixed following the previously described protocol (Cobb et al. 2017).

For localization assays, schizont-stage parasites were smeared onto a slide, fixed and
permeabilized with acetone for 10 min at room temperature.

For rhoptry secretion assays, synchronous parasites were allowed to develop into
mature schizonts to be isolated using a Percoll gradient (Genesee Scientific). Enriched
parasites were then incubated for 4 hours at 37 °C in pre-warmed RPMI media
supplemented with 25 nM ML10 compound. After incubation, parasites were washed
twice with pre-warmed RPMI media and transferred immediately to RBCs at 1%
hematocrit. Cultures were then allowed to egress for 30 min at 37 °C in media
supplemented with 1 µM cytochalasin D (Invitrogen). After incubation, parasites were
fixed with 4% paraformaldehyde (PFA) (Electron Microscopy Sciences) and 0.03%
glutaraldehyde, before being permeabilized with 0.1% Triton-X100.

Primary antibodies used in the IFAs included rabbit-anti-HA (SG77; ThermoFisher
Scientific; 1:100), mouse-anti-RAP1 (2.29; from Jana McBride via the European Malaria
Reagent Repository; 1:500) (Hall et al. 1983), mouse-anti-RON4 (10H11; from Alan
Cowman; 1:200) (Richard et al. 2010). Secondary antibodies used were Alexa Fluor
488 and Alexa Fluor 546 (Life Technologies, 1:1000).

After mounting the cells using ProLong Diamond with 4',6'-diamidino-2-phenylindole
(DAPI) (Invitrogen), they were imaged using a DeltaVision II microscope system with an
Olympus Ix-71 inverted microscope. Images were collected as a Z-stack and
deconvolved using SoftWorx (GE HealthCare), then displayed as a maximum intensity
projection. Adjustments to brightness and contrast were made for display purposes
using Adobe Photoshop. Colocalization was analyzed using the Pearson's correlation
coefficient calculated by the Cell Profiler software (Broad Institute Inc.).

For growth assays, synchronous schizont-stage parasites were washed five times and
split into two cultures, one without and one with aTc. Parasites were allowed to develop
during two life cycles and sample aliquots were collected at different time points.

For attachment assays, parasites were collected as previously described in the rhoptry
secretion assay with a slight change. After ML10 removal, schizont-stage parasites

were allowed to egress during 2 h at 37 °C in the presence and absence of 1 μM
cytochalasin D. Samples were collected after incubation.

For light microscopy imaging, aliquots were smeared into glass slides and stained using
Hema3 Fixative solutions (Fischer Healthcare). Slides were imaged using a Zeiss Axio
Scope A1 microscope with a Zeiss Axiocam 305 color camera. Images were processed
using Adobe Photoshop.

Ultrastructural expansion microscopy (U-ExM)

Cultures for U-ExM were synchronized to mature schizont parasites, following the
previously described methods. Ultrastructure expansion microscopy (U-ExM) was
performed as described previously (Liffner and Absalon 2021), with minor modifications.

To start, 12 mm round coverslips were treated with poly-D-lysine for 1 hour at 37 °C.
They were then washed three times with MilliQ water and placed in a 24-well plate.
Parasite cultures with approximately 5% parasitemia were adjusted to 0.5% hematocrit.
Then, 1 mL of parasite culture was added to the well containing the treated coverslip
and incubated for 1 h at 37 °C.

After the incubation, the supernatant was carefully removed, and a fixative solution (4%
v/v PFA in PBS) was added, followed by a 20 min incubation at 37 °C. The coverslips
were washed three times with 1X PBS and incubated overnight at 37 °C in 500 μL of
1.4% formaldehyde/2% acrylamide (FA/AA) in PBS.

The monomer solution (19% sodium acrylate, 10% acrylamide, 0.1% N,N'-
methylenebisacrylamide in PBS) was prepared a day prior and stored at -20 °C. Before
gelation, coverslips were removed from FA/AA solution and washed three times in 1X
PBS.

For gelation, 5 μL of 10% tetramethylenediamine (TEMED) and 5 μL of 10% ammonium
persulfate (APS) were added to 90 μL of the monomer solution, briefly vortexed, and 35
μL of the monomer mixture were pipetted onto parafilm. The coverslips were placed on
top with the cell-side facing down, and the gels were incubated at 37 °C for 30 min.

Next, the gels were transferred into a 6-well plate containing denaturing buffer (200 mM
sodium dodecyl sulfate (SDS), 200 mM NaCl, 50 mM Tris, pH 9) and incubated for 15
min incubation at room temperature. Afterward, the gels were separated from the
coverslips and transferred to 1.5 mL tubes with the denaturing buffer for 90-min
incubation at 95 °C.

Subsequently, the gels were incubated with secondary antibodies diluted in 1X PBS for
2.5 hours. After denaturation, gels were transferred to Petri dishes containing 25 mL of
MilliQ water and incubated three times for 30 min at room temperature with shaking,
changing the water in between. The gels were measured and subsequently shrunk
using two washes with 1X PBS. They were then transferred to a 24-well plate for
blocking in 3% BSA in PBS at room temperature for 30 min. After blocking gels were
incubated with primary antibodies diluted in 3% BSA overnight at room temperature.

Following primary antibody incubation, the gels were washed three times in 0.5% PBS-
Tween 20 for 10 min before incubation with secondary antibodies diluted in 1X PBS for
2.5 hours.

After secondary antibody staining, the gels were washed three times with 0.5% PBS-
Tween 20. Then, gels were transferred back to 10 cm Petri dishes for the second round
of expansion, involving three incubations with MilliQ water. After re-expansion, the gels
were either imaged immediately or stored in 0.2% propyl gallate in water until imaging.

The primary antibodies used were rat-anti-HA 3F10 (Roche, 1:25), mouse-anti-RAP1
(2.29; from Jana McBride via the European Malaria Reagent Repository; 1:500) (Hall et
al. 1983), mouse-anti-RON4 (10H11; from Alan Cowman; 1:200) (Richard et al. 2010).
The secondary antibodies used were Alexa Fluor 488 and Alexa Fluor 546 (Life
Technologies, 1:500), NHS-ester 405 (Thermofisher, 1:250). The gels were imaged
using a Zeiss LSM 980 microscope with Airyscan 2. Images were collected as a Z-
stack, processed by Airyscan, and then displayed as a maximum intensity projection.
Adjustments to brightness and contrast were made using ZEN Blue software for display
purposes.

Expression and purification of recombinant RON11

A *E.coli* codon optimized DNA fragment corresponding to amino acids 32-442 of RON11 was synthesized (Genscript) and cloned into pMAL vector (NEB) in frame with an N-terminal maltose binding protein (MBP) tag and c-terminal His6 tag. Protein expression was induced using 0.4mM IPTG at 16°C for 12-14h. The cell pellet was resuspended in a lysis buffer (300mM NaCl, 10mM imidazole, 500mM NaH_2PO_4 and 1µM PMSF, 10mM BME, and 0.1% Tween20). Following sonication, the soluble fraction containing recombinant MBP-RON11-His6 protein was affinity purified twice on immobilized-metal affinity chromatography (IMAC) column using FPLC (Biorad) and further purified on CHT Ceramic Hydroxyapatite column. Fractions containing rMBP-RON11 were pooled and dialyzed into phosphate buffered saline (PBS) containing 1µM PMSF, 10mM BME and 0.1% Tween20 and kept at -80°C.

Animal immunization, IgG purification and neutralization assay

Animal experiments were approved by the Johns Hopkins Animal Care and Use Committee (ACUC, protocol# RA22H291). Four female Sprague Dawley rats (Charles River Laboratory), 5-6 weeks old, were used for immunizations. Recombinant MBP-RON11 (25µg/rat/immunization) was mixed 1:1 with Addavax adjuvant and injected subcutaneously three times in two-week intervals. Two weeks following the last immunization blood was collected by terminal heart bleed. Total rat IgG was purified from serum using pre-equilibrated protein G column (GE health sciences), dialyzed against RPMI 1640 medium, and sterile filtered.

Synchronized trophozoite stage 3D7 strain parasites were adjusted to 0.5% parasitemia in 4% hematocrit in a 96-well plate and incubated with anti-RON11 IgG at the indicated concentrations with or without pre-incubation with 3µM rMBP protein. After 72 hours incubation at 37°C, cells were stained with SYBR Green I and parasitemia was measured using AttuneNxT flow cytometer (ThermoScientific).

ACKNOWLEDGEMENTS

We thank Dan Goldberg for anti-EF1α, Alan Cowman for anti-RON4, The European Malaria Reagent Repository for anti-RAP1 antibodies; Julie Nelson and Juan Bustamante at the CTEGD Cytometry Shared Resource Laboratory for help with flow cytometry and analysis; and Muthugapatti Kandasamy at the Biomedical Microscopy Core at the University of Georgia for help with microscopy. The study was funded by NIH/NIAID R56 AI73133 (V.M.), T32 AI060546 (G.S.V), T32 AI138953 (S.Y.), Office of the Vice-President for Research at UGA (D.A.), the CTEGD TIPS Fellowship and Daniel G. Colley Training in Parasitology Fund (D.A.), Johns Hopkins Malaria Research Institute predoctoral fellowship (O.A-F), and in part by the Johns Hopkins Malaria Research Institute and the Bloomberg Philanthropies (P.S.).

REFERENCES

Anaguano, David, Watcharatip Dedkhad, Carrie F. Brooks, David W. Cobb, and Vasant Muralidharan. 2023. "Time-Resolved Proximity Biotinylation Implicates a Porin Protein in Export of Transmembrane Malaria Parasite Effectors." *Journal of Cell Science*, September. https://doi.org/10.1242/jcs.260506.

Bannister, L. H., J. M. Hopkins, R. E. Fowler, S. Krishna, and G. H. Mitchell. 2000. "Ultrastructure of Rhoptry Development in Plasmodium Falciparum Erythrocytic Schizonts." *Parasitology* 121 (Pt 3) (September): 273–87.

Bantuchai, Sirasate, Mamoru Nozaki, Amporn Thongkukiatkul, Natcha Lorsuwannarat, Mayumi Tachibana, Minami Baba, Kazuhiro Matsuoka, Takafumi Tsuboi, Motomi Torii, and Tomoko Ishino. 2019. "Rhoptry Neck Protein 11 Has Crucial Roles during Malaria Parasite Sporozoite Invasion of Salivary Glands and Hepatocytes." *International Journal for Parasitology* 49 (9): 725–35.

Beeson, James G., Damien R. Drew, Michelle J. Boyle, Gaoqian Feng, Freya J. I. Fowkes, and Jack S. Richards. 2016. "Merozoite Surface Proteins in Red Blood Cell

Invasion, Immunity and Vaccines against Malaria." *FEMS Microbiology Reviews* 40 (3):

343–72.

Blake, Thomas C. A., Silvia Haase, and Jake Baum. 2020. "Actomyosin Forces and the

Energetics of Red Blood Cell Invasion by the Malaria Parasite Plasmodium Falciparum."

PLoS Pathogens 16 (10): e1009007.

Boyle, Michelle J., Danny W. Wilson, Jack S. Richards, David T. Riglar, Kevin K. A.

Tetteh, David J. Conway, Stuart A. Ralph, Jake Baum, and James G. Beeson. 2010.

"Isolation of Viable *Plasmodium Falciparum* Merozoites to Define Erythrocyte Invasion

Events and Advance Vaccine and Drug Development." *Proceedings of the National*

Academy of Sciences of the United States of America 107 (32): 14378–83.

Bozdech, Zbynek, Manuel Llinás, Brian Lee Pulliam, Edith D. Wong, Jingchun Zhu, and

Joseph L. DeRisi. 2003. "The Transcriptome of the Intraerythrocytic Developmental

Cycle of Plasmodium Falciparum." *PLoS Biology* 1 (1): E5.

Counihan, Natalie A., Scott A. Chisholm, Hayley E. Bullen, Anubhav Srivastava, Paul R.

Sanders, Thorey K. Jonsdottir, Greta E. Weiss, et al. 2017. "Plasmodium Falciparum

Parasites Deploy RhopH2 into the Host Erythrocyte to Obtain Nutrients, Grow and

Replicate." *ELife* 6 (March). https://doi.org/10.7554/eLife.23217.

Counihan, Natalie A., Ming Kalanon, Ross L. Coppel, and Tania F. de Koning-Ward.

2013. "Plasmodium Rhoptry Proteins: Why Order Is Important." *Trends in Parasitology*

29 (5): 228–36.

Cova, Marta Mendonça, Mauld H. Lamarque, and Maryse Lebrun. 2022. "How

Apicomplexa Parasites Secrete and Build Their Invasion Machinery." *Annual Review of*

Microbiology 76 (1): 619–40.

Cowman, Alan F., Christopher J. Tonkin, Wai-Hong Tham, and Manoj T. Duraisingh.

2017. "The Molecular Basis of Erythrocyte Invasion by Malaria Parasites." *Cell Host &*

Microbe 22 (2): 232–45.

Dvorin, Jeffrey D., and Daniel E. Goldberg. 2022. "Plasmodium Egress Across the
Parasite Life Cycle." *Annual Review of Microbiology* 76 (September): 67–90.

Favuzza, Paola, Manuel de Lera Ruiz, Jennifer K. Thompson, Tony Triglia, Anna Ngo,
Ryan W. J. Steel, Marissa Vavrek, et al. 2020. "Dual Plasmepsin-Targeting Antimalarial
Agents Disrupt Multiple Stages of the Malaria Parasite Life Cycle." *Cell Host & Microbe*
27 (4): 642-658.e12.

Florentin, Anat, David W. Cobb, Jillian D. Fishburn, Michael J. Cipriano, Paul S. Kim,
Manuel A. Fierro, Boris Striepen, and Vasant Muralidharan. 2017. "PfClpC Is an
Essential Clp Chaperone Required for Plastid Integrity and Clp Protease Stability in
Plasmodium Falciparum." *Cell Reports* 21 (7): 1746–56.

Florentin, Anat, Dylon R. Stephens, Carrie F. Brooks, Rodrigo P. Baptista, and Vasant
Muralidharan. 2020. "Plastid Biogenesis in Malaria Parasites Requires the Interactions
and Catalytic Activity of the Clp Proteolytic System." *Proceedings of the National*
Academy of Sciences of the United States of America 117 (24): 13719–29.

Frénal, Karine, Jean-François Dubremetz, Maryse Lebrun, and Dominique Soldati-
Favre. 2017. "Gliding Motility Powers Invasion and Egress in Apicomplexa." *Nature*
Reviews. Microbiology 15 (11): 645–60.

Ganesan, Suresh M., Joanne M. Morrisey, Hangjun Ke, Heather J. Painter, Kamal
Laroiya, Margaret A. Phillips, Pradipsinh K. Rathod, Michael W. Mather, and Akhil B.
Vaidya. 2011. "Yeast Dihydroorotate Dehydrogenase as a New Selectable Marker for
Plasmodium Falciparum Transfection." *Molecular and Biochemical Parasitology* 177 (1):
29–34.

Geoghegan, Niall D., Cindy Evelyn, Lachlan W. Whitehead, Michal Pasternak, Phoebe
McDonald, Tony Triglia, Danushka S. Marapana, et al. 2021. "4D Analysis of Malaria
Parasite Invasion Offers Insights into Erythrocyte Membrane Remodeling and
Parasitophorous Vacuole Formation." *Nature Communications* 12 (1): 1–16.

Ghosh, Sreejoyee, Kit Kennedy, Paul Sanders, Kathryn Matthews, Stuart A. Ralph,
Natalie A. Counihan, and Tania F. de Koning-Ward. 2017. "The Plasmodium Rhoptry

Associated Protein Complex Is Important for Parasitophorous Vacuole Membrane
Structure and Intraerythrocytic Parasite Growth." *Cellular Microbiology* 19 (8).
https://doi.org/10.1111/cmi.12733.

González-Sanz, Marta, Pedro Berzosa, and Francesca F. Norman. 2023. "Updates on
Malaria Epidemiology and Prevention Strategies." *Current Infectious Disease Reports*,
June, 1–9.

Groomes, Patrice V., Usheer Kanjee, and Manoj T. Duraisingh. 2022. "RBC Membrane
Biomechanics and Plasmodium Falciparum Invasion: Probing beyond Ligand-Receptor
Interactions." *Trends in Parasitology* 38 (4): 302–15.

Hallée, Stéphanie, Natalie A. Counihan, Kathryn Matthews, Tania F. de Koning-Ward,
and Dave Richard. 2018. "The Malaria Parasite Plasmodium Falciparum Sortilin Is
Essential for Merozoite Formation and Apical Complex Biogenesis." *Cellular
Microbiology* 20 (8): e12844.

Hanssen, Eric, Chaitali Dekiwadia, David T. Riglar, Melanie Rug, Leandro Lemgruber,
Alan F. Cowman, Marek Cyrklaff, et al. 2013. "Electron Tomography of Plasmodium
Falciparum Merozoites Reveals Core Cellular Events That Underpin Erythrocyte
Invasion." *Cellular Microbiology* 15 (9): 1457–72.

Hart, Melissa N., Franziska Mohring, Sophia M. DonVito, James A. Thomas, Nicole
Muller-Sienerth, Gavin J. Wright, Ellen Knuepfer, Helen R. Saibil, and Robert W. Moon.
2023. "Sequential Roles for Red Blood Cell Binding Proteins Enable Phased
Commitment to Invasion for Malaria Parasites." *Nature Communications* 14 (1): 4619.

Ito, Daisuke, Marc A. Schureck, and Sanjay A. Desai. 2017. "An Essential Dual-
Function Complex Mediates Erythrocyte Invasion and Channel-Mediated Nutrient
Uptake in Malaria Parasites." *ELife* 6 (February). https://doi.org/10.7554/eLife.23485.

Krai, Priscilla, Seema Dalal, and Michael Klemba. 2014. "Evidence for a Golgi-to-
Endosome Protein Sorting Pathway in Plasmodium Falciparum." *PloS One* 9 (2):
e89771.

Kudyba, Heather M., David W. Cobb, Anat Florentin, Michelle Krakowiak, and Vasant
Muralidharan. 2018. "CRISPR/Cas9 Gene Editing to Make Conditional Mutants of
Human Malaria Parasite P. Falciparum." *Journal of Visualized Experiments: JoVE*, no.
139 (September). https://doi.org/10.3791/57747.

Lasonder, Edwin, Chris J. Janse, Geert-Jan van Gemert, Gunnar R. Mair, Adriaan M.
W. Vermunt, Bruno G. Douradinha, Vera van Noort, et al. 2008. "Proteomic Profiling of
Plasmodium Sporozoite Maturation Identifies New Proteins Essential for Parasite
Development and Infectivity." *PLoS Pathogens* 4 (10): e1000195.

Liffner, Benjamin, and Sabrina Absalon. 2021. "Expansion Microscopy Reveals
Plasmodium Falciparum Blood-Stage Parasites Undergo Anaphase with A Chromatin
Bridge in the Absence of Mini-Chromosome Maintenance Complex Binding Protein."
Microorganisms 9 (11): 2306.

Liffner, Benjamin and Absalon, Sabrina. 2023. "Expansion Microscopy of Apicomplexan
Parasites." *Molecular Microbiology*, August. https://doi.org/10.1111/mmi.15135.

Liffner, Benjamin, Juan Miguel Balbin, Gerald J. Shami, Ghizal Siddiqui, Jan Strauss,
Sonja Frölich, Gary K. Heinemann, et al. 2022. "Cell Biological Analysis Reveals an
Essential Role for Pfcerli2 in Erythrocyte Invasion by Malaria Parasites."
Communications Biology 5 (1): 1–17.

Liffner, Benjamin, Juan Miguel Balbin, Jan Stephan Wichers, Tim-Wolf Gilberger, and
Danny W. Wilson. 2021. "The Ins and Outs of Plasmodium Rhoptries, Focusing on the
Cytosolic Side." *Trends in Parasitology* 37 (7): 638–50.

Liffner, Benjamin, Ana Karla Cepeda Diaz, James Blauwkamp, David Anaguano, Sonja
Frölich, Vasant Muralidharan, Danny W. Wilson, Jeffrey Dvorin, and Sabrina Absalon.
2023. "Atlas of Plasmodium Falciparum Intraerythrocytic Development Using Expansion
Microscopy." https://doi.org/10.7554/elife.88088.1.

Liffner, Benjamin, Ana Karla Cepeda Diaz, James Blauwkamp, David Anaguano, Sonja
Frolich, Vasant Muralidharan, Danny W. Wilson, Jeffrey D. Dvorin, and Sabrina

Absalon. 2023. "Atlas of Plasmodium Falciparum Intraerythrocytic Development Using
Expansion Microscopy." *ELife* 12 (December). https://doi.org/10.7554/eLife.88088.

Llinás, Manuel, Zbynek Bozdech, Edith D. Wong, Alex T. Adai, and Joseph L. DeRisi.
2006. "Comparative Whole Genome Transcriptome Analysis of Three Plasmodium
Falciparum Strains." *Nucleic Acids Research* 34 (4): 1166–73.

Mageswaran, Shrawan Kumar, Amandine Guérin, Liam M. Theveny, William David
Chen, Matthew Martinez, Maryse Lebrun, Boris Striepen, and Yi-Wei Chang. 2021. "In
Situ Ultrastructures of Two Evolutionarily Distant Apicomplexan Rhoptry Secretion
Systems." *Nature Communications* 12 (1): 4983.

Mahajan, Babita, Angamuthu Selvapandiyan, Noel J. Gerald, Victoria Majam, Hong
Zheng, Thilan Wickramarachchi, Jawahar Tiwari, et al. 2008. "Centrins, Cell Cycle
Regulation Proteins in Human Malaria Parasite Plasmodium Falciparum." *The Journal
of Biological Chemistry* 283 (46): 31871–83.

Martinez, Matthew, William David Chen, Marta Mendonça Cova, Petra Molnár, Shrawan
Kumar Mageswaran, Amandine Guérin, Audrey R. Odom John, Maryse Lebrun, and Yi-
Wei Chang. 2022. "Rhoptry Secretion System Structure and Priming in Plasmodium
Falciparum Revealed Using in Situ Cryo-Electron Tomography." *Nature Microbiology*.
https://doi.org/10.1038/s41564-022-01171-3.

Nasamu, Armiyaw S., Svetlana Glushakova, Ilaria Russo, Barbara Vaupel, Anna
Oksman, Arthur S. Kim, Daved H. Fremont, et al. 2017. "Plasmepsins IX and X Are
Essential and Druggable Mediators of Malaria Parasite Egress and Invasion." *Science*
358 (6362): 518–22.

Ooij, Christiaan van, Pamela Tamez, Souvik Bhattacharjee, N. Luisa Hiller, Travis
Harrison, Konstantinos Liolios, Taco Kooij, et al. 2008. "The Malaria Secretome: From
Algorithms to Essential Function in Blood Stage Infection." *PLoS Pathogens* 4 (6):
e1000084.

Perrin, Abigail J., Christine R. Collins, Matthew R. G. Russell, Lucy M. Collinson, David
A. Baker, and Michael J. Blackman. 2018. "The Actinomyosin Motor Drives Malaria

Parasite Red Blood Cell Invasion but Not Egress." *MBio* 9 (4).
https://doi.org/10.1128/mBio.00905-18.

Pino, Paco, Reto Caldelari, Budhaditya Mukherjee, Juha Vahokoski, Natacha Klages,
Bohumil Maco, Christine R. Collins, et al. 2017. "A Multistage Antimalarial Targets the
Plasmepsins IX and X Essential for Invasion and Egress." *Science* 358 (6362): 522–28.

Rajaram, Krithika, Hans B. Liu, and Sean T. Prigge. 2020. "Redesigned TetR-Aptamer
System To Control Gene Expression in Plasmodium Falciparum." *MSphere* 5 (4).
https://doi.org/10.1128/mSphere.00457-20.

Ressurreição, Margarida, James A. Thomas, Stephanie D. Nofal, Christian Flueck,
Robert W. Moon, David A. Baker, and Christiaan van Ooij. 2020. "Use of a Highly
Specific Kinase Inhibitor for Rapid, Simple and Precise Synchronization of Plasmodium
Falciparum and Plasmodium Knowlesi Asexual Blood-Stage Parasites." *PloS One* 15
(7): e0235798.

Riglar, David T., Dave Richard, Danny W. Wilson, Michelle J. Boyle, Chaitali Dekiwadia,
Lynne Turnbull, Fiona Angrisano, et al. 2011. "Super-Resolution Dissection of
Coordinated Events during Malaria Parasite Invasion of the Human Erythrocyte." *Cell*
Host & Microbe 9 (1): 9–20.

Rudlaff, Rachel M., Stephan Kraemer, Jeffrey Marshman, and Jeffrey D. Dvorin. 2020.
"Three-Dimensional Ultrastructure of Plasmodium Falciparum throughout Cytokinesis."
PLoS Pathogens 16 (6): e1008587.

Schureck, Marc A., Joseph E. Darling, Alan Merk, Jinfeng Shao, Geervani Daggupati,
Prakash Srinivasan, Paul Dominic B. Olinares, et al. 2021. "Malaria Parasites Use a
Soluble RhopH Complex for Erythrocyte Invasion and an Integral Form for Nutrient
Uptake." *ELife* 10 (January): e65282.

Segev-Zarko, Li-Av, Peter D. Dahlberg, Stella Y. Sun, Daniël M. Pelt, Chi Yong Kim,
Elizabeth S. Egan, James A. Sethian, Wah Chiu, and John C. Boothroyd. 2022. "Cryo-
Electron Tomography with Mixed-Scale Dense Neural Networks Reveals Key Steps in
Deployment of Toxoplasma Invasion Machinery." *PNAS Nexus* 1 (4): gac183.

Sherling, Emma S., Abigail J. Perrin, Ellen Knuepfer, Matthew R. G. Russell, Lucy M. Collinson, Louis H. Miller, and Michael J. Blackman. 2019. "The Plasmodium Falciparum Rhoptry Bulb Protein RAMA Plays an Essential Role in Rhoptry Neck Morphogenesis and Host Red Blood Cell Invasion." *PLoS Pathogens* 15 (9): e1008049.

Srinivasan, Prakash, Wandy L. Beatty, Ababacar Diouf, Raul Herrera, Xavier Ambroggio, J. Kathleen Moch, Jessica S. Tyler, et al. 2011. "Binding of Plasmodium Merozoite Proteins RON2 and AMA1 Triggers Commitment to Invasion." *Proceedings of the National Academy of Sciences of the United States of America* 108 (32): 13275–80.

Voß, Yannik, Severina Klaus, Julien Guizetti, and Markus Ganter. 2023. "Plasmodium Schizogony, a Chronology of the Parasite's Cell Cycle in the Blood Stage." *PLoS Pathogens* 19 (3): e1011157.

Vulliez-Le Normand, Brigitte, Michelle L. Tonkin, Mauld H. Lamarque, Susann Langer, Sylviane Hoos, Magali Roques, Frederick A. Saul, et al. 2012. "Structural and Functional Insights into the Malaria Parasite Moving Junction Complex." *PLoS Pathogens* 8 (6): e1002755.

Wang, Kevin, Eric D. Peng, Amy S. Huang, Dong Xia, Sarah J. Vermont, Gaelle Lentini, Maryse Lebrun, Jonathan M. Wastling, and Peter J. Bradley. 2016. "Identification of Novel O-Linked Glycosylated Toxoplasma Proteins by Vicia Villosa Lectin Chromatography." *PloS One* 11 (3): e0150561.

Weiss, Greta E., Paul R. Gilson, Tana Taechalertpaisarn, Wai-Hong Tham, Nienke W. M. de Jong, Katherine L. Harvey, Freya J. I. Fowkes, et al. 2015. "Revealing the Sequence and Resulting Cellular Morphology of Receptor-Ligand Interactions during Plasmodium Falciparum Invasion of Erythrocytes." *PLoS Pathogens* 11 (2): e1004670.

World Health Organization. 2023. "World Malaria Report 2023." 2023. https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2023.

Yahata, Kazuhide, Melissa N. Hart, Heledd Davies, Masahito Asada, Samuel C. Wassmer, Thomas J. Templeton, Moritz Treeck, Robert W. Moon, and Osamu Kaneko.

2021. "Gliding Motility of *Plasmodium* Merozoites." *Proceedings of the National*
Academy of Sciences of the United States of America 118 (48).
https://doi.org/10.1073/pnas.2114442118.

Yeh, Ellen, and Joseph L. DeRisi. 2011. "Chemical Rescue of Malaria Parasites Lacking
an Apicoplast Defines Organelle Function in Blood-Stage Plasmodium Falciparum."
PLoS Biology 9 (8): e1001138.

1036

1037 **Tables**

1038 **Table 3.1.** List of primers used in the study to generate the cell lines RON11[apt].

1039

Amplicon	Primer	Sequence (5' – 3')
RON11-Cterm integration	P1	TGTTTAGGAATTTTTGGAGGTATTTGATAG
Apt integration	P2	CTAGACTAGGTTCCAAGATCTCCC
RON11-Cterm	P3	ATTGTGTATCCCGATATCGATGTTAATAGTTTAAAAAAAATACTAACAGAAGTAAGATG
	P4	CGTCATAAGGGTATCCGGAGACGTCTTTCGAATCAGATGCTGACTGTAAATAAG
RON11-3'UTR	P5	CCAATGGCCCCTTTCCGGGCGCGCCCTTTTTTGTTTGAAATTGGCAAAAATATTAGTAG
	P6	ACTATTAACATCGATATCGGGATACACAATAGAATATTAATTAAGTGTATTATTAAGTG
RON11 gRNA	P7	CATATTAAGTATATAATATTGATATGGATGTCGAAAGAAAGTTTTAGAGCTAGAAATAGC

1040

1041

1042 **Supplemental Figures**

1043

1044

Supplemental figure 1. RON11 localizes to the parasite periphery during early-ring stages. IFAs
showing the localization of RON11 in RON11^{apt} rings with respect to the rhoptry markers RON4 and
RAP1. Synchronous parasites were fixed with PFA and stained with specific antibodies. Images from left
to right are phase-contrast, DAPI (nucleus, cyan), anti-HA (RON11, magenta), anti-RON4 or RAP1
(yellow), and fluorescence merge. Z stack images were deconvolved and projected as a combined single
image. Representative images of 2 biological replicates.

1051

1052

1053

Supplemental figure 2. Expression and purification of recombinant RON11, and validation of anti-
RON11 antibody. (A) Expression of rMBP-RON11 (aa 32-442) protein in *E.coli*. Soluble (S) and insoluble
(I) fractions from cell lysates without and without IPTG induction. M; Marker. **(B)** Western blot analysis of
the soluble fraction using anti-His antibody and detected using anti-mouse HRP conjugated secondary
antibody. **(C)** rMBP-RON11 following IMAC purification and subsequent clean-up using CHT column. **(D)**
Western blot analysis of P.falciparum schizont lysates using anti-RON11 antibody. **(E)**
Immunofluorescence microscopy of mature schizont stained with DAPI, anti-RON11 (N-term), anti-HA (c-
term) and anti-RAP1 antibodies.

1062

1063

1064

1065

1066

1067

1068

Supplemental figure 3. RON11[apt] merozoites attached to RBCs in the presence or absence of aTc.
(A) Duration between first contact of merozoites with RBCs and observation of RBC deformation. (n = 3 biological replicates, 20 merozoites for +RON11 and 13 for -RON11; error bars = SEM; *p < 0.05 by unpaired two-tailed t-test). **(B)** Duration between merozoite first merozoite-induced RBC deformation and echinocytosis. (n = 3 biological replicates, 20 merozoites for +RON11 and 13 for -RON11; error bars = SEM; ns = non-significant by unpaired two-tailed t-test). **(C)** Quantification of RBCs showing echinocytosis after merozoite-induced deformation. (n = 3 biological replicates, 34 merozoites for +RON11 and 44 for -RON11). **(D)** Deformation scores based on the strength of merozoite-RBC interaction. 0 = no deformation; 1 = shallow indentation/membrane pinching; 2 = deeper indentation to the side of RBC/intermediate level of host cell membrane pinching around parasite. (n = 3 biological replicates, 44 merozoites for each condition; *p < 0.05 by chi-squared test). **(E)** Schematic of the actin-polymerization inhibitor assay. Schizonts were tightly synchronized incubating for 4 hours with the PKG inhibitor, Compound 1. After incubation, schizonts were washed twice and then transferred to fresh red blood cells in the presence of the actin inhibitor, cytochalasin D, for 30 minutes. Created with BioRender.com. **(F and G)** Quantification of RON11[apt] merozoites attached to RBCs in the presence or absence of aTc, after incubation with **(F)** or without **(G)** cytochalasin D. Parasites were smeared 30 minutes after Compound 1 removal,, stained with Hema 3, and scored by light microscopy. Attached-merozoites were blindly scored and represented as the percentage of events per 100 RBCs (n = 5 and 3 biological replicates respectively; error bars = SEM; ns = non-significant; **p < 0.01 by unpaired two-tailed t-test).

1091

Supplemental figure 4. RON11 knockdown can generate multiple rhoptries but does not affect the number of merozoites. (A) Representative images of RON11[apt] parasite showing multiple rhoptries accumulating within fully developed schizonts in the absence of aTc. ML10-arrested parasites were expanded by U-ExM, fixed with PFA, and stained with NHS-Ester (greyscale), anti-HA (magenta), and the DNA dye SYTOX (cyan). Selected Z stack images were projected as a combined single image. Number on image = Z-axis thickness of projection in μm. **(B and C)** Quantification of **(B)** rhoptries per merozoites and **(C)** merozoites per schizont in the presence and absence of aTc. (n = 4 biological replicates, 28 schizonts for +RON11 and 36 for -RON11; error bars = SEM; ns = non-significant; ***p < 0.001 by unpaired two-tailed *t*-test)

1101

1102

Supplemental figure 5. RON11 knockdown does not affect the localization of other rhoptry proteins in single rhoptry merozoites. Representative images of RON11[apt] free merozoites showing the localization of RON11 with respect to RON4 and RAP1 in the presence and absence of aTc. Late-schizonts were expanded by U-ExM, fixed with PFA, and stained with NHS-Ester (greyscale), anti-HA (magenta), and anti-RON4 or RAP1 (yellow). Selected Z stack images were projected as a combined single image. Number on image = Z-axis thickness of projection in μm.

1109

Supplemental figure 6. RON11 knockdown does not impact the expression and secretion of MSP1
and AMA1. (A) Representative IFAs of RON11[apt] schizonts showing secretion of MSP1 proteins in the
presence or absence of aTc. E64-arrested schizonts were fixed with PFA and stained with specific
antibodies. Images from left to right are phase-contrast, DAPI (nucleus, blue), anti-HA (RON11, green),
anti-MSP1 (red), and fluorescence merges. Z stack images were deconvolved and projected as a
combined single image. Representative images of 3 biological replicates. **(B)** (Top) Western blot of
parasite lysates isolated from E64-arrested RON11[apt] parasites in the presence and absence of aTc.
Samples were probed with antibodies against MSP1 and EF1α (loading control). The protein marker sizes
are shown on the left. Representative blot of 4 biological replicates shown. (Bottom) Quantification of the
relative level of MSP1 total protein content in E64-arrested RON11[apt] parasites in the presence and
absence of aTc. Band intensities were normalized to the loading control, EF1α, and are presented as
normalized arbitrary units (A.U.) (n = 4 biological replicates; error bars = SD; ns = non-significant by
unpaired two-tailed t-test). **(C)** Representative IFAs of RON11[apt] schizonts showing AMA1 localization in
the presence or absence of aTc. E64-arrested schizonts were fixed with PFA and stained with specific
antibodies. Images from left to right are phase-contrast, DAPI (nucleus, blue), anti-HA (RON11, green),
anti-AMA1 (red), and fluorescence merges. Z stack images were deconvolved and projected as a
combined single image. Representative images of 3 biological replicates.