Evidence for interacting but decoupled controls of decisions and movements in non-human primates

Clara SALERI[1,2] & David THURA[1,2]

[1] University Claude Bernard Lyon 1, F-69000, France
[2] INSERM U1028, CNRS UMR5292, Lyon Neuroscience Research Center, Impact Team, Lyon, F-69000, France

Running head:

Integrated control of decisions and movements in monkeys

Corresponding author information:

David Thura

Lyon Neuroscience Research Center – Impact team
INSERM U1028 – CNRS UMR5292 – University Claude Bernard Lyon 1
16 avenue du Doyen Jean Lépine, 69675 Bron, France

E-mail : david.thura@inserm.fr

Manuscript information:

Abstract: 244 words; Main text: ~8000 words
26 pages – 4 main figures - 5 supplemental figures

Acknowledgements/Funding:

The authors have no financial or non-financial interests to declare. The authors wish to thank Sonia Alouche and Jean-Louis Borach for their administrative assistance, Gislène Gardechaux and Amélie Reynaud for helping to train the monkeys and collect data, Frédéric Volland for his expertise during the technical preparation and maintenance of this experiment, and Julien Dessenne for creating the illustration shown in Figure 1A. This work is supported by a CNRS/Inserm ATIP/Avenir grant to DT.

Abstract

Many recent studies indicate that control of decisions and actions is integrated during interactive behavior. Among these, several carried out in humans and monkeys conclude that there is a co-regulation of choices and movements. Another perspective, based on human data only, proposes a decoupled control of decision duration and movement speed, allowing for instance to trade decision duration for movement duration when time pressure increases. Crucially, it is not currently known whether this ability to flexibly dissociate decision duration from movement speed is specific to humans, whether it can vary depending on the context in which a task is performed, and whether it is stable over time. These are important questions to address, especially to rely on monkey electrophysiology to infer the neural mechanisms of decision-action coordination in humans. To do so, we trained two macaque monkeys in a perceptual decision-making task and analyzed data collected over multiple behavioral sessions. Our findings reveal a strong and complex relationship between decision duration and movement vigor. Decision duration and action duration can co-vary but also "compensate" each other. Such integrated but decoupled control of decisions and actions aligns with recent studies in humans, validating the monkey model in electrophysiology as a means of inferring neural mechanisms in humans. Crucially, we demonstrate for the first time that this control can evolve with experience, in an adapted manner. Together, the present findings contribute to deepening our understanding of the integrated control of decisions and actions during interactive behavior.

New & noteworthy

The mechanism by which the integrated control of decisions and actions occurs, coupled or interactive but decoupled, is debated. In the present study, we show in monkeys that decisions and actions influence each other in a decoupled way. For the first time, we also demonstrate that this control can evolve depending the subject's experience, allowing to trade movement time for decision time and limit the temporal discounting of reward value.

Keywords

Decision-making ; Arm movement ; Coordination; Non-human primates ; Reward value

Introduction

Recent theories and studies argue for an integrated control of decisions and actions during interactive behavior, at least when decisions are rapidly expressed through movements (1–4). According to this framework, decisions, or factors guiding choices such as reward, time and effort, influence action parameters (5–11), and conversely, actions influence decisions (12–15), even if the choice is about a sensory stimulus or reward harvest during foraging (16–21).

These observations are not compatible with a strict serial organization of behavior, where cognition, including decision-making is separate from perception and action, and lie between the two (22). Instead, they support a dynamic control system operating in a closed loop where movements shape current and future potential actions (i.e. affordances), assigning various importance to the stimuli given such opportunities, and thus directly influencing decisions (23–27).

While the integrated control of decisions and actions is an increasingly established hypothesis, a lively ongoing debate concerns the mechanism by which this integrated control occurs. On the one hand, proposals have suggested that decisions and actions are jointly regulated, possibly by a common source (6, 9, 28, 29). Such "co-regulation" hypothesis, demonstrated in both humans and monkeys, is compatible with the older observations that reward and effort invigorate and slow down both decisions and actions, respectively (8, 30–34). It has been proposed that a regulation signal computed in the basal ganglia allows to maximize the rate of correct responses by jointly adjusting decisions urgency and action vigor during tasks involving multiple successive choices between actions (6, 35–37).

A second line of research suggests however that decision duration and movement speed are controlled independently of each other, allowing for instance to trade decision time for movement time (19, 20, 38, 39). In support of this possibility, we recently showed that human subjects can shorten their decisions when an accurate and time-consuming movement is requirement (19, 20, 38). We proposed that such adjustment is a "compensatory" mechanism established to decrease the temporal discounting of a positive outcome in each trial, and/or to limit a drop of success rate on a more global time scale. This ability to exchange decision duration for movement duration, i.e. the flexibility to dissociate decision urgency from movement vigor, argues for interacting but decoupled decision and motor control systems.

Importantly, it is not currently known whether this ability to flexibly dissociate decision duration from movement speed is specific to humans, maybe made possible by a

sophistication of decision and motor neural circuits through evolution, or whether non-human
primates like old-world monkeys, who diverged from the human lineage about 25 MA ago,
already have this skill. Likewise, it is currently not known if the regulation mechanism of
decisions and actions is exclusive and stable over time or whether it can vary depending on
the context in which a task is performed, and whether or not it can evolve with practice.

These are crucial questions to address because a significant part of what we know about the
neural substrates of decision-making and motor control comes from (non-human) animal and,
more specifically, monkey neurophysiology (40, 41). It is thus important to figure out whether
or not the mechanism underlying the integrated control of decision and movement (i.e. joint
or compensatory) is similar in both species, stable across contexts (for example, by
prioritizing either speed or precision of behavior) and over time, first at the behavioral level in
order to then rely on monkey electrophysiology to infer the neural mechanisms in humans.

In the present study, we trained two macaque monkeys to perform a decision between actions
task and analyzed decision and movement data collected over dozens of sessions. We focused
our analyses on the relationship between decision duration and movement vigor (speed and/or
duration), both at the single trial level and between blocks of trials in which motor constraints
were varied. We also investigated these relationships as a function of animals' experience in
the task. The analyses strongly indicate that decisions and actions influence each other, but in
a decoupled way: decision duration and action vigor could co-vary but also "compensate"
each other. Crucially, this integrated control can evolve with practice in an adapted manner.
Our data indeed suggest that monkeys can trade movement time for decision time, and vice
versa, when the total duration of behavior increases, possibly allowing to limit a drop of
reward value due to temporal discounting.

Methods

Subjects and ethics statement

The present report describes data collected from two rhesus macaque monkeys (Macaca
mulatta - monkey G, female, 8 kg, 10 years old, right-handed; monkey B, male, 5 kg, 4 years
old, left-handed). Ethical permission was provided by "Comité d'Éthique Lyonnais pour les
Neurosciences Expérimentales" (CELYNE), C2EA #42, ref: C2EA42-11-11-0402-004.
Monkey housing and care was in accordance with European Community Council Directive
(2010) and the Weatherall report, "The use of non-human primates in research." Laboratory

authorization was provided by the "Préfet de la Région Rhône-Alpes" and the "Directeur

départemental de la protection des populations" under Approval Number: D69 029 06 01.

Experimental apparatus

The monkeys sat on a primate chair and made planar reaching movements using a lever held

in their dominant hand (Fig. 1A). A digitizing tablet (GTCO CalComp) continuously recorded

the lever horizontal and vertical positions (~100 Hz with 0.013-cm accuracy). Target stimuli

and cursor feedback were projected by a VIEWPixx monitor (VPixx Technologies, 120 Hz

refresh rate) onto a half-silvered mirror suspended 25 cm above and parallel to the digitizer

plane, creating the illusion that targets floated on the plane of the tablet.

Behavioral tasks

The monkeys performed the same task, a modified version of the "tokens task" (see 42 for the

original version), as that performed by human subjects and for which data were reported

recently (19) (Fig. 1B). They were faced with a visual display consisting of three blue circles

(radius: 2.25 cm for monkey G; 2 cm for monkey B) placed horizontally at a distance of 6.375

cm (monkey G) or 7 cm (monkey B) from each other (the "decision" circles). In the central

blue circle, 15 small tokens were randomly arranged. Positioned 5 cm (monkey G) or 5.5 cm

(monkey B) below, three black circles, organized horizontally as well, defined the

"movement" targets. The central black circle, the starting circle, remained constant at a radius

of 1.5 cm, while the lateral black circles, positioned at a distance of 8 cm (monkey G) or 7 cm

(monkey B) from the center, varied in size (radius: 1.25 -1.75 cm; and 2.25 cm) in separate

blocks of trials, resulting in two motor conditions ("small" vs. "large" targets).

A trial started when the animal moved and held the cursor in the black central circle for

500ms. At this point, the tokens began to jump one by one, every 200ms, from the central

blue circle into one of the two possible lateral blue circles. The monkeys had to determine

which of the two lateral blue circles would contain the majority of tokens before they had all

jumped. They reported their decisions by moving the cursor into the corresponding lateral

movement target (Fig. 1B). Importantly, monkeys could make and report their choices at any

time between the first and last token jumps. However, the movement duration could not

exceed 800ms, regardless of the motor condition.

Figure 1 - Experimental setup, task and data analysis. *A* - Experimental apparatus. *B* - Time course of a trial in the task. *C* - Temporal profile of success probability in one example trial of the choice task. At the beginning of the trial, each target has the same success probability (0.5). When the first token jumps into one of the two potential targets (the most leftward vertical dotted line), success probability of that target increases to ~0.6. Success probability then evolves with every jump. Subjects execute a reaching movement (red trace) to report their choices. Movement onset and offset times are used to compute movement duration. Peak velocity is determined as the maximum value between these two events. Movement offset marks the moment when the tokens that remain in the central decision circle jump more quickly to their assigned target (gray trace). The estimated time of the decision is computed by subtracting the subject's mean non-decision delay estimated in a simple delayed reaching task from movement onset time, allowing computation of the choice success probability at that moment. Only 10 out of 15 jumps are illustrated on this success probability profile.

The feedback about timing provided to the monkey was determined online: LabVIEW logged timestamps when the monkey left the start position and when it entered the chosen target, then compared this duration to the maximum allowed duration (800 ms) to ensure the movement was not too long. This differs from the offline analysis, which uses velocity thresholds to determine the onset and offset of movements (see below). If the movement exceeded 800ms

(too slow) or reached the target but failed to stop within 800ms (inaccurate), the trial was

considered as a movement error and the movement target turned orange. When the choice was

properly reported, the remaining tokens jumped more quickly to their final circles (Fig. 1C).

This post-decision interval was set to 100-150ms and this acceleration of the remaining

tokens implicitly encouraged animals to decide before all tokens had jumped into their

respective lateral circles, aiming to save time and increase their reward rate. Note that this

feature entails that movement duration carries a temporal cost with respect to the monkey's

rate of correct decisions because the remaining tokens accelerate at movement offset. The

visual feedback regarding trial success (both correct decision and correct movement) or

failure (correct movement but wrong decision) was provided after the last token jump, with

the chosen decision circle turning green or red, respectively. A reward (drops of fruit juice)

was given to the monkey after each correct trial performed. Using a pressure dispenser (Crist

Instrument) with 35 ms opening duration and 2.0 PSI pressure, 100 rewards equated to 15 ml

of juice, corresponding to approximately 0.15 ml per drop. Therefore, monkeys received

between 2 and 3 drops of juice per correct trial, depending on their motivation, amounting to

approximately 0.30 ml to 0.45 ml of juice. Finally, the next trial began after a 1500ms inter-

trial interval.

To ensure that the difficulty of decisions was homogeneous among experimental conditions,

especially between the two motor conditions, we controlled the sequence of trials experienced

by animals in each session. We interspersed among fully random trials (30% of the trials in

which each token is 50% likely to jump into the right or the left lateral circle) three special

types of trials characterized by particular patterns of token jumps. Between 30 and 40% of

trials were so-called "easy" trials, in which tokens tended to move consistently toward one of

the circles. Around 20% of trials were "ambiguous," in which the initial token movements

were balanced, making the success probability of each target close to 0.5 until later in the

trial. The last special trial type was called "misleading" trials (7-20%) in which the two to

three first tokens jumped into the incorrect circle and the remaining ones into the correct

circle (please refer to Fig. S1A for a complete description of trial types criteria). In all cases,

even when the temporal profile of token jumps of a trial was predesigned, the actual correct

target was randomly selected on each trial.

In addition to the tokens task, each animal had to complete trials in a delayed reaching task in

each session. The delayed reaching task was similar to the tokens task, but with only one

lateral decision circle displayed at the beginning of each trial (on the right or left side of the

central circle with a 50% probability). The tokens all moved at the same time (go signal) from

the central circle to this unique lateral circle after a variable delay (400-550ms). In this task

the reward was given to the monkey if the movement is initiated after the go signal and is

accurately executed (endpoint within the target and duration under 800 ms). The target turns

orange if the movement is inaccurate or too long. Animals performed this task in the two

different motor conditions (small and large targets). The delayed reaching task was employed

to estimate the non-decision delay (i.e. the sum of the delays attributable to sensory

processing of the stimulus display) as well as to response initiation in each motor condition

(Fig. 1C).

Procedure

Sessions included in the present report where those during which monkeys were proficient in

the task, typically following several months of training (see the stability of monkeys'

performance in Fig. 2A). Data included in the analyses were collected over a total of 71

sessions for monkey G and 46 for monkey B. Each session consisted of alternating blocks

(between 1 and 5) of 10-60 correct trials in each motor condition whose order alternated on a

daily basis. Importantly, the mean success probability profiles of trials in each motor

condition were similar (Fig. S1B), ensuring that a potential difference in decision strategy is

not due to a difference in the overall difficulty of the trials encountered by each monkey.

Monkeys had also to complete between 45 and 60 correct trials of the delayed reaching task in

each condition. Then, each session had a predefined goal for the number of correct trials to

perform in the main task (often 180 or 240 for monkey G and 240 for monkey B). Depending

on the motivation of the animal on a given day, the goal could be adjusted, being either

increased or decreased (ranging from 60-240 trials for monkey G and 180-400 for monkey B).

Sessions were stopped if cooperation was poor. On average, monkey G performed 174 ± 62

correct trials (both correct decisions and correct movements) by session while monkey B

made 253 ± 43 trials.

Data analysis

All arm movement data were analyzed off-line using MATLAB (MathWorks). Reaching

characteristics were assessed using the animals' movement kinematics. Horizontal and

vertical position data were first filtered using a tenth-degree polynomial filter and then

differentiated to obtain a velocity profile. Onset and offset of movements were determined

using a 3.75 cm/s velocity threshold. Peak velocity was determined as the maximum value

between these two events. To estimate the time at which monkeys committed to their choice

(decision timeon each trial, we detected the time of movement onset, defining the animal's

reaction time (RT), and subtracted from it her/his mean non decision delays estimated based

on her/his reaction times in the same motor condition (large or small targets) of the delayed

reaching task performed the same day. Decision duration was computed as the duration

between the decision time and the first token jump (Fig. 1C). The task design allows to

calculate, at each moment in time during a trial, the success probability $pi(t)$ associated with

choosing each target i (Eq. 1). For instance, for a total of 15 tokens, if at a particular moment

in time the right target contains N_R tokens, whereas the left target contains N_L tokens, and

there are N_C tokens remaining in the center, then the probability that the target on the right

will ultimately be the correct one, i.e., the success probability of guessing right is as follows:

$$p(\text{R}|N_R, N_L, N_C) = \frac{N_C!}{2^{N_C}} \sum_{k=0}^{\min(N_C, 7-N_L)} \frac{1}{k!\,(N_C - k)!} \tag{1}$$

253   The rate of correct trials was computed for each session as the number of correct trials to

254   complete in that given session divided by the sum of the durations of all trials performed to

255   reach this requested number of correct trials.

Statistics

Wilcoxon rank sum tests are used to assess the difference in decision duration between

categories of trials (e.g. easy versus ambiguous decisions, decisions made in early versus late

sessions, etc.). Pearson linear correlations are computed to assess the relationship between

decision duration and movement parameters (speed and duration) at the single trial level and

between sessions. Pearson linear correlations are also computed the assess the relationship

between the degree of coordination between decision duration and movement speed and the

duration of decisions between sessions. For all statistical tests, the significance level is set a

0.05.

Results

General observations

Over the 71 sessions performed by monkey G, $63 \pm 6\%$ of the trials (mean $\pm$ standard

deviation between the sessions) were successful (both correct choices and correct

movements). Without taking into account the success of the movements, $74 \pm 4\%$ of her

decisions were correct. Movement errors, too slow or inaccurate, were relatively sparse,

representing $7 \pm 5\%$ and $8 \pm 6\%$ of the total trials, respectively. Monkey B's performance

over the 46 sessions he completed was comparable to that of monkey G. 62 ± 7% of his trials

were successful (correct decisions and correct movements). Regardless of movement

outcome, 73 ± 5% of his choices were correct. 4 ± 4% of his movements were too slow, 10 ±

4% were inaccurate. For both monkeys, performance was stable across the sessions included

in the present report (Fig. 2A).

Figure 2 - Global performance, decisional and movement vigor in the task. *A* - Performance of each monkey
across sessions. *B* - Decision duration, reaching movement duration and speed for each of the two monkeys.
Bars illustrate mean values across sessions, and dots illustrate individual values for each session. *C* – Success
rate of each monkey across sessions.

Although their performance in the task is comparable, the analysis of animals' duration of

decisions, duration and speed of arm movements suggests that they used different strategies to

achieve these similar outcomes (Fig. 2B). On average, the duration of monkey G's decision

was 1026 ± 242ms. As expected, she decided more quickly when trials were easy compared to

when they were ambiguous (872 ± 232ms vs. 1154 ± 259ms, Wilcoxon rank sum test, z = -

5.7, p<0.0001). To report her decisions, monkey G made arm movements whose duration was

527 ± 21ms and their speed was 23 ± 2 cm/s. Monkey B was on average much faster to decide

and act compared to monkey G. His decision duration was 249 ± 43ms, with faster choices

when trials were easy compared to when they were ambiguous (230 ± 40ms vs. 369 ± 73ms,

Wilcoxon rank sum test, z = -6.6, p<0.0001). To report his decisions, monkey B made

movements whose speed was 32 ± 3 cm/s and their duration was351 ± 14ms. As a

consequence, the rate of correct trials, defined for each session as the number of correct trials

per minute (Fig. 2C), was much higher in monkey B than in monkey G (9.7 ± 1 correct

trials/min vs. 5.8 ± 0.7 correct trials/min).

Relationship between decision duration and movement vigor at the single trial level

To study the relationship between decision duration and arm movement vigor (speed and

duration) *within trials*, we first grouped all trials (correct and error trials, regardless of motor

condition) across sessions, separately for each monkey, according to decision time computed

in bins of 200ms, and calculated the mean (± standard error) values of movement parameters

in each bin.

The analysis of monkey G movement speed as a function of decision duration (including

decision durations until 2200ms, which corresponds to 99.6% of the trials) shows a slight but

continuous *increase* of movement speed with decision duration (from 22.5 to 23.9 cm/s, Fig.

3A, left panel). A linear regression through the data indicates a very strong and significant

relationship between the two variables (Pearson correlation, r=0.98, p<0.0001), with a

positive slope (0.14) and an intercept at 22.4 cm/s. No significant relationship between

decision duration and movement duration is observed (Pearson correlation, r=0.02, p=0.9).

Figure 3 - Relation between decision duration and arm movement speed within trials. *A – Left panel*: Speed
of monkey G's arm movements (reaching) as a function of decision duration, computed in bins of 200ms,
averaged (± standard error, shaded areas) across all trials (black), trials of the 20 first sessions ("early" sessions,
blue) and trials of the 20 last sessions ("late" sessions, orange). Histograms at the bottom of the panel show the
distributions of the duration of monkey G's decisions in the two groups, "early" and "late" sessions. *Center panel*:
Pearson linear correlation between the duration of monkey G's decisions and arm movement speed (velocity
peak) as a function of the session performed by the monkey. Each dot illustrates the value for a given session.
The blue and orange shaded areas highlight the 20 first and the 20 last sessions performed by monkey G,
respectively. A negative (positive) r value means that the longest the decision, the slowest (fastest) the movement.

By contrast, the analysis of the relationship between the duration of decisions and the speed

of monkey B's movement within trials (including decision durations until 2200ms, which

corresponds to 99.9% of the trials) shows a continuous *decrease* of movement speed with

decision duration (from 32.2 to 25 cm/s, Fig. 3B, left panel). A linear regression through the

data indicates a significant relationship between the two variables (Pearson correlation, r=-

0.85, p=0.0008), with a negative slope (-0.62) and an intercept at 34 cm/s. As for monkey G,

no significant relationship between monkey B decision duration and movement duration is

observed (Pearson correlation, r=-0.05, p=0.9).

We then asked whether this relationship between decision duration and movement speed

within trials was stable over extended periods of time, depending on the monkeys' experience

in the task, or if it could possibly evolve depending on this experience. To do this, we first

carried out the same analyzes as those described above but with trials grouped according to

the number of sessions performed by each monkey. For both animals, we compared trials

collected during their first 20 sessions ("early" sessions) with those performed in their last 20

sessions ("late" sessions).

The relationship between the duration of decisions and the speed of monkey G's movements

is strong and significant in both groups of trials ("early" sessions: r=-0.9, p=0.001; "late"

sessions: r=0.82, p=0.0021), but the slope is *negative* for "early" sessions (-0.22) and *positive*

for "late" sessions (0.19). This means that the longest decisions were followed by the slowest

movements during monkey G's first sessions and then, with practice, this relationship

reversed (Fig. 3A, left panel; Fig. S2A). To better capture this evolution, we analyzed the

session-by-session relationship between decision duration and action speed in monkey G and

we found that a linear regression shows a strong and significant fit (Pearson correlation,

r=0.61, p<0.0001), with a positive slope, meaning that the link between decision duration and

movement speed continuously evolved through monkey G's practice (Fig. 3A, middle panel).

In particular, among the first 20 sessions, decision durations were significantly correlated with

movement speed in 8 sessions, always with long decisions followed by slow movements.

Among the last 20 sessions, we found a significant correlation between decision duration and

movement speed in 8 sessions too, but this time always with long decisions followed by rapid

movements. Finally, we analyzed the duration of monkey G's decision in the two groups of

trials and noticed that with practice, she became more conservative and decided more slowly

(mean ± standard error; "early" sessions: 760 ± 6ms vs. "late" sessions: 1101 ± 7ms,

Wilcoxon rank sum test, z = -35, p<0.0001) (Fig. 3A, left panel). Interestingly, we found a

positive and significant correlation between the degree of coordination between decision

duration and movement speed and the duration of decisions between sessions (Pearson

correlation, r=0.6, p<0.001) (Fig. 3A, right panels). In other words, the emergence of the

positive correlation between decision duration and movement speed (the longest the decision,

the fastest the movement) within trials through practice coincides with the increase of the

duration of monkey G's decisions through practice. In the supplementary section of the

present report, we describe the relationship between the evolution of the correlation of

decision duration and movement speed and the evolution of decision duration in two other

monkeys who performed a similar version of the task (Fig. S3), strongly supporting the same

conclusion (Fig. S4).

As for monkey G, the relationship between the duration of decisions and the speed of monkey

B's movements is significant in both groups of trials ("early" sessions: r=-0.78, p=0.0044;

"late" sessions: r=-0.9, p=0.0002). But contrary to monkey G, the slope is negative in both

groups: -0.65 for "early" and -0.64 for "late" sessions (Fig. 3B, left panel; Fig. S2B). This

means that regardless of the monkey's level of experience in the task, the longest decisions

were followed by the slowest movements. We did not find a particular link between the

relationship between decision duration and action speed and the sessions performed by

monkey B through his practice (Fig. 3B, middle panel). Among the first 20 sessions, decision

durations were significantly correlated with movement speed in 7 sessions, and always with

long decisions followed by slow movements. Among the last 20 sessions, the correlation

between decision duration and movement speed was significant in 3 sessions, still with long

decisions followed by rapid movements. Interestingly though, despite the fact that the

relationship between decision duration and movement speed *within trials* did not change with

practice, we found that movement speed was overall higher in the "late" sessions group

(intercept = 35 cm/s) compared to the "early" sessions one (intercept = 32.9 cm/s) (Fig. 3B,

left panel). The analysis of the duration of monkey B's decision in the two groups showed that

he was slightly more conservative in the "late" sessions compared to the "early" sessions (302

± 3ms vs. 259 ± 3ms, Wilcoxon rank sum test, z = -17, p<0.0001) (Fig. 3B, left panel).

Contrary to monkey G, no significant relationship between the degree of coordination
between decision duration and movement speed and the duration of monkey B's decisions
between sessions is observed (Fig. 3B, right panels).

Effect of motor constraints on the relationship between decision and action

Another way to study the coordination between decision duration and movement vigor is to
encourage animals to vary the vigor of their movements in blocks of trials and assess the
consequence of this adjustment on decision duration. A co-regulatory mechanism predicts that
slowing movements will be accompanied by slowing decisions, while a more flexible
mechanism could be demonstrated by observing shortened decisions when movements are
longer, as we have seen in humans (19, 20, 38).

Across all sessions, when movements were correctly executed (decisions could be correct or
wrong), the speed of monkey G's movements was slower (mean speed ± standard error: 22.8
± 0.05 cm/s vs. 23.3 ± 0.05 cm/s, Wilcoxon rank sum test, z = -6, p<0.0001) and their
duration longer (536 ± 1ms vs. 525 ± 1ms, z = 8.4, p<0.0001) when they were executed
toward the small targets ("small" target block) compared to when they aimed to the large
targets ("large" target block). We found that the duration of monkey G's decisions was overall
slower when movements were directed toward small targets compared to when they reached
the large targets (1020 ± 6ms vs. 953 ± 6ms, z = 8.6, p<0.0001). At the session level, we
found a significant effect of the motor condition on the speed and duration of monkey G's
movements in 16 and 19 out of 71 sessions, respectively (Wilcoxon rank sum test, p<0.05),
with movements being faster and shorter when directed to the large targets in 15/16 and 18/19
cases, respectively (Fig. 4A, left and middle panels). The duration of monkey G's decisions
was modulated by the motor condition in 34 out of 71 sessions (Wilcoxon rank sum test,
p<0.05), with decisions being slower when movements were slower in the majority of cases
(27/34) (Fig. 4A, right panel). Interestingly, we found that this tendency to slow down
decisions when a time consuming movement was necessary diminished with practice. Indeed,
while the effect of the motor condition on movement speed and duration was stable over
monkey G's practice (Fig. 4A, left and middle panels), the number of sessions during which
long decisions preceded long movements decreased (20 over the first 35 sessions; 5 for the
last 35 sessions) (Fig. 4A, right panel).

Figure 4 - Effect of motor constraints on arm movement vigor and decision duration. *A – Left panel*: Difference of arm movement duration between trials of the small target blocks and those of the large target blocks, as a function of the session performed by monkey G. Each dot illustrates data of a given session, and the black dots mark sessions in which the difference is statistically significant (the number of statistically significant cases is reported for each difference direction). A positive value indicates that movements were longer when they were executed toward the small targets compared to when they were executed toward the large targets. The red line illustrates the result of a linear regression through the data. *Center panel*: Same as in A for the difference of the speed of monkey G's arm movement between the two motor conditions. *Right panel*: Same as A for the difference of the duration of monkey G's decisions between the two motor conditions. *B* – Same as in A for trials performed by monkey B.

The effect of target size on the speed and duration of monkey B's movement is less clear, at

least when trials are grouped across sessions. Movement speed was not significantly

modulated as a function of the motor condition (small vs. large target: 32 ± 0.07 cm/s vs. 32.2

± 0.07 cm/s, Wilcoxon rank sum test, z = -0.2, p=0.8). Movement duration was slightly longer

when movement reached the large targets compared to when they reached a small target (349

± 0.7ms vs. 351 ± 0.7ms; z = -2.3, p=0.02). At the session level, the effect of target size on the

kinematics of monkey B's movements (duration and speed) was mixed, as can be seen in

figure 4B, left and middle panels. However, target size significantly modulated decision

duration in 20 out of 46 sessions (Wilcoxon rank sum test, p<0.05). In the majority of cases

(16/20), decision duration was longer when movements reached the large targets (Fig. 4B,

right panel). Interestingly, with practice, this monkey progressively decreased his movement

speed in one condition ("large" target block) compared to the other one ("small" target block)

(Fig. 4B, middle panel), and this relative decrease of speed was accompanied by a session-by-

session lengthening of decisions in the large target blocks compared to the small target blocks

(Fig. 4B, right panel). The link between these two adjustments from one session to the next is

significant (Pearson correlation, r=-0.3, p=0.03).

Discussion

In the present report, we show in non-human primates that the duration of perceptual

decisions and the speed of the arm movements executed to report these choices are often

correlated at the single-trial level. The results support an interacting but decoupled control of

the decision-making and movement execution processes: decision duration and action speed

are sometimes co-regulated (the slower the decision, the slower the movement), and they

sometimes "compensate" each other (the slower the decision, the faster the movement). We

also demonstrate that the degree and direction of coordination between decision duration and

movement vigor can evolve, particularly as a function of the overall duration of the animals'

decisions in each session: the "compensatory" mode of regulation comes into play as duration

of choices increases. Finally, we show that when the motor context encourages time-

consuming movements, decision duration and movement vigor are often co-regulated. This

context-dependent relationship also appears flexible depending on the animals' experience in

the task.

These observations agree with many recent experimental results that challenge the classic

view of behavior organization (44), in which perception, decision, and action are considered

as independent, temporally separate and serial processes (22). More and more studies indeed

show that decisions and actions are closely interconnected and influence each other, at least

during rapid interactive behavior (5, 7–21, 28, 29, 38, 45).

A particularly interesting question concerns the mechanism by which such integrated control

of decisions and actions occurs. On the one hand, decisions and actions might be coupled,

jointly regulated by a common source (6, 9, 28, 29). According to this hypothesis, in a context

encouraging decision speed, the fastest choices will be reported with the fastest movements.

Similarly, if a decision needs to be reported with a slow and careful movement, this decision

will tend to be longer than that expressed by a rapid movement. A second line of research

suggests however that the control of decision duration and movement speed is decoupled,

allowing for instance to trade decision time for movement time if necessary (19, 20, 38, 39,

46, 47). At the origin of this second hypothesis, we showed that human subjects can shorten

their decisions when an accurate and time-consuming movement is required (19, 20, 38). We

proposed that such adjustment is a compensatory mode possibly established to prevent the

temporal discounting of a positive outcome in each trial, and/or to limit a drop of success rate

on a more extended time scale. Interestingly, in two recent studies specifically designed to

address the mode of regulation of decisions and actions (28, 29), and whose results support a

co-regulation mechanism, decision and movement durations had no impact on the

participants' success rate (i.e. it was not adaptive in terms of success rate to speed up

movements after a long decision for instance), unlike the design of the experiments described

in Reynaud et al. (19) and Saleri Lunazzi et al. (20, 38).

The data presented in this report suggest for the first time that both modes of coordination can

be used by the same subject. At the single trial level, the slowest decisions made by monkey B

were expressed with the slowest movements (Fig. 3B, left panel), a result consistent with a

co-regulation of decisions and actions. Monkey G also co-regulated but mostly during the

first sessions she performed. With experience, the relationship between the duration of her

decisions and the speed of her movements reversed, consistent with a decoupled,

"compensatory" mode of regulation of choices and actions (Fig. 3A, left panel). With trials

grouped as a function of animals' experience, we found in both monkeys that decisions were

slower in the late sessions compared to the early sessions, and these slower decisions were

reported with faster movements (Fig. 3A-B, left panels), a pattern of results also consistent

with a "compensatory" mode of regulation of decisions and actions. Finally, with trials

grouped according to the motor context in which decisions were made (i.e. small vs. large

targets), we found evidence for a co-regulation of decisions and actions, with slow

movements often preceded by slow decisions, with a trend for an evolution toward a

"compensatory" mode in monkey G (Fig. 4A). The way in which the duration of the decision

and the vigor of the action are coordinated thus appears complex, flexible and dynamic

depending on the context in which the task is carried out (especially the level of experience in

the task). In the following paragraph, we discuss the possible reasons which could explain the

use of these different modes of coordination.

It is well-established that individuals tend to exhibit a preference for policies that maximize

their rate of correct responses when faced with multiple successive decisions (48, 49). Several

factors influence this success rate, with time being a crucial one, as evidenced by the well-

known concept of temporal discounting of reward value (50). During movement execution for

instance, action duration and speed are two (strongly related, Fig. S5) parameters that partially

describe a single latent variable, the vigor, that reflects movement's utility and that the brain

seeks to optimize as a function of the economic context (speed-accuracy trade-off, experience

in the task, etc.) in which a behavior occurs (4, 34, 51, 52). Because both decisions and

actions are time consuming processes, minimizing the time required to obtain a reward likely

leads to adaptations concerning both decisions and movements. A decoupled yet

interconnected control of decisions and actions provides individuals with the ability to adapt

both deliberation and movement speed together, minimizing the time it takes to achieve a

given goal. Moreover, this mode also possesses the flexibility to selectively adjust one aspect

while preserving the integrity of the other. For instance, during easy decisions, deliberation

time can be slightly sacrificed when accurate and slow response movements are required (19,

20, 38). The present results indicate that just like humans, monkeys also have the ability to

flexibly adjust the way they coordinate their decisions with their actions, in particular

according to their experience in the task.

The evolution of the coordination between decision duration and action vigor depending on

animals' experience is particularly useful in revealing the principles underlying how these two

processes are controlled. Crucially, the emergence of the compensatory mode of coordination

always coincides with an increase of the duration of the animal's overall behavior (Fig. 3A,

right panel; Fig. S4). This is particularly salient in monkey G who showed an important

increase in her decision durations over the course of the sessions carried out (Fig. 3A, right

panel). Interestingly, the subtle and gradual decoupling of decision and action observed across

sessions between large and small target conditions (Fig. 4A, right panel) occurs

simultaneously with the overall lengthening of her decision durations across sessions (Fig.

3A, top right panel). She may have wanted to seek to compensate for her long movements by

limiting the increase in the duration of her decisions in the most constrained motor blocks (i.e.

small target). By contrast, monkey B, who was constantly very fast to decide and act, showed

almost no sign of compensation between decision and movement durations. In monkey S and

monkey Z who were even slower to decide than monkey G, the coordination within trials

between decision duration and action speed is only of the compensatory type, but it still

depends on the duration of the animals' overall decisions (Fig. S4). We thus propose that the

decoupled, compensatory mode of regulation is highly adaptive in the sense that it allows to

deal with deadlines and/or makes it possible to limit the temporal discounting of the reward

value in each trial. In the longer term, it could also help optimize the overall success rate of

the animals in each session. Between these three possibilities, we lean toward the limitation of

the temporal discounting of the reward value in each trial because (1) monkeys almost never

failed their trials because of responding after 3s (i.e. the deadline of deliberation) and (2) the

overall success rate of monkey G does not appear to improve despite the emergence of the

compensatory mode of regulation across sessions (Fig. 2C).

The different strategies adopted by the two monkeys can be explained by their very different

personalities and backgrounds. Monkey B who faced his first experimental task is much

younger than monkey G. He adopted such a fast and impulsive behavior that there was almost

no temporal discounting of reward in each (correct) trial, and no room for either decision or

movement duration shortening. Consequently, a decoupled mode of regulation that would

compensate for slightly longer decisions or movements to meet some reward rate

maximization principles was probably irrelevant to this monkey. Monkey G however, who is

older and more experienced in laboratory experiments, exhibited a more conservative and less

vigorous behavior. Therefore, she had to face higher temporal discounting of reward in each

trial and lower reward rate at a more global scale (Fig. 2C). Compensation between decision

and movement duration becomes beneficial in this case.

Human subjects having performed the same task show a compensatory mode of coordination

of decision and action when the trials are grouped according to the motor constraints imposed

on the subjects (19). By contrast, with the same comparison (decision durations between

small versus large target conditions), we report here that monkeys tend to primarily exhibit a

co-regulation of decisions and actions. This difference between monkeys and human subjects

can be explained by at least two factors. Firstly, monkeys received drops of juice at each

correct trial as a reward, so their strategy was probably primarily driven by the prospect of

receiving this strong primary reinforcer. By contrast, humans did not receive instant rewards.

Their motivation was likely more oriented toward the completion of the required number of

correct trials as quickly as possible, aiming to maximize their overall success rate at the

sessions level. Another, not mutually exclusive explanation is that taking the motor context

into account to compensate for movement duration during the decision-making process

requires a significant level of abstraction that monkeys can only gradually acquire with

training and experience, to eventually exploit it in an adapted manner. Interestingly, we report

that with practice, the slowdown of monkey G's decisions reported with slow movements in

specific blocks of trials tends to decrease (Fig. 4A). By mastering the timing parameters of the

task over the sessions, she may have realized that beyond accuracy at the single trial level, she

could also optimize her time spent in a session by slightly reducing the duration of her

decisions when movements were slow, just as human subjects did, with the difference that
they were able to do it after just one session.

In previous experiments in which humans and monkeys were tested in a task similar to that
described in the present report, we proposed that both decision urgency and movement vigor
(6, 9) were co-regulated. This conclusion was based on two observations. At the single trial
level first, early decisions (usually made on the basis of strong sensory evidence but low
urgency) were followed by long duration movements whereas later decisions (relying on
weak sensory evidence but stronger urgency) were followed by shorter and faster movements,
as if they were influenced by the strong level of urgency encountered when a significant
amount of time has elapsed during a trial. Second, when subjects were encouraged to make
earlier and less accurate decisions in blocks of trials (fast speed-accuracy trade-off regime),
we observed faster movements compared to a condition where accuracy was emphasized. The
multiple modes of regulation of decisions and actions that we report in the present study
suggest however that the mechanism underlying the control of decision duration and
movement speed is more elaborated than being under the influence of a single regulation
signal. Indeed, while a shared signal may control decision and movement durations at the
single trial level (especially when fast movements follow long decisions), and between
different decisional speed-accuracy trade-off regimes, a unique signal cannot account for the
fact that the movements executed following the slower decisions made during the late
sessions were faster compared to those made during the early sessions during which decision
were faster. Together, these observations suggest the involvement of multiple sources of
regulation of decisions and actions, likely implemented by diverse brain circuits, and used
differently and specifically depending on the context in which a task takes place (level of
experience in the task, speed-accuracy trade-off, etc.).

It is important to note that the present results are all correlational. A next very interesting
question concerns the relationship between the average duration of behavior and the decision-
action coordination mode used by the subjects. In this report we propose, on the basis of
correlational data, that this mode is determined by the duration of decisions but this must be
tested directly, if possible in a way that tends toward the establishment of a causal link
between the two phenomena. A possibility of intervention concerns electrical
microstimulation in the monkey cerebral cortex, especially in areas involved in the
determination of choice. For example, we have shown in the past that electrical stimulation in
the dorsal part of the premotor cortex and, to a lesser extent, in the primary motor cortex,

could lengthen decisions if this stimulation was delivered shortly before choice commitment occurs (53). By experimentally prolonging the duration of decisions, it would then be possible to observe movements executed rapidly in order to compensate for this additional decisional duration.

Electrophysiological data have indeed highlighted an overlap of brain regions involved in decision-making and those of action processes, particularly in the sensorimotor cortex and the basal ganglia (1, 40, 54). Studies suggest, for example, that activity in the internal section of the globus pallidus, the output nucleus of the basal ganglia, determines decision urgency (35, 36) as well as movement vigor (55–57). In the present study, we describe a within trial level of coordination between decisions and actions possibly established to deal with deadlines and promote the value of the immediate reward. This mechanism might be implemented within the cortico-basal ganglia-thalamo-cortex circuits, with the dopamine as a transmitter dedicated to the exploitation of the local reward opportunities (58). Interestingly, a recent study conducted by Herz and colleagues (39) shows that the subthalamic nucleus (STN) can independently control movement and decision speed in distinct time windows, a result that is compatible with the contribution of this basal ganglia nucleus to the decoupled coordination of decisions and actions that we report at the behavioral level in the present study. We also propose that the different modes of coordination of decisions and actions may serve more global, abstract goals such as the optimization of the rate of successes at the session level. This mechanism might be implemented in broader circuits under the regulation of neuromodulation systems, such as the noradrenergic circuits (59–61). Another possibility, not mutually exclusive, is that structures known to perform predictive computing and build internal models of the world, such as the cerebellum, would play a role in such long term goal establishment (26, 62), by building internal models of behavior utility in each task, allowing to optimize the global value of behavior over extended time scales.

Supplemental material

Supplemental Figs. S1-S5: https://doi.org/10.6084/m9.figshare.27125388

References

1. **Cisek P**, **Kalaska JF**. Neural Mechanisms for Interacting with a World Full of Action Choices. *Annual Review of Neuroscience* 33: 269–298, 2010. doi: 10.1146/annurev.neuro.051508.135409.

2. **Shadmehr R**. Control of movements and temporal discounting of reward. *Current Opinion in Neurobiology* 20: 726–730, 2010. doi: 10.1016/j.conb.2010.08.017.

3. **Gallivan JP**, **Chapman CS**, **Wolpert DM**, **Flanagan JR**. Decision-making in sensorimotor control. *Nat Rev Neurosci* 19: 519–534, 2018. doi: 10.1038/s41583-018-0045-9.

4. **Shadmehr R**, **Reppert TR**, **Summerside EM**, **Yoon T**, **Ahmed AA**. Movement Vigor as a Reflection of Subjective Economic Utility. *Trends in Neurosciences* 42: 323–336, 2019. doi: 10.1016/j.tins.2019.02.003.

5. **Haith AM**, **Reppert TR**, **Shadmehr R**. Evidence for Hyperbolic Temporal Discounting of Reward in Control of Movements. *Journal of Neuroscience* 32: 11727–11736, 2012. doi: 10.1523/JNEUROSCI.0424-12.2012.

6. **Thura D**, **Cos I**, **Trung J**, **Cisek P**. Context-dependent urgency influences speed-accuracy trade-offs in decision-making and movement execution. *J Neurosci* 34: 16442–16454, 2014. doi: 10.1523/JNEUROSCI.0162-14.2014.

7. **Reppert TR**, **Lempert KM**, **Glimcher PW**, **Shadmehr R**. Modulation of Saccade Vigor during Value-Based Decision Making. *Journal of Neuroscience* 35: 15369–15378, 2015. doi: 10.1523/JNEUROSCI.2621-15.2015.

8. **Summerside EM**, **Shadmehr R**, **Ahmed AA**. Vigor of reaching movements: reward discounts the cost of effort. *Journal of Neurophysiology* 119: 2347–2357, 2018. doi: 10.1152/jn.00872.2017.

9. **Thura D**. Decision urgency invigorates movement in humans. *Behavioural Brain Research* 382: 112477, 2020. doi: 10.1016/j.bbr.2020.112477.

10. **Korbisch CC**, **Apuan DR**, **Shadmehr R**, **Ahmed AA**. Saccade vigor reflects the rise of decision variables during deliberation. *Current Biology* 32: 5374-5381.e4, 2022. doi: 10.1016/j.cub.2022.10.053.

11. **Calalo JA**, **Ngo TT**, **Sullivan SR**, **Strand K**, **Buggeln JH**, **Lokesh R**, **Roth AM**, **Carter MJ**, **Kurtzer IL**, **Cashaback JGA**. Online movements reflect ongoing deliberation. bioRxiv: 2024.08.19.608669, 2024.

12. **Cos I**, **Bélanger N**, **Cisek P**. The influence of predicted arm biomechanics on decision making. *Journal of Neurophysiology* 105: 3022–3033, 2011. doi: 10.1152/jn.00975.2010.

13. **Burk D**, **Ingram JN**, **Franklin DW**, **Shadlen MN**, **Wolpert DM**. Motor Effort Alters Changes of Mind in Sensorimotor Decision Making. *PLoS ONE* 9: e92681, 2014. doi: 10.1371/journal.pone.0092681.

14.   **Morel P**, **Ulbrich P**, **Gail A**. What makes a reach movement effortful? Physical effort
discounting supports common minimization principles in decision making and motor
control. *PLoS Biol* 15: e2001323, 2017. doi: 10.1371/journal.pbio.2001323.

15.   **Michalski J**, **Green AM**, **Cisek P**. Reaching decisions during ongoing movements.
Journal of Neurophysiology 123: 1090–1102, 2020. doi: 10.1152/jn.00613.2019.

16.   **Marcos E**, **Cos I**, **Girard B**, **Verschure PFMJ**. Motor Cost Influences Perceptual
Decisions. *PLoS ONE* 10: e0144841, 2015. doi: 10.1371/journal.pone.0144841.

17.   **Hagura N**, **Haggard P**, **Diedrichsen J**. Perceptual decisions are biased by the cost to
act. *eLife* 6: e18422, 2017. doi: 10.7554/eLife.18422.

18.   **Yoon T**, **Geary RB**, **Ahmed AA**, **Shadmehr R**. Control of movement vigor and
decision making during foraging. *Proc Natl Acad Sci USA* 115: E10476–E10485, 2018.
doi: 10.1073/pnas.1812979115.

19.   **Reynaud AJ**, **Saleri Lunazzi C**, **Thura D**. Humans sacrifice decision-making for action
execution when a demanding control of movement is required. *Journal of*
Neurophysiology 124: 497–509, 2020. doi: 10.1152/jn.00220.2020.

20.   **Saleri Lunazzi C**, **Reynaud AJ**, **Thura D**. Dissociating the Impact of Movement Time
and Energy Costs on Decision-Making and Action Initiation in Humans. *Front Hum*
Neurosci 15: 715212, 2021. doi: 10.3389/fnhum.2021.715212.

21.   **Sukumar S**, **Shadmehr R**, **Ahmed A**. Effects of reward and effort history on decision-
making and movement vigor during foraging. .

22.   **Hurley S**. Perception And Action: Alternative Views. *Synthese* 129: 3–40, 2001. doi:
10.1023/A:1012643006930.

23.   **Cisek P**. Cortical mechanisms of action selection: the affordance competition
hypothesis. *Philosophical Transactions of the Royal Society B: Biological Sciences* 362:
1585–1599, 2007. doi: 10.1098/rstb.2007.2054.

24.   **Buzsáki G**, **Peyrache A**, **Kubie J**. Emergence of Cognition from Action. *Cold Spring*
Harb Symp Quant Biol 79: 41–50, 2014. doi: 10.1101/sqb.2014.79.024679.

25.   **Yin HH**. Action, time and the basal ganglia. *Philosophical Transactions of the Royal*
Society B: Biological Sciences 369: 20120473, 2014. doi: 10.1098/rstb.2012.0473.

26.   **Pezzulo G**, **Cisek P**. Navigating the Affordance Landscape: Feedback Control as a
Process Model of Behavior and Cognition. *Trends in Cognitive Sciences* 20: 414–424,
2016. doi: 10.1016/j.tics.2016.03.013.

27.   **Yoo SBM**, **Hayden BY**, **Pearson JM**. Continuous decisions. *Philosophical*
Transactions of the Royal Society B: Biological Sciences 376: 20190664, 2021. doi:
10.1098/rstb.2019.0664.

28.   **Carsten T**, **Fievez F**, **Duque J**. Movement characteristics impact decision-making and
vice versa. *Sci Rep* 13: 3281, 2023. doi: 10.1038/s41598-023-30325-4.

29. **Kita K**, **Du Y**, **Haith AM**. Evidence for a common mechanism supporting invigoration
of action selection and action execution. *Journal of Neurophysiology* 130: 238–246,
2023. doi: 10.1152/jn.00510.2022.

30. **Kawagoe R**, **Takikawa Y**, **Hikosaka O**. Expectation of reward modulates cognitive
signals in the basal ganglia. *Nat Neurosci* 1: 411–416, 1998. doi: 10.1038/1625.

31. **Wickler SJ**, **Hoyt DF**, **Cogger EA**, **Hirschbein MH**. Preferred speed and cost of
transport: the effect of incline. *Journal of Experimental Biology* 203: 2195–2200, 2000.
doi: 10.1242/jeb.203.14.2195.

32. **Milstein DM**, **Dorris MC**. The influence of expected value on saccadic preparation. *J*
Neurosci 27: 4810–4818, 2007. doi: 10.1523/JNEUROSCI.0577-07.2007.

33. **Xu-Wilson M**, **Zee DS**, **Shadmehr R**. The intrinsic value of visual information affects
saccade velocities. *Exp Brain Res* 196: 475–481, 2009. doi: 10.1007/s00221-009-1879-
1.

34. **Reppert TR**, **Rigas I**, **Herzfeld DJ**, **Sedaghat-Nejad E**, **Komogortsev O**, **Shadmehr**
R. Movement vigor as a traitlike attribute of individuality. *Journal of Neurophysiology*
120: 741–757, 2018. doi: 10.1152/jn.00033.2018.

35. **Thura D**, **Cabana J-F**, **Feghaly A**, **Cisek P**. Integrated neural dynamics of
sensorimotor decisions and actions. *PLoS Biol* 20: e3001861, 2022. doi:
10.1371/journal.pbio.3001861.

36. **Thura D**, **Cisek P**. The Basal Ganglia Do Not Select Reach Targets but Control the
Urgency of Commitment. *Neuron* 95: 1160-1170.e5, 2017. doi:
10.1016/j.neuron.2017.07.039.

37. **Carland MA**, **Thura D**, **Cisek P**. The Urge to Decide and Act: Implications for Brain
Function and Dysfunction. .

38. **Saleri Lunazzi C**, **Thura D**, **Reynaud AJ**. Impact of decision and action outcomes on
subsequent decision and action behaviours in humans. *European Journal of*
Neuroscience 57: 1098–1113, 2023. doi: 10.1111/ejn.15932.

39. **Herz DM**, **Bange M**, **Gonzalez-Escamilla G**, **Auer M**, **Ashkan K**, **Fischer P**, **Tan H**,
Bogacz R, **Muthuraman M**, **Groppa S**, **Brown P**. Dynamic control of decision and
movement speed in the human basal ganglia. *Nat Commun* 13: 7530, 2022. doi:
10.1038/s41467-022-35121-8.

40. **Gold JI**, **Shadlen MN**. The Neural Basis of Decision Making. *Annu Rev Neurosci* 30:
535–574, 2007. doi: 10.1146/annurev.neuro.29.051605.113038.

41. **Shenoy KV**, **Sahani M**, **Churchland MM**. Cortical control of arm movements: a
dynamical systems perspective. *Annu Rev Neurosci* 36: 337–359, 2013. doi:
10.1146/annurev-neuro-062111-150509.

42. **Cisek P**, **Puskas GA**, **El-Murr S**. Decisions in Changing Conditions: The Urgency-
Gating Model. *Journal of Neuroscience* 29: 11560–11571, 2009. doi:
10.1523/JNEUROSCI.1844-09.2009.

43.   **Thura D**, **Cisek P**. Deliberation and commitment in the premotor and primary motor
cortex during dynamic decision making. *Neuron* 81: 1401–1416, 2014. doi:
10.1016/j.neuron.2014.01.031.

44.   **Pylyshyn Z**. *Computation and Cognition: Toward a Foundation for Cognitive Science*.
Cambridge, MA: MIT Press, 1984.

45.   **Leroy É**, **Koun É**, **Thura D**. Integrated control of non-motor and motor efforts during
perceptual decision-making and action execution: a pilot study. *Sci Rep* 13: 9354, 2023.
doi: 10.1038/s41598-023-36443-3.

46.   **Fievez F**, **Cos I**, **Carsten T**, **Derosiere G**, **Zénon A**, **Duque J**. Task goals shape the
relationship between decision and movement speed. bioRxiv: 2023.12.29.573524, 2023.

47.   **Morvan T**, **Eloy C**, **Robbe D**. Running, Fast and Slow: The Dorsal Striatum Sets the
Cost of Movement During Foraging. bioRxiv: 2024.05.31.596850, 2024.

48.   **Bogacz R**, **Hu PT**, **Holmes PJ**, **Cohen JD**. Do humans produce the speed–accuracy
trade-off that maximizes reward rate? *Quarterly Journal of Experimental Psychology* 63:
863–891, 2010. doi: 10.1080/17470210903091643.

49.   **Balci F**, **Simen P**, **Niyogi R**, **Saxe A**, **Hughes JA**, **Holmes P**, **Cohen JD**. Acquisition of
decision making criteria: reward rate ultimately beats accuracy. *Atten Percept*
Psychophys 73: 640–657, 2011. doi: 10.3758/s13414-010-0049-7.

50.   **Myerson J**, **Green L**. Discounting of delayed rewards: Models of individual choice. *J*
Exp Anal Behav 64: 263–276, 1995. doi: 10.1901/jeab.1995.64-263.

51.   **Shadmehr R**, **Ahmed AA**. Vigor: Neuroeconomics of Movement Control [Online]. The
MIT Press. https://doi.org/10.7551/mitpress/12940.001.0001.

52.   **Thura D**. Reducing behavioral dimensions to study brain–environment interactions.
Behavioral and Brain Sciences 44, 2021. doi: 10.1017/S0140525X21000169.

53.   **Thura D**, **Cisek P**. Microstimulation of dorsal premotor and primary motor cortex
delays the volitional commitment to an action choice. *Journal of Neurophysiology* 123:
927–935, 2020. doi: 10.1152/jn.00682.2019.

54.   **Robbe D**, **Dudman JT**. The Basal Ganglia invigorate actions and decisions. In: *The*
Cognitive Neurosciences. MIT Press, 2020.

55.   **Turner RS**, **Desmurget M**. Basal ganglia contributions to motor control: a vigorous
tutor. *Current Opinion in Neurobiology* 20: 704–716, 2010. doi:
10.1016/j.conb.2010.08.022.

56.   **Dudman JT**, **Krakauer JW**. The basal ganglia: from motor commands to the control of
vigor. *Current Opinion in Neurobiology* 37: 158–166, 2016. doi:
10.1016/j.conb.2016.02.005.

57.   **Yttri EA**, **Dudman JT**. Opponent and bidirectional control of movement velocity in the
basal ganglia. *Nature* 533: 402–406, 2016. doi: 10.1038/nature17639.

58.   **Niv Y**, **Daw ND**, **Joel D**, **Dayan P**. Tonic dopamine: opportunity costs and the control of
response vigor. *Psychopharmacology (Berl)* 191: 507–520, 2007. doi: 10.1007/s00213-
006-0502-4.

59.   **Murphy PR**, **Boonstra E**, **Nieuwenhuis S**. Global gain modulation generates time-
dependent urgency during perceptual choice in humans. *Nat Commun* 7: 13526, 2016.
doi: 10.1038/ncomms13526.

60.   **Steinemann NA**, **O'Connell RG**, **Kelly SP**. Decisions are expedited through multiple
neural adjustments spanning the sensorimotor hierarchy. *Nat Commun* 9: 3627, 2018.
doi: 10.1038/s41467-018-06117-0.

61.   **Kaduk K**, **Henry T**, **Guitton J**, **Meunier M**, **Thura D**, **Hadj-Bouziane F**.
Atomoxetine and reward size equally improve task engagement and perceptual decisions
but differently affect movement execution. *Neuropharmacology* 241: 109736, 2023. doi:
10.1016/j.neuropharm.2023.109736.

62.   **Kruithof ES**, **Klaus J**, **Schutter DJLG**. The human cerebellum in reward anticipation
and outcome processing: An activation likelihood estimation meta-analysis. *Neurosci
Biobehav Rev* 149: 105171, 2023. doi: 10.1016/j.neubiorev.2023.105171.