Title: Recurrent drought increases grassland community seasonal synchrony

Running head: Recurrent drought increases synchrony

Lena M. Müller[1], Michael Bahn[1], Maximillian Weidle[1], Georg Leitinger[1], Dina in 't Zandt[1,2]

[1]Department of Ecology, Universität Innsbruck, Innsbruck, Austria

[2]Department of Population Ecology, Institute of Botany, The Czech Academy of Sciences, Průhonice, Czech

Republic

Corresponding author: Lena Marie Müller, 00491636954280, contact@lenamueller.com, orchid:

0000-0001-6193-0672

Abstract

Climate change leads to an increase in the frequency and severity of drought events with

strong repercussions on grassland ecosystems. While the effects of single drought events

on ecosystem structure and functioning are well understood, it is largely unknown whether

and how drought frequency modifies ecosystem responses to drought. Here, we assessed

how the increase in frequency of severe, annual summer drought affected grassland

communities. We examined these effects in a species-rich sub-alpine mountain meadow

with a drought frequency of one, three, and 13 years, as well as ambient conditions. We

found that high drought frequency increased seasonal plant community synchrony through a

reduction in species richness, a shift of plant functional groups, a loss of early-seasonal plant

species, and the constrained establishment of new species and individuals throughout the

growing season. These changes are associated with a decreased fraction of biomass as

drought frequency increased. We conclude that with the ongoing increase in drought

frequency due to climate change, grassland plant communities will increase in seasonal

synchrony. We suggest that this will leave ecosystems highly vulnerable to future

disturbances, as asynchrony is a critical component of stability. Furthermore, we show that

negative drought effects were enhanced with an increasing drought frequency. We therefore

conclude that single and low-frequency drought studies may not adequately predict longer-

term changes in our rapidly shifting climate. Moreover, we found that negative drought

effects on plant communities outweighed the weak adaptive effects of species. We therefore

conclude that plant communities are unlikely to be able to adapt to recurrent drought.

Keywords: recurrent drought, drought frequency, seasonal synchrony, synchrony,

asynchrony, grassland, species richness, early-seasonal species, functional groups, drought

1. Introduction

Grasslands cover 40 % of the Earth's surface and are crucial for carbon and nutrient cycling, biodiversity conservation, and ecosystem services (Petermann & Buzhdygan 2021; Blair *et al.* 2014; Dixon *et al.* 2014). However, up to 49 % of the world's grasslands are currently degraded, with the climate crisis predicted to exacerbate this number (Bardgett *et al.* 2021; Gibbs & Salmon 2015). Among the detrimental climate scenarios, the increase in frequency and severity of drought events stand out as particularly damaging (IPCC 2023, 2021; Wang *et al.* 2021; Spinoni *et al.* 2018). Drought events alter grassland nutrient cycling, disrupt interactions between species, result in species extinction, and therewith cause severe disruptions in ecosystem structure and function (Smith *et al.* 2024; Oram *et al.* 2023; Bardgett & Caruso 2020; Ingrisch *et al.* 2020; Tello-García *et al.* 2020). In the long-term, the predicted increase in the frequency of drought events therefore threatens to irrevocably change grassland ecosystems, including the fundamental services and biodiversity they conserve (Müller & Bahn 2022; Xu *et al.* 2021; Schirpke *et al.* 2019). At the same time, potential adaptations of species and communities may dampen these detrimental effects to some extent (Müller & Bahn 2022; Canarini *et al.* 2021; Coleman & Wernberg 2020). This occurs for example through drought-induced legacy effects that alter ecosystem structure and function after drought has subsided, and which results in plant communities being able to cope better with a subsequent drought event (Müller & Bahn 2022). Yet, while effects of single drought events are comparatively well studied, our understanding of the effects of a long-term increase in drought frequency on grassland community structure and function remains limited.

Multiple interlinked drought-induced changes, for example in plant species composition, diversity, complementarity, as well as species loss may have detrimental effects by affecting the **synchrony** of grassland communities (He *et al.* 2022; Muraina *et al.* 2021; de Mazancourt *et al.* 2013). Single drought was found to increase the synchrony of plant communities between years (Zhang *et al.* 2019), similarly as recurrent drought of three

(He *et al.* 2022) and four years (Muraina *et al.* 2021). With greater synchrony between years,

species communities exhibit increased similarity in their response to environmental changes,

the swiftness of their responses, or their functional complementarity throughout various

seasons (Loreau & de Mazancourt 2013). These changes in grassland community

synchrony likely have detrimental effects on long-term ecological functioning, especially

under a future climate with a high drought frequency. Increases in grassland community

synchrony between years has been shown to decrease community stability (Wilcox *et al.*

2017; Xu *et al.* 2015; Loreau & de Mazancourt 2013). This is because asynchronous

communities stabilise ecosystems through different responses to disturbances (de

Mazancourt *et al.* 2013). For drought disturbances, species asynchrony was suggested to be

the dominant driver to stabilise community productivity under drought (Luo *et al.* 2023b;

Muraina *et al.* 2021; Craven *et al.* 2018). Community synchrony between years is thus

influenced by drought and is suggested to shape a communities' response to subsequent

droughts by decreasing its stability. However, besides changing community synchrony

between seasons, drought is also likely to affect community synchrony *within* the growing

season. Therewith, it might be a critical mechanism in shaping responses to recurrent

drought. However, we currently lack evidence on the effects of recurrent drought on

community *seasonal* synchrony. In the presence of more frequent droughts, we hypothesize

that the plant community will become more synchronous within the season, which also leads

to reduced stability in the face of disturbances.

79   Drought events alter plant community composition, phenology, and productivity

(Smith *et al.* 2023; Müller & Bahn 2022). Drought leads to a decrease in plant community

productivity, by limiting plant growth through reduced water availability, or by mortality of

species under severe drought conditions (Sippel *et al.* 2018; Frank *et al.* 2015). Additionally,

after a drought event, soil rewetting results in a temporarily increased availability of nitrogen

(Ingrisch *et al.* 2018; Karlowsky *et al.* 2018). This surge in nitrogen supports the recovery

and, in some cases, leads to a short-term overshoot in productivity (Griffin-Nolan *et al.* 2018;

Hofer *et al.* 2016), which can compensate for the lower plant community productivity during

the drought event (Hahn *et al.* 2021). This response is mainly based on fast-growing grass

species, which makes the community sensitive to changes in composition post drought

(Mackie *et al.* 2019; Stampfli *et al.* 2018). An important issue is that drought-induced

changes in productivity and composition in grasslands may take a long time to come back to

pre-disturbance level (Müller & Bahn 2022). If a subsequent drought falls within this recovery

timespan of a previous drought, it may lead to constantly impaired ecosystems (Schwalm *et*

al. 2017), and consequently to a more profound effect on productivity under a high frequency

of drought (Xu *et al.* 2021). Hence, a potential accumulation of effects can emerge when

multiple droughts occur annually. However, as recurrent drought events are typically studied

in two-year experiments and only rarely extend beyond this timeframe, we lack critical

knowledge on the productivity loss that builds up under a high drought frequency.

Drought shifts plant community **composition** towards more slower growing and more

drought adapted-species (Wilcox *et al.* 2021). For example, drought may lead to a

dominance of grasses over forbs (Xu *et al.* 2021; DeBoeck *et al.* 2018; Hoover *et al.* 2014).

This may be driven by a higher post-drought resource use of grasses compared with the

conservative resource use of forbs (Stampfli *et al.* 2018) or by a more efficient resource

acquisition of water of grasses compared with forbs (Tello-García *et al.* 2020). Furthermore,

drought-induced changes in plant community composition are driven by a reduction in plant

species performance and an increase in mortality of plants both during and after a drought

event (Sippel *et al.* 2018; Frank *et al.* 2015). In the long term, plant compositional shifts due

to high drought frequency are likely to relate additionally to drought-induced changes in plant

reproductive effects, e.g. reduction in reproductive shoots, seeds (Zeiter *et al.* 2016),

belowground bud density (Qian *et al.* 2022), as well as altered seed banks (Basto *et al.*

2018), and regeneration from seed (Stampfli & Zeiter 2004). Through these drought-induced

changes in plant species composition, recurrent drought events may in the long-term result

in permanent loss of plant species with potentially irreversible consequences for the

functioning of plant communities. Furthermore, under recurrent drought, plant species also

have the capacity to adapt (Müller & Bahn 2022), which dampens drought effects under a

higher drought frequency (Backhaus *et al.* 2014; Walter *et al.* 2011). Therefore, a high

drought frequency is expected to result in plant communities consisting of drought-tolerant

species, a dominance of grasses, plant species profiting from the recovery dynamics of

drought events, as well as plant species that adapt to recurrent drought conditions. However,

due to the lack of studies investigating high drought frequency, long-term compositional

shifts might have been overlooked to date.

Drought alters **phenology** by shifting the beginning or end of the growing season

(Yuan *et al.* 2023; Ge *et al.* 2022; Rihan *et al.* 2022; Wang *et al.* 2022; Ji *et al.* 2021; Zeng *et*

al. 2021; Yuan *et al.* 2020; Berwaers *et al.* 2019; Kang *et al.* 2018) and advancing end-of-

season leaf senescence (Hoover *et al.* 2021; Berwaers *et al.* 2019; Peng *et al.* 2019; Kang

et al. 2018). Furthermore, drought alters species flowering phenology by advancing the

flowering date and extending the flowering length, whereby effects differ strongly between

species (Castillioni *et al.* 2022; Vorkauf *et al.* 2021; Forrest & Miller-Rushing 2010; Jentsch

et al. 2009). Hence, recurrent drought is expected to move the duration and timing of the

growing season and flowering with effects on community composition, though the directions

and magnitudes of these changes under recurrent drought are still poorly understood.

We lack a profound understanding of the implications under a high frequency of drought

events on the temporal shifts in community composition, productivity, and phenology. Here,

we assessed how the increase in frequency of severe summer drought affected grassland

community structure, function, and seasonal dynamics. We examined this in a species-rich

sub-alpine mountain meadow with increasing annual summer drought frequencies of up to

13 years. We quantified species abundances in the field throughout the growing season and

sampled plant species-specific biomass and necromass at peak drought and drought

recovery. We asked the following questions: How did an increase in drought frequency affect

(1) plant community productivity, species richness, and flower production during peak

drought and recovery, respectively, as well as (2) plant community seasonal synchrony. We

examined whether these patterns were underlain by shifts in (3) plant community

composition, (4) plant functional groups, and (5) plant community traits. Taken together, we

present the effects of a high frequency of drought events on plant communities and discuss

underlying mechanisms.

2.   Material and Methods

2.1 Study Site

The studied sub-alpine grassland is located in the Austrian Central Alps in the Stubai valley

at 1820 m a.s.l. (47°70 45″N, 11°180 20″E). The site is exposed southeast with an inclination

of ~ 20° and has an average annual temperature of 3 °C and an annual precipitation of 1100

mm. The soil type is a dystric cambisol (top soil pH = 5.5). The vegetation community is

classified as *Trisetetum flavescentis*, species-rich with on average 49 species per m^2 and

consists largely of perennial grasses and forbs. The growing season lasts from the end of

April until mid-October. The sub-alpine grassland is manured every 2-3 years, mown once in

early August at peak biomass, and sporadically grazed in spring and autumn. The studied

area, however, has not been fertilised and is excluded from grazing since 2007. For further

details on the site see Ingrisch *et al.* (2018), Hasibeder *et al.* (2015), and Schmitt *et al.*

(2010).

2.2 Experimental set-up

Four summer drought treatments were established in the enclosed area at the study site: a

single year summer drought, three years of recurrent summer drought, 13 years of recurrent

summer drought, and ambient conditions (n = 4, respectively). The treatment of 13 years of

recurrent drought was established in 2008, the three-year recurrent drought in 2018, and the

single-year drought in 2020. Summer drought was simulated for seven weeks from 17 June

until 7 August 2020 using rainout shelters (Fig 1). This is the warmest period in the growing

season and therefore the period where a natural drought is most likely to occur. The timing

and intensity of the annual drought were kept as similar as possible in previous years since

2008. The ambient treatment was not covered with rainout shelters and thus exposed to

ambient precipitation. The rainout shelters of the single and three years of recurrent summer

drought had a base area of 3 x 3.5 m^2 and 2.5 m height. For the rainout shelters of the 13

years of recurrent drought, two plots are combined, respectively, and as a result, the shelters

had a base area of 3 x 7 m^2. All shelters were covered with a transparent UV-A and UV-B

transmissive plastic foil (Lumisol clear AF, Folitec, Westerburg, Germany, light transmittance

ca. 90 %). To allow air circulation, the shelters were open at the bottom (up to 0.5 m) and at

the top of the face sides. Within each of the treatments, one permanent 20 cm x 20 cm plot

was installed four weeks before summer drought simulation (n = 4; in total 16). Each plot

was located at least 50 cm away from the rainout shelter to avoid edge effects. The drought

period was ended by removing the shelters and watering of all 20 x 20 cm^2 plots with 31 mm

of distilled water. At this time point, the entire site was also mown, and the cuttings were

removed.

Fig 1 The course of the experiment including setup and removal of the rainout shelters, and peak

drought and recovery harvest. The pictures show the drought intensity at peak drought of ambient

conditions, one year, three years, and 13 years of recurrent drought. One replicate per treatment is

shown.

2.3 Measurements

2.3.1 Microclimate

A microclimate station at the study site recorded precipitation, air temperature, and relative

air humidity (see details in Hasibeder *et al.* (2015)). During the whole season, soil water

content (SWC) (Decagon EC-5 and 5TM, Decagon Devices, Pullman, WA, USA; SM300,

Delta-T Devices, Cambridge, UK; Hobo S-SMD-M005, Onset Computer corporation,

Bourne, MA, USA) was measured continuously in the main rooting horizon (30-min interval)

at 5, 10, and 20 cm of soil depth. The measurements were replicated twice in the single-,

and three-years treatment, and three to four times in the ambient treatment. Due to partial

sensor failures, no data is available for the 13-years treatment. For results on the

microclimate and SWC, see Fig S1.

2.3.2 Measurements throughout the growing season

In all 16 plots, species composition and vegetation height were measured every two weeks

from 22 May to 22 October 2020. Plant species composition was estimated by noting for

each species present the number of modules and flowering stems. Modules refer to the

number of tillers for grasses and graminoids, the number of rosettes for small dicots, and the

number of leaves for larger dicots per rooted species, following (in 't Zandt *et al.* 2021;

Herben *et al.* 2013; Herben *et al.* 1997). The data was used to calculate the total number of

modules, plant species number, community evenness, and Shannon-diversity index

throughout the growing season. Counting was accompanied by measurements of mean

vegetation height (in cm). Seedling composition in the plots was counted separately every

three weeks between the 23 June and the 22 October 2020. A seedling was defined as a

plant individual with less than two leaves, and was not part of the module counts. Seedlings

were counted per plant functional group: forbs, grasses, and legumes.

2.3.3 Biomass, necromass, and leaf traits

Aboveground biomass (living plant material) and necromass (dead plant material) was

sampled destructively in the plots at peak drought on 6 August 2020 and during recovery on

22 October 2020 (Fig 1). For this, plant material was cut at 2 cm above the soil and frozen at

-18 °C until further analysis. Samples were split into plant species and separated into leaves,

stems, and flowers. This was done for biomass and necromass, respectively. For each

species of each plot, the leaf area was determined from thawed leaves that were saturated

with water and scanned (V700 Photo, Epson, WinRHIZO Pro 2012, Regent Instruments).

Scanned leaves were dried in the subsequent three days at 60 °C and were weighed.

Specific leaf area (SLA) was calculated by dividing the scanned leaf area by the dry weight

of the leaves for each species and treatment, respectively. Leaf area index (LAI) was

calculated from the leaf surface of all leaves divided by the total unit of ground area of the

plot.

We also assessed drought frequency effects on aboveground phytomass fractions.

These were calculated as the proportion of biomass, necromass, and flower mass of the

total aboveground phytomass in each plot. We did this, because phytomass fractions were in

most cases more informative concerning drought frequency effects than absolute values of

phytomass (Fig S2) as well as biomass and necromass (data not shown). This is because,

most phytomass at the alpine meadow is produced early in the season between May and

mid-July (data not shown), which is before the peak drought. Hence, in our study, drought

does not impair plant development and therefore has few effects on phytomass production

(Fig S2), as is typically the case in shorter-term drought experiments. Increasing drought

rather resulted in phytomass loss and/or plant mortality, which is visible in *fractions* of the

phytomass, as it illustrates the redistribution of mass. Notably, while biomass refers to

productivity, biomass fraction reflects the maintained productivity.

We assessed plant species seasonality, indicating their growth timing during the

season. For this, we determined the time point at which the plant species peaked in its

number of modules. The typical seasonal behaviour of each plant species was determined

based on the ambient treatment plots only. Early seasonal species were defined as species

that peaked in modules before the biomass cut and thus reached peak growth between the

end of May to end of July. Late seasonal species were defined as species that peaked in

modules after the biomass cut and thus reached peak growth between August and October.

Shifts in early and late seasonal patterns of species were assessed using values from all

treatments.

Because of the low occurrence of flowers, seasonal shifts in flowering was calculated

based on flower biomass. For this, we calculated the ln-response ratio between a species'

flower dry mass at peak drought and its flower dry mass after the 11-week recovery period.

A positive value signifies plant species flowering early in the growing season, i.e., during the

drought period. A negative value signifies plant species flowering late in the growing season,

i.e., during the recovery phase.

2.3.4 Plant community metrics and composition

Plant community evenness was calculated as Pilou's index. For this, we divided the plant

diversity Shannon index by the ln-transformed species richness using the function diversity

and specnumber of the vegan package (R Core Team 2021; Oksanen *et al.* 2020). Plant

community asynchrony was calculated following Loreau & de Mazancourt (2008), using the

package codyn (Hallett *et al.* 2016). Using the same package, plant species loss between

subsequent time points was calculated to indicate species loss *within* the growing season.

To determine whether communities were dominated by either early or late plant species, we

calculated the proportional number of early plant species present in each community at each

time point.

We assessed plant community composition using Non-Metric Multidimensional

Scaling (NMDS) using metaMDS from the vegan package (Oksanen *et al.* 2020). NMDS on

both module counts and biomass were run on Bray-Curtis dissimilarity matrices and with four

dimensions to minimise stress values. Significant separation of communities resulting from

'drought frequency' (factor), date (scaled, continuous), and their interaction was tested using

a restricted permutation test: samples were permuted *within* plots to take the repeated

sampling structure over time into account. The test was performed using adonis from the

vegan package with 999 permutations (Oksanen *et al.* 2020). A passive overlay of 'drought

frequency' (ln-transformed + 1), 'species richness', 'proportion early species', 'synchrony',

species loss between subsequent time points (for NMDS based on counts only), 'LAI',

functional group total abundance, flower abundance, and seedling abundance was created

and tested for significance using the function envfit of the vegan package with 999

permutations (Oksanen *et al.* 2020).

2.3.5 Statistics

All calculations and statistical analyses were done in R 4.1.1 (R Core Team 2021). Effects of

drought frequency on percentage data were performed using Generalized Linear Mixed

Models (GLMM) with a beta distribution and logit-link using the package glmmTMB

(Magnusson *et al.* 2017). Drought frequency effects on species richness were tested using

Generalised Linear Mixed-Effects Models (GLMER) with a poisson distribution and log-link

function using the lme4 package (Bates *et al.* 2015). Drought frequency effects on

continuous variables were tested using Linear Mixed Effects Models (LMER) using the lme4

package (Bates *et al.* 2015). Models including 'date' (continuous, scaled) were run with 'plot'

as a random effect to take the repeated sampling design into account. Models on plant

species-specific traits were only run for plant species for which at least two data points

occurred in each treatment.

Heterogeneity of the mountain meadow was taken into account as a random effect in

statistical models. For models on plant community measures, 'block' was added as a

random effect. Models on plant species-specific traits included the mean vegetation height of

each plot before the setup of the rainout shelters as a random effect. This variable was

better at capturing variation *within* drought treatments than 'block' and, importantly, was not

significantly affected by previous drought events (data not shown).

For GLMM and GLMER models, we tested over- and under-dispersion of the model

residuals using the package DHARMa (Hartig 2022). For LMER models, we tested

assumptions of homogeneity of variances, and normal residual distributions following Zuur *et*

al. (2010). Data was ln- or sqrt-transformed when model residuals did not adhere to model

assumptions.

3. Results

3.1 Drought frequency decreased biomass and species richness and increased

necromass

In a species-rich sub-alpine mountain meadow, we assessed how the annual frequency of

severe summer drought affected plant community phytomass, composition, and seasonal

dynamics. At the peak of drought, 7 weeks after the setup of the rainout shelters, we found

that drought frequency significantly decreased the fraction of community biomass and

increased the fraction of community necromass (Fig 2AB). The increase in necromass was

primarily related to an increase in the necromass fraction of grass and legume species (Fig

S3). Eleven weeks after soil rewetting (recovery), these effects were still present, albeit to a

lesser extent (Fig S4AB). In the recovery phase, the fraction of forb biomass decreased,

while the fraction of grass biomass and necromass increased with drought frequency (Fig

S3).

Drought frequency did not significantly affect the fraction of flower biomass at peak

drought (Fig 2C). During recovery, however, the fraction of flower biomass was significantly

decreased with increasing drought frequency (Fig S4C).

Furthermore, drought frequency significantly decreased plant species richness at both peak

drought and recovery (Fig 2D; S4D). On average, 13 years of recurrent summer drought

resulted in a 35 % and 46 % loss of plant species compared with ambient conditions at peak

drought and recovery, respectively. Despite this loss in plant species, drought frequency did

not significantly affect plant community evenness at both peak drought and recovery (Fig 2E;

S4E).

Fig 2 Drought frequency effects at peak drought on (A) the fraction of plant community biomass (living

plant material), (B) the fraction of plant community necromass (dead plant material), (C) the fraction of

total flower mass, (D) plant community species richness, and (E) plant community evenness. A-C are

expressed as percentages of the phytomass (bio- and necromass) in each plot. For significant

relations, the black line indicates the mean relation with the shading showing the 95% confidence

interval. P-values of significant relations are shown (ns: no significant effect, p > 0.05, n = 4 per

drought frequency level). For effects of drought frequency on overall plant community parameters in

the recovery phase, see Fig S4.

3.2 Drought frequency decreased forb and legume seedlings

To assess the impact of drought frequency on plant community seasonal development, we

monitored the number of plant species modules and seedlings at biweekly intervals

throughout the season. Drought frequency affected the number of forb and legume

seedlings, but not of grass seedlings. During drought and recovery, the number of forb and

legume seedlings was consistently lowest with 13 years of recurrent summer drought (Fig 3).

Additionally, while in the ambient treatment and one- year drought plots, many forb and

legume seedlings appeared after the mowing event, these peaks were lower in plots with

three years and especially 13 years of recurrent summer drought (Fig 3). At the same time,

plant community LAI decreased with drought frequency at both peak drought and recovery

(Fig S5AB). The latter indicates that the soil surface was less covered by leaves of the

established plant community with an increase in drought frequency.

Fig 3 Drought frequency effects on the number of seedlings of (A) grass, (B) forb, and (C) legume

species throughout the growing season. The dotted line indicates the timepoint at which drought

ended, and the plots were mown and rewetted. Solid lines indicate mean seedling numbers, and

faded lines indicate seedling numbers of each plot (n = 4 per drought frequency treatment). Results

from analyses of variances are shown with '***' indicating p < 0.001, '*' indicating 0.01 > p < 0.05, and

'ns' indicating no significant effect (p > 0.05).

3.3 Drought frequency increased plant community seasonal synchrony

The total number of plant modules in all treatments fluctuated throughout the growing

season (Fig S6A). However, it was not significantly affected by drought frequency throughout

the season (Fig S6A). Furthermore, plant species richness before, during, and after drought

was consistently lowest in plots subjected to 13 years of drought, followed by three years,

and one year of drought (Fig S6B). Plant species richness across all treatments dropped

sharply after peak drought and the mowing event but increased to levels similar to their

respective pre-drought conditions within a month of recovery (Fig S6B).

Using the module counts of each plant species over the season, we calculated plant

community seasonal synchrony. We found that drought significantly increased seasonal

synchrony, that is, drought synchronised the timing of plant species growth during the

season. Increasing drought frequency enhanced this synchronising effect (Fig 4A).

Furthermore, seasonal synchrony was negatively related to the average plant species

richness throughout the season, that is, synchrony was high when average plant species

richness was low (Fig 4B).

Fig 4 (A) Effects of increasing drought frequency on seasonal synchrony (sensu Loreau & de

Mazancourt (2008)) based on plant module counts across the growing season from mid-May to mid-

October, and (B) seasonal synchrony in relation to the average plant species richness (n = 4 per

drought frequency treatment). The black line indicates the mean relation with the shading showing the

95% confidence interval. P-values of significant relations are shown.

3.4 Drought frequency shifts plant community composition towards less early

seasonal and more grass species

We examined the effect of drought frequency on plant community composition throughout

the season using non-metric multidimensional scaling (NMDS). We found significant

differentiation of plant communities based on drought frequency, with communities

experiencing no drought and 13 years of drought exhibiting the greatest dissimilarity (Fig 5A,

S7). This compositional separation, driven by drought frequency, was primarily captured

along the first axes of the NMDS plot. This first axis coincided with the decrease in plant

species richness and the increase in community synchrony as drought frequency increased

(Fig 5A). Moreover, we observed that the compositional shifts associated with increasing

drought frequency were characterised by a high grass abundance and a decrease in early-

seasonal plant species. In addition, LAI, the number of forb flowers, and the number of forb

seedlings decreased with drought frequency (Fig 5A). These patterns were similar when

basing the analyses on plant biomass collected at peak drought and recovery (Fig S8).

During the recovery phase, plant communities in the plots exposed to one and three

years of drought shifted along the second NMDS axis. After rewetting, these one- and three-

year droughted plots temporarily shifted into the top-right quadrant of the NMDS (Fig 5B).

This direction was related to a high loss of species, particularly early-seasonal species (for

specification see Table S1), and a decrease in forb and legume abundance (Fig 5B).

However, later in the recovery phase, the one- and three-year droughted communities

gradually returned to a similar position on the second axis of the NMDS plot as before the

recovery phase (Fig 5B). This recovery pattern indicates slower recovery of early-seasonal

forb, and legume species within the one- and three-year droughted plots (Fig 5B).

Fig 5 Non-metric multidimensional scaling (NMDS) of plant species module counts throughout the

growing season in permanent plots receiving no drought (0 years), one year, three years, and 13

years of recurrent summer drought. (A) Drought frequency effects on plant community composition

with a passive overlay of community variables significantly relating to the NMDS patterns. Each point

represents a plot at a point in time. Hulls group all plots and time points per drought frequency

treatment together. Length of arrows indicate the strength of the relation. 'Proportion early species'

refers to the proportion of early species versus the total number of species present in a plot. 'Grass',

'forb', and 'legume abundance' refer to the total number of modules summed per functional group.

'Synchrony' refers to community seasonal synchrony sensu Loreau & de Mazancourt (2008). 'Species

loss' refers to the percentage of species lost *within* the growing season between two subsequent time

points. 'LAI' refers to leaf area index. Results of a restricted permutation test is shown with '**'

indicating $0.001 > p < 0.01$, '*' indicating $0.01 > p < 0.05$, and 'ns' indicating no significant effect ($p >$

0.05) ($n = 4$ plots per drought frequency treatment). (B) Community compositional development over

time. Points are mean values ± standard errors. Note that axes scales differ between A and B to

improve readability. For NMDS axes 3 and 4, see Fig S7. For position of plant species along the axes,

see Fig S9.

3.5 Weak drought frequency effects on individual plant species responses

For 10 plant species that were present in sufficient abundances in all three drought

frequency treatments and the ambient treatment, we tested whether drought frequency

affected SLA at peak drought and recovery, as well as the timing of their growth in the

season. At peak drought, we found that drought frequency significantly decreased SLA of

Alchemilla vulgaris, Leontodon hispidus, Plantago lanceolata (marginally)*, Trifolium repens,*

and *Veronica chamaedrys* (Fig S10)*.* During recovery, these patterns had disappeared (Fig

S11)*.*

Drought frequency only significantly affected the timing of *Trifolium pratense,* which peaked

in number of modules earlier in the season with a high drought frequency (Fig S12). Drought

frequency only significantly affected the timing of flowering of *Leontodon hispidus,* which

shifted its flower production to earlier in the season with a high drought frequency (Fig S13).

4. Discussion

We investigated the effects of recurrent summer drought of up to 13 years on plant

communities in a sub-alpine mountain meadow, as well as their underlying mechanisms. We

found that drought frequency increased plant community seasonal synchrony. This increase

in synchrony with greater drought frequency was underpinned by a decrease in species

richness, a loss of early-seasonal plant species, a shift in plant functional groups, and the

constrained establishment of new species and individuals throughout the growing season.

Moreover, these changes were associated with a decreased biomass fraction as drought

frequency increased.

Higher seasonal synchrony with increasing drought frequency due to species loss,

compositional shifts, and lower species establishment

We found that with increasing drought frequency, species **synchrony** within the plant

community was enhanced. This indicates that the species in the plant community under

increasing drought frequency were more similar in their timing of plant species growth

throughout the season. Results on increasing synchrony *between* years for grasslands were

reported by other studies, finding an increase in synchrony with drought frequency of three

years in an alpine meadow (He *et al.* 2022), and four years across grasslands (Muraina *et al.*

2021). Our study suggests that with increasing drought frequency (to up to 13 years of

recurrent drought) seasonal synchrony becomes increasingly pronounced. In consequence,

multiple droughts may hamper the stability (de Mazancourt *et al.* 2013; Loreau & de

Mazancourt 2013), e.g. under future disturbances. This is because synchronous

communities are typically less stable to disturbances than asynchronous communities (Luo

et al. 2023b; Muraina *et al.* 2021; Craven *et al.* 2018; de Mazancourt *et al.* 2013). Hence, an

increase in seasonal synchrony with drought frequency suggests that multiple droughts may

hamper the stability of communities to future disturbances. We suggest that with the ongoing

and projected increases in the frequency of drought events due to climate change (IPCC

2021), grassland ecosystems become increasingly seasonally synchronous and vulnerable

to future disturbances.

We identified several underlying mechanisms that contributed to the increase in

synchrony with increasing drought frequency. First, the enhancement in synchrony was

associated with a **decrease in plant species richness**. This drought-induced species loss

aligns with findings from other studies that observed enhanced mortality during and following

single (Sippel *et al.* 2018; Stampfli *et al.* 2018; Frank *et al.* 2015), and recurrent drought

events (Luo *et al.* 2023a; Wagg *et al.* 2017). Species richness mostly tends to be negatively

associated with synchrony (Zhang *et al.* 2018; Hautier *et al.* 2014), but was also found to be

positively associated with synchrony (Valencia *et al.* 2020). Furthermore, the species that

remained in the community, were more similar in terms of their seasonal development.

Under increasing drought frequency, communities were dominated by species growing later

in the season, as **early-growing species were lost**. This may have been caused by a

reduced recovery potential of early-season species after the drought event: We found that

early-seasonal species were taking a longer time to recover under single- and three-year

drought. While it was suggested, that early-season species may be more susceptible to

drought (Mamolos *et al.* 2001), we found first evidence that early-growing species actually

get lost at high drought frequencies. As a result, under recurrent drought, more temporally

similar growing species lead to more seasonally synchronized communities.

Furthermore, the seasonal synchronization of species may have been driven by the

observed shift within plant **functional groups** under higher drought frequency. We found

that the community composition showed a reduction in forb and legume species, and more

grass species with increasing drought frequency. The loss of species within these functional

groups with increasing drought frequency may be explained by the observed lower forb and

legume seedlings throughout the season and the slower recovery of forbs and legumes post

drought. Hence, our findings support the notion that forb and legume species are less robust

to recurrent drought compared with grass species. This is in line with studies on single

drought (DeBoeck *et al.* 2018; Hoover *et al.* 2014) and recurrent drought events (Luo *et al.*

2023b; Luo *et al.* 2023a; Sun *et al.* 2023; Xu *et al.* 2021), which also suggest grass species

to be less sensitive to drought compared to forbs. Overall, the observed shifts under

recurrent drought towards less diversity in functional groups can enhance synchrony by

increasing the similarity of seasonal growth within the community. This similarity in functional

groups can furthermore decrease the variety in responses to drought throughout the growing

season (Zhang *et al.* 2020) and thereby contribute to an enhanced vulnerability under future

disturbances.

The increase in synchrony with higher drought frequency may also be driven by the

constrained establishment of new species and individuals throughout the growing

season: As drought frequency increased, we observed a **decline in seedling numbers**.

This occurred during the entire vegetation period and was most pronounced with high

drought frequency of 13 years. At the same time, less soil surface area was covered by

leaves under high drought frequency. Hence, the seedling decline took place despite a

diminished light limitation, and thus likely a higher chance of seedling survival in the 13-year

recurrent drought plots (Eskelinen *et al.* 2022). Moreover, we observed that plant species

reproductive output was inhibited with increasing drought frequency: In the recovery

phase, the fraction of flower biomass was decreased with increasing drought frequency. A

negative drought effect on reproductive output was also found by Zeiter *et al.* (2016).

Reasons might be, that species experiencing drought were unable to produce flowers due to

preceding and ongoing water limitation. Consequently, in the communities with these high

drought frequencies, fewer seedlings appeared in the recovery phase.

Overall, in our study, flower development and consequently seedlings, were severely

inhibited with higher drought frequencies. This reduces the establishment of new species

and individuals and leads to species loss, with effects on community composition and

synchrony. During long-term recurrent drought, effects of drought on reduced plant

reproductive output and resultant seedlings likely exacerbated plant species loss and the

increase in seasonal synchrony. However, single- and low-frequency experiments do not

cover drought effects that go beyond the lifetime of the plant species. We therefore propose

that, to comprehensively comprehend changes in plant communities arising from high

drought frequency, the implementation of long-term experiments is essential.

Enhanced loss of biomass fraction with increasing drought frequency

Drought did not reduce productivity in our study (Fig S2), which is consistent with findings on

productivity under five years of drought by Jentsch *et al.* (2011) and shoot biomass under

four years of drought by Fuchslueger *et al.* (2016). Nonetheless, other low drought frequency

studies reported a continuous decline in aboveground net primary productivity under

recurrent drought of up to four years (Luo *et al.* 2023a; Xu *et al.* 2021). The divergent results

may be attributed to the different frequencies of drought events or plant communities.

However, we found that increasing drought frequency gradually decreased the fraction of

plant community biomass and enhanced the fraction of necromass at peak drought and

recovery, respectively. Hence, during the growing season, including drought and recovery,

the maintained productivity is lower, and senescence is higher with increasing drought

frequency. The higher senescence at peak drought and recovery supports our observation of

increasing species loss throughout the growing season. Reductions in maintained

productivity might be caused by the lasting repercussions of drought on plant community

composition and synchrony.

Under recurrent drought, plants have the capacity to develop **adaptations** that can

dampen drought effects (Müller & Bahn 2022). This can occur e.g., by adjusting

photosynthesis, osmosis (Li *et al.* 2022), root biomass (Legay *et al.* 2018), epigenetics,

transcription factors and rate (Jacques *et al.* 2021; Ding *et al.* 2012; Bruce *et al.* 2007), as

well as transgenerational changes (Chen *et al.* 2022). Within our study, we found only weak

species adaptations under recurrent drought. For example, increasing drought frequency did

not result in adaptations of the duration of the growing season and only shifted the timing of

growth for one legume species to earlier in the season. Also, the timing of plant species

flowering was only shifted earlier with increasing drought frequency for one forb species.

Moreover, the lower SLA at peak drought of some of the most abundant forb species and

one legume species with increasing drought frequency can hint towards higher water-use

efficiency, resource conservation, and reduced plant growth, which can enhance short-term

stress tolerance (Wellstein *et al.* 2017). This short-term adaptation could lead to less

vulnerability under drought. Overall, adaptations did not play a major role in compensating

drought effects in our study.

Chen *et al.* (2022) suggested that adaptations *along generations* under recurrent

drought lead to improved ecosystem responses. However, we suggest that the lower

reproduction might overrule any potential adaptive effects of single plant species along

generations through species loss. Plant communities are therefore likely not able to adapt to

the predicted rapid increase in drought frequency with the ongoing change in climate. To

gain further understanding of the repercussions on plant community due to increasing

drought frequencies, long-term drought experiments, including transgenerational effects,

could yield important insights.

5. Conclusion

We show that long-term annual drought frequency in grasslands increases plant community

seasonal synchrony. Synchrony increased through a reduction in species richness, a loss of

early-seasonal plant species, a shift in plant functional groups, and the constrained

establishment of new species and individuals throughout the growing season. These

changes were associated with a decreased fraction of biomass with increasing drought

frequency.

Overall, we conclude that with the increase in frequency and severity of drought events due

to climate change, grassland plant communities will increase in seasonal synchrony. As

asynchrony is a critical component of community stability, we suggest that this will lead to

lasting effects far beyond the drought event, by increasing the vulnerability of ecosystems to

future disturbances. Moreover, we show that negative drought effects were enhanced with

an increasing in drought frequency. We therefore conclude that single and low-frequency

drought studies may not adequately predict longer-term changes in our rapidly shifting

climate. Furthermore, on the long-term, negative effects, mostly associated with species

loss, will likely overrule the comparatively weak adaptation of species. Hence, with the

current speed of climate change, communities will be severely and probably irrevocably

affected the moment species are lost from the ecosystem. We therefore conclude that plant

communities are unlikely to be able to adapt to recurrent drought. Importantly, long-term

experiments will be essential in understanding the severity of these climate change impacts.

6.   Sample CRediT author statement

Lena M. Müller: Conceptualization, Investigation, Project administration, Writing - Original

Draft (lead), Visualization, Funding acquisition; **Michael Bahn**: Conceptualization,

Resources, Funding acquisition, Writing - Review & Editing; **Maximillian Weidle**:

Investigation, Writing - Review & Editing; **Georg Leitinger**: Writing - Review & Editing,

Funding acquisition; **Dina in 't Zandt**: Conceptualization, Investigation, Formal Analysis,

Data Curation, Visualization, Writing - Review & Editing, Writing - Original Draft, Supervision

7.   Acknowledgements

We are grateful to numerous colleagues, students, and friends who have helped to maintain

the long-term experiment over the years, as well as to those that assisted in setting up and

conducting the sampling for the experiment presented in this study. Amongst others, we

thank Beverley Anderson, Mario Deutschmann, Andrew Giunta, Roland Hasibeder,

Johannes Ingrisch, Florian Oberleitner, Natalie Oram, David Reinthaler, Michael Schmitt,

Carmen Telser, and Herbert Wachter.

The study site is part of the LTSER platform Tyrolean Alps, belonging to the national and

international long term ecological research networks (LTER-Austria, LTER Europe and

ILTER). The work was funded by the Forschungszentrum Berglandwirtschaft (project no.

326572). The long-term drought treatment was maintained by projects funded by the

Austrian Science Fund (FWF, project no. P22214-B17, P31132, and I1056) and the Austrian

Academy of Sciences (ESS-programme, project ClimLUC).

LM was supported by the University Innsbruck through a doctoral scholarship

(Doktoratsstipendium aus der Nachwuchsförderung der Universität Innsbruck) and partially

funded by the Austrian Science Fund (FWF): [I4969-B]. DZ was supported by a PPLZ

Postdoctoral Fellowship awarded by the Czech Academy of Sciences (RVO 67985939).

8.   Data Availability Statement

Upon acceptance, all data and code (R scripts) will be made publicly available via the

Zenodo data repository, will be referenced using a stable object identifier (DOI) in the

Materials and Methods section, and cited in the reference list.

9. Conflict of interest

The authors declare no conflict of interest.

References

Backhaus S, Kreyling J, Grant K, Beierkuhnlein C, Walter J, Jentsch A (2014) Recurrent mild drought events increase resistance toward extreme drought stress. *Ecosystems*, **17**, 1068–1081.

Bardgett RD, Bullock JM, Lavorel S *et al.* (2021) Combatting global grassland degradation. *Nature Reviews Earth & Environment*, **2**, 720–735.

Bardgett RD, Caruso T (2020) Soil microbial community responses to climate extremes: resistance, resilience and transitions to alternative states. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **375**, 20190112.

Basto S, Thompson K, Grime JP, Fridley JD, Calhim S, Askew AP, Rees M (2018) Severe effects of long-term drought on calcareous grassland seed banks. *npj Climate and Atmospheric Science*, **1**, 5.

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, **67**, 1–48.

Berwaers S, DeBoeck HJ, Nijs I (2019) End-of-season senescence in grassland species can be traced to leaf temperature during preceding summer drought. *Perspectives in Plant Ecology, Evolution and Systematics*, **38**, 31–38.

Blair J, Nippert J, Briggs J (2014) Grassland Ecology. In: *Ecology and the Environment* (ed Monson RK), pp 389–423. Springer New York.

Bruce TJ, Matthes MC, Napier JA, Pickett JA (2007) Stressful "memories" of plants: Evidence and possible mechanisms. *Plant Science*, **173**, 603–608.

Canarini A, Schmidt H, Fuchslueger L *et al.* (2021) Ecological memory of recurrent drought modifies soil processes via changes in soil microbial community. *Nature Communications*, **12**, 5308.

Castillioni K, Newman GS, Souza L, Iler AM (2022) Effects of drought on grassland phenology depend on functional types. *New Phytologist*, **236**, 1558–1571.

Chen Y, Vogel A, Wagg C *et al.* (2022) Drought-exposure history increases complementarity between plant species in response to a subsequent drought. *Nature Communications*, **13**, 3217.

Coleman MA, Wernberg T (2020) The silver lining of extreme events. *Trends in Ecology & Evolution*, **35**, 1065–1067.

Craven D, Eisenhauer N, Pearse WD *et al.* (2018) Multiple facets of biodiversity drive the diversity-stability relationship. *Nature Ecology & Evolution*, **2**, 1579–1587.

de Mazancourt C, Isbell F, Larocque A *et al.* (2013) Predicting ecosystem stability from community composition and biodiversity. *Ecology Letters*, **16**, 617–625.

DeBoeck HJ, Hiltbrunner E, Verlinden M, Bassin S, Zeiter M (2018) Legacy effects of climate extremes in alpine grassland. *Frontiers in Plant Science*, **9**, 1586.

Ding Y, Fromm M, Avramova Z (2012) Multiple exposures to drought 'train' transcriptional responses in Arabidopsis. *Nature Communications*, **3**, 740.

Dixon AP, Faber-Langendoen D, Josse C, Morrison J, Loucks CJ (2014) Distribution mapping of world grassland types. *Journal of Biogeography*, **41**, 2003–2019.

Eskelinen A, Harpole WS, Jessen M-T, Virtanen R, Hautier Y (2022) Light competition drives herbivore and nutrient effects on plant diversity. *Nature*, **611**, 301–305.

Forrest J, Miller-Rushing AJ (2010) Toward a synthetic understanding of the role of phenology in ecology and evolution. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **365**, 3101–3112.

Frank D, Reichstein M, Bahn M *et al.* (2015) Effects of climate extremes on the terrestrial carbon cycle: Concepts, processes and potential future impacts. *Global Change Biology*, **21**, 2861–2880.

Fuchslueger L, Bahn M, Hasibeder R, Kienzl S, Fritz K, Schmitt M, Watzka M, Richter A (2016) Drought history affects grassland plant and microbial carbon turnover during and after a subsequent drought event. *Journal of Ecology*, **104**, 1453–1465.

Ge C, Sun S, Yao R, Sun P, Li M, Bian Y (2022) Long-term vegetation phenology changes and response to multi-scale meteorological drought on the Loess Plateau, China. *Journal of Hydrology*, **614**, 128605.

Gibbs HK, Salmon JM (2015) Mapping the world's degraded lands. *Applied Geography*, **57**, 12–21.

Griffin-Nolan RJ, Carroll CJ, Denton EM, Johnston MK, Collins SL, Smith MD, Knapp AK (2018) Legacy effects of a regional drought on aboveground net primary production in six central US grasslands. *Plant Ecology*, **219**, 505–515.

Hahn C, Lüscher A, Ernst-Hasler S, Suter M, Kahmen A (2021) Timing of drought in the growing season and strong legacy effects determine the annual productivity of temperate grasses in a changing climate. *Biogeosciences*, **18**, 585–604.

Hallett LM, Jones SK, MacDonald AA *et al.* (2016) codyn : An r package of community dynamics metrics. *Methods in Ecology and Evolution*, **7**, 1146–1151.

Hartig F DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models. Available via http://florianhartig.github.io/DHARMa/.

Hasibeder R, Fuchslueger L, Richter A, Bahn M (2015) Summer drought alters carbon allocation to roots and root respiration in mountain grassland. *New Phytologist*, **205**, 1117–1127.

Hautier Y, Seabloom EW, Borer ET *et al.* (2014) Eutrophication weakens stabilizing effects of diversity in natural grasslands. *Nature*, **508**, 521–525.

He YL, Wang JS, Tian DS *et al.* (2022) Long-term drought aggravates instability of alpine grassland productivity to extreme climatic event. *Ecology*, **103**, 1-12.

Herben T, Březina S, Hadincová V, Krahulec F, Skálová H (2013) Mutual replacement of species in space in a grassland community: is there an evidence for functional complementarity of replacement groups? *Oikos*, **122**, 111–121.

Herben T, Krahulec F, Hadincová V, Pecháčková S, Kováčiová M (1997) Fine-scale spatio-temporal patterns in a mountain grassland: do species replace each other in a regular fashion? *Journal of Vegetation Science*, **8**, 217–224.

Hofer D, Suter M, Haughey E, Finn JA, Hoekstra NJ, Buchmann N, Lüscher A (2016) Yield of temperate forage grassland species is either largely resistant or resilient to experimental summer drought. *Journal of Applied Ecology*, **53**, 1023–1034.

Hoover DL, Knapp AK, Smith MD (2014) Resistance and resilience of a grassland ecosystem to climate extremes. *Ecology*, **95**, 2646–2656.

Hoover DL, Pfennigwerth AA, Duniway MC (2021) Drought resistance and resilience: the role of soil moisture-plant interactions and legacies in a dryland ecosystem. *Journal of Ecology*, **109**, 3280–3294.

in 't Zandt D, Herben T, Brink A, Visser EJ, Kroon H (2021) Species abundance fluctuations over 31 years are associated with plant–soil feedback in a species-rich mountain meadow. *Journal of Ecology*, **109**, 1511–1523.

Ingrisch J, Karlowsky S, Anadon-Rosell A, Hasibeder R, König A, Augusti A, Gleixner G, Bahn M (2018) Land Use Alters the Drought Responses of Productivity and CO2 Fluxes in Mountain Grassland. *Ecosystems*, **21**, 689–703.

Ingrisch J, Karlowsky S, Hasibeder R, Gleixner G, Bahn M (2020) Drought and recovery
effects on belowground respiration dynamics and the partitioning of recent carbon in
managed and abandoned grassland. *Global Change Biology*, **26**, 4366–4378.
IPCC (2021) Climate Change 2021. The Physical Science Basis. Contribution of Working
Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate
Change. Cambridge, UK: Cambridge University Press.
IPCC Climate Change 2023. Synthesis Report. Contribution of Working Groups I, II and III to
the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core
Writing Team, H. Lee and J. Romero (eds.)]., IPCC, Geneva, Switzerland.
Jacques C, Salon C, Barnard RL, Vernoud V, Prudent M (2021) Drought Stress Memory at
the Plant Cycle Level: A Review. *Plants*, **10**, 1873.
Jentsch A, Kreyling J, Boettcher-Treschkow J, Beierkuhnlein C (2009) Beyond gradual
warming: Extreme weather events alter flower phenology of European grassland and
heath species. *Global Change Biology*, **15**, 837–849.
Jentsch A, Kreyling J, Elmer M *et al.* (2011) Climate extremes initiate ecosystem-regulating
functions while maintaining productivity. *Journal of Ecology*, **99**, 689–702.
Ji S, Ren S, Li Y, Dong J, Wang L, Quan Q, Liu J (2021) Diverse responses of spring
phenology to preseason drought and warming under different biomes in the North China
Plain. *The Science of the Total Environment*, **766**, 144437.
Kang W, Wang T, Liu S (2018) The response of vegetation phenology and productivity to
drought in semi-arid regions of northern China. *Remote Sensing*, **10**, 727.
Karlowsky S, Augusti A, Ingrisch J, Akanda MK, Bahn M, Gleixner G (2018) Drought-
induced accumulation of root exudates supports post-drought recovery of microbes in
mountain grassland. *Frontiers in Plant Science*, **9**, 1593.
Legay N, Piton G, Arnoldi C *et al.* (2018) Soil legacy effects of climatic stress, management
and plant functional composition on microbial communities influence the response of
Lolium perenne to a new drought event. *Plant and Soil*, **424**, 233–254.
Li X, Jimoh SO, Li Y, Duan J, Cui Y, Jin K, Wang Z, Zhang Y (2022) Stress memory and
phyllosphere/soil legacy underlie tolerance and plasticity of Leymus chinensis to periodic
drought risk. *Agricultural and Forest Meteorology*, **312**, 108717.
Loreau M, de Mazancourt C (2008) Species synchrony and its drivers: neutral and
nonneutral community dynamics in fluctuating environments. *The American naturalist*,
172, E48-66.
Loreau M, de Mazancourt C (2013) Biodiversity and ecosystem stability: a synthesis of
underlying mechanisms. *Ecology letters*, **16**, 106–115.
Luo W, Muraina TO, Griffin-Nolan RJ *et al.* (2023a) Responses of a semiarid grassland to
recurrent drought are linked to community functional composition. *Ecology*, **104**, e3920.
Luo W, Shi Y, Wilkins K *et al.* (2023b) Plant traits modulate grassland stability during drought
and post-drought periods. *Functional Ecology*, **37**, 2611–2620.
Mackie KA, Zeiter M, Bloor JM, Stampfli A (2019) Plant functional groups mediate drought
resistance and recovery in a multisite grassland experiment. *Journal of Ecology*, **107**,
937–949.
Magnusson A, Skaug HJ, Nielsen A *et al.* (2017) glmmTMB: Generalised Linear
MixedModels using Template Model Builder. https://github.com/glmmTMB.
Mamolos AP, Veresoglou DS, Noitsakis V, Gerakis A (2001) Differential drought tolerance of
five coexisting plant species in Mediterranean lowland grasslands. *Journal of Arid*
Environments, **49**, 329–341.

Müller LM, Bahn M (2022) Drought legacies and ecosystem responses to subsequent drought. *Global Change Biology*, **28**, 5086–5103.

Muraina TO, Xu C, Yu Q *et al.* (2021) Species asynchrony stabilises productivity under extreme drought across Northern China grasslands. *Journal of Ecology*, **109**, 1665–1675.

Oksanen J, Blanchet FG, Friendly M *et al.* (2020) vegan: Community Ecology Package. https://CRAN.R-project.org/package=vegan.

Oram NJ, Ingrisch J, Bardgett RD *et al.* (2023) Drought intensity alters productivity, carbon allocation and plant nitrogen uptake in fast versus slow grassland communities. *Journal of Ecology*, **111**, 1681–1699.

Peng J, Wu C, Zhang X, Wang X, Gonsamo A (2019) Satellite detection of cumulative and lagged effects of drought on autumn leaf senescence over the Northern Hemisphere. *Global Change Biology*, **25**, 2174–2188.

Petermann JS, Buzhdygan OY (2021) Grassland biodiversity. *Current Biology*, **31**, R1195-R1201.

Qian J, Guo Z, Muraina TO *et al.* (2022) Legacy effects of a multi-year extreme drought on belowground bud banks in rhizomatous vs bunchgrass-dominated grasslands. *Oecologia*, **198**, 763–771.

R Core Team (2021) A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

Rihan W, Zhao J, Zhang H, Guo X (2022) Preseason drought controls on patterns of spring phenology in grasslands of the Mongolian Plateau. *The Science of the Total Environment*, **838**, 156018.

Schirpke U, Kohler M, Leitinger G, Fontana V, Tasser E, Tappeiner U (2019) Future impacts of changing land-use and climate on ecosystem services of mountain grassland and their resilience. *Ecosystem services*, **26**, 79–94.

Schmitt M, Bahn M, Wohlfahrt G, Tappeiner U, Cernusca A (2010) Land use affects the net ecosystem CO_2 exchange and its components in mountain grasslands. *Biogeosciences*, **7**, 2297–2309.

Schwalm CR, Anderegg WR, Michalak AM *et al.* (2017) Global patterns of drought recovery. *Nature*, **548**, 202–205.

Sippel S, Reichstein M, Ma X, Mahecha MD, Lange H, Flach M, Frank D (2018) Drought, heat, and the carbon cycle: A review. *Current Climate Change Reports*, **4**, 266–286.

Smith MD, Wilkins KD, Holdrege MC *et al.* (2023) Extreme drought impacts have been underestimated in grasslands and shrublands globally. *Proceedings of the National Academy of Sciences*, in press.

Smith MD, Wilkins KD, Holdrege MC *et al.* (2024) Extreme drought impacts have been underestimated in grasslands and shrublands globally. *Proceedings of the National Academy of Sciences*, **121**, e2309881120.

Spinoni J, Vogt JV, Naumann G, Barbosa P, Dosio A (2018) Will drought events become more frequent and severe in Europe? *International Journal of Climatology*, **38**, 1718–1736.

Stampfli A, Bloor JM, Fischer M, Zeiter M (2018) High land-use intensity exacerbates shifts in grassland vegetation composition after severe experimental drought. *Global Change Biology*, **24**, 2021–2034.

Stampfli A, Zeiter M (2004) Plant regeneration directs changes in grassland composition after extreme drought: a 13-year study in southern Switzerland. *Journal of Ecology*, **92**, 568–576.

Sun J, Zhang B, Pan Q *et al.* (2023) Non-linear response of productivity to precipitation extremes in the Inner Mongolia grassland. *Functional Ecology*, **37**, 1663–1673.

Tello-García E, Huber L, Leitinger G, Peters A, Newesely C, Ringler M-E, Tasser E (2020) Drought- and heat-induced shifts in vegetation composition impact biomass production and water use of alpine grasslands. *Environmental and Experimental Botany*, **169**, 103921.

Valencia E, Bello F de, Galland T *et al.* (2020) Synchrony matters more than species richness in plant community stability at a global scale. *Proceedings of the National Academy of Sciences*, **117**, 24345–24351.

Vorkauf M, Kahmen A, Körner C, Hiltbrunner E (2021) Flowering phenology in alpine grassland strongly responds to shifts in snowmelt but weakly to summer drought. *Alpine Botany*, **131**, 73–88.

Wagg C, O'Brien MJ, Vogel A, Scherer-Lorenzen M, Eisenhauer N, Schmid B, Weigelt A (2017) Plant diversity maintains long-term ecosystem productivity under frequent drought by increasing short-term variation. *Ecology*, **98**, 2952–2961.

Walter J, Nagy L, Hein R, Rascher U, Beierkuhnlein C, Willner E, Jentsch A (2011) Do plants remember drought? Hints towards a drought-memory in grasses. *Environmental and Experimental Botany*, **71**, 34–40.

Wang M, Li P, Peng C, Xiao J, Zhou X, Luo Y, Zhang C (2022) Divergent responses of autumn vegetation phenology to climate extremes over northern middle and high latitudes. *Global Ecology and Biogeography*, **31**, 2281–2296.

Wang T, Tu X, Singh VP, Chen X, Lin K (2021) Global data assessment and analysis of drought characteristics based on CMIP6. *Journal of Hydrology*, **596**, 126091.

Wellstein C, Poschlod P, Gohlke A *et al.* (2017) Effects of extreme drought on specific leaf area of grassland species: A meta-analysis of experimental studies in temperate and sub-Mediterranean systems. *Global Change Biology*, **23**, 2473–2481.

Wilcox KR, Blumenthal DM, Kray JA, Mueller KE, Derner JD, Ocheltree T, Porensky LM (2021) Plant traits related to precipitation sensitivity of species and communities in semiarid shortgrass prairie. *New Phytologist*, **229**, 2007–2019.

Wilcox KR, Tredennick AT, Koerner SE *et al.* (2017) Asynchrony among local communities stabilises ecosystem function of metacommunities. *Ecology Letters*, **20**, 1534–1545.

Xu C, Ke Y, Zhou W *et al.* (2021) Resistance and resilience of a semi-arid grassland to multi-year extreme drought. *Ecological Indicators*, **131**, 108139.

Xu Z, Ren H, Li M-H *et al.* (2015) Environmental changes drive the temporal stability of semi-arid natural grasslands through altering species asynchrony. *Journal of Ecology*, **103**, 1308–1316.

Yuan M, Zhao L, Lin A, Wang L, Li Q, She D, Qu S (2020) Impacts of preseason drought on vegetation spring phenology across the Northeast China Transect. *The Science of the Total Environment*, **738**, 140297.

Yuan Z, Tong S, Bao G, Chen J, Yin S, Li F, Sa C, Bao Y (2023) Spatiotemporal variation of autumn phenology responses to preseason drought and temperature in alpine and temperate grasslands in China. *The Science of the Total Environment*, **859**, 160373.

Zeiter M, Sara Schärrer, Roman Zweifel, David M. Newbery, Stampfli A (2016) Timing of extreme drought modifies reproductive output in semi-natural grassland. *Journal of Vegetation Science*, **27**, 238–248.

Zeng Z, Wu W, Ge Q *et al.* (2021) Legacy effects of spring phenology on vegetation growth under preseason meteorological drought in the Northern Hemisphere. *Agricultural and Forest Meteorology*, **310**, 108630.

Zhang F, Quan Q, Ma F, Tian D, Hoover DL, Zhou Q, Niu S (2019) When does extreme
drought elicit extreme ecological responses? *Journal of Ecology*, **107**, 2553–2563.
Zhang J, Miao Y, Zhang T *et al.* (2020) Drought timing and primary productivity in a semiarid
grassland. *Land Degradation & Development*, 2185–2195.
Zhang Y, Loreau M, He N, Wang J, Pan Q, Bai Y, Han X (2018) Climate variability
decreases species richness and community stability in a temperate grassland. *Oecologia*,
188, 183–192.
Zuur AF, Ieno EN, Elphick CS (2010) A protocol for data exploration to avoid common
statistical problems. *Methods in Ecology and Evolution*, **1**, 3–14.