Recurrent drought increases grassland community seasonal synchrony

Lena M. Müller[1], Michael Bahn[1], Maximillian Weidle[1], Georg Leitinger[1], Dina in 't Zandt[1,2]

[1]Department of Ecology, Universität Innsbruck, Innsbruck, Austria

[2]Department of Population Ecology, Institute of Botany, The Czech Academy of Sciences, Průhonice, Czech Republic

Abstract

1. Climate change increases the frequency and severity of drought events with strong repercussions on grassland ecosystems. While the effects of single drought events on ecosystem structure and functioning are well understood, it is largely unknown whether and how drought frequency modifies ecosystem responses to drought.

2. Here, we assessed how the increase in frequency of severe, annual summer drought impacted grassland communities. We examined these effects in a species-rich sub-alpine mountain meadow with a drought frequency of one, three, and 13 years, as well as ambient conditions.

3. We found that high drought frequency increased seasonal plant community synchrony through a reduction in species richness, a shift of plant functional groups, a loss of early-seasonal plant species, and the constrained establishment of seedlings throughout the growing season. These changes were associated with a decreased fraction of biomass as drought frequency increased.

4. Furthermore, we show that negative drought effects were enhanced with an increasing drought frequency, and that negative drought effects on plant communities outweighed the weak adaptive effects of species.

5. Synthesis. We conclude that single and low-frequency drought studies may not adequately predict longer-term changes in our rapidly shifting climate. With the ongoing increase in drought frequency due to climate change, we predict that grassland plant communities will increase in seasonal synchrony. We suggest that this increase in synchrony will leave ecosystems highly vulnerable to future disturbances, because asynchrony is a critical

component of stability. Moreover, given the weak adaptive effects of plant species to long-term recurrent drought, we conclude that plant communities are unlikely to be able to adapt to the rapid increase in recurrent drought events.

Keywords: recurrent drought, drought frequency, seasonal synchrony, synchrony, asynchrony, grassland, species richness, early-seasonal species, functional groups, drought

Introduction

Grasslands cover 40 % of the Earth's surface and are crucial for carbon and nutrient cycling, biodiversity conservation, and ecosystem services (Petermann & Buzhdygan 2021; Blair *et al.* 2014; Dixon *et al.* 2014). However, up to 49 % of the world's grasslands are currently degraded, with the climate crisis predicted to exacerbate this number (Bardgett *et al.* 2021; Gibbs & Salmon 2015). Among the detrimental climate scenarios, the increase in frequency and severity of drought events stand out as particularly damaging (IPCC 2023, 2021; Wang *et al.* 2021; Spinoni *et al.* 2018). Drought events alter grassland nutrient cycling, disrupt interactions between species, result in species extinction, and therewith cause severe disruptions in ecosystem structure and function (Smith *et al.* 2024; Oram *et al.* 2023; Bardgett & Caruso 2020; Ingrisch *et al.* 2020; Tello-García *et al.* 2020). In the long-term, the predicted increase in the frequency of drought events therefore threatens to irrevocably change grassland ecosystems, including the fundamental services and biodiversity they conserve (Müller & Bahn 2022; Xu *et al.* 2021; Schirpke *et al.* 2019). At the same time, potential adaptations of species and communities may dampen these detrimental effects to some extent (Müller & Bahn 2022; Canarini *et al.* 2021; Coleman & Wernberg 2020). This occurs for example through drought-induced legacy effects that alter ecosystem structure and function after drought has subsided, and which results in plant communities being able to cope better with a subsequent drought event (Müller & Bahn 2022). Yet, while the effects of single drought events are comparatively well studied, our understanding of the effects of a long-term increase in drought frequency on grassland community structure and function remains limited.

Multiple interlinked drought-induced changes, for example in plant species composition, diversity, complementarity, as well as species loss may have detrimental effects by affecting the **synchrony** of grassland communities (He *et al.* 2022; Muraina *et al.* 2021; De Mazancourt *et al.* 2013). Community synchrony refers to the degree to which populations of different species within a community fluctuate in tandem. High community synchrony indicates that populations of different species increase and decrease at the same points in time, while low synchrony (or asynchrony) indicates that species populations fluctuate independently of each other (Loreau & De Mazancourt 2013). With greater synchrony between years, species communities exhibit increased similarity in their response to environmental changes, the

swiftness of their responses, or their functional complementarity throughout various seasons

(Loreau & De Mazancourt 2013). Single drought was found to increase the synchrony of plant

communities between years (Zhang *et al.* 2019), similarly as recurrent drought of three (He *et*

al. 2022) and four years (Muraina *et al.* 2021). These changes in grassland community

synchrony likely have detrimental effects on long-term ecological functioning, especially

under a future climate with a high drought frequency. Increases in grassland community

synchrony between years has been shown to decrease community stability (Wilcox *et al.* 2017;

Xu *et al.* 2015; Loreau & De Mazancourt 2013). This is because asynchronous communities

stabilise ecosystems through different responses to disturbances (De Mazancourt *et al.* 2013).

For drought disturbances, species asynchrony was suggested to be the dominant driver to

stabilise community productivity under drought (Luo *et al.* 2023b; Muraina *et al.* 2021; Craven

et al. 2018). Community synchrony between years is thus influenced by drought and is

suggested to shape a communities' response to subsequent droughts by decreasing its

stability. However, besides changing community synchrony *between* seasons, drought is also

likely to affect community synchrony *within* the growing season. Therewith, it might be a

critical mechanism in shaping responses to recurrent drought. However, we currently lack

evidence on the effects of recurrent drought on community *seasonal* synchrony. In the presence

of more frequent droughts, we hypothesize that the plant community will become more

synchronous within the season.

Drought events alter plant community composition, phenology, and productivity (Smith *et al.*

2024; Müller & Bahn 2022). Drought leads to a decrease in plant community **productivity**, by

limiting plant growth through reduced water availability, or by mortality of species under

severe drought conditions (Sippel *et al.* 2018; Frank *et al.* 2015). Additionally, after a drought

event, soil rewetting results in a temporarily increased availability of nitrogen (Ingrisch *et al.*

2018; Karlowsky *et al.* 2018). This surge in nitrogen supports the recovery and, in some cases,

leads to a short-term overshoot in productivity (Griffin-Nolan *et al.* 2018; Hofer *et al.* 2016),

which can compensate for the lower plant community productivity during the drought event

(Hahn *et al.* 2021). This response is mainly based on fast-growing grass species, which makes

the community sensitive to changes in composition post drought (Mackie *et al.* 2019; Stampfli

et al. 2018). An important issue is that drought-induced changes in productivity and

composition in grasslands may take a long time to come back to pre-disturbance level (Müller

& Bahn 2022). If a subsequent drought falls within this recovery timespan of a previous
drought, it may lead to constantly impaired ecosystems (Schwalm *et al.* 2017), and
consequently to a more profound effect on productivity under a high frequency of drought
(Xu *et al.* 2021). Hence, a potential accumulation of effects can emerge when multiple droughts
occur annually. However, as recurrent drought events are typically studied in two-year
experiments and only rarely extend beyond this timeframe, we lack critical knowledge on the
productivity loss that builds up under a high drought frequency (Knapp *et al.* 2023; Hoover *et*
al. 2018).

Drought shifts plant community **composition** towards more slower growing and more
drought adapted-species (Wilcox *et al.* 2021). For example, drought may lead to a dominance
of grasses over forbs (Xu *et al.* 2021; De Boeck *et al.* 2018; Hoover *et al.* 2014). This may be
driven by a higher post-drought resource use of grasses compared with the conservative
resource use of forbs (Stampfli *et al.* 2018) or by a more efficient resource acquisition of water
of grasses compared with forbs (Tello-García *et al.* 2020). Furthermore, drought-induced
changes in plant community composition are driven by a reduction in plant species
performance and an increase in mortality of plants both during and after a drought event
(Sippel *et al.* 2018; Frank *et al.* 2015). In the long term, plant compositional shifts due to high
drought frequency are likely to relate additionally to drought-induced changes in plant
reproductive effects, e.g. reduction in reproductive shoots, seeds (Zeiter *et al.* 2016),
belowground bud density (Qian *et al.* 2022), as well as altered seed banks (Basto *et al.* 2018),
and regeneration from seed (Stampfli & Zeiter 2004). Through these drought-induced changes
in plant species composition, recurrent drought events may in the long-term result in
permanent loss of plant species with potentially irreversible consequences for the functioning
of plant communities. Furthermore, under recurrent drought, plant species also have the
capacity to adapt (Müller & Bahn 2022), which dampens drought effects under a higher
drought frequency (Backhaus *et al.* 2014; Walter *et al.* 2011). Therefore, a high drought
frequency is expected to result in plant communities consisting of drought-tolerant species, a
dominance of grasses, plant species profiting from the recovery dynamics of drought events,
as well as plant species that adapt to recurrent drought conditions. However, due to the lack
of studies investigating high drought frequency, long-term compositional shifts might have
been overlooked to date.

Drought alters **phenology** by shifting the beginning or end of the growing season (Yuan *et al.*
2023; Ge *et al.* 2022; Rihan *et al.* 2022; Wang *et al.* 2022; Ji *et al.* 2021; Zeng *et al.* 2021; Yuan *et al.*
2020; Berwaers *et al.* 2019; Kang *et al.* 2018) and advancing end-of-season leaf senescence
(Hoover *et al.* 2021; Berwaers *et al.* 2019; Peng *et al.* 2019; Kang *et al.* 2018). Furthermore,
drought alters species flowering phenology by advancing the flowering date and extending
the flowering length, whereby effects differ strongly between species (Castillioni *et al.* 2022;
Vorkauf *et al.* 2021; Forrest & Miller-Rushing 2010; Jentsch *et al.* 2009). Hence, recurrent
drought is expected to move the duration and timing of the growing season and flowering
with effects on community composition, though the directions and magnitudes of these
changes under recurrent drought are still poorly understood.

We lack a profound understanding of the implications of a high frequency of drought events
on the temporal shifts in community composition, productivity, and phenology. Here, we
assessed how the increase in frequency of severe summer drought affected grassland
community structure, function, and seasonal dynamics. We examined this in a species-rich
sub-alpine mountain meadow with increasing annual summer drought frequencies of up to
13 years. We quantified species abundances in the field throughout the growing season and
sampled plant species-specific biomass and necromass at peak drought and drought recovery.
We asked the following questions: How did an increase in drought frequency affect (1) plant
community productivity, species richness, and flower production during peak drought and
recovery, respectively, as well as (2) plant community seasonal synchrony. We examined
whether these patterns were underlain by shifts in (3) seedling establishment (4), plant
community composition, (5) plant functional groups, and (6) plant community traits. Taken
together, we present the effects of a high frequency of drought events on plant communities
and discuss underlying mechanisms.

Material and Methods

Study Site

The studied sub-alpine grassland is located in the Austrian Central Alps in the Stubai valley at 1820 m a.s.l. (47°70 45″N, 11°180 20″E). The site is exposed southeast with an inclination of ~ 20° and has an average annual temperature of 3 °C and an annual precipitation of 1100 mm. The soil type is a dystric cambisol (top soil pH = 5.5). The vegetation community is classified as *Trisetetum flavescentis*, species-rich with on average 49 species per m² and consists largely of perennial grasses and forbs. The growing season lasts from the end of April until mid-October. The sub-alpine grassland is manured every 2-3 years, mown once in early August at peak biomass, and sporadically grazed in spring and autumn. The studied area, however, has not been fertilised and is excluded from grazing since 2007. For further details on the site see Ingrisch *et al.* (2018), Hasibeder *et al.* (2015), and Schmitt *et al.* (2010).

Experimental set-up

Four summer drought treatments were established in the enclosed area at the study site: a single year summer drought, three years of recurrent summer drought, 13 years of recurrent summer drought, and ambient conditions (n = 4, respectively). The treatment of 13 years of recurrent drought was established in 2008, the three-year recurrent drought in 2018, and the single-year drought in 2020. Summer drought was simulated for seven weeks from 17 June until 7 August 2020 using rainout shelters (Figure 1). This is the warmest period in the growing season and therefore the period where a natural drought is most likely to occur. The timing and intensity of the annual drought were kept as similar as possible in previous years since 2008. The ambient treatment was not covered with rainout shelters and thus exposed to ambient precipitation. The rainout shelters of the single and three years of recurrent summer drought had a base area of 3 x 3.5 m² and 2.5 m height. Each of these shelters contained a single 20 x 20 cm² plot. The two rainout shelters of the 13 years of recurrent drought were twice the size, resulting in a base area of 3 x 7 m² (shelter height was the same with 2.5 m). These larger shelters contained two 20 x 20 cm² plot, installed at either side of the shelter. In total, 4 plots for ambient, single, three, and 13 years of drought frequency were installed four weeks before summer drought simulation in 2020, respectively. Each plot was located at least

50 cm away from the rainout shelter to avoid edge effects. Rainout shelters with plots were

distributed in four blocks in a transect across the mountain meadow.

All shelters were covered with a transparent UV-A and UV-B transmissive plastic foil

(Lumisol clear AF, Folitec, Westerburg, Germany, light transmittance ca. 90 %). To allow air

circulation, the shelters were open at the bottom (up to 0.5 m) and at the top of the face sides.

The drought period was ended by removing the shelters and watering of all 20 x 20 cm² plots

with 31 mm of distilled water. At this time point, the entire site was also mown, and the

cuttings were removed.

Figure 1 The course of the experiment including setup and removal of the rainout shelters, and peak drought and

recovery harvest. The pictures show the plots exposed to ambient conditions, and to one year, three years, and 13

years of drought. All pictures were taken in early August at peak drought.

Measurements

Microclimate

A microclimate station at the study site recorded precipitation, air temperature, and relative

air humidity (see details in Hasibeder *et al.* (2015)). During the whole season, soil water

content (SWC) (Decagon EC-5 and 5TM, Decagon Devices, Pullman, WA, USA; SM300, Delta-

T Devices, Cambridge, UK; Hobo S-SMD-M005, Onset Computer corporation, Bourne, MA,

USA) was measured continuously in the main rooting horizon (30-min interval) at 5, 10, and

20 cm of soil depth. The measurements were replicated twice in the single-, and three-years

treatment, and three to four times in the ambient treatment. Due to partial sensor failures, no

data is available for the 13-years treatment. For results on the microclimate and SWC, see

Figure S1.

Measurements throughout the growing season

In all 16 plots, species composition and vegetation height were measured every two weeks

from 22 May to 22 October 2020. Plant species composition was estimated by noting for each

species present the number of modules and flowering stems. Modules refer to the number of

tillers for grasses and graminoids, the number of rosettes for small dicots, and the number of

leaves for larger dicots per rooted species, following (in 't Zandt *et al.* 2021; Herben *et al.* 2013;

Herben *et al.* 1997). Counting was accompanied by measurements of mean vegetation height

(in cm). Seedling composition in the plots was counted separately every three weeks between

the 23 June and the 22 October 2020. A seedling was defined as a plant individual with less

than two leaves, and was not part of the module counts. Seedlings were counted per plant

functional group: forbs, grasses, and legumes.

Biomass, necromass, and leaf traits

Aboveground biomass (living plant material) and necromass (dead plant material) was

sampled destructively in the plots at peak drought on 6 August 2020 and during recovery on

22 October 2020 (Figure 1). As the entire site is mown every summer at the same time point as

the peak drought sampling, there was no effect of peak drought biomass removal on recovery

sampling. For this, plant material was cut at 2 cm above the soil and frozen at -18 °C until

further analysis. Samples were split into plant species and separated into leaves, stems, and

flowers. This was done for biomass and necromass, respectively. For each species of each plot,

the leaf area was determined from thawed leaves that were saturated with water and scanned

(V700 Photo, Epson, WinRHIZO Pro 2012, Regent Instruments). Scanned leaves were dried in

the subsequent three days at 60 °C and were weighed. Specific leaf area (SLA) was calculated

by dividing the scanned leaf area by the dry weight of the leaves for each species and

treatment, respectively. Leaf area index (LAI) was calculated from the leaf surface of all leaves
divided by the total unit of ground area of the plot.

We also assessed drought frequency effects on aboveground phytomass fractions. These were
calculated as the proportion of biomass, necromass, and flower mass of the total aboveground
phytomass in each plot. We did this, because phytomass fractions were in most cases more
informative concerning drought frequency effects than absolute values of phytomass (Figure
S2) as well as biomass and necromass (data not shown). This is because, a substantial portion
of the phytomass at the subalpine meadow is produced early in the season between May and
mid-July (data not shown), which is before the peak drought period. Hence, in our study,
(recurrent) drought had little effect on this early-seasonal phytomass build-up. Instead,
increasing drought frequency resulted in phytomass loss and/or plant mortality, which is
visible in *fractions* of the phytomass, as it illustrates the redistribution of mass. Notably, while
biomass refers to productivity, biomass fraction reflects the maintained productivity.

Plant species seasonality and flowering

We assessed plant species seasonal growth to characterise plant species as early and late
seasonal species. For this, we determined the time point at which the plant species peaked in
its number of modules. The typical seasonal behaviour of each plant species was determined
based on the ambient treatment plots only. Early seasonal species were defined as species that
peaked in modules before the biomass cut and thus reached peak growth between the end of
May to end of July. Late seasonal species were defined as species that peaked in modules after
the biomass cut and thus reached peak growth between August and October. Shifts in early
and late seasonal patterns of species were assessed using values from all treatments.

Because of the low occurrence of flowers, seasonal shifts in flowering was calculated based on
flower biomass. For this, we calculated the ln-response ratio between a species' flower dry
mass at peak drought and its flower dry mass after the 11-week recovery period. A positive
value signifies plant species flowering early in the growing season, i.e., during the drought
period. A negative value signifies plant species flowering late in the growing season, i.e.,
during the recovery phase.

Plant community metrics and composition

Plant community evenness throughout the season was calculated as Pielou's index using the plant species module counts per time point and plot. For this, we divided the plant diversity Shannon index by the ln-transformed species richness using the function diversity and specnumber of the vegan package (R Core Team 2021; Oksanen *et al.* 2020). Plant community seasonal synchrony was calculated following Loreau & De Mazancourt (2008), using the package codyn (Hallett *et al.* 2016). In brief, we compared the variance of the total module counts of all species over the season (x_T) with the summed variances of the individual species' module counts over the season (x_i) to quantify seasonal synchrony:

$$Seasonal\ synchrony = \frac{\sigma(x_T)^2}{(\sum_i \sigma(x_i))^2}$$

Where:

$$x_T(t) = \sum_{i=1}^{N} x_i(t)$$

With $x_T(t)$ being the total module counts of all species combined at time t, $x_i(t)$ the module counts of species i at time t, $\sigma(x_T)^2$ the variance of the total module counts of all species over the season combined, and $\sigma(x_i)$ the variance of the module counts of species i over the season. Using the codyn package (Hallett *et al.* 2016), plant species loss between subsequent time points was calculated to indicate species loss *within* the growing season. To determine whether communities were dominated by either early or late plant species, we calculated the proportional number of early plant species present in each community at each time point (see section Plant species seasonality and flowering).

We assessed plant community composition using Non-Metric Multidimensional Scaling (NMDS) using metaMDS from the vegan package (Oksanen *et al.* 2020). NMDS on both module counts and biomass were run on Bray-Curtis dissimilarity matrices between plots and time points and with four dimensions to minimise stress values.

Statistics

All calculations and statistical analyses were done in R 4.1.1 (R Core Team 2021). Effects of

drought frequency on percentage data (phytomass fractions) were performed using

Generalized Linear Mixed Models (GLMM) with a beta distribution and logit-link using the

package glmmTMB (Magnusson *et al.* 2017). Drought frequency effects on species richness

and the number of seedlings were tested using Generalised Linear Mixed-Effects Models

(GLMER) with a poisson distribution and log-link function using the lme4 package (Bates *et*

al. 2015). Drought frequency effects on continuous variables (seasonal synchrony and

community evenness) were tested using Linear Mixed Effects Models (LMER) using the lme4

package (Bates *et al.* 2015). Models including 'date' (continuous, scaled) were run with 'plot'

as a random effect to take the repeated sampling design into account. Models on plant species-

specific traits were only run for plant species for which at least two data points occurred in

each treatment.

Heterogeneity of the mountain meadow was taken into account as a random effect in

statistical models. For models on plant community measures (phytomass fractions, seasonal

synchrony, species richness, community evenness and seedling numbers), 'block' was added

as a random effect. Models on plant species-specific traits included the mean vegetation height

of each plot before the setup of the rainout shelters as a random effect. This variable was better

at capturing variation *within* drought treatments than 'block' and, importantly, was not

significantly affected by previous drought events (data not shown).

For GLMM and GLMER models, we tested over- and under-dispersion of the model residuals

using the package DHARMa (Hartig 2022). For LMER models, we tested assumptions of

homogeneity of variances, and normal residual distributions following Zuur *et al.* (2010). Data

was ln- or sqrt-transformed when model residuals did not adhere to model assumptions.

For plant community composition based on NMDS analyses, significant separation of

communities resulting from 'drought frequency' (factor), date (scaled, continuous), and their

interaction was tested using a restricted permutation test: samples were permuted *within* plots

to take the repeated sampling structure over time into account. The test was performed using

adonis from the vegan package with 999 permutations (Oksanen *et al.* 2020). A passive overlay

of 'drought frequency' (ln-transformed + 1), 'species richness', 'proportion early species',

'seasonal synchrony', species loss between subsequent time points (for NMDS based on

counts only), 'LAI', functional group total abundance, flower abundance, and seedling

abundance was created and tested for significance using the function envfit of the vegan

package with 999 permutations (Oksanen *et al.* 2020).

Results

Drought frequency decreased biomass and species richness and increased necromass

In a species-rich sub-alpine mountain meadow, we asked how an increase in the annual frequency of severe summer drought affects plant community phytomass, species richness, and flower production during peak drought and recovery. At the peak of drought, 7 weeks after the setup of the rainout shelters, we found that drought frequency significantly decreased the fraction of community biomass and increased the fraction of community necromass. These effects were non-linear, with the strongest effects observed during low drought frequency, plateauing as drought frequency increased (Figure 2AB). The increase in necromass was primarily related to an increase in the necromass fraction of grass and legume species (Figure S3). Eleven weeks after soil rewetting (recovery), these effects were still present, albeit to a lesser extent (Figure S4AB). In the recovery phase, the fraction of forb biomass decreased, while the fraction of grass biomass and necromass increased with drought frequency (Figure S3).

Drought frequency did not significantly affect the fraction of flower biomass at peak drought (Figure 2C). During recovery, however, the fraction of flower biomass was significantly decreased with increasing drought frequency (Figure S4C).

Furthermore, drought frequency significantly decreased plant species richness at both peak drought and recovery (Figure 2D; S4D). On average, 13 years of recurrent summer drought resulted in a 35 % and 46 % loss of plant species compared with ambient conditions at peak drought and recovery, respectively. Despite this loss in plant species, drought frequency did not significantly affect plant community evenness at both peak drought and recovery (Figure 2E; S4E).

Figure *2 Drought frequency effects at peak drought on (A) the fraction of plant community biomass (living plant*
material), (B) the fraction of plant community necromass (dead plant material), (C) the fraction of total flower
mass, (D) plant community species richness, and (E) plant community evenness. A-C are expressed as
percentages of the phytomass (bio- and necromass) in each plot. For significant relations, the black line indicates
the mean relation with the shading showing the 95% confidence interval. P-values of significant relations are
shown (ns: no significant effect, p > 0.05, n = 4 per drought frequency level). For effects of drought frequency on
overall plant community parameters in the recovery phase, see Figure S4.

Drought frequency decreased forb and legume seedlings

To assess the impact of drought frequency on the establishment of plant seedlings throughout

the growing season, we monitored the number of seedlings per plant functional group at

biweekly intervals throughout the season. Drought frequency affected the number of forb and

legume seedlings, but not of grass seedlings. During drought and recovery, the number of

forb and legume seedlings was consistently lowest with 13 years of recurrent summer drought

(Figure 3). Additionally, while in the ambient treatment and one- year drought plots, many

forb and legume seedlings appeared after the mowing event, these peaks were lower in plots

with three years and especially 13 years of recurrent summer drought (Figure 3). At the same

time, plant community LAI decreased with drought frequency at both peak drought and

recovery (Figure S5AB). The latter indicates that the soil surface was less covered by leaves of

the established plant community with an increase in drought frequency.

Figure 3 *Drought frequency effects on the number of seedlings of (A) grass, (B) forb, and (C) legume species*
throughout the growing season. The dotted line indicates the timepoint at which drought ended, and the plots
were mown and rewetted. Solid lines indicate mean seedling numbers, and faded lines indicate seedling numbers
of each plot (n = 4 per drought frequency treatment). Results from analyses of variances are shown with '***'
indicating p < 0.001, '*' indicating 0.01 > p < 0.05, and 'ns' indicating no significant effect (p > 0.05).

Drought frequency increased plant community seasonal synchrony

We tested how drought frequency affected plant community seasonal dynamics based on

plant species module counts which we monitored throughout the growing season. The total

number of plant modules in all treatments fluctuated throughout the growing season (Figure

S6A). However, it was not significantly affected by drought frequency throughout the season

(Figure S6A). Furthermore, plant species richness before, during, and after drought was

consistently lowest in plots subjected to 13 years of drought, followed by three years, and one

year of drought (Figure S6B). Plant species richness across all treatments dropped sharply

after peak drought and the mowing event but increased to levels similar to their respective

pre-drought conditions within a month of recovery (Figure S6B).

Next, we tested drought frequency effects on plant community seasonal synchrony, which we calculated based on the module counts of each plant species over the season. We found that drought significantly increased seasonal synchrony, that is, drought synchronised the timing of plant species growth during the season. Increasing drought frequency enhanced this synchronising effect. This relationship was non-linear, with the strongest increase in synchrony occurring during low drought frequencies and the increase plateauing as drought frequency became higher (Figure 4A). Furthermore, seasonal synchrony was negatively related to the average plant species richness throughout the season, that is, synchrony was high when average plant species richness was low (Figure 4B).

Figure 4 (A) Effects of increasing drought frequency on seasonal synchrony (sensu Loreau & De Mazancourt (2008)) based on plant module counts across the growing season from mid-May to mid-October, and (B) seasonal synchrony in relation to the average plant species richness (n = 4 per drought frequency treatment). The black line indicates the mean relation with the shading showing the 95% confidence interval. P-values of significant relations are shown.

Drought frequency shifts plant community composition towards less early seasonal and more grass species

We examined the effect of drought frequency on plant community composition throughout the season using non-metric multidimensional scaling (NMDS). We found significant differentiation of plant communities based on drought frequency, with communities experiencing no drought and 13 years of drought exhibiting the greatest dissimilarity (Figure

5A, S7). This compositional separation, driven by drought frequency, was primarily captured

along the first axes of the NMDS plot. This first axis coincided with the decrease in plant

species richness and the increase in community synchrony as drought frequency increased

(Figure 5A). Moreover, we observed that the compositional shifts associated with increasing

drought frequency were characterised by a high grass abundance and a decrease in early-

seasonal plant species. In addition, LAI, the number of forb flowers, and the number of forb

seedlings decreased with drought frequency (Figure 5A). These patterns were similar when

basing the analyses on plant biomass collected at peak drought and recovery (Figure S8).

During the recovery phase, plant communities in the plots exposed to one and three years of

drought shifted along the second NMDS axis. After rewetting, these one- and three-year

droughted plots temporarily shifted into the top-right quadrant of the NMDS (Figure 5B). This

direction was related to a high loss of species, particularly early-seasonal species (for

specification see Table S1), and a decrease in forb and legume abundance (Figure 5B).

However, later in the recovery phase, the one- and three-year droughted communities

gradually returned to a similar position on the second axis of the NMDS plot as before the

recovery phase (Figure 5B). This recovery pattern indicates slower recovery of early-seasonal

forb, and legume species within the one- and three-year droughted plots (Figure 5B).

 Figure 5 Non-metric multidimensional scaling (NMDS) of plant species module counts throughout the growing
season in permanent plots receiving no drought (0 years), one year, three years, and 13 years of recurrent summer
drought. (A) Drought frequency effects on plant community composition with a passive overlay of community
variables significantly relating to the NMDS patterns. Each point represents a plot at a point in time. Hulls group
all plots and time points per drought frequency treatment together. Length of arrows indicate the strength of the
relation. 'Proportion early species' refers to the proportion of early species versus the total number of species
present in a plot. 'Grass', 'forb', and 'legume abundance' refer to the total number of modules summed per
functional group. 'Synchrony' refers to community seasonal synchrony sensu Loreau & De Mazancourt (2008).
'Species loss' refers to the percentage of species lost within the growing season between two subsequent time points
(species gain was not significantly related). 'LAI' refers to leaf area index. Results of a restricted permutation test
*is shown with '**' indicating 0.001 > p < 0.01, '*' indicating 0.01 > p < 0.05, and 'ns' indicating no significant*
effect (p > 0.05) (n = 4 plots per drought frequency treatment). (B) Community compositional development over
time. Points are mean values ± standard errors. Note that axes scales differ between A and B to improve
readability. For NMDS axes 3 and 4, see Figure S7. For position of plant species along the axes, see Figure S9.
For the plant community composition results based on plant species biomass, see Fig S8.

437 Weak drought frequency effects on individual plant species responses

Finally, for 10 plant species that were present in sufficient abundances in all three drought
frequency treatments and the ambient treatment, we tested whether drought frequency
affected SLA at peak drought and recovery, as well as the timing of their growth in the season.
At peak drought, we found that drought frequency significantly decreased SLA of *Alchemilla*
vulgaris, *Leontodon hispidus*, *Plantago lanceolata* (marginally), *Trifolium repens*, and *Veronica*
chamaedrys (Figure S10). During recovery, these patterns had disappeared (Figure S11).
Drought frequency only significantly affected the timing of *Trifolium pratense*, which peaked
in number of modules earlier in the season with a high drought frequency (Figure S12).
Drought frequency only significantly affected the timing of flowering of *Leontodon hispidus*,
which shifted its flower production to earlier in the season with a high drought frequency
(Figure S13).

Discussion

We investigated the effects of recurrent summer drought of up to 13 years on plant

communities in a sub-alpine mountain meadow, as well as their underlying mechanisms. We

found that drought frequency increased plant community seasonal synchrony. This increase

in synchrony with greater drought frequency was underpinned by a decrease in species

richness, a loss of early-seasonal plant species, a shift in plant functional groups, and the

constrained establishment of seedlings throughout the growing season. Moreover, these

changes were associated with a decreased biomass fraction as drought frequency increased.

Higher seasonal synchrony with increasing drought frequency due to species loss, compositional shifts, and lower seedling establishment

We found that with increasing drought frequency, species **synchrony** within the plant

community was enhanced in a non-linear fashion: the increase was strongest at low drought

frequency and plateaued at high drought frequencies. This indicates that the species in the

plant community under increasing drought frequency were more similar in their timing of

plant species growth throughout the season. Results on increasing synchrony *between* years

for grasslands were reported by other studies, finding an increase in synchrony with drought

frequency of three years in an alpine meadow (He *et al.* 2022), and four years across grasslands

(Muraina *et al.* 2021). Our study suggests that with increasing drought frequency (to up to 13

years of recurrent drought) seasonal synchrony becomes increasingly pronounced. In

consequence, multiple droughts may hamper the stability (De Mazancourt *et al.* 2013; Loreau

& De Mazancourt 2013), e.g. under future disturbances. This is because synchronous

communities are typically less stable to disturbances than asynchronous communities (Luo *et*

al. 2023b; Muraina *et al.* 2021; Craven *et al.* 2018; De Mazancourt *et al.* 2013). Hence, an increase

in seasonal synchrony with drought frequency suggests that multiple droughts may hamper

the stability of communities to future disturbances. We suggest that with the ongoing and

projected increases in the frequency of drought events due to climate change (IPCC 2021),

grassland ecosystems become increasingly seasonally synchronous and vulnerable to future

disturbances.

We identified several underlying mechanisms that contributed to the increase in synchrony

with increasing drought frequency. First, the enhancement in synchrony was associated with

a **decrease in plant species richness**. This drought-induced species loss aligns with findings

from other studies that observed enhanced mortality during and following single (Sippel *et*

al. 2018; Stampfli *et al.* 2018; Frank *et al.* 2015), and recurrent drought events (Luo *et al.* 2023a;

Wagg *et al.* 2017). Species richness mostly tends to be negatively associated with synchrony

(Zhang *et al.* 2018; Hautier *et al.* 2014), but was also found to be positively associated with

synchrony (Valencia *et al.* 2020). The negative association between plant species richness and

seasonal synchrony in our study likely occurred because the plant species that survived in the

community with recurrent drought, were more similar in terms of their seasonal

development. Moreover, under increasing drought frequency, communities were dominated

by species growing later in the season, as **early-growing species were lost**. This may have

been caused by a reduced recovery potential of early-season species after the drought event:

We found that early-seasonal species were taking a longer time to recover under single- and

three-year drought. While it was suggested, that early-season species may be more susceptible

to drought (Mamolos *et al.* 2001), we found first evidence that early-growing species actually

get lost at high drought frequencies. As a result, under recurrent drought, more temporally

similar growing species lead to more seasonally synchronized communities.

Furthermore, the seasonal synchronization of species may have additionally been driven by

the observed shift within plant **functional groups** under higher drought frequency. We found

that the community composition showed a reduction in forb and legume species, and more

grass species with increasing drought frequency. The loss of species within these functional

groups with increasing drought frequency may be explained by the observed lower forb and

legume seedlings throughout the season and the slower recovery of forbs and legumes post

drought. Hence, our findings support the notion that forb and legume species are less robust

to recurrent drought compared with grass species. This is in line with studies on single

drought (De Boeck *et al.* 2018; Hoover *et al.* 2014) and recurrent drought events (Luo *et al.*

2023a; Luo *et al.* 2023b; Sun *et al.* 2023; Xu *et al.* 2021), which also suggest grass species to be

less sensitive to drought compared to forbs. Overall, the observed shifts under recurrent

drought towards less diversity in functional groups can enhance synchrony by increasing the

similarity of seasonal growth within the community. This similarity in functional groups can

furthermore decrease the variety in responses to drought throughout the growing season

(Zhang *et al.* 2020) and thereby contribute to an enhanced vulnerability under future

disturbances.

The increase in synchrony with higher drought frequency may also be driven by the

constrained establishment of seedlings throughout the growing season: As drought

frequency increased, we observed a **decline in seedling numbers**. This occurred during the

entire vegetation period and was most pronounced with high drought frequency of 13 years.

At the same time, less soil surface area was covered by leaves under high drought frequency.

Hence, the seedling decline took place despite a diminished light limitation, and thus likely a

higher chance of seedling survival in the 13-year recurrent drought plots (Eskelinen *et al.*

2022). Moreover, we observed that plant species **reproductive output** was inhibited with

increasing drought frequency: In the recovery phase, the fraction of flower biomass was

decreased with increasing drought frequency. A negative drought effect on reproductive

output was also found by Zeiter *et al.* (2016). Reasons might be, that species experiencing

drought were unable to produce flowers due to preceding and ongoing water limitation.

Consequently, in the communities with these high drought frequencies, fewer seedlings

appeared in the recovery phase.

Overall, in our study, flower development and consequently seedlings, were severely

inhibited with higher drought frequencies. This reduces the establishment of seedlings and

leads to species loss, with effects on community composition and synchrony. During long-

term recurrent drought, effects of drought on reduced plant reproductive output and resultant

seedlings likely exacerbated plant species loss and the increase in seasonal synchrony.

However, single- and low-frequency experiments do not cover drought effects that go beyond

the lifetime of the plant species. We therefore propose that, to comprehensively comprehend

changes in plant communities arising from high drought frequency, the implementation of

long-term experiments is essential.

Enhanced loss of biomass fraction with increasing drought frequency

Drought did not reduce productivity in our study (Figure S2), which is consistent with

findings on productivity under five years of drought by Jentsch *et al.* (2011) and shoot biomass

under four years of drought by Fuchslueger *et al.* (2016). Nonetheless, other low drought

frequency studies reported a continuous decline in aboveground net primary productivity

under recurrent drought of up to four years (Luo *et al.* 2023a; Xu *et al.* 2021). The divergent

results may be attributed to the different frequencies of drought events or plant communities.

However, we found that increasing drought frequency gradually decreased the fraction of

plant community biomass and enhanced the fraction of necromass at peak drought and

recovery, respectively. These were non-linear effects, with strongest increases/decreases at

low drought frequency that plateaued as drought frequency became higher. Hence, during

the growing season, including drought and recovery, the maintained productivity is lower,

and senescence is higher with increasing drought frequency. The higher senescence at peak

drought and recovery supports our observation of increasing species loss throughout the

growing season. Reductions in maintained productivity might be caused by the lasting

repercussions of drought on plant community composition and synchrony.

Under recurrent drought, plants have the capacity to develop **adaptations** that can dampen

drought effects (Müller & Bahn 2022). This can occur e.g., by adjusting photosynthesis,

osmosis (Li *et al.* 2022), root biomass (Legay *et al.* 2018), epigenetics, transcription factors and

rate (Jacques *et al.* 2021; Ding *et al.* 2012; Bruce *et al.* 2007), as well as transgenerational changes

(Chen *et al.* 2022). Within our study, we found only weak species adaptations under recurrent

drought. For example, increasing drought frequency did not result in adaptations of the

duration of the growing season and only shifted the timing of growth for one legume species

to earlier in the season. Also, the timing of plant species flowering was only shifted earlier

with increasing drought frequency for one forb species. Moreover, the lower SLA at peak

drought of some of the most abundant forb species and one legume species with increasing

drought frequency can hint towards higher water-use efficiency, resource conservation, and

reduced plant growth, which can enhance short-term stress tolerance (Wellstein *et al.* 2017).

This short-term adaptation could lead to less vulnerability under drought. Overall,

adaptations did not play a major role in compensating drought effects in our study.

Chen *et al.* (2022) suggested that adaptations *along generations* under recurrent drought lead to

improved ecosystem responses. However, we suggest that the lower reproduction might

overrule any potential adaptive effects of single plant species along generations through

species loss. Plant communities are therefore likely not able to adapt to the predicted rapid

increase in drought frequency with the ongoing change in climate. To gain further

understanding of the repercussions on plant community due to increasing drought

frequencies, long-term drought experiments, including transgenerational effects, could yield

important insights.

Conclusion

We show that long-term annual drought frequency in grasslands increases plant community seasonal synchrony. Synchrony increased through a reduction in species richness, a loss of early-seasonal plant species, a shift in plant functional groups, and the constrained establishment of seedlings throughout the growing season. These changes were associated with a decreased fraction of biomass with increasing drought frequency.

Overall, we conclude that with the increase in frequency and severity of drought events due to climate change, grassland plant communities will increase in seasonal synchrony. As asynchrony is a critical component of community stability, we suggest that this will lead to lasting effects far beyond the drought event, by increasing the vulnerability of ecosystems to future disturbances. Moreover, we show that negative drought effects were enhanced with an increasing drought frequency, though effects may vary across ecosystems and climate conditions (Tielbörger *et al.* 2014). We conclude that single and low-frequency drought studies may not adequately predict longer-term changes in our rapidly shifting climate. Furthermore, on the long-term, negative effects, mostly associated with species loss, will likely overrule the comparatively weak adaptation of species. Hence, with the current speed of climate change, communities will be severely and probably irrevocably affected the moment species are lost from the ecosystem. We therefore conclude that plant communities are unlikely to be able to adapt to recurrent drought. Importantly, long-term experiments will be essential in understanding the severity of these climate change impacts.

Sample CRediT author statement

Lena M. Müller: Conceptualization, Investigation, Project administration, Writing - Original Draft (lead), Visualization, Funding acquisition; **Michael Bahn**: Conceptualization, Resources, Funding acquisition, Writing - Review & Editing; **Maximillian Weidle**: Investigation, Writing - Review & Editing; **Georg Leitinger**: Writing - Review & Editing, Funding acquisition; **Dina in 't Zandt**: Conceptualization, Investigation, Formal Analysis, Data Curation, Visualization, Writing - Review & Editing, Writing - Original Draft, Supervision

Acknowledgements

We are grateful to numerous colleagues, students, and friends who have helped to maintain the long-term experiment over the years, as well as to those that assisted in setting up and conducting the sampling for the experiment presented in this study. Amongst others, we thank Beverley Anderson, Mario Deutschmann, Andrew Giunta, Roland Hasibeder, Johannes Ingrisch, Florian Oberleitner, Natalie Oram, David Reinthaler, Michael Schmitt, Carmen Telser, and Herbert Wachter.

The study site is part of the LTSER platform Tyrolean Alps, belonging to the national and international long-term ecological research networks (LTER-Austria, LTER Europe and ILTER). The work was funded by the Forschungszentrum Berglandwirtschaft (project no. 326572). The long-term drought treatment was maintained by projects funded by the Austrian Science Fund (FWF, project no. P22214-B17, P31132, and I1056) and the Austrian Academy of Sciences (ESS-programme, project ClimLUC).

LM was supported by the University Innsbruck through a doctoral scholarship (Doktoratsstipendium aus der Nachwuchsförderung der Universität Innsbruck) and partially funded by the Austrian Science Fund (FWF): [I4969-B]. DZ was supported by a PPLZ Postdoctoral Fellowship awarded by the Czech Academy of Sciences (RVO 67985939).

Data Availability Statement

618   Upon acceptance, all data and code (R scripts) will be made publicly available via the Zenodo

619   data repository, will be referenced using a stable object identifier (DOI) in the Materials and

620   Methods section, and cited in the reference list.

621   **Conflict of interest**

622   The authors declare no conflict of interest.

References

Backhaus S, Kreyling J, Grant K, Beierkuhnlein C, Walter J, Jentsch A (2014) Recurrent mild drought events increase resistance toward extreme drought stress. *Ecosystems*, **17**, 1068–1081.

Bardgett RD, Bullock JM, Lavorel S *et al.* (2021) Combatting global grassland degradation. *Nature Reviews Earth & Environment*, **2**, 720–735.

Bardgett RD, Caruso T (2020) Soil microbial community responses to climate extremes: resistance, resilience and transitions to alternative states. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **375**, 20190112.

Basto S, Thompson K, Grime JP, Fridley JD, Calhim S, Askew AP, Rees M (2018) Severe effects of long-term drought on calcareous grassland seed banks. *npj Climate and Atmospheric Science*, **1**, 5.

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, **67**, 1–48.

Berwaers S, De Boeck HJ, Nijs I (2019) End-of-season senescence in grassland species can be traced to leaf temperature during preceding summer drought. *Perspectives in Plant Ecology, Evolution and Systematics*, **38**, 31–38.

Blair J, Nippert J, Briggs J (2014) Grassland Ecology. In: *Ecology and the Environment* (ed Monson RK), pp 389–423. Springer New York.

Bruce TJ, Matthes MC, Napier JA, Pickett JA (2007) Stressful "memories" of plants: Evidence and possible mechanisms. *Plant Science*, **173**, 603–608.

Canarini A, Schmidt H, Fuchslueger L *et al.* (2021) Ecological memory of recurrent drought modifies soil processes via changes in soil microbial community. *Nature Communications*, **12**, 5308.

Castillioni K, Newman GS, Souza L, Iler AM (2022) Effects of drought on grassland phenology depend on functional types. *New Phytologist*, **236**, 1558–1571.

Chen Y, Vogel A, Wagg C *et al.* (2022) Drought-exposure history increases complementarity between plant species in response to a subsequent drought. *Nature Communications*, **13**, 3217.

Coleman MA, Wernberg T (2020) The silver lining of extreme events. *Trends in Ecology & Evolution*, **35**, 1065–1067.

Craven D, Eisenhauer N, Pearse WD *et al.* (2018) Multiple facets of biodiversity drive the diversity-stability relationship. *Nature Ecology & Evolution*, **2**, 1579–1587.

De Boeck HJ, Hiltbrunner E, Verlinden M, Bassin S, Zeiter M (2018) Legacy effects of climate extremes in alpine grassland. *Frontiers in Plant Science*, **9**, 1586.

De Mazancourt C, Isbell F, Larocque A *et al.* (2013) Predicting ecosystem stability from community composition and biodiversity. *Ecology Letters*, **16**, 617–625.

Ding Y, Fromm M, Avramova Z (2012) Multiple exposures to drought 'train' transcriptional responses in Arabidopsis. *Nature Communications*, **3**, 740.

Dixon AP, Faber-Langendoen D, Josse C, Morrison J, Loucks CJ (2014) Distribution mapping of world grassland types. *Journal of Biogeography*, **41**, 2003–2019.

Eskelinen A, Harpole WS, Jessen M-T, Virtanen R, Hautier Y (2022) Light competition drives herbivore and nutrient effects on plant diversity. *Nature*, **611**, 301–305.

Forrest J, Miller-Rushing AJ (2010) Toward a synthetic understanding of the role of phenology in ecology and evolution. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **365**, 3101–3112.

Frank D, Reichstein M, Bahn M *et al.* (2015) Effects of climate extremes on the terrestrial
carbon cycle: Concepts, processes and potential future impacts. *Global Change Biology*,
21, 2861–2880.
Fuchslueger L, Bahn M, Hasibeder R, Kienzl S, Fritz K, Schmitt M, Watzka M, Richter A
(2016) Drought history affects grassland plant and microbial carbon turnover during and
after a subsequent drought event. *Journal of Ecology*, **104**, 1453–1465.
Ge C, Sun S, Yao R, Sun P, Li M, Bian Y (2022) Long-term vegetation phenology changes
and response to multi-scale meteorological drought on the Loess Plateau, China. *Journal*
of Hydrology, **614**, 128605.
Gibbs HK, Salmon JM (2015) Mapping the world's degraded lands. *Applied Geography*, **57**,
12–21.
Griffin-Nolan RJ, Carroll CJ, Denton EM, Johnston MK, Collins SL, Smith MD, Knapp AK
(2018) Legacy effects of a regional drought on aboveground net primary production in six
central US grasslands. *Plant Ecology*, **219**, 505–515.
Hahn C, Lüscher A, Ernst-Hasler S, Suter M, Kahmen A (2021) Timing of drought in the
growing season and strong legacy effects determine the annual productivity of temperate
grasses in a changing climate. *Biogeosciences*, **18**, 585–604.
Hallett LM, Jones SK, MacDonald AA *et al.* (2016) codyn : An r package of community
dynamics metrics. *Methods in Ecology and Evolution*, **7**, 1146–1151.
Hartig F DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression
Models. Available via http://florianhartig.github.io/DHARMa/.
Hasibeder R, Fuchslueger L, Richter A, Bahn M (2015) Summer drought alters carbon
allocation to roots and root respiration in mountain grassland. *New Phytologist*, **205**,
1117–1127.
Hautier Y, Seabloom EW, Borer ET *et al.* (2014) Eutrophication weakens stabilizing effects
of diversity in natural grasslands. *Nature*, **508**, 521–525.
He YL, Wang JS, Tian DS *et al.* (2022) Long-term drought aggravates instability of alpine
grassland productivity to extreme climatic event. *Ecology*, **103**, 1-12.
Herben T, Březina S, Hadincová V, Krahulec F, Skálová H (2013) Mutual replacement of
species in space in a grassland community: is there an evidence for functional
complementarity of replacement groups? *Oikos*, **122**, 111–121.
Herben T, Krahulec F, Hadincová V, Pecháčková S, Kováčiová M (1997) Fine-scale spatio-
temporal patterns in a mountain grassland: do species replace each other in a regular
fashion? *Journal of Vegetation Science*, **8**, 217–224.
Hofer D, Suter M, Haughey E, Finn JA, Hoekstra NJ, Buchmann N, Lüscher A (2016) Yield
of temperate forage grassland species is either largely resistant or resilient to
experimental summer drought. *Journal of Applied Ecology*, **53**, 1023–1034.
Hoover DL, Knapp AK, Smith MD (2014) Resistance and resilience of a grassland
ecosystem to climate extremes. *Ecology*, **95**, 2646–2656.
Hoover DL, Pfennigwerth AA, Duniway MC (2021) Drought resistance and resilience: the
role of soil moisture-plant interactions and legacies in a dryland ecosystem. *Journal of*
Ecology, **109**, 3280–3294.
Hoover DL, Wilcox KR, Young KE (2018) Experimental droughts with rainout shelters: a
methodological review. *Ecosphere*, **9**.
in 't Zandt D, Herben T, Brink A, Visser EJ, Kroon H (2021) Species abundance fluctuations
over 31 years are associated with plant–soil feedback in a species-rich mountain meadow.
Journal of Ecology, **109**, 1511–1523.

Ingrisch J, Karlowsky S, Anadon-Rosell A, Hasibeder R, König A, Augusti A, Gleixner G,
Bahn M (2018) Land Use Alters the Drought Responses of Productivity and CO2 Fluxes in
Mountain Grassland. *Ecosystems*, **21**, 689–703.
Ingrisch J, Karlowsky S, Hasibeder R, Gleixner G, Bahn M (2020) Drought and recovery
effects on belowground respiration dynamics and the partitioning of recent carbon in
managed and abandoned grassland. *Global Change Biology*, **26**, 4366–4378.
IPCC (2021) Climate Change 2021. The Physical Science Basis. Contribution of Working
Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate
Change. Cambridge, UK: Cambridge University Press.
IPCC (2023) Climate Change 2023. Synthesis Report. Contribution of Working Groups I, II
and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change
[Core Writing Team, H. Lee and J. Romero (eds.)]., IPCC, Geneva, Switzerland.
Jacques C, Salon C, Barnard RL, Vernoud V, Prudent M (2021) Drought Stress Memory at
the Plant Cycle Level: A Review. *Plants*, **10**, 1873.
Jentsch A, Kreyling J, Boettcher-Treschkow J, Beierkuhnlein C (2009) Beyond gradual
warming: Extreme weather events alter flower phenology of European grassland and
heath species. *Global Change Biology*, **15**, 837–849.
Jentsch A, Kreyling J, Elmer M *et al.* (2011) Climate extremes initiate ecosystem-regulating
functions while maintaining productivity. *Journal of Ecology*, **99**, 689–702.
Ji S, Ren S, Li Y, Dong J, Wang L, Quan Q, Liu J (2021) Diverse responses of spring
phenology to preseason drought and warming under different biomes in the North China
Plain. *The Science of the Total Environment*, **766**, 144437.
Kang W, Wang T, Liu S (2018) The response of vegetation phenology and productivity to
drought in semi-arid regions of northern China. *Remote Sensing*, **10**, 727.
Karlowsky S, Augusti A, Ingrisch J, Akanda MK, Bahn M, Gleixner G (2018) Drought-
induced accumulation of root exudates supports post-drought recovery of microbes in
mountain grassland. *Frontiers in Plant Science*, **9**, 1593.
Knapp AK, Condon KV, Folks CC *et al.* (2023) Field experiments have enhanced our
understanding of drought impacts on terrestrial ecosystems—But where do we go from
here? *Functional Ecology*, **38**, 76–97.
Legay N, Piton G, Arnoldi C *et al.* (2018) Soil legacy effects of climatic stress, management
and plant functional composition on microbial communities influence the response of
Lolium perenne to a new drought event. *Plant and Soil*, **424**, 233–254.
Li X, Jimoh SO, Li Y, Duan J, Cui Y, Jin K, Wang Z, Zhang Y (2022) Stress memory and
phyllosphere/soil legacy underlie tolerance and plasticity of Leymus chinensis to periodic
drought risk. *Agricultural and Forest Meteorology*, **312**, 108717.
Loreau M, De Mazancourt C (2008) Species synchrony and its drivers: neutral and
nonneutral community dynamics in fluctuating environments. *The American naturalist*,
172, E48-66.
Loreau M, De Mazancourt C (2013) Biodiversity and ecosystem stability: a synthesis of
underlying mechanisms. *Ecology letters*, **16**, 106–115.
Luo W, Muraina TO, Griffin-Nolan RJ *et al.* (2023a) Responses of a semiarid grassland to
recurrent drought are linked to community functional composition. *Ecology*, **104**, e3920.
Luo W, Shi Y, Wilkins K *et al.* (2023b) Plant traits modulate grassland stability during drought
and post-drought periods. *Functional Ecology*, **37**, 2611–2620.
Mackie KA, Zeiter M, Bloor JM, Stampfli A (2019) Plant functional groups mediate drought
resistance and recovery in a multisite grassland experiment. *Journal of Ecology*, **107**,
937–949.

Magnusson A, Skaug HJ, Nielsen A *et al.* (2017) glmmTMB: Generalised Linear
MixedModels using Template Model Builder. https://github.com/glmmTMB.
Mamolos AP, Veresoglou DS, Noitsakis V, Gerakis A (2001) Differential drought tolerance of
five coexisting plant species in Mediterranean lowland grasslands. *Journal of Arid*
Environments, **49**, 329–341.
Müller LM, Bahn M (2022) Drought legacies and ecosystem responses to subsequent
drought. *Global Change Biology*, **28**, 5086–5103.
Muraina TO, Xu C, Yu Q *et al.* (2021) Species asynchrony stabilises productivity under
extreme drought across Northern China grasslands. *Journal of Ecology*, **109**, 1665–1675.
Oksanen J, Blanchet FG, Friendly M *et al.* (2020) vegan: Community Ecology Package.
https://CRAN.R-project.org/package=vegan.
Oram NJ, Ingrisch J, Bardgett RD *et al.* (2023) Drought intensity alters productivity, carbon
allocation and plant nitrogen uptake in fast versus slow grassland communities. *Journal of*
Ecology, **111**, 1681–1699.
Peng J, Wu C, Zhang X, Wang X, Gonsamo A (2019) Satellite detection of cumulative and
lagged effects of drought on autumn leaf senescence over the Northern Hemisphere.
Global Change Biology, **25**, 2174–2188.
Petermann JS, Buzhdygan OY (2021) Grassland biodiversity. *Current Biology*, **31**, R1195-
R1201.
Qian J, Guo Z, Muraina TO *et al.* (2022) Legacy effects of a multi-year extreme drought on
belowground bud banks in rhizomatous vs bunchgrass-dominated grasslands. *Oecologia*,
198, 763–771.
R Core Team (2021) A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria.
Rihan W, Zhao J, Zhang H, Guo X (2022) Preseason drought controls on patterns of spring
phenology in grasslands of the Mongolian Plateau. *The Science of the Total Environment*,
838, 156018.
Schirpke U, Kohler M, Leitinger G, Fontana V, Tasser E, Tappeiner U (2019) Future impacts
of changing land-use and climate on ecosystem services of mountain grassland and their
resilience. *Ecosystem services*, **26**, 79–94.
Schmitt M, Bahn M, Wohlfahrt G, Tappeiner U, Cernusca A (2010) Land use affects the net
ecosystem CO_2 exchange and its components in mountain grasslands. *Biogeosciences*,
7, 2297–2309.
Schwalm CR, Anderegg WR, Michalak AM *et al.* (2017) Global patterns of drought recovery.
Nature, **548**, 202–205.
Sippel S, Reichstein M, Ma X, Mahecha MD, Lange H, Flach M, Frank D (2018) Drought,
heat, and the carbon cycle: A review. *Current Climate Change Reports*, **4**, 266–286.
Smith MD, Wilkins KD, Holdrege MC *et al.* (2024) Extreme drought impacts have been
underestimated in grasslands and shrublands globally. *Proceedings of the National*
Academy of Sciences, **121**, e2309881120.
Spinoni J, Vogt JV, Naumann G, Barbosa P, Dosio A (2018) Will drought events become
more frequent and severe in Europe? *International Journal of Climatology*, **38**, 1718–
1736.
Stampfli A, Bloor JM, Fischer M, Zeiter M (2018) High land-use intensity exacerbates shifts
in grassland vegetation composition after severe experimental drought. *Global Change*
Biology, **24**, 2021–2034.

Stampfli A, Zeiter M (2004) Plant regeneration directs changes in grassland composition
after extreme drought: a 13-year study in southern Switzerland. *Journal of Ecology*, **92**,
568–576.
Sun J, Zhang B, Pan Q *et al.* (2023) Non-linear response of productivity to precipitation
extremes in the Inner Mongolia grassland. *Functional Ecology*, **37**, 1663–1673.
Tello-García E, Huber L, Leitinger G, Peters A, Newesely C, Ringler M-E, Tasser E (2020)
Drought- and heat-induced shifts in vegetation composition impact biomass production
and water use of alpine grasslands. *Environmental and Experimental Botany*, **169**,
103921.
Tielbörger K, Bilton MC, Metz J *et al.* (2014) Middle-Eastern plant communities tolerate 9
years of drought in a multi-site climate manipulation experiment. *Nature Communications*,
5, 5102.
Valencia E, Bello F de, Galland T *et al.* (2020) Synchrony matters more than species
richness in plant community stability at a global scale. *Proceedings of the National*
Academy of Sciences, **117**, 24345–24351.
Vorkauf M, Kahmen A, Körner C, Hiltbrunner E (2021) Flowering phenology in alpine
grassland strongly responds to shifts in snowmelt but weakly to summer drought. *Alpine*
Botany, **131**, 73–88.
Wagg C, O'Brien MJ, Vogel A, Scherer-Lorenzen M, Eisenhauer N, Schmid B, Weigelt A
(2017) Plant diversity maintains long-term ecosystem productivity under frequent drought
by increasing short-term variation. *Ecology*, **98**, 2952–2961.
Walter J, Nagy L, Hein R, Rascher U, Beierkuhnlein C, Willner E, Jentsch A (2011) Do
plants remember drought? Hints towards a drought-memory in grasses. *Environmental*
and Experimental Botany, **71**, 34–40.
Wang M, Li P, Peng C, Xiao J, Zhou X, Luo Y, Zhang C (2022) Divergent responses of
autumn vegetation phenology to climate extremes over northern middle and high latitudes.
Global Ecology and Biogeography, **31**, 2281–2296.
Wang T, Tu X, Singh VP, Chen X, Lin K (2021) Global data assessment and analysis of
drought characteristics based on CMIP6. *Journal of Hydrology*, **596**, 126091.
Wellstein C, Poschlod P, Gohlke A *et al.* (2017) Effects of extreme drought on specific leaf
area of grassland species: A meta-analysis of experimental studies in temperate and sub-
Mediterranean systems. *Global Change Biology*, **23**, 2473–2481.
Wilcox KR, Blumenthal DM, Kray JA, Mueller KE, Derner JD, Ocheltree T, Porensky LM
(2021) Plant traits related to precipitation sensitivity of species and communities in
semiarid shortgrass prairie. *New Phytologist*, **229**, 2007–2019.
Wilcox KR, Tredennick AT, Koerner SE *et al.* (2017) Asynchrony among local communities
stabilises ecosystem function of metacommunities. *Ecology Letters*, **20**, 1534–1545.
Xu C, Ke Y, Zhou W *et al.* (2021) Resistance and resilience of a semi-arid grassland to
multi-year extreme drought. *Ecological Indicators*, **131**, 108139.
Xu Z, Ren H, Li M-H *et al.* (2015) Environmental changes drive the temporal stability of
semi-arid natural grasslands through altering species asynchrony. *Journal of Ecology*,
103, 1308–1316.
Yuan M, Zhao L, Lin A, Wang L, Li Q, She D, Qu S (2020) Impacts of preseason drought on
vegetation spring phenology across the Northeast China Transect. *The Science of the*
Total Environment, **738**, 140297.
Yuan Z, Tong S, Bao G, Chen J, Yin S, Li F, Sa C, Bao Y (2023) Spatiotemporal variation of
autumn phenology responses to preseason drought and temperature in alpine and
temperate grasslands in China. *The Science of the Total Environment*, **859**, 160373.

Zeiter M, Sara Schärrer, Roman Zweifel, David M. Newbery, Stampfli A (2016) Timing of
extreme drought modifies reproductive output in semi-natural grassland. *Journal of*
Vegetation Science, **27**, 238–248.
Zeng Z, Wu W, Ge Q *et al.* (2021) Legacy effects of spring phenology on vegetation growth
under preseason meteorological drought in the Northern Hemisphere. *Agricultural and*
Forest Meteorology, **310**, 108630.
Zhang F, Quan Q, Ma F, Tian D, Hoover DL, Zhou Q, Niu S (2019) When does extreme
drought elicit extreme ecological responses? *Journal of Ecology*, **107**, 2553–2563.
Zhang J, Miao Y, Zhang T *et al.* (2020) Drought timing and primary productivity in a semiarid
grassland. *Land Degradation & Development*, **31**, 2185–2195.
Zhang Y, Loreau M, He N, Wang J, Pan Q, Bai Y, Han X (2018) Climate variability
decreases species richness and community stability in a temperate grassland. *Oecologia*,
188, 183–192.
Zuur AF, Ieno EN, Elphick CS (2010) A protocol for data exploration to avoid common
statistical problems. *Methods in Ecology and Evolution*, **1**, 3–14.