Candida albicans' **inorganic phosphate transport and evolutionary adaptation to phosphate scarcity**

Maikel Acosta-Zaldívar*[1,2], Wanjun Qi*[1], Abhishek Mishra*[3], Udita Roy[1], William R. King[4], Jana Patton-
Vogt[4], Matthew Z. Anderson[3,5], Julia R. Köhler[1,6]

1 Division of Infectious Diseases, Boston Children's Hospital/Harvard Medical School, Boston, MA 02115,
USA
2 Current affiliation: Planasa, Valladolid, Spain
3 Center for Genomic Science Innovation, University of Wisconsin-Madison, Madison, WI
4 Department of Biological Sciences, Duquesne University, Pittsburgh, Pennsylvania, USA
5 Department of Medical Genetics, Laboratory of Genetics, University of Wisconsin-Madison, Madison,
WI

6 Corresponding author

* These authors contributed equally.

macosta@planasa.com
Wanjun.Qi@childrens.harvard.edu
abhishek.mishra@wisc.edu
Udita.roy@childrens.harvard.edu
kingw@duq.edu
pattonvogt@duq.edu
mzanderson@wisc.edu
Julia.Koehler@childrens.harvard.edu

Abstract

Phosphorus is essential in all cells' structural, metabolic and regulatory functions. For fungal cells that import inorganic phosphate (Pi) up a steep concentration gradient, surface Pi transporters are critical capacitators of growth. Fungi must deploy Pi transporters that enable optimal Pi uptake in pH and Pi concentration ranges prevalent in their environments. Single, triple and quadruple mutants were used to characterize the four Pi transporters we identified for the human fungal pathogen *Candida albicans*, which must adapt to alkaline conditions during invasion of the host bloodstream and deep organs. A high-affinity Pi transporter, Pho84, was most efficient across the widest pH range while another, Pho89, showed high-affinity characteristics only within one pH unit of neutral. Two low-affinity Pi transporters, Pho87 and Fgr2, were active only in acidic conditions. Only Pho84 among the Pi transporters was clearly required in previously identified Pi-related functions including Target of Rapamycin Complex 1 signaling and hyphal growth. We used in vitro evolution and whole genome sequencing as an unbiased forward genetic approach to probe adaptation to prolonged Pi scarcity of two quadruple mutant lineages lacking all 4 Pi transporters. Lineage-specific genomic changes corresponded to divergent success of the two lineages in fitness recovery during Pi limitation. In this process, initial, large-scale genomic alterations like aneuploidies and loss of heterozygosity were eventually lost as populations presumably gained small-scale mutations. Severity of some phenotypes linked to Pi starvation, like cell wall stress hypersensitivity, decreased in parallel to evolving populations' fitness recovery in Pi scarcity, while that of others like membrane stress responses diverged from these fitness phenotypes. *C. albicans* therefore has diverse options to reconfigure Pi management during prolonged scarcity. Since Pi homeostasis differs substantially between fungi and humans, adaptive processes to Pi deprivation may harbor small-molecule targets that impact fungal growth and virulence.

Author Summary

Fungi must be able to access enough phosphate in order to invade the human body. Virulence of *Candida albicans*, the most common invasive human fungal pathogen, is known to decrease when one of the proteins that brings phosphate into the fungal cell, called Pho84, is disabled. We identified three more proteins in *C. albicans* that transport phosphate into the cell. We found that Pho84 plays the largest role among them across the broadest range of environmental conditions. After eliminating all 4 of these

transporters, we let two resulting mutants evolve for two months in limited phosphate and analyzed the

growth and stress resistance of the resulting populations. We analyzed genomes of representative

populations and found that early adaptations to phosphate scarcity occurred together with major changes

to chromosome configurations. In later stages of the adaptation process, these large-scale changes

disappeared as populations presumably gained small-scale mutations that increased cells' ability to grow

in limited phosphate. Some but not all of these favorable mutations improved resistance of evolving

populations to stressors like membrane- and cell wall stress. Pinpointing distinct mutation combinations

that affect stress resistance differently in populations adapting to scarce phosphate, may identify useful

antifungal drug targets.

Introduction

Phosphorus is an essential macronutrient for living cells and a major component of chromosomes,

membranes and the transcription and translation machineries [1]. Inorganic phosphate (Pi) is required in

the production of ATP, the energy currency of the cell, that governs central metabolic processes and

intracellular signaling. Consequently, Pi is not only required for growth and proliferation but also for

survival: e.g., fission yeast cells starved for Pi initially become quiescent and then lose viability [2].

Osmotrophic organisms that take up soluble small-molecule nutrients from their immediate environment

must import Pi separately from molecular sources of nitrogen and carbon to acquire sufficient

phosphorus. Most soils and aquatic environments contain <1%, or ≤10 mM soluble Pi, so that Pi is a scarce

resource for plants and free-living microorganisms [3-5]; human serum Pi ranges from 0.8-1.3 mM [6],

suggesting that microbial invasive pathogens of humans also experience Pi deprivation. For this reason,

the Pi-homeostatic systems of small-molecule importing organisms like bacteria, plants and fungi have

much in common. Orthology of phosphate proton symporters among plants and fungi was first

determined by complementation of a *Saccharomyces cerevisiae* null mutant in *PHO84* with two

Arabidopsis thaliana Pi transporters [7, 8]. In contrast, human phosphate homeostasis regulation differs

fundamentally from that of osmotrophs [9, 10], and since abundant phosphorus-containing molecules are

present in all human food sources of protein, the major high-affinity Pi transporter of fungi has no human

homolog.

Saccharomyces cerevisiae Pi transporters were characterized over decades according to their Pi affinity

and their pH optima [11]. Kinetic studies in the 1980s identified two Pi transport systems, one with a low

K_m value of 8.4-21.4 µM, defined by the early investigators of these systems as high-affinity, and another

with a high K_m value of 0.77-1.7 mM, defined as low-affinity [12, 13]. Further analysis suggested two

separate transporters with distinct pH optima within the high-affinity uptake system [14]. Cloning and

functional characterization of the *PHO84* gene showed that its product is a component of the high-affinity

Pi transport system [15]. Heterologous expression of Pho84 and its incorporation into liposomes then

permitted kinetic studies that demonstrated a K_m for Pi of 24 µM [16]. *S. cerevisiae* Pho84 is a member of

the Phosphate: H^+ Symporter Family within the Major Facilitator Superfamily [17]; it uses the

chemiosmotic energy of proton symport to transport the Pi anion up a concentration gradient across the

plasma membrane.

Additional Pi import systems were subsequently identified and characterized. A separate high-affinity Pi

uptake system that was enhanced in the presence of sodium at pH 7.2 was described [18]; the gene

encoding this activity was later identified and its product, named Pho89, confirmed to have a Pi K_m of 0.5

µM [14]. Pho89 belongs to solute carrier family 20 as a sodium-dependent phosphate transporter [19, 20].

Low-affinity *S. cerevisiae* Pi transporters Pho87, Pho90 and Pho91 were subsequently characterized

genetically and functionally [21]. Pho91 was later shown to reside on the vacuolar membrane and

facilitate Pi export from the vacuole to the cytosol [22]. Further work showed distinct activities of the 2

low-affinity transporters: Pho87 versus Pho90 can sustain growth of *S. cerevisiae* down to Pi

concentrations of 5 mM versus 0.5 mM, respectively [23]. *S. cerevisiae* therefore has 2 high-affinity Pi

transporters, Pho84 and Pho89, whose energetic drivers are proton- and sodium symport, respectively,

and 2 paralogous low-affinity Pi transporters, Pho87 and Pho90 [21, 23].

The genome of the opportunistic fungal pathogen *Candida albicans* encodes 4 homologs of *S. cerevisiae*

Pi transporters. In a *mariner* transposon mutant screen we previously identified a mutant in the *C. albicans*

homolog of *S. cerevisiae PHO84* as hypersensitive to rapamycin [24]. We showed that *C. albicans PHO84*

is required in normal Target of Rapamycin Complex 1 (TORC1) signaling, oxidative- and cell wall stress

resistance, survival during exposure to amphotericin B and the echinocandin micafungin, and normal

virulence [24-26]. Given the presence of other Pi transporter homologs in the *C. albicans* genome, we

sought to understand how loss of just one, Pho84, could significantly impact important physiological functions and even virulence in *C. albicans*.

Pi-acquisition and -homeostatic systems (PHO regulons) of bacterial and other fungal human pathogens are required for virulence, implicating Pi scarcity as a prevalent condition in the host [27-30]: e.g., in a pathogen that is completely adapted to its human host, *Mycobacterium tuberculosis*, transcription of a secretion system for virulence factors is activated by Pi starvation [31, 32]. Expression of high-affinity Pi transporters is typically regulated according to ambient Pi concentrations [33]. In *S. cerevisiae*, high-affinity Pi transporter-encoding genes *PHO84* and *PHO89* are upregulated during Pi starvation [21, 34, 35]. In *C. albicans* ex vivo and in vivo infection models [36-40], the *PHO84* and *PHO89* homologs are similarly upregulated [14]. These findings suggest that in the host, *C. albicans* like *M. tuberculosis* experiences Pi starvation. We therefore set out to identify and characterize the other putative *C. albicans* Pi importers that can contribute to cytosolic Pi availability for the fungus' growth and interaction with the host [24-26, 41].

We then asked whether *C. albicans* can adapt to persistent Pi scarcity. Forward genetic screens of chemically or transposon-mutagenized cells subjected to specific selective conditions are powerful discovery tools because they provide unbiased and often unexpected information [42]. In vitro evolution and whole genome analysis has been used in other pathogens for the characterization of drug responses [43-47] and in *Candida* species for analysis of drug resistance development [48-52]. We questioned whether this approach might have benefits for analysis of adaptation to nutrient scarcity compared with mutant screens. Its possible advantages might be a higher likelihood of revealing illuminating gain-of-function mutations and the ability to uncover mutations involved in polygenic traits. We used in vitro evolution and genome analysis to begin uncovering the cellular processes linked to *C. albicans'* management of Pi scarcity.

Results

***PHO84* plays a central role in growth and filamentation.**

Homology searches identified four Pi transporters in the *C. albicans* genome. A *C. albicans* ortholog of *S.*

cerevisiae PHO89, which encodes a high-affinity Pi transporter with an alkaline optimum [53], resides on

chromosome 4. Homology searches using the low affinity *S. cerevisiae PHO87* and *PHO90* paralogs found

a single homolog named *PHO87* in the Candida Genome Database (CGD) [53]. A *PHO84* homolog named

FGR2 was also identified that shares 23% amino acid identity and 42% similarity with Pho84 (S1 Fig.)*. FGR2*

was first isolated in a screen for *C. albicans* mutants impaired in filamentous growth [54] and more

recently found to contribute to filamentation differences between *C. albicans* strains [55]. To delineate

the contribution of these 4 predicted transporters to Pi acquisition, we constructed single gene deletion

mutants for *PHO89, PHO87* and *FGR2* using our FLP-*NAT1* system [56] to compare with our previously

constructed *pho84-/-* null mutants [24].

Pi restriction reduced growth only of *pho84-/-* mutants at acidic pH (Fig. 1). Null mutants in *PHO84* were

unable to grow on synthetic complete (SC) agar medium containing low (0.05 mM) or moderate (0.5 mM)

Pi at pH 3 or 5. In contrast, null mutants in the 3 other predicted Pi transporters grew similarly to the

wildtype control strain (WT) under these conditions, indicating that only *PHO84* is required in moderate

to low Pi at lower than neutral pH (Fig. 1; shown in Fig. 1 are also triple and quadruple mutants, which

retain only one or none of the 4 Pi transporters, respectively; these are described below in detail). At pH

7, *pho84* null mutants grew at all tested Pi concentrations, indicating that one or more other Pi

transporters were able to uptake sufficient Pi to sustain growth at neutral pH. At 7.3 mM Pi, all single

mutants in the 4 predicted Pi transporters grew robustly, indicating that no single transporter is

indispensable at high Pi concentrations (Fig. 1).

C. albicans' ability to readily switch between growth as single budding yeast versus as multicellular

filamentous hyphae contributes to its virulence [57, 58]. Cells lacking *PHO84* were previously shown to be

defective in hyphal formation [25, 59]. To examine the role of the different Pi transporters in

morphogenesis, cell suspensions from each of the 4 Pi transporter mutants were spotted on

filamentation-inducing agar media. Most clearly on Spider medium, cells lacking *PHO84* had minimal or

absent hyphal growth (Fig. 2A). Single mutants of the other 3 Pi transporters showed more subtle

filamentation defects than *pho84-/-* cells. Similar filamentation phenotypes were observed on RPMI agar

at pH 5 and pH 7: *pho84-/-* mutants produced occasional thin wisps of peripheral hyphae only on RPMI at

pH 5 but not pH 7, while *pho89-/-*, *pho87-/-* and *fgr2-/-* mutants showed robust hyphal growth on these

media (Fig. 2B). Together, these findings show that *PHO84* is the most important predicted Pi transporter
for filamentation under these conditions.

Only *PHO84* impacted TORC1 signaling and oxidative stress endurance.

We previously found that mutants in *PHO84* are hypersensitive to rapamycin and show decreased TORC1
signaling when growing in limited Pi [24], and that TORC1 co-regulates *PHO84* expression in addition to
its known regulation by Pho4 [60, 61]. TORC1 activation was reduced in *pho84-/-* cells as determined by
the phosphorylation state of ribosomal protein S6 (P-S6) [62] (Fig. 3) while null mutants of the other Pi
transporters did not show reduced P-S6. We concluded that Pho84 specifically contributes to TORC1
activation among the Pi transporters. TORC1 contributes to managing oxidative stress responses in *C.*
albicans, and mutants in *PHO84* are known to be hypersensitive to oxidative stress [25, 26, 59]. Among
single null mutants in each of the Pi transporters*,* only the *pho84-/-* mutant showed hypersensitivity to
the superoxide inducer plumbagin (S2 Fig.).

Triple and quadruple mutants in predicted Pi transporters support a major role for Pho84.

To examine the Pi uptake characteristics of each transporter, we constructed triple mutants that retained
one predicted Pi transporter, Pho84, Pho89, Pho87 or Fgr2. Triple mutants retaining a single Pi transporter
are abbreviated as the name of the sole remaining transporter followed by a capital A for "alone among
predicted Pi transporters" (e.g.*,* Pho84-A is *pho89-/- pho87-/- fgr2-/-*). Among these triple mutants, only
Pho84-A grew under all tested conditions (Fig. 1). Under Pi limiting conditions, Pho89-A cells showed
significant growth only at neutral pH. These results support a role for Pho89 as a high-affinity transporter
with a more alkaline optimum as in *S. cerevisiae* (Fig. 1). Pho87-A and Fgr2-A cells grew only in high Pi (7.3
mM) at acidic pH (pH 5 and pH 3) (Fig. 1), suggesting that Pho87 and Fgr2 are low-affinity transporters
with an acidic optimum.

We engineered two quadruple Pi transporter mutants (Q-, *pho84-/- pho89-/- pho87-/- fgr2-/-*). These
strains grew on rich complex medium, YPD, that contains organic phosphate compounds. They were then
tested for growth on Pi as the sole phosphorus source at a range of concentrations. Q- cells were able to
grow on high (7.3 mM) Pi at acidic pH, while on moderate (0.5 mM) Pi their growth was barely detectable

(Fig. 1). They did not grow at pH 7 or at a low Pi concentration (0.05 mM, Fig. 1). These findings show that

a residual Pi transport capacity exists in cells lacking the 4 identified Pi transporters that is active at high

Pi concentrations and at an acidic pH.

***PHO84* supported growth under the broadest range of conditions.**

We defined the pH range at which each triple mutant retaining a single predicted Pi transporter was able

to grow in SC medium with low or high Pi. Growth was assayed by optical density and the area under each

growth curve (AUC) was depicted as a histogram (Fig. 4 and S3 Fig.). Substantial growth was defined as

growth ≥ AUC 5. Overall, we found that growth of Pho84-A (*pho89-/- pho87-/- fgr2-/-*) cells resembled

that of WT (Fig. 4), with growth optima between pH 2 and pH 7 in both high and low Pi conditions.

The role of Pho89 in growth was dependent on pH and Pi availability (Fig. 4). In low Pi, Pho89-A cells grew

equivalently to WT at pH 6 and above. In high Pi , Pho89 supported intermediate levels of growth at more

acidic pH and growth equivalent to WT at pH 6 and above. Pho89 could hence be described as a putative

high-affinity Pi transporter in neutral and alkaline conditions and a low-affinity transporter in acidic

conditions. *C. albicans*, like *S. cerevisiae*, therefore has two high-affinity Pi transporters, Pho84 and Pho89,

with the former having a broad pH activity range including in alkaline conditions, and the latter active at

pH ≥6.

Pho87 and Fgr2 were unable to support substantial growth in low ambient Pi and therefore are low-

affinity Pi transporters. Both supported growth only at acidic pH in high Pi. Pho87-A cells grew to an AUC

≥5 only between pH 2 and 6, and even at their optimal conditions supported only ~70% of the WT growth

(Fig. 4). Fgr2-A cells showed the weakest growth with similar optima to Pho87-A cells, growing to an AUC

≥5 only at pH 3 and 4 (Fig. 4). *C. albicans* therefore has two low-affinity Pi transporters, Pho87 and Fgr2,

with the latter, a Pho84 homolog, showing a narrow, acidic pH optimum.

Hyphal formation of triple Pi transporter mutants largely reflected the growth-sustaining properties of the

transporters. Q- mutants lacking all four Pi transporters failed to form hyphae under any conditions tested

(Fig 2). In contrast, Pho84-A cells encoding only Pho84 formed robust hyphae across all conditions (Fig. 2),

consistent with the strong hyphal defect of *pho84-/-* mutants. Pho89-A cells had severe hyphal growth

defects resembling those of *pho84-/-* mutants. Filamentation of the Pho87-A and Fgr2-A cells resembled

the Q- mutants (Fig. 2). These mutants did not grow sufficiently to form hyphae on RPMI buffered to pH

7 (Fig. 2C). Collectively, these findings are consistent with a concept that hyphal growth requires Pi uptake.

Pho84 showed the most active Pi uptake under all tested pH conditions.

In order to quantify the Pi transport capacity of each transporter, we performed Pi uptake experiments

with the triple mutants that each retained a single predicted transporter, by measuring Pi concentrations

remaining in culture medium over a time course. WT and Pho84-A cells removed Pi from the medium

rapidly and with almost identical kinetics (Fig. 5A,B). Pho89-A cells efficiently transported Pi at a narrow

range of pH 6-8, but their uptake dramatically slowed at pH 5 and below (Fig. 5C,D). These results support

a predominant role of Pho84 as the major Pi importer in *C. albicans,* while Pho89 makes a substantial

contribution around neutral pH.

Low-affinity Pi transporters Pho87 and Fgr2 showed slow Pi uptake under these conditions. Uptake of Pi

by Pho87-A cells was sluggish at pH 2-5 and almost undetectable at pH 6-9 (Fig. 5E). Despite known strong

induction of *FGR2* in low Pi conditions by Pho4, the transcriptional regulator of the PHO regulon [61], Pi

uptake by Fgr2-A cells was weak across the pH levels tested and these cells were unable to fully deplete

Pi from the medium at their pH 4 optimum (Fig. 5F). Both Pho87-A and Fgr2-A cells removed Pi from the

medium most rapidly at pH 4. Still, Pi uptake by Fgr2-A cells was significantly higher than that of Q- cells

at 30 hours (p=0.0001 by two-tailed t-test, Fig. 5G). These data support a role for Pho87 and Fgr2 as Pi

importers albeit with poor kinetics.

To identify any growth defects unrelated to Pi limitation in mutants containing a single Pi transporter, we

compared growth of triple mutants to WT in liquid YPD and Pi replete SC media. YPD contains organic as

well as inorganic phosphate sources, and SC contains high Pi concentrations (7.3 mM) and has an acidic

pH of 4-5 that favors activity of most Pi transporters. No triple mutants exhibited a growth defect in YPD

medium. In SC medium, Pho84-A grew as well as WT while Pho87-A, Pho89-A and Fgr2-A cells grew more

slowly (S4 Fig.), consistent with our previous results (Fig. 1 for SC medium with 7.3 mM Pi at pH 5). These

findings indicate that growth defects in these mutants correspond to a lack of Pi and not a nonspecific

fitness loss.

Glycerophosphocholine transporters provided a minor Pi import function.

To test whether *C. albicans* expresses additional Pi transporters that were not detected by our homology
searches, we next examined Q- cells for their ability to import Pi. At 30 hours, Q- mutants took up ~40%
of the Pi at pH 4 with efficiency being further reduced at pH 2, 5, 6 and 7 (Fig. 6A). These results
demonstrated that another low-capacity Pi transporting activity existed at low pH and high Pi
concentrations.

C. albicans GIT3 and *GIT4* are distant *PHO84* homologs whose products import glycerophosphocholine
(GPC), a phospholipid degradation product that can serve as an organic source of phosphorus [63]. Once
in the cytosol, GPC is metabolized to glycerol, choline and phosphate under Pi limiting conditions [63]. The
GPC transporters Git3 and Git4 share 23% amino acid identity with Pho84 (41% and 39% similarity,
respectively, S1 Fig.). To test whether Git3 and Git4 might contribute to the residual Pi transport in Q-
cells, we competed Pi uptake by Git3 and Git4 with excess GPC in the medium. Addition of a 10-fold excess
of GPC completely eliminated Pi uptake by Q- cells (Fig. 6B). These findings support a role for Git3/4 in Pi
import at high ambient Pi and acidic pH.

We asked whether *C. albicans* has another modality to import Pi from its surroundings, in addition to the
2 high-affinity Pi transporters Pho84 and Pho89, the 2 low-affinity Pi transporters Pho87 and Fgr2, and the
GPC transporters Git3 and Git4. We engineered a septuple mutant strain "*tetO-PHO87"* that lacked three
Pi transporter homologs (*pho84-/- pho89-/- fgr2-/-*), GPC transporters that occupy adjacent loci on
chromosome 5 (*git2-/- git3-/- git4-/-*), and had a single tetracycline-repressible allele of *PHO87* (*pho87-
/tetO-PHO87)*. Mutants were maintained without doxycycline to retain maximal expression of *PHO87*. A
role for Git2 in Pi import is currently not known; *GIT2* was deleted alongside the other two transporters
in their initial characterization and it was included in this construct to permit comparisons with mutants
described in Bishop et al. [63].

We reasoned that if a Pi-transporting activity remained in these cells, they would grow in Pi as their only
source of phosphorus, both in the absence and presence of doxycycline, i.e., during induction and
repression of *PHO87*. On the other hand, if we had mutated all transporters capable of importing sufficient

Pi to sustain growth, doxycycline exposure in SC media, devoid of organic phosphate sources, would

repress growth once internal Pi stores were depleted. We observed the latter result for the *tetO-PHO87*

septuple mutants (*pho87-/tetO-PHO87 pho84-/- pho89-/- fgr2-/- git2-/- git3-/- git4-/-*) (Fig. 6C). In

contrast at pH 3, an optimal pH for Pho87 activity, these mutants grew robustly in media without

doxycycline, YPD or normal SC (which contains high Pi, 7.3 mM) though they had a slight growth defect

compared to WT (Fig. 6C). Thus, the reduced growth of these mutants is largely attributable to Pi

starvation and not a consequence of a nonspecific fitness loss (S5A Fig.). We concluded that minimal Pi

import activity remains when *PHO84*, *PHO89*, *PHO87*, *FGR2* and *GIT2-4* have been genetically eliminated.

In vitro evolution during Pi starvation restored growth in two distinct quadruple Pi transporter mutant

lineages

We reasoned that in vitro evolution of *C. albicans* Q- mutants lacking all four Pi transporter homologs

(*pho84-/- pho89-/- pho87-/- fgr2-/-*) could reveal adaptive mechanisms that facilitate growth during Pi

scarcity. Our previous work showed that *pho84* null mutants' cell wall contained less phosphomannan and

had a thinner outer layer [26], suggesting that modifying cell wall structures while reserving scarce Pi for

essential processes like nucleic acid biosynthesis could sustain Pi-starved cells. We propagated two

lineages of the Q- mutants through 30 serial passages every other day in liquid SC medium with a

moderate concentration of 0.4 mM Pi. To reduce genetic bottlenecks, a large number of cells (~3.5x10^6

cells in 10 μl of saturated culture) were reinoculated into 10 ml of fresh medium. The population at each

passage was saved for DNA extraction and as a glycerol stock. During passaging, the growth rates of both

Q- lineages increased substantially and in distinct increments. Growth rates of the Q- lineages ultimately

plateaued before the end of the experiment. However, fitness recovery in the two lineages during

passaging differed significantly. Lineage 1 (Q- L1), evolving from JKC2830, recovered fitness more rapidly

and attained a growth rate similar to the WT by passage 24 (Fig. 7A). In contrast, growth rates of

populations derived from lineage 2 (Q- L2, evolving from JKC2860) remained lower than WT throughout

the experiment, plateauing at passage 20 (Fig. 7A). Fitness loss of Q- strains and fitness recovery of their

descendant passaged populations was specific to Pi scarcity. In YPD, both Q- strains and their 30[th] passage

descendant populations grew robustly (S5B Fig.). These findings indicated that *C. albicans* cells lacking Pi

transporters could recover fitness during prolonged Pi deprivation, and that this adaption did not lead to

fitness loss when the selective pressure was relieved.

Acquisition and loss of aneuploidies accompanied adaptation of evolving Q- lineages to Pi scarcity.

To uncover the underlying genomic changes associated with improved growth of evolving Q- strains, we

performed whole genome sequencing of the WT ancestral to the Pi transporter mutants (JKC915), as well

as cell populations of selected passages from both lineages, Q- L1 and Q- L2. Populations that bounded

significant fitness gains were chosen for sequencing, generating genomic snapshots of the evolved

population for passages 4, 13, 18, 19, and 30 of Q- L1, and passages 5, 7, 19, 24, and 30 for Q- L2.

Major chromosomal rearrangements occurred during construction of the Q- mutants and during their

passaging. The WT strain used to construct the Q- mutant lineages was diploid with no evidence of

segmental copy number variations or large loss of heterozygosity (LOH) tracts. The genomes of both Q-

strains were trisomic for chromosome 5 (Chr5, Fig. 7B), which includes *GIT2-4*. This aneuploidy was

accompanied by LOH tracts on the left arm of Chr2 (Chr2L) and Chr3R, neither of which include loci

disrupted in the Q- mutant. LOH of Chr2 resulted in homozygosis of the A allele for >50% of the

chromosome, and LOH of Chr3 produced a short tract of homozygosity for B alleles (Fig. 7B).

Both evolving lineages acquired similar large-scale genomic changes during adaptation to Pi scarcity. Each

lineage independently acquired a Chr2 trisomy, first seen in passage 13 and passage 19 for Q- L1 and Q-

L2, respectively (Fig. 8). The Chr2 trisomy was subsequently largely or partially lost in Q- L1 and Q- L2,

respectively, possibly as populations accumulated fitness-enhancing small-scale mutations. The Chr5

trisomy was also lost during passaging of Q- L1, likely due to fitness defects associated with aneuploidy

(Fig. 8).

Persistence of Q- L2's segmental aneuploidy on the left arm of Chr2 (Fig. 8) by the 30th passage suggests

potential adaptive contributions of the encoded loci. This amplified region of Chr2 ranged from nt

1,316,396 to 1,617,025 and encompassed 147 open reading frames. Manual annotation of these genes

(SI Table 1) identified *GIT1,* which encodes a glycerophosphoinositol permease [64] with 22% amino acid

sequence identity to Pho84. There is experimental evidence against a role of Git1 in Pi transport in *C.*

albicans [64] but its role in phosphorus homeostasis might be indirect, i.e. by facilitating

glycerophosphoinositol uptake as Pi starvation induces plasma membrane remodeling [65].

Specific stress phenotypes of Q- population passages did not consistently correspond to their fitness in Pi scarcity.

As we previously found hypersensitivity of *pho84* null mutants to rapamycin as well as oxidative-, cell wall-and membrane stress, induced through plumbagin, micafungin and amphotericin exposure respectively [24, 26], we investigated these responses in the Q- mutants and their evolved lineages. WT, both Q- strains and their passage 30 descendant populations were grown in liquid medium; these strains and in addition, populations from intermediate passages from the evolution experiment were spotted as serial dilutions onto solid medium in the absence or presence of these stressors. Q- mutants were hypersensitive to membrane stress induced by amphotericin and SDS and to cell wall stress induced by micafungin; they showed rapamycin sensitivity corresponding to their growth defects in vehicle (Fig. 9).

In contrast, the responses of evolved, late-passage Q- L1 and Q- L2 populations were distinct for each stressor. Q- L1 but not Q- L2 populations regained growth rates in amphotericin almost to WT levels by passage 30 (Fig. 9A,B). The Q- L2 passage 30 population had increased sensitivity to SDS, compared with its ancestral Q- L2 strain (Fig. 9A,B) while its Q- L1 counterpart regained the ability to grow in the presence of SDS almost to the level of the WT (Fig. 9A,B). In the presence of micafungin, growth of selected passages reflected their fitness in Pi-limited medium (Fig. 9C). However, Q- L1 but not Q- L2 passaged populations regained growth in plumbagin while conversely, Q- L2 populations evolved frank resistance to rapamycin by passage 30 (Fig. 9C). Like the different fitness levels in Pi scarcity reached by the end of the experiment, the distinct stress phenotypes of Pi scarcity-adapted populations suggest their evolutionary trajectories had diverged.

Discussion

In this work, we characterized 4 predicted *C. albicans* Pi transporters, Pho84, Pho89, Pho87 and Fgr2, identified by sequence homology, and determined their contributions to Pi acquisition. In brief, among the high-affinity Pi transporters, Pho84 was the most important for growth, filamentation, stress responses, and induction of TORC1 signaling and had the broadest pH range of Pi uptake capacity, while Pho89 was specialized for uptake in neutral and alkaline pH (Table 1). Among the low-affinity Pi

transporters, both of which were only active at acidic pH, Pho87 was more efficient and had a broader pH

range while Fgr2 functioned only between pH 3 and 5 (Figs. 4, 5). In contrast to *S. cerevisiae*, *C. albicans*

low-affinity Pi transporters are not paralogs; rather, the less efficient one, Fgr2, is a distant Pho84

homolog. The minor role for Fgr2 in *C. albicans* Pi import stands in contrast to *Cryptococcus neoformans*

where both *PHO84* homologs (*PHO84* and *PHO840*) make significant contributions to Pi transport [66]. In

the absence of all specific Pi transporters, glycerophosphocholine transporters were able to provide

residual Pi import to sustain growth. Cells lacking all Pi transporters were able to regain fitness during

sequential passaging in limited Pi, that plateaued at distinct levels for 2 populations in accordance with

previously described declining adaptability [67, 68], while engendering distinct responses to some Pi-

relevant stressors. During the evolution experiment, similar large-scale genomic changes were

sequentially acquired and partially or completely lost in the 2 independently evolving populations.

Retention of a segmental amplification within Chr2 in the less fit lineage suggests that genes in this

trisomic region may contribute to fitness in limited Pi but the precise loci responsible remain undefined.

The severe growth defect of Q- cells that lack all 4 identified Pi transporters argues against the presence

of other specific Pi transporters. Q- cells removed a small fraction of the Pi present in their medium. The

glycerophosphocholine transporters Git3 and 4 provided minor transport activity that was most evident

in the growth difference between a Q- mutant and the septuple mutant when grown in the presence of

doxycycline to repress *PHO87* (Fig. 6C). The ability of GPC provision to completely outcompete measurable

Pi uptake in Q- cells (Fig. 6B) also argues against other cell surface transporters beside Git3 and Git4 playing

a measurable role in Pi uptake under our experimental conditions.

pH sensitivity is a key feature of each transporter. Like WT, Pho84-A, Pho87-A, and Fgr2-A cells grew

optimally between pH 3 and 6. Only cells expressing *PHO89* alone showed a growth optimum between pH

6 and 8. The pH of the oral and pharyngeal mucosa as well as most of the gastrointestinal tract colonized

by *C. albicans* is broadly neutral or alkaline, though mucosal microenvironments may be acidic due to

bacterial metabolites. During acute invasive disease, *C. albicans* finds itself in mildly alkaline blood and

tissue environments between pH 7.35 and 7.45. Pi acquisition systems of *C. albicans* are therefore not

well adapted to host environments encountered during invasive disease. The reduced Pi transport activity

at alkaline pH of the bloodstream might explain why the PHO regulon is induced during systemic disease,

reflecting "alkaline pH-simulated nutrient deprivation" [69], despite the presence of abundant Pi and

organic phosphate compounds like GPC. Pi import is also critical for proliferation of other human fungal

pathogens [30, 70, 71] and unicellular parasites [72] each of which must contend with neutral to alkaline

conditions in host deep organs.

Loss of *PHO84* but not of the other Pi transporters had a substantial effect on TORC1 activity (Fig. 3) and

oxidative stress endurance (S2 Fig.). These experiments cannot distinguish between specific activities of

Pho84 in these cellular functions, versus the predominant role of Pho84 in providing Pi to the cell. We

found in another context that TORC1 activity depends on availability of Pi but not on the presence of

Pho84, while endurance of peroxide stress may require an activity specifically of Pho84 [73]. *C. albicans*

PHO84 transcription is co-regulated by TORC1 in addition to Pho4 [24], but how these systems interact

and modulate each others' outputs remains unknown.

Hyphal growth defects mirrored the severity of Pi transport deficiency among the constructed mutants.

Among the Pi transporters assayed in defined mutants, only Pho84-A cells produced robust hyphae on all

3 media examined (Fig. 2), while Pho87-A cells produced some hyphae on RPMI pH 5, and Pho89-A cells

had sparse, short hyphae on RPMI pH 7 (Fig. 2). The role of Pi uptake in filamentation might be indirect,

through activation of signaling systems like TORC1 required for hyphal morphogenesis [74]. Alternatively,

given the larger surface area of hyphal cells compared to yeast cells, hyphal cells must consume larger

amounts of phosphoric metabolites like nucleotide sugars required as building blocks for the cell wall. The

inability to produce sufficient phosphorus-containing intermediates might inhibit hyphal morphogenesis

to minimize cell wall surface area and preserve phosphorus for other vital functions.

To probe *C. albicans'* options for adaptation to Pi scarcity, we performed an in vitro evolution experiment

with Q- strains in which we had deleted the known Pi transporters. These strains had acquired, at an

unknown point during their construction, a trisomy of Chr5 where the genes encoding organic phosphate

transporters Git3 and Git4 reside. *C. albicans* GPC transporters are upregulated by the transcription factor

Pho4 during Pi starvation [61]. As Q- cells grew very poorly in medium with a moderate Pi content of 0.4

mM, increasing the gene dosage of *GIT3* and *GIT4* by retaining a third copy of Chr5 still did not restore

significant Pi uptake, as shown in Fig. 5G. During the evolution experiment, a further large-scale genomic

alteration appeared in both lineages: triploidy of Chr2 with simultaneous LOH reducing all 3 alleles to AAA

in a long segment on the left arm of the chromosome (cyan-colored segment of Chr2 in both lineages, Fig.

8).

When abruptly exposed to significant stress, *C. albicans* is known to frequently resort to aneuploidy and

LOH to provide a crude but rapid option to improve fitness specific to the particular stress [48-50, 75, 76],

reviewed in [77]. Ploidy increases that enhance fitness under specific stress conditions can be achieved

more rapidly than accumulation of advantageous (in the setting of the specific stress) point mutations

because chromosome missegregation occurs once every 5×10^5 cell divisions (in *S. cerevisia*e) [78] while

substitution of any particular base pair is estimated to occur once every 1.2×10^{10} cell divisions in *C.*

albicans [79] and every 1.67×10^{10} cell divisions in *S. cerevisiae* [80]. Populations under strong selection

are therefore more likely to initially become enriched for aneuploid mutants than for cells containing

constellations of advantageous point mutations [81]. Large-scale copy number variants like trisomies

however incur fitness costs due to increased transcription and translation of a multitude of genes that

result in excess protein production and protein complex stoichiometry imbalances [78, 82, 83]. These

fitness costs predominate when the selective pressure of the stress relents [48, 83]. In addition to

environmental changes that diminish stress intensity, small-scale genomic changes (like single nucleotide

mutations and small insertions or deletions) that promote adaptation to the specific stressor can relieve

selective pressure and favor loss of a trisomic chromosome. For example, a gain-of-function point

mutation of a transcriptional regulator [48, 84], may alleviate a specific stressors' selective pressure to the

point that trisomies resolve back to diploid, as we observed in the Q- L1 but not the Q- L2 lineage.

Evolving quadruple Pi transporter mutants similarly gained trisomies in both independent populations,

highlighting the strong selection experienced by these cells. Loss of trisomies during passaging may have

been enabled by accumulating fitness-enhancing small-scale variants. The distinct mutations that

permitted these adaptive solutions remain to be identified. Given their cell wall- and membrane stress

response phenotypes, it is tempting to speculate that Pi-sparing modifications of membranes and the cell

wall might increase fitness during Pi scarcity. The cyanobacterium *Prochlorococcus,* a dominant species in

waters of the North Pacific Subtropical Gyre, has a competitive advantage in this Pi-scarce ecosystem due

to its membrane composition largely of sulfo- and glycolipids, in which fatty acids are linked to

sulfate/sugar- or sugar-based, instead of phosphate-based, polar head groups [85]; other phytoplankton

use similar adaptations [4]. *C. albicans* also economizes on non-essential uses of Pi by remodeling its

membrane systems: the gene encoding a homolog of diacylglyceryl-*N,N,N*-trimethylhomoserine synthase

is strongly upregulated during Pi starvation in dependence on the PHO regulon [61] and like *Neurospora*

crassa and *C. neoformans, C. albicans* can replace membrane phospholipids with betaine-headgroup lipids

during Pi starvation [65, 86].

A caveat is that the distinct stress phenotypes of the two passage 30 populations could be incidental or

integral to their Pi management strategies. Detailed genotype comparisons and further genetic and cell

biologic analysis of distinct mutations in the two lineages will be required to test causal relationships

between the phenotypes. For example, while distinct SDS stress phenotypes and distinct strategies to Pi

scarcity adaptation could be unrelated, another possibility is that the overall genomic changes that

occurred in the Q- L2 during Pi scarcity adaptation led to membrane changes that rendered it susceptible

to detergent stress from SDS. In contrast, the adaptation trajectory of lineage Q- L1 may have required

less changes to membrane composition so that its SDS endurance was restored as it adapted to growth in

scarce Pi. Genetic, genomic and lipidomic analyses will be required to distinguish these possibilities.

These experiments have several limitations. As noted, there was a residual Pi transport activity in addition

to the predicted Pi transporters present in triple mutant cells due to presence of GPC transporters Git3

and Git4; this residual activity was minor, though, and only detectable at high ambient Pi and acidic pH.

Our experiments permit only semi-quantitative conclusions about the efficiency and optimum of each

transporter, since we did not assay binding affinity and maximal transport velocity with ^{32}P uptake

measurements. Pi uptake experiments in early timepoints are likely not confounded by different growth

rates of each triple mutant expressing a single Pi transporter "alone" because all experiments were

performed at an OD_{600} of 2. However, inefficient Pi uptake may be artifactually amplified by slower growth

of mutants at time points longer than 4-6 hours. Another potential confounder could be differential

expression levels of the 4 transporters, which are likely to vary between pH and Pi concentration

conditions; e.g. the alkali-responsive transcriptional regulator Rim101 induces transcription of *PHO89*

[87], and protein levels of these 4 *C. albicans* transporters remain to be defined. Our in vitro evolution

experiment was limited in that 2 transformants from a final strain-construction transformation step were

evolved without technical replicates; nevertheless, experimental testing of candidate variants will be

required in any case to ascribe fitness roles in Pi scarcity to mutations identified in both or in one of the

phenotypically distinct lineages.

In summary, *C. albicans'* Pi acquisition system is suboptimal for the neutral to alkaline host environments
it typically occupies during invasive disease. Differences in functional optima among transporters may
provide backup mechanisms for Pi transport as *C. albicans* moves through different host niches.
Redistribution of intracellular Pi among organelles and processes may sustain survival during "alkaline pH-
simulated nutrient deprivation" [69], in ways that remain to be elucidated. At the same time, this
redistribution might also render the fungus more sensitive to host-relevant stressors like membrane- and
cell wall stress. Pi acquisition and regulation in humans differs fundamentally from that in fungi; given the
crucial role of phosphorus in structural, metabolic and regulatory processes and in antifungal drug
endurance, definition of these systems could reveal potential fungus-specific drug targets.

Methods

Culture conditions.

Cells were grown in rich complex medium, YPD, defined media synthetic complete (SC) or yeast nitrogen
base (YNB) with 2 % glucose as described in [25, 26]. To provide defined Pi concentrations, Yeast Nitrogen
Base without Amino Acids, without Ammonium sulphate and without Phosphate, supplemented with KCl,
was used (CYN6802, Formedium, Swaffham, Norfolk, England) and supplemented with the indicated
concentrations of KH_2PO_4. Incubation temperatures were 30°C for liquid and solid media except for hyphal
growth assays which were incubated at 37°C.

Strain construction.

C. albicans mutants were generated as in [88]; details of strain construction are provided in S2 Table
(Strains used in this study), S3 Table (Plasmids used in this study) and S4 Table (Oligonucleotides used in
this study). At least 2 and typically 3 independent heterozygous lineages were constructed for each set of
deletion mutants. Only during construction of the Q- strain, 2 isolates were generated from a single *pho84-*
/- pho89-/- pho87-/- fgr2/FGR2 heterozygous parent. At each mutant construction step, we compared
growth phenotypes of independent transformants in synthetic media; isolates with outlier phenotypes
were eliminated from further use, since those presumably were due to unrelated mutations that arose
during transformation. Among isogenic mutants with similar phenotypes, we performed initial phenotypic
characterizations with 2 or more isolates to confirm their similar behaviors. The null mutants for each

predicted transporter were used to examine the role of that transporter in filamentous growth, stress

responses, susceptibility to antifungal agents and TORC1 signaling.

Growth curves.

Cells from glycerol stocks at -80°C were recovered on YPD agar medium for 2 days. Cells were scraped

from the plate and washed twice with NaCl 0.9% and diluted to a final OD$_{600}$ 0.01 in 150 μl medium in flat

bottom 96-well plates. OD$_{600}$ readings were obtained every 15 min in a Biotech Synergy 2 Multi-Mode

Microplate Reader (Winooski, VT, USA). Standard deviations of three technical replicates, representing

separate wells, were calculated, and graphed in Graphpad Prism Version 9.5.1 (528), and displayed as

error bars. At least 3 biological replicates were obtained on different days unless stated otherwise. For

some assays the area under the curve was calculated and graphed using the same software, and the

average from ≥3 biological replicates per condition was graphed.

Phosphate uptake.

Phosphate uptake measurement experiments were performed as in [23] with some modifications. In brief,

cells were incubated with a defined amount of Pi; the rate of Pi removal from the medium corresponds to

the strains' Pi transport capacity. Cells were recovered on YPD plates from glycerol stocks at -80°C for 2

days. SC medium without Pi, buffered at one-unit increments from pH 2 to 8 was inoculated with cells at

an OD$_{600}$ of 2. Cells were given 30 minutes to equilibrate, 1 mM Pi was then added, and the Pi remaining

in the medium was measured every 1-5 h for a period of 6 to 30 hours, depending on each strain or

condition. Samples were harvested at 20,000 rpm for 5 min in the cold room and two technical replicates

per sample (300 μl) were collected from each time point. The total concentration of Pi in the supernatant

was calculated according to Ames [89].

When the assay required pre-feeding the cells with GPC before the addition of Pi, 10 mM of GPC was

added during the 30-minute incubation without Pi. Then Pi was added at a concentration of 1 mM.

Growth of cell dilution spots on solid medium.

Cells recovered from glycerol stock at -80°C were grown on YPD plates for ≥2 days. They were washed in

0.9 % NaCl and diluted in 5- or 3-fold steps from a starting OD$_{600}$ of 0.5 in a microtiter plate, then pin

transferred to agar medium and incubated for 48 h at 30°C.

Hyphal morphogenesis assay.

Cells recovered from glycerol stock at -80°C were grown on YPD plates for 1-2 days, washed and resuspended in 0.9% NaCl to an OD_{600} 0.1. Variation between spots and spot density effects were minimized by spotting 3 µl cell suspensions at 6 equidistant points, using a template around the perimeter of an agar medium plate. Each agar plate contained a WT spot that served as a control to which the other strains on the plate must be compared. This method minimizes variation between colony filamentation within each genotype that occurs when colonies are streaked or plated at varying density and at uncontrolled distances from each other. By including a WT on each plate, we also control for the effects of different hydration states of the agar and slight variations in medium composition which cannot be excluded by other means. RPMI and Spider media were used; the latter is not buffered and has a slightly higher than neutral pH. RPMI medium pH 7 was buffered with 165 mM MOPS; and RPMI pH 5 was buffered with 100 mM MES. Plates were incubated at 37°C for the indicated durations. All panels shown represent at least 3 biological replicates.

Western blot.

Cell lysis and Western blotting were performed as described in [62]. Antibodies used are shown in S5 Table. For densitometry, ImageJ (imagej.net/welcome) software (opensource) was used to quantitate signals obtained from Azure biosystems c600.

Population evolution.

Two Q- mutants (*pho84-/- pho89-/- pho87-/- fgr2-/-*), distinct transformants (Q- L1 and Q- L2)for deletion of the second *FGR2* allele, derived from the same heterozygous strain (JKC2812 *pho84-/- pho89-/- pho87-/- fgr2-/FGR2*)*,* were inoculated into 10 ml SC 0.4 mM Pi at an OD_{600} of 0.05. Cultures were incubated at 30°C at 200 rpm. From each culture, 10 µl were transferred into 10 ml of the same fresh medium every 48 hours, for a total of 30 passages. The culture from each passage was saved as a glycerol stock. Populations from each passage were used in growth curve experiments and were always used after direct revival from frozen stock without further passaging.

Genomic DNA isolation and whole-genome sequencing.

For DNA extraction from cells grown on YPD plates from cells saved at the end of each passage, the Zymo

Quick DNA Fungal/Bacterial Miniprep kit was used according to the manufacturer's instructions.Library

preparation and sequencing was carried out by the Applied Microbiology Services Lab (AMSL) at The Ohio

State University, using the Illumina Nextseq 2000 platform to generate 150 basepair paired-end reads. All

samples were sequenced to a minimum depth of 175x. The reads were trimmed using trimmomatic 0.39

(with default parameters except slidingwindow:4:20, maxinfo:125:1, headcrop:20, and minlen:35) to

remove adaptors and poor quality sequences [90]. Using bowtie2 v2.2.6 [91], the trimmed reads were

aligned against the SC5314 reference genome (version A21-s02-m09-r10) obtained from the Candida

Genome Database (www.candidagenome.org). The aligned SAM files were then converted to the BAM

format using samtools v1.7 [92].

Copy number analysis.

To detect karyotypic changes, pileup data for each whole-genome sequenced strain was obtained using

bbMAP v39.01 [93]. Average read pileup depth highlighted any whole-chromosome aneuploidies. Copy

number variation, including aneuploidies, were further confirmed via visualization in YMAP [94].

Data availability.

The sequencing data are available at the Sequence Read Archive under BioProject Accession Number

PRJNA1035923.

Acknowledgements

We thank Yuping Li for helpful discussions. M.A.-Z. was funded in part by the Alfonso Martin Escudero

Foundation (Madrid, Spain). J.R.K. was funded by Boston Children's Hospital and by NIAID R21AI096054

and NIAID R21AI137716. M.Z.A. was funded by National Institutes of Health grant 1R01AI148788 and an

NSF CAREER Award 2046863. A.M. was supported by a President's Postdoctoral Scholars Program Award

at The Ohio State University. J.P.-V. was funded by NIH R15 GM104876. The funders played no role in the

study design, data collection and analysis, decision to publish, or preparation of the manuscript. The

authors declare no competing interests, financial or non-financial.

 Table 1. Pi transporter characteristics.

	Pho84	Pho89	Pho87	Fgr2
Chromosomal location	Chr1	Chr4	Chr1	Chr7
Pi affinity	high	high	low	low
Optimal pH range	pH 2-8	pH 6-8	pH 2-6	pH 3-4
Role in hyphal growth	+++	+	-	-
Role in Target of Rapamycin Complex1 signaling	Yes	No	No	No
Role in oxidative stress response	Yes	No	No	No

Figure Legends

Fig. 1. Among Pi transporters, Pho84 contributed to growth over the broadest range of tested
conditions. Fivefold dilutions of cells of indicated genotypes were spotted (top to bottom) onto YPD (top,
center) or SC (all others) agar media buffered to indicated pH (3, 5 or 7) with 100 mM MES and containing
indicated Pi concentrations (0.05, 0.5 or 7.3 mM) and grown at 30° C for 2 d. Strains are WT (JKC915);
pho84-/- (JKC1450); *pho89-/-* (JKC2585); *pho87-/-* (JKC2581); *fgr2-/-* (JKC2667); Pho84-A: *pho87-/- pho89-*
/- fgr2-/- PHO84+/+ (JKC2788); Pho89-A: *pho84-/- pho87-/- fgr2-/- PHO89+/+* (JKC2783); Pho87-A: *pho84-*
/- pho89-/- fgr2-/- PHO87+/+ (JKC2777); Fgr2-A: *pho84-/- pho87-/- pho89-/- FGR2+/+* (JKC2758); Q-:
pho84-/- pho87-/- pho89-/- fgr2-/- (JKC2830 and JKC2860). Representative of 3 biological replicates.

Fig. 2. Pho84 was the major contributor to hyphal growth among 4 Pi transporters. Cell suspensions of
indicated genotypes were spotted at equidistant points around the perimeter of Spider (**A**) and RPMI (**B,**
C) agar plates. Photomicrographs of the edge of spots were obtained at 4 days for RPMI and 11 days for
Spider plates. Spot edges were aligned with image frame corners to allow comparisons of hyphal fringes'
length. Strains are WT (JKC915); *pho84-/-* (JKC1450); Pho84-A: *pho87-/- pho89-/- fgr2-/- PHO84+/+*
(JKC2788); *pho89-/-* (JKC2585); Pho89-A: *pho84-/- pho87-/- fgr2-/- PHO89+/+* (JKC2783)*; pho87-/-*
(JKC2581); Pho87-A: *pho84-/- pho89-/- fgr2-/- PHO87+/+* (JKC2777); *fgr2-/-* (JKC2667); Fgr2-A: *pho84-/-*
pho87-/- pho89-/- FGR2+/+ (JKC2758); Q- L1: *pho84-/- pho87-/- pho89-/- fgr2-/-* (JKC2830); Q- L2: *pho84-*
/- pho87-/- pho89-/- fgr2-/- (JKC2860). Size bar 200 µm. Representative of 3 biological replicates.

Fig. 3. Pho84 was required for TORC1 activation. Cells were grown in YNB with indicated Pi concentrations for 90 min. Western blots were probed against phosphorylated Rps6 (P-S6) for monitoring TORC1 activity, and tubulin (Tub) as loading control. Dens: ratio between P-S6 and tubulin signals by densitometry. Representative of 3 biological replicates. Strains are 1: WT (JKC915); 2: *pho84-/-* (JKC1450); 3: *pho87-/-* (JKC2581); 4: *pho89-/-* (JKC2585) and 5: *fgr2-/-* (JKC2667)

Fig. 4. Pi transporters differed for their optimal pH and Pi concentration range while Pho84 was most active overall. Cells expressing the indicated Pi transporter alone among the 4 Pi transporters were inoculated to an OD_{600} of 0.2 into SC medium buffered to the indicated pH with 100 mM MES and grown in a plate reader at 30° C for 20 h. OD_{600} was measured every 15 min and the area under the growth curve was calculated in Graphpad Prism; means of 3 biological replicates are depicted in histograms. See S3 Fig. for the represented growth curves. Error bars represent SD of 3 biological replicates. **A.** SC containing 0.1 mM KH_2PO_4. **B.** SC containing 7.3 mM KH_2PO_4. Strains are WT (JKC915); Pho84-A: *pho87-/- pho89-/- fgr2-/- PHO84+/+* (JKC2788); Pho87-A: *pho84-/- pho89-/- fgr2-/- PHO87+/+* (JKC2777); Pho89-A: *pho84-/- pho87-/- fgr2-/- PHO89+/+* (JKC2783); Fgr2-A: *pho84-/- pho87-/- pho89-/- FGR2+/+* (JKC2758).

Fig. 5. Pi uptake of cells expressing single Pi transporters reflected their growth optima. Cells with indicated genotypes were inoculated into SC without Pi (buffered to pH 1-9) at OD_{600} 2. After 30 minutes, KH_2PO_4 was added to a final concentration of 1 mM, and the extracellular concentration of phosphate was measured in 2 technical replicates at indicated time points. Error bars SD. Representative of three biological replicates. **A.** WT (JKC915). **B.** Pho84-A: *pho87-/-pho89-/- fgr2-/- PHO84+/+* (JKC2788). **C.** Pho89-A in pH 1-5; **D.** Pho89-A in pH 6-9: *pho84-/-pho87-/- fgr2-/- PHO89+/+* (JKC2783). **E.** Pho87-A: *pho84-/-pho89-/- fgr2-/- PHO87+/+* (JKC2777). **F.** Fgr2-A: *pho84-/- pho87-/- pho89-/- FGR2+/+* (JKC2758). **G.** Q- cells (*pho84-/- pho87-/- pho89-/- fgr2-/-*; JKC2830) took up significantly less Pi at pH 4 than Fgr2-A cells at 30 h of incubation. Histograms depict average and SD of 3 biological replicates, *p*=0.0001 (two-tailed t-test).

Fig. 6. Cells lacking 4 Pi transporters showed residual Pi uptake ability that was outcompeted by glycerophosphocholine. A. Pi uptake experiments performed as in Fig. 5 showed that in Q- cells (*pho84-/- pho87-/- pho89-/- fgr2-/-*, JKC2830) residual Pi uptake occurred and was most efficient at pH 4. **B.** Tenfold excess glycerophosphocholine (GPC) inhibited Pi uptake in Q- cells at pH 4. As in Fig. 5, Q- cells

(JKC2830) were inoculated into SC 0 Pi with 10 mM GPC; after 30 minutes, KH_2PO_4 was added to a final

concentration of 1 mM; Pi concentration in the medium was measured with 3 technical replicates at each

time point. Graph shows mean of 3 biological replicates. Error bars SD. **C.** Cells in which *PHO87* is

expressed from repressible *tetO* while the other 3 Pi transporters and *GIT2-4* are deleted (*tetO-*

PHO87/pho87 pho84-/- pho89-/- fgr2-/- git2-4-/-, JKC2969), grew well in the absence of doxycycline but

grew minimally during *tetO* repression in 50 µg/ml doxycycline. WT (JKC915) and *tetO-PHO87* (JKC2969)

were starved for Pi in SC 0 Pi in the presence of 50 µg/ml doxycycline for 48 h. The medium and doxycyline

were replaced every 24 h. Cells were then inoculated at OD 0.01 into SC medium (7.3 mM Pi), buffered to

pH 3 with 100 mM MES, without and with 50 µg/ml Doxycycline. OD_{600} was recorded every 15 min. Error

bars SD of 3 technical replicates. Representative of 3 biological replicates.

Fig. 7. Whole genome sequencing revealed aneuploidies in Q- cells, and evolution of 2 Q- lineages during

Pi scarcity proceeded along distinct trajectories. A. Cells from selected passages of populations evolving

under Pi starvation from Q- isolates JKC2830 and JKC2860 were grown in SC 0.4 mM Pi. Representative

growth curves and corresponding area under the curve (AUC) shown for selected passages. P0 denotes

the Q- isolate before passaging; all experiments after P0 were performed with populations, not with

clones derived from single colonies. Growth curves are representative of 3 biological replicates, except for

passages 4 and 8, which are representative of 2 biological replicates. Error bars SD of 3 technical replicates.

B. YMAP [74] depictions of WGS results of 2 distinct Q- isolates (*pho84-/- pho87-/- pho89-/- fgr2-/-* ,

JKC2830, Q- L1 and JKC2860, Q- L2) showing Chr5 trisomy and loss of heterozygosity of Chr2 and Chr3.

Fig. 8. Whole genome sequencing of selected passages of 2 evolving Q- derived populations showed

distinct trajectories of acquisition and resolution of aneuploidies and loss of heterozygosity. YMAP [74]

depictions of WGS results of populations evolving from JKC2830, Q- L1 and JKC2860, Q- L2 (both *pho84-/-*

pho87-/- pho89-/- fgr2-/-).

Fig. 9. Two evolving lineages of Q- cells showed distinct stressor endurance. A. Growth area under the

curve (AUC) of cells grown in SC 1 mM Pi containing Vehicle (Veh, DMSO), 0.3 µg/ml Amphotericin B

(AmpB) or 0.005% SDS. AmpB representative of 2 and SDS of 3 biological replicates; error bar SD of 3

technical replicates. **B**. Amphotericin B (AmpB) and SDS growth area under the curve (AUC) from panel A

normalized to each strain's vehicle (Veh) control. WT (JKC915); Q- L1 (JKC2830); Q- L1 P30 (JKC2830

passage 30); Q- L2 (JKC2860); Q- L2 P30 (JKC2860 passage 30). AmpB average of 2 and SDS of 3 biological replicates; error bar SD of biological replicates. **C**. Threefold dilutions of cells of indicated genotypes, starting at OD_{600} 0.5, were spotted (top to bottom) onto SC medium containing Vehicle (Veh, H_2O) or 10 ng/ml micafungin (Mica), 15 µM plumbagin (Plum), 50 ng/ml rapamycin (Rapa), and grown at $30^{\circ}C$ for 1 d (Mica), 2 d (Plum), 4 d (Rapa), respectively. Strains are WT (JKC915); Q- L1 (JKC2830) passage 0, 4, 13, 19, 30 and Q- L2 (JKC2860) passages 0, 5, 7, 19, 30; *pho84-/-* (JKC1450). JKC2830 and JKC2860 genotypes are *pho84-/- pho87-/- pho89-/- fgr2-/-*.

S1 Fig. Structural comparisons of Pho84 and its homologs. AlphaFold structure prediction of *S. cerevisiae* Pho84 and *C. albicans* Fgr2 was obtained from the AlphaFold Protein Structure Database (https://alphafold.ebi.ac.uk/) and visualized in ChimeraX-1.4. The coloring of each model is based on a per-residue confidence score (pLDDT): dark blue – very high (pLDDT > 90), light blue – confident (90 > pLDDT > 70), yellow – low (70 > pLDDT > 50), orange – very low (pLDDT < 50). Inorganic phosphate transporters were aligned in MacVector; identical amino acid residues tinted with dark gray and chemically similar ones in light gray.

S2 Fig. Among Pi transporters, only Pho84 was required for oxidative stress endurance. Cell suspensions of the indicated genotypes WT JKC915; *pho84-/-* (JKC1450); *pho87-/-* (JKC2581); *pho89-/-* (JKC2585); *fgr2-/-* (JKC2667) and *git2-4-/-* (JKC2963) were spotted in 3-fold dilution steps onto SC medium with DMSO (Veh) or plumbagin 15 µM. Plates were incubated for 2 days at 30° C. Representative of 3 biological replicates. All spots were on the same plate.

S3 Fig. Individual growth curves summarized in Figure 4. Strains were grown as described in Fig. 4. Shown are 3 biological replicates performed on different days. Error bars SD of 2 or 3 technical replicates. Strains are WT (JKC915); Pho84-A: *pho87-/- pho89-/- fgr2-/- PHO84+/+* (JKC2788); Pho87-A: *pho84-/- pho89-/- fgr2-/- PHO87+/+* (JKC2777); Pho89-A: *pho84-/- pho87-/- fgr2-/- PHO89+/+* (JKC2783)*;* Fgr2-A: *pho84-/- pho87-/- pho89-/- FGR2+/+* (JKC2758).

S4 Fig. Pi transporter triple mutants had no growth defects in rich complex medium. Cells of indicated triple mutant genotypes were grown in YPD (left) and SC (right) and OD_{600} was monitored. Upper panels: strains expressing only one of 2 high-affinity transporters. Lower panels: Strains expressing only one of 2

low-affinity transporters. Pho84-A: *pho87-/- pho89-/- fgr2-/- PHO84+/+* (JKC2788); Pho89-A: *pho84-/-*

pho87-/- fgr2-/- PHO89+/+ (JKC2783); Pho87-A: *pho84-/- pho89-/- fgr2-/- PHO87+/+* (JKC2777); Fgr2-A:

pho84-/- pho87-/- pho89-/- FGR2+/+ (JKC2758). Representative of 3 biological replicates; error bars SD of

3 technical replicates.

S5 Fig. Cells whose single Pi transporter was expressed from *tetO*, as well as Q- cells and their Pi scarcity-

evolved descendant populations had no substantial growth defects in rich complex medium. Strains

were grown as in S4 Fig. **A.** Cells in which a single allele of one Pi transporter, *PHO87*, is expressed from

repressible *tetO*, were grown in YPD and SC without doxycycline and compared with WT and Q- cells. WT

(JKC915), Q- L1 (*pho84-/- pho89-/- pho87-/- fgr2-/-*, JKC2830), *tetO-PHO87* (*tetO-PHO87/pho87 pho84-/-*

pho89-/- fgr2-/- git2-4-/-, JKC2969). **B.** Growth in YPD of WT (JKC915); Q- L1 (JKC2830) and Q- L2 (*pho84-*

/- pho89-/- pho87-/- fgr2-/-, JKC2860). P30: population from the 30th Pi scarcity passage.

References

1.   Pontes MH, Groisman EA. Protein synthesis controls phosphate homeostasis. Genes &
development. 2018;32(1):79-92. Epub 2018/02/14. doi: 10.1101/gad.309245.117. PubMed PMID:
29437726; PubMed Central PMCID: PMCPMC5828397.
2.   Garg A, Sanchez AM, Miele M, Schwer B, Shuman S. Cellular responses to long-term phosphate
starvation of fission yeast: Maf1 determines fate choice between quiescence and death associated with
aberrant tRNA biogenesis. Nucleic acids research. 2023;51(7):3094-115. Epub 2023/02/17. doi:
10.1093/nar/gkad063. PubMed PMID: 36794724; PubMed Central PMCID: PMCPMC10123115.
3.   Ducousso-Detrez A, Fontaine J, Lounes-Hadj Sahraoui A, Hijri M. Diversity of Phosphate Chemical
Forms in Soils and Their Contributions on Soil Microbial Community Structure Changes. Microorganisms.
2022;10(3). Epub 2022/03/27. doi: 10.3390/microorganisms10030609. PubMed PMID: 35336184;
PubMed Central PMCID: PMCPMC8950675.
4.   Van Mooy BA, Fredricks HF, Pedler BE, Dyhrman ST, Karl DM, Koblizek M, et al. Phytoplankton in
the ocean use non-phosphorus lipids in response to phosphorus scarcity. Nature. 2009;458(7234):69-72.
Epub 2009/02/03. doi: 10.1038/nature07659. PubMed PMID: 19182781.
5.   Oliverio AM, Bissett A, McGuire K, Saltonstall K, Turner BL, Fierer N. The Role of Phosphorus
Limitation in Shaping Soil Bacterial Communities and Their Metabolic Capabilities. mBio. 2020;11(5). Epub
2020/10/29. doi: 10.1128/mBio.01718-20. PubMed PMID: 33109755; PubMed Central PMCID:
PMCPMC7593963.
6.   Crook M, Swaminathan R. Disorders of plasma phosphate and indications for its measurement.
Ann Clin Biochem. 1996;33 (Pt 5):376-96. Epub 1996/09/01. doi: 10.1177/000456329603300502.
PubMed PMID: 8888972.
7.   Muchhal US, Pardo JM, Raghothama KG. Phosphate transporters from the higher plant Arabidopsis
thaliana. Proceedings of the National Academy of Sciences of the United States of America.
1996;93(19):10519-23. Epub 1996/09/17. doi: 10.1073/pnas.93.19.10519. PubMed PMID: 8927627;
PubMed Central PMCID: PMCPMC38418.

8.   Mitsukawa N, Okumura S, Shirano Y, Sato S, Kato T, Harashima S, et al. Overexpression of an
Arabidopsis thaliana high-affinity phosphate transporter gene in tobacco cultured cells enhances cell
growth under phosphate-limited conditions. Proceedings of the National Academy of Sciences of the
United States of America. 1997;94(13):7098-102. Epub 1997/06/24. doi: 10.1073/pnas.94.13.7098.
PubMed PMID: 9192698; PubMed Central PMCID: PMCPMC21291.
9.   Berndt T, Kumar R. Novel mechanisms in the regulation of phosphorus homeostasis. Physiology
(Bethesda). 2009;24:17-25. Epub 2009/02/07. doi: 10.1152/physiol.00034.2008. PubMed PMID:
19196648; PubMed Central PMCID: PMCPMC3963411.
10.   Figueres L, Beck-Cormier S, Beck L, Marks J. The Complexities of Organ Crosstalk in Phosphate
Homeostasis: Time to Put Phosphate Sensing Back in the Limelight. Int J Mol Sci. 2021;22(11). Epub
2021/06/03. doi: 10.3390/ijms22115701. PubMed PMID: 34071837; PubMed Central PMCID:
PMCPMC8199323.
11.   Persson BL, Petersson J, Fristedt U, Weinander R, Berhe A, Pattison J. Phosphate permeases of
Saccharomyces cerevisiae: structure, function and regulation. Biochimica et biophysica acta.
1999;1422(3):255-72. Epub 1999/11/05. PubMed PMID: 10548719.
12.   Nieuwenhuis BJ, Borst-Pauwels GW. Derepression of the high-affinity phosphate uptake in the
yeast Saccharomyces cerevisiae. Biochimica et biophysica acta. 1984;770(1):40-6. Epub 1984/02/29. doi:
10.1016/0005-2736(84)90071-3. PubMed PMID: 6365165.
13.   Tamai Y, Toh-e A, Oshima Y. Regulation of inorganic phosphate transport systems in
Saccharomyces cerevisiae. Journal of bacteriology. 1985;164(2):964-8. Epub 1985/11/01. doi:
10.1128/jb.164.2.964-968.1985. PubMed PMID: 3902805; PubMed Central PMCID: PMCPMC214353.
14.   Martinez P, Persson BL. Identification, cloning and characterization of a derepressible Na+-coupled
phosphate transporter in Saccharomyces cerevisiae. Molecular & general genetics : MGG.
1998;258(6):628-38. Epub 1998/07/22. PubMed PMID: 9671031.
15.   Bun-Ya M, Nishimura M, Harashima S, Oshima Y. The PHO84 gene of Saccharomyces cerevisiae
encodes an inorganic phosphate transporter. Molecular and cellular biology. 1991;11(6):3229-38. Epub
1991/06/01. PubMed PMID: 2038328; PubMed Central PMCID: PMC360175.
16.   Berhe A, Fristedt U, Persson BL. Expression and purification of the high-affinity phosphate
transporter of Saccharomyces cerevisiae. Eur J Biochem. 1995;227(1-2):566-72. Epub 1995/01/15. doi:
10.1111/j.1432-1033.1995.tb20426.x. PubMed PMID: 7851439.
17.   Saier MH, Reddy VS, Moreno-Hagelsieb G, Hendargo KJ, Zhang Y, Iddamsetty V, et al. The
Transporter Classification Database (TCDB): 2021 update. Nucleic acids research. 2021;49(D1):D461-D7.
Epub 2020/11/11. doi: 10.1093/nar/gkaa1004. PubMed PMID: 33170213; PubMed Central PMCID:
PMCPMC7778945.
18.   Roomans GM, Blasco F, Borst-Pauwels GW. Cotransport of phosphate and sodium by yeast.
Biochimica et biophysica acta. 1977;467(1):65-71. Epub 1977/05/16. doi: 10.1016/0005-2736(77)90242-
5. PubMed PMID: 16650.
19.   Kanehisa M, Furumichi M, Sato Y, Kawashima M, Ishiguro-Watanabe M. KEGG for taxonomy-based
analysis of pathways and genomes. Nucleic acids research. 2023;51(D1):D587-D92. Epub 2022/10/28. doi:
10.1093/nar/gkac963. PubMed PMID: 36300620; PubMed Central PMCID: PMCPMC9825424.
20.   Harding SD, Armstrong JF, Faccenda E, Southan C, Alexander SPH, Davenport AP, et al. The
IUPHAR/BPS Guide to PHARMACOLOGY in 2024. Nucleic acids research. 2024;52(D1):D1438-D49. Epub
2023/10/29. doi: 10.1093/nar/gkad944. PubMed PMID: 37897341; PubMed Central PMCID:
PMCPMC10767925.
21.   Wykoff DD, O'Shea EK. Phosphate transport and sensing in Saccharomyces cerevisiae. Genetics.
2001;159(4):1491-9. Epub 2002/01/10. PubMed PMID: 11779791; PubMed Central PMCID: PMC1450841.

22.   Hurlimann HC, Stadler-Waibel M, Werner TP, Freimoser FM. Pho91 Is a vacuolar phosphate
transporter that regulates phosphate and polyphosphate metabolism in Saccharomyces cerevisiae.
Molecular biology of the cell. 2007;18(11):4438-45. Epub 2007/09/07. doi: 10.1091/mbc.E07-05-0457.
PubMed PMID: 17804816; PubMed Central PMCID: PMC2043573.
23.   Ghillebert R, Swinnen E, De Snijder P, Smets B, Winderickx J. Differential roles for the low-affinity
phosphate transporters Pho87 and Pho90 in Saccharomyces cerevisiae. The Biochemical journal.
2011;434(2):243-51. Epub 2010/12/15. doi: 10.1042/BJ20101118. PubMed PMID: 21143198.
24.   Liu NN, Flanagan PR, Zeng J, Jani NM, Cardenas ME, Moran GP, et al. Phosphate is the third nutrient
monitored by TOR in Candida albicans and provides a target for fungal-specific indirect TOR inhibition.
Proceedings of the National Academy of Sciences of the United States of America. 2017. Epub 2017/06/02.
doi: 10.1073/pnas.1617799114. PubMed PMID: 28566496.
25.   Liu NN, Uppuluri P, Broggi A, Besold A, Ryman K, Kambara H, et al. Intersection of phosphate
transport, oxidative stress and TOR signalling in Candida albicans virulence. PLoS pathogens.
2018;14(7):e1007076. Epub 2018/07/31. doi: 10.1371/journal.ppat.1007076. PubMed PMID: 30059535;
PubMed Central PMCID: PMCPMC6085062.
26.   Liu NN, Acosta-Zaldivar M, Qi W, Diray-Arce J, Walker LA, Kottom TJ, et al. Phosphoric Metabolites
Link Phosphate Import and Polysaccharide Biosynthesis for Candida albicans Cell Wall Maintenance. mBio.
2020;11(2). Epub 2020/03/19. doi: 10.1128/mBio.03225-19. PubMed PMID: 32184254; PubMed Central
PMCID: PMCPMC7078483.
27.   Lamarche MG, Wanner BL, Crepin S, Harel J. The phosphate regulon and bacterial virulence: a
regulatory network connecting phosphate homeostasis and pathogenesis. FEMS microbiology reviews.
2008;32(3):461-73. Epub 2008/02/06. doi: 10.1111/j.1574-6976.2008.00101.x. PubMed PMID: 18248418.
28.   Santos-Beneit F. The Pho regulon: a huge regulatory network in bacteria. Frontiers in microbiology.
2015;6:402. Epub 2015/05/20. doi: 10.3389/fmicb.2015.00402. PubMed PMID: 25983732; PubMed
Central PMCID: PMC4415409.
29.   Chekabab SM, Harel J, Dozois CM. Interplay between genetic regulation of phosphate homeostasis
and bacterial virulence. Virulence. 2014;5(8):786-93. Epub 2014/12/09. doi: 10.4161/viru.29307. PubMed
PMID: 25483775; PubMed Central PMCID: PMC4601331.
30.   Ikeh M, Ahmed Y, Quinn J. Phosphate Acquisition and Virulence in Human Fungal Pathogens.
Microorganisms. 2017;5(3). Epub 2017/08/23. doi: 10.3390/microorganisms5030048. PubMed PMID:
28829379; PubMed Central PMCID: PMCPMC5620639.
31.   Elliott SR, Tischler AD. Phosphate Starvation: A Novel Signal that Triggers ESX-5 Secretion in
Mycobacterium tuberculosis. Molecular microbiology. 2016. Epub 2016/01/23. doi: 10.1111/mmi.13332.
PubMed PMID: 26800324.
32.   Groschel MI, Sayes F, Simeone R, Majlessi L, Brosch R. ESX secretion systems: mycobacterial
evolution to counter host immunity. Nature reviews Microbiology. 2016;14(11):677-91. Epub 2016/09/27.
doi: 10.1038/nrmicro.2016.131. PubMed PMID: 27665717.
33.   Nussaume L, Kanno S, Javot H, Marin E, Pochon N, Ayadi A, et al. Phosphate Import in Plants: Focus
on the PHT1 Transporters. Front Plant Sci. 2011;2:83. Epub 2011/01/01. doi: 10.3389/fpls.2011.00083.
PubMed PMID: 22645553; PubMed Central PMCID: PMCPMC3355772.
34.   Auesukaree C, Homma T, Kaneko Y, Harashima S. Transcriptional regulation of phosphate-
responsive genes in low-affinity phosphate-transporter-defective mutants in Saccharomyces cerevisiae.
Biochemical and biophysical research communications. 2003;306(4):843-50. Epub 2003/06/25. PubMed
PMID: 12821119.
35.   Wykoff DD, Rizvi AH, Raser JM, Margolin B, O'Shea EK. Positive feedback regulates switching of
phosphate transporters in S. cerevisiae. Molecular cell. 2007;27(6):1005-13. Epub 2007/09/25. doi:
10.1016/j.molcel.2007.07.022. PubMed PMID: 17889672; PubMed Central PMCID: PMC2034509.

36.   Fradin C, De Groot P, MacCallum D, Schaller M, Klis F, Odds FC, et al. Granulocytes govern the
transcriptional response, morphology and proliferation of Candida albicans in human blood. Molecular
microbiology. 2005;56(2):397-415. doi: 10.1111/j.1365-2958.2005.04557.x. PubMed PMID: 15813733.
37.   Thewes S, Kretschmar M, Park H, Schaller M, Filler SG, Hube B. In vivo and ex vivo comparative
transcriptional profiling of invasive and non-invasive Candida albicans isolates identifies genes associated
with tissue invasion. Molecular microbiology. 2007;63(6):1606-28. Epub 2007/03/21. doi: 10.1111/j.1365-
2958.2007.05614.x. PubMed PMID: 17367383.
38.   Zakikhany K, Naglik JR, Schmidt-Westhausen A, Holland G, Schaller M, Hube B. In vivo transcript
profiling of Candida albicans identifies a gene essential for interepithelial dissemination. Cellular
microbiology. 2007;9(12):2938-54. Epub 2007/07/25. doi: 10.1111/j.1462-5822.2007.01009.x. PubMed
PMID: 17645752.
39.   Walker LA, Maccallum DM, Bertram G, Gow NA, Odds FC, Brown AJ. Genome-wide analysis of
Candida albicans gene expression patterns during infection of the mammalian kidney. Fungal genetics and
biology : FG & B. 2009;46(2):210-9. Epub 2008/11/27. doi: 10.1016/j.fgb.2008.10.012. PubMed PMID:
19032986; PubMed Central PMCID: PMCPMC2698078.
40.   Hebecker B, Vlaic S, Conrad T, Bauer M, Brunke S, Kapitan M, et al. Dual-species transcriptional
profiling during systemic candidiasis reveals organ-specific host-pathogen interactions. Scientific reports.
2016;6:36055. Epub 2016/11/04. doi: 10.1038/srep36055. PubMed PMID: 27808111; PubMed Central
PMCID: PMCPMC5093689.
41.   Köhler JR, Acosta-Zaldivar M, Qi W. Phosphate in Virulence of Candida albicans and Candida
glabrata. Journal of fungi (Basel, Switzerland). 2020;6(2). Epub 2020/04/01. doi: 10.3390/jof6020040.
PubMed PMID: 32224872; PubMed Central PMCID: PMCPMC7344514.
42.   Forsburg SL. The art and design of genetic screens: yeast. Nature reviews Genetics. 2001;2(9):659-
68. Epub 2001/09/05. doi: 10.1038/35088500. PubMed PMID: 11533715.
43.   Ottilie S, Goldgof GM, Calvet CM, Jennings GK, LaMonte G, Schenken J, et al. Rapid Chagas Disease
Drug Target Discovery Using Directed Evolution in Drug-Sensitive Yeast. ACS Chem Biol. 2017;12(2):422-
34. Epub 2016/12/16. doi: 10.1021/acschembio.6b01037. PubMed PMID: 27977118; PubMed Central
PMCID: PMCPMC5649375.
44.   Luth MR, Gupta P, Ottilie S, Winzeler EA. Using in Vitro Evolution and Whole Genome Analysis To
Discover Next Generation Targets for Antimalarial Drug Discovery. ACS Infect Dis. 2018;4(3):301-14. Epub
2018/02/17. doi: 10.1021/acsinfecdis.7b00276. PubMed PMID: 29451780; PubMed Central PMCID:
PMCPMC5848146.
45.   Cowell AN, Istvan ES, Lukens AK, Gomez-Lorenzo MG, Vanaerschot M, Sakata-Kato T, et al.
Mapping the malaria parasite druggable genome by using in vitro evolution and chemogenomics. Science.
2018;359(6372):191-9. Epub 2018/01/13. doi: 10.1126/science.aan4472. PubMed PMID: 29326268;
PubMed Central PMCID: PMCPMC5925756.
46.   Antonova-Koch Y, Meister S, Abraham M, Luth MR, Ottilie S, Lukens AK, et al. Open-source
discovery of chemical leads for next-generation chemoprotective antimalarials. Science. 2018;362(6419).
Epub 2018/12/14. doi: 10.1126/science.aat9446. PubMed PMID: 30523084; PubMed Central PMCID:
PMCPMC6516198.
47.   Andries K, Verhasselt P, Guillemont J, Gohlmann HW, Neefs JM, Winkler H, et al. A diarylquinoline
drug active on the ATP synthase of Mycobacterium tuberculosis. Science. 2005;307(5707):223-7. Epub
2004/12/14. doi: 10.1126/science.1106753. PubMed PMID: 15591164.
48.   Selmecki A, Forche A, Berman J. Aneuploidy and isochromosome formation in drug-resistant
Candida albicans. Science. 2006;313(5785):367-70. Epub 2006/07/22. doi: 10.1126/science.1128242.
PubMed PMID: 16857942; PubMed Central PMCID: PMCPMC1717021.

49. Selmecki A, Gerami-Nejad M, Paulson C, Forche A, Berman J. An isochromosome confers drug
resistance in vivo by amplification of two genes, ERG11 and TAC1. Molecular microbiology.
2008;68(3):624-41. Epub 2008/03/28. doi: 10.1111/j.1365-2958.2008.06176.x. PubMed PMID: 18363649.
50. Selmecki AM, Dulmage K, Cowen LE, Anderson JB, Berman J. Acquisition of aneuploidy provides
increased fitness during the evolution of antifungal drug resistance. PLoS genetics. 2009;5(10):e1000705.
Epub 2009/10/31. doi: 10.1371/journal.pgen.1000705. PubMed PMID: 19876375; PubMed Central
PMCID: PMCPMC2760147.
51. Huang M, McClellan M, Berman J, Kao KC. Evolutionary dynamics of Candida albicans during in
vitro evolution. Eukaryotic cell. 2011;10(11):1413-21. Epub 2011/09/06. doi: 10.1128/EC.05168-11.
PubMed PMID: 21890821; PubMed Central PMCID: PMCPMC3209058.
52. Carolus H, Pierson S, Munoz JF, Subotic A, Cruz RB, Cuomo CA, et al. Genome-Wide Analysis of
Experimentally Evolved Candida auris Reveals Multiple Novel Mechanisms of Multidrug Resistance. mBio.
2021;12(2). Epub 2021/04/07. doi: 10.1128/mBio.03333-20. PubMed PMID: 33820824; PubMed Central
PMCID: PMCPMC8092288.
53. Inglis DO, Arnaud MB, Binkley J, Shah P, Skrzypek MS, Wymore F, et al. The Candida genome
database incorporates multiple Candida species: multispecies search and analysis tools with curated gene
and protein information for Candida albicans and Candida glabrata. Nucleic acids research.
2012;40(Database issue):D667-74. Epub 2011/11/09. doi: 10.1093/nar/gkr945. PubMed PMID: 22064862;
PubMed Central PMCID: PMC3245171.
54. Uhl MA, Biery M, Craig N, Johnson AD. Haploinsufficiency-based large-scale forward genetic
analysis of filamentous growth in the diploid human fungal pathogen C.albicans. The EMBO journal.
2003;22(11):2668-78. Epub 2003/05/30. doi: 10.1093/emboj/cdg256. PubMed PMID: 12773383; PubMed
Central PMCID: PMC156753.
55. Wang JM, Woodruff AL, Dunn MJ, Fillinger RJ, Bennett RJ, Anderson MZ. Intraspecies
Transcriptional Profiling Reveals Key Regulators of Candida albicans Pathogenic Traits. mBio. 2021;12(2).
Epub 2021/04/22. doi: 10.1128/mBio.00586-21. PubMed PMID: 33879584; PubMed Central PMCID:
PMCPMC8092256.
56. Shen J, Guo W, Köhler JR. CaNAT1, a heterologous dominant selectable marker for transformation
of Candida albicans and other pathogenic Candida species. Infection and immunity. 2005;73(2):1239-42.
Epub 2005/01/25. doi: 10.1128/IAI.73.2.1239-1242.2005. PubMed PMID: 15664973; PubMed Central
PMCID: PMC547112.
57. Lo HJ, Köhler JR, DiDomenico B, Loebenberg D, Cacciapuoti A, Fink GR. Nonfilamentous C. albicans
mutants are avirulent. Cell. 1997;90(5):939-49. Epub 1997/09/23. PubMed PMID: 9298905.
58. Uppuluri P, Chaturvedi AK, Jani N, Pukkila-Worley R, Monteagudo C, Mylonakis E, et al. Physiologic
expression of the Candida albicans pescadillo homolog is required for virulence in a murine model of
hematogenously disseminated candidiasis. Eukaryotic cell. 2012;11(12):1552-6. Epub 2012/10/30. doi:
10.1128/EC.00171-12. PubMed PMID: 23104566; PubMed Central PMCID: PMC3536276.
59. Wang Y, Zhou J, Zou Y, Chen X, Liu L, Qi W, et al. Fungal commensalism modulated by a dual-action
phosphate transceptor. Cell Rep. 2022;38(4):110293. Epub 2022/01/27. doi:
10.1016/j.celrep.2021.110293. PubMed PMID: 35081357.
60. Qi W, Acosta-Zaldivar M, Flanagan PR, Liu NN, Jani N, Fierro JF, et al. Stress- and metabolic
responses of Candida albicans require Tor1 kinase N-terminal HEAT repeats. PLoS pathogens.
2022;18(6):e1010089. Epub 2022/06/11. doi: 10.1371/journal.ppat.1010089. PubMed PMID: 35687592;
PubMed Central PMCID: PMCPMC9223334.
61. Ikeh MA, Kastora SL, Day AM, Herrero-de-Dios CM, Tarrant E, Waldron KJ, et al. Pho4 mediates
phosphate acquisition in Candida albicans and is vital for stress resistance and metal homeostasis.

Molecular biology of the cell. 2016. Epub 2016/07/08. doi: 10.1091/mbc.E16-05-0266. PubMed PMID: 27385340.

62.   Chowdhury T, Köhler JR. Ribosomal protein S6 phosphorylation is controlled by TOR and modulated by PKA in Candida albicans. Molecular microbiology. 2015;98(2):384-402. Epub 2015/07/16. doi: 10.1111/mmi.13130. PubMed PMID: 26173379; PubMed Central PMCID: PMC4631378.

63.   Bishop AC, Ganguly S, Solis NV, Cooley BM, Jensen-Seaman MI, Filler SG, et al. Glycerophosphocholine utilization by Candida albicans: role of the Git3 transporter in virulence. The Journal of biological chemistry. 2013;288(47):33939-52. Epub 2013/10/12. doi: 10.1074/jbc.M113.505735. PubMed PMID: 24114876; PubMed Central PMCID: PMCPMC3837134.

64.   Bishop AC, Sun T, Johnson ME, Bruno VM, Patton-Vogt J. Robust utilization of phospholipase-generated metabolites, glycerophosphodiesters, by Candida albicans: role of the CaGit1 permease. Eukaryotic cell. 2011;10(12):1618-27. Epub 2011/10/11. doi: 10.1128/EC.05160-11. PubMed PMID: 21984707; PubMed Central PMCID: PMCPMC3232722.

65.   Riekhof WR, Naik S, Bertrand H, Benning C, Voelker DR. Phosphate starvation in fungi induces the replacement of phosphatidylcholine with the phosphorus-free betaine lipid diacylglyceryl-N,N,N-trimethylhomoserine. Eukaryotic cell. 2014;13(6):749-57. Epub 2014/04/15. doi: 10.1128/EC.00004-14. PubMed PMID: 24728191; PubMed Central PMCID: PMCPMC4054272.

66.   Kretschmer M, Reiner E, Hu G, Tam N, Oliveira DL, Caza M, et al. Defects in phosphate acquisition and storage influence virulence of Cryptococcus neoformans. Infection and immunity. 2014;82(7):2697-712. Epub 2014/04/09. doi: 10.1128/IAI.01607-14. PubMed PMID: 24711572; PubMed Central PMCID: PMC4097646.

67.   Couce A, Tenaillon OA. The rule of declining adaptability in microbial evolution experiments. Front Genet. 2015;6:99. Epub 2015/03/31. doi: 10.3389/fgene.2015.00099. PubMed PMID: 25815007; PubMed Central PMCID: PMCPMC4356158.

68.   Kryazhimskiy S, Rice DP, Jerison ER, Desai MM. Microbial evolution. Global epistasis makes adaptation predictable despite sequence-level stochasticity. Science. 2014;344(6191):1519-22. Epub 2014/06/28. doi: 10.1126/science.1250939. PubMed PMID: 24970088; PubMed Central PMCID: PMCPMC4314286.

69.   Lev S, Djordjevic JT. Why is a functional PHO pathway required by fungal pathogens to disseminate within a phosphate-rich host: A paradox explained by alkaline pH-simulated nutrient deprivation and expanded PHO pathway function. PLoS pathogens. 2018;14(6):e1007021. Epub 2018/06/22. doi: 10.1371/journal.ppat.1007021. PubMed PMID: 29928051; PubMed Central PMCID: PMCPMC6013017.

70.   He BZ, Zhou X, O'Shea EK. Evolution of reduced co-activator dependence led to target expansion of a starvation response pathway. eLife. 2017;6. Epub 2017/05/10. doi: 10.7554/eLife.25157. PubMed PMID: 28485712; PubMed Central PMCID: PMCPMC5446240.

71.   Bhalla K, Qu X, Kretschmer M, Kronstad JW. The phosphate language of fungi. Trends in microbiology. 2022;30(4):338-49. Epub 2021/09/05. doi: 10.1016/j.tim.2021.08.002. PubMed PMID: 34479774; PubMed Central PMCID: PMCPMC8882699.

72.   Lacerda-Abreu MA, Dick CF, Meyer-Fernandes JR. The Role of Inorganic Phosphate Transporters in Highly Proliferating Cells: From Protozoan Parasites to Cancer Cells. Membranes (Basel). 2022;13(1). Epub 2023/01/22. doi: 10.3390/membranes13010042. PubMed PMID: 36676849; PubMed Central PMCID: PMCPMC9860751.

73.   King WR, Acosta-Zaldivar M, Qi W, Cherico N, Cooke L, Kohler JR, et al. Glycerophosphocholine provision rescues Candida albicans growth and signaling phenotypes associated with phosphate limitation. mSphere. 2023:e0023123. Epub 2023/10/16. doi: 10.1128/msphere.00231-23. PubMed PMID: 37843297.

74.   Cutler NS, Pan X, Heitman J, Cardenas ME. The TOR signal transduction cascade controls cellular
differentiation in response to nutrients. Molecular biology of the cell. 2001;12(12):4103-13. Epub
2001/12/12. PubMed PMID: 11739804; PubMed Central PMCID: PMC60779.
75.   Anderson MZ, Saha A, Haseeb A, Bennett RJ. A chromosome 4 trisomy contributes to increased
fluconazole resistance in a clinical isolate of Candida albicans. Microbiology (Reading). 2017;163(6):856-
65. Epub 2017/06/24. doi: 10.1099/mic.0.000478. PubMed PMID: 28640746; PubMed Central PMCID:
PMCPMC5737213.
76.   Yang F, Gritsenko V, Slor Futterman Y, Gao L, Zhen C, Lu H, et al. Tunicamycin Potentiates
Antifungal Drug Tolerance via Aneuploidy in Candida albicans. mBio. 2021;12(4):e0227221. Epub
2021/09/02. doi: 10.1128/mBio.02272-21. PubMed PMID: 34465026; PubMed Central PMCID:
PMCPMC8406271.
77.   Bennett RJ, Forche A, Berman J. Rapid mechanisms for generating genome diversity: whole ploidy
shifts, aneuploidy, and loss of heterozygosity. Cold Spring Harbor perspectives in medicine. 2014;4(10).
Epub 2014/08/02. doi: 10.1101/cshperspect.a019604. PubMed PMID: 25081629; PubMed Central PMCID:
PMCPMC4200206.
78.   Torres EM, Sokolsky T, Tucker CM, Chan LY, Boselli M, Dunham MJ, et al. Effects of aneuploidy on
cellular physiology and cell division in haploid yeast. Science. 2007;317(5840):916-24. Epub 2007/08/19.
doi: 10.1126/science.1142210. PubMed PMID: 17702937.
79.   Ene IV, Farrer RA, Hirakawa MP, Agwamba K, Cuomo CA, Bennett RJ. Global analysis of mutations
driving microevolution of a heterozygous diploid fungal pathogen. Proceedings of the National Academy
of Sciences of the United States of America. 2018;115(37):E8688-E97. Epub 2018/08/29. doi:
10.1073/pnas.1806002115. PubMed PMID: 30150418; PubMed Central PMCID: PMCPMC6140516.
80.   Zhu YO, Siegal ML, Hall DW, Petrov DA. Precise estimates of mutation rate and spectrum in yeast.
Proceedings of the National Academy of Sciences of the United States of America. 2014;111(22):E2310-8.
Epub 2014/05/23. doi: 10.1073/pnas.1323011111. PubMed PMID: 24847077; PubMed Central PMCID:
PMCPMC4050626.
81.   Yona AH, Manor YS, Herbst RH, Romano GH, Mitchell A, Kupiec M, et al. Chromosomal duplication
is a transient evolutionary solution to stress. Proceedings of the National Academy of Sciences of the
United States of America. 2012;109(51):21010-5. Epub 2012/12/01. doi: 10.1073/pnas.1211150109.
PubMed PMID: 23197825; PubMed Central PMCID: PMCPMC3529009.
82.   Tang YC, Amon A. Gene copy-number alterations: a cost-benefit analysis. Cell. 2013;152(3):394-
405. Epub 2013/02/05. doi: 10.1016/j.cell.2012.11.043. PubMed PMID: 23374337; PubMed Central
PMCID: PMCPMC3641674.
83.   Yang F, Todd RT, Selmecki A, Jiang YY, Cao YB, Berman J. The fitness costs and benefits of trisomy
of each Candida albicans chromosome. Genetics. 2021;218(2). Epub 2021/04/11. doi:
10.1093/genetics/iyab056. PubMed PMID: 33837402; PubMed Central PMCID: PMCPMC8225349.
84.   Coste A, Turner V, Ischer F, Morschhauser J, Forche A, Selmecki A, et al. A mutation in Tac1p, a
transcription factor regulating CDR1 and CDR2, is coupled with loss of heterozygosity at chromosome 5 to
mediate antifungal resistance in Candida albicans. Genetics. 2006;172(4):2139-56. Epub 2006/02/03. doi:
10.1534/genetics.105.054767. PubMed PMID: 16452151; PubMed Central PMCID: PMCPMC1456413.
85.   Van Mooy BA, Rocap G, Fredricks HF, Evans CT, Devol AH. Sulfolipids dramatically decrease
phosphorus demand by picocyanobacteria in oligotrophic marine environments. Proceedings of the
National Academy of Sciences of the United States of America. 2006;103(23):8607-12. Epub 2006/05/30.
doi: 10.1073/pnas.0600540103. PubMed PMID: 16731626; PubMed Central PMCID: PMCPMC1482627.
86.   Lev S, Rupasinghe T, Desmarini D, Kaufman-Francis K, Sorrell TC, Roessner U, et al. The PHO
signaling pathway directs lipid remodeling in Cryptococcus neoformans via DGTS synthase to recycle

phosphate during phosphate deficiency. PloS one. 2019;14(2):e0212651. Epub 2019/02/23. doi:
10.1371/journal.pone.0212651. PubMed PMID: 30789965; PubMed Central PMCID: PMCPMC6383925.
87.   Bensen ES, Martin SJ, Li M, Berman J, Davis DA. Transcriptional profiling in Candida albicans reveals
new adaptive responses to extracellular pH and functions for Rim101p. Molecular microbiology.
2004;54(5):1335-51. Epub 2004/11/24. doi: 10.1111/j.1365-2958.2004.04350.x. PubMed PMID:
15554973.
88.   Shen J, Cowen LE, Griffin AM, Chan L, Köhler JR. The Candida albicans pescadillo homolog is
required for normal hypha-to-yeast morphogenesis and yeast proliferation. Proceedings of the National
Academy of Sciences of the United States of America. 2008;105(52):20918-23. Epub 2008/12/17. doi:
10.1073/pnas.0809147105. PubMed PMID: 19075239; PubMed Central PMCID: PMC2634893.
89.   Ames BN. Assay of inorganic phosphate, total phosphate and phosphatases. Methods Enzymol.
1966;8:115-8.
90.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30(15):2114-20. Epub 2014/04/04. doi: 10.1093/bioinformatics/btu170. PubMed
PMID: 24695404; PubMed Central PMCID: PMCPMC4103590.
91.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature methods.
2012;9(4):357-9. Epub 2012/03/06. doi: 10.1038/nmeth.1923. PubMed PMID: 22388286; PubMed Central
PMCID: PMCPMC3322381.
92.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map
format and SAMtools. Bioinformatics. 2009;25(16):2078-9. Epub 2009/06/10. doi:
10.1093/bioinformatics/btp352. PubMed PMID: 19505943; PubMed Central PMCID: PMCPMC2723002.
93.   Bushnell B. BBMap. Available from: sourceforge.net/projects/bbmap/.
94.   Abbey DA, Funt J, Lurie-Weinberger MN, Thompson DA, Regev A, Myers CL, et al. YMAP: a pipeline
for visualization of copy number variation and loss of heterozygosity in eukaryotic pathogens. Genome
Med. 2014;6(11):100. Epub 2014/12/17. doi: 10.1186/s13073-014-0100-8. PubMed PMID: 25505934;
PubMed Central PMCID: PMCPMC4263066.
1109

Fig. 1. Among Pi transporters, Pho84 contributed to growth over the broadest range of tested conditions. Fivefold dilutions of cells of indicated genotypes were spotted (top to bottom) onto YPD (top, center) or SC (all others) agar media buffered to indicated pH (3, 5 or 7) with 100 mM MES and containing indicated Pi concentrations (0.05, 0.5 or 7.3 mM) and grown at 30° C for 2 d. Strains are WT (JKC915); *pho84-/-* (JKC1450); *pho89-/-* (JKC2585); *pho87-/-* (JKC2581); *fgr2-/-* (JKC2667); Pho84-A: *pho87-/- pho89-/- fgr2-/- PHO84+/+* (JKC2788); Pho89-A: *pho84-/- pho87-/- fgr2-/- PHO89+/+* (JKC2783); Pho87-A: *pho84-/- pho89-/- fgr2-/- PHO87+/+* (JKC2777); Fgr2-A: *pho84-/- pho87-/- pho89-/- FGR2+/+* (JKC2758); Q-: *pho84-/- pho87-/- pho89-/- fgr2-/-* (JKC2830 and JKC2860). Representative of 3 biological replicates.

Fig. 2. Pho84 was the major contributor to hyphal growth among 4 Pi transporters. Cell suspensions of indicated genotypes were spotted at equidistant points around the perimeter of Spider (**A**) and RPMI (**B, C**) agar plates. Photomicrographs of the edge of spots were obtained at 4 days for RPMI and 11 days for Spider plates. Spot edges were aligned with image frame corners to allow comparisons of hyphal fringes' length. Strains are WT (JKC915); *pho84-/-* (JKC1450); Pho84-A: *pho87-/- pho89-/- fgr2-/- PHO84+/+* (JKC2788); *pho89-/-* (JKC2585); Pho89-A: *pho84-/- pho87-/- fgr2-/- PHO89+/+* (JKC2783)*; pho87-/-* (JKC2581); Pho87-A: *pho84-/- pho89-/- fgr2-/- PHO87+/+* (JKC2777); *fgr2-/-* (JKC2667); Fgr2-A: *pho84-/- pho87-/- pho89-/- FGR2+/+* (JKC2758); Q-L1: *pho84-/- pho87-/- pho89-/- fgr2-/-* (JKC2830); Q-L2: *pho84-/- pho87-/- pho89-/- fgr2-/-* (JKC2860). Size bar 200 µm. Representative of 3 biological replicates.

Fig. 3. Pho84 was required for TORC1 activation. Cells were grown in YNB with indicated Pi concentrations for 90 min. Western blots were probed against phosphorylated Rps6 (P-S6) for monitoring TORC1 activity, and tubulin (Tub) as loading control. Dens: ratio between P-S6 and tubulin signals by densitometry. Representative of 3 biological replicates. Strains are 1: WT (JKC915); 2: *pho84-/-* (JKC1450); 3: *pho87-/-* (JKC2581); 4: *pho89-/-* (JKC2585) and 5: *fgr2-/-* (JKC2667)

Fig. 4. Pi transporters differed for their optimal pH and Pi concentration range while Pho84 was most active overall. Cells expressing the indicated Pi transporter alone among the 4 Pi transporters were inoculated to an OD_{600} of 0.2 into SC medium buffered to the indicated pH with 100 mM MES and grown in a plate reader at 30° C for 20 h. OD_{600} was measured every 15 min and the area under the growth curve was calculated in Graphpad Prism; means of 3 biological replicates are depicted in histograms. See S3 Fig. for the represented growth curves. Error bars represent SD of 3 biological replicates. **A.** SC containing 0.1 mM KH_2PO_4. **B.** SC containing 7.3 mM KH_2PO_4. Strains are WT (JKC915); Pho84-A: *pho87-/- pho89-/- fgr2-/- PHO84+/+* (JKC2788); Pho87-A: *pho84-/- pho89-/- fgr2-/- PHO87+/+* (JKC2777); Pho89-A: *pho84-/- pho87-/- fgr2-/- PHO89+/+* (JKC2783); Fgr2-A: *pho84-/- pho87-/- pho89-/- FGR2+/+* (JKC2758).

Fig. 5. Pi uptake of cells expressing single Pi transporters reflected their growth optima. Cells with indicated genotypes were inoculated into SC without Pi (buffered to pH 1-9) at OD_{600} 2. After 30 minutes, KH_2PO_4 was added to a final concentration of 1 mM, and the extracellular concentration of phosphate was measured in 2 technical replicates at indicated time points. Error bars SD. Representative of three biological replicates. **A.** WT (JKC915). **B.** Pho84-A: *pho87-/-pho89-/- fgr2-/- PHO84+/+* (JKC2788). **C.** Pho89-A in pH 1-5; **D.** Pho89-A in pH 6-9: *pho84-/-pho87-/- fgr2-/- PHO89+/+* (JKC2783). **E.** Pho87-A: *pho84-/-pho89-/- fgr2-/- PHO87+/+* (JKC2777). **F.** Fgr2-A: *pho84-/- pho87-/- pho89-/- FGR2+/+* (JKC2758). **G.** Q- cells (*pho84-/-pho87-/- pho89-/- fgr2-/-;* JKC2830) took up significantly less Pi at pH 4 than Fgr2-A cells at 30 h of incubation. Histograms depict average and SD of 3 biological replicates, *p*=0.0001 (two-tailed t-test).

Fig. 6. Cells lacking 4 Pi transporters showed residual Pi uptake ability that was outcompeted by glycerophosphocholine. A. Pi uptake experiments performed as in Fig. 5 showed that in Q- cells (*pho84-/- pho87-/- pho89-/- fgr2-/-*, JKC2830) residual Pi uptake occurred and was most efficient at pH 4. **B.** Tenfold excess glycerophosphocholine (GPC) inhibited Pi uptake in Q- cells at pH 4. As in Fig. 5, Q- cells (JKC2830) were inoculated into SC 0 Pi with 10 mM GPC; after 30 minutes, KH_2PO_4 was added to a final concentration of 1 mM; Pi concentration in the medium was measured with 3 technical replicates at each time point. Graph shows mean of 3 biological replicates. Error bars SD. **C.** Cells in which *PHO87* is expressed from repressible *tetO* while the other 3 Pi transporters and *GIT2-4* are deleted (*tetO-PHO87/pho87 pho84-/- pho89-/- fgr2-/- git2-4-/-,* JKC2969), grew well in the absence of doxycycline but grew minimally during *tetO* repression in 50 μg/ml doxycycline. WT (JKC915) and *tetO-PHO87* (JKC2969) were starved for Pi in SC 0 Pi in the presence of 50 μg/ml doxycycline for 48 h. The medium and doxycyline were replaced every 24 h. Cells were then inoculated at OD 0.01 into SC medium (7.3 mM Pi), buffered to pH 3 with 100 mM MES, without and with 50 μg/ml Doxycycline. OD_{600} was recorded every 15 min. Error bars SD of 3 technical replicates. Representative of 3 biological replicates.

Fig. 7. Whole genome sequencing revealed aneuploidies in Q- cells, and evolution of 2 Q- lineages during Pi scarcity proceeded along distinct trajectories. A. Cells from selected passages of populations evolving under Pi starvation from Q- isolates JKC2830 and JKC2860 were grown in SC 0.4 mM Pi. Representative growth curves and corresponding area under the curve (AUC) shown for selected passages. P0 denotes the Q- isolate before passaging; all experiments after P0 were performed with populations, not with clones derived from single colonies. Growth curves are representative of 3 biological replicates, except for passages 4 and 8, which are representative of 2 biological replicates. Error bars SD of 3 technical replicates. **B**. YMAP [74] depictions of WGS results of 2 distinct Q- isolates (*pho84-/- pho87-/- pho89-/- fgr2-/-* , JKC2830, Q-L1 and JKC2860, Q-L2) showing Chr5 trisomy and loss of heterozygosity of Chr2 and Chr3.

Fig. 8. Whole genome sequencing of selected passages of 2 evolving Q- derived populations showed distinct trajectories of acquisition and resolution of aneuploidies and loss of heterozygosity. YMAP [74] depictions of WGS results of populations evolving from JKC2830, Q-L1 and JKC2860, Q-L2 (both *pho84-/- pho87-/- pho89-/- fgr2-/-*).

Fig. 9. Two evolving lineages of Q- cells showed distinct stressor endurance. A. Growth area under the curve (AUC) of cells grown in SC 1 mM Pi containing Vehicle (Veh, DMSO), 0.3 μg/ml Amphotericin B (AmpB) or 0.005% SDS. AmpB representative of 2 and SDS of 3 biological replicates; error bar SD of 3 technical replicates. **B**. Amphotericin B (AmpB) and SDS growth area under the curve (AUC) from panel A normalized to each strain's vehicle (Veh) control. WT (JKC915); Q-L1 (JKC2830); Q-L1 P30 (JKC2830 passage 30); Q-L2 (JKC2860); Q-L2 P30 (JKC2860 passage 30). AmpB average of 2 and SDS of 3 biological replicates; error bar SD of biological replicates. **C**. Threefold dilutions of cells of indicated genotypes, starting at OD$_{600}$ 0.5, were spotted (top to bottom) onto SC medium containing Vehicle (Veh, H$_2$O) or 10 ng/ml micafungin (Mica), 15 μM plumbagin (Plum), 50 ng/ml rapamycin (Rapa), and grown at 30°C for 1 d (Mica), 2 d (Plum), 4 d (Rapa), respectively. Strains are WT (JKC915); Q-L1 (JKC2830) passage 0, 4, 13, 19, 30 and Q-L2 (JKC2860) passages 0, 5, 7, 19, 30; *pho84-/-* (JKC1450). JKC2830 and JKC2860 genotypes are *pho84-/- pho87-/- pho89-/- fgr2-/-*.

S1 Fig. Structural comparisons of Pho84 and its homologs. AlphaFold structure prediction of *S. cerevisiae* Pho84 and *C. albicans* Fgr2 was obtained from the AlphaFold Protein Structure Database (https://alphafold.ebi.ac.uk/) and visualized in ChimeraX-1.4. The coloring of each model is based on a per-residue confidence score (pLDDT): dark blue – very high (pLDDT > 90), light blue – confident (90 > pLDDT > 70), yellow – low (70 > pLDDT > 50), orange – very low (pLDDT < 50). Inorganic phosphate transporters were aligned in MacVector; identical amino acid residues tinted with dark gray and chemically similar ones in light gray.

S2 Fig. Among Pi transporters, only Pho84 was required for oxidative stress endurance. Cell suspensions of the indicated genotypes WT JKC915; *pho84-/-* (JKC1450); *pho87-/-* (JKC2581); *pho89-/-* (JKC2585); *fgr2-/-* (JKC2667) and *git2-4-/-* (JKC2963) were spotted in 3-fold dilution steps onto SC medium with DMSO (Veh) or plumbagin 15 µM. Plates were incubated for 2 days at 30° C. Representative of 3 biological replicates. All spots were on the same plate.

S3 Fig. Individual growth curves summarized in Figure 4. Strains were grown as described in Fig. 4. Shown are 3 biological replicates performed on different days. Error bars SD of 2 or 3 technical replicates. Strains are WT (JKC915); Pho84-A: *pho87-/- pho89-/- fgr2-/- PHO84+/+* (JKC2788); Pho87-A: *pho84-/- pho89-/- fgr2-/- PHO87+/+* (JKC2777); Pho89-A: *pho84-/- pho87-/- fgr2-/- PHO89+/+* (JKC2783); Fgr2-A: *pho84-/- pho87-/- pho89-/- FGR2+/+* (JKC2758).

S4 Fig. Pi transporter triple mutants had no growth defects in rich complex medium. Cells of indicated triple mutant genotypes were grown in YPD (left) and SC (right) and OD_{600} was monitored. Upper panels: strains expressing only one of 2 high-affinity transporters. Lower panels: Strains expressing only one of 2 low-affinity transporters. Pho84-A: *pho87-/- pho89-/- fgr2-/- PHO84+/+* (JKC2788); Pho89-A: *pho84-/- pho87-/- fgr2-/- PHO89+/+* (JKC2783); Pho87-A: *pho84-/- pho89-/- fgr2-/- PHO87+/+* (JKC2777); Fgr2-A: *pho84-/- pho87-/- pho89-/- FGR2+/+* (JKC2758). Representative of 3 biological replicates; error bars SD of 3 technical replicates.

S5 Fig. Cells whose single Pi transporter was expressed from *tetO*, as well as Q- cells and their Pi scarcity-evolved descendant populations had no substantial growth defects in rich complex medium. Strains were grown as in S4 Fig. **A.** Cells in which a single allele of one Pi transporter, *PHO87*, is expressed from repressible *tetO*, were grown in YPD and SC without doxycycline and compared with WT and Q-cells. WT (JKC915), Q-L1 (*pho84-/- pho89-/- pho87-/- fgr2-/-*, JKC2830), *tetO-PHO87* (*tetO-PHO87/pho87 pho84-/- pho89-/- fgr2-/- git2-4-/-*, JKC2969). **B.** Growth in YPD of WT (JKC915); Q-L1 (JKC2830) and Q-L2 (*pho84-/- pho89-/- pho87-/- fgr2-/-*, JKC2860). P30: population from the 30[th] Pi scarcity passage.