The collective application of shorebird tracking data to conservation

Abstract

Addressing urgent conservation issues, like the drastic declines of North American migratory birds,

requires creative, evidence-based, efficient, and collaborative approaches. Over 50% of monitored North

American shorebird populations have lost over 50% of their abundance since 1980. To address these

declines, we developed a partnership of scientists and practitioners called the Shorebird Science and

Conservation Collective (hereinafter "the Collective"). Here, we present this successful case study as an

example for others engaged in translational science. The Collective acts as an intermediary whereby

dedicated staff collate and analyze data contributions from scientists to support knowledge requests from

conservation practitioners. Data contributions from 74 organizations include over 6.7 million shorebird

locations forming movement paths of 3,345 individuals representing 36 species tracked across the

Americas. We describe the founding and structure of the Collective and conservation activities we

supported in our first two years. As the volume of scientific data on animal movements continues to grow,

groups like the Collective can be vital liaisons to rapidly integrate and interpret research to support

conservation action.

Autumn-Lynn Harrison[1*†], Candace Stenzel[1‡], Alexandra Anderson[1‡], Jessica Howell[1‡], Richard B. Lanctot[2†§],

Marley Aikens[3¶], Joaquín Aldabe[4,5†], Liam A. Berigan[6†], Joël Bêty[7†], Erik Blomberg[6†], Juliana Bosi de Almeida[8†],

Andy J. Boyce[1†], David W. Bradley[9†], Stephen Brown[10†§], Jay Carlisle[11†], Edward Cheskey[12¶], Katherine Christie[13†],

Sylvain Christin[14¶], Rob Clay[15†§¶], Ashley Dayer[16§], Jill L. Deppe[17§], Willow English[18†], Scott A. Flemming[19†],

Olivier Gilg[20, 21†], Christine Gilroy[14†], Susan Heath[22†], Jason M. Hill[23†], J. Mark Hipfner[19†], James A. Johnson[2†],

Luanne Johnson[24†], Bart Kempenaers[25†], Paul Knaga[3¶], Eunbi Kwon[25†], Benjamin J. Lagassé[26†], Jean-François

Lamarre[27†], Christopher Latty[28†], Don-Jean Léandri-Breton[29†], Nicolas Lecomte[30†], Pam Loring[31†], Rebecca

McGuire[32, 33†], Scott Moorhead[34¶], Juan G. Navedo[35, 36, 37†], David Newstead[38†¶], Erica Nol[39†], Alina Olalla-

Kerstupp[40†], Bridget Olson[41†], Elizabeth Olson[24†], Julie Paquet[14†], Allison K. Pierce[42†], Jennie Rausch[43†], Kevin

Regan[44†], Matt Reiter[45§], Amber M. Roth[46†], Mike Russell[47†], Sarah T. Saalfeld[2†], Amy L. Scarpignato[1†], Shiloh

Schulte[48†], Nathan R. Senner[49†], Joseph A. M. Smith[50†], Paul A. Smith[51†§], Zach Spector[52¶], Kelly Srigley Werner[53§],

Michelle L. Stantial[54†], Audrey R. Taylor[55†], Mihai Valcu[25†], Walter Wehtje[56†], Brad Winn[49†], Michael B. Wunder[42 †]

[*] Corresponding Author/Principal Investigator. HarrisonAL@si.edu
[†] Data Contributor to the Collective
[‡] Collective Staff Member
[§] Member of the Collective's Advisory Group
[¶] Practitioner Contributor to the Shorebird Collective
[1] Smithsonian's National Zoo and Conservation Biology Institute, Migratory Bird Center, 3001 Connecticut Ave
NW, MRC 5503, Washington, D.C. 20008, USA

[2] U.S. Fish and Wildlife Service, Division of Migratory Bird Management, 1011 E Tudor Rd, MS 201, Anchorage,
AK 99503, USA
41
42 [3] Environment and Climate Change Canada, Canadian Wildlife Service, 45 Alderney Dr, Dartmouth, NS B2Y 2N6,
43 Canada
44
45 [4] Centro Universitario Regional del Este, Universidad de la República, Departamento de Sistemas Agrarios y
46 Paisajes Culturales, Ruta 15 y Ruta 9, Km 28.500, Departamento de Rocha, Uruguay
47
48 [5] Manomet Inc., Flyways Program, Rocha, Uruguay
49
50 [6] University of Maine, Department of Wildlife, Fisheries, and Conservation Biology, 5755 Nutting Hall, Room 210,
51 Orono, ME 04469, USA
52
53 [7] Université du Québec à Rimouski, Département de biologie et Centre d'études Nordiques, 300, allée des Ursulines,
54 C.P. 3300, succ. A, Rimouski, QC G5L 3A1, Canada
55
56 [8] Manomet Inc., Brasília, Brazil
57
58 [9] Birds Canada, 115 Front Rd, Port Rowan, ON N0E 1M0, Canada
59
60 [10] Manomet Inc., P.O. Box 545, Saxtons River, VT 05154, USA
61
62 [11] Boise State University, Intermountain Bird Observatory, 1910 University Dr, Boise, ID 83725, USA
63
64 [12] Nature Canada, 300-240 Bank St, Ottawa, ON K2P 1X4, Canada
65
66 [13] State of Alaska, Department of Fish and Game, Threatened, Endangered, and Diversity Program, 333 Raspberry
67 Rd, Anchorage, AK 99518, USA
68
69 [14] Environment and Climate Change Canada, Canadian Wildlife Service, 17 Waterfowl Ln, Sackville, NB E4L 4N1,
70 Canada
71
72 [15] Manomet Inc., WHSRN Executive Office, Asunción, Paraguay
73
74 [16] Virginia Tech, Department of Fish and Wildlife Conservation, Cheatham Hall, RM108, Blacksburg, VA 24061,
75 USA
76
77 [17] National Audubon Society, Migratory Bird Initiative, 1828 L St NW, Washington, D.C. 20036, USA
78
79 [18] Carleton University, 1125 Colonel By Dr, Ottawa, ON K1S 5B6, Canada
80
81 [19] Environment and Climate Change Canada, Wildlife Research Division, 5421 Robertson Rd, Delta, BC V4K 3N2,
82 Canada
83
84 [20] Laboratoire Chrono-environnement, UMR 6249/CNRS-Université de Franche-Comté, 16 route de Gray, 25030
85 Besançon, Bourgogne-Franche-Comté, France
86
87 [21] Groupe de Recherche en Ecologie Arctique, 16 rue de Vernot, 21440 Francheville, Bourgogne-Franche-Comté,
88 France
89
90 [22] Gulf Coast Bird Observatory, 299 Hwy 332 West, Lake Jackson, TX 77566, USA
91
92 [23] Vermont Center for Ecostudies, 20 Palmer Ct, White River Junction, VT 05001, USA
93
94 [24] BiodiversityWorks, 18 Helen Ave, Vineyard Haven, MA 02568, USA

[25] Max Planck Institute for Biological Intelligence, Department of Ornithology, Eberhard-Gwinner-Straße, 82319 Seewiesen, Bayern, Germany

[26] University of Alaska Fairbanks, Department of Biology and Wildlife, 505 S Chandalar, Fairbanks, AK 99775, USA

[27] Polar Knowledge Canada, Canadian High Arctic Research Station Campus, 1 Uvajuq Rd, P.O. Box 2150, Cambridge Bay, NU X0B 0C0, Canada

[28] U.S. Fish and Wildlife Service, Arctic National Wildlife Refuge, 101 12th Ave, Fairbanks, AK 99701, USA

[29] McGill University, Department of Natural Resource Sciences, 21111 Lakeshore Rd, Sainte-Anne-de-Bellevue, QC H9X 3V9, Canada

[30] Centre d'Études Nordiques and Canada Research Chair in Polar and Boreal Ecology, Université de Moncton, Département de Biologie, Pavillon Rémi-Rossignol, 18, Antonine-Maillet, Moncton, NB E1A 3E9, Canada

[31] U.S. Fish and Wildlife Service, Migratory Birds Program, Populations Branch, 50 Bend Rd, Charlestown, RI 02813, USA

[32] ABR, Inc. – Environmental Research & Services, P.O. Box 80410, Fairbanks, AK 99708, USA

[33] Wildlife Conservation Society, 302 Cushman Street, Suite 203, Fairbanks, AK 99701, USA

[34] Texas Water Trade, 611 S Congress Ave, Austin, TX 78704, USA

[35] Universidad Austral de Chile, Bird Ecology Lab, Instituto de Ciencias Marinas y Limnológicas, Independencia 641, Valdivia, Los Ríos, Chile

[36] Universidad Austral de Chile, Estación Experimental Quempillén, Ancud, Chiloé, Chile

[37] Millennium Institute Biodiversity of Antarctic and Subantarctic Ecosystems, Las Palmeras 3425, Ñuñoa, Santiago, Chile

[38] Coastal Bend Bays & Estuaries Program, 1305 N Shoreline Blvd #205, Corpus Christi, TX 78401, USA

[39] Trent University, Biology Department, 1600 West Bank Dr, Peterborough, ON K9L 0G2, Canada

[40] Universidad Autónoma de Nuevo León, Facultad de Ciencias Biológicas, Laboratorio de Biología de la Conservación y Desarrollo Sustentable, Ave. Universidad s/n. Ciudad Universitaria, San Nicolás de los Garza, N.L. México 66455

[41] U.S. Fish and Wildlife Service, St. Croix Wetland Management District, 1764 95th St, New Richmond, WI 54017, USA

[42] University of Colorado Denver, Department of Integrative Biology, 1201 Larimer St, Denver, CO 80204, USA

[43] Environment and Climate Change Canada, Canadian Wildlife Service, 5019 52nd St, P.O. Box 2310, Yellowknife, NT X1A 2P7, Canada

[44] Biodiversity Research Institute, 276 Canco Rd, Portland, ME 04103, USA

[45] Point Blue Conservation Science, 3820 Cypress Drive, Suite 11, Petaluma, CA 94954, USA

[46] University of Maine, School of Forest Resources and Department of Wildlife, Fisheries, and Conservation Biology, 5755 Nutting Hall, Orono, ME 04469, USA

[47] Government of Alberta, Alberta Environment and Protected Areas, 1601 Provincial Building, 10320-99 ST, Grande Prairie, AB T8V 6J4, Canada

[48] Manomet Inc., P.O. Box 1770, Manomet, MA 02345, USA

[49] University of Massachusetts Amherst, Department of Environmental Conservation, 160 Holdsworth Way, Amherst, MA 01003, USA

[50] U.S. Fish and Wildlife Service, Edwin B. Forsythe National Wildlife Refuge, 800 Great Creek Rd, Oceanville, NJ 08231, USA

[51] Environment and Climate Change Canada, National Wildlife Research Centre, 1125 Colonel By Dr, Ottawa, ON K1S 5B6, Canada

[52] Texas Parks and Wildlife Foundation, 2914 Swiss Ave, Dallas, TX 75204, USA

[53] U.S. Fish and Wildlife Service, Missouri Private Lands Office, 101 Park DeVille Drive, Suite B, Columbia, MO 65203, USA

[53] SUNY College of Environmental Science and Forestry, Department of Environmental and Forest Biology, 1 Forestry Dr, Syracuse, NY 13210, USA

[54] University of Alaska Anchorage, Department of Biological Sciences, 3211 Providence Dr, Anchorage, AK 99508, USA

[55] Ricketts Conservation Foundation, 607 Upper Hoback Road, P.O. Box 300, Little Jackson Hole, WY 82922, USA

The collective application of shorebird tracking data to conservation

Introduction

Conservation efforts are often location-specific (Stewart et al. 2013), presenting challenges for managing migratory species that traverse vast distances. Shorebirds (Charadriiformes, suborder Charadrii) exemplify these challenges as many travel thousands of kilometers biannually across continents and oceans (e.g. Gill et al. 2009). The flocking behavior of shorebirds gives the illusion of abundance, leading to shifting baselines (*sensu* Pauly 1995) and masking major declines. Shorebirds have the largest declines (net and % loss in abundance; greatest proportion of species in decline) of nearly 3 billion birds lost in North America since 1970 (compared to landbirds, waterbirds, and waterfowl, Rosenberg et al. 2019). Declines show no sign of abating (Smith et al. 2020), emphasizing the urgent need for evidence-based conservation action for shorebirds. Here we present a partnership of scientists and practitioners (defined here as those directly guiding conservation activites) responding to this need for action by linking a relatively new scientific resource—shorebird tracking data—with conservation practice.

Scientists worldwide use electronic tracking devices (hereinafter "tags", reviewed by Robinson et al. 2010) to collect animal movement data (hereinafter "tracking data"). Tracking data have become instrumental in unveiling migration routes; identifying important habitats; inferring animal behavior, and understanding population connectivity (Kays et al. 2015; Hussey et al. 2015). Aggregating tracking data across species has led to significant insights, for example documenting patterns in timing of seasonal events in relation to climate change in the Arctic (Davidson et al. 2020). Insights from tracking data have also led to significant conservation achievements. For example, tracking data collected from marine animals were used to designate marine protected areas, reduce fisheries bycatch and vessel strikes, and minimize impacts of extractive industries (Hays et al. 2019).

There has been significant growth in shorebird tracking studies over the last 20 years, unlocking the potential for tracking data to contribute to shorebird conservation. In a review of avian tracking studies (Scarpignato et al. 2023), shorebirds and songbirds constituted <1% of studies before 2006, but these groups constituted 30–75% of tracking studies in subsequent years. However, the conservation potential of these increasingly abundant data is limited by four

major analytical and conservation challenges (Table 1): only 35% of shorebird tracking data are
open access (Scarpignato et al. 2023); analyzing tracking data requires specialized knowledge;
shorebirds are not the management priority in many of their important habitats; and due to their
migratory nature, there is a need for strategic integration across organizations and habitats.

To overcome challenges limiting science translation for conservation, both linear and
interactive approaches are used (Beier et al. 2017; Toomey et al. 2017; Saunders et al. 2021).
Linear methods involve creating scientific products to be accessed by practitioners (e.g. data
portals, online tools, publications, and manuals). Despite conservation benefits (Seavy & Howell
2010; Sullivan et al. 2017), linear delivery has limitations. Tools may not suit practical purposes
due to formatting or language incompatibilities, incomplete data, or mismatched questions, and
can be time-consuming to find and synthesize (Vogel et al. 2007). In contrast, interactive
approaches range from focused workshops to long-term co-production where research and
products are collaboratively designed, planned, and implemented by practitioners and scientists
(Young et al. 2014; Roux et al. 2006; Saunders et al. 2021). These approaches allow participants
to ask clarifying questions (Gerber et al. 2020), build and maintain relationships (Lindell &
Dayer 2022), and engage in active collaborations (Young et al. 2014). Interactive approaches are
generally more successful (Cooke et al. 2020), but there are sometimes participation barriers
(Seavy & Howell 2010; Oliver et al. 2019).

We combined elements of both linear and interactive knowledge-sharing approaches to
form the Shorebird Science and Conservation Collective (hereinafter, 'the Collective'). The
Collective is a partnership of scientists and practitioners working to advance shorebird
conservation in the Americas (Figure 1). Our goal in sharing the origin, structure, and
achievements of the Collective is to provide a case study that can help guide others who are
interested in increasing the impact of existing science on conservation. The Collective acts as an
intermediary (or boundary group, Cash et al. 2006) whereby dedicated staff collate and analyze
data contributions from scientists to support the conservation needs of practitioners. We describe
how the Collective a) raised funds, b) initiated a focused proof-of-concept, c) requested data,
established processes to address privacy concerns and to report to contributors, d) developed a
co-production process to listen and respond to the needs of practitioners, and e) contributed to
conservation in its first two years.

Forming the idea for the Shorebird Science and Conservation Collective

Private foundations can greatly impact conservation funding (Zavaleta et al. 2008;

Bakker et al. 2010). In our case, the Knobloch Family Foundation catalyzed the Collective by

inviting the Principal Investigator to propose a "transformative idea for data-driven shorebird

conservation in North America." The idea to collate shorebird tracking studies for conservation

was inspired by the successful example of BirdLife International's Seabird Tracking Database in

informing conservation (BirdLife International 2004)—but adapted for unique considerations of

how and where shorebird conservation is implemented (Table 1, Table 3). Given existing

limitations (Table 1), a linear approach facilitating data discovery or access, but requiring

practitioners to interpret data and literature themselves, seemed unlikely on its own to maximize

knowledge transfer. Instead, we proposed a Collective that would extend beyond a repository to

include staff for data analysis and co-development of conservation products to support direct

requests from practitioners.

A funding proposal was developed with feedback from shorebird researchers and

conservation practitioners through open-invitation webinar discussions. Many participants were

already sharing, co-developing or applying research to conservation. As conceived, the

Collective would add value by expanding conservation opportunities, rather than compete with

or replace existing partnerships. Webinars were instrumental for assessing support, refining

ideas, and identifying existing gaps in translating science to shorebird conservation.

A proof-of-concept for the Collective

We proposed a three-year proof-of-concept focused on the midcontinent of North

America to evaluate the Collective's structure and gauge demand for shorebird tracking data.

The Midcontinent has been elevated as a priority landscape by multiple conservation entities

(www.grasslandsroadmap.org, www.midamericasshorebirds.org) and provides habitat for 42 of

the 50 shorebird species that regularly breed in North America (Donaldson et al. 2000; Brown et

al. 2001). For example, the Texas Gulf Coast provides migratory and wintering habitat for more

than 1 million shorebirds annually (Withers 2002) while the Prairie Pothole Region's wetlands

and grasslands (Doherty et al. 2018) support millions of migrants each spring and fall (Skagen

2006; Steen et al. 2018). These landscapes have experienced major wetland loss (up to 90% at

the U.S. state level, (Dahl 1990; Dahl 2014)) and have high spatial and temporal climatic

variability from droughts to flooding, making cohesive shorebird management challenging

across the region and among years (Fellows et al. 2001; Russell et al. 2016; Skagen & Thompson

2013). Many areas are privately owned (e.g. nearly 90% of the Great Plains is non-public (NRCS

2021), requiring coordination with landowner-focused agencies, which deliver voluntary

incentive programs and/or provide technical assistance to manage habitat on working lands (e.g.,

NRCS, USFWS Partners for Fish and Wildlife Program), as well as state and local governments,

land trusts, and other conservation entities.

280   The proposal was funded in late 2020, allowing implementation to begin in early 2021.

282   **The Collective's structure and systems**

283   The Collective is comprised of 4 groups of people (Figure 1): a core operational team

(P.I., staff, and Chair of the Advisory Group), an Advisory Group (Appendix I), data

contributors, and practitioner partners. All groups are represented as co-authors. The P.I. and

staff are based at the Smithsonian Migratory Bird Center. Staff include a quantitative ecologist

who aggregates, processes, and analyzes data; and conservation specialists who work with

practitioners to inform them of the tracking data resource, assess their information needs, and

help prepare conservation products. The 16-person Advisory Group advises on maximizing the

conservation benefit of the Collective, promotes its value to potential data contributors, and

connects it with organizations engaged in similar efforts or those that would benefit from

tracking data. Data contributors typically are researchers from within public, private, or non-

profit organizations. Practitioners directly guide conservation actions from within management

agencies, trusts, and non-profit groups (Table 3). Some data contributors are also practitioners.

295   With guidance from all involved, the core team developed clear operational processes

296   including a Data Sharing Agreement (Appendix II), a conservation use request form (Appendix

297   III), and a conservation reporting system (Appendices IV-VI) to ensure data privacy agreements

298   are upheld and to increase accountability to data contributors and practitioners.

300   *Crafting a Data Sharing Agreement*

301   Data contributors are the Collective's foundation; their data drive evidence-based

recommendations and products. Feedback indicated that careful crafting of a Data Sharing

Agreement would ensure maximum trust and participation by the scientific community. The

Collective aims for shorebird tracking data to be findable, accessible, interoperable, and reusable

(FAIR Guiding Principles, Wilkinson et al. 2016) while respecting contributor rights. Many

factors influence data contribution (e.g., Roche et al. 2020), with some policies mandating public

sharing. Conversely, certain datasets, like recently-collected data or those restricted by

endangered species status, may require embargo or organizational approval before public release.

We thus drafted a flexible agreement (Appendix II, https://nationalzoo.si.edu/migratory-

birds/shorebird-science-and-conservation-collective-data-sharing-agreement-form). All

contributors allow the Collective's core team to explore data to assess applicability to a

conservation request. Contributors further customize usage permissions (either providing full

permission or requiring a permission request before each use) within use categories:

conservation, education, funder reports, scientific publications, and/or research requests.

Compiling data

We invited data contributions through email, a webinar, social networks, and word-of-

mouth. Data were received either through repositories (e.g., Movebank) or were sent directly to

the Collective's staff. We prioritized processing data with higher spatial accuracy (e.g., Argos

and GPS; Table 2) that we anticipated would be most relevant to conservation requests from

practitioners. We developed a set of R scripts that first apply pre-filters (e.g. to determine when a

tag has become stationary, indicating it has either been shed by the bird or the bird has died).

Then a continuous-time state-space model (Jonsen et al. 2023) estimates the most probable

movement path accounting for the 1) spatial error of the tracking technology and 2) irregular

sampling across individuals and species. The Collective is currently integrating data collected by

the Motus Wildlife Tracking System (Taylor et al. 2017) and light-level geolocators (Table 2)

that were contributed either pre-processed using various methods, or in raw, unprocessed form,

requiring additional processing or standardization to ensure appropriate interpretation (Lisovski

et al. 2018). Data are linked to their associated Data Sharing Agreements to ensure that

permissions and attributions are closely followed by staff when considering use for conservation

applications.

Reporting conservation uses to contributors

Contributors are notified when their data are relevant to a conservation request (examples

in Appendices III-V). The notification describes the project and provides a map of the data

involved. For any data use that the contributor did not pre-approve in the Data Sharing

Agreement, the Collective requests permission before proceeding. While this process adds to the

administrative work of the Collective's staff, it incentivizes and rewards data contribution,

increases accountability, safeguards privacy settings, and promotes dialogue to prevent

misinterpretations of contributed data.

Engaging conservation partners

Information requests from practitioners are submitted via a Conservation Use Request

Form (Appendix III and https://nationalzoo.si.edu/migratory-birds/request-for-use-shorebird-

tracking-data), or through emails, informational meetings, conference presentations, and

referrals. Conservation specialists currently prioritize outreach to practitioners within the

Midcontinent given the geographic focus of our proof-of-concept. However, requests have been

received from across North America. Knowledge-sharing typically begins with a 1-hr virtual

meeting between practitioners and members of the Collective's core team. The team describes

the Collective's approach for data-driven conservation, and practitioners describe their work,

priorities, and shorebird-related questions. If there is opportunity for the Collective to inform a

project, further conversations help refine questions, information needs, and preferred formats of

data products. Data contributors are then engaged to notify them that their data are relevant to a

conservation request and/or to request further permissions.

Throughout the project, the core team communicates with practitioners and data

contributors as needed. The team presents findings to practitioners, allowing questions and

feedback regarding the delivery format and recommendations. At the project's conclusion, a

Conservation Contribution Report summarizes recommendations, acknowledges data

contributors, and provides information about shorebird ecology and best management practices.

A public version (with any sensitive data removed) is also shared online

(https://nationalzoo.si.edu/migratory-birds/shorebird-collective). The Collective classifies

conservation contributions according to the IUCN Conservation Actions Classification Scheme

(Salafsky et al. 2008; IUCN 2012) to align with standard conservation evidence reporting

(Sutherland et al. 2019).

The wealth of shorebird tracking data available for conservation in the Americas

Data contributions include 6.7 million locations from 3,345 individual shorebirds of 36 species (Figure 2). Since January 2021, when data were first requested, contributors have signed 58 agreements for data originating from 74 organizations (e.g., academic institutions, nonprofits, and government agencies) and 136 data co-owners from 14 countries.

Data span the Americas with areas in the Midcontinent demonstrating clear importance to multiple species, including the Prairie Pothole Region and Gulf of Mexico coast (Figure 2). Shorebirds were tracked via light-level geolocator, Argos, GPS, and Motus technologies, as well as with visual observations of uniquely marked birds (i.e. with flags or bands) (Table 2, Figure 2). Data come from the smallest (Least Sandpiper, *Calidris minutilla,* 19-30g, tracked with 0.67 g Motus tags; (Anderson et al. 2019)) to the largest (Long-billed Curlew, *Numenius americanus,* 490-950 g, tracked with 9.5-22 g satellite tags; (Page et al. 2014)) shorebirds in North America (Species list, Appendix VII). Detections came from varied habitats including fresh and saline wetlands, uplands, grasslands, agricultural fields, coastal beaches and mudflats, tundra, forest, and over open ocean.

Contributing to conservation

Since our first conservation request in November, 2021 to October, 2023, the Collective has supported 10 conservation projects (with another 4 ongoing). We have sent 93 notifications to 23 contributors informing them that their data collected from 17 species were relevant to conservation requests. Fifty-six notifications were for datasets pre-approved to be used for conservation, and 37 were requests to contributors requiring further permission (see Appendices II-IV for examples). Of the latter set, 92% were approved. Below, we provide examples of ways the Collective applied tracking data to support: 1) land conservation, 2) land and species management, and 3) education and outreach efforts.

1. Private Land Conservation

Both public and private lands support shorebird conservation in North America, but within the Midcontinent where there is high private ownership, private lands are an especially crucial component of environmental stewardship and land protection (Merenlender et al. 2004).

Trusts, foundations, and government agencies can use insights from tracking data to help guide

conservation investments and secure funds for voluntary land purchase or conservation. These

datasets are especially useful when they provide the only evidence of shorebirds using areas (for

example, when systematic ground surveys or eBird observations are unavailable).

Funding and siting of conservation easements

The Texas Parks and Wildlife Foundation's Buffer Lands Incentive Program (BLIP)

provides financial assistance for conservation easements (Merenlender et al. 2004) on privately

owned "buffer lands" bordering Texas Parks and Wildlife Department (TPWD) public lands.

The BLIP requested evidence for shorebird use of buffer lands bordering 12 priority TPWD

lands as one metric in their rubric to prioritize applications for conservation easement funding;

half of BLIP funds are allocated for shorebird and grassland bird projects. Because many parcels

under consideration are inaccessible to the public, direct evidence of use by shorebirds comes

almost exclusively from tracking data. Fifty-two individuals of nine species in the Collective's

dataset stopped or overwintered on buffer lands. For 83% of individuals, tags were deployed out-

of-state. The Collective summarized when birds stopped or wintered in buffer lands, the habitats

used, and global migration links ranging from Russia to Argentina to assist with landowner

engagement. Finally, we shared a list of all TPWD lands visited by tracked shorebirds in the

Collective's dataset, and state-wide density maps of individuals and species tracked.

Purchasing land for conservation

The Conservation Fund (TCF) and The Nature Conservancy (TNC), two non-

governmental conservation organizations, separately sought information to support funding

efforts to acquire lands for permanent protection in the Texas coast (TCF) and the Texas

panhandle (TNC). Although no shorebirds within the Collective's dataset visited the parcels

under consideration, 23 individuals of four species were tracked nearby (within 3–18 km). We

used habitat layers and standardized eBird relative abundance (Fink et al. 2022) to provide

additional context about the potential value of the parcels to shorebirds. Some habitats in the

parcels seemed promising and we recommended on-the-ground surveys to confirm shorebird use.

Both organizations shared these data with funders to demonstrate the potential value of these

landscapes for shorebirds.

2. Land and Species Management

Many shorebird conservation activities manage for habitat or species of concern (Iglecia
& Winn 2021). Land management includes activities like deploying water to create habitat,
alternating wetting and drying of rice fields timed with shorebird usage, and prescribed grazing
and burning of grasslands or rangelands. Species management includes activities like species
recovery planning and status assessments, harvest management, and beach closures. Tracking
data can reveal important sites, connectivity among sites, individual arrival and departure timing
at particular sites, and species population structure to inform management actions.

Informing the timing and location of freshwater deployments

The Galveston Bay Foundation and Texas Water Trade, two Texas-based non-
governmental organizations, recently purchased over 5,000 acre-feet of water rights. They
wanted to know when and where to deploy water along the east Texas coast to benefit
shorebirds. "Focused flows" offer many benefits including habitat creation and restoration,
salinity management, and continued crop yields during drought (Culp et al. 2014; Garmany
2020). Tracking data from six shorebird species provided information on local habitat use and
timing through the focal area (Figure 3a, 3b). Standardized relative abundance data from eBird
(Fink et al. 2022) added insights for species with limited tracking data and provided peaks of
occurrence across multiple species (Figure 3b). We provided partners with recommendations for
when to deploy water (preferred and alternate times, Figure 3c) and shorebird habitat
management best practices.

Offshore wind energy development

Wind energy is important for reducing carbon emissions, but may have cumulative
negative effects on birds through collisions or displacement (Goodale & Milman 2016; Fox &
Petersen 2019). Tracking data can provide valuable insight about movement routes and timing of
passage in proposed and active offshore wind areas to better understand shorebird vulnerability,
inform siting decisions, and/or encourage "smart curtailment" strategies during high-risk
migratory periods (Loring et al. 2021; Schwemmer et al. 2023). The Canadian Wildlife Service,
Canada's federal wildlife agency, requested information for two rapid (18-month) regional

assessments offshore Nova Scotia and Newfoundland and Labrador to understand the temporal

and spatial distribution of shorebirds moving through the focal areas. We provided maps of 41

individuals of eight species that migrated through the offshore areas of interest and highlighted

areas of high activity (i.e., where track lines congregated for multiple individuals/species). The

U.S. Fish and Wildlife Service, the United States' federal wildlife agency, has requested similar

data to incorporate into a collision risk model and to inform official comments on environmental

impact statements. The Collective is actively supporting this request.

Improving length of stay estimates and bird use days

Population abundance and trends are critical metrics that guide conservation actions and

targets. However, accurate estimates are challenging to obtain for many shorebird species that

aggregate in large groups with regular inflow and outflow of individuals (Vallecillo et al. 2021;

Farmer & Durbian 2006; Gillings et al. 2009). Tracking data provide reliable estimates of length

of stay (time between arrival and departure) for individuals at migratory stopover sites that can

improve survey-based estimates of passage populations (Farmer & Durbian 2006). This

information in turn, facilitates population monitoring, guides habitat management objectives and

helps assess management effectiveness, and informs other planning tools such as abundance

targets (Environment and Climate Change Canada 2016; U.S. Fish and Wildlife Service 2023)

and harvest and caloric models (Loesch et al. 2000; Vermillion & Lancaster 2022). The

Collective is applying movement models to efficiently classify stopover locations, estimate

length of stay, and determine bird use days to inform requests from managers with Migratory

Bird Joint Ventures and U.S. State and Canadian Federal wildlife management agencies.

3. Education and Outreach

Despite increasing public awareness of declining bird populations, many audiences may

not be aware of the broad array of habitats shorebirds need during their migrations. The term

"shorebird" may limit conservation efforts in the Midcontinent, as many people do not connect

them to grasslands or interior wetlands. Shorebirds are sometimes wrongly assumed to have

similar habitat needs as other waterbirds like gulls, egrets, rails, or waterfowl. Environmental

education can lead to indirect and direct conservation outcomes (Ardoin et al. 2020) and as an

outreach tool, tracking data offer strong visual and hands-on learning opportunities to gain

support for shorebird conservation efforts.

School curriculum

Educators at Grays Harbor National Wildlife Refuge in Hoquiam, Washington, USA

requested geographic coordinates and track lines to support their 3rd and 4th grade Shorebird

Education Programs. The Collective provided data from 17 individuals of five species detected

in or near Grays Harbor Estuary. Educators incorporated these data into four lessons, allowing

students to map and explore shorebird movements within the Grays Harbor estuary and across

the Pacific Flyway. The activity allowed the students to enhance skills in geography and ecology

while learning how small actions in their communities can affect a shorebird's entire migration.

Refuge educators visited multiple classrooms every month, reaching a total of 870 students

across 14 schools with this lesson over the 2022–23 school year.

Community outreach

Nature Canada, a non-governmental conservation organization, requested tracking data

for a comic following the movements of Bico, a fictional Hudsonian Godwit, as she travels from

Canada to Chile. The comic's goal is to raise awareness and generate excitement about

shorebirds that visit James Bay, Canada, a globally important stopover area. The Collective

provided two Hudsonian Godwit tracks and summary statistics (e.g., stopover sites and tenures,

average flight speeds) for an infographic at the end of the comic. Nature Canada posted the

comic online (link) and distributed paper copies in six First Nations communities around James

Bay. The Collective provided similar information to the Minnesota Board of Water and Soil

Resources, a state soil and water management agency, to communicate the role that wetland

restoration and water management projects have in protecting shorebird populations in

Minnesota. The Board shared this with their staff and conservation partners.

Connecting communities with shorebird movements

The Western Hemisphere Shorebird Reserve Network, a science-based, partnership-

driven, conservation initiative for protecting shorebirds through a network of key sites, wanted to

show how shorebird movements connect "sister sites" within their reserve network. The

Collective created an online storytelling series using ArcGIS StoryMaps to follow three tracked

individuals as they migrated across the Americas (http://tiny.cc/ssccstorymap). Each StoryMap

features a migratory bird flyway (Pacific, Midcontinent, Atlantic) and cultural theme (food,

music, crafts), highlighting the people and places each shorebird might encounter along their

migrations.

Successes and future

The Shorebird Science and Conservation Collective is advancing shorebird conservation

through collaborative data sharing and analysis in support of needs identified by practitioners.

Such on-the-ground conservation is especially crucial given long-term and, now, accelerating

shorebird population declines (Rosenberg et al. 2019; Smith et al. 2023). During the first two

years we worked to address limitations to data access and analysis, and to conservation

implementation (Table 1) in order to advance data-driven shorebird conservation.

We created a central resource of over 3,300 individual movement paths of shorebirds and

provided staff to facilitate data access, and interpret and analyze shorebird science. New data

become available as research projects develop and new contributors join. Staff share scientific

expertise with practitioners, including specialized knowledge of shorebird ecology and statistical

approaches to analyzing tracking data. Through these free scientific services offered to

practitioners, and through the Collective's extensive outreach throughout the Midcontinent, we

have made it easier for practitioners to learn how shorebirds use their jurisdictions and to

consider shorebirds when implementing their job duties. Our structure facilitates interaction,

leading to co-production of analyses to inform land conservation, species management, and

education and outreach.

The Collective is also providing a more complete understanding of shorebird ecology

across the Americas, but some data limitations remain. For example, data contributed to the

Collective are limited in some places or for some species and some datasets have high spatial

uncertainty or low sampling frequency. Small sample sizes can limit generalizability and

statistical inference (Hebblewhite & Haydon 2010). These factors are inherent to collated

tracking datasets (Block et al. 2011), requiring careful evaluation of the suitability of available

data to specific conservation questions, and communication of uncertainty (something the

Collective emphasizes with partners). However, even small datasets can provide meaningful

insights (Sequeira et al. 2019), and tracking data are often the only data available for shorebirds

on private lands. Tracking datasets are the foundation of the Collective's ability to implement

bird conservation, but monitoring programs (e.g., International Shorebird Survey,

https://www.manomet.org/project/international-shorebird-survey/; Migratory Shorebird Project,

https://migratoryshorebirdproject.org/) and eBird participatory science data complement tracking

data to inform conservation, especially where tracking data are limited (often for small-bodied

species). Integrating across data types and facilitating communication with data registry efforts

(Rutz 2022; Nightingale 2023) to create efficient and effective pipelines from data to action is a

priority.

Due to privacy settings for some datasets, the Collective has not made a public mapping

tool. Complementary tools are available that allow practitioners to explore tracking maps (e.g.,

Bird Migration Explorer, Smith et al. 2022). Data repositories (e.g. Movebank, Davidson et al.

2022; Motus, Taylor et al. 2017) allow those with time and motivation to find and download

open access data, or to request permission to download private data, but see limitations (Figure 1,

Scarpignato et al. 2023). However, practitioners do not often request raw data from us, or

typically have staff capacity, time, motivation, or background to find and analyze data, even if all

data were open access. The Collective adds value by analyzing and interpreting a complex

hemispheric-scale dataset of shorebird movements in a way that is understandable, co-developed,

and delivered in preferred formats. This lowers the activation energy for practitioners to consider

and use shorebird science.

Most projects supported by the Collective are in response to direct requests, referrals, or

targeted outreach. There is a need for future contributions to inform regional, national, and

international priorities through strategic analyses designed to have the biggest effect on reversing

shorebird population declines. We recognize potential to inform policy and funding decisions

with shorebird science. For example, in the United States, federal agencies are administering

landmark investments in climate and conservation programs through the 2022 Inflation

Reduction Act (H.R.5376). The Collective's consolidated dataset provides spatial and temporal

evidence of shorebird habitat use that is also critical for informing international conservation

conventions and agreements (Navedo & Piersma 2023).

Our funding prioritized the Midcontinent as a proof-of-concept, but since the Collective's

inception, partners have emphasized the importance of taking a full annual cycle approach to

shorebird conservation. While our highest engagement came from practitioners in Texas, USA,

the Collective also supported projects in the Pacific and Atlantic Flyways of North America, and

data were contributed from scientists in 14 countries. We recognize an urgent need to secure

additional funding to continue existing support of conservation, and to expand the Collective's

impact in the Caribbean region and Central and South America. Priorities include hiring

specialists focused on supporting conservation initiatives in these regions, and who are fluent in

Spanish, Portuguese, and French, and the inclusion of tracking data from species endemic to

Latin America and the Caribbean. The Collective's synthesized dataset can also help prioritize

research to fill knowledge gaps. For example, tracking data are already being used to help inform

shorebird surveys of the Amazon Basin, where eBird and survey data are extremely limited.

Innovations in tracking technology continue to enhance our understanding of shorebird

migrations and habitat use—patterns that are changing with persistent habitat loss and climate

change (Sutherland et al. 2012; Galbraith et al. 2014). Ongoing data collection and sharing are

thus vital for liaisons like the Collective to rapidly integrate and interpret research for

conservation in a changing world.

Acknowledgments

Support for the Shorebird Science and Conservation Collective is provided by the

Knobloch Family Foundation. Additional support is provided from ConocoPhillips Charitable

Investments in support of the Migratory Connectivity Project (A.-L.H.) and from the U.S. Fish

and Wildlife Service (R.B.L.). Thanks to the Advisory Group (Appendix I) and for data or

conservation contributions from B. Ballard, B. Corey, G. Castresana, K. De Santiago, P.

Erdmann, J. Pinnix, D. Ruthrauff, R. Summers, J. Shackelford, D. Shaw, M. Singer, T.L.

Tibbitts, M. Weegman, L. Wright. Thanks to all who joined listening sessions and provided early

feedback. Finally, thanks to funders, collaborators, volunteers, and staff who collected or

supported the collection of shorebird tracking data. Any use of trade, product or firm names is

for descriptive purposes only and does not imply endorsement by the U.S. Government.

Bibliography

Anderson, AM Duijns, S Smith, PA Friis, C, Nol, E (2019) Migration distance and body

condition influence shorebird migration strategies and stopover decisions during

southbound migration. *Frontiers in Ecology and Evolution* 7:Article 251

Ardoin, NM Bowers, AW, Gaillard, E (2020) Environmental education outcomes for

conservation: A systematic review. *Biological Conservation* 241:108224

Bakker, VJ Baum, JK Brodie, JF Salomon, AK and others (2010) The changing landscape of

conservation science funding in the United States. *Conservation Letters* 3(6):435–444

Beier, P Hansen, LJ Helbrecht, L, Behar, D (2017) A how-to guide for coproduction of

actionable science. *Conservation Letters* 10(3):288–296

BirdLife International (2004) Tracking ocean wanderers: The global distribution of albatrosses

and petrels, BirdLife International, Cambridge, United Kingdom

Blickley, JL Deiner, K Garbach, K Lacher, I and others (2013) Graduate student's guide to

necessary skills for nonacademic conservation careers. *Conserv Biol* 27(1):24–34

Block, BA Jonsen, ID Jorgensen, SJ Winship, AJ and others (2011) Tracking apex marine

predator movements in a dynamic ocean. *Nature* 475(7354):86–90

Boere, GC, Stroud, DA (2006) The flyway concept: what it is and what it isn't. In: Boere, GC

Galbraith, CA, Stroud, DA (eds) Waterbirds around the world. The Stationery Office,

Edinburgh, pp 40–47

Brown, S Hickey, C Harrington, B, Gill, R (2001) United States Shorebird Conservation Plan,

2nd. ed., Manomet Center for Conservation Sciences, Manomet, Massachusetts

Cash, DW Borck, JC, Patt, AG (2006) Countering the loading-dock approach to linking science

and decision making. *Science, Technology, & Human Values* 31(4):465–494

Ciuzio, E Hohman, WL Martin, B Smith, MD Stephens, S Strong, AM, Vercauteren, T (2013)

Opportunities and challenges to implementing bird conservation on private lands. *Wildlife*

Society Bulletin 37(2):267–277

CLS America (2016) Argos User's Manual: Worldwide tracking and environmental monitoring

by satellite.

Culp, PW Glennon, RJ, Libecap, G (2014) Shopping for water: How the market can mitigate

water shortages in the American West. 36

Dahl, TE (1990) Wetlands losses in the United States 1780's to 1980's. Report to Congress. 13

Dahl, TE (2014) Status and trends of prairie wetlands in the United States 1997 to 2009, US Fish
and Wildlife Service Washington, DC,

Davidson, SC Bohrer, G Gurarie, E LaPoint, S and others (2020) Ecological insights from three
decades of animal movement tracking across a changing Arctic. *Science* 370(6517):712–
715

Doherty, KE Howerter, DW Devries, JH, Walker, J (2018) Prairie pothole region of North
America. The Wetland Book. Springer Netherlands, Dordrecht, pp 679–688

Donaldson, GM Hyslop, C Morrison, RIG Dickson, HL, Davidson, I (2000) Canadian Shorebird
Conservation Plan. 27

Environment and Climate Change Canada (2016) Recovery strategy and management plan for
the Red Knot (*Calidris canutus*) in Canada [Proposed]. *Species at Risk Act* Recovery
Strategy Series. ix + 54

Farmer, A, Durbian, F (2006) Estimating shorebird numbers at migration stopover sites. *The
Condor* 108(4):792–807

Fellows, S Stone, K Jones, S Damude, N, Brown, S (2001) Central Plains/Playa Lakes regional
shorebird conservation plan. *Version* 1:26

Fernández, RJ (2016) How to be a more effective environmental scientist in management and
policy contexts. *Environmental Science & Policy* 64:171–176

Fink, D Auer, T Johnston, A Strimas-Mackey, M and others (2022) eBird Status and Trends.
Data Version: 2021; Released: 2022.

Fox, AD, Petersen, IK (2019) Offshore wind farms and their effects on birds. *Dansk Orn Foren
Tidsskr* 113:86–101

Galbraith, H DesRochers, DW Brown, S, Reed, JM (2014) Predicting vulnerabilities of North
American shorebirds to climate change. *PLoS One* 9(9):e108899

Garmany, K (2020) Market strategies for addressing water scarcity: An introduction to
environmental water transactions in texas. *A thesis in partial fulfillment of the requirements
for a degree of master of applied geography.* Texas State University

Gerber, LR Barton, CJ Cheng, SH, Anderson, D (2020) Producing actionable science in
conservation: Best practices for organizations and individuals. *Conservation Science and
Practice* 2(12):2:e295

Gill, RE Tibbitts, TL Douglas, DC Handel, CM and others (2009) Extreme endurance flights by
landbirds crossing the Pacific Ocean: ecological corridor rather than barrier. *Proceedings of*
the Royal Society B: Biological Sciences 276(1656):447–457

Gillings, S Atkinson, PW Baker, AJ Bennett, KA and others (2009) Staging behavior in Red
Knot (*Calidris canutus*) in Delaware Bay: implications for monitoring mass and population
size. *The Auk* 126(1):54–63

Goodale, MW, Milman, A (2016) Cumulative adverse effects of offshore wind energy
development on wildlife. *Journal of Environmental Planning and Management* 59(1):1–21

Gupte, PR Beardsworth, CE Spiegel, O Lourie, E Toledo, S Nathan, R, Bijleveld, AI (2022) A
guide to pre-processing high-throughput animal tracking data. *J Anim Ecol* 91(2):287–307

Hays, GC Bailey, H Bograd, SJ Bowen, WD and others (2019) Translating marine animal
tracking data into conservation policy and management. *Trends in Ecology & Evolution*
34(5):459–472

Hebblewhite, M, Haydon, DT (2010) Distinguishing technology from biology: a critical review
of the use of GPS telemetry data in ecology. *Philos Trans R Soc Lond B Biol Sci*
365(1550):2303–2312

Hussey, NE Kessel, ST Aarestrup, K Cooke, SJ and others (2015) Aquatic animal telemetry: A
panoramic window into the underwater world. *Science* 348(6240):1255642

Iglecia, M, Winn, B (2021) A shorebird management manual. *Massachusetts, USA: Manomet*

IUCN (2012) Conservation Actions Classification Scheme (Version 2.0).

Jonsen, ID Grecian, WJ Phillips, L Carroll, G and others (2023) aniMotum, an R package for
animal movement data: Rapid quality control, behavioural estimation and simulation.
Methods in Ecology and Evolution 14(3):806–816

Jonsen, ID Patterson, TA Costa, DP Doherty, PD and others (2020) A continuous-time state-
space model for rapid quality control of argos locations from animal-borne tags. *Movement*
Ecology 8:31

Joo, R Boone, ME Clay, TA Patrick, SC Clusella-Trullas, S, Basille, M (2020) Navigating
through the r packages for movement. *Journal of Animal Ecology* 89(1):248–267

Kays, R Crofoot, MC Jetz, W, Wikelski, M (2015) Terrestrial animal tracking as an eye on life
and planet. *Science* 348(6240):aaa2478

Kays, R Davidson, SC Berger, M Bohrer, G and others (2022) The Movebank system for
studying global animal movement and demography. *Methods in Ecology and Evolution*
13(2):419–431

Lindell, CA, Dayer, AA (2022) Six principles for working effectively with landowners to
advance bird conservation. *Ornithological Applications* 124(4)

Lisovski, S Bauer, S Briedis, M Davidson, SC and others (2020) Light-level geolocator analyses:
A user's guide. *Journal of Animal Ecology* 89:221–236

Lisovski, S Hewson, CM Klaassen, RHG Korner-Nievergelt, F Kristensen, MW, Hahn, S (2012)
Geolocation by light: accuracy and precision affected by environmental factors. *Methods in*
Ecology and Evolution 3:603–612

Lisovski, S Schmaljohann, H Bridge, ES Bauer, S and others (2018) Inherent limits of light-level
geolocation may lead to over-interpretation. *Current Biology* 28(3):R99–R100

Loesch, CR Twedt, DJ Tripp, K Hunter, WC, Woodrey, MS (2000) Development of
management objectives for waterfowl and shorebirds in the Mississippi Alluvial Valley.
Proceedings of the 3rd partners in flight workshop. USDA Forest Service Proceedings
RMRS-P-16, Cape May, NJ:8–11

Loring, PH Lenske, AK McLaren, JD Aikens, M and others (2021) Tracking movements of
migratory shorebirds in the US Atlantic Outer Continental Shelf Region. *Sterling (VA): US*
Department of the Interior, Bureau of Ocean Energy Management OCS Study BOEM 8:104

Merenlender, AM Huntsinger, L Guthey, G, Fairfax, SK (2004) Land trusts and conservation
easements: Who is conserving what for whom? *Conservation Biology* 18(1):65–76

Navedo, JG, Piersma, T (2023) Do 50-year-old Ramsar criteria still do the best possible job? A
plea for broadened scientific underpinning of the global protection of wetlands and
migratory waterbirds. *Conservation Letters* 16(2)

Nightingale, J (2023) Conserving wader tracking data for the future: Introducing the Global
Wader project. *Wader Study* 130(3)

Noonan, MJ Fleming, CH Akre, TS Drescher-Lehman, J and others (2019) Scale-insensitive
estimation of speed and distance traveled from animal tracking data. *Mov Ecol* 7:35

NRCS (2021) A framework for conservation action in the Great Plains Grasslands Biome.

O'Toole, M Queiroz, N Humphries, NE Sims, DW, Sequeira, AMM (2021) Quantifying effects
of tracking data bias on species distribution models. *Methods in Ecology and Evolution*
12(1):170–181

Oliver, K Kothari, A, Mays, N (2019) The dark side of coproduction: do the costs outweigh the
benefits for health research. *Health Res Policy Syst* 17(1):33

Page, GW Warnock, N Tibbitts, TL Jorgensen, D Hartman, CA, Stenzel, LE (2014) Annual
migratory patterns of Long-billed Curlews in the American West. *The Condor* 116(1):50–
61

Rakhimberdiev, E Senner, NR Verhoeven, MA Winkler, DW Bouten, W, Piersma, T (2016)
Comparing inferences of solar geolocation data against high-precision GPS data: annual
movements of a double-tagged black-tailed godwit. *Journal of Avian Biology* 47(4):589–
596

Reichman, OJ Jones, MB, Schildhauer, MP (2011) Challenges and opportunities of open data in
ecology. *Science* 331(6018):703–705

Robinson, WD Bowlin, MS Bisson, I Shamoun-Baranes, J and others (2010) Integrating
concepts and technologies to advance the study of bird migration. *Frontiers in Ecology and
the Environment* 8(7):354–361

Roche, DG Granados, M Austin, CC Wilson, S and others (2020) Open government data and
environmental science: a federal Canadian perspective. *FACETS* 5(1):942–962

Rosenberg, KV Dokter, AM Blancher, PJ Sauer, JR and others (2019) Decline of the North
American avifauna. *Science* 366(6461):120–124

Roux, DJ Rogers, KH Biggs, HC Ashton, PJ, Sergeant, A (2006) Bridging the science–
management divide: moving from unidirectional knowledge transfer to knowledge
interfacing and sharing. *Ecology and society* 11(1):4

Russell, RP Koch, KE, Lewis, SJ (2016) Upper Mississippi Valley/Great Lakes Regional
Shorebird Conservation Plan. Version 2.0. 64

Rutz, C (2022) Register animal-tracking tags to boost conservation. *Nature* 609(7926):221–221

Salafsky, N Salzer, D Stattersfield, AJ Hilton-Taylor, C and others (2008) A standard lexicon for
biodiversity conservation: unified classifications of threats and actions. *Conserv Biol*
22(4):897–911

Saunders, SP Wu, JX Gow, EA Adams, E and others (2021) Bridging the research-

implementation gap in avian conservation with translational ecology. *Ornithological*

Applications 123(3):1–13

Scarpignato, AL Huysman, AE Jimenez, MF Witko, CJ and others (2023) Shortfalls in tracking

data available to inform North American migratory bird conservation. *Biological*

Conservation 286:110224

Schwemmer, P Pederson, R Haecker, K Bocher, P and others (2023) Assessing potential

conflicts between offshore wind farms and migration patterns of a threatened shorebird

species. *Animal Conservation* 26(3):303–316

Seavy, NE, Howell, CA (2010) How can we improve information delivery to support

conservation and restoration decisions. *Biodiversity and Conservation* 19(5):1261–1267

Sequeira, AMM Heupel, MR Lea, M-A Eguíluz, VM and others (2019) The importance of

sample size in marine megafauna tagging studies. *Ecological Applications* 29(6):e01947

Skagen, SK (2006) Migration stopovers and the conservation of arctic-breeding calidridine

sandpipers. *The Auk* 123(2):313–322

Skagen, SK, Thompson, G (2013) Northern Plains/Prairie Potholes regional shorebird

conservation plan. *US Shorebird Conservation Partnership, Lakewood, Colorado, USA*

Smith, MA Mahoney, J Knight, EJ Taylor, L and others (2022) Bird Migration Explorer.
National Audubon Society, New York, NY. birdmigrationexplorer.org | exploraves.org
778
Smith, PA McKinnon, L Meltofte, H Lanctot, RB and others (2020) Status and trends of tundra

birds across the circumpolar Arctic. *Ambio* 49(3):732–748

Smith, PA Smith, AC Andres, B Francis, CM and others (2023) Accelerating declines of North

America's shorebirds signal the need for urgent conservation action. *Ornithological*

Applications 125(2)

Steen, V Skagen, SK, Noon, BR (2018) Preparing for an uncertain future: migrating shorebird

response to past climatic fluctuations in the Prairie Potholes. *Ecosphere* 9(2):e02095

Stewart, WP Williams, DR, Kruger, LE (2013) Place-Based Conservation. Perspectives from the

Social Sciences, Springer Netherlands, Dordrecht

Sullivan, BL Phillips, T Dayer, AA Wood, CL and others (2017) Using open access

observational data for conservation action: A case study for birds. *Biological Conservation*

208:5–14

Sutherland, WJ Alves, JA Amano, T Chang, CH and others (2012) A horizon scanning
assessment of current and potential future threats to migratory shorebirds. *Ibis* 154(4):663–
679

Sutherland, WJ Taylor, NG MacFarlane, D Amano, T and others (2019) Building a tool to
overcome barriers in research-implementation spaces: The Conservation Evidence
database. *Biological Conservation* 238:108199

Taylor, PD Crewe, TL Mackenzie, SA Lepage, D and others (2017) The Motus Wildlife
Tracking System: a collaborative research network to enhance the understanding of wildlife
movement. *Avian Conservation and Ecology* 12(1):Article 8

Texas Parks and Wildlife Department (2012) Texas conservation action plan 2012-2016.

Toomey, AH Knight, AT, Barlow, J (2017) Navigating the space between research and
implementation in conservation. *Conservation Letters* 10(5):619–625

U.S. Fish and Wildlife Service (2023) Recovery plan for the Rufa Red Knot (*Calidris canutus*
rufa). 22

Vallecillo, D Gauthier-Clerc, M Guillemain, M Vittecoq, M Vandewalle, P Roche, B,
Champagnon, J (2021) Reliability of animal counts and implications for the interpretation
of trends. *Ecol Evol* 11(5):2249–2260

Vermillion, WG, Lancaster, JD (2022) Habitat objectives for priority gulf coast joint venture
shorebird species. 10

Wilkinson, MD Dumontier, M Aalbersberg, IJ Appleton, G and others (2016) The FAIR Guiding
Principles for scientific data management and stewardship. *Sci Data* 3:160018

Withers, K (2002) Shorebird use of coastal wetland and barrier island habitat in the Gulf of
Mexico. *ScientificWorldJournal* 2:514–536

Young, JC Waylen, KA Sarkki, S Albon, S and others (2014) Improving the science-policy
dialogue to meet the challenges of biodiversity conservation: having conversations rather
than talking at one-another. *Biodiversity and Conservation* 23(2):387–404

Zavaleta, E Miller, DC Salafsky, N Fleishman, E and others (2008) Enhancing the engagement
of U.S. private foundations with conservation science. *Conserv Biol* 22(6):1477–1484

Figure 1. The process by which the Shorebird Science and Conservation Collective distributes

knowledge generated from electronic tracking studies of shorebirds to conservation partners

across the Americas to enhance land and species conservation, management, and outreach and

education activities. In the future, this resource also has potential to inform conservation policy.

The Collective engages with practitioners in service to community-driven management and

conservation needs, with the ultimate goal of reversing population declines.

Figure 2. Distribution of Argos and GPS tracking data (Table 2; n=1,771 individuals)

contributed to the Shorebird Science and Conservation Collective as of October 2023, shaded by

the number of species with tracking locations in 100 km x100 km hexagonal cells.

Figure 3. Example figure from a conservation project report informing the timing of freshwater

deployments to provide habitat for migratory and wintering shorebirds in Texas. A) Example

tracking data from Whimbrel, shared by Jennie Rausch, Environment and Climate Change

Canada (middle) and overlaid on a focal area of interest in Texas (right). B) Time series of when

tracked birds traveled through the Texas Gulf Coast (black histogram bars) together with median

weekly relative abundance (gray lines) in the area of interest predicted by eBird community

science data (Fink et al. 2022). Individuals tracked in more than one year contributed to

histogram counts more than once because timing varied for individuals between years. Fall

migration is shaded orange and spring migration is shaded green. Species with asterisks are

Texas State Species of Greatest Conservation Need (Texas Parks and Wildlife Department

2012). Information from A) and B) were used to determine time periods during which water

deployments could benefit peak diversity and abundances of migratory shorebirds, leading to C)

Recommendation to conservation partners for deploying water to benefit migrating and

overwintering shorebirds under recommended and alternate scenarios.

867   Table 1. Factors limiting the application of shorebird tracking data to conservation include

868   limitations to science accessibility and analysis, and limitations to conservation implementation.

Science Limitations	**Data Access** Finding and securing access to relevant data is time-consuming • Despite increasing use of data repositories (e.g. Movebank, Kays et al. 2022), only 35% of shorebird tracking datasets (reviewed by Scarpignato et al. 2023) were accessible for conservation. • Within repositories, data are spread across hundreds of unique studies led by many scientists. • Obtaining private data involves asking permission from many owners.	**Data Analysis** Analyzing tracking data for applications is not straight-forward • Analyzing tracking data requires specialized statistical and software knowledge (O'Toole et al. 2021; Joo et al. 2020; Gupte et al. 2022). • Data integration challenges are common in ecology and limit the use of datasets for the public good (Reichman et al. 2011). • Practitioners typically do not have analysis as a primary job function (Blickley et al. 2013).
Implementation Limitations	**Competing Priorities** Shorebirds are not the priority within many of their important habitats • Although interior wetland management in North America has moved from a waterfowl focus to an "all birds" approach, the extent to which shorebirds are integrated into wetland planning still varies (Doherty et al. 2018). • Many shorebirds rely on working lands where management may focus on food commodities production or on game or economically important species. • Some management programs need updating to reflect changing conservation circumstances (Sutherland et al. 2012). • Practitioners come from diverse professional backgrounds (Fernández 2016; Ciuzio et al. 2013) that may not include a background in shorebird ecology, or how to manage for them alongside primary objectives.	**Need for Strategic Integration** Diverse groups undertake shorebird conservation • Groups work at all geographic scales (Table 3), have varied funding mechanisms, operate under a variety of legal frameworks, and use varied approaches to science-based conservation implementation. • Within groups, distinct individuals can be responsible for research, regulations, program development, or technical assistance. • The lack of a central scientific resource focused on shorebirds limits the ability to integrate knowledge across groups to help target local actions to support hemispheric priorities (Brown et al. 2001).

Table 2. Description and sample size of data types submitted to the Shorebird Science and Conservation Collective as of October 2023.

TECHNOLOGY	DESCRIPTION	DATA FORMAT	SPATIAL ACCURACY	SAMPLING FREQUENCY	MONITORING DURATION	NUMBER OF INDIVIDUALS CONTRIBUTED (% of individuals)	NUMBER OF SPECIES CONTRIBUTED (% of species)
GPS	Electronic tags attached to individual birds. Locations estimated from GPS satellites and obtained either by retrieving the device and downloading stored data (i.e., archival tags) or transmitting the data to satellites.	Latitude and longitude coordinates	Typically 10-100 m (Noonan et al. 2019)	Minutes to days or pre-programmed schedules such as every 2 weeks during overwintering and 2 days during migration depending on study goal and tag battery life	Months to years depending on tag type and attachment method	1,106* (33.1 %)	17 (47.2 %)
Argos Doppler Shift	Electronic satellite tags (called Platform Terminal Transmitters or PTTs) attached to individual birds. Tags send signals to orbiting satellites that re-transmit location data back to a receiving station on Earth for processing.	Latitude and longitude coordinates with "location class" codes or error ellipses that indicate the accuracy of the estimated location	Generally 250—1,500 m (CLS America 2016) but individual locations can have errors larger than 25km (Jonsen et al. 2020)	Minutes to days, often irregular; tags may use pre-programmed on/off schedules to allow for charging (solar models), or to conserve energy (battery models)	1+ years depending on tag type and attachment method	853* (25.5 %)	14 (38.9%)
Motus	Electronic VHF radio tags attached to individual birds that	Raw detection data (e.g., timestamp,	Typically 15 – 20 km from receiving stations (Taylor et	3 - 30 seconds when tag nears receiving station,	Days to months	1,091 (32.6 %)	4 (11.1%)

	are detected by a network of receiving stations. Detection data are stored on receivers, synched over the internet, or sent via satellite or cellular network.	signal strength) associated with the location of each receiving station	al. 2017) but up to 180 km (Anderson et al. 2019); ~500 m if omnidirectional antennas are used (Taylor et al. 2017)	generally, though gaps can be minutes to days		
Light-level Geolocator	Electronic devices attached to individual birds that record ambient light levels, which can be compared to global light patterns to estimate the animal's location. Devices must be retrieved to download the data.	Raw ambient light levels which require substantial processing and analysis to estimate locations (Lisovski et al. 2020)	50 - 200 km with much greater error in latitude during spring and fall equinox (Lisovski et al. 2012; Rakhimberdiev et al. 2016)	1-2 locations per day	1+ years	483 (14.4 %)	15 (41.7 %)
Resight	Observations of individual birds marked with unique color band / flag combinations, or alphanumeric bands or flags	Latitude and longitude coordinates provided by the observer	Depends on observer proximity but usually < 100 m.	Limited to where and when observers search for birds, marker retention, and animal lifespan	Limited to observer effort, marker retention, and animal lifespan	Submitted data currently not integrated into the Collective dataset	15 (41.7%)
						* Includes 188 individuals tracked with both GPS and Argos technologies	

Table 3. Examples of organizations, initiatives, and individuals engaged in shorebird conservation and management and the geographic scale (hemispheric to local) at which these organizations operate.

Hemispheric / Continental	
Western Hemisphere Shorebird Reserve Network	A science-based, partnership-driven initiative dedicated to protecting shorebirds through a network of key sites. The Executive Office of the network is housed by Manomet, Inc., a non-governmental organization.
Shorebird Flyway Conservation Initiatives	A network of governmental and non-governmental partners working to protect shorebirds and their habitats along three shorebird flyways (Boere & Stroud 2006) of the Americas (Pacific, Midcontinent [combining the Central and Mississippi Flyways], Atlantic). Each initiative has a conservation strategy to address threats and opportunities respective to that flyway. The Arctic Migratory Bird Initiative is an initiative of the Conservation of Flora and Fauna working group of the Arctic Council, and operates at a pan-Arctic level.
Flyway Councils	A partnership of representatives from each state, province, and/or territory that facilitates the management of migratory birds and their habitats in each of four administrative migratory bird flyways in North America (Pacific, Central, Mississippi, Atlantic). Flyway Councils can establish priorities and coordinate collaborative conservation efforts to benefit shorebirds and their habitats.
Country	
Federal Wildlife and Environmental Agencies	Federal agencies focused on protecting and managing migratory birds, species at risk, and other wildlife; such agencies have the potential to target efforts towards shorebirds and their habitat (e.g., enforce policies, write management/recovery plans, monitor, conduct research). Examples include U.S. Fish and Wildlife Service and Environment and Climate Change Canada.
Region	
Migratory Bird Joint Ventures (JVs)	Cooperative, regional partnerships that work to conserve habitat for birds, people, and other wildlife. Many JVs have wetland and grassland programs that could affect or create shorebird habitat. There are 22 JVs spread across North America.
Collaborative Alliances	The Central Grasslands Roadmap (multi-country, North America), and the Southern Cone Grassland Alliance (multi-country, South America) focus on the private land management/conservation interface with potential to strengthen commitments to manage grasslands for shorebirds and to sustain human livelihoods.
State or Province	
State/Provincial Environmental Agencies	State and provincial agencies focused on the protection and management of habitats and wildlife have potential to target efforts towards shorebirds and their habitats (e.g., enforce policies, create/protect habitat, monitor species, conduct research). Examples include State Departments of Fish and Game or Natural Resources and Provincial Ministries of the Environment. Often also have local assistance programs.
County / Local	

Soil & Water Conservation Districts	U.S. county-level programs that conserve soil and water resources, many of which could affect or create shorebird habitat if targeted appropriately.
Private Landowners	Owners of lands associated with shorebird use, including agricultural lands (particularly rice fields), aquaculture (e.g., crayfish, shrimp), salt ponds, and rangelands. These landscapes have potential to affect or create shorebird habitat if managed appropriately.
Local Non-profit Groups	Non-profits that are engaged in shorebird conservation locally (sometimes connected to international initiatives). Examples from major flyways include Corbidi (Peru, Pacific Flyway), Coastal Bend Bays and Estuaries Coastal Bird Program (Texas, USA, Midcontinent), SAVE Brasil (Atlantic), and many more.
Multi-level	
Public Land Managers	Managers of protected areas can operate at local, state, or even regional and transnational levels. Protected areas are sometimes created specifically for birds, sometimes mixed recreational or human use, and sometimes strictly protected.
Natural Resources Conservation Service (NRCS)	U.S. Department of Agriculture branch that provides technical and financial assistance to farmers, ranchers, and other private landowners and managers. NRCS hosts a variety of conservation programs, typically delivered at the county-level, intended to help landowners and other partners protect, conserve, and sustainably manage their private lands. Many NRCS programs have potential to support shorebirds and/or create habitat if managed and targeted appropriately.
U.S. Fish and Wildlife Service Partners for Fish and Wildlife Program	Provides free technical and financial assistance to landowners, managers, tribes, corporations, schools and nonprofits interested in improving wildlife habitat on their lands. Projects are voluntary and customized to meet landowners' needs. Participating landowners continue to own and manage their land while they improve conditions for wildlife. Staff plan, design, supervise and monitor customized habitat restoration projects with a focus on habitats of concern like wetlands and native grasslands. Every U.S. State and Territory has a coordinator who helps deliver the program.
Land Trusts	Non-governmental organizations that place easements on or take ownership of private lands at the request of the landowner. Private lands with wetland, grassland, or working agricultural habitats have potential to support shorebirds and/or create habitat if managed appropriately. They can operate at local, regional, and national levels.
Non-profit / Non-governmental Conservation Organizations	Any non-profit / non-governmental program or organization not mentioned above that may operate at multiple levels from local to hemispheric. Examples include Birds Canada, Birds Caribbean, BirdLife International, Ducks Unlimited, National Audubon Society, Point Blue, Wetlands International, Wildlife Conservation Society, and more. Through programs to create or manage wetland, grassland, or agricultural habitat, these groups are (or could be) providing shorebird habitat.