The collective application of shorebird tracking data to conservation

Abstract

Addressing urgent conservation issues, like the drastic declines of North American migratory birds, requires creative, efficient, and collaborative approaches. Over 50% of monitored North American shorebird populations have lost over 50% of their abundance since 1980. To address these declines, we developed a partnership of scientists and practitioners called the Shorebird Science and Conservation Collective (hereinafter "the Collective"). Here, we describe the Collective's founding, structure, conservation contributions, and lessons learned to assist others engaged in translational science. The Collective acts as an intermediary whereby dedicated staff collate and analyze data contributions from scientists to support knowledge requests from conservation practitioners. Data contributions from 75 organizations include over 7.1 million shorebird locations forming movement paths of 3,420 individuals representing 36 species tracked across the Americas and have informed 13 conservation projects. As the volume of scientific data continues to grow, groups like the Collective can be vital liaisons to rapidly integrate and interpret research to support conservation action.

Autumn-Lynn Harrison[1*†], Candace Stenzel[1‡], Alexandra Anderson[1‡], Jessica Howell[1‡], Richard B. Lanctot[2†§], Marley Aikens[3¶], Joaquín Aldabe[4,5†], Liam A. Berigan[6†], Joël Bêty[7†], Erik Blomberg[6†], Juliana Bosi de Almeida[8†], Andy J. Boyce[1†], David W. Bradley[9†], Stephen Brown[10†§], Jay Carlisle[11†], Edward Cheskey[12¶], Katherine Christie[13†], Sylvain Christin[14¶], Rob Clay[15†§¶], Ashley Dayer[16§], Jill L. Deppe[17§], Willow English[18†], Scott A. Flemming[19†], Olivier Gilg[20, 21†], Christine Gilroy[14†], Susan Heath[22†], Jason M. Hill[23†], J. Mark Hipfner[19†], James A. Johnson[2†], Luanne Johnson[24†], Bart Kempenaers[25†], Paul Knaga[3¶], Eunbi Kwon[25†], Benjamin J. Lagassé[26†], Jean-François Lamarre[27†], Christopher Latty[28†], Don-Jean Léandri-Breton[29†], Nicolas Lecomte[30†], Pam Loring[31†], Laura A. McDuffie[32†], Rebecca McGuire[33, 34†], Scott Moorhead[35¶], Juan G. Navedo[36, 37, 38†], David Newstead[39†¶], Erica Nol[40†], Alina Olalla-Kerstupp[41†], Bridget Olson[42†], Elizabeth Olson[24†], Julie Paquet[14†], Allison K. Pierce[43†], Jennie Rausch[44†], Kevin Regan[45†], Matt Reiter[46§], Amber M. Roth[47†], Mike Russell[48†], Daniel Ruthrauff[32†], Sarah T. Saalfeld[2†], Amy L. Scarpignato[1†], Shiloh Schulte[49†], Nathan R. Senner[50†], Joseph A. M. Smith[51†], Paul A. Smith[52†§], Zach Spector[53¶], Kelly Srigley Werner[54§], Michelle L. Stantial[55†], Audrey R. Taylor[56†], T. Lee Tibbitts[32†], Mihai Valcu[25†], Nils Warnock[57†], Walter Wehtje[58†], Brad Winn[49†], Michael B. Wunder[43†]

[*] Corresponding Author/Principal Investigator. HarrisonAL@si.edu
[†] Data Contributor to the Collective
[‡] Collective Staff Member
[§] Member of the Collective's Advisory Group
[¶] Practitioner Contributor to the Shorebird Collective

[1] Smithsonian's National Zoo and Conservation Biology Institute, Migratory Bird Center, 3001 Connecticut Ave NW, MRC 5503, Washington, D.C. 20008, USA

[2] U.S. Fish and Wildlife Service, Division of Migratory Bird Management, 1011 E Tudor Rd, MS 201, Anchorage, AK 99503, USA

[3] Environment and Climate Change Canada, Canadian Wildlife Service, 45 Alderney Dr, Dartmouth, NS B2Y 2N6, Canada

[4] Centro Universitario Regional del Este, Universidad de la República, Departamento de Sistemas Agrarios y Paisajes Culturales, Ruta 15 y Ruta 9, Km 28.500, Departamento de Rocha, Uruguay

[5] Manomet Inc., Flyways Program, Rocha, Uruguay

[6] University of Maine, Department of Wildlife, Fisheries, and Conservation Biology, 5755 Nutting Hall, Room 210, Orono, ME 04469, USA

[7] Université du Québec à Rimouski, Département de biologie et Centre d'études Nordiques, 300, allée des Ursulines, C.P. 3300, succ. A, Rimouski, QC G5L 3A1, Canada

[8] Manomet Inc., Brasília, Brazil

[9] Birds Canada, 115 Front Rd, Port Rowan, ON N0E 1M0, Canada

[10] Manomet Inc., P.O. Box 545, Saxtons River, VT 05154, USA

[11] Boise State University, Intermountain Bird Observatory, 1910 University Dr, Boise, ID 83725, USA

[12] Nature Canada, 300-240 Bank St, Ottawa, ON K2P 1X4, Canada

[13] State of Alaska, Department of Fish and Game, Threatened, Endangered, and Diversity Program, 333 Raspberry Rd, Anchorage, AK 99518, USA

[14] Environment and Climate Change Canada, Canadian Wildlife Service, 17 Waterfowl Ln, Sackville, NB E4L 4N1, Canada

[15] Manomet Inc., WHSRN Executive Office, Asunción, Paraguay

[16] Virginia Tech, Department of Fish and Wildlife Conservation, Cheatham Hall, RM108, Blacksburg, VA 24061, USA

[17] National Audubon Society, Migratory Bird Initiative, 1828 L St NW, Washington, D.C. 20036, USA

[18] Carleton University, 1125 Colonel By Dr, Ottawa, ON K1S 5B6, Canada

[19] Environment and Climate Change Canada, Wildlife Research Division, 5421 Robertson Rd, Delta, BC V4K 3N2, Canada

[20] Laboratoire Chrono-environnement, UMR 6249/CNRS-Université de Franche-Comté, 16 route de Gray, 25030 Besançon, Bourgogne-Franche-Comté, France

[21] Groupe de Recherche en Ecologie Arctique, 16 rue de Vernot, 21440 Francheville, Bourgogne-Franche-Comté, France

[22] Gulf Coast Bird Observatory, 299 Hwy 332 West, Lake Jackson, TX 77566, USA

[23] Vermont Center for Ecostudies, 20 Palmer Ct, White River Junction, VT 05001, USA

[24] BiodiversityWorks, 18 Helen Ave, Vineyard Haven, MA 02568, USA

[25] Max Planck Institute for Biological Intelligence, Department of Ornithology, Eberhard-Gwinner-Straße, 82319 Seewiesen, Bayern, Germany

[26] University of Alaska Fairbanks, Department of Biology and Wildlife, 505 S Chandalar, Fairbanks, AK 99775, USA

[27] Polar Knowledge Canada, Canadian High Arctic Research Station Campus, 1 Uvajuq Rd, P.O. Box 2150, Cambridge Bay, NU X0B 0C0, Canada

[28] U.S. Fish and Wildlife Service, Arctic National Wildlife Refuge, 101 12th Ave, Fairbanks, AK 99701, USA

[29] McGill University, Department of Natural Resource Sciences, 21111 Lakeshore Rd, Sainte-Anne-de-Bellevue, QC H9X 3V9, Canada

[30] Centre d'Études Nordiques and Canada Research Chair in Polar and Boreal Ecology, Université de Moncton, Département de Biologie, Pavillon Rémi-Rossignol, 18, Antonine-Maillet, Moncton, NB E1A 3E9, Canada

[31] U.S. Fish and Wildlife Service, Migratory Birds Program, Populations Branch, 50 Bend Rd, Charlestown, RI 02813, USA

[48] U.S. Geological Survey, Alaska Science Center, 4210 University Dr, Anchorage, AK 99508, USA

[33] ABR, Inc. – Environmental Research & Services, P.O. Box 80410, Fairbanks, AK 99708, USA

[34] Wildlife Conservation Society, 302 Cushman Street, Suite 203, Fairbanks, AK 99701, USA

[35] Texas Water Trade, 611 S Congress Ave, Austin, TX 78704, USA

[36] Universidad Austral de Chile, Bird Ecology Lab, Instituto de Ciencias Marinas y Limnológicas, Independencia 641, Valdivia, Los Ríos, Chile

[37] Universidad Austral de Chile, Estación Experimental Quempillén, Ancud, Chiloé, Chile

[38] Millennium Institute Biodiversity of Antarctic and Subantarctic Ecosystems, Las Palmeras 3425, Ñuñoa, Santiago, Chile

[39] Coastal Bend Bays & Estuaries Program, 1305 N Shoreline Blvd #205, Corpus Christi, TX 78401, USA

[40] Trent University, Biology Department, 1600 West Bank Dr, Peterborough, ON K9L 0G2, Canada

[41] Universidad Autónoma de Nuevo León, Facultad de Ciencias Biológicas, Laboratorio de Biología de la Conservación y Desarrollo Sustentable, Ave. Universidad s/n. Ciudad Universitaria, San Nicolás de los Garza, N.L. México 66455

[42] U.S. Fish and Wildlife Service, St. Croix Wetland Management District, 1764 95th St, New Richmond, WI 54017, USA

[43] University of Colorado Denver, Department of Integrative Biology, 1201 Larimer St, Denver, CO 80204, USA

[44] Environment and Climate Change Canada, Canadian Wildlife Service, 5019 52nd St, P.O. Box 2310, Yellowknife, NT X1A 2P7, Canada

[45] Biodiversity Research Institute, 276 Canco Rd, Portland, ME 04103, USA

[46] Point Blue Conservation Science, 3820 Cypress Drive, Suite 11, Petaluma, CA 94954, USA

[47] University of Maine, School of Forest Resources and Department of Wildlife, Fisheries, and Conservation Biology, 5755 Nutting Hall, Orono, ME 04469, USA

[48] Government of Alberta, Alberta Environment and Protected Areas, 1601 Provincial Building, 10320-99 ST, Grande Prairie, AB T8V 6J4, Canada

[49] Manomet Inc., P.O. Box 1770, Manomet, MA 02345, USA

[50] University of Massachusetts Amherst, Department of Environmental Conservation, 160 Holdsworth Way, Amherst, MA 01003, USA

[51] U.S. Fish and Wildlife Service, Edwin B. Forsythe National Wildlife Refuge, 800 Great Creek Rd, Oceanville, NJ 08231, USA

[52] Environment and Climate Change Canada, National Wildlife Research Centre, 1125 Colonel By Dr, Ottawa, ON K1S 5B6, Canada

[53] Texas Parks and Wildlife Foundation, 2914 Swiss Ave, Dallas, TX 75204, USA

[54] U.S. Fish and Wildlife Service, Missouri Private Lands Office, 101 Park DeVille Drive, Suite B, Columbia, MO 65203, USA

[55] SUNY College of Environmental Science and Forestry, Department of Environmental and Forest Biology, 1 Forestry Dr, Syracuse, NY 13210, USA

[56] University of Alaska Anchorage, Department of Biological Sciences, 3211 Providence Dr, Anchorage, AK 99508, USA

[57] Audubon Canyon Ranch, 20545 Shoreline Highway 1, P.O. Box 808, Marshall, CA 94940, USA

[58] Ricketts Conservation Foundation, 607 Upper Hoback Road, P.O. Box 300, Little Jackson Hole, WY 82922, USA

Name, Email, and Mailing Address of the Person to whom Correspondence should be Sent:
Autumn-Lynn Harrison
HarrisonAL@si.edu
Smithsonian's National Zoo and Conservation Biology Institute, Migratory Bird Center, 3001 Connecticut Ave NW, MRC 5503, Washington, D.C. 20008, United States

Acknowledgments:
Support for the Shorebird Science and Conservation Collective is provided by the Knobloch Family Foundation. Additional support is provided from ConocoPhillips Charitable Investments in support of the Migratory Connectivity Project (A.-L.H.) and from the U.S. Fish and Wildlife Service (R.B.L.). Thanks to the Advisory Group (Appendix I) and for data or conservation contributions from B. Ballard, B. Corey, G. Castresana, K. De Santiago, P. Erdmann, J. Pinnix, R. Summers, J. Shackelford, D. Shaw, M. Singer, M. Weegman, L. Wright. Thanks to all who provided early feedback. Thanks to funders, collaborators, volunteers, and staff who collected or supported the collection of shorebird tracking data. Any use of trade, firm, or products is for descriptive purposes only and does not imply endorsement by the U.S. Government.

The collective application of shorebird tracking data to conservation

Abstract

Addressing urgent conservation issues, like the drastic declines of North American migratory
birds, requires creative, efficient, and collaborative approaches. Over 50% of monitored North
American shorebird populations have lost over 50% of their abundance since 1980. To address
these declines, we developed a partnership of scientists and practitioners called the Shorebird
Science and Conservation Collective (hereinafter "the Collective"). Here, we describe the
Collective's founding, structure, conservation contributions, and lessons learned to assist others
engaged in translational science. The Collective acts as an intermediary whereby dedicated staff
collate and analyze data contributions from scientists to support knowledge requests from
conservation practitioners. Data contributions from 75 organizations include over 7.1 million
shorebird locations forming movement paths of 3,420 individuals representing 36 species tracked
across the Americas and have informed 13 conservation projects. As the volume of scientific
data continues to grow, groups like the Collective can be vital liaisons to rapidly integrate and
interpret research to support conservation action.

Introduction

Conservation efforts are often location specific (Stewart et al., 2013), presenting
challenges for managing migratory species that traverse vast distances. Shorebirds
(Charadriiformes, suborder Charadrii) exemplify these challenges as many travel thousands of
kilometers biannually across continents and oceans (e.g., Gill et al., 2009). The flocking behavior
of shorebirds gives the illusion of abundance, leading to shifting baselines (*sensu* Pauly 1995)
and masking major declines. Shorebirds have the largest declines (net and % loss in abundance;
greatest proportion of species in decline) of nearly 3 billion birds lost in North America since
1970 (compared to landbirds, waterbirds, and waterfowl, Rosenberg et al., 2019). Declines are
not abating (Smith et al., 2020), emphasizing the urgent need for conservation action for
shorebirds. Here we present a partnership of scientists and practitioners (defined here as those
directly guiding conservation) responding to this need for action by linking a growing scientific
resource—shorebird tracking data—with conservation practice.

Scientists worldwide use electronic tracking devices attached to animals (hereinafter

"tags", reviewed by Robinson et al., 2010) to record movements, identify important habitats,

infer animal behavior, understand population connectivity, and monitor the environment (Kays et

al., 2015; Hussey et al., 2015). Aggregating tracking data across species has led to significant

insights, for example documenting climatic influences on animal movement patterns in the

Arctic (Davidson et al., 2020). Insights from tracking data have also yielded significant

conservation achievements. For example, tracking data collected from marine animals were used

to designate marine protected areas, reduce fisheries bycatch and vessel strikes, and minimize

impacts of extractive industries (Hays et al., 2019). Tracking data are an invaluable resource to

integrate with other occurrence datasets like local surveys and community science data.

Significant growth in shorebird tracking studies over the last 20 years offers opportunity

for tracking data to contribute to conservation. In a review of avian tracking studies (Scarpignato

et al., 2023), shorebirds and songbirds constituted <1% of studies before 2006, but these groups

constituted 30–75% of tracking studies in subsequent years. The conservation potential of these

increasingly abundant data is evident through recent examples (Chan et al., 2019; Sanders et al.,

2021; Verhoeven, 2021), but scaling up is limited by four major analytical and conservation

challenges (Table 1): limits to data availability; limits to data usability; a lack of management

focus on shorebirds in many of their important habitats; and the need for strategic integration

across organizations and habitats. While the first two challenges apply to many types of

occurrence data, they may be particularly relevant to animal tracking data (Table 1), requiring a

special effort to link these data with conservation.

To overcome challenges limiting science translation for conservation, linear and

interactive approaches are used (Beier et al., 2017; Toomey et al., 2017; Saunders et al., 2021).

Each offers benefits under different scenarios. Linear methods involve creating scientific

products to be accessed by practitioners (e.g., data portals, online tools, publications, and

manuals). Despite conservation benefits (Seavy & Howell, 2010; Sullivan et al., 2017) these

tools may not suit practical purposes due to formatting or language incompatibilities, incomplete

data, or mismatched questions, and can be time-consuming to find and synthesize (Vogel et al.

2007). Interactive approaches range from focused workshops to long-term coproduction where

research and products are collaboratively designed, planned, and implemented by practitioners

and scientists (Young et al., 2014; Roux et al., 2006; Saunders et al., 2021). These approaches

allow participants to ask clarifying questions (Gerber et al., 2020), build and maintain

relationships (Lindell & Dayer, 2022), and engage in active collaborations (Young et al., 2014).

Interactive approaches are generally more successful (Cooke et al. 2020), but there are

sometimes participation barriers (Seavy & Howell, 2010; Oliver et al., 2019).

We combined elements of both linear and interactive knowledge-sharing approaches to

form the Shorebird Science and Conservation Collective (hereinafter, 'the Collective'). The

Collective is a partnership of scientists and practitioners working to advance shorebird

conservation in the Americas (Figure 1). Our goal in sharing the origin, structure, and impact of

the Collective is to provide a case study and lessons learned to help others increase the impact of

existing science on conservation. The Collective is an intermediary whereby dedicated staff

collate and analyze data contributions from scientists to support the conservation needs of

practitioners. We describe how the Collective a) raised funds, b) initiated a proof-of-concept, c)

requested data and established operational processes, d) developed a coproduction process to

listen and respond to the needs of practitioners, and e) contributes to conservation.

Forming the idea for the Shorebird Science and Conservation Collective

The idea to collate shorebird tracking studies for conservation was inspired by the

conservation successes of BirdLife International's Seabird Tracking Database (BirdLife

International, 2004) but adapted for unique considerations of how and where shorebird

conservation is implemented (Table 1, Table 2). Given existing limitations (Table 1), a solely

linear approach facilitating data discovery or access but requiring practitioners to interpret data

and literature themselves, seemed unlikely on its own to maximize knowledge transfer. Instead,

we proposed a Collective that would extend beyond a repository to include staff for data analysis

and co-development of conservation products to support direct requests from practitioners.

A funding proposal was developed with feedback from researchers and conservation

practitioners through open-invitation webinar discussions. Many participants were already

sharing, co-developing or applying research to conservation. As conceived, the Collective would

add value by expanding conservation opportunities, rather than compete with or replace existing

partnerships. Webinars were instrumental for assessing support, refining ideas, and identifying

existing gaps in translating science to shorebird conservation.

A proof-of-concept for the Collective

We developed a three-year proof-of-concept focused on North America's interior (hereinafter, "Midcontinent", groups the Central and Mississippi migratory bird flyways) to evaluate the proposed structure and gauge demand for shorebird tracking data. The Midcontinent has been elevated as a priority landscape by multiple conservation entities (Midcontinent Shorebird Conservation Initiative, 2021; The Central Grasslands Roadmap, 2022) and provides habitat for 42 of the 50 shorebird species that regularly breed in North America (Donaldson et al., 2000; Brown et al., 2001). The Texas Gulf Coast provides migratory and wintering habitat for more than 1 million shorebirds annually (Withers, 2002) while the Prairie Pothole Region's wetlands and grasslands (Doherty et al., 2018) support millions of migrants each spring and fall (Skagen, 2006; Steen et al., 2018). These landscapes have experienced major wetland loss (up to 90% at the U.S. state level, (Dahl, 1990; Dahl, 2014)) and have high climatic variability, making cohesive shorebird management challenging across the region and among years (Fellows et al., 2001; Russell et al., 2016; Skagen & Thompson, 2013). Many areas are privately owned (e.g., nearly 90% of the Great Plains is non-public (NRCS, 2021)), requiring coordination with landowner-focused agencies which deliver voluntary incentive programs and provide technical assistance to manage habitat on working lands (e.g., U.S. Department of Agriculture Natural Resources Conservation Service, the U.S. Fish and Wildlife Service Partners for Fish and Wildlife Program), as well as state and local governments, land trusts, and other conservation entities.

The Collective's structure and systems

The Collective comprises 4 groups of people (Figure 1): a core operational team (P.I., staff, and Chair of the Advisory Group), an Advisory Group (Appendix I), data contributors, and conservation practitioners. Staff include a quantitative ecologist who aggregates, processes, and analyzes data; and conservation specialists who work with practitioners to inform them of the tracking data resource, assess their information needs, and prepare conservation products. The 16-person Advisory Group advises on maximizing the conservation benefit of the Collective and promotes its value to potential data contributors and conservation practitioner partners. Data contributors typically are researchers from public, private, or non-profit organizations.

Practitioners directly guide conservation actions within management agencies, trusts, and non-
profit groups (Table 2). Some data contributors are also practitioners.

With guidance from all involved, the core team developed operational processes
including a Data Sharing Agreement (DSA, SNZCBI, 2024a), a conservation use request form
(SNZCBI, 2024b), and a reporting system (Anderson, Stenzel and Harrison, 2024) to uphold data
privacy agreements and increase accountability to data contributors and practitioners.

Crafting a Data Sharing Agreement

Feedback indicated that careful crafting of a DSA would ensure maximum trust and
participation by the scientific community. Many factors influence data sharing (e.g., Roche et al.,
2020), with some policies mandating public sharing (e.g., the Wildlife Tracking Data Collection,
U.S. Geological Survey Alaska Science Center, 2019). Conversely, recent data or those restricted
by endangered status, may require embargo. We thus drafted a flexible agreement to facilitate
maximum data available for conservation (SNZCBI, 2024a). All contributors allow the
Collective's core team to explore data for applicability to a conservation request and to show
summary products (e.g., density grids) to conservation partners. Contributors customize
permissions (either providing pre-approval or requiring a permission request before each use)
within use categories: conservation, education, funder reports, or Collective-led scientific
publications. Contributors also indicate whether their contact information can be shared for
scholarly requests, for example a data request for a thesis or peer-reviewed publication that is not
directly tied to conservation.

For data that are publicly available, a DSA is not required for the Collective (or any
individual) to use the data and should not be viewed as restrictive. However, the Collective
requests a DSA be completed to help guide the values shaping the collaboration, provide fair
attribution, enable reporting of conservation uses to data creators, and provide engagement
opportunities to assist with analytical interpretation.

Compiling data

We invited data contributions through email, a webinar, social networks, and word-of-
mouth. Data were received either through repositories such as Movebank (Kays et al. 2022) or
were sent directly to the Collective's staff. We prioritized processing data with higher spatial

accuracy (e.g., Argos and GPS; Table 3) that we anticipated would be most relevant to

conservation requests. We developed a code pipeline (Anderson, Stenzel, and Harrison, 2024)

that first applies pre-filters (e.g., to determine when a tag has become stationary, indicating it has

either been shed or the bird has died). Then a state-space model (Jonsen et al., 2023) estimates

the most probable movement path accounting for spatial error and irregular sampling across

individuals and species. The Collective is currently integrating data from the Motus Wildlife

Tracking System (Taylor et al., 2017) and light-level geolocators (Table 3) that were contributed

either pre-processed, or in raw, unprocessed form, requiring additional processing or

standardization to ensure appropriate interpretation (Lisovski et al., 2018).

Shared data are harmonized from their sources and stored in a private, password-

protected, database. This dataset is accessible to the Collective's core team. Data are linked to

their associated DSAs to ensure that permissions and attributions are followed. Contributors are

invited annually to share new data, to review their DSA, and to review any proposed

amendments to the agreement. Outreach to new contributors occurs via presentations, email

invitations to authors of new publications or new Movebank studies, or conversations as a part of

professional collaborations. The Collective's database is updated approximately biannually to

incorporate new contributions and to ingest new data from active tags in repositories associated

with existing DSAs.

Reporting conservation uses to contributors

Contributors are notified when their data are relevant to a conservation request

(examples, Anderson, Stenzel, and Harrison, 2024). The notification describes the project and

provides a map of associated data. For any use that the contributor did not pre-approve, the

Collective requests permission before proceeding. While this step adds to the administrative

work of the Collective's staff, it incentivizes and rewards data contribution, increases

accountability, safeguards privacy settings, and promotes dialogue to prevent misinterpretations

of contributed data.

Engaging conservation partners

An initial email campaign to practitioner organizations within the Midcontinent increased

awareness of the Collective, and we continue to conduct outreach via email, presentations, and

word-of-mouth. Introductory emails describe the resources and analytical support the Collective

provides. Outreach often leads to a 1-hr virtual knowledge-sharing session between interested

practitioners and members of the Collective's core team. The team describes the Collective's

approach for data-driven conservation, and practitioners describe their work, priorities, and

shorebird-related questions. If there is opportunity to inform a project, further conversations

refine questions, information needs, and preferred formats of data products. After questions are

refined, practitioners submit a formal Conservation Use Request Form (SNZCBI, 2024b). Data

contributors are then engaged either to notify that their data are relevant to a conservation request

or to request further permission. Increasingly, the Collective receives unsolicited requests via our

online form or through emails, and from practitioners beyond our original geographic focus.

Throughout the project, the core team communicates with practitioners and data

contributors as needed. At the project's conclusion, the team presents findings to practitioners,

allowing questions and feedback regarding the delivery format, analyses, and recommendations.

A Conservation Contribution Report summarizes findings, acknowledges data contributors, and

provides information about shorebird ecology and best management practices. A public version

(with any sensitive data removed) is shared online (SNZCBI, 2024c). The Collective's core team

classifies conservation contributions following the IUCN Conservation Actions Classification

Scheme (Salafsky et al., 2008; IUCN, 2012) to align with standard evidence reporting

(Sutherland et al., 2019).

The wealth of shorebird tracking data available for conservation in the Americas

As of June 2024, data contributions include 7.1 million locations from 3,420 individual

shorebirds of 36 species (Figure 2). Since January 2021 when data were first requested,

contributors have signed 58 agreements for data originating from 75 organizations and 137 data

co-owners. Data span the Americas with areas in the Midcontinent demonstrating clear

importance to multiple species, (Figure 2). Shorebirds were tracked via light-level geolocator,

Argos, GPS, Motus technologies, and through visual observations of marked birds (Table 3,

Figure 2). A portion of tags (17% of tracked species and 22% of tracked individuals) also

provided environmental or behavioral measures (Table 3). Data come from the smallest (least

sandpiper, *Calidris minutilla,* 19–30g, tracked with 0.67-g Motus tags; (Anderson et al., 2019))

to the largest (long-billed curlew, *Numenius americanus,* 490–950 g, tracked with 9.5–22-g

satellite tags (Page et al., 2014)) shorebirds in North America (Species list, Appendix II).

Data for over half of tags (56% of tags, 26 species) were shared via Movebank (Kays et

al. 2022, 84% of birds tracked by Argos or GPS) and 32% of these (622 tags [18% of total], 10

species) currently also are open access. The remainder of data were sent to the Collective

directly, including data from the Motus Tracking Network (Taylor et al. 2017) which are

viewable online, but are not currently downloadable. Thirty-seven data sharing agreements

(63.8%) preapproved conservation uses (1,661 [49%]) tags of 23 [64%] species). Data from

only 23.6% of Argos/GPS tags are pre-approved for conservation uses. Some agreements cover

multiple datasets.

Contributing to conservation

Since our first conservation request in November 2021 to June 2024, the Collective has

supported 13 conservation projects (with another 3 ongoing). We have sent 115 notifications to

24 contributors informing them that their data collected from 17 species were relevant to

conservation requests. Sixty-six notifications were for datasets pre-approved to be used for

conservation, and 49 were requests to contributors requiring further permission. Of the latter set,

93% were approved. Below, we provide examples of ways the Collective applied tracking data to

support: 1) land conservation, 2) land and species management, and 3) education and outreach

efforts. Conservation Contribution Reports are available online as they are finalized (NZCBI,

2024c).

1. Private Land Conservation

Public and private lands support shorebird conservation in North America, but within the

Midcontinent where there is high private ownership, private lands are a crucial component of

environmental stewardship and land protection (Merenlender et al., 2004). Trusts, foundations,

and government agencies can use tracking data to help guide conservation investments and

secure funds for voluntary land purchase or conservation. Tracking datasets are especially

valuable when they provide the only evidence of shorebirds using areas (for example, when

systematic ground surveys or community science observations are unavailable).

Funding and siting of conservation easements

The Texas Parks and Wildlife Foundation's Buffer Lands Incentive Program (BLIP)

provides financial assistance for conservation easements on privately owned "buffer lands"

bordering Texas Parks and Wildlife Department (TPWD) public lands. The BLIP requested

evidence for shorebird use of buffer lands bordering 12 priority TPWD lands as one metric in

their rubric to prioritize applications for conservation easement funding; half of BLIP funds are

allocated for shorebird and grassland bird projects. Fifty-two individuals of nine species in the

Collective's dataset stopped or overwintered on buffer lands (83% of tags were deployed outside

Texas). In a report to BLIP, the Collective summarized when birds used buffer lands, the habitats

used, global migration links ranging from Russia to Argentina, a list of all TPWD lands visited

by tracked shorebirds, and state-wide density maps of individuals and species tracked.

Because many parcels under consideration are inaccessible to the public, direct evidence of

shorebird use comes almost exclusively from tracking data. This information would have likely

been unattainable by BLIP without collaborative data integration and facilitation by the

Collective.

Purchasing land for conservation

The Conservation Fund (TCF) and The Nature Conservancy (TNC), two non-

governmental conservation organizations, separately sought information to support funding

efforts to acquire lands for permanent protection in the Texas coast (TCF) and the Texas

panhandle (TNC). Although no shorebirds within the Collective's dataset visited the parcels

under consideration, 23 individuals of four species were tracked nearby (within 3–18 km). We

used habitat layers and standardized eBird relative abundance (Fink et al., 2022) to provide

additional context about the potential value of the parcels to shorebirds. Some habitats in the

parcels seemed promising, suggesting that on-the-ground surveys could be used to confirm

shorebird use. Both organizations shared these data with funders to demonstrate the potential

value of the lands for shorebirds.

2. Land and Species Management

Many shorebird conservation actions manage for habitat or species of concern (Iglecia &

Winn, 2021). Land management includes activities like deploying water to create habitat,

alternating wetting and drying of rice fields timed with shorebird usage, and prescribed grazing

and burning of grasslands or rangelands. Species management includes activities like species

recovery planning and status assessments, harvest management, and beach closures. Tracking

data can reveal important sites, connectivity among sites, individual arrival and departure timing

at particular sites, and species population structure to inform management actions.

Informing the timing and location of freshwater deployments

The Galveston Bay Foundation, the Nature Conservancy, and Texas Water Trade, three

non-governmental organizations, recently entered into multiple multiyear agreements with water

authorities to purchase water for wildlife management and environmental flows. The allocated

water volumes range from 200 to greater than 2,000 acre-feet transacted annually. They wanted

to know when and where to deploy water on private and public lands along the Texas coast to

benefit shorebirds. Water transactions offer many benefits including habitat creation and

restoration, salinity management, and continued crop yields during drought (Culp et al., 2014;

Garmany, 2020). Tracking data from six shorebird species provided information on local habitat

use and timing through the focal area (Figure 3a, 3b). Standardized relative abundance data from

eBird (Fink et al., 2022) added insights for species with limited tracking data and provided peaks

of occurrence across multiple species (Figure 3b). The Collective provided partners with

recommendations for when to deploy water (preferred and alternate times, Figure 3c) and

shorebird habitat management best practices.

Offshore wind energy development

Wind energy is important for reducing carbon emissions, but may have negative effects

on birds through collisions or displacement (Goodale & Milman, 2016; Fox & Petersen, 2019).

Tracking data can provide valuable insight about routes and passage timing in proposed and

active offshore wind areas to help understand shorebird vulnerability, inform siting decisions, or

encourage "smart curtailment" strategies during high-risk migratory periods (Loring et al., 2021;

Schwemmer et al., 2023). The Canadian Wildlife Service requested information for two rapid

(18-month) assessments offshore Nova Scotia and Newfoundland and Labrador to understand

the temporal and spatial distribution of shorebirds moving through the focal areas. The

Collective provided maps of 41 individuals of eight species that migrated through the offshore

areas of interest and highlighted places of high activity (i.e., where track lines congregated for
multiple individuals/species).

3. Education and Outreach

Despite increasing public awareness of declining bird populations, many audiences may
not be aware of the broad array of habitats shorebirds need during their migrations. The term
"shorebird" may limit conservation efforts in the Midcontinent, as many people do not connect
them to grasslands or interior wetlands. Shorebirds are sometimes wrongly assumed to have
similar habitat needs as other waterbirds like gulls, egrets, rails, or waterfowl. Environmental
education has indirect and direct conservation outcomes (Ardoin et al., 2020) and as an outreach
tool, tracking data offer strong visual and hands-on learning opportunities to gain support for
shorebird conservation efforts.

School curriculum

Educators at Grays Harbor National Wildlife Refuge in Hoquiam, Washington, USA
requested example geographic coordinates and track lines to support their 3rd and 4th grade
education programs. Educators incorporated data from 17 individuals of five species into four
lessons, allowing students to map and explore shorebird movements within the Grays Harbor
estuary and across the Pacific Flyway. The activities allowed the students to enhance skills in
geography and ecology while learning how small actions in their communities can affect a
shorebird's entire migration. Refuge educators visited multiple classrooms every month,
reaching a total of 870 students across 14 schools with these lessons over the 2022–23 school
year.

Community outreach

Nature Canada, a non-governmental conservation organization, requested tracking data
for a comic book following Bico, a fictional Hudsonian godwit (*Limosa haemastica*), as she
travels from Canada to Chile. The book's goal is to raise awareness and generate excitement
about shorebirds that visit James Bay, Canada, a globally important stopover area. The
Collective provided two godwit tracks and summary statistics (e.g., stopover sites and tenures,
average flight speeds) for an infographic at the end of the book. Nature Canada posted the comic

book online (Nature Canada, 2022) and distributed paper copies in six First Nations communities

around James Bay.

Connecting communities with shorebird movements

The Western Hemisphere Shorebird Reserve Network, a science-based, partnership-

driven, conservation initiative for protecting shorebirds through a network of key sites, wanted to

show how shorebird movements connect "sister sites" within their reserve network. The

Collective created an online storytelling series in three languages using ArcGIS StoryMaps to

follow three tracked individuals as they migrated across the Americas (SMBC and WHSRN,

2024). Each StoryMap features a migratory bird flyway (Pacific, Midcontinent, Atlantic) and

cultural theme (food, music, crafts), highlighting the people and places each shorebird might

encounter along their migrations.

Lessons Learned

The Shorebird Science and Conservation Collective is advancing conservation through

collaborative data sharing and analysis in support of needs identified by practitioners. Such on-

the-ground conservation is imperative given long-term and accelerating shorebird population

declines (Rosenberg et al., 2019; Smith et al., 2023). During the first two years we addressed

limitations to data access, analysis, and practitioners use of data insights (Table 1). As templates

for others to use, the Collective has shared forms used to communicate with practitioners and

contributors and the code pipeline used to process contributed data (Anderson, Stenzel, and

Harrison, 2024). The Collective's development offers insights for translational science initiatives

globally. Five lessons emerged that are transferable to other initiatives.

Lesson 1: Establishing access to processed usable data is crucial to meet the urgency of

our crisis discipline (Soulé, 1985). In its first year, the Collective built trust and facilitated

agreements with data owners, compiled and standardized data, and created reproducible

analytical pipelines. Having data pre-processed in multiple formats (e.g., points, density grids,

grouped by priority region) helps quickly inform varied conservation and management

discussions. However, barriers to efficiency remain and we agree with the recommendations of

Urbano and Cagnacci (2021) to streamline data sharing and harmonization (Urbano and

Cagnacci, 2021) including: supporting policy measures to improve the FAIRness of data

authorized by funding and permitting agencies (Sansone et al., 2019); facilitating communication

with data registries (Rutz, 2022; Nightingale, 2023); use of data and metadata standards to

improve datasets and increase interoperability (Campbell et al., 2016; Sequeira et al., 2021);

code sharing (Culina et al., 2020); and efforts to integrate multiple data types (Urbano et al.,

2010; Binley et al., 2023).

Lesson 2: Science translators and subject matter experts are a valuable resource for

practitioners, providing guidance on the use and interpretation of science to meet real-world

needs. Collaborative engagement and co-production require significant investment in staff salary

and time but provide significant benefit. Our impression from approximately 50 hours of

listening sessions with practitioners from 35 organizations is that interactive knowledge-sharing

significantly influences whether and how deeply shorebird science is trusted and considered for

their programs. The Collective's model of engaging data contributors with each conservation

request also increases trust with data sharing, gives reporting benefit to contributors, and

increases the network of science translators available to each conservation project. Online

decision support services or artificial intelligence tools could streamline data access and science

translation for some projects but also requires investment and continued maintenance and

development and may lack the benefits of trust and interpretability provided by direct

engagement.

Lesson 3: A focused collective can elevate a specific group (here, shorebirds) within the

priorities of on-the-ground conservation while informing science strategy. Outreach to

practitioners has raised awareness of opportunities to support shorebirds among multiple

competing priorities. While conservation initiatives vary in their prioritization of biodiversity,

biomes, function, species / guilds, and human communities, guild-specific conservation efforts

like those for waterfowl can be greatly successful (Anderson et al., 2018; Brasher et al., 2019).

The added capacity and consistent leadership of a collective becomes a powerful incentive for

agencies to prioritize group-specific conservation initiatives. The Collective's synthesized

dataset also leverages the partnership to identify conservation knowledge gaps and prioritize new

data collection, reduces the costs to achieve robust sampling designs (by pooling smaller studies

across organizations), and reduces fiscal and administrative barriers for agencies.

Lesson 4: Multiple years of support during an initiative's founding are critical. The

Collective's core team founding budget of 1.3 million USD provided three years of salary (51%)

and benefits (16%) for 3 staff members, travel (conferences and site visits, 8%), computing

equipment (1%), contract services (language translations, strategic planning, 12%), supplies and

printing (2%) support plus 10% overhead. This gave time to develop the Collective's structure,

attract strong candidates by guaranteeing three years of support, standardize data, and learn from

diverse conservation partners. Sustainability is a challenge for any new initiative. Well-resourced

stable leadership is important to maintain momentum, coordinate within and amongst scientists

and practitioners, and provide accountability. As the Collective moves from a founding to a

sustaining phase, we anticipate a model where some partners can contribute to the Collective's

sustainability through contracts or inter-agency agreements. This would ensure the Collective

can continue providing free scientific services to financially constrained groups.

Lesson 5: A proof-of-concept stage helps balance the need to develop efficient and

scalable operations for a nascent initiative with tangible mission progress. There is enough

demand for the Collective's services within the the proof-of-concept region to fill staff time.

However, migratory species conservation demands action across boundaries (Harrison et al.

2018). Requests for the Collective's support now regularly come from other regions. To expand

conservation impact, the Collective's vision is to scale up to better support practitioners in Latin

America and the Caribbean through the inclusion of tracking data from species endemic to these

regions, and through a conservation specialist based outside North America. We also hope to

inform regional, national, and international priorities through strategic analyses designed to have

the biggest effect on reversing shorebird population declines. However, the sustainability of the

Collective is not guaranteed (Lesson 4), creating a tension between maintenance and growth.

The lessons above are transferrable to groups with similar conservation aims.

Translational models of data sharing simplify complex datasets in efficient, understandable,

trusted, co-developed, and actionable ways. This lowers the activation energy for practitioners

and policymakers to use science. For migratory species, innovations in tracking technology

continue to enhance our understanding of behaviors and habitat use—patterns that are changing

with persistent habitat loss and climate change (Sutherland et al., 2012; Galbraith et al., 2014).

Ongoing data collection and sharing are vital for liaisons like the Collective to rapidly integrate

and interpret research for conservation in a changing world.

Bibliography

Anderson, A., Stenzel & Harrison, A.-L. (2024). Tracking data processing pipeline used by the

Shorebird Science and Conservation Collective (Version 0.1) [Computer software].

https://github.com/autumnlynn/ShorebirdCollective

Anderson, A. M., Duijns, S., Smith, P. A., Friis, C., & Nol, E. (2019). Migration distance and

body condition influence shorebird migration strategies and stopover decisions during

southbound migration. *Frontiers in Ecology and Evolution*, *7*, Article 251.

Anderson, M. G., Alisauskas, R. T., Batt, B. D. J., Blohm, R. J., Higgins, K. F., Perry, M. C.,

Ringelman, J. K., Sedinger, J. S., Serie, J. R., Sharp, D. E., Trauger, D. L., & Williams, C.

K. (2018). The migratory bird treaty and a century of waterfowl conservation. *The Journal*

of Wildlife Management, *82*(2), 247-259.

Ardoin, N. M., Bowers, A. W., & Gaillard, E. (2020). Environmental education outcomes for

conservation: A systematic review. *Biological Conservation*, *241*, 108224.

Bakker, V. J., Baum, J. K., Brodie, J. F., Salomon, A. K., Dickson, B. G., Gibbs, H. K., Jensen,

O. P., & McIntyre, P. B. (2010). The changing landscape of conservation science funding

in the United States. *Conservation Letters*, *3*(6), 435-444.

Beier, P., Hansen, L. J., Helbrecht, L., & Behar, D. (2017). A how-to guide for coproduction of

actionable science. *Conservation Letters*, *10*(3), 288-296.

Binley, A. D., Edwards, B. P. M., Dansereau, G., Knight, E. C., & Momeni-Dehaghi, I. (2023).

Minimizing Data Waste: Conservation in the Big Data Era. *The Bulletin of the Ecological*

Society of America, *104*(2), 1.

BirdLife International. (2004). *Tracking ocean wanderers: The global distribution of albatrosses*

and petrels. BirdLife International.

Blickley, J. L., Deiner, K., Garbach, K., Lacher, I., Meek, M. H., Porensky, L. M., Wilkerson, M.

L., Winford, E. M., & Schwartz, M. W. (2013). Graduate student's guide to necessary

skills for nonacademic conservation careers. *Conservation Biology*, *27*(1), 24-34.

Block, B. A., Jonsen, I. D., Jorgensen, S. J., Winship, A. J., Shaffer, S. A., Bograd, S. J., Hazen,

E. L., Foley, D. G., Breed, G. A., Harrison, A.-L., Ganong, J. E., Swithenbank, A.,

Castleton, M., Dewar, H., Mate, B. R., Shillinger, G. L., Schaefer, K. M., Benson, S. R.,

Weise, M. J., . . . Costa, D. P. (2011). Tracking apex marine predator movements in a

dynamic ocean. *Nature*, *475*(7354), 86-90.

Boere, G. C., & Stroud, D. A. (2006). The flyway concept: what it is and what it isn't. In G. C.

Boere, C. A. Galbraith, & D. A. Stroud (Eds.), *Waterbirds around the world* (pp. 40-47).

The Stationery Office.

Brasher, M. G., Giocomo, J. J., Azure, D. A., Bartuszevige, A. M., Flaspohler, M. E., Harrigal,

D. E., Olson, B. W., Pitre, J. M., Renner, R. W., Stephens, S. E., & Vest, J. L. (2019). The

history and importance of private lands for North American waterfowl conservation.

Wildlife Society Bulletin, *43*(3), 338-354.

Brown, S., Hickey, C., Harrington, B., & Gill, R. (Eds.). (2001). *United States Shorebird*

Conservation Plan, 2nd. ed. Manomet Center for Conservation Sciences.

Campbell, H. A., Urbano, F., Davidson, S., Dettki, H., & Cagnacci, F. (2016). A plea for

standards in reporting data collected by animal-borne electronic devices. *Animal*

Biotelemetry, *4*(1), 25.

The Central Grasslands Roadmap. 2022. Guiding us toward resilient and sustainable grasslands

and human communities. Available from https://www.grasslandsroadmap.org.

Chan, Y., Tibbitts, T. L., Lok, T., Hassell, C. J., Peng, H., Ma, Z., Zhang, Z., & Piersma, T.

(2019). Filling knowledge gaps in a threatened shorebird flyway through satellite tracking.

Journal of Applied Ecology, *56*(10), 2305-2315.

Ciuzio, E., Hohman, W. L., Martin, B., Smith, M. D., Stephens, S., Strong, A. M., &

Vercauteren, T. (2013). Opportunities and challenges to implementing bird conservation on

private lands. *Wildlife Society Bulletin*, *37*(2), 267-277.

CLS America. (2016). Argos User's Manual: Worldwide tracking and environmental monitoring

by satellite.

Culina, A., van den Berg, I., Evans, S., & Sánchez-Tójar, A. (2020). Low availability of code in

ecology: A call for urgent action. *PLoS Biology*, *18*(7), e3000763.

Culp, P. W., Glennon, R. J., & Libecap, G. (2014). Shopping for water: How the market can

mitigate water shortages in the American West [White Paper]. The Hamilton Project,

Brookings, Stanford Woods Institute for the Environment.

https://www.brookings.edu/articles/shopping-for-water-how-the-market-can-mitigate-

water-shortages-in-the-american-west/

Dahl, T. E. (1990). *Wetlands losses in the United States 1780's to 1980's. Report to Congress*.

US Department of the Interior, Fish and Wildlife Service.

Dahl, T. E. (2014). *Status and trends of prairie wetlands in the United States 1997 to 2009*. US

Fish and Wildlife Service Washington, DC.

Davidson, S. C., Bohrer, G., Gurarie, E., LaPoint, S., Mahoney, P. J., Boelman, N. T., Eitel, J. U.

H., Prugh, L. R., Vierling, L. A., Jennewein, J., Grier, E., Couriot, O., Kelly, A. P.,

Meddens, A. J. H., Oliver, R. Y., Kays, R., Wikelski, M., Aarvak, T., Ackerman, J.

T., . . . Hebblewhite, M. (2020). Ecological insights from three decades of animal

movement tracking across a changing Arctic. *Science*, *370*(6517), 712-715.

Doherty, K. E., Howerter, D. W., Devries, J. H., & Walker, J. (2018). Prairie pothole region of

North America. In *The Wetland Book* (pp. 679-688). Springer Netherlands.

https://doi.org/10.1007/978-94-007-4001-3_15

Donaldson, G. M., Hyslop, C., Morrison, R. I. G., Dickson, H. L., & Davidson, I. (2000).

Canadian Shorebird Conservation Plan [White Paper]. Canadian Wildlife Service.

https://publications.gc.ca/collections/collection_2011/ec/CW69-15-5-2000-eng.pdf

Environment and Climate Change Canada. (2016). Recovery strategy and management plan for

the Red Knot (*Calidris canutus*) in Canada [Proposed]. *Species at Risk Act* Recovery

Strategy Series. [White Paper]. Environment and Climate Change Canada.

https://publications.gc.ca/collections/collection_2018/eccc/En3-4-260-2017-eng.pdf

Farmer, A., & Durbian, F. (2006). Estimating shorebird numbers at migration stopover sites. *The*

Condor, *108*(4), 792-807.

Fellows, S., Stone, K., Jones, S., Damude, N., & Brown, S. (2001). Central Plains/Playa Lakes

regional shorebird conservation plan. *Version*, *1*, 26.

Fernández, R. J. (2016). How to be a more effective environmental scientist in management and

policy contexts. *Environmental Science & Policy*, *64*, 171-176.

Fink, D., Auer, T., Johnston, A., Strimas-Mackey, M., Robinson, O., Ligocki, S., Hochachka,

W., Jaromczyk, L., Rodewald, A., Wood, C., Davies, I., & Spencer, A. (2022). eBird Status

and Trends. Data Version: 2021; Released: 2022.

Fox, A. D., & Petersen, I. K. (2019). Offshore wind farms and their effects on birds. *Dansk*

Ornitologisk Forenings Tidsskrift, *113*, 86-101.

Galbraith, H., DesRochers, D. W., Brown, S., & Reed, J. M. (2014). Predicting vulnerabilities of

North American shorebirds to climate change. *PLoS One*, *9*(9), e108899.

Garmany, K. (2020). *Market Strategies for Addressing Water Scarcity: An Introduction to*

Environmental Water Transactions in Texas. A thesis in partial fulfillment of the

requirements for a degree of Master of Applied Geography. Texas State University].

https://digital.library.txstate.edu/bitstream/handle/10877/11993/Garmany-

Kyle.pdf?sequence=1.

Gerber, L. R., Barton, C. J., Cheng, S. H., & Anderson, D. (2020). Producing actionable science

in conservation: Best practices for organizations and individuals. *Conservation Science and*

Practice, *2*(12), 2:e295.

Gill, R. E., Tibbitts, T. L., Douglas, D. C., Handel, C. M., Mulcahy, D. M., Gottschalck, J. C.,

Warnock, N., McCaffery, B. J., Battley, P. F., & Piersma, T. (2009). Extreme endurance

flights by landbirds crossing the Pacific Ocean: ecological corridor rather than barrier.

Proceedings of the Royal Society B: Biological Sciences, *276*(1656), 447-457.

Gillings, S., Atkinson, P. W., Baker, A. J., Bennett, K. A., Clark, N. A., Cole, K. B., González,

P. M., Kalasz, K. S., Minton, C. D. T., & Niles, L. J. (2009). Staging behavior in Red Knot

(*Calidris canutus*) in Delaware Bay: implications for monitoring mass and population size.

The Auk, *126*(1), 54-63.

Goodale, M. W., & Milman, A. (2016). Cumulative adverse effects of offshore wind energy

development on wildlife. *Journal of Environmental Planning and Management*, *59*(1), 1-

21.

Gupte, P. R., Beardsworth, C. E., Spiegel, O., Lourie, E., Toledo, S., Nathan, R., & Bijleveld, A.

I. (2022). A guide to pre-processing high-throughput animal tracking data. *Journal of*

Animal Ecology, *91*(2), 287-307.

Haila, Y., & Kouki, J. (1994). *The phenomenon of biodiversity in conservation biology*. In

Annales Zoologici Fennici (pp. 5-18). JSTOR.

https://www.jstor.org/stable/pdf/23735495.pdf?casa_token=v-

LlhTxkBSMAAAAA:gWRnI_bmQ4XHjwyImdZQOgVIkdHp4N3Ed2UtmXzqkMlzu7pp

FFtowXQJSROT-AXboajpR3QgmKjk3fpfpQmLZSLjLAGIAf3WZ6yv8Ct21-

WlXYKgtIzS

Harrison, A.-L., Costa, D. P., Winship, A. J., Benson, S. R., Bograd, S. J., Antolos, M., Carlisle,

A. B., Dewar, H., Dutton, P. H., Jorgensen, S. J., Kohin, S., Mate, B. R., Robinson, P. W.,

Schaefer, K. M., Shaffer, S. A., Shillinger, G. L., Simmons, S. E., Weng, K. C., Gjerde, K.

M., . . . Block, B. A. (2018). The political biogeography of migratory marine predators.

Nature Ecology & Evolution, *2*(10), 1571-1578.Hays, G. C., Bailey, H., Bograd, S. J.,

Bowen, W. D., Campagna, C., Carmichael, R. H., Casale, P., Chiaradia, A., Costa, D. P.,

Cuevas, E., Bruyn, P. J. N. D., Dias, M. P., Duarte, C. M., Dunn, D. C., Dutton, P. H.,

Esteban, N., Friedlaender, A., Goetz, K. T., Godley, B. J., . . . Sequeira, A. M. M. (2019).

Translating marine animal tracking data into conservation policy and management. *Trends*

in Ecology & Evolution, *34*(5), 459-472.

Hebblewhite, M., & Haydon, D. T. (2010). Distinguishing technology from biology: a critical

review of the use of GPS telemetry data in ecology. *Philosophical Transactions of the*

Royal Society B, *365*(1550), 2303-2312.

Hussey, N. E., Kessel, S. T., Aarestrup, K., Cooke, S. J., Cowley, P. D., Fisk, A. T., Harcourt, R.

G., Holland, K. N., Iverson, S. J., Kocik, J. F., Mills Flemming, J. E., & Whoriskey, F. G.

(2015). Aquatic animal telemetry: A panoramic window into the underwater world.

Science, *348*(6240), 1255642.

Iglecia, M., & Winn, B. (2021). A shorebird management manual [White Paper].

https://www.manomet.org/wp-

content/uploads/2021/01/Iglecia_and_Winn_2021_AShorebirdManagementManual-

012021-web.pdf

IUCN. (2012). *Conservation Actions Classification Scheme (Version 2.0)*.

https://www.iucnredlist.org/resources/conservation-actions-classification-scheme

Jonsen, I. D., Grecian, W. J., Phillips, L., Carroll, G., McMahon, C., Harcourt, R. G., Hindell, M.

A., & Patterson, T. A. (2023). aniMotum, an R package for animal movement data: Rapid

quality control, behavioural estimation and simulation. *Methods in Ecology and Evolution*,

14(3), 806-816.

Jonsen, I. D., Patterson, T. A., Costa, D. P., Doherty, P. D., Godley, B. J., Grecian, W. J., Guinet,

C., Hoenner, X., Kienle, S. S., Robinson, P. W., Votier, S. C., Whiting, S., Witt, M. J.,

Hindell, M. A., Harcourt, R. G., & McMahon, C. R. (2020). A continuous-time state-space

model for rapid quality control of argos locations from animal-borne tags. *Movement*

Ecology, *8*, 31.

Joo, R., Boone, M. E., Clay, T. A., Patrick, S. C., Clusella-Trullas, S., & Basille, M. (2020).
Navigating through the r packages for movement. *Journal of Animal Ecology*, *89*(1), 248-
267.

Kays, R., Crofoot, M. C., Jetz, W., & Wikelski, M. (2015). Terrestrial animal tracking as an eye
on life and planet. *Science*, *348*(6240), aaa2478.

Kays, R., Davidson, S. C., Berger, M., Bohrer, G., Fiedler, W., Flack, A., Hirt, J., Hahn, C.,
Gauggel, D., Russell, B., Kölzsch, A., Lohr, A., Partecke, J., Quetting, M., Safi, K., Scharf,
A., Schneider, G., Lang, I., Schaeuffelhut, F., . . . Wikelski, M. (2022). The Movebank
system for studying global animal movement and demography. *Methods in Ecology and*
Evolution, *13*(2), 419-431.

Lindell, C. A., & Dayer, A. A. (2022). Six principles for working effectively with landowners to
advance bird conservation. *Ornithological Applications*, *124*(4).

Lisovski, S., Bauer, S., Briedis, M., Davidson, S. C., Dhanjal-Adams, K. L., Hallworth, M. T.,
Karagicheva, J., Meier, C. M., Merkel, B., Ouwehand, J., Pedersen, L., Rakhimberdiev, E.,
Roberto-Charron, A., Seavy, N. E., Sumner, M. D., Taylor, C. M., Wotherspoon, S. J., &
Bridge, E. S. (2020). Light-level geolocator analyses: A user's guide. *Journal of Animal*
Ecology, *89*, 221-236.

Lisovski, S., Hewson, C. M., Klaassen, R. H. G., Korner-Nievergelt, F., Kristensen, M. W., &
Hahn, S. (2012). Geolocation by light: accuracy and precision affected by environmental
factors. *Methods in Ecology and Evolution*, *3*, 603-612.

Lisovski, S., Schmaljohann, H., Bridge, E. S., Bauer, S., Farnsworth, A., Gauthreaux, S. A.,
Hahn, S., Hallworth, M. T., Hewson, C. M., Kelly, J. F., Liechti, F., Marra, P. P.,
Rakhimberdiev, E., Ross, J. D., Seavy, N. E., Sumner, M. D., Taylor, C. M., Winkler, D.
W., Wotherspoon, S. J., . . . Wunder, M. B. (2018). Inherent limits of light-level
geolocation may lead to over-interpretation. *Current Biology*, *28*(3), R99-R100.

Loesch, C. R., Twedt, D. J., Tripp, K., Hunter, W. C., & Woodrey, M. S. (2000). *Development*
of management objectives for waterfowl and shorebirds in the Mississippi Alluvial Valley.
In *Proceedings of the 3rd partners in flight workshop. USDA Forest Service Proceedings*
RMRS-P-16, Cape May, NJ (pp. 8-11).

Loring, P. H., Lenske, A. K., McLaren, J. D., Aikens, M., Anderson, A. M., Aubrey, Y., Dalton,
E., Dey, A., Friis, C., Hamilton, D., Holberton, R. L., Kriensky, D., Mizrahi, D., Niles, L.

J., Parkins, K. L., Paquet, J., Sanders, F., Smith, A., Turcotte, Y., . . . Smith, P. A. (2021).

Tracking movements of migratory shorebirds in the US Atlantic Outer Continental Shelf

Region (8). US Department of the Interior, Bureau of Ocean Energy Management.

Merenlender, A. M., Huntsinger, L., Guthey, G., & Fairfax, S. K. (2004). Land trusts and

conservation easements: Who is conserving what for whom? *Conservation Biology*, *18*(1),

65-76.

Midcontinent Shorebird Conservation Intiative (2024). Available from

https://midamericasshorebirds.org.

Nature Canada, (2022), The story of Bico, the Hudsonian Godwit: Available from

https://naturecanada.ca/discover-nature/about-our-birds/rufus-red-knot/

Navedo, J. G., & Piersma, T. (2023). Do 50-year-old Ramsar criteria still do the best possible

job? A plea for broadened scientific underpinning of the global protection of wetlands and

migratory waterbirds. *Conservation Letters*, *16*(2).

Nightingale, J. (2023). Conserving wader tracking data for the future: Introducing the Global

Wader project. *Wader Study*, *130*(3).

Noonan, M. J., Fleming, C. H., Akre, T. S., Drescher-Lehman, J., Gurarie, E., Harrison, A. L.,

Kays, R., & Calabrese, J. M. (2019). Scale-insensitive estimation of speed and distance

traveled from animal tracking data. *Movement Ecology*, *7*, 35.

NRCS. (2021). A framework for conservation action in the Great Plains Grasslands Biome

[White Paper]. Working Lands for Wildlife, USDA-Natural Resources Conservation

Service (NRCS). https://www.nrcs.usda.gov/sites/default/files/2022-

06/greatPlainsFramework.pdf

O'Toole, M., Queiroz, N., Humphries, N. E., Sims, D. W., & Sequeira, A. M. M. (2021).

Quantifying effects of tracking data bias on species distribution models. *Methods in*

Ecology and Evolution, *12*(1), 170-181.

Oliver, K., Kothari, A., & Mays, N. (2019). The dark side of coproduction: do the costs outweigh

the benefits for health research. *Health Research Policy and Systems*, *17*(1), 33.

Page, G. W., Warnock, N., Tibbitts, T. L., Jorgensen, D., Hartman, C. A., & Stenzel, L. E.

(2014). Annual migratory patterns of Long-billed Curlews in the American West. *The*

Condor, *116*(1), 50-61.

Rakhimberdiev, E., Senner, N. R., Verhoeven, M. A., Winkler, D. W., Bouten, W., & Piersma,

T. (2016). Comparing inferences of solar geolocation data against high-precision GPS data:

annual movements of a double-tagged black-tailed godwit. *Journal of Avian Biology*,

47(4), 589-596.

Reichman, O. J., Jones, M. B., & Schildhauer, M. P. (2011). Challenges and opportunities of

open data in ecology. *Science*, *331*(6018), 703-705.

Robinson, W. D., Bowlin, M. S., Bisson, I., Shamoun-Baranes, J., Thorup, K., Diehl, R. H.,

Kunz, T. H., Mabey, S., & Winkler, D. W. (2010). Integrating concepts and technologies to

advance the study of bird migration. *Frontiers in Ecology and the Environment*, *8*(7), 354-

361.

Roche, D. G., Granados, M., Austin, C. C., Wilson, S., Mitchell, G. M., Smith, P. A., Cooke, S.

J., & Bennett, J. R. (2020). Open government data and environmental science: a federal

Canadian perspective. *FACETS*, *5*(1), 942-962.

Rosenberg, K. V., Dokter, A. M., Blancher, P. J., Sauer, J. R., Smith, A. C., Smith, P. A.,

Stanton, J. C., Panjabi, A., Helft, L., Parr, M., & Marra, P. P. (2019). Decline of the North

American avifauna. *Science*, *366*(6461), 120-124.

Roux, D. J., Rogers, K. H., Biggs, H. C., Ashton, P. J., & Sergeant, A. (2006). Bridging the

science–management divide: moving from unidirectional knowledge transfer to knowledge

interfacing and sharing. *Ecology and Society*, *11*(1), 4.

Russell, R. P., Koch, K. E., & Lewis, S. J. (2016). *Upper Mississippi Valley/Great Lakes*

Regional Shorebird Conservation Plan. Version 2.0. U.S. Fish & Wildlife Service,

Division of Migratory Birds.

Rutz, C. (2022). Register animal-tracking tags to boost conservation. *Nature*, *609*(7926), 221-

221.

Salafsky, N., Salzer, D., Stattersfield, A. J., Hilton-Taylor, C., Neugarten, R., Butchart, S. H.,

Collen, B., Cox, N., Master, L. L., O'Connor, S., & Wilkie, D. (2008). A standard lexicon

for biodiversity conservation: unified classifications of threats and actions. *Conservation*

Biology, *22*(4), 897-911.

Salafsky, N., Margoluis, R., Redford, K. H., & Robinson, J. G. (2002). Improving the Practice of

Conservation: a Conceptual Framework and Research Agenda for Conservation Science.

Conservation Biology, *16*(6), 1469-1479.

Sanders, F. J., Handmaker, M. C., Johnson, A. S., & Senner, N. R. (2021). Nocturnal roost on
South Carolina coast supports nearly half of Atlantic coast population of Hudsonian
Whimbrel Numenius hudsonicus during northward migration. *Wader Study*, *128*(2), 4.

Sansone, S.-A., McQuilton, P., Rocca-Serra, P., Gonzalez-Beltran, A., Izzo, M., Lister, A. L.,
Thurston, M., & Community, F. A. I. R. (2019). FAIRsharing as a community approach to
standards, repositories and policies. *Nature biotechnology*, *37*(4), 358-367.

Saunders, S. P., Wu, J. X., Gow, E. A., Adams, E., Bateman, B. L., Bayard, T., Beilke, S., Dayer,
A. A., Fournier, A. M. V., Fox, K., Heglund, P., Lerman, S. B., Michel, N. L., Paxton, E.
H., Şekercioğlu, Ç. H., Smith, M. A., Thogmartin, W., Woodrey, M. S., & van Riper, C.
(2021). Bridging the research-implementation gap in avian conservation with translational
ecology. *Ornithological Applications*, *123*(3), 1-13.

Scarpignato, A. L., Huysman, A. E., Jimenez, M. F., Witko, C. J., Harrison, A.-L., Seavy, N. E.,
Smith, M. A., Deppe, J. L., Wilsey, C. B., & Marra, P. P. (2023). Shortfalls in tracking data
available to inform North American migratory bird conservation. *Biological Conservation*,
286, 110224.

Schwemmer, P., Pederson, R., Haecker, K., Bocher, P., Fort, J., Mercker, M., Jiguet, F., Elts, J.,
Marja, R., Piha, M., Rousseau, P., & Garthe, S. (2023). Assessing potential conflicts
between offshore wind farms and migration patterns of a threatened shorebird species.
Animal Conservation, *26*(3), 303-316.

Seavy, N. E., & Howell, C. A. (2010). How can we improve information delivery to support
conservation and restoration decisions. *Biodiversity and Conservation*, *19*(5), 1261-1267.

Sequeira, A. M. M., Heupel, M. R., Lea, M. -. A., Eguíluz, V. M., Duarte, C. M., Meekan, M. G.,
Thums, M., Calich, H. J., Carmichael, R. H., Costa, D. P., Ferreira, L. C., Fernandéz-
Gracia, J., Harcourt, R., Harrison, A. -. L., Jonsen, I., McMahon, C. R., Sims, D. W.,
Wilson, R. P., & Hays, G. C. (2019). The importance of sample size in marine megafauna
tagging studies. *Ecological Applications*, *29*(6), e01947.

Sequeira, A. M. M., O'Toole, M., Keates, T. R., McDonnell, L. H., Braun, C. D., Hoenner, X.,
Jaine, F. R. A., Jonsen, I. D., Newman, P., Pye, J., Bograd, S. J., Hays, G. C., Hazen, E. L.,
Holland, M., Tsontos, V. M., Blight, C., Cagnacci, F., Davidson, S. C., Dettki,
H., . . . Weise, M. (2021). A standardisation framework for bio-logging data to advance
ecological research and conservation. *Methods in Ecology and Evolution*, *12*(6), 996-1007.

705 Skagen, S. K. (2006). Migration stopovers and the conservation of arctic-breeding calidridine

706   sandpipers. *The Auk*, *123*(2), 313-322.

707 Skagen, S. K., & Thompson, G. (2013). U.S. shorebird conservation plan: Northern

708   Plains/Prairie Potholes regional shorebird conservation plan [White Paper].

709   https://www.shorebirdplan.org/wp-content/uploads/2013/01/NORPLPP2.pdf

710 Smith, M. A., Mahoney, J., Knight, E. J., Taylor, L., Seavy, N. E., Bailey, C. H., Carbone, M.,

711   DeLuca, W. V., Gonzalez, N. S., Jimenez, M. F., O'Bryan, G. M. W., Rao, N., Witko, C.

712   J., Wilsey, C. B., & Deppe, J. L. (2022). *Bird Migration Explorer*.

713   birdmigrationexplorer.org / exploraves.org

714 Smith, P. A., McKinnon, L., Meltofte, H., Lanctot, R. B., Fox, A. D., Leafloor, J. O., Soloviev,

715   M., Franke, A., Falk, K., Golovatin, M., Sokolov, V., Sokolov, A., & Smith, A. C. (2020).

716   Status and trends of tundra birds across the circumpolar Arctic. *Ambio*, *49*(3), 732-748.

717 Smith, P. A., Smith, A. C., Andres, B., Francis, C. M., Harrington, B., Friis, C., Morrison, R. I.

718   G., Paquet, J., Winn, B., & Brown, S. (2023). Accelerating declines of North America's

719   shorebirds signal the need for urgent conservation action. *Ornithological Applications*,

720   *125*(2).

721 Smithsonian Migratory Bird Center (SMBC) and Western Hemisphere Shorebird Group. (2024).

722   Cultural connections across migratory flyways. Available from

723   https://storymaps.arcgis.com/collections/8b0faf990fc84e70b7aa09900634a560.

724 Smithsonian's National Zoo & Conservation Biology Institute (SNZCBI). (2024a). Shorebird

725   Science and Conservation Collective Data Sharing Agreement Form. Available from

726   https://nationalzoo.si.edu/migratory-birds/shorebird-science-and-conservation-collective-

727   data-sharing-agreement-form

728 Smithsonian's National Zoo & Conservation Biology Institute (SNZCBI). (2024b). Request for

729   use of shorebird tracking data. Available from https://nationalzoo.si.edu/migratory-birds/

730   request-for-use-shorebird-tracking-data

731 Smithsonian's National Zoo & Conservation Biology Institute (SNZCBI). (2024c). Shorebird

732   Science and Conservation Collective Website. Available from

733   https://nationalzoo.si.edu/migratory-birds/shorebird-collective

734 Soulé, M. E. (1985). What is conservation biology? *BioScience*, *35*(11), 727-734.

Steen, V., Skagen, S. K., & Noon, B. R. (2018). Preparing for an uncertain future: migrating

shorebird response to past climatic fluctuations in the Prairie Potholes. *Ecosphere*, *9*(2),

e02095.

Stewart, W. P., Williams, D. R., & Kruger, L. E. (Eds.). (2013). *Place-Based Conservation.*

Perspectives from the Social Sciences. Springer Netherlands. https://doi.org/10.1007/978-

94-007-5802-5

Sullivan, B. L., Phillips, T., Dayer, A. A., Wood, C. L., Farnsworth, A., Iliff, M. J., Davies, I. J.,

Wiggins, A., Fink, D., Hochachka, W. M., Rodewald, A. D., Rosenberg, K. V., Bonney, R.,

& Kelling, S. (2017). Using open access observational data for conservation action: A case

study for birds. *Biological Conservation*, *208*, 5-14.

Sutherland, W. J., Alves, J. A., Amano, T., Chang, C. H., Davidson, N. C., Max Finlayson, C.,

Gill, J. A., Gill Jr, R. E., González, P. M., & Gunnarsson, T. G. (2012). A horizon scanning

assessment of current and potential future threats to migratory shorebirds. *Ibis*, *154*(4), 663-

679.

Sutherland, W. J., Taylor, N. G., MacFarlane, D., Amano, T., Christie, A. P., Dicks, L. V.,

Lemasson, A. J., Littlewood, N. A., Martin, P. A., Ockendon, N., Petrovan, S. O.,

Robertson, R. J., Rocha, R., Shackelford, G. E., Smith, R. K., Tyler, E. H. M., & Wordley,

C. F. R. (2019). Building a tool to overcome barriers in research-implementation spaces:

The Conservation Evidence database. *Biological Conservation*, *238*, 108199.

Taylor, P. D., Crewe, T. L., Mackenzie, S. A., Lepage, D., Aubry, Y., Crysler, Z., Finney, G.,

Francis, C. M., Guglielmo, C. G., Hamilton, D. J., Holberton, R. L., Loring, P. H.,

Mitchell, G. W., Norris, D. R., Paquet, J., Ronconi, R. A., Smetzer, J. R., Smith, P. A.,

Welch, L. J., . . . Woodworth, B. K. (2017). The Motus Wildlife Tracking System: a

collaborative research network to enhance the understanding of wildlife movement. *Avian*

Conservation and Ecology, *12*(1), Article 8.

Texas Parks and Wildlife Department. (2012). *Texas conservation action plan 2012-2016*.

Toomey, A. H., Knight, A. T., & Barlow, J. (2017). Navigating the space between research and

implementation in conservation. *Conservation Letters*, *10*(5), 619-625.

U.S. Fish and Wildlife Service. (2023). *Recovery plan for the Rufa Red Knot (*Calidris canutus

rufa*)*. U.S. Fish and Wildlife Service, Northeast Region.

Urbano, F., Cagnacci, F., Calenge, C., Dettki, H., Cameron, A., & Neteler, M. (2010). Wildlife
tracking data management: a new vision. *Philos Trans R Soc Lond B Biol Sci*, *365*(1550),
2177-2185.

Urbano, F., Cagnacci, F., & Euromammals, C. I. (2021). Data Management and Sharing for
Collaborative Science: Lessons Learnt From the Euromammals Initiative. *Frontiers in*
Ecology and Evolution, *9*, 15.

U.S. Geological Survey, Alaska Science Center, 2019, USGS Alaska Science Center wildlife
tracking data collection: U.S. Geological Survey data release,
http://doi.org/10.5066/P9VYSWEH

Vallecillo, D., Gauthier-Clerc, M., Guillemain, M., Vittecoq, M., Vandewalle, P., Roche, B., &
Champagnon, J. (2021). Reliability of animal counts and implications for the interpretation
of trends. *Ecology and Evolution*, *11*(5), 2249-2260.

Verhoeven, M. (2021). Wader community – The power of collaboration. *Wader Study*, *128*(2), 0.

Vermillion, W. G., & Lancaster, J. D. (2022). Habitat objectives for priority gulf coast joint
venture shorebird species [White Paper]. Gulf Coast Joint Venture.
https://gcjv.org/docs/conservation_plans/habitat_objectives_for_gcjv_shorebirds_2022.pdf

Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A.,
Blomberg, N., Boiten, J. W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes,
A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers,
R., . . . Mons, B. (2016). The FAIR Guiding Principles for scientific data management and
stewardship. *Scientific Data*, *3*, 160018.

Withers, K. (2002). Shorebird use of coastal wetland and barrier island habitat in the Gulf of
Mexico. *The Scientific World Journal*, *2*, 514-536.

Young, J. C., Waylen, K. A., Sarkki, S., Albon, S., Bainbridge, I., Balian, E., Davidson, J.,
Edwards, D., Fairley, R., Margerison, C., McCracken, D., Owen, R., Quine, C. P., Stewart-
Roper, C., Thompson, D., Tinch, R., Van den Hove, S., & Watt, A. (2014). Improving the
science-policy dialogue to meet the challenges of biodiversity conservation: having
conversations rather than talking at one-another. *Biodiversity and Conservation*, *23*(2),
387-404.

Zavaleta, E., Miller, D. C., Salafsky, N., Fleishman, E., Webster, M., Gold, B., Hulse, D.,
Rowen, M., Tabor, G., & Vanderryn, J. (2008). Enhancing the engagement of U.S. private
foundations with conservation science. *Conservation Biology*, *22*(6), 1477-1484.

Table 1. Factors limiting the application of shorebird tracking data to conservation include

limitations to science accessibility and analysis, and limitations to conservation implementation.

	Data Access Finding and securing access to relevant data is time-consuming	**Data Integration and Analysis** Analyzing tracking data for applications is not straight-forward
Science Limitations	• Despite increasing use of data repositories (e.g., Movebank, Kays et al., 2022), many tracking datasets are not open access (reviewed by Scarpignato et al., 2023). • Within repositories, data are spread across hundreds of unique studies led by many scientists. • Obtaining private data involves asking permission from many owners.	• Preparing, integrating, and analyzing tracking data requires specialized statistical and software knowledge (O'Toole et al., 2021; Joo et al., 2020; Gupte et al., 2022). • Data integration challenges are common in ecology and limit the use of datasets for the public good (Reichman et al., 2011). • Practitioners typically do not have data analysis as a primary job function (Blickley et al., 2013).
	Competing Priorities Shorebirds are not the priority within many of their important habitats	**Need for Strategic Integration** Diverse groups undertake shorebird conservation
Implementation Limitations	• Although interior wetland management in North America has moved from a waterfowl focus to an "all birds" approach, the extent to which shorebirds are integrated into wetland planning still varies (Doherty et al., 2018). • Many shorebirds rely on working lands where management may focus on food commodities production or on game or economically important species. • Some management programs need updating to reflect changing conservation circumstances (Sutherland et al., 2012). • Practitioners come from diverse professional backgrounds (Fernández, 2016; Ciuzio et al., 2013) that may not include a background in shorebird ecology, or how to manage for them alongside primary objectives.	• Groups work at all geographic scales (Table 2), have varied funding mechanisms, operate under a variety of legal frameworks, and use varied approaches to science-based conservation implementation. • Within groups, distinct individuals can be responsible for research, regulations, program development, or technical assistance. • The lack of a central scientific resource focused on shorebirds limits the ability to integrate knowledge across groups to help target local actions to support hemispheric priorities (Brown et al., 2001).

  Table 2. Examples of organizations, initiatives, and individuals engaged in shorebird
803 conservation and management and the geographic scale (hemispheric to local) at which these
804 organizations operate.
805

Hemispheric / Continental	
Western Hemisphere Shorebird Reserve Network	A science-based, partnership-driven initiative dedicated to protecting shorebirds through a network of key sites. The Executive Office of the network is housed by Manomet, Inc., a non-governmental organization.
Shorebird Flyway Conservation Initiatives	A network of governmental and non-governmental partners working to protect shorebirds and their habitats along three shorebird flyways (Boere & Stroud, 2006) of the Americas (Pacific, Midcontinent [combining the Central and Mississippi Flyways], Atlantic). Each initiative has a conservation strategy to address threats and opportunities respective to that flyway. The Arctic Migratory Bird Initiative is an initiative of the Conservation of Flora and Fauna working group of the Arctic Council and operates at a pan-Arctic level.
Flyway Councils	A partnership of representatives from states, provinces, and territories that facilitates the management of migratory birds and their habitats in each of four administrative migratory bird flyways in North America (Pacific, Central, Mississippi, Atlantic). Flyway Councils can establish priorities and coordinate collaborative conservation efforts to benefit shorebirds and their habitats.
Country	
Federal Wildlife and Environmental Agencies	Federal agencies focused on protecting and managing migratory birds, species at risk, and other wildlife; such agencies have the potential to target efforts towards shorebirds and their habitat (e.g., enforce policies, write management/recovery plans, monitor, conduct research). Examples include U.S. Fish and Wildlife Service and Environment and Climate Change Canada.
Region	
Migratory Bird Joint Ventures (JVs)	Cooperative, regional partnerships that work to conserve habitat for birds, people, and other wildlife. Many JVs have wetland and grassland programs that could affect or create shorebird habitat. There are 22 JVs spread across North America.
Collaborative Alliances	The Central Grasslands Roadmap (multi-country, North America), and the Southern Cone Grassland Alliance (multi-country, South America) focus on the private land management/conservation interface with potential to strengthen commitments to manage grasslands for shorebirds and to sustain human livelihoods.
State or Province	
State/Provincial Environmental Agencies	State and provincial agencies focused on the protection and management of habitats and wildlife have potential to target efforts towards shorebirds and their habitats (e.g., enforce policies, create/protect habitat, monitor species, conduct research). Examples include State Departments of Fish and Game or Natural Resources and Provincial Ministries of the Environment. Often also have local assistance programs.
County / Local	

Soil & Water Conservation Districts	U.S. county-level programs that conserve soil and water resources, many of which could affect or create shorebird habitat if targeted appropriately.
Private Landowners	Owners of lands associated with shorebird use, including agricultural lands (particularly rice fields), aquaculture (e.g., crayfish, shrimp), salt ponds, and rangelands. These landscapes have potential to affect or create shorebird habitat if managed appropriately.
Local Non-profit Groups	Non-profits that are engaged in shorebird conservation locally (sometimes connected to international initiatives). Examples from major flyways include Corbidi (Peru, Pacific Flyway), Coastal Bend Bays & Estuaries Coastal Bird Program (Texas, USA, Midcontinent), SAVE Brasil (Atlantic), and many more.
Multi-level	
Public Land Managers	Managers of protected areas can operate at local, state, or even regional and transnational levels. Protected areas are sometimes created specifically for birds, sometimes mixed recreational or human use, and sometimes strictly protected.
Natural Resources Conservation Service (NRCS)	U.S. Department of Agriculture branch that provides technical and financial assistance to farmers, ranchers, and other private landowners and managers. NRCS hosts a variety of conservation programs, typically delivered at the county-level, intended to help landowners and other partners protect, conserve, and sustainably manage their private lands. Many NRCS programs have potential to support shorebirds and/or create habitat if managed and targeted appropriately.
U.S. Fish and Wildlife Service Partners for Fish and Wildlife Program	Provides free technical and financial assistance to landowners, managers, tribes, corporations, schools and nonprofits interested in improving wildlife habitat on their lands. Projects are voluntary and customized to meet landowners' needs. Participating landowners continue to own and manage their land while they improve conditions for wildlife. Staff plan, design, supervise and monitor customized habitat restoration projects with a focus on habitats of concern like wetlands and native grasslands. Every U.S. State and Territory has a coordinator who helps deliver the program.
Land Trusts	Non-governmental organizations that place easements on or take ownership of private lands at the request of the landowner. Private lands with wetland, grassland, or working agricultural habitats have potential to support shorebirds and/or create habitat if managed appropriately. They can operate at local, regional, and national levels.
Non-profit / Non-governmental Conservation Organizations	Any non-profit / non-governmental program or organization not mentioned above that may operate at multiple levels from local to hemispheric. Examples include Birds Canada, Birds Caribbean, BirdLife International, Ducks Unlimited, National Audubon Society, Point Blue, Wetlands International, Wildlife Conservation Society, and more. Through programs to create or manage wetland, grassland, or agricultural habitat, these groups are (or could be) providing shorebird habitat.

806
807
808

809   Table 3. Description and sample size of data types submitted to the Shorebird Science and Conservation Collective as of June 2024.

TECHNOLOGY	DESCRIPTION	DATA FORMAT**	SPATIAL ACCURACY	SAMPLING FREQUENCY	MONITORING DURATION	NUMBER OF INDIVIDUALS CONTRIBUTED (% of individuals)	NUMBER OF SPECIES CONTRIBUTED (% of species)
GPS	Electronic tags attached to individual birds. Locations estimated from GPS satellites and obtained either by retrieving the device and downloading stored data (i.e., archival tags) or transmitting the data to satellites.	Latitude and longitude coordinates	Typically 10-100 m (Noonan et al., 2019)	Minutes to days or pre-programmed schedules such as every 2 weeks during overwintering and 2 days during migration depending on study goal and tag battery life	Months to years depending on tag type and attachment method	1,213* (35.5 %)	16 (44.4 %)
Argos Doppler Shift	Electronic satellite tags (called Platform Terminal Transmitters or PTTs) attached to individual birds. Tags send signals to orbiting satellites that re-transmit location data back to a receiving station on Earth for processing.	Latitude and longitude coordinates with "location class" codes or error ellipses that indicate the accuracy of the estimated location	Generally 250—1,500 m (CLS America, 2016) but individual locations can have errors larger than 25km (Jonsen et al., 2020)	Minutes to days, often irregular; tags may use pre-programmed on/off schedules to allow for charging (solar models), or to conserve energy (battery models)	1+ years depending on tag type and attachment method	938* (27.4 %)	15 (41.7%)
Motus	Electronic VHF radio tags attached to individual birds that are detected by a	Raw detection data (e.g., timestamp, signal strength)	Typically 15 – 20 km from receiving stations (Taylor et al., 2017) but up to	3 - 30 seconds when tag nears receiving station, generally, though	Days to months	1,091 (31.9 %)	4 (11.1%)

	network of receiving stations. Detection data are stored on receivers, synched over the internet, or sent via satellite or cellular network.	associated with the location of each receiving station	180 km (Anderson et al., 2019); ~500 m if omnidirectional antennas are used (Taylor et al., 2017)	gaps can be minutes to days			
Light-level Geolocator	Electronic devices attached to individual birds that record ambient light levels, which can be related to global light patterns to estimate the animal's location. Devices must be retrieved to download the data.	Raw ambient light levels which require substantial processing and analysis to estimate locations (Lisovski et al., 2020)	50 - 200 km with much greater error in latitude during spring and fall equinox (Lisovski et al., 2012; Rakhimberdiev et al., 2016)	1-2 locations per day	1+ years	362 (10.6 %)	15 (41.7 %)
Resight	Observations of individual birds marked with unique color band / flag combinations, or alphanumeric bands or flags	Latitude and longitude coordinates provided by the observer	Depends on observer proximity but usually < 100 m.	Limited to where and when observers search for birds, marker retention, and animal lifespan	Limited to observer effort, marker retention, and animal lifespan	Submitted data currently not integrated into the Collective dataset	15 (41.7%)

* Includes 184 individuals tracked with both GPS and Argos technologies
** Tags also collected barometric pressure (0.5% of individuals, 1 species), external temperature (1.4%, 5 species), altitude (12.3%, 5 species, generally of
limited accuracy), acceleration (0.7%, 1 species), magnetic field values (0.6%, 1 species), gyroscope data (0.5% of individuals, 1 species), an indication that the
tag was wet or dry (0.2%, 1 species), or that the tag became immobile (17.3%, 4 species).

Figure 1. The process by which the Shorebird Science and Conservation Collective distributes

knowledge generated from electronic tracking studies of shorebirds to conservation partners

across the Americas to enhance land and species conservation, management, and outreach and

education activities. In the future, this resource also has potential to inform conservation policy.

The Collective engages with practitioners in service to community-driven management and

conservation needs, with the ultimate goal of reversing population declines.

Figure 2. Distribution of Argos and GPS tracking data from tags deployed in the Americas

(Table 3; n=1,883 individuals of 22 species) contributed to the Shorebird Science and

Conservation Collective as of June 2024, shaded by the number of species with tracking

locations in 100 km x100 km hexagonal cells.

Figure 3. Example figure from a conservation project report informing the timing of freshwater

deployments to provide habitat for migratory and wintering shorebirds in Texas. A) Example

tracking data from whimbrel (*Numenius phaeopus)*, shared by Jennie Rausch, Environment and

Climate Change Canada (middle) and overlaid on a focal area of interest in Texas (right). B)

Time series of when tracked birds traveled through the Texas Gulf Coast (black histogram bars)

together with median weekly relative abundance (gray lines) in the area of interest predicted by

eBird community science data (Fink et al., 2022). Individuals tracked in more than one year

contributed to histogram counts more than once because timing varied for individuals between

years. Fall migration is shaded orange and spring migration is shaded green. Species with

asterisks are Texas State Species of Greatest Conservation Need (Texas Parks and Wildlife

Department, 2012). Information from A) and B) were used to determine time periods during

which water deployments could benefit peak diversity and abundances of migratory shorebirds,

leading to C) Recommendation to conservation partners for deploying water to benefit migrating

and overwintering shorebirds under recommended and alternate scenarios.