1

LukProt: A database of eukaryotic predicted proteins designed for investigations of animal origins

Łukasz F. Sobala*[1]

[1]Laboratory of Glycobiology, Hirszfeld Institute of Immunology and Experimental Therapy, PAS – Wrocław, Poland

*Corresponding author

Correspondence: lukasz.sobala@hirszfeld.pl

ABSTRACT

The origins and early evolution of animals is a subject with many outstanding questions. One problem faced by researchers trying to answer them is the absence of a comprehensive database of sequences from non-bilaterians. Publicly available data is plentiful but scattered and often not associated with proper metadata. A new database presented in this paper, LukProt, is an attempt at solving this issue. The database contains protein sequences obtained mostly from genomic, transcriptomic and metagenomic studies and is an extension of EukProt (a collection of eukaryotic sequences from all known clades). LukProt adopts and extends EukProt naming conventions and includes data from 216 additional animals. The database is associated with a taxogroup scheme suitable for studying early animal evolution. Minor updates to the database will contain species additions or metadata corrections and major updates will synchronize LukProt to each new version of EukProt. All public LukProt releases are permanently stored on Zenodo. A BLAST server to search the database is available at https://lukprot.hirszfeld.pl/. The community is invited to participate

in maintaining and correcting LukProt. As it can be searched without downloading locally, the database can be a convenient resource not only for evolutionary biologists, but the broader scientific community as well.

Keywords: EukProt, LukProt, database, evolution, animals, Metazoa, eukaryote, proteome, dataset

                                                    Introduction

Animals (Metazoa) are a diverse group of organisms whose deep evolutionary history has been a
subject of a long debate (Ruiz-Trillo et al., 2023). Many early branching lineages of animals are extinct, for
example much of the Ediacaran fauna is not similar to any living animals. Animal tissues were not often
amenable to fossilization pre-Ediacaran (Murdock & Donoghue, 2011). Recent advancements in
phylogenomics helped resolve fundamental questions in early history of Metazoa, for example the
identity of the earliest extant animal branch (Dunn et al., 2008; Schultz et al., 2023) and the timing of the
two whole genome duplications which occurred in the ancestors of all living jawed vertebrates (Simakov
et al., 2022; Marlétaz et al., 2024). These developments were enabled by more advanced bioinformatic
tools, broader access to processing power and broader availability of sequencing data.
As animals are members of Eukaryota, knowledge about eukaryotic evolution can provide context for
evolution of animals. In the last decade, the topology of the eukarotic tree of life has been undergoing
revisions as major as the animal tree. The discovery and characterization of Asgard archaea (Zaremba-
Niedzwiedzka et al., 2017; Eme et al., 2023) helped place the trunk of the tree. Due to the recent
sequencing boom, new supergroups of protists are still being defined (Tikhonenkov et al., 2022). A
comprehensive database which collects predicted proteomes of eukaryotes, EukProt, plays a role in these
developments (Richter et al., 2022). It is integrated with UniEuk, a recently created reference taxonomy
of eukaryotes (Berney et al., 2017). The broad taxon sampling and careful curation of EukProt makes it
useful for many applications. The EukProt database, however, compromises on the number of included
species in the clades with a high number of species sequenced (plants, fungi or animals).
Investigating animal evolution requires broad taxon sampling of early diverging animal clades, which
EukProt, NCBI or UniProt databases do not provide. One attempt at a more even animal taxon sampling,
the Animal Proteome Database – AniProtDB (Barreira et al., 2021), only encompasses data from 100
species. Another large database of non-vertebrate animal sequences focuses on protostomes (Fernández
et al., 2022). A more comprehensive database, which includes data from early diverging lineages, could
serve as a tool for researchers to study the origins and evolution of uniquely animal characters, such as
obligatory multicellularity or neurons.
To fill this gap, I have created LukProt. It is a synthesis of the latest available EukProt database, the
AniProtDB and a significant number of additional collected sequences. The database focuses on
holozoans (a clade of opisthokonts sister to Nucletmycea) and animal species from the clades:
Ctenophora, Porifera, Placozoa and Cnidaria. Bilaterian sampling is also increased. The LukProt database
follows the conventions established by EukProt: each dataset has a unique identifier and each protein has
a persistent ID composed of the dataset identifier, the species/strain name and protein number. In case a
superior dataset from the same species replaces the previous one, the previous ID is retired in a new
version of the database. A more detailed characterization of LukProt, as well as EukProt data or metadata
modifications made for LukProt, is presented below.

                                                     Methods

Dataset naming and species selection
In most cases, species and strain names from EukProt were conserved. To aid further analyses,
extremely long species/strain names were shortened so that no species+strain name is longer than 32
characters (including underscores). In these cases, the EukProt dataset identifier and protein ID were not
changed. EukProt version 3 is used in the current LukProt revision (v1.5.1.rev1).
Newly downloaded datasets were assigned names after reading the associated publications, if any
referred to the name. Otherwise, names were adapted from the websites from which the datasets were

downloaded. As in EukProt IDs, spaces are replaced with underscores in individual FASTA files and
additionally periods were removed from the names. After the protein ID, the sequence identifier from the
original file is included.

Distribution of the database
LukProt is available as an archive containing FASTA files placed inside a directory tree which
recapitulates the taxonomic relationships between the species. Additionally, an archive with BLAST+ v5
database files (one per species, one per taxogroup and one full database) is available to download.
Miscellaneous files, such as the coloring scheme, the changelog, the metadata spreadsheet and helper
scripts, are available in the associated Zenodo repository. Datasets that were not included or
untranslated transcriptomes are not provided. The repository is designed to be easy to use and the
provided files can be implemented in local workflows with minimal user modification required.

Data processing – the LukProt database
The data processing steps are detailed in the associated metadata file. Briefly, unassembled
transcriptomes were assembled using Trinity v2.12 to v2.15.1 (Grabherr et al., 2011; Haas et al., 2013) or
TransAbyss v2.0.1 (Robertson et al., 2010). Trinity version is indicated in all cases (the software was
updated after some datasets were created). TransAbyss was used in only 1 case where Trinity could not
complete the run due to an error. Proteins were predicted using TransDecoder.LongOrfs (Haas, BJ.
https://github.com/TransDecoder/TransDecoder, versions v5.5.0 or v5.7.0, indicated in metadata) with
default options (minimum protein length: 100). Predicted peptides were then clustered (in most cases)
using cd-hit v4.8.1 (Fu et al., 2012) using the command: cd-hit -s 0.5 -g 1 -d 0 -T 6 -M 16000 -c 0.99 -i
input.fa -o output.fa. Strain merging was done by concatenating the protein FASTA files to a single file
and running cd-hit as above. To extract protein sequences from a genome, gffread v0.12.7 (Pertea &
Pertea, 2020) was run using the following parameters: gffread -g genome.fasta annotation.gff -y
output.fa -C -J –no-pseudo. The R package taxonomizr (Sherrill-Mix, S.
https://github.com/sherrillmix/taxonomizr v0.10.6) and manual curation was used to find the NCBI
taxonomy IDs of each species. For dataset quality analyses, BUSCO 5.7.1 (Manni et al., 2021) was used
with dataset "eukaryota_odb10" (255 BUSCOs) created on 2024-01-08. BUSCO utilized hmmsearch
version 3.1 (http://hmmer.org/). The OMArk (Nevers et al., 2024) version was 2.0.3 and the OMAmer
version was 0.3.0, both were used with database LUCA.h5 (Nov 2022 release). The NCBI taxdb used in the
OMArk analysis was downloaded on 2024-05-12. BUSCO, OMAmer and OMArk were run with default
parameters, with the species NCBI taxid specified for OMArk where it was available.
In a number of cases, the dataset names were shortened with respect to EukProt, without changing
the dataset ID or protein ID number. The mapping file is available in the Zenodo repository. Additional
processing steps that were done to data from EukProt were: removing any asterisk (*) characters and line
breaks from sequences. This was done to broaden compatibility and reduce file size. Sequence
manipulation was done using SeqKit (Shen et al., 2016) and GNU coreutils.

Data processing software – 20 BUSCOs example
Firstly, 20 most common single BUSCOs in the LukProt database were selected. Sequences of these 20
BUSCOs were collected into a separate FASTA file for each species. Species which possessed fewer than
10 of these BUSCOs were filtered out from subsequent analyses. BUSCO sequences from the remaining
species were then concatenated, merged into a single multi-FASTA file and aligned. Protein sequence
alignment was performed using FAMSA v2.2.2- (Deorowicz et al., 2016) (-refine-mode on). The alignment
was trimmed using trimAl v1.4.rev15 (Capella-Gutierrez et al., 2009) with a gap threshold of 50%.
Phylogenies were inferred using IQ-TREE (2.3.3). IQ-TREE branch support values were calculates using SH-

aLRT (Shimodaira–Hasegawa-like approximate likelihood ratio test, 5000 replicates) (Guindon et al.,
2010), aBayes (approximate Bayes) (Anisimova et al., 2011) and UFBoot2 (Ultrafast bootstrap 2, 5000
replicates) (Hoang et al., 2018) algorithms. IQ-TREE model was selected using ModelFinder
(Kalyaanamoorthy et al., 2017). FigTree 1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/) was used for
phylogeny visualization and editing.

129   **Results and Discussion**

General purpose of LukProt
The purpose of the database release is to create a resource which one could consult on whether a
given domain/protein/protein family exists within a clade of eukaryotes, with a focus of animals and their
closest relatives, holozoans. To this end, individual dataset quality is of lower importance. No
decontamination procedures were applied to any datasets and none were excluded because of possible
contamination. Contamination was, in fact, used to extract a new dataset – LP00177 – containing
choanoflagellate-derived sequences found in EP00436 from EukProt v2 (superseded by EP00913 in
EukProt v3). Many of the LukProt transcriptome-derived proteomic datasets from cnidarians and sponges
are in fact holobiont transcriptomes which include bacterial transcripts and those from other associated
organisms. It is assumed that phylogeny reconstructions will be the guide when drawing conclusions from
database searches. Phylogeny-guided analysis, using a suitable sister database, can also be the basis for
studies of lateral gene transfers.
The purpose of scripts associated with the database is to accelerate phylogeny inference and
interpretation of the results. The documentation, which includes usage examples, should guide a less
experienced user and let them perform independent investigations.

Characterization of the LukProt database
The current version of the database (v1.5.1.rev1) contains 1281 datasets, with 990 coming from
EukProt, 41 from AniProtDB and 250 newly added. One dataset represents a proteome from a single
species or strain. The newly added datasets have been collected from numerous studies and publicly
available repositories, or generated from publicly available data using *in silico* methods. Details of each
dataset generation can be found in the supporting metadata. Overall database statistics are given in
Table 1. Taxon sampling of the early branching animal clades, especially ctenophores, sponges and
cnidarians, is improved in comparison to EukProt. For example, placozoans account for six datasets and
this constitutes 26% of all currently known, potential placozoan species (23) (Tessler et al., 2022).

Table 1 – Comparison of the number of taxa in selected clades of LukProt and other databases.

Taxogroup	EukProt v3 – included in the stock database	AniProtDB within LukProt	LukProt v1.5.1.rev1 added (+)/excluded (-)	LukProt v1.5.1.rev1 total
Eukaryota	993	41	+250/-3	1281
Diaphoretickes	632	0	+28/-0	660
Holozoa (excluding Metazoa)	105 (40)	41 (0)	+221/-2 (+5/-1)	365 (44)
Metazoa (excluding Bilateria)	65 (14)	41 (3)	+216/-1 (+162/-0)	321 (179)
Ctenophora	2	1	+35/-0	38
Porifera	5	1	+41/-0	47
Placozoa	2	0	+4/-0	6
Cnidaria	5	1	+82/-0	88
Bilateria	51	38	+54/-1	142

A consensus taxonomy

Included in the database is a consensus taxonomy of eukaryotes. To aid analyses of animal evolution,
some UniEuk (Berney et al., 2017) taxogroups outside Podiata were merged, and Podiata was split into
more granular groups (mostly monophyletic). This treatment should make deep animal phylogenetic
analyses more accessible. Clades are explained in Supplementary Information and the full eukaryotic tree
can be found in Figure 1. Each internal node was assigned a unique name, following UniEuk or recent
literature where possible. A number of unnamed clades were assigned provisional names (Figure 1). The
LukProt taxonomic framework is necessarily different from UniEuk but will be adjusted following the
updates that are implemented in the latter and in future publications, as well as community suggestions.
It encompasses 81 groups of organisms, most of which are monophyletic. Based on this taxonomy, 59
groups to limit the number of species were created to aid analyses focused on narrower clades. These
groups can be found in the database metadata.

Figure 1 – The eukaryotic tree of life used in LukProt. Each group is assigned a unique color which makes visual interpretation of phylogenies easier. Groups with blue dots are paraphyletic. Internal node names written in red were invented for the database.

Source data traceability

Each dataset is associated with a URL and a DOI for the source article – if it exists – following the EukProt conventions. Considerable effort was made to apply the traceability guidelines set by EukProt to the metadata included with LukProt. As an improvement over EukProt, the metadata file now includes the NCBI taxids for the species for which they were possible to find (including EukProt-derived species).

Release schedule

Version 1.4.1 of LukProt of was made available on Zenodo in 2022 and was downloaded 38 times (as of 2024-05-21). It was based on EukProt v2. Since then, many additions and improvements were introduced, which resulted in the publication-ready version 1.5.1.rev1, based on EukProt v3. LukProt will receive major updates to trace each new release of the EukProt database; it should be considered its

extension. Minor updates will be released as needed. The release version number will be incremented up
only if the underlying sequence files are changed. If only metadata changes, a "revision" will be
introduced. In case of EukProt including any of the LukProt datasets, the LukProt ID will be retired and the
EukProt ID will be used instead. Future releases will include the improvements in EukProt and contain all
the additional data provided by LukProt. Changes will be documented and explained in readable form in
the changelog. All public versions of the database are archived in the Zenodo repository and a unique DOI
is generated for each new version (as well as a DOI which always resolves to the newest version).

BLAST server and convenience scripts
A publicly available BLAST+ (Camacho et al., 2009) server for LukProt is available at
https://lukprot.hirszfeld.pl/ using SequenceServer 3.0.1 (Priyam et al., 2019) and NCBI BLAST 2.15.0+ as
backends. The taxonomy in Figure 1 is applied to the database and broad or narrow clades can be readily
selected from the tree (or alphabetically). In comparison, currently the EukProt BLAST server
(https://evocellbio.com/SAGdb/EukProt/) only provides an alphabetical list of taxogroups, without tree of
life topology. The LukProt BLAST server will be updated as new versions of the database or the underlying
software are released.
The functions of the executable scripts that accompany the database include, but are not limited to:
extracting sequences from preselected groups of organisms (for example: Diaphoretickes, Holozoa
excluding Metazoa), extracting LukProt sequences from the full BLAST database or using a list of
identifiers and tree tip recoloring. The scripts are documented with usage examples in the Zenodo
repository of the database.

'LukProt Comparative Set' (LCS): the EukProt TCS extended for LukProt
By analogy to 'The Comparative Set' of EukProt – a reduced set of species of high sequencing quality
and phylogenetic importance – the LCS is proposed. The set consists of 233 datasets, a similar proportion
(18%) of the number of selected species for TCS versus the full EukProt database (19%). The additional
species selected reflect the clade upon which LukProt focuses – the Holozoa. It is available to search using
the BLAST server. As of LukProt v1.5.1.rev1 the LCS should be considered a beta version and is subject to
change following community suggestions. An included script supports extracting LCS sequences from any
multi-FASTA file.

A sample of LukProt data: top single BUSCOs analysis
One example of the utility of the database can be seen in my previous work, where an earlier version
of the database (v1.4.1) was used (Sobala, 2023). The example below is a phylogeny reconstructed from
20 most often found single eukaryotic BUSCOs in the database. The sequences were concatenated and
aligned; details can be found in the methods and the Supplementary Data. Only initial species filtering
was conducted. No post-tree reconstruction sequence removal was done and the example should
represent a fairly unfiltered view of data one can expect to find in LukProt for the selected clade, Filozoa.

Figure 2 – (**A**) Consensus IQ-TREE phylogeny reconstruction of the 20 most prevalent LukProt BUSCOs in Filozoans. UFBoot2 branch support values are indicated. LBA – long branch attraction artifact. (**B**) Reference, colored cladogram of the group Filozoa. Calcarea is shown in brackets because of its absence from the main phylogeny.

The obtained phylogeny (Figure 2A) mostly agrees with the current understanding of filozoan/early animal evolution (Figure 2B). The consensus phylogeny (.contree) was selected for the figure due to a slightly better log likelihood than the main output tree (.treefile). Ctenophores were recovered as the earliest branching animal clade with full support. Most sponge sequences formed a single clade, as well as the cnidarian and bilaterian sequences. However, the long the branch attraction phenomenon (Susko & Roger, 2021) caused formation of a branch with animals from clades that are known to have highly divergent sequences (Myxozoa, Xenacoelomorpha, Nematoda, Tardigrada, Platyhelminthes). What is more, the somewhat controversial clade Placozoa + Cnidaria (here called "Placnidia") was not recovered, but the Placozoa-out UFBoot2 support value was low (69%).

For targeted evolutionary analyses, sequence pruning using user-selected criteria might improve tree topology. Other examples of downstream analysis of LukProt data include: inferring single gene trees, single domain trees and gene family presence analysis. The concatenated and trimmed alignment used in the example contained 299 species and 5428 sites (5229 parsimony informative) which means that the number of informative sites was over 17x the number of sequences. In cases of short sequences, the number of sequences found in LukProt may be too high for maximum likelihood algorithms, resulting in overparametrized trees. To deal with this problem, alternative approaches, such as sequence similarity networks (SSNpipe – https://github.com/ahvdk/SSNpipe), cd-hit or MMSEQS2 (Mirdita et al., 2019) clustering, can be helpful. Depending on the question asked, database users may need to find other ways of reducing the complexity of the problem.

Limitations of the LukProt database and future plans

As an extension of EukProt, the limitations of the parent database apply to LukProt as well. One further limitation results from an editorial decision not to predict proteins shorter than 100 amino acids (50 in EukProt). The reduced "comparative set" was introduced as a beta version and the community is invited to comment on the species selection and suggest changes. A number of NCBI taxids are missing; these can be added in future database revisions.

Acknowledgements

I would like to thank the creators and maintainers of the EukProt database as well as the teams and individuals who generated the source data. Members of the Multicellgenome Lab (IBE-CSIC, Barcelona): Marta Álvarez, Elena Casacuberta, Michelle Leger, Koryu Kin and Iñaki Ruiz-Trillo, are thanked for helpful discussions and the permission to release the protein dataset from *Corallochytrium limacisporum* (strain India). Michelle Leger and Eduard Ocaña-Pallarès are acknowledged for suggestions about the taxonomic framework. I also acknowledge Marcin Czerwiński for helpful comments about the manuscript.

Data, scripts, code, and supplementary information availability

The LukProt database and associated files are available on Zenodo: https://doi.org/10.5281/zenodo.7089120 (Sobala, 2022).

Supplementary information is available online: https://doi.org/10.5281/zenodo.11324807 (Sobala, 2024). Users can contact the corresponding author by email if any additional questions, or correction suggestions, arise.

Conflict of interest disclosure

The author declares no financial conflicts of interest in relation to the content of the article.

Funding

This work was supported by National Science Centre of Poland (NCN) as part of the grant SONATINA no. 2020/36/C/NZ8/00081 entitled "The role of glycosylation in the emergence of animal multicellularity".

References

Anisimova M, Gil M, Dufayard J-F, Dessimoz C, Gascuel O (2011) Survey of Branch Support Methods Demonstrates Accuracy, Power, and Robustness of Fast Likelihood-based Approximation Schemes. *Systematic Biology*, **60**, 685–699. https://doi.org/10.1093/sysbio/syr041

Barreira SN, Nguyen A-D, Fredriksen MT, Wolfsberg TG, Moreland RT, Baxevanis AD (2021) AniProtDB: A Collection of Consistently Generated Metazoan Proteomes for Comparative Genomics Studies. *Molecular Biology and Evolution*, **38**, 4628–4633. https://doi.org/10.1093/molbev/msab165

Berney C, Ciuprina A, Bender S, Brodie J, Edgcomb V, Kim E, Rajan J, Parfrey LW, Adl S, Audic S, Bass D, Caron DA, Cochrane G, Czech L, Dunthorn M, Geisen S, Glöckner FO, Mahé F, Quast C, Kaye JZ, Simpson AGB, Stamatakis A, Campo J del, Yilmaz P, Vargas C de (2017) UniEuk: Time to Speak a Common Language in Protistology! *Journal of Eukaryotic Microbiology*, **64**, 407–411. https://doi.org/10.1111/jeu.12414

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL (2009) BLAST+: architecture and applications. *BMC Bioinformatics*, **10**, 421. https://doi.org/10.1186/1471-2105-10-421

Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T (2009) trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, **25**, 1972–1973. https://doi.org/10.1093/bioinformatics/btp348

Deorowicz S, Debudaj-Grabysz A, Gudyś A (2016) FAMSA: Fast and accurate multiple sequence alignment of huge protein families. *Scientific Reports*, **6**, 33964. https://doi.org/10.1038/srep33964

Dunn CW, Hejnol A, Matus DQ, Pang K, Browne WE, Smith SA, Seaver E, Rouse GW, Obst M, Edgecombe GD, Sørensen MV, Haddock SHD, Schmidt-Rhaesa A, Okusu A, Kristensen RM, Wheeler WC, Martindale MQ, Giribet G (2008) Broad phylogenomic sampling improves resolution of the animal tree of life. *Nature*, **452**, 745–749. https://doi.org/10.1038/nature06614

Eme L, Tamarit D, Caceres EF, Stairs CW, De Anda V, Schön ME, Seitz KW, Dombrowski N, Lewis WH, Homa F, Saw JH, Lombard J, Nunoura T, Li W-J, Hua Z-S, Chen L-X, Banfield JF, John ES, Reysenbach A-L, Stott MB, Schramm A, Kjeldsen KU, Teske AP, Baker BJ, Ettema TJG (2023) Inference and reconstruction of the heimdallarchaeial ancestry of eukaryotes. *Nature*, 1–8. https://doi.org/10.1038/s41586-023-06186-2

Fernández R, Tonzo V, Simón Guerrero C, Lozano-Fernandez J, Martínez-Redondo GI, Balart-García P, Aristide L, Eleftheriadi K, Vargas-Chávez C (2022) MATEdb, a data repository of high-quality metazoan transcriptome assemblies to accelerate phylogenomic studies. *Peer Community Journal*, **2**. https://doi.org/10.24072/pcjournal.177

Fu L, Niu B, Zhu Z, Wu S, Li W (2012) CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics*, **28**, 3150–3152. https://doi.org/10.1093/bioinformatics/bts565

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A, Rhind N, di Palma F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A (2011) Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology*, **29**, 644–652. https://doi.org/10.1038/nbt.1883

Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New Algorithms and
Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the Performance of PhyML 3.0.
Systematic Biology, **59**, 307–321. https://doi.org/10.1093/sysbio/syq010

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li B, Lieber
M, MacManes MD, Ott M, Orvis J, Pochet N, Strozzi F, Weeks N, Westerman R, William T, Dewey CN,
Henschel R, LeDuc RD, Friedman N, Regev A (2013) De novo transcript sequence reconstruction from
RNA-seq using the Trinity platform for reference generation and analysis. *Nature Protocols*, **8**, 1494–
1512. https://doi.org/10.1038/nprot.2013.084

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS (2018) UFBoot2: Improving the Ultrafast
Bootstrap Approximation. *Molecular Biology and Evolution*, **35**, 518–522.
https://doi.org/10.1093/molbev/msx281

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017) ModelFinder: fast model
selection for accurate phylogenetic estimates. *Nature Methods*, **14**, 587–589.
https://doi.org/10.1038/nmeth.4285

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM (2021) BUSCO Update: Novel and Streamlined
Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic,
Prokaryotic, and Viral Genomes. *Molecular Biology and Evolution*, **38**, 4647–4654.
https://doi.org/10.1093/molbev/msab199

Marlétaz F, Timoshevskaya N, Timoshevskiy VA, Parey E, Simakov O, Gavriouchkina D, Suzuki M,
Kubokawa K, Brenner S, Smith JJ, Rokhsar DS (2024) The hagfish genome and the evolution of
vertebrates. *Nature*, 1–3. https://doi.org/10.1038/s41586-024-07070-3

Mirdita M, Steinegger M, Söding J (2019) MMseqs2 desktop and local web server app for fast, interactive
sequence searches. *Bioinformatics*, **35**, 2856–2858. https://doi.org/10.1093/bioinformatics/bty1057

Murdock DJE, Donoghue PCJ (2011) Evolutionary Origins of Animal Skeletal Biomineralization. *Cells
Tissues Organs*, **194**, 98–102. https://doi.org/10.1159/000324245

Nevers Y, Warwick Vesztrocy A, Rossier V, Train C-M, Altenhoff A, Dessimoz C, Glover NM (2024) Quality
assessment of gene repertoire annotations with OMArk. *Nature Biotechnology*, 1–10.
https://doi.org/10.1038/s41587-024-02147-w

Pertea G, Pertea M (2020) GFF Utilities: GffRead and GffCompare. *F1000Research*, **9:304**.
https://doi.org/10.12688/f1000research.23297.2

Piñeiro C, Abuín JM, Pichel JC (2020) Very Fast Tree: speeding up the estimation of phylogenies for large
alignments through parallelization and vectorization strategies. *Bioinformatics*, **36**, 4658–4659.
https://doi.org/10.1093/bioinformatics/btaa582

Priyam A, Woodcroft BJ, Rai V, Moghul I, Munagala A, Ter F, Chowdhary H, Pieniak I, Maynard LJ, Gibbins
MA, Moon H, Davis-Richardson A, Uludag M, Watson-Haigh NS, Challis R, Nakamura H, Favreau E,
Gómez EA, Pluskal T, Leonard G, Rumpf W, Wurm Y (2019) Sequenceserver: A Modern Graphical User
Interface for Custom BLAST Databases. *Molecular Biology and Evolution*, **36**, 2922–2924.
https://doi.org/10.1093/molbev/msz185

Richter DJ, Berney C, Strassert JFH, Poh Y-P, Herman EK, Muñoz-Gómez SA, Wideman JG, Burki F, de
Vargas C (2022) EukProt: A database of genome-scale predicted proteins across the diversity of
eukaryotes. *Peer Community Journal*, **2**. https://doi.org/10.24072/pcjournal.173

Robertson G, Schein J, Chiu R, Corbett R, Field M, Jackman SD, Mungall K, Lee S, Okada HM, Qian JQ,
Griffith M, Raymond A, Thiessen N, Cezard T, Butterfield YS, Newsome R, Chan SK, She R, Varhol R,
Kamoh B, Prabhu A-L, Tam A, Zhao Y, Moore RA, Hirst M, Marra MA, Jones SJM, Hoodless PA, Birol I
(2010) De novo assembly and analysis of RNA-seq data. *Nature Methods*, **7**, 909–912.
https://doi.org/10.1038/nmeth.1517

Ruiz-Trillo I, Kin K, Casacuberta E (2023) The Origin of Metazoan Multicellularity: A Potential Microbial
Black Swan Event. *Annual Review of Microbiology*, **77**, 499–516. https://doi.org/10.1146/annurev-
micro-032421-120023
Schultz DT, Haddock SHD, Bredeson JV, Green RE, Simakov O, Rokhsar DS (2023) Ancient gene linkages
support ctenophores as sister to other animals. *Nature*, **618**, 110–117.
https://doi.org/10.1038/s41586-023-05936-6
Shen W, Le S, Li Y, Hu F (2016) SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File
Manipulation. *PLOS ONE*, **11**, e0163962. https://doi.org/10.1371/journal.pone.0163962
Simakov O, Bredeson J, Berkoff K, Marletaz F, Mitros T, Schultz DT, O'Connell BL, Dear P, Martinez DE,
Steele RE, Green RE, David CN, Rokhsar DS (2022) Deeply conserved synteny and the evolution of
metazoan chromosomes. *Science Advances*, **8**, eabi5884. https://doi.org/10.1126/sciadv.abi5884
Sobala ŁF (2022) LukProt – an animal evolution-centric eukaryotic protein database.
https://doi.org/10.5281/zenodo.7089120
Sobala ŁF (2023) Evolution and phylogenetic distribution of endo-α-mannosidase. *Glycobiology*, **33**, 687–
699. https://doi.org/10.1093/glycob/cwad041
Sobala ŁF (2024) Supplementary Data associated with the article "LukProt: A database of eukaryotic
predicted proteins designed for investigations of animal origins."
https://doi.org/10.5281/zenodo.11324807
Susko E, Roger AJ (2021) Long Branch Attraction Biases in Phylogenetics. *Systematic Biology*, **70**, 838–843.
https://doi.org/10.1093/sysbio/syab001
Tessler M, Neumann JS, Kamm K, Osigus H-J, Eshel G, Narechania A, Burns JA, DeSalle R, Schierwater B
(2022) Phylogenomics and the first higher taxonomy of Placozoa, an ancient and enigmatic animal
phylum. *Frontiers in Ecology and Evolution*, **10**. https://doi.org/10.3389/fevo.2022.1016357
Tikhonenkov DV, Mikhailov KV, Gawryluk RMR, Belyaev AO, Mathur V, Karpov SA, Zagumyonnyi DG,
Borodina AS, Prokina KI, Mylnikov AP, Aleoshin VV, Keeling PJ (2022) Microbial predators form a new
supergroup of eukaryotes. *Nature*, **612**, 714–719. https://doi.org/10.1038/s41586-022-05511-5
Zaremba-Niedzwiedzka K, Caceres EF, Saw JH, Bäckström D, Juzokaite L, Vancaester E, Seitz KW,
Anantharaman K, Starnawski P, Kjeldsen KU, Stott MB, Nunoura T, Banfield JF, Schramm A, Baker BJ,
Spang A, Ettema TJG (2017) Asgard archaea illuminate the origin of eukaryotic cellular complexity.
Nature, **541**, 353–358. https://doi.org/10.1038/nature21031