Gastric microbiomes of *Cassiopea* in the Florida Keys are low diversity and *Endozoicomonas-*,

Vibrio- and *Mycoplasma-dominated*

Kaden M Muffett[1]*, Jessica M Labonté[2], Maria Pia Miglietta[2]

1   University of California Merced, Merced, CA, 95343

2   Texas A&M University at Galveston, Galveston, TX, 77554

*Corresponding email: kmmuffett@gmail.com

Running Page Head: Bacteria of *Cassiopea* in the Florida Keys

Abstract

Interactions with microbial communities fundamentally shape metazoans' physiology,

development, and health across marine ecosystems. This is especially true in zooxanthellate

(symbiotic algae-containing) cnidarians. In photosymbiotic anthozoans (eg. shallow water

anemones and corals), the key members of the associated microbiota are increasingly well

studied, however there is limited data on photosymbiotic scyphozoans (true jellyfish). Using 16S

rRNA barcoding, we sampled the internal and external mucus of the zooxanthellate Upside-

Down Jellyfish, *Cassiopea xamachana* throughout eight sites covering the full length of the

Florida Keys. We find that across sites, these medusae to have low-diversity internal microbial

communities distinct from the communities of their external surfaces and their environment.

These internal communities are dominated by only three taxa: *Endozoicomonas* cf. *atrinae*, an

uncultured novel *Mycoplasma*, and *Vibrio* cf. *coralliilyticus*. *Cassiopea* bell mucosal samples

conform largely to the communities of surrounding sediment with the addition of

Endozoicomonas cf. *atrinae*. The microbial taxa we identify associated with wild Florida Keys

Cassiopea bear a strong resemblance to those found within photosymbiotic anthozoans,

increasing the known links in ecological position between these groups.

Key words: Cnidaria, microbiome, symbiosis, *Cassiopea*, Scyphozoa

Introduction

Cnidarian-photosymbiont interactions are central to ecosystem function in coral reefs and beyond. Algal photosymbionts, or zooxanthellae, are present in the well-known stony corals, octocorals, and anemones, but also play important roles in a variety of lesser-known jellyfishes, chief among them, jellyfish of the genus *Cassiopea* (1–4). For symbiotic anthozoans, microbial communities play a key role in maintaining health within this photosymbiont partnership, with community members directly conferring disease resistance and heat tolerance on hosts (5–7). These microbial communities are often specialized, with distinct corals demonstrating co-evolution with members of their bacterial communities (8), but retain several prevalent genera, such as *Endozoicomonas*, across many hosts (9). While anthozoan communities are under active investigation, few analyses have documented the stability and nature of microbiome communities associated with zooxanthellate jellyfishes, a group with divergent life history traits from Anthozoa but with a comparable obligate symbiosis with dinoflagellates (clade Symbiodiniaceae).

The Upside-Down Jellyfish, *Cassiopea* spp., is a model organism for cnidarian-photosymbiont interactions (10,11). These medusae are residents of tropical and subtropical waters globally and can survive in water that is frequently shallow (as little as 8 cm depth) and hot (up to 33°C for *C. andromeda*) (12,13). *Cassiopea* obligatorily hosts algal symbionts, losing mass even when well fed while in aposymbiotic conditions (14). As in corals and other scyphozoans, mucus plays a central role for *Cassiopea*, presenting a sticky interface between the host and their benthic environment (15,16). In contrast to other symbiotic cnidarians, such as stony corals, *Cassiopea* can tolerate heat and intense temperature fluctuations(17).

48   Understanding the *Cassiopea* natural microbial community could provide new potential insights

49   into the function and diversity of cnidarian holobionts.

50     Past jellyfish (class Scyphozoa) microbial studies have focused on the microbiome of lab-

reared specimens (18,19) or limited sample sizes (20–22). While these types of studies may be

able to track state changes across life stages (18) and between condition types

(symbiotic/aposymbiotic) (19), there is little guarantee that the microbial communities of

jellyfish kept under prolonged laboratory conditions translate bear resemblance to *in situ*

microbiomes. Recent *in situ* studies have provided insights into microbial taxa associated with

some scyphozoans ("true jellyfish") but have not included species of the genus *Cassiopea*. In the

moon jellyfish, *Aurelia* spp.*,* individuals from the Chinese coast were *Vibrionaceae*-dominated,

while samples from the northwestern Atlantic were *Mycoplasma*-dominated(22,23). Between the

Indonesian marine lakes Kakaban, Haji Buang, and Tanah Bamban, *Mastigias* microbial

community composition varied significantly by lake, but consistently included *Oceanospirillales*

(primarily *Endozoicomonas*-like) (20). In samples across taxa, *Mycoplasma* is a common

component of the identified gut and tissue microbiome of jellyfishes, though its function remains

unknown (22–27). Vertebrate pathogen research is often a focus of scyphozoan microbiome

discovery—*Tenacibaculum* has been identified in several scyphozoan species, e.g. *Cotyllorhiza*

tuberculata, Aurelia aurita, and *Pelagia noctiluca*, and, as a fish pathogen, which can have

important fisheries ramifications (22,23,25,28).

67     As most studied jellyfish populations occur in coastal waters only sporadically, the

relative consistency of *Cassiopea* assemblages provides a system in which location effects can

be analyzed at single time points. *Cassiopea* spp. presents an opportunity to act as a comparison

group for the microbial communities associated with both other scyphozoans and symbiotic

anthozoans. The ubiquity of *Cassiopea* assemblages within the Florida Keys allows for the

examination of differences in the microbial community of one population (Florida Keys

Cassiopea) at one time point across a broad geographic area.

The aim of this work is to identify the core components of the internal and external

mucosal microbiomes of *Cassiopea* and compare the microbial communities of *Cassiopea* across

sites of the Florida Keys.

Materials and Methods

Sampling

A total of 55 medusae were collected by wading in eight sites along the length of the

Florida Keys in late August 2021 over eight days (see Table 1 for a list of sites). Species

identification for each individual is reported in Muffett & Miglietta 2023 and summarized in

Supplementary Table S1 (29). At each site, coordinates, temperature, time of day, depth, salinity,

and pH were recorded (Table 1). Salinity was measured with a refractometer, and temperature

and pH were measured with an Apera A1209. A minimum of two and a maximum of 10

medusae were sampled per site. At each site and in between each medusa, all equipment was

cleaned with absolute ethanol and 10% bleach solution. A 1 L water sample was collected from

above (>10 cm above the benthos) each medusa aggregation and filtered (0.2 um pore size,

Thermo Scientific Cat No.09-740-30G) with a manual sampler as described in the USGS manual

water sampling protocol (30), replicate filters were placed in 15 mL of absolute ethanol or

dimethyl sulfoxide ethylenediamine tetraacetic acid saturated salt storage solution (1 L pH 7.5:

93.06 g EDTA, 60 mL 20 % NaOH solution, 20 mL 25 % HCl, 40 mL DMSO, 800 mL water,

NaCl to oversaturation; see ref. 30) (commonly referred to as DESS), one filter in each buffer. At

each *Cassiopea* medusa aggregation, duplicate scrapings of ~10 g of sediment collections

(substrate sample) were placed in either 15 mL of absolute ethanol or DESS storage solution.

Table 1: List of collection sites. Table with latitude, longitude, salinity, pH, surface temperature,
average diameter of collected medusae from the site, and date of collection. Number of paired
GVC and bell swab samples per collection are for *Cassiopea xamachana* (CX). For collections
that included *C. andromeda* mitotyped medusae, the number of *C. andromeda* samples is
presented after the dash (CA). These values only include samples collected in DESS.

Site Name	Latitude	Longitude	Salinity (ppt)	pH	Temp (C)	Average depth (m)	Average col. medusa diam (cm)	# of GVC Samps	# of Bell Samps	Date (AM/PM)
Key West	24.561526	-81.7881727	36	8.2	30.4	0.56	6.62	CX 3/ CA 1*	CX 3/ CA 1*	15.8.2021 PM
Cudjoe Key	24.6775378	-81.4991819	36	8.4	32.6	0.36	5.91	CX 4/ CA 2*	CX 4/ CA 2*	17.8.2021 AM
Big Pine Key	24.6978731	-81.3572574	35	7.7	29.1	0.15	4.43	CX 4	CX 4	15.8.2021 PM
Bahia Honda Key TP1	24.6825793	-81.2293673	37	8.3	31.8	0.86	12.41	CX 2	CX 2	13.8.2021 PM
Bahia Honda Key TP2	24.6825793	-81.2293673	36	8	29.4	0.51	10.28	CX 2	CX 2	15.8.2021 AM
Marathon Key	24.69396	-81.0980507	31	8.1	35.2	0.08	6.53	CX 7	CX 7	11.8.2021 PM
Lower Matacumbe Key	24.8582157	-80.7267927	35	8.1	32.9	0.89	11.40	CX 2	CX 2	16.8.2021 PM
Tavernier	25.0233783	-80.4940316	31	8.1	33.3	0.35	12.26	CX 3	CX 3	16.8.2021 AM
Key Largo	25.0872104	-80.4415797	35	7.8	32.0	0.56	4.30	CX 3/ CA 1*	CX 3/ CA 1*	18.8.2021 AM

101   Each *Cassiopea* medusa was collected from shallow waters <1.5 m within a single

assemblage at each site. Medusae were gently grasped by the oral arms, lifted from the water,

swabbed along the apex of the bell, then placed bell-down onto a pre-prepared dissection plate

(Figure 1). External bell swabs were done in replicate and placed in 3 mL DESS or absolute

ethanol. Medusa oral arm base was bisected between oral arm groups, and replicate swabs were

inserted and circled the digestive cavity. One internal swab was placed in DESS, the other in

ethanol, as with external swabs. A section of bell tissue was then collected and placed in ethanol

for mitochondrial species confirmation (see 28). The size of each medusa was measured, and

individual medusae were photographed. As sampling was done in a nonsterile environment,

blanks for buffer with swab and buffer with filter were also taken.

Figure 1: Sampling strategy for medusae collection onsite. Demonstration of the typical
orientation of medusa (right upper) and swab locations (apical bell surface and internal to
medusa). GVC access strategy is displayed on the left and relevant environmental samples (water
filter, substrate sample) are displayed on the right.

DNA Extraction and Sequencing

Samples in DESS and EtOH were maintained at room temperature for three to ten days

and then frozen. DESS can be stored at room temperature for at least thirty days without

significantly compromising community integrity (31). DNA from all samples (see

Supplementary Table S1) was then extracted using Zymobiomics DNA Kits (PN# D4300T). All

samples were pre-tested through the ampliciation of the 16S rRNA gene (variable region V4)

amplifications. Only samples that produced visible bands for the V4 region were sent for 16S

rRNA V3-V4 preparation and sequencing. While DESS preserved samples yielded good

amplification results, EtOH-preserved samples largely failed to produce adequate amplification.

As such, only DESS-stored samples were analyzed. Sequencing was performed by the University

of Texas Medical Branch Sequencing Lab. Library was prepared with Zymo Quick 16S V3-V4

kit (PN #D6400) in 20 uL reactions (10 uL Quick-16S qPCR premix, 4 uL Quick-16S V3-V4

primers (341f: (CCTACGGGDGGCWGCAG, CCTAYGGGGYGCWGCAG) and 806r:

(GACTACNVGGGTMTCTAATCC)), 4 uL water and 2 uL sample). qPCR was run using

Roche LightCycler 480 Instrument II (10min 95C, [30sec 95C, 30sec 55C, 3min 72C]x20).

PCRs were enzymatically cleaned according to Zymo Quick-16S kit protocols, and barcodes

were added in a secondary reaction at 20 uL (10 min 95˚C, [30 sec 95˚C, 30 sec 55˚C, 3 min

72˚C]x5). Plates included two replicate microbial community standards (Zymobiomics Cat. No.

D6300) and two PCR negative controls. Samples were loaded at 10 pM, with 15% 10 pM PhiX

sequencing control, and run on Illumina MiSeq with a 600 cycle v3 kit.

Data preprocessing

Data generated was run through the DADA2 v 1.26 pipeline on R v 4.4 with only limited

deviations from the standard protocols for ASV generation(32,33). Briefly, sequences were

filtered and trimmed ("filterAndTrim(truncLen=c(285,230), maxN=0, maxEE=c(4,4),

truncQ=2,trimLeft = c(17,21)"), errors were error corrected ("learnErrors", "dada"), and

sequences were merged ("mergePairs"). Reads between 380 and 458 bp were retained. Chimeric

reads were removed ("removeBimeraDenovo"), then taxonomy was assigned using the Silva v

138.1 database ("assignTaxonomy"). Retention metrics were retained and these metrics along

with the full code can be found on the github (https://github.com/kadenmuffett/Cassiopea-Code-

Microbiome-Repository). Sequence and metadata tables were imported into phyloseq v 1.42,

where reads assigned to mitochondria and chloroplast were removed. Contaminants were

removed using the decontam package v. 1.18, identifying and removing 47 ASVs based on

differential abundance in swab and buffer controls(34,35).

For all downstream analyses, dataset was limited to those stored in DESS and medusae

without a *C. andromeda* mitotype. This includes 60 paired bell and GVC samples (30 of each), 8

water samples and 8 substrate samples with 6.47 mil sequences across 34808 taxa.The complete

dataset (NCBI Bioproject Accession number PRJNA1020388) contains 12.78 million reads.

Statistical analyses

Refraction curves were generated in vegan 2.6.6.1 ("rarecurve") (SFIG. 1)(36). Shannon

diversity was computed with vegan and observed diversity was ploted with phyloseq

("estimate_richness", "plot_richness")(34). Faith's phylogenetic diversity was computed using

the MiscMetabar v 0.10.1, phanghorn v 2.12.1, metagMisc v 0.5.0 and PhyloseqMeasures v 2.1

packages by first generating an ML tree from all sequences over 300 reads across the dataset,

then computing per sample Faith's diversity (37–42). Faith's diversity was displayed with

ggstatsplot v 0.13.0 (43). Bray curtis and Euclidean distance were computed in phyloseq with

"distance" to "ordinate" to "plot_ordination". Between group comparisons were completed with

the pairwiseAdonis package v 0.4.1 (" pairwise.adonis2") on Euclidean distance measurements

(44). Statistical comparisons were computed with Kruskal-Wallis using the Kruskal.test and

pairwise comparisons using the "kruskal.test" and "dunn.test" commands with holm p

adjustment (45).

Data transmutation for starplots and display was assisted by the speedyseq package v

0.5.3.9021 (46). Star plots were made using the stars package v 0.6-7 (47). Differential

abundance was calculated with ALDEx2 v 1.38.0 following established pipelines for

microbiome data (48,49). Dirichlet Multinomial mixture analysis, as described by Holmes and

implemented in the DirichletMultinomial package v 1.48 (50,51). Core microbiome was

computed with the microbiome package v 1.28.0(52).

Ethics Statement

173   This work was performed on lower level invertebrates not subject to Animal Use

174   Protocols or within the purview of Texas A&M University IRB approval. Permitting for these

175   specimens was waived by Florida Fish and Wildlife Conservation Commision.

Results

Data description

We analyzed the microbiome associated with 34 medusae from eight sites within the

Florida Keys. For each medusa, we analyzed the microbial composition of their gastrovascular

cavity (GVC) and external or bell mucus (herein referred to as "bell"), as well as their substrate

(Sub) or water environment (WTR). All medusae were identified as *Cassiopea xamachana,* with

the exception of four medusae identified as mitochondrial haplotype *C. andromeda* found in Key

West, Cudjoe Key, and Key Largo (see Supplementary Table S1). 10.13 and 9.07 million reads

were retained after filtering ("filterAndTrim") and merging ("mergePairs") respectively (for all

retention statistics, see supplementary table S2) . There were 8.91 million sequences after

chimera removal. After primary phyloseq filtration, the dataset included 6.47 million sequences

retained in 34808 unique ASVs across 76 total samples (30 bell, 30 GVC, 8 water, 8 substrate).

Reference sequences for top ASVs are also available in the supplementary material (table S3).

Distribution of sequences in full dataset

190   The dataset generated was predominantly bacteria with few archael reads. At the phylum

191   level, 57% of sequences were Proteobacteria, 7% were Firmicutes, 13% were Bacteroidota, and

192   4% were Cyanobacteria (Fig 2). No other phylum had greater than 3% relative abundance in the

193   dataset.

Figure 2. Top ten (top) phyla and (bottom) class relative abundances of sequences in samples aggregated by sample type.

At the class level, Gammaproteobacteria comprised 41% of all sequences. Alphaproteobacteria, the next most abundant, comprised 17%. Bacteriodia (12%), Bacilli (6%), Cyanobacteriia (4%), and Desulfobacteria (3%), were the other classes with a relative abundance above 2%. No Archaean group was prevalent above a relative abundance of 0.25%, with the three most abundant classes being Halobacterota (0.25%), Thermoplasmatota (0.21%) and Nanoarchaeota (0.22%).

Distribution of sequences within gastrovascular cavity (GVC) samples

Within gastrovascular cavity samples, 75% of all sequences belonged to just three orders, Pseudomonadales (39%), Mycoplasmatales (24%), and Enterobacterales (13%). The majority of Pseudomonadales sequences identified within the GVC were dispersed across very few OTUs, with 66% identified as *Endozoicomonas atrinae* ASV1 and 8% as *E. atrinae* ASV4. This was true of the Mycoplasmatales as well, with 77% of associated gut sequences identified as *Mycoplasma* sp. ASV2 or *Mycoplasma* sp. ASV9 (15%).

Distribution of sequences within bell mucus samples

At the phylum level, within the bell mucosal swabs, 55% of the sequences were Proteobacteria, 14.5% Bacteroidota, 5.1% Cyanobacteria, and 6% Desulfobacterota.

At the order level, the most abundant groups were Pseudomonadales (23%), Chromatiales (6%), Rhodobacterales (7%), Desulfobacterales (5%), and Flavobacteriales (5%). *Endozoicomonas*

216 *atrinae* ASV1 was the single most abundant surface mucus ASV at a mean of 12%. The next

217 most abundant were Synechococcus CC9902 ASV15 (1.3%), *Endozoicomonas atrinae* ASV4

218 (1.1%) and Chromatiales *Candidatus Thiobios* ASV6 (0.8%).

Substrate and water samples

Water samples shared some ASVs with the bell mucus of medusae, the largest

contributing ASV was Chromatiales *Candidatus Thiobios* sp. ASV6 (8%). Additional prominent

taxa were Rhodobacteraceae *HIMB11* ASV12 (6%), *HIMB11* ASV19 (5%), and SAR11 Clade

III ASV 10 (4%). By contrast, substrate samples had no ASVs surpassing a mean of 1%

abundance across sites, with the highest two ASVs Sandaracinaceae ASV94 and *Actibacter*

ASV105 reaching 0.6% mean abundance across sediment samples (see table 2; extended table is

table S4).

Table 2: Top 5 ASVs for each sample type. Table with ASVs ordered from highest to
lowest average proportion in samples of each type dispayed as a percentage of composition
within that sample type. Includes Silva 138 taxanomic assignment for each ASV.
230

	1	2	3	4	5
	ASV1 *Endozoicomonas atrinae*	ASV15 *Synechococcus* CC9902	ASV4 *Endozoicomonas atrinae*	ASV6 *Candidatus Thiobios*	ASV10 SAR11 *Clade III*
Bell	11.585 %	1.272 %	1.105 %	0.832 %	0.709 %
	ASV94 Sandaracinaceae	ASV105 *Actibacter*	ASV95 Desulfocapsaceae	ASV169 *Robiginitalea*	ASV358 *Clostridiisalibacter*
Sub	0.625 %	0.605%	0.566%	0.492%	0.474%
	ASV6 *Candidatus Thiobios*	ASV12 Rhodobacteracea e *HIMB11*	ASV19 Rhodobacteraceae *HIMB11*	ASV10 SAR11 *Clade III*	ASV18 *Marinobacterium*
WTR	7.521 %	5.839%	4.725%	4.497%	2.711%
	ASV1 *Endozoicomonas atrinae*	ASV2 *Mycoplasma*	ASV9 *Mycoplasma*	ASV4 *Endozoicomonas atrinae*	ASV11 *Vibrio coralliilyticus*
GVC	34.133%	20.099%	2.823%	2.327%	2.094%

Diversity

Shannon diversity varied by sample type (Kruskal-Wallis: chi-squared = 48.8, df = 3, p-value =1.412e-10). Shannon diversity had a mean of 2.30 in GVCs, 4.01 in water samples, 5.63 in substrate samples and 6.56 in bell samples. Shannon diversity indices for the gastrovascular samples varied from substrate samples and bell samples (Kruskal-Wallis Dunn test with Holm correction: GVC-Bell, p-val=0.00; GVC-Sub, p-val= 0.00; GVC-Water, p-val=0.07; Bell-Sub, p-val=0.07; Bell-Water, p-val=0.06; Water-Sub, p-val=0.01). Shannon diversity was not explained by location (Kruskal-Wallis: chi-squared = 9.7, df = 7, p-value =0.21)(Fig. 3).

Figure 3. Shannon diversity (a) by sample type and (b + c) location. GVC, bell, substrate and water Shannon diversity are shown across both the entire dataset or when segmented by sample type across sites with (b) GVC and (c) bell alpha diversity across sites. All p values of holm-adj Dunn comparison of means tests below a = 0.05 are displayed, with statistical test details displayed above each plot.

Beta diversity

Euclidean distance showed differences between medusa and non-medusa samples (pairwise.adonis, GVC vs bell: p=0.001, bell vs sub: p=0.049, bell vs water: p=0.022, GVC vs sub: p=0.001, GVC vs water: p=0.001, sub v water p=0.001). Variability along the primary axis was found in both GVC and bell samples, while variability along the secondary axis was found mainly in GVC samples. The location of collection had no clear impact on beta diversity.

Figure 4. PCA of Euclidean distance of samples. PCA is colored by sample collection latitude
from low (orange) to high (purple). Sample type is designated by shape, circle (Bell), triangle
(GVC), square (Substrate) and cross (Water). The plots show the same PCA faceted by sample
type.

Faith's diversity of communities were distinct from each other (Kruskal-Wallis test, chi-

squared = 44.1, df = 3, p-value = 1.4e-09). GVC communities were distinct from bell and

substrate comparison groups, while bells were distinct from water and gastrovascular cavities

(Dunn test with holm correction) (Fig. 5).

Figure 5. Violin plot of Faith's diversity. Samples are colored by sample type. Kruskal-Wallis
chi-sq, pvalue and pairwise dunn test with holm correction p-values are displayed. Differences
between GVC-water and substrate-bell are not significant.

Core microbiome

Core microbiome for the purposes of this study was evaluated based on 70% occurrence

of an ASV and a relative abundance of > 0.1% of reads in all 34 gastrovascular cavity samples or

external bell samples. This is in line with thresholds for other core microbiome work (53). The

most abundant ASV, *Endozoicomonas* ASV1, was present in 90% of GVC samples at a relative

abundance of > 0.1% of reads, ASV2 and ASV4 were present in 80% . Using less stringent

criteria (>70% of samples, any abundance), ASV 11, ASV 132 and ASV 6 become part of the

core microbiome (Fig 6). These ASVs are across only three genera- *Endozoicommonas*,

Mycoplasma and *Vibrio* (full taxonomic details of ASVs can be found in table S5).

Figure 6. Abundance of Cassiopea core GVC microbiome across dataset. The top five ASVs
from the gut core microbiome across all sample types (Bell, GVC, Substrate (Sub) and Water
(WTR)) organized from furthest south to furthest north sites. Note the difference in y axis
maxima between bell and GVC samples (top) and substrate and water samples (bottom).

In bell samples, ASV1, ASV104, and ASV86 have 73%, 70% and 70% relative

abundance in samples at a 0.1% threshold. ASV6, ASV95, ASV37, ASV105, and ASV135 are

present at any level in >70% of bell samples. These ASVs cover the genera *Endozoicommonas*,

Candidatus Thiobios, *Methyloceanibacter, Ruegeria* and unknown genera from

Desulfocapsaceae and Actibacter (Fig. 7).

Figure 7. Core microbiome prevalence in GVC and bell. Heatmap of gastrovascular cavity
(top) and bell (bottom) shaded by prevalence (0-1 scale) vs detection threshold (proportions of
reads in samples) across the ASVs identified as core in each group.

In terms of abundance, internal core microbiome components are not just present, but

highly dominant in *Cassiopea*'s gut. Some of these taxa are highly restricted to GVC samples

(*Mycoplasma*), while others are prevalent across body surfaces (*Endozoicomonas*) (Fig. 7).

The vast majority of ASVs were not differentially abundant between bell and GVC

samples. Differential abundance (Aldex2, min effect size absolute value >[±2]) between the

gastric and bell surfaces is restricted to the novel *Mycoplasma* group (ASV2), which is

completely lacking in external samples (Fig 8). No features were differentially abundant between

substrate and bell samples (Sfig 2).

Figure 8. Aldex2 plot of effect size of all ASVs between bell and GVC samples. Only ASV2 has an effect >[± 2].

Dirichlet multinomial mixture (DMM) analysis identified specific proportions of many of

these core microbiome as valuable for category differentiation. Within animal-associated

samples, model probability value was optimal for two communities, one GVC and one bell

without subcommunities. False assignment rate (mismatch between predicted community and

true sample type) was present in 5/60 samples. ASV1 amount was a core group differentiator,

with fewer meaningful ASVs defining the GVC community than the bell community (Fig. 9).

Figure 9. Most impactful ASVs for Dirichlet multinomial modeling. Top 0.5% most
impactful drivers of body cavity prediction in GVC (top) and bell (bottom) Dirichlet multinomial
models ordered from largest to smallest effect size.

Despite DMM identifying all gastric samples as belonging to the same community,

relative proportions of those community members differed dramatically, varying substantially in

abundance of *Endozoicommonas*, *Mycoplasma* and *Vibrio*. Variation within bell samples

primarily occurred with proportion of *Endozoicommonas,* but was not reliably distinguishable

from substrate samples by DMM (Fig. 10).

Figure 10. Star plots of community composition across sites. Star plots for internal (GVC),
external (Bell) mucus swabs, water (WTR) and substrate (Sub) sample averages at each site
separated by order (three most prominent orders and "other"). Each spoke of the star plot ranges
from 0% (center) to 100%. **Internal** group, starting from the 3' position and going
counterclockwise: Pseudomonadales, Mycoplasmatales, Enterobacteriales and "Other". **External**
group spokes (going counterclockwise): Pseudomonadales, Rhodobacterales, Chromatiales, and
"Other". **Water** group spokes (going counterclockwise): Pseudomonadales, Rhodobacterales,

Flavobacteriales, and "Other". **Substrate** group spokes (going counterclockwise): Rhodobacterales, Flavobacteriales, Desulfobacterales, and "Other". Each site's measured environmental conditions (salinity, water surface temperature, and pH) are presented below star plots, colored by intensity (low to high represented by pale to dark) . Sites are mapped onto their location within the Florida Keys.

Vibrio

ASVs identical to *Vibrio coralliilyticus* were found across *Cassiopea* gastric samples,

with abundance of up to 10% of reads within the sample. This far surpasses rates in the

surrounding water and substrate samples (ASV11 averages: Bell: 0.05%, WTR: 0.03%, Sub: 0%,

GVC: 2.09%) (Fig. 11).

Figure 11. Percent of reads (0%-10%) assigned to *V. coralliilyticus* ASVs in bell, GVC, substrate and water samples averaged by site. Organized by body cavity**.**

Host medusa species identity

The mitochondrial haplotype of the non-native *Cassiopea*, *Cassiopea andromeda,* a

cryptic sister species to *C. xamachana*, was found in four of the 34 medusae sampled. The four *C.*

andromeda mitotype individuals included were from the Key Largo (1), Cudjoe Key (2), and

Key West (1) sites. While these were excluded from the primary analysis, *Cassiopea andromeda*

showed no divergence from the trends of the *C. xamachana* simultaneously collected (Fig. 12).

Figure 12. PCA of Euclidean distance between *Cassiopea* species. *Cassiopea andromeda* (CA: grey) and *Cassiopea xamachana* (CX: black) samples at the sites were they cooccurred (Cudjoe Key, Garrison Bight and Key Largo).

Discussion

Cassiopea medusae show consistent differences in the microbial communities present on their bell and gastric surfaces. Gastrovascular cavity samples have lower diversity than the sediment, external surfaces, and host a novel *Mycoplasma* lineage. This low diversity is consistent with results for multiple other scyphozoans— including *Cotylorhiza tuberculata* (25). *Cassiopea* also appears to have some variation in community composition between sites, a feature consistent with results from other species, such as *Aurelia, Mastigias,* and *Tipedalia* (20,24).

A bifurcated outer and inner microbiome

Across all sites, the gastrovascular cavities of *Cassiopea* medusae showed a core microbiome comprised of taxa previously found in photosymbiotic cnidarians. The external microbial community of *Cassiopea* from the Florida Keys adheres fairly closely to the epibenthic microbial community at each site, with the addition of *Endozoicomonas*. ASV1 *Endozoicomonas* cf. *atrinae* may be a primary mucosal component and extend to the exterior of the body through mucus exchange, as it is less prevalent on the bell surface than internal for most individuals. Alternitavely, ASV1 may be a common mucosal resident with no function for the host.

The abundant *Mycoplasma* and *Vibrio* cf. *coralliilyticus* from the gastric cavity of *Cassiopea* were not found in any external mucus samples and found at far lower abundances in external mucosal samples respectively.

The high diversity of the *Cassiopea* bell microbiome found in this study may be unique to *Cassiopea*. Previous work has reported in the low-diversity mucosa, oral arms and umbrella of

a previously studied pelagic rhizostome, *Rhizostoma pulmo,* a likely result of the constant

contact of *Cassiopea* with the epibenthos (26). The limited overlap between diverse bell

communities and low diversity gut communities in *Cassiopea* is notable and consistent with the

significant differences in body compartment microbiome of other scyphozoans and anthozoans

(21,24,54). While the lack of diversity in scyphozoan microbiomes may seem distinct from many

described stony coral microbial communities, these stony coral microbial profiles are often

produced using whole body sampling. Lower diversity gastrovascular cavities with distinct

community composition may also be a feature of stony corals (55–57).

Microbial communities across sites within this study were not significantly different from

each other, as gastrovascular cavity components remained dominated by three taxa throughout

the Keys. This microbiome bears limited resemblance to those sampled in common garden

experiments and no resemblance to lab-held lines, demonstrating that "core" here should not be

misconstrued as obligate (19,58). The lack of obligate bacterial taxa in the *Cassiopea*

gastrovascular cavity is also consistent with other microbial hosts, notably the anemone

Exaiptasia diaphana, another successful model cnidarian (1). While *Cassiopea* require their

algal symbiont, bacterial partners may be transient, and bacteria may have far less significance to

host health for *Cassiopea* than for specialized obligate hosts (e.g., leafhoppers) (59–62). Without

a better understanding of specific metabolic function of different bacterial associates,

microbiome importance cannot be determined. We identified *Endozoicomonas*, *Mycoplasma,*

and *Vibrio* as the three most common bacterial genera found within the gastrovascular cavity of

Cassiopea. These are all commonly associated with coral microbial communities (8,9,63). Below,

we provide information on these common bacterial components of cnidarian holobionts.

Endozoicomonas functions in microbiomes

The large proportion of *Endozoicomonas* in internal and external samples fits within a

broader framework of both *Endozoicomonas* and known cnidarian-associated microbial

communities; *Endozoicomonas* strains are present in a variety of invertebrates, including many

corals, sponges, sea anemones, scyphozoans and cubozoans (9,63–66). Some hosts house

multiple distinct *Endozoicomonas* genotypes, suggesting that different *Endozoicomonas* clades

may provide different metabolic services to a host (e.g., Vitamin B12 metabolism) (67).

Endozoicomonas is more commonly identified in healthy scleractinian corals than bleached and

diseased ones (68). For these corals, it may play a role in a variety of fundamental

Symbiodinium–host interactions, including nutrient transport and symbiont cell breakdown (67).

Notably, *Endozoicomonas* is not a consistent marker of microbiome health, its disappearance and

reappearance without apparent host health declines has been identified in octocorals (2,66). In

some coral, members of *Endozoicomonas* are found in close proximity to *Symbiodinium,* it is

unknown whether this feature is shared with *Cassiopea* (67).

Within scyphozoa, *Endozoicomonas* is abundant in the microbial community of the

rhizostome scyphozoan *Mastigias* (20). While *Mastigias*, like *Cassiopea*, host zooxanthellae

from the family *Symbiodiniaceae*, not all symbiotic rhizostomes harbor the microbe, and not all

taxa with the microbe are symbiotic (20,25). *Endozoicomonas* may be common across medusae,

but the difficulty of culturing some strains may have excluded them from the results of past

culture-based analyses (66,67). Greater clarity is needed on the function of *Endozoicomonas* for

these medusae and other taxa.

Mycoplasmataceae: a poorly understood cnidarian microbiome

resident

Mycoplasma, like *Endozoicomonas*, is present in studied coral and scyphozoan

microbiomes (9,22,23,25). While intracellular parasitism is common for members of

Mycoplasmataceae, those associated with cnidarians may be located extracellularly and exist as

opportunistic commensals (2,69). *Mycoplasma* sp. is noted as a potential endosymbiont in lab-

raised *Aurelia* polyps, while adults from many jellyfish populations are known to have them

(22,24,27). The high divergence between this *Mycoplasma* and others of the genus (closest

sequence identity on NCBI at 89%: Accn. No. NR_115937) suggests that this may be a hitherto

unknown group. Whether this *Mycoplasma* represents an endosymbiont, a result of prey capture,

or a parasite, its exclusivity to the gastrovascular tract in *Cassiopea* sampled within the Keys is

notable, as other studies have not demonstrated gastrovascular cavity localization (24,27).

Vibrionaceae

Vibrionaceae is both a core part of coral microbiomes (9,70) and known to facilitate

metamorphosis within *Cassiopea* (71,72)*,* however, it is a highly diverse family. While *Vibrio*

spp. are found in gut communities of presumed healthy scyphozoans (Daley et al. 2016, Peng et

al. 2021), they can also be associated with senescence and disease (Kramar et al., 2019; Tinta et

al., 2012). For *Cassiopea* specifically, an increase in *Vibrio* has previously been associated with

an aposymbiotic or stressed state (58). Given the variation in Vibrionaceae species and strains,

family-wide statements on the beneficial or harmful nature of Vibrionaceae within *Cassiopea*

gastrovascular cavities should not be made.

Vibrio coralliilyticus

In this dataset we found ample signal of a 16S V3-V4 with 100% sequence match to

multiple *Vibrio* strains, including *V. coralliilyticus* and the oyster pathogen, *V. tubiashii*. As

Vibrio species are highly varied, the sequence match does not mean assured pathogenicity

towards any taxa (75–77). The apparent health of the medusae from whom these samples were

collected suggest a either a lack of impact of this specific group on the host or a host-microbe

facultative interaction between this *Vibrio* and *Cassiopea.* While there is data on *Cassiopea*

microbiome antimicrobial isolates and *Cassiopea* mortality when faced with *Serratia marcescens*

infection, we lack adequate details on the *Vibrio – Cassiopea* system to determine whether this

represents a lowlying widespread infection or a mutualism (78,79).

Drivers of external microbiome

While the external microbiome was not distinct from the sediment around it, it is worth

discussing the major groups more common externally than internally. While the core bell

microbiome represents a substrate-like community, the bell core microbiome includes

representatives from families and genera with a variety of potential metabolic capacities. The

most plentiful of these ASVs was *Candidatus Thiobios*, known as a sulfur-oxidizing epibiont of

the *Zoothamnium niveum* (80,81). As this bacteria can be found in the Florida Keys, their

appearance on *Cassiopea* may be circumstantial, but is consistent across medusae (80). Isolates

from *Methyloceanibacter* have been demonstrated to be methane oxidizers from marine

environments (82). These may be sediment associated or may recruit to the host to utilize algal

products. The potential sulfate reducers within Desulfocapaceae may similarly recruit to the bell

and its access to sediment pore waters (83). While *Pseudomonas* cf. *stuzeri* was not as high

abundance as the other metabolizers listed above, it may have a role in denitrification or coral

health for some species (84–86)

In addition to microbes with known metabolic functions that align with the *Cassiopea*

holobiont, the genus *Ruegeria* includes many marine species and has been identified in the

microbiomes of photsymbiotic and nonphotosymbiotic cnidarians alike (9,73,87). As Dirichlet

modeling for bell microbiomes gave similar weight to many bacteria, the relative impact of all of

these bacterial epibionts on the host is unclear.

Laboratory cultures

Cassiopea are primarily used for in-lab studies on symbiosis (12,14,88). Some of these

use wild-caught individuals, and others use lab-bred cultures, many of which have been held for

years. The lab-raised *Cassiopea* used in Rothig et al. 2021 appear to have a completely different

microbiome than the wild individuals caught here (19). This potential discrepancy should be

considered when using wild-caught or captive-bred individuals, as captive-bred symbiosis and

tolerance levels may differ from medusae in the wild. These discrepancies in microbiome are

found in other model cnidarians, such as *Exaiptasia* (1). The information on wild *Cassiopea*

microbial communities from the Florida Keys presented here may allow researchers working

with lab-raised individuals to better contextualize the taxa that they find in cultured medusae.

Conclusion

The microbiome of *Cassiopea* within the Florida Keys demonstrates strong body

compartment discrimination and includes a central group of ASVs overrepresented in the gut and

bell mucus. All of these families, Pseudomonadales, Vibrionaceae, and Mycoplasmataceae*,* are

found in other medusae and anthozoans. Despite this overrepresentation of some microbial taxa

within the gut, the *Cassiopea* bell is highly diverse, and overlaps with the communities of the

water and substrate in and on which it resides. The consistency of *Cassiopea* microbial

communities across the Florida Keys demonstrates that specific taxa are strongly associated with

Cassiopea and that this can extend across many sites with diverse physical conditions. This work

provides a greater understanding of the wild microbiome of the common lab organism *Cassiopea*.

Acknowledgements

490 The present work was part of KMM's PhD dissertation. Thank you to Dr. David

491 Retchless for input on figure design. We thank the Florida Fish and Wildlife Conservation

492 Commission for their help and guidance.

References

1. Brown T, Otero C, Grajales A, Rodriguez E, Rodriguez-Lanetty M. Worldwide

exploration of the microbiome harbored by the cnidarian model, Exaiptasia pallida

(Agassiz in Verrill, 1864) indicates a lack of bacterial association specificity at a lower

taxonomic rank. PeerJ. 2017;2017(5).

2. Van De Water JAJM, Allemand D, Ferrier-Pagès C. Host-microbe interactions in

octocoral holobionts - recent advances and perspectives. Microbiome. 2018;6(1):1–28.

3. Meyer JL, Castellanos-Gell J, Aeby GS, Häse CC, Ushijima B, Paul VJ. Microbial

Community Shifts Associated With the Ongoing Stony Coral Tissue Loss Disease

Outbreak on the Florida Reef Tract. Front Microbiol. 2019;10(September):1–12.

4. Djeghri N, Pondaven P, Stibor H, Dawson MN. Review of the diversity, traits, and

ecology of zooxanthellate jellyfishes. Mar Biol [Internet]. 2019;166(11):1–19. Available

from: https://doi.org/10.1007/s00227-019-3581-6

5.   Baldassarre L, Ying H, Reitzel AM, Franzenburg S, Fraune S. Microbiota mediated

507   plasticity promotes thermal adaptation in the sea anemone Nematostella vectensis. Nat

508   Commun. 2022;13(1).

509   6.   Ushijima B, Gunasekera SP, Meyer JL, Tittl J, Pitts KA, Thompson S, et al. Chemical and

510   genomic characterization of a potential probiotic treatment for stony coral tissue loss

511   disease. Commun Biol. 2023;6(1).

512   7.   Connelly MT, McRae CJ, Liu PJ, Martin CE, Traylor-Knowles N. Antibiotics Alter

513   Pocillopora Coral-Symbiodiniaceae-Bacteria Interactions and Cause Microbial Dysbiosis

514   During Heat Stress. Front Mar Sci. 2022;8(January):1–18.

515   8.   Pollock FJ, McMinds R, Smith S, Bourne DG, Willis BL, Medina M, et al. Coral-

516   associated bacteria demonstrate phylosymbiosis and cophylogeny. Nat Commun [Internet].

517   2018;9(1):1–13. Available from: http://dx.doi.org/10.1038/s41467-018-07275-x

518   9.   Huggett MJ, Apprill A. Coral microbiome database□: Integration of sequences reveals

519   high diversity and relatedness of coral- associated microbes. Environ Microbiol Rep. 2019

520   Jun 1;11(3):372–85.

521   10.   Puntin G, Sweet M, Fraune S, Medina M, Sharp K, Weis VM, et al. Harnessing the Power

522   of Model Organisms To Unravel Microbial Functions in the Coral Holobiont. Microbiol

523   Mol Biol Rev [Internet]. 2022 Oct 26 [cited 2023 Feb 24];86(4). Available from:

524   https://journals.asm.org/doi/10.1128/mmbr.00053-22

11.   Ohdera AH, Abrams MJ, Ames CL, Baker DM, Suescún-Bolívar LP, Collins AG, et al.

Upside-Down but Headed in the Right Direction: Review of the Highly Versatile

Cassiopea xamachana System. Front Ecol Evol. 2018 Apr 9;6:1–15.

12.   Mammone M, Ferrier-Pages C, Lavorano S, Rizzo L, Piraino S, Rossi S. High

photosynthetic plasticity may reinforce invasiveness of upside-down zooxanthellate

jellyfish in Mediterranean coastal waters. PLoS One [Internet]. 2021 Mar 1 [cited 2021

Jun 30];16(3 March):e0248814. Available from:

https://doi.org/10.1371/journal.pone.0248814

13.   Béziat P, Kunzmann A. Under pressure: Cassiopea andromeda jellyfish exposed to

increasing water temperature or lead, cadmium and anthropogenic gadolinium

contamination. Mar Biol Res [Internet]. 2022;18(1–2):48–63. Available from:

https://doi.org/10.1080/17451000.2022.2066132

14.   McGill CJ, Pomory CM. Effects of bleaching and nutrient supplementation on wet weight

in the jellyfish Cassiopea xamachana (Bigelow) (Cnidaria: Scyphozoa). Mar Freshw

Behav Physiol. 2008;41(3):179–89.

15.   Ames CL, Klompen AML, Badhiwala K, Muffett K, Reft AJ, Kumar M, et al.

Cassiosomes are stinging-cell structures in the mucus of the upside-down jellyfish

Cassiopea xamachana. Commun Biol [Internet]. 2020;3(1):1–15. Available from:

http://dx.doi.org/10.1038/s42003-020-0777-8

16.   Bakshani CR, Morales-Garcia AL, Althaus M, Wilcox MD, Pearson JP, Bythell JC, et al.

Evolutionary conservation of the antimicrobial function of mucus: A first defence against

infection. npj Biofilms Microbiomes [Internet]. 2018;4(1):1–12. Available from:

http://dx.doi.org/10.1038/s41522-018-0057-2

17.   Klein SG, Pitt KA, Lucas CH, Hung S, Schmidt-roach S, Aranda M, et al. Night-Time

549   Temperature Reprieves Enhance the Thermal Tolerance of a Symbiotic Cnidarian.

550   2019;6(July):1–16.

551   18.   Lee MD, Kling JD, Araya R, Ceh J. Jellyfish Life Stages Shape Associated Microbial

552   Communities, While a Core Microbiome Is Maintained Across All. Front Microbiol. 2018

553   Jul 12;0(JUL):1534.

554   19.   Röthig T, Puntin G, Wong JCY, Burian A, McLeod W, Baker DM. Holobiont nitrogen

555   control and its potential for eutrophication resistance in an obligate photosymbiotic

556   jellyfish. Microbiome. 2021;9(1):1–16.

557   20.   Cleary DFR, Becking LE, Polónia ARM, Freitas RM, Gomes NCM. Jellyfish-associated

558   bacterial communities and bacterioplankton in Indonesian Marine lakes. FEMS Microbiol

559   Ecol. 2016;92(5):1–14.

560   21.   Hao W, Gerdts G, Holst S, Wichels A. Bacterial communities associated with

561   scyphomedusae at Helgoland Roads. Mar Biodivers. 2019;49(3):1489–503.

562   22.   Peng S, Hao W, Li Y, Wang L, Sun T, Zhao J, et al. Bacterial Communities Associated

563   With Four Blooming Scyphozoan Jellyfish: Potential Species-Specific Consequences for

564   Marine Organisms and Humans Health. Front Microbiol. 2021 May 5;12(May):1–16.

565   23.   Daley MC, Urban-Rich J, Moisander PH. Bacterial associations with the hydromedusa

566   Nemopsis bachei and scyphomedusa Aurelia aurita from the North Atlantic Ocean. Mar

567   Biol Res. 2016;12(10):1088–100.

24.   Weiland-Bräuer N, Neulinger SC, Pinnow N, Künzel S, Baines JF, Schmitz RA.

Composition of bacterial communities associated with Aurelia aurita changes with

compartment, life stage, and population. Appl Environ Microbiol. 2015;81(17):6038–52.

25.   Viver T, Orellana LH, Hatt JK, Urdiain M, Díaz S, Richter M, et al. The low diverse

gastric microbiome of the jellyfish Cotylorhiza tuberculata is dominated by four novel

taxa. Environ Microbiol. 2017;19(8):3039–58.

26.   Basso L, Rizzo L, Marzano M, Intranuovo M, Fosso B, Pesole G, et al. Jellyfish summer

outbreaks as bacterial vectors and potential hazards for marine animals and humans health?

The case of Rhizostoma pulmo (Scyphozoa, Cnidaria). Sci Total Environ [Internet].

2019;692:305–18. Available from: https://doi.org/10.1016/j.scitotenv.2019.07.155

27.   Stabili L, Rizzo L, Basso L, Marzano M, Fosso B, Pesole G, et al. The microbial

community associated with Rhizostoma pulmo: Ecological significance and potential

consequences for marine organisms and human health. Mar Drugs. 2020;18(9).

28.   Delannoy CMJ, Houghton JDR, Fleming NEC, Ferguson HW. Mauve Stingers (Pelagia

noctiluca) as carriers of the bacterial fish pathogen Tenacibaculum maritimum.

Aquaculture [Internet]. 2011;311(1–4):255–7. Available from:

http://dx.doi.org/10.1016/j.aquaculture.2010.11.033

29.   Muffett K, Miglietta MP. Demystifying Cassiopea species identity in the Florida Keys:

Cassiopea xamachana and Cassiopea andromeda coexist in shallow waters. PLoS One

[Internet]. 2023;18(3):e0283441. Available from:

http://dx.doi.org/10.1371/journal.pone.0283441

30.   Laramie MB, Pilliod DS, Goldberg CS, Strickler KM. Environmental DNA Sampling

Protocol — Filtering Water to Capture DNA from Aquatic Organisms. In: US Geological

Survey Techniques and Methods, book 2. Reston, VA: U.S. Geological Survey; 2015. p.

1–15.

31.   Pavlovska M, Prekrasna I, Parnikoza I, Dykyi E. Soil sample preservation strategy affects

the microbial community structure. Microbes Environ. 2021;36(1):1–7.

32.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:

High-resolution sample inference from Illumina amplicon data. Nat Methods.

2016;13(7):581–3.

33.   R-Core-Team (Foundation for Statistical Computing). R: A Language and Environment

for Statistical Computing [Internet]. 2022. Available from: https://www.r-project.org/

34.   McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible Interactive Analysis

and Graphics of Microbiome Census Data. PLoS One. 2013;8(4).

35.   Davis NM, Proctor DiM, Holmes SP, Relman DA, Callahan BJ. Simple statistical

identification and removal of contaminant sequences in marker-gene and metagenomics

data. Microbiome. 2018;6(1):1–14.

36.   Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, et al. _vegan:

Community Ecology Package_. 2024.

37.   Paradis E, Schliep K. ape 5 . 0□: an environment for modern phylogenetics and

evolutionary analyses in R. 2019;35(July 2018):526–8.

38.   Mikryukov V. metagMisc: Miscellaneous functions for metagenomic analysis [Internet].

Available from: https://github.com/vmikk/metagMisc

39.   Taudiere A. MiscMetabar: Miscellaneous Functions for Metabarcoding Analysis

[Internet]. 2024. Available from: https://cran.r-project.org/package=MiscMetabar

40.	Tsirogiannis C, Sandel B. PhyloMeasures: Fast and Exact Algorithms for Computing

Phylogenetic Biodiversity Measures. 2017.

41.	Schliep KP. phangorn□: phylogenetic analysis in R. 2011;27(4):592–3.

42.	Battaglia T. btools: A suite of R function for all types of microbial diversity analyses.

2025.

43.	Patil I. Visualizations with statistical details: The "ggstatsplot" approach. J Open Source

Softw [Internet]. 2021;6(61)(3167). Available from: doi:10.21105/joss.03167

44.	Martinez Arbizu P. pairwiseAdonis: Pairwise Multilevel Comparison using Adonis

[Internet]. 2017. Available from: https://github.com/pmartinezarbizu/pairwiseAdonis

45.	Kassambara A. rstatix: Pipe-Friendly Framework for Basic Statistical Tests [Internet].

2023. Available from: https://cran.r-project.org/package=rstatix

46.	McLaren M. speedyseq: Faster implementations of phyloseq functions [Internet]. 2025.

Available from: https://github.com/mikemc/speedyseq

47.	Pebesma E, Bivand R. Spatial Data Science: With applications in R [Internet]. London:

Chapman and Hall/CRC; 2023. 352 p. Available from: https://r-spatial.org/book/

48.	Macvittie S, Doroodian S, Alberto A, Sogin M, Macvittie S, Doroodian S, et al. Exaiptasia

diaphana , following antibiotic exposure. 2024;9(6).

49.	Fernandes AD, Reid JNS, Macklaim JM, Mcmurrough TA, Edgell DR, Gloor GB.

Unifying the analysis of high-throughput sequencing datasets□: characterizing RNA-seq ,

16S rRNA gene sequencing and selective growth experiments by compositional data

analysis. 2014;1–13.

50.   Holmes I, Harris K, Quince C. Dirichlet Multinomial Mixtures□: Generative Models for

Microbial Metagenomics. 2012;7(2).

51.   Morgan M. DirichletMultinomial: Dirichlet-Multinomial Mixture Model Machine

Learning for Microbiome Data [Internet]. 2024. Available from:

https://doi.org/10.18129/B9.bioc.DirichletMultinomial

52.   Lahti L, Shetty S. microbiome R package. 2019.

53.   Neu AT. Defining and quantifying the core microbiome□: Challenges and prospects.

2021;118(51):1–10.

54.   Bonacolta AM, Connelly MT, Rosales SM, del Campo J, Traylor-knowles N. The starlet

sea anemone , Nematostella vectensis , possesses body region-specific bacterial

associations with spirochetes dominating the capitulum. FEMS Microbiol Lett.

2021;(August 2020):1–9.

55.   van Oppen MJH, Blackall LL. Coral microbiome dynamics, functions and design in a

changing world. Nat Rev Microbiol [Internet]. 2019;17(9):557–67. Available from:

http://dx.doi.org/10.1038/s41579-019-0223-4

56.   Agostini S, Yoshimi S, Higuchi T, Casareto BE. Biological and chemical characteristics

of the coral gastric cavity. Coral Reefs. 2011;(March).

57.   Bollati E, Hughes DJ, Suggett DJ, Raina J-B, Kühl M. Microscale sampling of the coral

gastric cavity reveals a gut-like microbial community. bioRxiv [Internet].

2024;2024.05.20.594925. Available from:

https://www.biorxiv.org/content/10.1101/2024.05.20.594925v1%0Ahttps://www.biorxiv.o

rg/content/10.1101/2024.05.20.594925v1.abstract

58. Carabantes N, Cerqueda-García D, García-Maldonado JQ, Thomé PE. Changes in the

Bacterial Community Associated With Experimental Symbiont Loss in the Mucus Layer

of Cassiopea xamachana Jellyfish. Front Mar Sci. 2022;9(May):1–13.

59. Wu W, Lei J-N, Mao Q, Tian Y-Z, Shan H-W, Chen J-P. Distribution, Vertical

Transmission, and Cooperative Mechanisms of Obligate Symbiotic Bacteria in the

Leafhopper Maiestas dorsalis (Hemiptera, Cicadellidea). Insects. 2023;1–13.

60. Sharp KH, Pratte ZA, Kerwin AH, Rotjan RD, Stewart FJ. Season, but not symbiont state,

drives microbiome structure in the temperate coral Astrangia poculata. Microbiome.

2017;5(1):120.

61. Hammer TJ, Sanders JG, Fierer N. Not all animals need a microbiome. FEMS Microbiol

Lett. 2019;366(10):1–11.

62. Riviere M La, Roumagnac M, Garrabou J, Bally M. Transient Shifts in Bacterial

Communities Associated with the Temperate Gorgonian Paramuricea clavata in the

Northwestern Mediterranean Sea. PLoS One. 2013;8(2).

63. Neave MJ, Apprill A, Ferrier-Pagès C, Voolstra CR. Diversity and function of prevalent

symbiotic marine bacteria in the genus Endozoicomonas. Appl Microbiol Biotechnol

[Internet]. 2016;100(19):8315–24. Available from: http://dx.doi.org/10.1007/s00253-016-

7777-0

64. Tinta T, Kogovšek T, Klun K, Malej A, Herndl GJ, Turk V. Jellyfish-associated

microbiome in the marine environment: Exploring its biotechnological potential. Mar

Drugs. 2019;17(2):1–33.

65. Maher RL, Schmeltzer ER, Meiling S, McMinds R, Ezzat L, Shantz AA, et al. Coral

Microbiomes Demonstrate Flexibility and Resilience Through a Reduction in Community

Diversity Following a Thermal Stress Event. Front Ecol Evol. 2020;8(October):1–15.

66. Pogoreutz C, Oakley CA, Rädecker N, Cárdenas A, Perna G, Xiang N, et al. Coral

holobiont cues prime Endozoicomonas for a symbiotic lifestyle. ISME J.

2022;16(8):1883–95.

67. Neave MJ, Michell CT, Apprill A, Voolstra CR. Endozoicomonas genomes reveal

functional adaptation and plasticity in bacterial strains symbiotically associated with

diverse marine hosts. Sci Rep. 2017;7(December 2016):1–12.

68. McDevitt-Irwin JM, Baum JK, Garren M, Vega Thurber RL. Responses of coral-

associated bacterial communities to local and global stressors. Front Mar Sci.

2017;4(AUG):1–16.

69. Neulinger SC, Gärtner A, Järnegren J, Ludvigsen M, Lochte K, Dullo WC. Tissue-

associated "Candidatus mycoplasma corallicola" and filamentous bacteria on the cold-

water coral Lophelia pertusa (Scleractinia). Appl Environ Microbiol. 2009;75(5):1437–44.

70. Sweet MJ, Bulling MT. On the importance of the microbiome and pathobiome in coral

health and disease. Front Mar Sci. 2017;4(January):1–11.

71. Hofmann DK, Neumann R, Henne K. Strobilation, budding and initiation of scyphistoma

morphogenesis in the rhizostome Cassiopea andromeda (Cnidaria: Scyphozoa). Mar Biol.

1978;47(2):161–76.

72.   Neumann R. Bacterial Induction of Settlement and Metamorphosis in the Planula Larvae

of Cassiopea andromeda (Cnidaria: Scyphozoa, Rhizostomeae). Vol. 1. 1979.

73.   Kramar MK, Tinta T, Lučić D, Malej A, Turk V. Bacteria associated with moon jellyfish

during bloom and post-bloom periods in the Gulf of Trieste (northern Adriatic). PLoS One.

2019;14(1):1–21.

74.   Tinta T, Kogovsek T, Malej A, Turk V. Jellyfish Modulate Bacterial Dynamic and

Community Structure. PLoS One. 2012;7(4):1–10.

75.   Pollock FJ, Wilson B, Johnson WR, Morris PJ, Willis BL, Bourne DG. Phylogeny of the

coral pathogen Vibrio coralliilyticus. Environ Microbiol Rep. 2010;2(1):172–8.

76.   Ben-Haim Y, Thompson FL, Thompson CC, Cnockaert MC, Hoste B, Swings J, et al.

Vibrio coralliilyticus sp. nov., a temperature-dependent pathogen of the coral Pocillopora

damicornis. Int J Syst Evol Microbiol. 2003;53(1):309–15.

77.   Richards GP, Watson MA, Needleman DS, Church KM, Häse CC. Mortalities of Eastern

And Pacific oyster larvae caused by the pathogens Vibrio coralliilyticus and Vibrio

tubiashii. Appl Environ Microbiol. 2015;81(1):292–7.

78.   Emery MA, Beavers KM, Van Buren EW, Batiste R, Dimos B, Pellegrino MW, et al.

Trade-off between photosymbiosis and innate immunity influences cnidarian's response to

pathogenic bacteria. Proc R Soc B Biol Sci. 2024;291(2032):20240428.

79.   Thomé PE, Rivera-Ortega J. Specific antibacterial activity against potential pathogens and

re-straining of larvae settlement from a pigmented Pseudoalteromonas strain isolated from

the jellyfish Cassiopea xamachana. Int Aquat Res. 2022;14(2):95–105.

80.   Bright M, Espada-hinojosa S, Lagkouvardos I, Volland J, Hole W. The giant ciliate

Zoothamnium niveum and its thiotrophic epibiont Candidatus Thiobios zoothamnicoli□: a

model system to study interspecies cooperation. 2014;5(April):1–13.

81.   Rinke C, Schmitz-esser S, Stoecker K, Nussbaumer AD, Vanura K, Wagner M, et al.

" Candidatus Thiobios zoothamnicoli ," an Ectosymbiotic Bacterium Covering the Giant

Marine Ciliate Zoothamnium niveum. 2021;72(3):2014–21.

82.   Vekeman B. Methanotrophic microbiomes from North Sea sediment.

83.   Song J, Hwang J, Kang I, Cho JC. OPEN A sulfate □ reducing bacterial genus ,

Desulfosediminicola gen . nov ., comprising two novel species cultivated from tidal □ flat

sediments. Sci Rep [Internet]. 2021;1–13. Available from: https://doi.org/10.1038/s41598-

021-99469-5

84.   Engelen AH, Aires T, Vermeij MJA, Herndl GJ, Serrão EA, Frade PR. Host

differentiation and compartmentalization of microbial communities in the azooxanthellate

cupcorals Tubastrea coccinea and Rhizopsammia goesi in the Caribbean. Front Mar Sci.

2018;9(NOV):1–9.

85.   Meirelles LA, Newman DK. Both toxic and beneficial effects of pyocyanin contribute to

the lifecycle of Pseudomonas aeruginosa. Mol Microbiol. 2018;110(6):995–1010.

86.   Carlson CA, Ingraham JL. Comparison of Denitrification by Pseudomonas stutzeri ,

Pseudomonas aeruginosa , and Paracoccus denitrificans. 1983;45(4):1247–53.

87.   Huo Y, Xu X, Li X, Liu C, Cui H, Wang C, et al. Ruegeria marina sp . nov ., isolated from

Marine Sediment. 2011;347–50.

88.   Klein SG, Pitt KA, Carroll AR. Reduced salinity increases susceptibility of zooxanthellate

jellyfish to herbicide toxicity during a simulated rainfall event. Environ Pollut [Internet].

2016;209:79–86. Available from: http://dx.doi.org/10.1016/j.envpol.2015.11.012

89.   Hyun DW, Shin NR, Kim MS, Oh SJ, Kim PS, Whon TW, et al. Endozoicomonas atrinae

sp. nov., isolated from the intestine of a comb pen shell Atrina pectinata. Int J Syst Evol

Microbiol. 2014;64(PART 7):2312–8.

90.   Sunagawa S, Woodley CM, Medina M. Threatened corals provide underexplored

microbial habitats. PLoS One. 2010;5(3):1–7.

Supplementary Information

Supplement A1: Full length 16S rRNA gene identities

Three samples were sent to CD Genomics for full length 16S on PacBio Sequel using

universal 16S primers (F: AGRGTTTGATYNTGGCTCAG;

R:TASGGHTACCTTGTTASGACTT). Samples sent for full length 16S rRNA sequencing

included gastrovascular cavity swab extractions from medusae Marathon Key GVC 30 and Key

West GVC 96 and GVC 120. As the usable reads returned from this were low (597-12520 per

sample), these full-length identities are not discussed in the main text. For further details, please

refer to the appendix (A1).

Full length 16S rRNA gene sequences (BioProject: PRJNA1020446) were run on mothur

v1.48.0 with deviation from the MiSeq SOP on trim length. ASV cutoff was set to five

differences, and ASV representative sequences were compared against the NCBI GenBank (Sept

24, 2023) database using BLAST to find nearest matches by e-value.

In order to identify core microbial groups to a lower taxonomic level, three internal

medusa gastrovascular swabs (medusae Kmuffet_96, Kmuffet_120 and Kmuffet_30) from

Marathon Key (MK) and Key West (GB) were sequence for full length 16S rRNA gene (see S1

for location of samples). As the primers were not chloroplast exclusionary, the Key West sites

produced few useable sequences (Kmuffet_96: 597 and Kmuffet_120: 1097). The Marathon Key

medusa (Kmuffet_30), likely bleached at time of collection, returned 12,520 non-chloroplast

sequences. The Mycoplasma-like full length 16S rRNA gene sequence (Genbank Acc. No.

OR592270) from Marathon Key and Key West has no closely related sequences currently

identified (greatest identity 88.43% to sequence HQ393440). The *Endozoicomonas* strain from

both Marathon Key and Key West (Genbank Acc. No. OR592271) has 99.93% identity to

Endozoicomonas atrinae strain WP70(T) collected from the intestine of South Korean *Atrina*

pectinata (Accession NR_134024)(89) and 99.80% identity (<5 nucleotide differences) with

isolates collected from *Gorgonia ventalina* (GenBank Acc. No. GU118345) in Bocas del Toro,

Panama (90). *Endozoicomanas* and *Mycoplasma*-like isolates were the only two groups to have

ASVs with over 100 sequences.

S1 Tables: (in Supplementary Excel Workbook)

Supplementary Table S1: Roster of collected medusae

Supplementary Table S2: Read processing tracking table

Supplementary Table S3: Representative sequences of all ASVs from the dataset

Supplementary Table S4: ASVs orderd by abundance in each sample type

Supplementary Table S5: Taxonomy of top 1000 ASVs in dataset

Sfig 1: Rarefaction curves of all samples

Internal

Water and sediment
blanks

External

Top drivers of DMM: GVC community

Top drivers of DMM: Bell community

V. coralliilyticus by average of site and sample type

Euclidean distance by sample type and species