LSD Modulates Proteins Involved in Cell Proteostasis, Energy Metabolism and Neuroplasticity in Human Brain Organoids

Marcelo N. Costa, Livia Goto-Silva, Juliana M. Nascimento, Ivan Domith, Karina Karmirian, Amanda Feilding, Pablo Trindade, Daniel Martins-de-Souza, and Stevens K. Rehen*

ABSTRACT: The effects of psychedelics encompass modulation of subjective experience, neuronal plasticity, brain activity and connectivity, constituting a complex phenomenon. Underlying these effects, molecular changes at the protein level are expected. Proteomic analysis of human brain cells can elicit a comprehensive view of proteins and biological processes regulated within the central nervous system. To explore the molecular pathways influenced by lysergic acid diethylamide (LSD), we utilized mass spectrometry-based proteomics on human brain organoids. This approach allowed for an in-depth analysis of the proteomic alterations induced by LSD, providing insights into its effects at the molecular level within a brain-like environment. Alterations in proteostasis and energy metabolism, which are required for neural plasticity, were observed. Alongside, we identified changes in protein synthesis, folding, autophagy, and proteasomal degradation, as well as in glycolysis, oxidative phosphorylation, cytoskeleton regulation, and neurotransmitter release, providing a comprehensive view of the regulation of cellular process by LSD exposure. Furthermore, the ability of LSD to induce plasticity in human brain cells was corroborated through complementary *in vitro* experiments focusing on neurite outgrowth. This study sheds light on the specific proteins that LSD influences, thereby enhancing neurite extension and plasticity.

KEYWORDS: LSD, psychedelics, human brain organoids, proteomics, proteostasis, energy metabolism, neuroplasticity, structural and functional plasticity.

INTRODUCTION

Lysergic acid diethylamide (LSD) is a classic psychedelic substance that induces altered states of consciousness characterized by changes in sensory perception, mood, and thought patterns (1). LSD's psychedelic effects are primarily attributed to its agonist actions at brain serotonin 2A receptors (5-HT2ARs) (2,3). LSD also binds to dopaminergic, adrenergic, and other subtypes of serotonergic receptors (4). Furthermore, studies have demonstrated that psychedelics can penetrate cellular membranes, allowing interaction with intracellular 5-HT2A receptors (5), and LSD exhibits allosteric binding to the tropomyosin receptor kinase B (TrkB), enhancing TrkB's interaction with brain-derived neurotrophic factor (BDNF) (6). The interaction with multiple receptors underscores the complex pharmacology of psychedelics.

Emerging evidence suggests that LSD could cause lasting changes in neuronal plasticity and brain function (7). LSD exhibited potential therapeutic effects for anxiety, depression, and addiction (8–10), conditions associated with impaired neuroplasticity (11–13). It is hypothesized that psychedelics like LSD exert long-term psychotropic effects by acutely inducing heightened plasticity in brain circuits (9). In rodents, LSD can stimulate structural remodeling, such as increased dendritic arborization and spinogenesis (14).

Despite important progress, the molecular mechanisms underlying LSD's pro-plasticity properties remain incompletely defined (15). There is a lack of data on how LSD impacts human neurons (16,17). Advances in stem cell technology enabled the generation of human brain organoids that model aspects of human neurobiology (18,19). Additionally, analyzing the proteomic profiling of these organoids can provide insights into drug-induced functional changes in the human brain (20–22).

In this study, we utilized proteomics to examine human brain organoids exposed to LSD. By adopting an unbiased methodology, our objective was to uncover proteomic patterns and biological pathways that are indicative of the neural plasticity induced by LSD in human brain cells. Furthermore, to validate our proteomic findings, we carried out a functional neurite outgrowth assay on these cells.

RESULTS AND DISCUSSION

Human cells within brain organoids express serotonin 2A receptors. The characterization of brain organoids was conducted through immunofluorescence, revealing distinct zones and diverse cell populations. The analysis revealed the presence of neurons and neural progenitors, as evidenced by positive immunostaining for β-tubulin III (TUJ1) and PAX6, respectively. The distribution of TUJ1 was widespread throughout the organoids, while the expression of PAX6 was concentrated near the ventricles, delineating the ventricle-like zone (proliferative zones) (Figure 1A). GFAP-positive cells, representing radial glia or astrocytes, were present throughout the organoids, including within the ventricular zones (Figure 1B). The expression of serotonin receptors was evidenced by positive labeling of 5-HT2A receptors, known for their key role in mediating psychedelic effects (2,3). This labeling demonstrated a clear colocalization with the neuronal marker MAP2, indicating the presence of these receptors within neurons (Figure 1C). Together, these results establish our brain organoid model as a suitable platform for investigating the effects of LSD on living human neural tissue.

Figure 1. Characterization of the human brain organoids. Nuclei are stained blue with 4',6-Diamidino-2-Phenylindole, Dihydrochloride (DAPI) in all images. (A) TUJ1 immunostaining (green), a neuronal marker, accompanied by PAX6 immunostaining in red, which highlights neural progenitors; (B) GFAP immunostaining identifies radial glia and astrocytes; (C) Immunostaining for 5-HT2A receptors, which colocalize with the neuronal marker MAP2. Scale bars represent 400 μm for whole organoid images and 60 μm for zoomed-in images.

LSD changes the proteome of brain organoids. In a previous study, we demonstrated that LSD enhances performance in novel object recognition tasks in rats and visuospatial memory tasks in humans. In the same study, proteomic data of brain organoids exposed to 10 nM LSD for 24 hours provided molecular insights into the mechanisms underlying the observed improvements. Specifically, we identified a notable modulation of metabolic pathways associated with neural plasticity, including the mTOR pathway (23).

In this study, our primary focus was on understanding the impacts of a higher LSD concentration on human brain cells, specifically 100 nM (32.34 ng/mL). This concentration was chosen based on a pharmacokinetic analysis of plasma LSD concentrations in healthy human participants (24), as well as from an investigation on postmortem tissues, including peripheral blood and brain (25).

In the pharmacokinetic study, after oral administration of 200 μg LSD, the geometric mean maximum plasma concentration was 3.1 ng/mL (24). However, concentrations of LSD in the brain tissue surpassed those found in the bloodstream. Analysis of postmortem brain samples revealed LSD concentrations ranging from 8.6 to 42.5 times higher than those detected in peripheral blood (25). It is important to highlight that macrodoses (hallucinogenic doses) administered to study subjects in clinical studies have typically exhibited a tenfold range, spanning from 20 to 200 μg (26–28). Considering the significant variation in LSD dosing, understanding the mechanisms influenced by different concentrations is essential. The exposure period was chosen considering the pharmacokinetic study mentioned above, which showed that LSD remained in

the plasma of most human participants for up to 24 hours after a 200 µg oral dose (24).

With this perspective, we selected a concentration tenfold higher than our previous research (23). This choice allowed us to explore the effects of an increased concentration and, at the same time, provided a framework for drawing meaningful comparisons with our earlier findings. This methodical increase in concentration aims to broaden our understanding of LSD's impact at varying levels.

After exposing the brain organoids to 100 nM LSD for 24 hours, the protein extracts from these organoids were subjected to liquid chromatography-tandem mass spectrometry (LC-MS/MS)-based shotgun proteomic analysis (Figure 2A). When comparing LSD-exposed and control organoids proteomes, we identified and quantified a total of 3195 proteins (false discovery rate [FDR] = 1%) (Supplementary Table 1), with 239 showing significant differences in their abundance (p-value <0.05). The names of all the differentially abundant proteins (DAPs) are shown in Supplementary Table 2. Among these proteins, 158 were downregulated and 81 upregulated (Figure 2B). The proteins that exhibited the highest increases were the tumor-suppressor protein involved in DNA repair ABRAXAS1, the calcium sensor for endo- and exocytosis of synaptic vesicles SYT1; the mitochondrial RNA polymerase POLRMT; the eukaryotic translation initiation factor EIF5A2 and TTC7B, an isoform of a subunit of the PI4KIIIα complex, responsible for the first step in plasma membrane phosphoinositide synthesis. Among the proteins that were found to decrease in abundance, the ones that exhibited substantial downregulation were the serine/threonine kinase BRAF, the transcriptional corepressor TLE3, the transcriptional coactivator TRIP12 (a.k.a. MED1), the linker Histone H1.4 (HIST1H1E) and the α-tubulin isoform TUBA3E (Figure 2B and 2C).

Figure 2. LSD alters the proteome of human brain organoids. In the experimental setup (**A**), 45-day-old brain organoids were exposed to either LSD or vehicle for 24 hours. Subsequently, label-free quantitative proteomic analysis was performed on the samples. The volcano plot (**B**) presents proteins identified by mass spectrometry-based shotgun proteomics in the organoids, highlighting DAPs induced by LSD. The significance threshold is marked by a horizontal line (p-value = 0.05), while two vertical lines denote the log fold change cut-off (−1.0 fold on the left and +1.0 fold on the right), distinguishing between minor and major alterations. Cold-colored circles represent proteins with significant decreases (dark blue for $log_2FC < −1.0$; light blue for $−1.0 ≤ log_2FC > 0$), whereas warm-colored circles indicate proteins with significant increases (orange for $0 < log_2FC ≤ 1.0$; red for $log_2FC > 1.0$). Proteins without significant differences are shown in gray. The top five most increased and decreased proteins are also displayed. In part (**C**), a horizontal bar chart shows the proteins with the most notable abundance changes, along with their corresponding fold changes (log_2) and p-values, which are indicated within the bars.

LSD regulates proteins involved in proteostasis, energetic metabolism, and neuronal plasticity. LSD-induced DAPs were subjected to enrichment analysis in Metascape (using KEGG and Reactome databases). The top 10 statistically enriched terms are shown in Figure 3A. The result exhibits a predominance of terms associated with cellular proteostasis (selective autophagy [$-\log_{10}P$ = 7.77]; apoptosis [$-\log_{10}P$ = 7.62]; and cellular responses to stress [$-\log_{10}P$ = 6.46]), energetic metabolism (glycolysis [$-\log_{10}P$ = 5.81]) and neuronal plasticity (signaling by Rho GTPases, Miro GTPases and RHOBTB3 [$-\log_{10}P$ = 16.97]; membrane trafficking [$-\log_{10}P$ = 12.63]; axon guidance [$-\log_{10}P$ = 8.41]; RHOQ GTPase cycle [$-\log_{10}P$ = 5.13]; RHOBTB GTPase cycle [$-\log_{10}P$ = 5.10]; and transmission across chemical synapses [$-\log_{10}P$ = 4.94]). Supplementary Table 3 shows the complete results derived from the pathway and process enrichment analysis.

We also constructed a protein-protein interaction (PPI) network with LSD-induced DAPs (Figure 3B) using the STRING database. The resulting network comprised 237 nodes (proteins) and 159 edges (interactions). Notably, CDC42 displayed the highest degree of connection (14 degrees), followed by SRC (11 degrees), RPL9 (9 degrees), PAK2 (8 degrees), RBBP7 (8 degrees), and PXN (8 degrees). Among these highly connected nodes, RPL9 and RBBP7 were associated with proteostasis. At the same time, CDC42, SRC, PAK2, and PXN were linked to neuroplasticity and are part of a significant cluster related to this process. Altogether, we identified ten significant clusters using MCODE analysis. Three clusters (2, 3, and 10) were related to cellular proteostasis, two clusters (4 and 8) to energy metabolism, and three clusters (1, 7, and 9) to neuronal plasticity. Specifically, clusters 2, 3, and 10 exhibited enrichments in translation (p = 6.0E-6), neddylation (p = 1.1E-5), and the proteasome complex (p = 5.5E-3), respectively. Clusters 4 and 8 significantly correlated with the glycolytic process (p = 1.1E-8) and oxidative phosphorylation (p = 2.6E-4), respectively. Clusters 1, 7, and 9 were associated with the regulation of actin cytoskeleton (p = 5.9E-7), microtubule cytoskeleton (p = 8.6E-7), and synaptic vesicle cycle (p = 9.5E-3), in that order.

Regarding proteostasis, selective autophagy is crucial for neuronal homeostasis (29) and plays key roles in guidance signaling, dendritic spine

architecture, spine pruning, vesicular release, and synaptic plasticity in neurons (30). Neddylation is a post-translational modification, wherein a particular family of E3 ubiquitin ligases are the best characterized substrates. This modification within these E3 ligases facilitates the ubiquitination of their respective targets (31,32). Furthermore, the enrichment of 'cellular responses to stress' and 'apoptosis' may stem from the abundant number of proteins involved in the proteostasis network in those categories. Various components of this network respond to proteotoxic stress, significantly influencing cellular decisions between apoptosis and survival (33–35). Thus, fundamentally, the regulation of the proteostasis network profoundly affects both the composition and functionality of the cellular proteome.

In reference to cellular energy metabolism, it encompasses the metabolic pathways responsible for ATP synthesis through NADH turnover. The two primary pathways involved in these processes are glycolysis/fermentation and oxidative phosphorylation (36). Notably, when investigating the functional enrichment and network analyses of DAPs induced by LSD, glycolysis prominently emerges as a modulated pathway in both analyses, while oxidative phosphorylation surfaces in the last one. Consequently, it appears that LSD induces changes in proteins associated with cellular energy metabolism.

Neuroplasticity can be classified into structural and functional plasticity. Structural plasticity refers to changes in neuronal morphology (37,38), while functional plasticity encompasses modifications in synaptic transmission strength. Alterations in presynaptic neurotransmitter release and postsynaptic responses mediated by specific receptors can influence such changes (39). As part of structural plasticity, signaling by Rho GTPases and axon guidance emerged as potential pathways elicited by LSD in the functional enrichment and network analyses (Figures 3A and 3B). The precise control of neuronal cytoskeletal dynamics plays a pivotal role in orchestrating structural plasticity (37), whereby the Rho family of small GTPases emerge as critical modulators (40,41). Many signaling cascades implicated in neuronal structural modifications converge upon actin and microtubule cytoskeletal networks as shared terminal effectors (37). Moreover, Rho GTPases and their regulatory factors serve as critical downstream constituents of guidance signaling pathways (42), thereby

establishing a connection between axon guidance molecules, pivotal for nervous system development, and their significance in governing synaptic plasticity in the mature brain (43,44). In terms of functional plasticity, our analysis uncovered enriched terms related to neurotransmitter release, membrane trafficking, and synaptic transmission, besides a cluster associated with the synaptic vesicle cycle in the interaction analysis. Modulation of the structural and functional aspects of neuroplasticity appears to be highly relevant when assessing the effects of LSD on human brain organoids.

Figure 3. LSD influences proteins associated with proteostasis, cellular energy metabolism, and neuronal plasticity, as evidenced by: (**A**) The functional distribution of DAPs, which are categorized based on KEGG Pathway and Reactome Gene Sets; (**B**) A PPI network of DAPs induced by 100 nM LSD. For this network, the interaction score was set to a high confidence level (0.7). The node sizes represent the degrees (number of connections), with significant clusters highlighted in gray. The node colors indicate the extent of fold change: dark blue for more than two-fold decrease, light blue for less than two-fold decrease, orange for less than two-fold increase, and red for more than two-fold increase.

The proteostasis network dynamically adapts the proteome, enabling cellular responses to LSD stimuli. Given that our enrichment and network analysis revealed terms, hubs, and clusters associated with the molecular machinery responsible for protein quantity and quality control, we focused on the proteostasis network pathway to understand the modulation of this machinery. In Figure 4, LSD-modulated proteins in each step of the network are identified and color-coded based on their regulation. We noted in this pathway that LSD-induced DAPs, which are predominantly reduced, potentially impact protein synthesis, folding, maturation, transport, targeting, and degradation via the ubiquitin-proteasome system and autophagy. A reduction in degradative pathways may extend the lifespan of key synaptic proteins, as their metabolic turnover is dependent on synthesis and degradation rates (45). Additionally, autophagy is known to be inhibited by mTOR pathway activation (46), a pathway significantly implicated in LSD's neuroplastic effects (14). However, further research is necessary to confirm whether LSD specifically influences any aspect of the proteostasis network, or if these observations are merely homeostatic responses to a stimulus. Neuroplasticity itself poses a challenge to the proteostasis network (47), as signals inducing plasticity necessitate the adaptation of protein content to elicit specific responses (48).

Figure 4. LSD modulates proteins within the proteostasis network. The schematic diagram illustrates the components of this network, comprising the synthesis, folding, maturation, targeting, and degradation of proteins. In this diagram, the DAPs affected by LSD are color-coded, reflecting the extent of their modulation: dark blue for more than two-fold decrease, light blue for less than two-fold decrease, orange for less than two-fold increase, and red for more than two-fold increase.

Changes in neural energy metabolism to support neuroplasticity. Our enrichment and network analyses suggest that LSD modulates key pathways in energetic metabolism, including glycolysis, the tricarboxylic acid (TCA) cycle, and oxidative phosphorylation, as depicted in Figure 5. We observed that glycolysis is a pathway with four out of ten regulated steps. Modulated targets include phosphofructokinase PFKM and pyruvate kinase PKLR, representing two out of

the three key steps responsible for the irreversible reactions of glycolysis. Furthermore, several proteins critical to the TCA cycle and oxidative phosphorylation were found to be downregulated. Upregulation of PKLR ($log_2FC = 0.37$), coupled with the downregulation of proteins in the subsequent steps of mitochondrial energy metabolism, may favor lactate formation. In the brain, astrocytes emerge as the primary producers of this metabolite (49). Considering that periods of heightened neuronal activity and plasticity demand increased energy (50) and recognizing astrocyte-derived lactate as an alternative energy source for neurons (51), an intriguing hypothesis arises. LSD might induce a metabolic shift in astrocytes, potentially affecting the astrocyte-neuron lactate shuttle and, consequently, impacting neuroplasticity (9,26,52). This hypothesis presents an interesting avenue for further research.

Figure 5. LSD exerts regulatory effects on proteins involved in key cellular energy metabolism pathways. The schematic representation highlights the most critical pathways: glycolysis, the TCA cycle, and oxidative phosphorylation. Within this schematic, the DAPs influenced by LSD in each pathway are displayed. These are color-coded to indicate the extent of their change: dark blue for more than two-fold decrease, light blue for less than two-fold decrease, orange for less than two-fold increase, and red for more than two-fold increase. 1,3BPG, 1,3-bisphosphoglycerate; F1,6PP, Fructose 1,6-bisphosphate; F2,6PP, Fructose 2,6-bisphosphate; Fru(6)P, Fructose 6-phosphate; GA3P, Glyceraldehyde 3-phosphate; OXPHOS, oxidative phosphorylation; PEP, Phosphoenolpyruvate.

LSD modulates proteins involved in cytoskeleton regulation and release of synaptic vesicles pathways. LSD exerts modulation on several crucial biological processes involved in neuroplasticity, with two noteworthy examples being the actin cytoskeleton pathway (KEGG ID: hsa04810; p= 8.6E-3) and the synaptic vesicle cycle pathway (KEGG ID: hsa04721; p= 3.2E-2). We present a simplified representation of their regulation in response to DAPs induced by 100 nM LSD (Figure 6A and 6B), as determined through KEGG pathway analysis. These findings underpin the previously described neuroplastic effects of LSD (14). Table 1 displays the representation names of the DAPs within pathways alongside their corresponding encoding genes.

According to the regulation of actin cytoskeleton map (Figure 6A), signaling to the cytoskeleton can be mediated through G protein-coupled receptors (GPCRs), integrins, and receptor tyrosine kinases (RTKs), leading to a wide range of effects, including alterations in cell shape. Indeed, it is well known that LSD can activate various G protein-coupled serotonin receptors, also increasing the levels of BDNF, which acts through the tyrosine kinase receptor TrkB (3,26). Additionally, it was recently shown that LSD can directly bind TrkB receptors, thereby facilitating the action of BDNF (6). Furthermore, it was also demonstrated that LSD regulates the expression of extracellular matrix proteins, including fibronectin (53).

Cellular responses to these external cues are regulated intracellularly through numerous signaling cascades, which include the Rho family of small GTPases and their downstream protein kinase effectors. Arp2/3 complex is a key factor in actin filament branching and polymerization, essential for dendritic spine structural plasticity and stability (54) which was upregulated in LSD exposed organoids. LSD also modulated Rho GTPase CDC42, the protein kinase PAK1, and WASF1, a member of the actin regulatory WAVE complex. Therefore, our findings suggest modulation of actin cytoskeleton, including focal adhesions, adherens junctions, actomyosin, and stress fibers (Figure 6A).

In the synaptic vesicle cycle pathway, we observed upregulation of proteins involved in the fusion of synaptic vesicles to the plasma membrane (Figure 6B). Rab proteins located on the vesicle membrane can form complexes with effector proteins, such as rabphilin and rab-interacting molecule (RIM), to facilitate the docking of synaptic vesicles (55). SYT1, one of the most upregulated proteins in the analysis, functions as the primary calcium sensor for neurotransmitter release. Upon Ca^{2+} binding, SYT1 triggers the complete assembly of the soluble NSF adaptor protein receptor (SNARE) complex, which leads to rapid and synchronized membrane fusion (56,57). The upregulation of proteins implicated in the release machinery implies an augmented likelihood of neurotransmitter release, indicating enhanced synaptic transmission capabilities (39,58–60).

Additionally, proteins associated with the recovery and recycling of synaptic vesicles, particularly in clathrin-mediated endocytosis (CME), exhibited significant alterations in abundance following exposure to LSD (Figure 6B). SYT1, up-regulated, not only functions as the Ca^{2+} sensor for neurotransmitter release but also acts in endocytosis (57). The soluble N-ethylmaleimide-sensitive factor (NSF) attachment protein α (α-SNAP, encoded by NAPA) is downregulated. α-SNAP, along with NSF, disassembles the SNARE complex after fusion (61). One isoform of the clathrin heavy chain was upregulated, whereas one subunit of the clathrin adaptor protein 2 (AP2) complex (AP2B1) was downregulated (Figure 6B). The downregulation of α-SNAP and the primary clathrin adaptor protein AP2 suggests an unfavorable impact on synaptic vesicle recovery via CME. The disassembly of the SNARE complex is crucial for the

retrieval of synaptic vesicles through CME (61), while clathrin depends on adaptor proteins to effectively bind to cargo (62–64).

Figure 6. Signaling pathway diagrams exhibit LSD-modulated proteins involved in cytoskeleton regulation and neurotransmitter release. The LSD-induced DAPs involved in each step are depicted with distinct colors: dark blue (higher than two-fold decreased), light blue (less than two-fold decreased), orange (less than two-fold increased), or red (higher than two-fold increased): **(A)** LSD-induced modulation of proteins involved in the regulation of the actin cytoskeleton pathway. Figure adapted from KEGG (KEGG ID: hsa04810; p= 8.6E-3). ACTN, actinin; ERM, ezrin, radixin or moesin; F2, coagulation factor II; F2R, coagulation factor II receptor; FN1, fibronectin 1; GF, growth factor; GPCR, G protein-coupled

receptors; Gα12,13, G protein subunit alpha 12, 13; ITG, integrin; MLC, myosin light chain; NT, neurotransmitter; ROCK, Rho kinase; **(B)** LSD-induced changes in proteins involved in synaptic vesicle cycle. Figure modified from KEGG (KEGG ID: hsa04721; p= 3.2E-2). AP2, adaptor protein 2 complex; NSF, N-ethylmaleimide-sensitive fusion protein; NT, neurotransmitter; RIM, Rab3 interacting molecule; SNAP, soluble NSF attachment proteins; SNARE, soluble NSF adaptor protein receptor; SYT, synaptotagmin; t-SNARE, target SNARE; VAMP, vesicle-associated membrane protein; VGCC, voltage-gated calcium channel; V-ATPase, vacuolar ATPase; v-SNARE, vesicle SNARE.

Table 1. DAP-encoding gene symbols and protein names in Figures 6A and 6B and their respective representations in the pathways. The upper section corresponds to Figure 6A, while the lower section refers to Figure 6B.

Representation	Gene symbol	Protein name	log$_2$FC
ACTN	ACTN1	Actinin alpha 1	-0.543033
Arp2/3	ACTR3	Actin related protein 3	-0.395243
	ARPC2	Actin related protein 2/3 complex subunit 2	1.40449
Cdc42	CDC42	Cell division cycle 42	-1.27146
c-Src	SRC	SRC proto-oncogene, non-receptor tyrosine kinase	-0.296145
MLC	MYL12B	Myosin light chain 12B	-0.675952
PAK	PAK2	p21 (RAC1) activated kinase 2	-0.559334
WAVE1	WASF1	WASP family member 1	-1.06246
α-SNAP	NAPA	NSF attachment protein alpha	-0.39629
AP2	AP2B1	Adaptor related protein complex 2 subunit beta 1	-1.20892
Clathrin	CLTCL1	Clathrin heavy chain like 1	1.23632
Rab3A	RAB3A	RAB3A, member RAS oncogene family	1.42592
SYT	SYT1	Synaptotagmin 1	4.44077

Two distinct LSD concentrations share modulation of proteins involved in regulating cell morphology and synaptic processes. Human brain organoids exposed to different concentrations of LSD (10 and 100 nM) exhibit similarities in the modulation of proteins and biological processes. The proteomic dataset of human brain organoids exposed to 10 nM LSD was obtained from the previous study by Ornelas et al. (23).

The comparison showed that distinct LSD concentrations resulted in a similar number of significantly modulated proteins, being 234 when exposed to 10 nM LSD and 239 to 100 nM LSD. Upon exploring the datasets, we identified 26 proteins that were modulated in both concentrations, as shown in figure 7A (purple lines). These persistently modulated proteins are potential candidates for a molecular pattern of LSD action in human brain. These 26 proteins are listed in Supplementary Table 4.

Among the consistently modulated proteins, the α6 subunit of GABA$_A$ receptors (GABRA6) and the protein O-GlcNAcase (OGA) emerge as commonly downregulated proteins. Besides the well-established roles of GABA$_A$ receptors, emerging evidence has identified this very specific subunit as a promising therapeutic target for addressing a wide spectrum of neurological and neuropsychiatric disorders (65,66). Interestingly, functional magnetic resonance imaging (fMRI) studies suggest that LSD perturbs the excitation/inhibition balance of the brain (67). On the other hand, OGA is an enzyme that mediates the removal of O-GlcNAc from target proteins (68). This modification was described to occur in many biological processes such as regulation of gene expression, signal transduction, cellular stress response, metabolism (69), and, more recently, in synaptic function (70). Reduction of OGA levels can decrease dendritic spine density in primary neurons (70). Although the exact roles of O-GlcNAcylation in synaptic function remain unclear, research has extensively investigated its role in aging-associated neurodegenerative diseases, such as Alzheimer's and Parkinson's disease. Elevated O-GlcNAcylation has been shown to prevent protein aggregation and slow neurodegeneration, making it a promising therapeutic target for these conditions (68).

Another noteworthy modulation is the LSD-induced upregulation of copine-1 (CPNE1) in both concentrations. Studies have demonstrated that CPNE1 activates the AKT-mTOR signaling pathway in neural stem cells (NSCs) (71).

Given the established association between mTOR and psychedelic-induced neural plasticity (14,23), CPNE1 and other copines may be potential upstream regulators of LSD-induced mTOR-mediated plasticity in other cell types as well. Despite few proteins being affected by both concentrations, numerous proteins modulated in one group shared common ontology with different proteins from the other group (Figure 7A – blue lines). This functional overlap underscores the remarkable similarity in the elicited biological processes, despite the modulation of several distinct proteins. And, by comparing the top 20 enriched terms from both analyses, we were able to identify processes that were commonly modulated. We noted a significant enrichment of terms associated with cellular stress at both concentrations (Supplementary Figure 1). However, this enrichment may result from proteostatic responses to LSD. It is noteworthy that certain mechanisms involved in protein turnover also play a role in responding to proteotoxicity. Interestingly, most proteins involved in cellular responses to stress and apoptosis are reduced under the 100 nM LSD condition, including proteins with pro-apoptotic functions like DAPK1 and MAPK8 (JNK1) (72,73). This observation could indicate a potential modulation of neuroprotective mechanisms. Indeed, recent research has revealed that psychedelics, such as LSD, can enhance the survival of dentate granule cells within the hippocampus of mice. These cells can integrate into the local circuitry of the dentate gyrus following maturation, thereby influencing plasticity (6). Lastly, the presence of several terms associated with the regulation of cell morphology (Figure 7B) and synaptic-related processes (Figure 7C), which are crucial for structural and functional plasticity, respectively, emerges as a prominent characteristic of LSD action at both concentrations.

Figure 7. Different LSD concentrations share modulation of several biological processes: (**A**) Overlap between genes encoding DAPs induced by 10 and 100 nM LSD. On the outside, the light gray arch corresponds to the 10 nM LSD gene list, and the dark gray to 100 nM LSD. On the inside, light orange represents genes uniquely modulated by one condition, and in dark orange those regulated in both conditions. The purple lines link the same genes common in both 10 nM and 100 nM LSD conditions. Blue lines link different genes with the same ontology term; (**B**) and (**C**) Significant terms associated with alterations in cell morphology and synaptic processes, respectively. The light gray bars correspond to data from the 10 nM LSD concentration, while the dark gray bars represent data from the 10 nM LSD concentration.

LSD induces neurite outgrowth in human brain cells. To specifically investigate the impact of LSD on structural plasticity, we conducted a neurite outgrowth assay, and the results are presented in Figure 8. In this assay, neurospheres were initially placed on poly-ornithine/laminin-coated plates for 24 hours. They were then exposed to LSD at concentrations of 10 nM and 100 nM for an equivalent duration. Representative images of neurospheres are displayed in Figure 8A. In our research, we employed the Sholl analysis to quantify changes on arbor complexity induced by LSD. This method focuses on the evaluation of neurite intersections against a sequence of concentric circles of gradually

increased radius, drawn around the neurospheres' core. A visual representation of the Sholl analysis conducted on these images is available in Supplementary Figure 2.

Neurite intersections of circles were analyzed by mean crossings in groups of five successive circles. Both concentrations, 10 and 100 nM LSD, led to an increase in the average number of neurite crossings in circles 6 to 10 (control: 236.77 ± 8.65; 10 nM LSD: 285.33 ± 12.04, $p<0.01$; 100 nM LSD: 272.84 ± 8.65, $p<0.05$), and 11 to 15 (control: 139.13 ± 11.75; 10 nM LSD: 218.46 ± 20.00, $p<0.001$; 100 nM LSD: 179.01 ± 16.62, $p<0.05$). The concentration of 10 nM LSD still increased the number of crossings in circles 16 to 20 (control: 39.63 ± 5.92; 10 nM LSD: 109.16 ± 16.44, $p<0.001$; 100 nM LSD: 57.07 ± 12.44). These findings suggest that both concentrations led to increased arbor complexity (branching and/or elongation). Interestingly, different LSD concentrations can impact neurite outgrowth dynamics in varying ways. Therefore, further studies are essential to comprehend these dose-dependent variations and temporally distinct responses (Figure 8B).

The lack of significant changes in the number of crossings in circles 1 to 5 (control: 253.63 ± 7.66; 10 nM LSD: 262.09 ± 7.28; 100 nM LSD: 266.85 ± 7.70) suggests that the observed effects weren't influenced by alterations in the number of primary neurites (Figure 8B). One the other hand, the absence of significant changes in intersections of circles 21 to 25 (control: 9.49 ± 1.78; 10 nM LSD: 28.04 ± 6.83; 100 nM LSD: 9.39 ± 2.30) and 26 to 30 (control: 5.19 ± 1.14; 10 nM LSD: 9.42 ± 3.18; 100 nM LSD: 2.61 ± 0.38) indicates that the effect was more pronounced in the proximal segments of the neurites (Figure 8B).

These findings are consistent with the results of Ly et al. (2018) in primary cortical rat cultures, where LSD was also found to promote neurite outgrowth. Similar to our observations, Ly et al. reported that LSD had a limited impact on the number of primary neurites and did not cause changes in the length of the longest dendrite (14). This parallel in outcomes across different experimental models underscores the potential of LSD in modulating neural structure, hinting at a consistent biological response irrespective of the model organism.

Figure 8. Neurite Outgrowth in Response to LSD: Human neurospheres were initially plated for 24 hours and then subjected to exposure to either 10 nM or 100 nM LSD for an additional 24 hours. For this analysis, the control group included 18 neurospheres, while the groups exposed to 10 nM and 100 nM LSD consisted of 17 and 15 neurospheres, respectively. (**A**) The neurospheres were analyzed using immunofluorescence staining for β-tubulin III (TUJ1; depicted in green) and DAPI staining to highlight nuclei (shown in blue). (**B**) Sholl analysis was employed to assess neurite outgrowth. An inset in this section provides a schematic of the Sholl analysis method. The analysis was conducted on the mean numbers of crossings for each group of five circles. The quantification was carried out over five independent experiments. Statistical significance is indicated as follows: *$p<0.05$, **$p<0.01$, ***$p<0.001$, in comparison to the control group. A scale bar indicating 1000 μm is included for reference.

CONCLUSIONS

Our study reveals that LSD exposure leads to a significant alteration in the abundance of numerous proteins in human brain organoids, marking a shift in the proteomic profile of human brain cells. The enrichment analysis of these DAPS indicates that LSD affects processes such as proteostasis, energy metabolism, and neuroplasticity.

Specifically, LSD modulates proteins involved in various aspects of the proteostasis network, including protein synthesis, folding, maturation, transport, autophagy, and proteasomal degradation. A notable observation is the reduction in most proteostasis proteins, potentially extending the lifespan of synaptic proteins by decelerating turnover rates reliant on a balance between synthesis and degradation (45). Additionally, LSD appears to inhibit autophagy, possibly through activation of the mTOR pathway (46), a known stimulant of LSD-induced neuroplasticity (14). However, it remains to be investigated whether LSD's regulation of proteostasis is a direct effect or an indirect homeostatic response. The adaptation in proteostasis is crucial for proteome remodeling and cellular plasticity (47,48).

Furthermore, LSD impacts the abundance of proteins involved in glycolysis, the TCA cycle, and oxidative phosphorylation. This suggests that psychedelics may modulate energy metabolism to accommodate the increased demands during neural excitation and plasticity (50). Interestingly, our data points to a metabolic shift favoring lactate production, a primary energy source from astrocytes supporting neuronal plasticity (49,51).

Our analysis also implicates LSD in pathways essential for structural and functional neuroplasticity, including cytoskeletal regulation and neurotransmitter release. The remodeling of dendrites requires precise control over actin and microtubule dynamics, typically mediated by Rho GTPases (37,40). Our findings suggest a phase in LSD's mechanism of action characterized by reduced actin polymerization and stabilization of the existing cytoskeleton, aligning with a transition from the initiation to the maintenance of structural plasticity changes (74). Additionally, LSD seems to enhance synaptic vesicle fusion proteins while

reducing components of clathrin-mediated endocytosis, hinting at increased neurotransmitter release, though its implications for reuptake warrant further investigation.

Interestingly, a noteworthy observation emerges from the comparison of proteins modulated in human brain organoids exposed to 100 nM LSD and those exposed to 10 nM LSD (23). A significant overlap in ontology is evident among the modulated proteins at both concentrations. This overlap is particularly pronounced in terms associated with cellular stress, regulation of cell morphology, and synaptic-related processes. The prevalence of terms linked to cellular stress may arise from proteostatic responses triggered by LSD or could potentially indicate a modulation of neuroprotective mechanisms. The presence of multiple terms pertaining to the regulation of cell morphology and synaptic-related processes may point towards events encompassing structural and functional plasticity, respectively. These biological processes, consistently regulated at both concentrations, appear to be important hallmarks of LSD action in the human brain. Furthermore, our research revealed that LSD stimulates neurite outgrowth in iPSC-derived brain cells. We observed this effect at both concentrations, 10 nM and 100 nM, where LSD was found to enhance the complexity of the neurites. This finding suggests a broader spectrum of LSD biological activity on neuronal plasticity.

In conclusion, our proteomic analysis uncovers potential mechanisms behind the LSD-induced plasticity previously reported (14). So, neuroplasticity induced by LSD was demonstrated in both proteomics and neurite outgrowth assay. Overall, these findings validate neuroplastic effects induced by LSD in human models and underscores the potential of psychedelics in treating conditions associated with impaired plasticity. Our study also highlights the value of human brain organoids as a tool for characterizing neural responses to psychedelics and deciphering aspects of neuroplasticity.

METHODS

hiPSC-derived human brain organoids. GM23279A, an hiPSC line from a healthy female donor obtained from the NIGMS Repository of the Coriell Institute, was used in this study. Cells were cultured on Matrigel (Corning, USA) coated plates with mTeSR1 (STEMCELL Technologies, Canada) at 37°C and in a 5% CO_2 atmosphere. Colonies were manually passed when 80% confluence was reached. hiPSCs were differentiated into brain organoids as described by Goto-Silva et al. (75). Shortly afterward, hiPSCs were detached and dissociated with Accutase (MP Biomedicals, USA) for 5 min at 37°C, generating single cells in suspension. Then, 9,000 cells were added in each well of an ultra-low binding 96-well plate (Corning, USA) in human embryonic stem cells (hESC) media (Dulbecco's modified eagle medium/Ham's F12 [DMEM/F12; Gibco, USA], 20% KnockOut Serum Replacement [KSR; Gibco, USA], 3% fetal bovine serum, certified, United States [Gibco, USA], 1% Glutamax [Gibco, USA], 1% minimum essential media-nonessential amino acids [MEM-NEAA; Gibco, USA], 0.7% β-mercaptoethanol [Gibco, USA] and 1% Penicillin-Streptomycin [Pen-Strep; Gibco, USA]) with 4 ng/ml basic fibroblast growth factor (bFGF; Invitrogen, USA) and 50 μM Rho-associated protein kinase inhibitor (ROCKi; Calbiochem, USA). This media was changed every other day for 6 days, and the embryoid bodies transferred to low-adhesion 24-well plates (Corning, USA) in neural induction media (DMEM/F12, 1% N2 supplement [Gibco, USA], 1% Glutamax, 1% MEM-NEAA, 1 μg/ml heparin [Sigma-Aldrich, USA]) and 1% Pen-Strep. The neural induction media was changed every other day for 4 days. On day 10, tissues were immersed, for 1h at 37°C, in a solution of Matrigel diluted in DMEM/F-12, according to the dilution factor given by the manufacturer. After this step, coated organoids were returned to the 24-well ultra-low-attachment plates with differentiation media without vitamin A [1:1 mixture of DMEM/F12 and Neurobasal (Gibco, USA), 0.5% N2 supplement, 1% B27 supplement without vitamin A (Gibco, USA), 3.5 ml/l 2-mercaptoethanol, 1:4,000 insulin (Sigma-Aldrich, USA), 1% Glutamax, 0.5% MEM-NEAA and 1% Pen-Strep] for 4 days, changing the medium every 48h. After the stationary growth, the organoids were transferred to 6 well plates under agitation (90 RPM), containing differentiation media with vitamin A [same composition of differentiation media above, except

B27 supplement with vitamin A (Gibco, USA)]. These media were replaced every 4 days until day 45.

LSD exposure. High-purity LSD was dissolved in ultrapure water (18.2 MΩ.cm) at room temperature, protected from light exposure. 45-day-old human brain organoids were exposed to 100 nM LSD for 24 hours, and the control group received culture medium.

Immunofluorescence in organoid sections. Five brain organoids were collected per condition (control and LSD) from each of the three independent experiments. Organoids were fixed overnight in 4% paraformaldehyde (PFA), rinsed with phosphate-buffered saline (PBS), dehydrated, cryoprotected in a 30% sucrose solution, and kept at 4°C for 48 – 72h. They were transferred into Tissue-Tek O.C.T. Compound (Sakura Finetek Japan, Japan), snap-frozen on dry ice, and stored at -80°C. Organoids were sectioned with 20 µM thickness in a cryostat (Leica Biosystems, Germany), and the slides were maintained at -80°C until staining with specific markers.

For immunofluorescence, slides were thawed for 30 min at 37°C, washed three times with PBS, and permeabilized with 0.3% Triton X-100 (Sigma-Aldrich, USA) diluted in 1X PBS for 15 min. For specific stainings (5-HT2A and PAX6), antigen retrieval procedure was performed before the permeabilization step. Sections were incubated in 10mM citrate buffer, 0.05% Tween 20, pH=6 for 10 min at 98°C. Following this step, the sections were blocked with 1% BSA + 10% normal goat serum (NGS) in PSB for 2 hours. All antibodies were diluted in this blocking solution. Cryosections were incubated overnight with primary antibody at 4°C, washed three times for 5 min with PBS, and incubated with secondary antibody for 2h. The antibodies used in this study are indicated in Supplementary Table 5. After three more washes with PBS for 5 min, the sections were incubated with 300 nM DAPI (Invitrogen, USA) for 5 min, for nuclear staining. Then, a last round of three washes with PBS was performed, and the slides were cover-slipped with Aqua-Poly/Mount (Polysciences, USA).

Images of organoid slices were acquired in a Leica TCS SP8 confocal microscope. For the immunofluorescence images, a 20x oil-immersion objective lens was used.

Liquid chromatography-mass spectrometry. Ten 45-day-old human brain organoids were collected from each of three independent experimental batches. Five of these organoids were exposed to 100 nM LSD for 24 hours, and the other five received only medium (control condition). After exposure, organoids were pelleted and frozen at −80 °C until sample processing for mass spectrometry-based label-free shotgun proteomics. Organoids were lysed in a buffer containing 7 M urea, 2 M thiourea, 1% CHAPS, 70 mM DTT, and Complete Protease Inhibitor Cocktail (Roche, Switzerland). Total protein content was measured using the bicinchoninic acid (BCA) method. The protein extracts (50 µg) were then in-gel digested with trypsin overnight (1:100, w/w, Sigma-Aldrich, USA). Peptides were applied to a reverse-phase liquid chromatographer [Acquity UPLC M-Class System (Waters Corporation, USA)], coupled to a Synapt G2-Si mass spectrometer (Waters Corporation, USA). Data-independent acquisition (DIA) strategy was used with ion mobility separation (high-definition data-independent mass spectrometry; HDMSE). Peptides were loaded onto a first-dimension chromatography on an M-Class BEH C18 Column (130 Å, 5 µm, 300 µm X 50 mm, Waters Corporation, USA) and eluted by discontinuous fractionation steps (13%, 18%, and 50% acetonitrile). After each step, the peptide loads were directed to a second-dimension chromatography on a nanoACQUITY UPLC HSS T3 Column (1.8 µm, 75 µm X 150 mm; Waters Corporation, USA) and eluted with acetonitrile [7 to 40% gradient (v/v), for 54 min, at a flow rate of 0.4 µL/min] into a Synapt G2-Si. MS/MS analysis was performed using nano-electrospray ionization in positive ion mode [nanoESI (+)] and a NanoLock Spray ionization source (Waters Corporation, UK). The lock mass channel was sampled every 30 seconds. Mass spectrometer calibration was performed with a [Glu1]-fibrinopeptide B human (Glu-Fib) (Sigma-Aldrich, USA) solution with an MS/MS spectrum reference from the NanoLock Spray source. Samples were run in technical duplicates of biological triplicates.

Database search and quantification. HDMSE raw data was imported to Progenesis® QI for Proteomics software, version 4.0 (Waters Corporation, USA), where protein identification and quantification was processed using dedicated algorithms (including Apex3D, peptide eD, and ion accounting informatics). Using default parameters for ion accounting and quantitation, peptide identification was carried out using UniProt's human proteomic database (version 2018/09, reviewed and non-reviewed). During the database queries, the databases used were reversed "on the fly" and appended to the original database to assess the false positive identification rate. Parameters for peptide identification: 1) in trypsin digestion; 2) variable modifications by oxidation (M) and fixed modification by carbamidomethyl (C); and 3) FDR less than 1%. Identifications not satisfying these criteria were not included in the analysis. Relative quantitation of proteins was done with Hi-N method (N=3). The quantitative analysis was carried out on the $\log_2$ values of the measured intensities. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (76) partner repository with the dataset identifiers PXD027369 and PXD037814.

In silico **analysis.** The fold change in the abundance of each protein was determined as a ratio of the intensity values from LSD-exposed and control organoids in each batch. Statistical analysis, aiming for the identification of the DAPs between control and LSD-exposed organoids, was carried out in Perseus software (version 2.0.7.0, Max-Planck-Gesellschaft, München) (77,78). Firstly, the rows with the same gene name were combined, and the ratios' median was used to calculate the resulting combined abundance values. Fold change values were $\log_2$ transformed, and a one sample t-test was performed to evaluate whether the abundance of each protein was significantly changed. Significant hits ($p < 0.05$) were considered DAPs induced by LSD and subjected to the bioinformatic analysis.

For biological process and pathway enrichment analysis, given DAPs were launched on the Metascape platform (http://metascape.org/) (79). The analysis considered the following ontology databases: Reactome Gene Sets and KEGG Pathway. Terms with a minimum overlap of 3.0, P value cutoff 0.01, and minimum enrichment of 1.5 were collected and groupe into clusters. For PPI analysis, a

network with given DAPs was constructed using the STRING 11.5 database (https://string-db.org/) (80) with a confidence score > 0.7 and visualized in Cytoscape 3.9.1 (81). Molecular Complex Detection (MCODE; Network scoring: loops not included, degree cutoff = 2; Cluster finding: haircut parameter, node score cutoff = 0.2, k-core = 2, and max. depth = 100) and CytoHubba (degree ranking method) (82) plugins of Cytoscape software were applied to obtain clusters and hub proteins, respectively. The functional annotation of the clusters was obtained in the version 2021 of the Database for Annotation, Visualization, and Integrated Discovery (DAVID 2021) (https://david.ncifcrf.gov/). The thresholds were count = 2 and EASE = 0.1. For an in-depth study of possibly affected pathways, schemes were constructed, showing where the given DAPs would act in each pathway. The pathways were chosen based on previous analysis. Some schemes were constructed based on the literature, and for others, the Kyoto Encyclopedia of genes and genomes (KEGG) pathway enrichment in DAVID 2021 was employed.

In the comparative analyses, we contrasted the data obtained from the analysis of this paper (100 nM LSD) with the data from Ornelas et al.'s study (23) (10 nM LSD). To assess the similarity level in the ontology, we employed Metascape to perform an overlap correlation. Furthermore, we compared the top 20 most enriched terms in both analyses, aiming to identify the commonly modulated processes.

hiPSC-derived human neurospheres. hiPSCs were differentiated into NSCs using PSC Neural Induction Medium (Thermo Fisher Scientific, USA), following the manufacturer's guidelines. Media was changed every other day until day 7. At this stage, NSCs were split and expanded in neural expansion medium, composed of a 1:1 ratio of Advanced DMEM/F12 and Neurobasal medium supplemented with neural induction supplement (Thermo Fisher Scientific, USA). To form neurospheres (83), 8.0×10^4 NSCs in 150 µL were seeded into each well of round-bottom ultra-low attachment 96-well plates (Corning, USA) and centrifuged at 300 g for 3 minutes to facilitate settling. After 72 hours, the medium was switched to a differentiation medium (1:1 mixture of Neurobasal medium and DMEM, enriched with B27 and N2 supplements), with subsequent media

replacements every other day. By day 10 of aggregation, the neurospheres were ready for the neurite outgrowth assay.

Neurite outgrowth assay: Each well in 96-well plates (Perkin Elmer, USA) was seeded with a single neurosphere, coated with 100 µg/ml poly-ornithine and 20 µg/ml laminin. After settling for 24 hours, the neurospheres were exposed to 10 and 100 nM LSD for an additional 24 hours. Subsequently, neurospheres were submitted to both immunofluorescence and Sholl analysis.

Immunofluorescence in plated neurospheres. Following a 15-minute fixation in a 4% paraformaldehyde solution, neurospheres were washed with PBS and then permeabilized using 0.3% Triton X-100 in PBS for 15 minutes. Subsequently, a blocking step was carried out by incubating the neurospheres in a solution containing 3% fetal calf serum in PBS for 1 hour. Following blocking, the neurospheres were subjected to overnight incubation at 4˚C with the primary antibody TUJ1 (Neuromics, USA) at a dilution of 1:2000 in the blocking solution. After incubation, the neurospheres underwent three washes with PBS and were re-blocked using 3% fetal calf serum for 20 minutes. Next, the neurospheres were incubated with the secondary antibody, specifically goat anti-mouse Alexa Fluor 594 (Thermo Fisher Scientific, USA) at a dilution of 1:400, for 60 minutes. To visualize nuclei, a counterstain using 0.5 µg/mL DAPI was performed for 5 minutes. Finally, a solution composed of a 1:1 mixture of glycerol and PBS was added to the plates, and images were captured using a 20x objective on an Agilent BioTek Cytation 1 imaging system.

Sholl analysis in plated neurospheres. This study used Sholl analysis to assess neurite outgrowth in plated neurospheres. The Neuroanatomy plugin within Fiji ImageJ software (version 1.54f) was employed for this purpose (84). The analysis involved setting up 30 circles originated from the neurosphere body, each spaced at 31.5 µm intervals. Neurite outgrowth was quantified by analyzing the mean crossings in groups of five circles. Statistical analysis was performed

using two-way ANOVA, complemented by Dunnets's post-hoc test for multiple comparisons. A visual representation of the Sholl analysis can be found in Supplementary Figure 2.

AUTHOR INFORMATION

Corresponding Author

Stevens K. Rehen – D'Or Institute for Research and Education (IDOR), Rio de Janeiro, Brazil; Department of Genetics, Institute of Biology, Federal University of Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil; stevens.rehen@idor.org**.**

Authors

Marcelo N. Costa – D'Or Institute for Research and Education (IDOR), Rio de Janeiro, Brazil; Department of Genetics, Institute of Biology, Federal University of Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil.

Lívia Goto-Silva – D'Or Institute for Research and Education (IDOR), Rio de Janeiro, Brazil.

Juliana M. Nascimento – D'Or Institute for Research and Education (IDOR), Rio de Janeiro, Brazil; Institute of Biology, University of Campinas (UNICAMP), Campinas, Brazil.

Ivan Domith – D'Or Institute for Research and Education (IDOR), Rio de Janeiro, Brazil; Rio de Janeiro, Brazil.

Karina Karmirian – D'Or Institute for Research and Education (IDOR), Rio de Janeiro, Brazil.

Amanda Feilding – The Beckley Foundation, Oxford, UK.

Pablo Trindade – College of Pharmacy, Federal University of Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil.

Daniel Martins-de-Souza – Laboratory of Neuroproteomics, Department of Biochemistry and Tissue Biology, Institute of Biology, University of Campinas, Campinas, Brazil; D'Or Institute for Research and Education (IDOR), Rio de Janeiro, Brazil; Experimental Medicine Research Cluster (EMRC), University of Campinas, Campinas, SP 13083-862, Brazil; Instituto Nacional de Biomarcadores em Neuropsiquiatria (INBION), Conselho Nacional de Desenvolvimento Científico e Tecnológico, São Paulo, Brazil; INCT in Modelling Human Complex Diseases with 3D Platforms (Model3D), São Paulo, Brazil.

Author Contributions

S.K.R., and M.N.C. conceived and designed the study; M. N. C. performed all cell cultures; J.M.N. carried out the proteomics and assisted in the interpretation; D.M.S. supervised proteomic experiments and data interpretation; M.N.C., L.G.S., and P.T. analyzed the majority of the proteomic data; L. G. S. and I. D. undertook conducted the experiments related to neurite outgrowth; M.N.C. drafted the manuscript and figures; K. K. undertook the characterization of cerebral organoids; S.K.R coordinated all the study; A.F. contributed to the conceptual framework of the proposal and provided critical review and feedback on the manuscript; All authors reviewed and approved the final manuscript prior to submission.

Funding

This project was supported by the Beckley Foundation, Conselho Nacional de Desenvolvimento Científico e Tecnológico – Brasil (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), grants 2017/25588-1 and 2019/00098-7, Instituto Serrapilheira, Pioneer Science Initiative

Notes

The authors declare no competing interest.

ACKNOWLEDGEMENTS

We acknowledge the contributions on technical support of Beatriz Luzia de Mello Lima Guimarães, Gabriela Lopes Vitória, Ismael Carlos da Silva Gomes, and Jhonata de Sousa do Nascimento. We also thank Joana Cardoso for her contribution to the immunofluorescence experiments and the acquisition of images of these experiments. We used the GPT-3.5 language model, developed by OpenAI, as a spelling correction tool. The use of ChatGPT in this specific context had no significant impact on the content or substance of the text.

ABBREVIATIONS

α-SNAP, soluble NSF attachment protein α; 5-HT, 5-hydroxytryptamine; ACTN, actinin; AP2, adaptor protein 2 complex; ATP, adenosine triphosphate; BDNF, brain-derived neurotrophic factor; BSA, bicinchoninic acid; BSA, bovine serum albumin; cDNA, Complementary DNA; CME, clathrin-mediated endocytosis; CNS, central nervous system; CPNE1, copine-1; DAP, differentially abundant proteins; DAPI, 4',6-Diamidino-2-Phenylindole; DAVID, Database for Annotation, Visualization, and Integrated Discovery; DIA, data-independent acquisition; DMEM/F12, Dulbecco's Modified Eagle Medium/Nutrient Mixture F-12; DNA, deoxyribonucleic acid; DTT, dithiothreitol; ERM, ezrin, radixin or moesin; F2, coagulation factor II; F2R, coagulation factor II receptor; FC, fold change; FDR, false discovery rate; fMRI, functional magnetic resonance imaging; FN1, fibronectin 1; Gα12,13, G protein subunit alpha 12, 13; GABA, *gamma-*

aminobutyric acid; GF, growth factor; GPCR, G protein-coupled receptor; GTPase, guanosine triphosphatases; Glu-Fib, [Glu1]-fibrinopeptide B human; HDMSE, high-definition data-independent mass spectrometry; hESC, human embryonic stem cell; HGNC, HUGO Gene Nomenclature Committee; hiPSC, human induced pluripotent stem cell; HUGO, Human Genome Organization; ITG, integrin; KEGG, Kyoto Encyclopedia of Genes and Genomes; KSR, KnockOut Serum Replacement; LC-MS/MS, liquid chromatography tandem mass spectrometry; LSD, lysergic acid diethylamide; M-MLV, moloney murine leukemia virus; MAP2, microtubule-associated protein 2; MCODE, Molecular Complex Detection; MEM-NEAA, minimum essential media-nonessential amino acids; MLC, myosin light chain; mRNA, messenger ribonucleic acid; mTOR, mammalian target of rapamycin; mTeSR1, modified TeSR1 medium; NADH, nicotinamide adenine dinucleotide; NSF, N-ethylmaleimide-sensitive fusion protein; NSC, neural stem cell; NT, neurotransmitter; OGA, O-GlcNAcase; PBS, phosphate-buffered saline; PCR, polymerase chain reaction; PFA, paraformaldehyde; Pen-Strep, Penicillin-Streptomycin; PPI, protein-protein interaction; Rho, Rat sarcoma virus homolog; ROCK, Rho kinase; ROCKi, Rho-associated protein kinase inhibitor; RIM, Rab-interacting molecule; RNA, ribonucleic acid; SNAP, soluble NSF attachment proteins; SNARE, soluble NSF adaptor protein receptor; RPM, rotation per minute; RTK, Receptor Tyrosine Kinase; STRING, Search Tool for the Retrieval of Interacting Genes; t-SNARE, target SNARE; TCA, tricarboxylic acid; Taq, *Thermus aquaticus*; TeSR1, serum-free medium for human pluripotent stem cells; TrkB, tropomyosin receptor kinase B; VAMP, vesicle-associated membrane protein; VGCC, voltage-gated calcium channel; V-ATPase, vacuolar ATPase; v-SNARE, vesicle SNARE.

REFERENCES

1. Preller KH, Vollenweider FX. Phenomenology, Structure, and Dynamic of Psychedelic States. In 2016. p. 221–56.

2. De Gregorio D, Aguilar-Valles A, Preller KH, Heifets BD, Hibicke M, Mitchell J, et al. Hallucinogens in Mental Health: Preclinical and Clinical Studies on LSD, Psilocybin, MDMA, and Ketamine. The Journal of Neuroscience. 2021 Feb 3;41(5):891–900.

3. Nichols DE. Psychedelics. Pharmacol Rev. 2016 Apr;68(2):264–355.

4. Ray TS. Psychedelics and the Human Receptorome. PLoS One. 2010 Feb 2;5(2):e9019.

5. Vargas M V., Dunlap LE, Dong C, Carter SJ, Tombari RJ, Jami SA, et al. Psychedelics promote neuroplasticity through the activation of intracellular 5-HT2A receptors. Science (1979). 2023 Feb 17;379(6633):700–6.

6. Moliner R, Girych M, Brunello CA, Kovaleva V, Biojone C, Enkavi G, et al. Psychedelics promote plasticity by directly binding to BDNF receptor TrkB. Nat Neurosci. 2023 Jun 5;26(6):1032–41.

7. Grieco SF, Castrén E, Knudsen GM, Kwan AC, Olson DE, Zuo Y, et al. Psychedelics and Neural Plasticity: Therapeutic Implications. The Journal of Neuroscience. 2022 Nov 9;42(45):8439–49.

8. Appelbaum LG, Shenasa MA, Stolz L, Daskalakis Z. Synaptic plasticity and mental health: methods, challenges and opportunities. Neuropsychopharmacology. 2023 Jan 9;48(1):113–20.

9. Calder AE, Hasler G. Towards an understanding of psychedelic-induced neuroplasticity. Neuropsychopharmacology. 2023 Jan 19;48(1):104–12.

10. Fuentes JJ, Fonseca F, Elices M, Farré M, Torrens M. Therapeutic Use of LSD in Psychiatry: A Systematic Review of Randomized-Controlled Clinical Trials. Front Psychiatry. 2020 Jan 21;10.

11. Price RB, Duman R. Neuroplasticity in cognitive and psychological mechanisms of depression: an integrative model. Mol Psychiatry. 2020 Mar 4;25(3):530–43.

12. Rezayof A, Ghasemzadeh Z, Sahafi OH. Addictive drugs modify neurogenesis, synaptogenesis and synaptic plasticity to impair memory formation through neurotransmitter imbalances and signaling dysfunction. Neurochem Int. 2023 Oct;169:105572.

13. Sha Z, Xu J, Li N, Li O. Regulatory Molecules of Synaptic Plasticity in Anxiety Disorder. Int J Gen Med. 2023 Jul;Volume 16:2877–86.

14. Ly C, Greb AC, Cameron LP, Wong JM, Barragan E V., Wilson PC, et al. Psychedelics Promote Structural and Functional Neural Plasticity. Cell Rep. 2018 Jun;23(11):3170–82.

15. Mller F, Borgwardt S. Acute effects of lysergic acid diethylamide (LSD) on resting brain function. Swiss Med Wkly. 2019 Sep 30;

16. Lee K, Park TIH, Heppner P, Schweder P, Mee EW, Dragunow M, et al. Human in vitro systems for examining synaptic function and plasticity in the brain. J Neurophysiol. 2020 Mar 1;123(3):945–65.

17. Liechti ME. Modern Clinical Research on LSD. Neuropsychopharmacology. 2017 Oct 27;42(11):2114–27.

18. Porciúncula LO, Goto-Silva L, Ledur PF, Rehen SK. The Age of Brain Organoids: Tailoring Cell Identity and Functionality for Normal Brain Development and Disease Modeling. Front Neurosci. 2021 Aug 13;15.

19. Benito-Kwiecinski S, Lancaster MA. Brain Organoids: Human Neurodevelopment in a Dish. Cold Spring Harb Perspect Biol. 2020 Aug;12(8):a035709.

20. Nascimento JM, Saia-Cereda VM, Sartore RC, da Costa RM, Schitine CS, Freitas HR, et al. Human Cerebral Organoids and Fetal Brain Tissue Share Proteomic Similarities. Front Cell Dev Biol. 2019 Nov 28;7.

21. Frantzi M, Latosinska A, Mischak H. Proteomics in Drug Development: The Dawn of a New Era? Proteomics Clin Appl. 2019 Mar 25;13(2).

22. Wang Y, Chiu JF. Proteomic Approaches in Understanding Action Mechanisms of Metal-Based Anticancer Drugs. Met Based Drugs. 2008 Jul 22;2008:1–9.

23. Ornelas IM, Cini FA, Wießner I, Marcos E, Araújo DB, Goto-Silva L, et al. Nootropic effects of LSD: Behavioral, molecular and computational evidence. Exp Neurol. 2022 Oct;356:114148.

24. Dolder PC, Schmid Y, Steuer AE, Kraemer T, Rentsch KM, Hammann F, et al. Pharmacokinetics and Pharmacodynamics of Lysergic Acid Diethylamide in Healthy Subjects. Clin Pharmacokinet. 2017 Oct 14;56(10):1219–30.

25. Mardal M, Johansen SS, Thomsen R, Linnet K. Advantages of analyzing postmortem brain samples in routine forensic drug screening—Case series of three non-natural deaths tested positive for lysergic acid diethylamide (LSD). Forensic Sci Int. 2017 Sep;278:e14–8.

26. de Vos CMH, Mason NL, Kuypers KPC. Psychedelics and Neuroplasticity: A Systematic Review Unraveling the Biological Underpinnings of Psychedelics. Front Psychiatry. 2021 Sep 10;12.

27. Petranker R, Anderson T, Maier LJ, Barratt MJ, Ferris JA, Winstock AR. Microdosing psychedelics: Subjective benefits and challenges, substance testing behavior, and the relevance of intention. Journal of Psychopharmacology. 2022 Jan 8;36(1):85–96.

28. Bonnelle V, Smith WJ, Mason NL, Cavarra M, Kryskow P, Kuypers KP, et al. Analgesic potential of macrodoses and microdoses of classical psychedelics in chronic pain sufferers: a population survey. Br J Pain. 2022 Dec 14;16(6):619–31.

29. Evans CS, Holzbaur ELF. Quality Control in Neurons: Mitophagy and Other Selective Autophagy Mechanisms. J Mol Biol. 2020 Jan;432(1):240–60.

30. Hwang JY, Yan J, Zukin RS. Autophagy and synaptic plasticity: epigenetic regulation. Curr Opin Neurobiol. 2019 Dec;59:207–12.

31. Zou T, Zhang J. Diverse and pivotal roles of neddylation in metabolism and immunity. FEBS J. 2021 Jul 20;288(13):3884–912.

32. Zhou L, Jiang Y, Luo Q, Li L, Jia L. Neddylation: a novel modulator of the tumor microenvironment. Mol Cancer. 2019 Dec 3;18(1):77.

33. Wang M, Law ME, Law BK. Proteotoxicity and endoplasmic reticulum stress-mediated cell death. In: Mechanisms of Cell Death and Opportunities for Therapeutic Development. Elsevier; 2022. p. 119–74.

34. Brancolini C, Iuliano L. Proteotoxic Stress and Cell Death in Cancer Cells. Cancers (Basel). 2020 Aug 23;12(9):2385.

35. Iuliano L, Dalla E, Picco R, Mallavarapu S, Minisini M, Malavasi E, et al. Proteotoxic stress-induced apoptosis in cancer cells: understanding the susceptibility and enhancing the potency. Cell Death Discov. 2022 Oct 4;8(1):407.

36. Rigoulet M, Bouchez CL, Paumard P, Ransac S, Cuvellier S, Duvezin-Caubet S, et al. Cell energy metabolism: An update. Biochimica et Biophysica Acta (BBA) - Bioenergetics. 2020 Nov;1861(11):148276.

37. Weerasinghe-Mudiyanselage PDE, Ang MJ, Kang S, Kim JS, Moon C. Structural Plasticity of the Hippocampus in Neurodegenerative Diseases. Int J Mol Sci. 2022 Mar 20;23(6):3349.

38. Urbanska M, Swiech L, Jaworski J. Developmental Plasticity of the Dendritic Compartment: Focus on the Cytoskeleton. In 2012. p. 265–84.

39. Wu X, Hu S, Kang X, Wang C. Synaptotagmins: Beyond Presynaptic Neurotransmitter Release. The Neuroscientist. 2020 Feb 3;26(1):9–15.

40. Duman JG, Blanco FA, Cronkite CA, Ru Q, Erikson KC, Mulherkar S, et al. Rac-maninoff and Rho-vel: The symphony of Rho-GTPase signaling at excitatory synapses. Small GTPases. 2022 Dec 31;13(1):14–47.

41. Zhang H, Ben Zablah Y, Zhang H, Jia Z. Rho Signaling in Synaptic Plasticity, Memory, and Brain Disorders. Front Cell Dev Biol. 2021 Oct 4;9.

42. Niftullayev S, Lamarche-Vane N. Regulators of Rho GTPases in the Nervous System: Molecular Implication in Axon Guidance and Neurological Disorders. Int J Mol Sci. 2019 Mar 25;20(6):1497.

43. Charron F. Axon Guidance: Gained in Translation. Neuron. 2018 Jul;99(1):1–2.

44. Thompson-Steckel G, Kennedy TE. Maintaining and Modifying Connections: Roles for Axon Guidance Cues in the Mature Nervous System. Neuropsychopharmacology. 2014 Jan 9;39(1):246–7.

45. Rosenberg T, Gal-Ben-Ari S, Dieterich DC, Kreutz MR, Ziv NE, Gundelfinger ED, et al. The roles of protein expression in synaptic plasticity and memory consolidation. Front Mol Neurosci. 2014 Nov 12;7.

46. Wang MM, Feng YS, Yang SD, Xing Y, Zhang J, Dong F, et al. The Relationship Between Autophagy and Brain Plasticity in Neurological Diseases. Front Cell Neurosci. 2019 May 24;13.

47. Hetz C. Adapting the proteostasis capacity to sustain brain healthspan. Cell. 2021 Mar;184(6):1545–60.

48. Giandomenico SL, Alvarez-Castelao B, Schuman EM. Proteostatic regulation in neuronal compartments. Trends Neurosci. 2022 Jan;45(1):41–52.

49. Horvat A, Zorec R, Vardjan N. Lactate as an Astroglial Signal Augmenting Aerobic Glycolysis and Lipid Metabolism. Front Physiol. 2021 Sep 30;12.

50. Watts ME, Pocock R, Claudianos C. Brain Energy and Oxygen Metabolism: Emerging Role in Normal Function and Disease. Front Mol Neurosci. 2018 Jun 22;11.

51. Roumes H, Dumont U, Sanchez S, Mazuel L, Blanc J, Raffard G, et al. Neuroprotective role of lactate in rat neonatal hypoxia-ischemia. Journal of Cerebral Blood Flow & Metabolism. 2021 Feb 24;41(2):342–58.

52. Olson DE. Biochemical Mechanisms Underlying Psychedelic-Induced Neuroplasticity. Biochemistry. 2022 Feb 1;61(3):127–36.

53. Nardou R, Sawyer E, Song YJ, Wilkinson M, Padovan-Hernandez Y, de Deus JL, et al. Psychedelics reopen the social reward learning critical period. Nature. 2023 Jun 22;618(7966):790–8.

54. Liu CCS, Cheung PW, Dinesh A, Baylor N, Paunescu TC, Nair A V., et al. Actin-related protein 2/3 complex plays a critical role in the aquaporin-2 exocytotic pathway. American Journal of Physiology-Renal Physiology. 2021 Aug 1;321(2):F179–94.

55. Wang Y qiang, Lin W wei, Wu N, Wang S yi, Chen M zi, Lin Z hua, et al. Structural insight into the serotonin (5-HT) receptor family by molecular docking, molecular dynamics simulation and systems pharmacology analysis. Acta Pharmacol Sin. 2019 Sep 27;40(9):1138–56.

56. Gruget C, Bello O, Coleman J, Krishnakumar SS, Perez E, Rothman JE, et al. Synaptotagmin-1 membrane binding is driven by the C2B domain and assisted cooperatively by the C2A domain. Sci Rep. 2020 Oct 22;10(1):18011.

57. Chen Y, Hu S, Wu X, Xie Z, Wang Y, Wang B, et al. Synaptotagmin-1 is a bidirectional Ca $^{2+}$ sensor for neuronal endocytosis. Proceedings of the National Academy of Sciences. 2022 May 17;119(20).

58. Quiñones-Frías MC, Littleton JT. Function of Drosophila Synaptotagmins in membrane trafficking at synapses. Cellular and Molecular Life Sciences. 2021 May 22;78(9):4335–64.

59. Baker K, Gordon SL, Melland H, Bumbak F, Scott DJ, Jiang TJ, et al. SYT1-associated neurodevelopmental disorder: a case series. Brain. 2018 Sep 1;141(9):2576–91.

60. Körber C, Kuner T. Molecular Machines Regulating the Release Probability of Synaptic Vesicles at the Active Zone. Front Synaptic Neurosci. 2016 Mar 2;8.

61. Sauvola CW, Littleton JT. SNARE Regulatory Proteins in Synaptic Vesicle Fusion and Recycling. Front Mol Neurosci. 2021 Aug 6;14.

62. Gulbranson DR, Crisman L, Lee M, Ouyang Y, Menasche BL, Demmitt BA, et al. AAGAB Controls AP2 Adaptor Assembly in Clathrin-Mediated Endocytosis. Dev Cell. 2019 Aug;50(4):436-446.e5.

63. Kovtun O, Dickson VK, Kelly BT, Owen DJ, Briggs JAG. Architecture of the AP2/clathrin coat on the membranes of clathrin-coated vesicles. Sci Adv. 2020 Jul 24;6(30).

64. Kadlecova Z, Spielman SJ, Loerke D, Mohanakrishnan A, Reed DK, Schmid SL. Regulation of clathrin-mediated endocytosis by hierarchical allosteric activation of AP2. Journal of Cell Biology. 2017 Jan 2;216(1):167–79.

65. Sieghart W, Chiou LC, Ernst M, Fabjan J, M. Savić M, Lee MT. α 6-Containing GABA A Receptors: Functional Roles and Therapeutic Potentials. Pharmacol Rev. 2022 Jan;74(1):238–70.

66. Fan PC, Lai TH, Hor CC, Lee MT, Huang P, Sieghart W, et al. The α6 subunit-containing GABAA receptor: A novel drug target for inhibition of trigeminal activation. Neuropharmacology. 2018 Sep;140:1–13.

67. Bedford P, Hauke DJ, Wang Z, Roth V, Nagy-Huber M, Holze F, et al. The effect of lysergic acid diethylamide (LSD) on whole-brain functional and effective connectivity. Neuropsychopharmacology. 2023 Jul 25;48(8):1175–83.

68. Yang YR, Song S, Hwang H, Jung JH, Kim SJ, Yoon S, et al. Memory and synaptic plasticity are impaired by dysregulated hippocampal O-GlcNAcylation. Sci Rep. 2017 Apr 3;7(1):44921.

69. Martinez MR, Dias TB, Natov PS, Zachara NE. Stress-induced O-GlcNAcylation: an adaptive process of injured cells. Biochem Soc Trans. 2017 Feb 8;45(1):237–49.

70. Wheatley EG, Albarran E, White CW, Bieri G, Sanchez-Diaz C, Pratt K, et al. Neuronal O-GlcNAcylation Improves Cognitive Function in the Aged Mouse Brain. Current Biology. 2019 Oct;29(20):3359-3369.e4.

71. Kim TH, Sung SE, Cheal Yoo J, Park JY, Yi G su, Heo JY, et al. Copine1 regulates neural stem cell functions during brain development. Biochem Biophys Res Commun. 2018 Jan;495(1):168–73.

72. Sulaiman Alsaadi M. Role of DAPK1 in neuronal cell death, survival and diseases in the nervous system. International Journal of Developmental Neuroscience. 2019 May 11;74(1):11–7.

73. Hollville E, Romero SE, Deshmukh M. Apoptotic cell death regulation in neurons. FEBS J. 2019 Sep 12;286(17):3276–98.

74. Jędrzejewska-Szmek J, Blackwell KT. From membrane receptors to protein synthesis and actin cytoskeleton: Mechanisms underlying long lasting forms of synaptic plasticity. Semin Cell Dev Biol. 2019 Nov;95:120–9.

75. Goto-Silva L, Ayad NME, Herzog IL, Silva NP, Lamien B, Orlande HRB, et al. Computational fluid dynamic analysis of physical forces playing a role in brain organoid cultures in two different multiplex platforms. BMC Dev Biol. 2019 Dec 7;19(1):3.

76. Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ, et al. The PRIDE database and related tools and resources in 2019: improving support for quantification data. Nucleic Acids Res. 2019 Jan 8;47(D1):D442–50.

77. Cox J, Mann M. 1D and 2D annotation enrichment: a statistical method integrating quantitative proteomics with complementary high-throughput data. BMC Bioinformatics. 2012 Nov 5;13(S16):S12.

78. Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, et al. The Perseus computational platform for comprehensive analysis of (prote)omics data. Nat Methods. 2016 Sep 27;13(9):731–40.

79. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al. Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. Nat Commun. 2019 Apr 3;10(1):1523.

80. Szklarczyk D, Morris JH, Cook H, Kuhn M, Wyder S, Simonovic M, et al. The STRING database in 2017: quality-controlled protein–protein association networks, made broadly accessible. Nucleic Acids Res. 2017 Jan 4;45(D1):D362–8.

81. Kohl M, Wiese S, Warscheid B. Cytoscape: Software for Visualization and Analysis of Biological Networks. In 2011. p. 291–303.

82. Chin CH, Chen SH, Wu HH, Ho CW, Ko MT, Lin CY. cytoHubba: identifying hub objects and sub-networks from complex interactome. BMC Syst Biol. 2014 Dec 8;8(S4):S11.

83. Goto-Silva L, Martins M, Murillo JR, Souza LRQ, Vitória G, Oliveira JT, et al. Quantitative profiling of axonal guidance proteins during the differentiation of human neurospheres. Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics. 2021 Aug;1869(8):140656.

84. Ferreira TA, Blackman A V, Oyrer J, Jayabal S, Chung AJ, Watt AJ, et al. Neuronal morphometry directly from bitmap images. Nat Methods. 2014 Oct 29;11(10):982–4.