A deep-learning strategy to identify cell types across species from high-density extracellular recordings

Maxime Beau[*,1], David J. Herzfeld[*,2], Francisco Naveros[*,3,4], Marie E. Hemelt[*,2], Federico D'Agostino[*,1], Marlies Oostland[*,1,5], Alvaro Sánchez-López[*,3], Young Yoon Chung[1], Michael Maibach[1], Hannah N. Stabb[1], M. Gabriela Martínez Lopera[1], Agoston Lajko[1], Marie Zedler[1], Shogo Ohmae[3], Nathan J. Hall[2], Beverley A. Clark[&,1], Dana Cohen[&,6], Stephen G. Lisberger[&,2], Dimitar Kostadinov[&,1,7], Court Hull[&,2], Michael Häusser[&,1], Javier F. Medina[&,3]

[1]Wolfson Institute for Biomedical Research, University College London, London, UK
[2]Department of Neurobiology, Duke University School of Medicine, Durham, NC, USA
[3]Department of Neuroscience, Baylor College of Medicine, Houston, TX, USA
[4]Department of Computer Engineering, Automation and Robotics, Research Centre for Information and Communication Technologies, University of Granada, Granada, Spain
[5]Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam, the Netherlands
[6]The Leslie and Susan Gonda Multidisciplinary Brain Research Center, Bar-Ilan University, Ramat-Gan, Israel
[7]Centre for Developmental Neurobiology, King's College London, London, UK

[*]Equal contribution as first authors
[&]Equal contribution as senior authors

Figures: 8, total; 8, color

Word counts: Abstract, 150; Introduction, 538; Discussion, 1,597

Keywords: cerebellum, cerebellar cortex, cell-type identification, circuit mapping, Neuropixels, variational autoencoder, machine learning

Acknowledgements: We thank Bonnie Bowell, Soyon Chun, Wenjuan Kong, Caroline Reuter, and Stefanie Tokiyama for technical support. Funding provided by: NIH grants R01-NS112917 (SGL, JFM, CH), K99-EY030528 (DJH), R01-NS092623 (SGL), R01-MH093727 (JFM); ERC AdG 695709 (MH); Wellcome Trust PRF 201225/Z/16/Z and 224668/Z/21/Z (MH) and 225951/Z/22/Z (DK); EMBO ALTF 914-2015 (DK); European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 844318 (MO) and No 891774 (FN); SYNCH project funded by the European Commission under the H2020 FET Proactive program-Grant agreement ID 824162 (DC)

Correspondence to:

Javier F. Medina, PhD
Department of Neuroscience, S709
Baylor College of Medicine
Houston, TX 77030
Email: jfmedina@bcm.edu
Phone: (713) 798-8141

<u>Abstract</u>

High-density probes allow electrophysiological recordings from many neurons simultaneously
across entire brain circuits but fail to determine each recorded neuron's cell type. Here, we
develop a strategy to identify cell types from extracellular recordings in awake animals, opening
avenues to unveil the computational roles of neurons with distinct functional, molecular, and
anatomical properties. We combine optogenetic activation and pharmacology using the
cerebellum as a testbed to generate a curated ground-truth library of electrophysiological
properties for Purkinje cells, molecular layer interneurons, Golgi cells, and mossy fibers. We
train a semi-supervised deep-learning classifier that predicts cell types with greater than 95%
accuracy based on waveform, discharge statistics, and layer of the recorded neuron. The
classifier's predictions agree with expert classification on recordings using different probes, in
different laboratories, from functionally distinct cerebellar regions, and across animal species.
Our approach provides a general blueprint for cell-type identification from extracellular
recordings across the brain.

The nervous system comprises many molecularly, anatomically, and physiologically defined cell
types[1–6]. Powerful modern molecular techniques now have revealed multiple sub-types even
within known anatomical cell classes[7–13]. Identification of cell type at multiple levels will be
crucial to understand how the brain works and to develop selective, targeted therapeutics for
brain dysfunction. Therefore, it is crucial to develop strategies to determine cell type and to
cross-reference different formulations of cell type across levels of analysis[3,5,6,12,14,15].

With the advent of high-density multi-contact recording probes[16,17], it is now possible to record
from hundreds of neurons simultaneously and characterize their activity during specific,
quantified behaviors. Simultaneous large-scale electrophysiological recordings coupled with
cell-type identification *in vivo* would facilitate characterization of circuit-level processing in the
service of behavior. Yet, identification of cell type is a particularly difficult challenge for
extracellular recording technologies that cannot access the transcriptional or anatomical
properties of neurons[18]. Efforts to classify neurons based on specific features of their spike
waveform and firing statistics have not proven robust across laboratories[19,20]. Moreover,
optogenetic approaches to cell-type identification[21–24] currently are routine only in mice and
bring the technical challenges of (i) off-target expression of opsins[25], (ii) disambiguating direct
responses versus those due to recurrent connectivity within circuits[26], and (iii) the ability to
target only one or two cell types at a time in a given preparation[27].

We assembled a collaboration of four laboratories with the single-minded goal of enabling cell-
type identification solely from extracellular recordings in awake animals by developing a
strategy that could scale across labs, probes, species, and in the future maybe across brain areas.
We chose to pioneer the strategy in the cerebellar cortex, which provides key advantages.
Specifically, the cerebellum has a crystalline architecture with well-defined neuronal
connectivity and a small number of anatomically-defined cell types[1,28] that are consistent across
species[29,30], allowing direct comparison of recordings in monkeys, mice and other species. The
cerebellum has a range of neuron sizes from among the smallest and most densely packed
(granule cells) to the largest (Purkinje cells), allowing us to test the resolution of our recording
approaches. The cerebellum has many spontaneously firing neurons[31–33], some with high
spontaneous rates, allowing us to extract rigorous information about their electrophysiological
properties. Genetically-defined mouse Cre-lines are available for all major cell types in the
cerebellum[34–38], allowing us to leverage optogenetic strategies for cell-type identification[21].
Finally, the cerebellum has a long history of neurophysiological recording[39], allowing us to
reference our measurements and automated cell-type classifications against hard-won human
expertise. Strategies to solve the challenges of cell-type identification in such a testbed should
provide a roadmap for application to other structures, including the cerebral cortex, the
hippocampus, and the basal ganglia.

Our approach succeeded. We created a ground-truth library of identified cerebellar cell types
recorded in unanesthetized mice by combining rigorous spike sorting and unit curation with
identification through combined optogenetic activation and pharmacological synaptic blockade.
We demonstrate that a semi-supervised deep-learning classifier accurately predicts cell type for
the ground-truth library based on the waveform, discharge statistics, and anatomical layer of the
recording. Importantly, the classifier identifies cell type with high confidence in a high fraction
of expert-labeled cerebellar recordings from two different laboratories, in behaving mice and
macaque monkeys.

Results

General approach
We start by creating a ground-truth library of extracellular recordings from neurons whose cell
type is established unequivocally. The analysis and classification pipeline (Figure 1) begins with
several data curation steps to ensure high-quality recordings and allow us to characterize with
high confidence each neuron's waveform, resting discharge properties, and anatomical location.
We use well-characterized mouse Cre-lines to identify cell types by optogenetic activation of
specific cell types in the presence of synaptic blockers. We then develop a semi-supervised deep-
learning classifier with performance evaluated with leave-one-out cross-validation. Finally, we
use the classifier to predict the cell types of an independent dataset of recordings made in mice
and macaque monkeys and we compare the performance of the classifier against cell-type
identification by human experts. Below, we develop the details of our strategy one step at a time.

Figure 1: A strategy for cell type identification from extracellular recordings in neural circuits. The
strategy comprises three steps: data acquisition and curation to build a ground-truth cell type library, selection of
features from the ground-truth library to train a machine-learning based classifier, and tests of the classifier using
additional datasets, including from other species. The first step is to create a ground-truth library of cell types
based on optogenetic activation of genetically-defined neurons during electrophysiological recordings in awake
mice. Neurons in the ground-truth library must be activated directly, as confirmed by a combination of synaptic
blocker pharmacology and electrophysiological criteria, followed by careful data curation. The second step is to
identify features in the dataset that can be used to train a semi-supervised deep-learning classifier. The third step
is to test the generality of the classifier by asking it to predict cell types in independent datasets of expert-
classified recordings from mice and monkeys.

Multi-contact probe recordings and data curation
We develop and deploy the general strategy for cell-type classification (Figure 1) in the
cerebellum, based on ground-truth recordings with Neuropixels probes in two laboratories
(Häusser and Hull labs). In the cerebellar cortex, morphologically distinct cell types reside in
different layers (Figure 2A). Purkinje cells comprise a monolayer and extend their planar
dendrites through the molecular layer. Molecular layer interneurons reside across the extent of
the molecular layer and include basket cells that innervate the Purkinje cell's soma and stellate
cells that innervate the Purkinje cell's dendrites. The granule cell layer includes mossy fiber
terminals, Golgi cells, and granule cells. Recent studies have identified other, less-common cell

types in the different layers[37], but we have elected to focus on the main cell types from the
cerebellar circuit (Figure 2A).

Figure 2: Curation of Neuropixels recordings in the mouse cerebellar cortex.

A. Schematic diagram of the canonical cerebellar circuit.

B. Traces on the left show example simple spikes (light blue) and complex spikes (black) in a Purkinje cell. Histogram on the right documents a complex-spike-triggered pause in simple spikes.

C. Example recordings from many channels of a Neuropixels probe with magenta, blue, black, and green used to highlight a single unit recorded in the molecular layer, a Purkinje cell's simple spikes, the same Purkinje cell's complex spikes, and a unit recorded in the granule cell layer.

D. Comparison of example histology labeled with DiI and Hoechst to show the excellent agreement of histological determination of layers and the layers predicted by Phyllum from the electrical recordings.

Different colors on the Neuropixels schematic show: magenta, molecular layer; blue, Purkinje cell layer; green, granule cell layer; gray, unknown layer; black, outside cerebellar cortex.

E. Autocorrelograms plotting a neuron's firing rate as a function of time from one of its own trigger spikes for two neurons with very few refractory period violations (RPVs). Note that the spike counts in the autocorrelograms have been divided by the width of the bin so that the y-axis is in spikes/s.

F. Analysis of quality of isolation as a function of time during a recording session. From top to bottom graphs show the percentage of refractory period violations, the estimated percentage of missed spikes, and spike amplitude. Horizontal dashed lines show thresholds for acceptance. Gray regions show periods that were rejected from analysis. Blue, green, and red symbols indicate spikes that came from intervals that had too many missed spikes, acceptable isolation, and too many refractory period violations. Marginal histograms on the right show the distribution of spike amplitudes to document clipping at the noise level in the blue histogram that would be cause for rejection of a time interval.

G. Example recording traces and spatial footprint of a representative recording with a signal-to-noise ratio (SNR) of 9.33, with the waveforms numbered according to their channel. Asterisk (*) denotes the channel with the largest peak-to-trough amplitude, used to compute the SNR.

H. Distribution of percentage of refractory period violations across neurons accepted to the ground-truth library.

I. Distribution of estimates of percentage of spikes that were missed across neurons accepted to the ground-truth library.

J. Distribution of signal-to-noise ratios on the channel with the largest-amplitude waveform across neurons accepted into the ground-truth library.

Purkinje cells are the one cell type in the cerebellum that allows ground-truth identification from
its extracellular electrical signature. Purkinje cells show two types of action potentials (Figure
2B, left panel): "simple spikes" that fire at high rates and "complex spikes" driven directly by
climbing fiber input[40–42]. Complex spikes occur only at ~1 Hz, and trigger a characteristic 10-50
ms pause in simple spikes[43]. Thus, Purkinje cells can be identified unequivocally, and admitted
into the ground-truth library, if they show a pause in a complex-spike-triggered histogram of
simple-spike firing (Figure 2B, right panel).

Recordings with Neuropixels probes detect neural activity on many of the 384 channels and
spike sorting yields many units including non-Purkinje cells. The magenta waveforms in Figure
2C arise from a neuron in the molecular layer that would be a candidate to be a molecular layer
interneuron. The green waveforms come from a neuron recorded in the granule cell layer that
could be a mossy fiber, a Golgi cell, or a granule cell. The blue and black waveforms are the
simple spikes and complex spikes of an identified Purkinje cell.

Given that the soma of each cell type resides in one of the three different layers of the cerebellar
cortex, the first step in our analysis pipeline was an objective procedure to identify the layer of
each recording. The cerebellum is a foliated structure so that a single penetration with a
Neuropixels probe usually records from neurons in multiple repetitions of each of the 3 layers of
the cerebellar cortex. For example, the recording trajectory documented with DiI staining in
Figure 2D crossed 3 molecular layers, 5 Purkinje cell layers, and 3 granule cell layers. We
assigned each channel to a layer using Phyllum, a Phy plugin that analyzes recordings across the
channels on a probe to infer the layer recorded by each channel (see ***Methods***).

The layer structure inferred by Phyllum agreed well with histological data based on simultaneous
DiI and cell body staining (Figure 2D). We validated Phyllum across 21 histologically confirmed
penetrations and found that its conclusions agree with the histology at 99, 95, and 98% of 776,
367, and 1140 recording sites respectively in the molecular, Purkinje cell, and granule cell layers.

The layer assignments from Phyllum were also consonant with the finding from single electrodes
of (i) clear complex-spike activity and well-isolated simple spikes in the Purkinje cell layer, (ii)
relative silence and abundant dendritic Purkinje cell complex spikes[44,45] in the molecular layer,
and (iii) a jungle of high-intensity activity with many units in the granule cell layer.

We next ensured that each unit we admitted for further analysis was a well-isolated single neuron
with credible waveform and resting discharge properties, two of the three features we ultimately
would use, along with layer, to classify units. We manually curated the output from Kilosort2
with Phy and subsequently performed automated quality checks to ensure the quality of isolation
and the veracity of the waveforms and resting discharge statistics of neurons that would become
part of our ground-truth library. We strove to ensure that we neither missed many spikes from
the neuron under study nor included electrical artifacts or spikes from neighboring neurons.

● We analyzed the refractory periods from each isolated neuron to assess the level of
contamination from other neurons or noise[46]. The examples of autocorrelograms in Figure 2E
have vanishingly small numbers of refractory period violations and respectively represent the
mean (0.25%) and median (0.01%) in our dataset. We rejected from the ground-truth library
autocorrelograms with greater than 5% period violations (Figure 2F, red symbols and
histogram). Almost all accepted neurons had fewer than 1% refractory period violations with
a mean of 0.25% (Figure 2H).

● We estimated the number of missed spikes by fitting the spike amplitude distribution with a
Gaussian function and quantifying the fraction of the area under the curve that was clipped at
noise threshold[47,48] (Figure 2F). We estimated that few spikes were missed if the distribution
of spike amplitudes for the entire recording was continuous and not clipped at noise
threshold. In Figure 2F, we estimated that more than 5% of spikes were missed in the first
~150 s of the recording (blue symbols and histogram), causing us to exclude those intervals
from further analysis. Among the recordings we accepted, the percentage of missed spikes
averaged 0.26% and almost all neurons showed fewer than 1% missed spikes (Figure 2I).

The requirement for few violations of the refractory period and small numbers of missed spikes
ensured that the units we accepted had high signal-to-noise ratios. The mean signal-to-noise ratio
in our accepted sample, measured as the signal-to-noise ratio on the channel with the largest unit
potential, was 9.3, almost identical to that of the example recording in Figure 2G. Over 90% of
the neurons had signal-to-noise ratios larger than 4 (Figure 2J).

We identified and resolved two other issues that impaired consistent and reliable estimates of
waveform. The _first_ issue is related to an on-board hardware high-pass filter on Neuropixels
probes. The filter distorts the shape of waveforms and therefore hinders comparison across
recordings made when the filter was on versus off. We used the technical description of the
analog filter in the Neuropixels documentation to apply an equivalent digital filter to data
recorded with the filter disengaged or using other probes in monkeys (Supplementary Figure 1).
The _second_ issue concerns temporal alignment of individual spikes, which is unreliable in
Kilosort's output when the signal-to-noise ratio is low to medium or when a unit drifts across
channels during a recording session. We resolve the alignment issue with an iterative procedure
we called "drift-shift-matching" to minimize waveform distortion from the averaging of
individual action potentials (see Supplementary Figure 1 and ***Methods***).

Combination of optogenetics and pharmacology for ground-truth cell-type identification
For optogenetic activation, we combined genetic and viral approaches to cause expression of an
opsin in a specific cell type, thereby allowing these neurons to be selectively activated by light
and identified by photostimulation[21]. We added synaptic blockers to our experimental
preparations to ensure that light activates an opsin-expressing neuron directly, and not indirectly
by optogenetic activation of its pre-synaptic inputs. The standard criterion of short-latency
activation by optogenetic stimulation[49] (e.g., <10 ms) is inadequate on its own. We found ample
examples of short-latency responses (some <5 ms) that disappeared with synaptic blockade[21].

We introduced Neuropixels probes into the lateral cerebellar cortex or the vermis of mice
expressing opsins (usually Channelrhodopsin-2, ChR2) in specific cell types and allowed the
probe to settle at a location where we recorded activity across much of its length. As illustrated
in Figure 3A, we performed experiments with optogenetic stimulation in multiple phases. In a
baseline phase, we recorded spontaneous activity. In a *control phase*, we applied light externally
to the cerebellum to activate opsins in the cell types that expressed them. We also introduced a
tapered optic fiber that ran alongside the recording probe in some experiments, to deliver light in
closer proximity to cells that expressed opsins. In an *infusion phase*, we continued to deliver
light to the cerebellum while we added synaptic blockers (see ***Methods***) to the surface of the
cerebellum. In the *blockade phase*, when the synaptic blockers had permeated well into the
tissue, we assayed neurons for direct responses to optogenetic stimulation. The approach of
applying synaptic blockers on the surface of the cerebellum, instead of trying to inject them deep
into the tissue, has the advantage of preserving the integrity of the tissue but the disadvantage of
relatively slow diffusion.

We accepted neurons as activated directly by photostimulation only if we had strong evidence
that they were within the locus of successful synaptic blockade and they continued to have
reliable, short-latency responses to light. To determine whether synaptic blockade was effective
at a given recording depth, we evaluated recordings at or below that depth and looked for any of
the indications in the top row of Figure 3B:

1. Loss during the *blockade phase* of responses to optogenetic stimulation present in the *control*
phase (Figure 3B, top row, left). Neurons were excluded from the ground-truth library if they
retained their response in the *blockade phase* but were outside the region of synaptic
blockade (Figure 3B, bottom row, left).

2. Putative mossy fibers with loss of negative afterwaves in the *blockade phase* (Figure 3B, top
row, middle). Here, we rest partly on prior evidence that the negative afterwave is a post-
synaptic response of granule cells[50,51] and that analogous negative afterwaves have been
shown to correspond to post-synaptic responses in other brain regions[52]. Also, our finding of
the effect of synaptic blockade on the negative afterwave in recordings from single putative
mossy fibers provides the strongest evidence to date that the negative afterwave represents
post-synaptic depolarization.

3. Substantial changes in a neuron's autocorrelogram or "coefficient of variation 2" (CV2),
usually due to an increase in regularity caused by a shift from synaptically- and intrinsically-
driven spiking to purely intrinsically-generated spiking[31] (Figure 3B, top row, right).

To illustrate our strategy, we provide detailed examples from one experiment in a transgenic
mouse line that expresses ChR2 in mossy fibers[37,53] (Thy1-ChR2 line 18). Here, some neurons
lost their responses to optogenetic activation with synaptic blockade (Figure 3C), while others
recorded nearby retained their responses (Figure 3D). We evaluated the effect of synaptic
blockade along the electrode penetration that yielded these two units (Figure 3E, region above
the dashed line on the Neuropixels schematic) to confirm that the neural responses in Figure 3D
were activated directly by optogenetic stimulation. At sites near and deeper than the neuron in D,
neurons lost their responses with synaptic blockade, indicating that they were within the region
of successful blockade. Still deeper in the penetration, we recorded two putative mossy fiber
waveforms that retained their negative afterwave with synaptic blockade, indicating that they
were outside the region of successful blockade. Finally, the extracellular waveforms of the
activated neuron were constant across the entire experiment (Figure 3F), indicating that we had a
stable recording. Therefore, we concluded that the neuron in Figure 3D was an optogenetically-
activated neuron, namely a mossy fiber.

Figure 3: Strategy for ground-truth identification of cell type.

A. Schematic showing the sequential phases in an experiment designed to test for optogenetic activation in the presence of synaptic blockers.

B. Examples of the results used to verify the region of synaptic blockade. Examples above versus below the horizontal dashed line were taken as evidence for versus against blockade at that site. From left to right, we assayed the effect of blockade on the response to optogenetic stimulation, the negative afterwave of a

> putative mossy fiber waveform, and the discharge statistics defined by autocorrelograms and the value of CV2.
>
> C. Raster and peri-stimulus time histogram for a neuron that lost its response to optogenetic stimulation with synaptic blockade. Trial numbers on the y-axis align with the cartoon showing the periods in the experiment to the right of D. Black versus orange histograms show responses before versus during synaptic blockade. Blue shading indicates the time of photostimulation.
>
> D. Same as C except for a neuron that retained its response to optogenetic stimulation during synaptic blockade.
>
> E. Example of how we determined whether the recordings in C and D were within the region of synaptic blockade. The cartoon schematizes a Neuropixels probe, the top histograms on the right show sites that were within the region of blockade because they lost their responses to optogenetic stimulation, and the lower waveforms show mossy fibers that were below the region of blockade because they retained their negative afterwaves.
>
> F. Spatial footprint of the neuron in D. Black, orange, and blue traces show the similarity of the waveforms recorded during the baseline period, during synaptic blockade without optogenetic stimulation, and during synaptic blockade with optogenetic stimulation.
>
> G. Distribution of neural response latencies to optogenetic stimulation of directly-activated neurons in presence of synaptic blockade.

Ground-truth neurons generally responded to optogenetic stimulation with latencies shorter than 5 ms (Figure 3G). However, we recommend against using latency as the sole criterion to accept or reject neurons as optogenetically-activated[21]. Latency is strongly dependent on illumination intensity and the density of opsin expression. We frequently observed short-latency responses to optogenetic stimulation that disappeared with synaptic blockade, for example the neuron in Figure 3C where the latency was less than 3 ms. Also, the neuron in the top row of Figure 3B responded to optogenetic stimulation with a latency shorter than 5 ms, but lost its response with synaptic blockade, indicating that it was driven by synaptic activation rather than direct optogenetic stimulation.

Strategy to mitigate off-target expression in transgenic mouse lines

To varying degrees, off-target expression is a common feature of transgenic mouse lines. Often, there is no 'clean' line available where expression is limited to a given cell-type of interest. Our strategy to obtain ground-truth cell-type identification despite off-target expression was to (i) characterize the anatomical specificity of opsin expression for all mouse lines under study and (ii) combine identification of the recording layer based on Phyllum with optogenetic activation in the confirmed presence of synaptic blockers to establish cell-type unambiguously.

The problem of off-target expression was most pronounced in the GlyT2-Cre line used previously to image activity in Golgi cells[35]. The GlyT2-Cre line has substantial off-target expression in molecular layer interneurons and, very occasionally, Purkinje cells (Figure 4A, Supplementary Figure 2): the relative density of molecular layer interneurons was higher than that of Golgi cells (Figure 4B, 79 vs. 20%). Accordingly, we recorded neurons directly responsive to optogenetic stimulation in both the granule cell layer and the molecular layer (Figure 4C). We used Phyllum to identify the recording layers and labeled units in the granule cell layer that were directly activated by optogenetic stimulation as Golgi cells (Figure 4C, green PSTH). We labeled activated units in the molecular layer as molecular layer interneurons (Figure 4C, magenta PSTH).

Figure 4: Analysis and mitigation of off-target expression in mouse optogenetic lines.

A. Double stained section of cerebellum in the GlyT2-Cre line showing expression in both Golgi cells in the granule cell layer and molecular layer interneurons. Red arrows point to cells that express Td-Tomato. Blue cells express parvalbumin (PV). MoL, molecular layer; PCL, Purkinje cell layer; GCL, granule cell layer.

B. Cartoon of cerebellar circuit and histogram showing density of TdT-positive somata in each of the three layers in a GlyT2-Cre mouse: GoC, Golgi cell; GrC, granule cell; PC, Purkinje cell; MLI, molecular layer interneuron; CF, climbing fiber; MF, mossy fiber.

C. Representative recordings from a Neuropixels probe using optogenetics to activate neurons that express opsins in the GlyT2 line. Magenta, blue, and green waveforms on the right show the spatial footprint of neurons in the MoL, PCL, and GCL. Histograms below the voltage traces show that both the MoL and GCL layer neurons were activated by optogenetic stimulation at the time indicated by the blue shading.

Other mouse lines also showed some off-target expression. For example, the Math1-Cre line
used to label granule cells was generally specific (Figure 4D) but exhibited rare labeling of
Purkinje cells. The c-kit-Cre line also labeled a small number of Golgi cells[34], and the Nos1-Cre
line exhibited occasional labeling of non-neuronal cells in addition to molecular layer
interneurons (Supplementary Figure 2). By contrast, other lines we used were cleaner, such as
the Thy1-ChR2-YFP line 18 and Pcp2-Cre lines used to label mossy fibers and Purkinje cells,
respectively (Supplementary Figure 2). Crucially, we did not observe multiple labeled cell-types
within a single cerebellar layer in any of our lines. Thus, the combination of an identified layer
with direct optogenetic activation (Figure 4E) allowed us to disambiguate cell type for all
experiments.

The ground-truth library
Across 188 Neuropixels recordings in two laboratories, we recorded a total of 3652 neurons that
survived the spike-sorting and curation pipeline (Figure 5A). Of these, 562 exhibited a response
to optogenetic stimulation but only 97 passed our rigorous criteria for direct rather than synaptic
activation based on reliable, short-latency responses in the presence of synaptic blockade. We
added the simple spikes and complex spikes of 62 Purkinje cells identified by a complex-spike
triggered pause in simple spikes. We removed 6 units recorded in Cre-lines with off-target
expression where the layer of the recording was ambiguous, and 13 units that, on final closer
inspection, did not have sufficiently long baseline periods due to intervals that violated our
missed/extra spikes criteria (Figure 2E, F). The resulting ground-truth library contained 202
units: 69 Purkinje cell simple spikes, 58 Purkinje cell complex spikes recorded at the same time
as the simple spikes, 27 molecular layer interneurons, 18 Golgi cells, and 30 mossy fibers. For
comparison with previous reports[20,54,55], Supplementary Figure 4 provides the
electrophysiological signatures of different cell types in our ground-truth library of cell types
using a range of metrics.

In an attempt to obtain ground-truth recordings from granule cells, we made 82 recordings with
Neuropixels probes in mice with the Math1-Cre or BACα6Cre-C lines either crossed to Cre-
dependent ChR2 (Ai32) or injected with AAV to confer ChR2 expression (see **Methods**). We did
record multiple unit activity that was responsive to photostimulation in the region of confirmed
synaptic blockade, but almost all putative single units found by KiloSort failed one or more of
our criteria for good isolation (Figure 2). After careful curation, we retained zero likely granule
cells from 32 recordings in the Hull lab and at most 3 from 50 recordings in the Häusser lab. The
yield of fewer than 0.04 granule cells per recording was significantly lower than for the other cell
types (Figure 5B). We conclude that it is challenging to record from granule cells using the
current generation of Neuropixels probes; our sample is far too small to include them in the
classifier we will develop next. A combination of factors may contribute to the inability to record
regularly from granule cells: their comparatively small size[56,57], the likelihood that they generate
a spatially-restricted closed-field extracellular potential, and the low electrode impedance[58] of
Neuropixels[16] (150 kOhms).

Figure 5: Selection criteria and properties of the ground-truth library of cerebellar cell types.

A. Curation criteria used to decide which neurons to include in the ground truth library, including the numbers that were retained or deleted at each stage of the curation.

B. Histogram showing the number of ground-truth units of each cell type normalized for the number of recordings: MLIs, molecular layer interneurons; GoCs, Golgi cells; MFs, mossy fibers; GrCs, granule cells.

C. Superimposed waveforms for each cell type in the ground truth library. Abbreviations as in B, plus: PC$_{SS}$, Purkinje cell simple spikes; PC$_{CS}$, Purkinje cell complex spikes. The bold trace indicates the neuron that has an example 3D-ACG in Supplementary Figure 5. Waveforms are normalized and flipped to ensure the largest peak is negative (see **_Methods_**).

D. Same as C but showing autocorrelograms of ground-truth neurons. Note that the spike counts in the autocorrelograms have been divided by the width of the bin so that the y-axis is in spikes/s.

E. Failure of traditional measurements of waveform or discharge statistics to differentiate cell types. Each symbol shows Z-scored values of different features from a single neuron; different colors indicate different cell types, per the key in the upper right. Z-scores were computed separately for each feature but across cell types within each feature. Abbreviations as in B.

Armed with a ground truth dataset, the next challenge was to develop an accurate classification
method based on consistent differences in electrophysiological features across cell types[59]. To
maximize the success of classification, we strove to use both waveform[20,60,61] and discharge
statistics[62–64] as features for cell-type classification.

Waveform: We anticipated that the different cellular properties and morphology of different
cell types would lead to different waveforms[60,61,65,66]. Patch clamp recordings *in vitro*
confirmed that biophysical differences across neuron classes are manifest as consistent
variations in the shape of the waveform (Supplementary Figure 3). Yet, as shown by the
comparison of Figure 5B with Supplementary Figure 3, waveforms are much more variable
in extracellular recordings *in vivo* than *in vitro*, and waveforms alone do not cleanly
distinguish cell type.

Discharge statistics: It is common for different cell types to have different discharge
statistics throughout the brain[67,68] and the same is true in the cerebellum of anesthetized
animals[62–64]. In awake animals, discharge statistics are likely to vary across cerebellar regions
and to depend on the specific behavior or sensory input[20,69]. Therefore, a robust
classification strategy should harness additional information that normalizes for the factors
that contribute to variation in awake animals.

Cell-type identification from a semi-supervised deep-learning classifier
Our deep-learning classifier strategy takes advantage of the rich information contained in the
diverse waveforms and firing statistics in the ground-truth library (Figure 5C, D), along with the
layer information that also provides information about cell type. Rather than using a potentially
biased set of investigator-chosen measurements from waveform and firing statistics, we chose to
use raw data because they (i) contain richer information, (ii) provide unbiased inputs for cell-
type identification, and (iii) are likely to generalize across regions, tasks, and species. Further,
Figure 5E and Supplementary Figure 4 reveal that it is difficult to guess which specific measures
of waveform and firing statistics would be most informative to successfully distinguish cell types
in awake animals.

We represent spike waveforms as the full time-course of the average, drift- and shift-corrected
waveform on the channel with the largest signal. We represent firing statistics as
autocorrelograms (ACGs) that assess the firing rate of a neuron as a function of time relative to
each spike. Because the traditional two-dimensional-ACG (2D-ACG) is subject to artifacts when
neural firing rate varies across a recording session or in relation to behavior, we developed
"three-dimensional autocorrelograms" (3D-ACGs) that normalize for firing rate (Figure 6A, see
Methods). Supplementary Figure 5 shows example 3D-ACGs for each ground-truth cell type.
We developed a strategy to avoid the potential issue of overfitting that is inherent in a deep-
learning classifier given the high dimensionality of the waveforms and 3D-ACGs and the
relatively small number of training examples in the ground-truth library. To address the
mismatch of training data relative to input dimensionality, we used an unsupervised dimension-
reduction technique[70] that took advantage of 3090 unlabeled single units recorded with
Neuropixels probes during the optogenetics experiments. We trained two variational
autoencoders (Figure 6B), one each for waveforms and 3D-ACGs, to reduce the input
dimensionality to 10 for both features (see *Methods*), thereby minimizing the number of
parameters in the ultimate classifier that needed to be trained *de novo*.

Our classifier (Figure 6C) consists of: (i) a multi-headed, normalized input layer that accepts the
10-dimensional representations of the waveform and 3D-ACG produced by the variational
autoencoders, along with a "one-hot" 3-bit binary code of the unit's cerebellar layer; (ii) a hidden
layer that processes the 3 normalized inputs, and (iii) an output layer with one output unit for

each of the 5 cell types. The value of the output units sums to 1 so that the output of the classifier
is the probability that a given set of inputs are from each of the 5 cell types. We trained the
weights in the classifier on the data in the ground-truth library using gradient descent with a
leave-one-out cross-validation strategy.

Figure 6: Performance of a deep-learning classifier on cell type identification for the ground-truth library.

A. Method for normalizing effects of mean firing rate on firing statistics through three-dimensional autocorrelograms (3D-ACGs). Left graphs show the consensus ACG for an example neuron without regard for firing rate on top and 3 ACGs for different mean firing rates on the bottom. The heatmap on the right plots 10 rows that show 2D-ACGs as heatmaps for 10 different deciles of mean firing rate. Arrows indicate the row in the 3D-ACG for each 2D-ACG.

B. Schematic of autoencoders used in unsupervised learning to reduce the dimensionality of the waveform and 3D-ACG inputs to the classifier.

C. Classifier architecture. Note that we ran the classifier with 10 different initializations for each of the 202 ground-truth units, symbolized by the 202 pages in the classifier.

D. Histograms showing the predictions of the classifier on 10 repetitions of training starting with different initial conditions to develop an estimate of confidence from the means of the probabilities assigned to each cell type.

E. Percentage of units classified as a function of the ratio we chose as a threshold for confidence in the assignment of cell type. Different colors show data for different ground-truth cell types.

F. Confusion matrix showing the agreement between the predictions of the classifier on a single left-out testing unit and the ground-truth cell type of that testing unit. The numbers in each cell indicate the percentage of ground-truth cell types on the y-axis for each prediction of the classifier on the x-axis, where confidence was required to be greater than 2. The rightmost column shows the percentage of ground-truth neurons that received a confidence greater than 2.

G. Same as F, but for neurons in the ground-truth library regardless of confidence, i.e. confidence threshold = 0.

We evaluated not only the accuracy, but also the "confidence" of the output from the classifier.
For each leave-one-out sample (n = 202 ground-truth units), we trained an ensemble of 10
models with random initial conditions. We then averaged the classifier-predicted probability for
each cell type across model instantiations. For the example illustrated in Figure 6D, the
distributions of cell-type probability reveal repeated predictions that the held-out unit was a
Purkinje cell simple spike. The average probability assigned to the simple spike was 0.89 while
the average probability assigned to each of the other cell types was less than 0.1. However, that
need not have been the case: if the data for a given unit were compatible with more than one cell
type, then the classifier might classify the unit as highly-probable to be cell type #1 in one model
instance and highly-probable to be cell type #2 in another instance: the average probabilities
across 10 runs of the classifier might be similar and therefore closer to 0.5 for these two cell
types, indicative of low classifier confidence.

We quantified the classifier confidence for each neuron with the "confidence ratio", computed as
the ratio of the mean probability of the most-likely cell type to the mean probability of the
second-most-likely cell type. As expected, the percentage of ground-truth units that could be
classified decreased as a function of the value of the confidence ratio we chose as the confidence
threshold (Figure 6E). Classifier confidence in general was higher for Purkinje cell simple
spikes, Purkinje cell complex spikes, and mossy fibers compared to Golgi cells or molecular
layer interneurons. Higher confidence thresholds increase the likelihood that cell-type
classification is correct, but also decrease the number of units that get classified. We chose a
confidence threshold of 2 in the remainder of our analysis because it allowed the majority of
neurons to be classified while providing excellent cross-validated classification performance.

The classifier showed impressive accuracy when applied to the units in the ground-truth library.
For each held-out neuron that exceeded the chosen confidence threshold, we assigned it the cell
type that had the highest probability, averaged across the 10 classifier runs. The classifier
assigned cell types to 78% of ground-truth molecular layer interneurons and 74% of ground-truth
Golgi cells at a confidence threshold of 2 (rightmost column of Figure 6F), almost all correctly
as demonstrated by the values of 100% along the diagonal of the confusion matrix (Figure 6F).
The classifier exceeded the confidence threshold for more than 90% of mossy fibers, Purkinje
cell simple spikes, and complex spikes and again it classified nearly all such units correctly. The
accuracy of the classifier degraded without a confidence threshold, but still performed quite well:
it exceeded 90% accuracy on all cell types (Figure 6G). The fact that the classifier was more
accurate when we required higher confidence means that 1) the classifier has a good internal
model of true neuron classes and 2) the choice to set a confidence threshold improves the
performance of the classifier. We note that there is some confusion between neurons in different
layers despite the use of layer as an input because their waveforms and/or autocorrelograms look
ambiguous to a classifier that makes a statistical prediction based on multiple inputs.The
classifier performed less well without layer information, mainly because of greater conflation of

Golgi cells and molecular layer interneurons (Supplementary Figure 6). Molecular layer
interneurons exhibited two distinctly different waveforms with either a small or a large
repolarization phase. The latter were nearly indistinguishable from the waveforms of Golgi cells.
The two types of waveforms in molecular layer interneurons do not map onto the known
subtypes of molecular layer interneurons[71].

Classifier validation of expert-labeled datasets
We next evaluated how well the ground-truth classifier (Figure 7A) generalized by attempting to
predict the cell type for neurons in a sample of expert-classified, non-ground-truth recordings
from mice (Medina lab) and from the monkey's floccular complex (Lisberger lab).

Figure 7: Ground-truth classifier performance on expert-classified datasets from mice and monkeys.

A. Schematic of the ground-truth classifier, repeated from Figure 6C, but now making predictions based on non-ground-truth data from mouse or monkey. The n = 2020 instantiations of the classifier arise from training the classifier 10 times with different initial conditions for each of 202 left-out ground-truth units: 10 x 202 = 2020.

Confidence is a particularly important metric for non-ground-truth data. We took advantage of
the large number of differentially trained and instantiated models from the ground-truth cross-
validation analysis to improve our computation of confidence in the expert-classified datasets.
The ground-truth cross-validation of the ground-truth library resulted in 2020 versions of our
classifier (202 ground-truth units with 10 instantiations per cross-validation). For each unit in the
expert-classified datasets, we averaged the cell-type probabilities predicted by the 2020
instantiations of the classifier and created plots of the probability assigned by the classifier as a
function of cell type (Figure 7B, C). Units appear in exactly one of 5 different plots, chosen
according to the cell type assigned by the classifier as the highest probability, not according to
the expert-assigned cell type. For example, the leftmost graph reports probability versus cell type
for all units that were classified as most probable to be simple spikes of Purkinje cells, colored
according to whether the confidence ratio was below or above 2 (gray versus colored lines). The
collection of confidence plots in Figures 7B and C underscores the points of confusion for the
classifier, perhaps due to subtle differences in the properties of probes, the behaviors performed
by the animals, or the cerebellar sites of recording. For both the mouse data and the monkey data,
classifier confidence was greater than 2 for the majority of units, except that only 45% of the for
units classified by the experts as molecular layer interneurons in the monkey data were classified
"correctly". The gray curves in Figure 7C suggest that the classifier often conflated molecular
layer interneurons and simple spikes in the monkey data, resulting in lower confidence for these
neurons.

The ground-truth classifier agreed with the human experts about the cell types in mice and
monkeys of almost all units that were above confidence threshold, as demonstrated by the large
percentages along the diagonal in the correspondence matrices of Figure 7D and E. Further, the
ground-truth classifier was quite confident about the cell-types in the non-ground-truth data, as
illustrated by the high percentage of units above a confidence threshold of 2 in the rightmost
columns of Figure 7D and E. Here, it is important to explain a subtle difference in the numbers in
Figures 7B and C versus Figures 7D and E. The leftmost graph of Figure 7C indicates that 75%
of the monkey units classified as simple spikes exceeded confidence threshold. Because the units
included in Figure 7C are not necessarily classified by experts as simple spikes, 75% is not
inconsistent with the 37.9% of expert-classified simple spikes that were below confidence

threshold in Figure 7E. The difference in the numbers results from analysis of the space across
orthogonal axes.

Figure 8: Multiple forms of evidence for the similarity of waveforms and resting discharge statistics of different cell types across the ground-truth library and the expert-labeled data from mouse and monkey.

A. Comparison of percentage of classified units as a function of confidence threshold for 3 preparations. Faint colored traces show the same curves for the ground-truth library, from Figure 5E. Bold black and gray traces show results for unclassified mouse and monkey data, respectively.

B. Congruence of the output from the autoencoders for identically labeled ground-truth versus expert-classified neurons across preparations. Each row corresponds to a single ground-truth identified neuron. Each column

corresponds to a single classifier-identified neuron from mouse (left) or monkey (right). Colors at the intersections for each row and column indicate the cosine similarity of the concatenated outputs from the autoencoders for waveform and autocorrelograms, where redder colors indicate greater similarity.

C. Waveforms of different cell-types across laboratories and species. In the first row, waveforms are divided according to ground-truth cell type in mice. In the second and third rows, cell types are divided according to classifier predictions of cell type for non-ground-truth neurons recorded in mice and monkeys.

D. Same as C, except showing 2D-autocorrelograms. Note that the spike counts in the autocorrelograms have been normalized by the width of the bin so that the y-axis is in spikes/s.

The ground-truth classifier identified correctly, with confidence greater than 2, the mouse and
monkey Purkinje cell simple spikes and complex spikes from recordings with a complex-spike-
triggered pause in simple spikes (100/86.9% and 100/100% in mice/monkeys for simple and
complex spikes, respectively. Figure 7F). "Unknown" units, defined as those not identified
definitively as Purkinje cells, were distributed across cell-types by the classifier, as expected
given that they included recordings from all cell types.

Similar properties within cell types across species and cerebellar regions
Three additional analyses indicate that the success of the ground-truth classifier on the expert-
classified data is based on true statistical similarity of the waveforms and firing statistics of each
cell type across datasets. *First*, the percentage of units that we classified with confidence
decreased similarly as a function of the confidence threshold for the two samples of expert-
classified cells (Figure 8A, thick gray and black traces) and the ground-truth data set (Figure 8A,
colored traces). *Second*, analysis of the output of the classifier's autoencoders revealed excellent
agreement between the reduced-dimension representation of expert-classified and ground-truth
data (Figures 8B). Here, the neurons in the ground-truth library plot along the y-axis and the
expert-classified neurons plot along the x-axis. Warmer colors in the heatmap indicate greater
alignment of the 20-dimensional vectors defined by the concatenated outputs of the two auto-
encoders in the classifier. *Third*, inspection of the waveforms (Figure 8C) and the 2D-ACGs
(Figure 8D) reveals impressive similarity across the ground-truth data, the non-ground-truth
mouse data, and the monkey recordings. Here, we have included only the neurons that were
classified with confidence greater than 2. The only real exception to the visual impression of
similarity is a few of the 2D-ACGs for the ground-truth Golgi cells. The similarity of resting
discharge properties across preparations also appears in the 3D-ACGs (Supplementary Figure 4).

The large fraction of non-ground-truth neurons that can be classified with confidence, and the
agreement with the experts, is unexpected evidence that the properties of different cerebellar cell
types are consistent across species and cerebellar regions. It certainly was possible, *a priori*, that
a different outcome might have emerged because of genuine differences in waveform or
discharge statistics across species and cerebellar regions, differences in data collection and
analysis across labs, or a failure of rigor in our procedures for curating the ground-truth and
expert-classified data. We anticipate that Figure 8 will serve as a useful resource for other
cerebellar labs to be confident of their own rigor as they assign cell types in their own data.

Discussion

Identification of cell type from *in vivo* extracellular recordings is a fundamental issue in systems
neuroscience[5,6,20,55,60,62,63,68,72–75]. Our approach delivers a highly-reliable ground-truth library of
the electrophysiological properties of cerebellar cell types in awake mice based on identification
through optogenetic stimulation in the presence of synaptic blockers. The ground-truth library
consists of the waveform of the electrical recording, the statistics of the spike train, and the layer
of the cerebellum where we recorded each unit, information that comes from well-isolated neural
recordings using high-density multi-contact probes. Our semi-supervised deep-learning classifier
performs well in identifying the cell types in the ground-truth library, while also reporting its
confidence in each identification. The internal representations in the classifier reveal similar
statistics in the ground-truth library and in independent datasets of expert-classified recordings
from the mouse and monkey cerebellum. The cell types predicted by the classifier for the mouse
and monkey data agree with the experts' assessments. We are encouraged by the accuracy and
precision of our classifier and expect that it will be possible in the future to align the cell type
obtained from extracellular recordings with that obtained from other levels of analysis, including
anatomical and molecular fingerprints.

The strategy we developed may be more useful and important than the exact classifier. Our goal
at the outset of our project was to achieve cell-type identification from extracellular recordings in
the cerebellar cortex across laboratories and species. We think that the strategies inherent in our
classifier, and the classifier itself, can be used with confidence by any cerebellar recording lab
that is curating their electrophysiological data with sufficient rigor. However, we also point out
that a failure of rigorous curation will lead to noisy and unnecessarily variable inputs to the
classifier and will contaminate the cell-type identifications provided as its output[61].

Others have attempted to identify the distinct signatures of discrete populations of cerebellar
cortical neurons[20,62–64,76,77]. Past attempts identified either (i) neurons of interest by qualitative
agreement with spiking signatures found *in vitro*[64] or (ii) recordings in anesthetized preparations
with neurons identified anatomically via juxtacellular labeling[20,58,62,63,76,77]. Our recordings in
awake and behaving mice demonstrate large variance in the discrete metrics used for
summarizing spiking activity both within and across ground-truth classes (Figure 5E,
Supplemental Figure 4). Thus, classification schemes reliant on a finite set of specific features
probably will not well generalize to other tasks or regions[60], or from anesthetized to behaving
preparations[20]. We think our approach is more likely to generalize because it leverages the
informativeness of the full waveform[60,61] and 3D-ACG to classify cell types across a wide range
of behaviors and stimulus-driven responses.

Several features of the strategy embedded in our classifier were critical to its success:

● *Raw waveforms and 3D-ACGs*. Raw features are an unbiased input[60] and allow the classifier
to take advantage of extensive information in waveform[60,61] and discharge statistics. 3D-
ACGs normalize for variations in firing rate and create a statistic that can be compared across
cerebellar areas, experimental tasks, and species. Similarly, the choice to use single-channel
waveforms allows the classifier to generalize across electrode types. We think our strategies
are likely to generalize because they use raw features that can be measured readily in other
brain areas.

● *Mitigation of overfitting*. We developed a semi-supervised[78–80] deep-learning strategy (see
Methods) to train the classifier with a relatively small number of ground-truth neurons. The
unsupervised training of variational autoencoders reduces the chances of overfitting[81]. The
use of a large unlabeled dataset to train the autoencoders also reduces the chances of
overfitting by ensuring that their architecture was designed independently from the ground-
truth dataset. Successful predictions of cell type that agree with two expert-classified datasets
supports the generalizability of the classifier on other data. Our choice of how to mitigate
overfitting should allow our strategy to generalize to other datasets where the dimensionality
of the inputs is high and the number of ground-truth neurons is comparatively small.

● *Confidence*. We were particularly cognizant of making our classifier trustworthy. To do so,
we established confidence by training multiple models on the same data[82] and by using a
Bayesian method to calibrate confidence at the single model level[83,84] (see ***Methods***). By
requiring confidence above a given threshold[85,86], we improved the accuracy of the model on
the ground-truth data as well as for non-ground-truth recordings. The ability to choose a
confidence threshold allows the user to balance whether to include all neurons even if some
cell type assignments might be incorrect or to include fewer neurons with greater certainty in
the cell type assignment.

The classifier was more successful when it included layer information as an input. With layer
information, it classified a higher percentage of the units and, in the ground-truth data, classified
them with greater accuracy. However, the use of layer as an input does not make classification
trivial. Rather, it creates a platform that will become even more useful as we are able to achieve
ground-truth identification of other cell-types in the cerebellum, for example of granule cells
with improved recording probes. Also, because waveform and firing statistics are necessary to
distinguish cells that are in the same layer, the classifier makes a statistical decision about cell
type rather than relying solely on layer for cell identification[87]. Layer is defined in a specific way
for the cerebellum[1], but we think of layer information more generally as a specific example of
"local electrical properties". We imagine that there are other ways to quantify those properties,
for example LFPs and current-source-density analysis[88,89], that will work in brain areas without a
laminar structure.

The biggest challenge, and our bigger goal, is to see deployment of the strategy outlined here in
other brain areas. The use of layer information to improve classification should be relevant to
other structures – cerebral cortex[90], hippocampus[91], superior colliculus[92] – that have layers with
measurable local electrical properties. We also think that the strategies used in our classifier
enable generalization by showing how to reduce the dimensions of raw data used as inputs while
mitigating the challenges of small numbers of neurons in training sets. We hope that application
of our strategy in other brain areas will enable cell-type identification from extracellular
recordings, a key element in our collective long-term goal of understanding how neural circuits
work and how they generate behavior.

Limitations of the study
One class of limitations is related to our procedures for data collection and curation. While we
were scrupulous about spike sorting and criteria for data inclusion, we relied on indirect
measures and logic to determine whether a recording was within the region of synaptic blockade.
Any errors might have allowed inclusion of certain mis-identified cell types due to off-target

opsin expression. We assessed off-target expression thoroughly, revealing, for example,
extensive off-target expression in the GlyT2 line[35,93,94]. Less obvious off-target expression could
have escaped our histological analysis and introduced a small fraction of incorrectly-identified
cells, for instance in the Math1 line[95].

A second class of limitations is related to the ground-truth classifier's performance for labeling
neurons outside of those we focused on in this study. While we trained the classifier on 5
identified cell types, other neuronal subtypes exist in the cerebellar cortex[28]. We may have
recorded from the rare cell types in our expert-labeled datasets. Were they misclassified as one of
the 5 cell types in our ground-truth library? Or were they correctly recognized as "other" cell
types and relegated to the ~15-20% of neurons that failed to reach confidence threshold? In due
course, we expect to be able to augment our ground-truth library with unipolar brush cells[96],
candelabrum cells[97,98], and other Purkinje layer interneurons[99,100] as ever more specific Cre lines
and viruses become available. We anticipate that advancements in recording probes with higher
impedance and/or more dense recording sites will enable reliable recordings from granule cells.
If waveforms, firing properties, and layer prove insufficient to segregate identity as other cell
types are incorporated, we anticipate that additional information about synaptic and electrical
connectivity can further improve the accuracy of our classifier.

A third possible limitation is related to the electrophysiological features we chose as inputs for
our classifier. Because our goal was to identify cell type based on extracellular recordings with
increasingly popular high-density probes, we used waveform, discharge statistics, and layer as
inputs. The use of discharge statistics might be problematic in structures with little or no
spontaneous discharge in some cell types, although 3D-ACGs enable assessment of discharge
statistics from firing related to sensory inputs or behavior. Also, our use of physiological
characteristics as inputs to the classifier limits our ability to align electrophysiological cell-type
identifications with those provided by single-cell RNAseq[101–103], juxta-cellular labeling[62,63], or
combinations of *in vivo* recording and single-cell imaging[104]. For example, extracellular
electrophysiology cannot 'mark' recorded cells for post-hoc analysis of molecular identity using
approaches such as RNAseq. However, our approach affords several advantages over recording
methods that are compatible with genetic profiling but sacrifice experimental throughput, tissue
accessibility, temporal resolution, and/or the number of cell-types that can be simultaneously
recorded and identified. Thus, it is necessary to weigh the trade-offs of different strategies for
cell-type identification according to the particular experimental question at hand.

Despite its limitations, our study demonstrates a strategy that allows different cell types to be
identified robustly and reliably, even when analyzing independently collected data. Thus, the
strategy should be of great value to the growing community of cerebellar researchers using high-
density silicon probes. Furthermore, our strategy provides a template for principled semi-
automated detection of cell type, based on assembly of a ground-truth library, that can be applied
across other neural circuits in the brain.

Methods

We conducted experiments in four laboratories and on two species, mice and macaque monkeys.
All mouse procedures in the Häusser lab were approved by the local Animal Welfare and Ethical
Review Board at University College London and performed under license from the UK Home
Office in accordance with the Animals (Scientific Procedures) Act 1986 and in line with the
European Directive 2010/63/EU on the protection of animals used for experimental purposes.
Mouse procedures in the Hull and Medina labs were approved in advance by the *Institutional*
Animal Care and Use Committees at Duke University and the Baylor College of Medicine,
respectively, based on the guidelines of the United States' *National Institutes of Health*. Monkey
procedures in the Lisberger lab were approved in advance by the *Institutional Animal Care and*
Use Committee at Duke University. Every effort was made to minimize both the number of
animals required and any possible distress they might experience.

Mouse experimental procedures
Mouse lines. All transgenic mice were maintained on the C57BL/6J background. Both male and
female mice were used and results were pooled.

● *Häusser:* Mice expressing Channelrhodopsin-2 (ChR2) in various cerebellar cell types were
generated primarily by crossing Cre lines to a Cre-dependent ChR2-eYFP reporter line[105]
(Ai32, B6.Cg-Gt(ROSA)26Sortm32(CAG-COP4*H134R/EYFP)Hze/J), or, in a subset of
experiments, by injecting Cre-dependent ChR2 virus (AAV1.CAGGS.Flex.ChR2-tdTomato
[UPenn]). Cre lines were: BAC-Pcp2-IRES-Cre (B6.Cg-Tg(Pcp2-cre)3555Jdhu/J), intended
to label Purkinje cells[36]; Nos1-Cre (B6.129-Nos1tm1(cre)Mgmj/J), intended to label
molecular layer interneurons[106]; Glyt2-Cre (Tg(Slc6a5-cre)1Uze), intended to label Golgi
cells[35]; and Math1-Cre (B6.Cg-Tg(Atoh1-cre)1Bfri/J), intended to label granule cells[95]. In
addition to the transgenic crosses and viral ChR2 expression, we used the Thy1-ChR2 line 18
(B6.Cg-Tg(Thy1-COP4/EYFP)18Gfng/J) to express ChR2 in mossy fibers[37]. Recordings
using each strategy were performed as follows: L7-Cre x Ai32 – 1 recording (1 mouse), Nos1-
Cre x Ai32 – 40 recordings (34 mice), Nos1-Cre + AAV1.CAGGS.Flex.ChR2-tdTomato – 3
recordings (3 mice), GlyT2-Cre x Ai32 – 32 recording (31 mice), GlyT2-Cre +
AAV1.CAGGS.Flex.ChR2-tdTomato – 3 recordings (3 mice), Math1-Cre x Ai32 – 47
recording (38 mice), Math1-Cre + AAV1.CAGGS.Flex.ChR2-tdTomato – 3 recordings (3
mice), and Thy1-ChR2 line 18 – 26 recordings (22 mice). The specificity of opsin expression
in the cerebellum of our Cre transgenic crosses was further investigated by crossing the listed
Cre lines to a Cre-dependent tdTomato reporter line, Ai9 (B6.Cg-Gt(ROSA)26Sortm9(CAG-
tdTomato)Hze/J)[107], so that we could evaluate expression specificity through cytosolic, rather
than membrane-bound, fluorescence.

● *Hull:* Mice expressing ChR2 or the inhibitory opsin GtACR2 were generated by either
crossing the *c-kit*[IRES-Cre], intended to label molecular layer interneurons[34] or BACα6Cre-C,
intended to label granule cells[108], to Ai32[105] or a Cre-dependent ArchT-GFP reporter line,
Ai40 (B6.Cg-Gt(ROSA)26Sortm40.1(CAG-aop3/EGFP)Hze/J)[109]. Alternatively, we injected
the same lines with Cre-dependent viruses: (AAV1.CAGGS.Flex.ChR2-tdTomato [UPenn]
and AAV1.Ef1a.Flex.GtACR2.eYFP [Duke]). In addition, we used the Thy1-ChR2 line 18 to
express ChR2 in mossy fibers. Recordings using each strategy were performed as follows: *c-*
kit[IRES-Cre] + AAV1.CAGGS.Flex.ChR2-tdTomato – 8 recordings (2 mice), *c-kit*[IRES-Cre] x Ai40

– 3 recordings (1 mouse), *c-kit*[IRES-Cre] + AAV1.Ef1a.Flex.GtACR2.eYFP – 11 recordings (6
mice), BACα6Cre-C x Ai32 – 22 recordings (12 mice), BACα6Cre-C +
AAV1.CAGGS.Flex.ChR2-tdTomato – 10 recordings (4 mice), and Thy1-Chr2 line 18 – 13
recordings (4 mice).

●   *Medina:* All experiments were performed in C57BL/6J mice of at least 10 weeks of age,
obtained from Jackson Laboratories.

Experimental preparation. To prepare mice for awake in vivo recordings, in all labs they were
implanted with a headplate/headpost under isoflurane anesthesia in sterile conditions. Pre-
operative and post-operative analgesia were administered, and mice were allowed to recover
from surgery for at least one week before being habituated to head-fixation and prepared for
recordings. Lab-specific details are as follows:

●   *Häusser:* We installed a custom-made aluminum headplate with a 5 mm long and 9 mm wide
oval inner opening over the cerebellum. Mice received a steroid anti-inflammatory drug at
least 1 hour before surgery (Dexamethasone, 0.5 mg/kg), followed by an analgesic NSAID
(Meloxicam, 5mg/kg) immediately before surgery. Anesthesia was induced and maintained
with 5% and 1-2% isoflurane, respectively. The headplate was positioned over the lobule
simplex of the left cerebellar hemisphere, angled at approximately 26° with respect to the
transverse plane, and attached to the skull with dental cement (Super-Bond C&B, Sun-
Medical). Post-operative analgesia (Carprieve, 5 mg/kg) was given for 3 days. After several
days of habituation on the recording apparatus, a 1 mm-diameter craniotomy and durotomy
were performed to allow access for Neuropixels probes into the lobule simplex (3 mm lateral
to the midline, anterior to the interparietal-occipital fissure). Before the craniotomy, a conical
nitrile rubber seal (Stock no. 749-581, RS components) was attached to the headplate with
dental cement to serve as a bath chamber. The exposed brain was then covered with a humid
gelatinous hemostatic sponge (Surgispon) and silicone sealant (Kwik-Cast, WPI) until the
experiment was performed (1-2 h after recovery). At the beginning of the experiment, mice
were head-fixed, the silicone sealant was removed, and physiological HEPES-buffered saline
solution was immediately applied to keep the craniotomy hydrated.

●   *Hull:* We installed a titanium headpost (HE Palmer, 32.6x19.4 mm) to the skull and a
stainless-steel ground screw (F.S. Tools) over the left cerebellum, both secured with
Metabond (Parkell). Mice received dexamethasone (3 mg/kg) 4-24 hours before surgery and
an initial dose of ketamine/xylazine (50 mg/kg and 5 mg/kg, IP) and carprofen (5 mg/kg) 20
min before induction with isoflurane anesthesia. Isoflurane was administered at 1-2%
throughout surgery to maintain appropriate breathing rates and prevent toe pinch response,
which were monitored throughout the duration of the surgery. Body temperature was
maintained with a heating pad (TC-111 CWE). Mice received buprenex and cefazolin (0.05
mg/kg and 50 mg/kg respectively, subq) twice daily for 48 hours after surgery and were
monitored daily for 4 days. After 2+ weeks of recovery, mice received dexamethasone (3
mg/k) 4-24 hours before recordings. Craniotomies (approx. 0.5-1.5 mm) were opened over
vermis or lateral cerebellum (relative to bregma: between -6.0 and -7.0mm AP, and between
1.0 and 2.8mm ML) on the first day of recording, under 1-2% isoflurane anesthesia, and were
sealed between recordings using Kwik-Cast (WPI) covered by Metabond. Craniotomies

could be re-opened for subsequent recordings under brief (<30 min) 1-2% isoflurane
anesthesia.

•   *Medina:* Preoperative analgesia was provided (5g/kg meloxicam, subq, 0.02mL 0.5%
bupivacaine and 2% lidocaine, subq) and surgery was carried out under sterile conditions.
Mice were anesthetized with isoflurane (5% by volume in O_2 for induction and 1-2% by
volume for maintenance; SurgiVet) and kept on a heating pad to maintain body temperature.
The skull was exposed and leveled to match the stereotaxic plane before two stainless steel
screws were implanted (relative to bregma: AP -0.3mm, ML ±1.4mm) to anchor the whole
preparation. A custom-made stainless steel headplate was placed over the screws and the
whole preparation was secured to the skull with Metabond cement (Parkell). Additionally, a
craniotomy was performed (relative to bregma: AP -5.5mm) consisting of a 5x2 mm section
of bone removed to expose the cerebellar vermis and the right anterior and posterior lobes. A
chamber was then built with Metabond to cover the exposed bone around the craniotomy, the
dura was protected with a thin layer of biocompatible silicone (Kwik-Cast, WPI) and the
whole chamber sealed with silicone adhesive (Kwik-Sil, WPI). Mice were monitored until
fully recovered from anesthesia and analgesia was provided during the three days following
the surgical procedure.

Recording procedures. All labs followed the same general procedures for mouse cerebellar
recordings. Mice were progressively habituated to head fixation prior to Neuropixels recordings.
Recordings from the cerebellar cortex were made using Neuropixels 1.0 probes. Probes were
coated with DiI, DiO, or DiD (Cat.Nos.V22885, V22886, and V22887; Thermo Fisher
Scientific) by repeatedly moving a drop of dye along the probe shank using a pipette until a dye
residue was visible along its entire length (~20 passes). Probes were inserted into the brain at a
speed of 1-4 µm/s while monitoring electrophysiological signals. The recording chamber
surrounding the craniotomy was bathed in ACSF, with or without blockers. After each recording,
the probe was removed and soaked in Tergazyme, then soaked in distilled water, and finally
washed with isopropyl alcohol. After the last recording session, the brains of most mice were
fixed and processed for histology to verify recording locations.

In all three laboratories, Neuropixels data were acquired using SpikeGLX
(https://github.com/billkarsh/SpikeGLX). Following data acquisition, automated spike sorting
was performed using Kilosort 2.0[17,110] and manual curation was performed using Phy
(https://github.com/cortex-lab/phy). Across all labs, signals were digitized at 30 kHz. Onboard
filtering was turned on in some but not all cases.

Optogenetic stimulation and pharmacology. The same general procedures were followed for
optogenetic stimulation in both the Häusser and Hull labs. This procedure consisted of four parts:
(1) a baseline recording period without stimulation or drug block, (2) a period of optogenetic
stimulation without drug block, (3) a period during which synaptic blockers were applied and
diffused into the brain, and (4) a period of optogenetic stimulation with blockers present. The
details of the procedures for ground-truth identification of cell-type varied slightly between the
two labs.

•   *Häusser:* Optogenetic stimulation was performed using 1 or 2 blue LEDs (470 nm, Thorlabs
M470F3) and in some experiments a blue laser for surface illumination (Stradus 472,

Voltran). Surface illumination was performed by coupling the laser or the LED via a patch
cable (M95L01, Thorlabs) to a cannula (CFMXB05, Thorlabs) positioned in contact with the
brain surface near the probe. In some experiments a second illumination source - a tapered
fiber (Optogenix 0.39NA/200μm) glued directly to the head of the Neuropixels probe - was
inserted into the brain. Total power at the fiber tip (surface fiber) and coupling cannula
(tapered fiber) was 1-6.9 mW. Each recording session consisted of: (1) a 20 minute baseline
period of spontaneous activity, (2) a set of 50 optogenetic stimuli (stimulation duration: every
10 seconds, 1 stimulation of 250 ms or a train of 5 stimulations of 50ms at 5 Hz, depending
on the experiment), (3) an application of a synaptic blocker cocktail (Gabazine 0.2-0.8 mM,
NBQX 0.8 mM, APV 1.6 mM, MCPG 0-1.3 mM) to the surface of the cerebellum followed
by a 20 minute incubation, and (4) a second set of 50 optogenetic stimuli in the presence of
synaptic blockers. We note that we did not record any neurons in the ground-truth library
with the blue laser as a source of photostimulation.

● *Hull:* Neurons expressing ChR2 were activated and neurons expressing GtACR2 were
inhibited using a 450 nm laser (MDL-III, OptoEngine) using a 400 micron optic patch fiber
(FT400 EMT, Thorlabs) that was positioned 4-10 mm from the brain surface. Power at the
brain surface was approximately 2-30 mW and was calibrated for each experiment to
produce neuronal responses with minimal artifact. Laser stimulations lasted 50 or 100 ms and
were delivered at 0.1 Hz throughout the recording after the 20 minute baseline period, with
brief pauses to replenish ACSF or apply blockers (Gabazine 0.2-0.8 mM, NBQX 0.6-1.2
mM, AP-5 0.15-0.6 mM, MCPG 1-2.5 mM).

Histology
● *Häusser:* Mice were deeply anesthetized with ketamine/xylazine and perfused transcardially
with PBS followed by 4% PFA in PBS. The brains were dissected and post-fixed overnight
in 4% PFA, then embedded in 5% agarose (for electrode tract reconstruction) or sectioned at
100 μm (for immunohistochemistry). To reconstruct electrode tracts, we imaged full 3D
stacks of the brains in a custom-made serial two-photon tomography microscope coupled to a
microtome[111], controlled using ScanImage (2017b, Vidrio Technologies) and BakingTray
(https://github.com/SainsburyWellcomeCentre/BakingTray, extension for serial sectioning).
The entire brain was acquired with the thickness of physical slices set at 40 μm and that of
the optical sections at 20 μm (2 optical sections/slice) using a piezo objective scanner
(PIFOC P-725, Physik Instrumente) in two channels (green channel: 500–550 nm, ET525/50;
red channel: 580–630 nm, ET605/70; Chroma). Each section was imaged in 1025 x 1025 μm
tiles at 512x512-pixel identification with 7% overlap using a Nikon 16x/0.8NA objective.

After slicing, samples for immunohistochemistry were blocked with 2.5% normal donkey
serum / 2.5% normal goat serum / 0.5% Triton X-100/PBS for 4-6 hours at room
temperature, primary antibodies for 4-6 days at 4°C, and secondary antibodies overnight at
4°C. Antibodies were diluted in blocking solution. The following antibodies were used: rat
anti-mCherry (1:250, ThermoFisher M11217), Mouse anti-Parvalbumin (1:1000, Millipore
MAB1572), Donkey anti-Rat-Alexa 594 (1:1000, Invitrogen), and Goat anti-Mouse-Alexa
633 (1:1000, Invitrogen). Neurotrace 435/455 (1:250, ThermoFisher N21479) was added to
the secondary antibody solution. Sections were mounted and imaged on a Zeiss LSM 880
using a 20x objective in 425x425 μm tiles at 1024x1024-pixel identification.

● *Hull:* After the last day of recording, mice were deeply anesthetized with ketamine/xylezine
(350 mg/kg and 35 mg/kg, IP) and perfused with PBS followed by 4% PFA in PBS. Brains
were extracted and post-fixed in 4% PFA in PBS overnight, then sectioned at 100 mm using
a vibratome (Pelco 102). Before sectioning, some brains were encased in a 2% agar block for
stability. Slices were either stained with DAPI (DAPI, Dihydrochloride, 268298, EMD
Millipore) and then mounted with mounting medium (Fluoromount-G, Southern Biotech) or
were mounted with a DAPI-containing mounting medium (DAPI Fluoromount-G, Southern
Biotech). Electrode tracts were visualized using a confocal microscope (Leica SP8).

● *Medina:* After perfusion with 4% PFA in PBS, brains were extracted, post-fixed in the same
solution for at least 12h and then cryoprotected in 30% sucrose solution in PBS for 48h. The
brains were aligned so the coronal sections would match the track angle and sectioned at 50
µm on a cryostat (Leica CM1950). Free floating sections were recovered in PBS and
incubated in Hoechst solution for 3 minutes (Hoechst 33342, 2µg/mL in PBS-TritonX
0.25%, Thermo Fisher Scientific). Sections were then washed in PBS three times using
fluorescence protectant medium (ProLong Diamond Antifade, Thermo Fisher Scientific).
Epifluorescence was acquired at 10x magnification on an Axio Imager Z1 microscope
(Zeiss), track reconstruction and measurements were made on specific microscopy analysis
software (ZEN software, Zeiss).

Validation of ChR2 specificity. To identify the classes of cerebellar neurons that expressed
optogenetic actuators, we determined the layer in which fluorescent neurons were present and
whether they expressed parvalbumin (PV), which should be present in all molecular layer
interneurons and Purkinje cells[112]. The location of cerebellar layers in each image were identified
in the Neurotrace (fluorescent Nissl) channel. The soma locations of neurons expressing
tdTomato (as a proxy for Cre expression) and PV were marked manually in grayscale images
using Fiji (NIH). Neurons were deemed to express both tdTomato and PV if their somatic
locations were less than 5 µm apart, and the layer of each neuron was determined by overlaying
the Neurotrace laminar mask to cell locations.

Macaque experimental procedures
Recordings in non-human primates were conducted in the *Lisberger* lab on three male rhesus
monkeys (*Macaca mulatta*) weighing 10-15 kg. A portion of the primate dataset reported here
have been published previously along with corresponding detailed methods[113]. Briefly, monkeys
underwent several surgical procedures under isoflurane in preparation for neurophysiological
recordings. In succession, we (i) affixed a head-holder to the calvarium, (ii) sutured a small coil
of wire to the sclera of one eye to monitor eye position and velocity using the search coil
technique[114] and (iii) implanted a recording cylinder aimed at the floccular complex. Analgesics
were provided to the monkeys after each surgery until they had recovered.

Each day, we acutely inserted either tungsten single electrodes (FHC) or, for the majority of our
data, custom-designed Plexon s-Probes into the cerebellar floccular complex. Plexon s-Probes
included 16 recording contacts (tungsten, 7.5 µm diameter) spaced in two columns separated by
50 µm. Adjacent rows of contacts were also separated by 50 µm. Once we had arrived in the
ventral paraflocculus, we allowed the electrode to settle for a minimum of 30 minutes. We
recorded continuous wideband data from all contacts at a sampling rate of 40 kHz using the
Plexon Omniplex system. We used a 4th order Butterworth low-pass hardware filter with a cutoff

frequency of 6 kHz prior to digitization to eliminate distortion of the recorded signal by the
electrical field produced by the eye coils. All recordings were performed while the monkey
tracked discrete trials of smooth motion of a single black target (0.5° diameter) on a light grey
background in exchange for liquid reward. All analyses of the primate neurons utilized the entire
recording period and were not contingent on the animal's behavior.

Data processing and analysis
Assignment of layers with Phyllum. For recordings in the mouse, we assigned each channel of
the Neuropixels probe to a layer using *Phyllum*, a custom-designed plugin for the curation
software *Phy*. The algorithm for layer identification in *Phyllum* starts by automatically setting
'anchor' channels whose recorded layer can be unambiguously identified by the presence of
Purkinje cell units with simple and complex spikes (Purkinje layer anchor), mossy fiber units
with triphasic waveforms (granule layer anchor), or low 1-2 Hz frequency units with wide
waveforms indicative of dendritic complex spikes (molecular layer anchor). Then, *Phyllum* fills
in the layer of the remaining channels via an iterative procedure based on (1) proximity to the
nearest Purkinje cell anchor and (2) allowed layer transitions. Every channel assigned to the
Purkinje cell layer must contain at least a Purkinje cell recording within 100 μm, but the channel
may also contain additional units located in the neighboring granule or molecular layers. If none
of the channels between two consecutive Purkinje cell anchors contain another anchor unit, their
layer is set to 'Unknown'. On average, *Phyllum* assigns 82% of all the channels on the
Neuropixels probe to a specific layer. Histological reconstruction of 21 recording tracks
confirmed that for channels that are assigned a specific layer, the assignment is highly accurate:
>99% for molecular layer channels, >98% for granule layer channels, and >95% for Purkinje
layer channels.

Curation procedures
● **Mouse**: After automated sorting with Kilosort and initial manual curation with Phy, we
implemented checks to ensure that the resulting clusters selected for further analysis
corresponded to single units with physiological waveforms, good isolation properties, and
few or no refractory period violations. Rigorous curation was especially important for our
long recordings, which could have periods of good isolation intermixed with periods of drift
or poor unit isolation. We divided our recordings into overlapping segments (30 seconds
segments computed every 10 seconds) and computed the 'false-positive' and 'false-negative'
rates in each segment. False positives were defined as spikes that fell within the refractory
period of a unit (± 0.8 ms from a given spike) and termed refractory period violations
(RPVs). The proportion of false-positives was estimated as the quotient between the RPV
rate and the mean firing rate[46]. False negatives were defined as spikes that were not detected
because they fell below the noise threshold of the recording. They were estimated by fitting
each unit's spike amplitude distribution with a Gaussian function[47,48] and quantifying the
fraction of area under the curve clipped at the noise threshold. A 30-second segment was
deemed acceptable if it had less than 5% of false positive rate <u>and</u> less than 5% of false
negative rate. Acceptable intervals were concatenated and used for subsequent classifier
training. A unit was required to have 3 minutes of acceptable isolation during the baseline
period to be included in the sample.

● **Monkey**: Following each recording session, individual action potentials were assigned to
putative neural units using the semi-automated "Full Binary Pursuit" sorter[115], designed to

distinguish temporally and spatially overlapping spikes from different neurons. Following
automated sorting, we manually curated our dataset, removing neurons with significant
interspike interval violations or low signal-to-noise ratios. The majority of units in our
primate dataset significantly exceeded the metrics used for automated curation of the mouse
data, which potentially biases our sample of primate units towards those that are easier to
record.

Data harmonization. To achieve consistency of data acquired across labs and setups, we
implemented several procedures (Supplementary Figure 1):

1. We reprocessed the wideband voltage recordings from monkeys and mice where the
SpikeGLX acquisition filter was off with a causal first-order Butterworth high-pass filter
(300 Hz cutoff) to agree with the hardware filter used by Neuropixels probes. Following
filtering, we used a drift- and shift-matching algorithm to generate mean waveforms for each
recorded unit.

2. We sought to remove one source of waveform variability by flipping the waveform, if
necessary, so that the largest peak was always negative. We did so with the knowledge that
the polarity of the action potential waveform depends on a number of factors including the
proximity of the recording electrode to the dendrites, soma, and axon[88,116] and relative
orientation of the recording contact and the reference.

3. We preprocessed all waveform templates by selecting the mean waveform from the highest
amplitude channel, resampling it to 30 kHz (if necessary), aligning it to the peak, and
inverting it if necessary (see #2 above) to ensure the most prominent peak in the waveform
was always negative. We used the harmonized waveforms to compute summary statistics
(Figure 5, Supp Figure 4), which have been previously used to classify cerebellar
neurons[19,20,62,63].

4. We sub-sampled the spikes of each neuron by grouping waveforms with a similar amplitude
on the principal channel, and therefore the same drift-state (i.e. position of probe relative to
the recorded neuron): "drift-matching".

5. We re-aligned the spikes in time by maximizing the cross-correlation of each spike to a high
amplitude template: "shift-matching". After alignment, the individual spikes were averaged,
resulting in the final mean waveform for the neuron under study. Neuropixels data processing
(non-manual curation, filtering, drift-shift-matching) was performed using the NeuroPyxels
library[117].

Identification of units directly responsive to optogenetic stimulation
Units recorded during optogenetic activation experiments were deemed to be directly
responsive to photostimulation if they met the following conditions: (1) their firing rate increased
(ChR2) or decreased (GtACR2) more than 3.3 standard deviations from the pre-stimulus baseline
within 10 ms of stimulation onset in the 'post-blocker' trials (computed using 0.1 ms bins
smoothed with a causal Gaussian filter with a standard deviation of 0.5 ms), (2) they were
recorded at a depth at which pharmacological blockade was confirmed, and (3) the spike

waveforms evoked in the 'post-blocker' optogenetic stimulation trials matched those recorded
during the pre-stimulation 'baseline' period.

Construction of 3D autocorrelograms
All recordings were performed in awake animals that were either head-fixed but otherwise free
to move on a wheel (mice) or performing discrete trials of smooth pursuit (primates), so that
firing rates modulated across the experimental session. To account for the impact of changes in
firing rate on measures of firing statistics, we constructed "three-dimensional autocorrelograms"
(3D-ACGs). At each point in time, we estimated the instantaneous firing rate of the neuron as the
inverse interspike interval[118]. We smoothed firing rates using a boxcar filter (250 ms width) and
evaluated the smoothed firing rate at each spike. Finally, we determined the distribution of firing
rates from all interspike intervals in a recording, stratified firing rate into 10 deciles, and
computed separate 2D-ACGs for the spikes in each decile. We visualized the resulting 3D-ACGs
as a surface where the color axis corresponds to the probability of firing, the y-axis stratifies the
firing rate deciles so that each 3D-ACG contains 10 rows, and the x-axis represents time from
the trigger spike. Note that the spike counts in the autocorrelograms have been divided by the
width of the bin so that the y-axis or color map is calibrated in spikes/s.

As input to the classifier, we used log distributed bins relative to $t=0$ in contrast to the linearly
spaced bins shown in the Figures and Supplemental material.

Human expert labeling of cerebellar units
● **Mouse**. We used Phyllum to identify the layer of each recording. Most Purkinje cells were
identified by the presence of both simple spikes and complex spikes and complex-spike-
triggered histograms that showed a characteristic pause in the simple spike firing rate
following the complex spike. We identified a number of recordings as Purkinje cells by the
presence of simple spikes without a complex spike, location in a Purkinje cell layer, and
regular firing rate resulting in characteristic "shoulders" present in the autocorrelogram.
Putative molecular layer interneurons were identified by their presence in a molecular layer
with firing rates above 5 spikes/s, incompatible with the firing properties of the dendritic
complex spikes. A subset of putative molecular layer interneurons yielded a properly timed
spike-triggered inhibition of an identified Purkinje cell simple spike. Putative mossy fibers
were in a granular cell layer and some displayed a characteristic triphasic shape due to the
negative afterwave recorded near the glomerulus[50,51]. Putative Golgi cells were in the
granular cell layer and had broad waveforms and relatively regular firing rates. In addition,
some pairs of putative Golgi cells showed a double peak in the millisecond range in their
cross-correlograms, indicative of gap-junction coupling[119].

● **Monkey.** We classified recordings as ground-truth Purkinje cells if they demonstrated the
characteristic post-complex-spike pause in simple-spike firing. Units that exhibited known
characteristics of Purkinje cell simple spikes but lacked a complex spike were treated as
"putative" Purkinje cells and used in the comparison of classifier-predicted and expert-
predicted labels. We included molecular layer interneurons only if they showed spike-
triggered inhibition of an identified Purkinje cell's simple spikes at short latency, leaving
some potential molecular layer interneurons out of our sample. We included units as putative
mossy fibers only if the waveform showed a negative after-wave, characteristic of recording
near a single glomerulus[50,51]. We note that our classification of mossy fibers is highly

conservative and likely leaves a large subset of mossy fiber recordings not near a glomerulus
as unlabeled. Putative Golgi cells were identified by their presence in the granule cell layer,
broad waveforms, and highly regular firing, consistent with previous recordings[63]. Expert
labeling of units in the monkey were performed before collection and analysis of the ground-
truth units in the mouse.

Cross-validated cell-type classification
We began the design of our cell-type classifier by selecting the feature space passed to the
model, the model class, and model characteristics such as number of units and learning rate,
collectively the features that define the model's "hyperparameters". Our decision to select
hyperparameters independently from the ground truth dataset was critical to ensure
generalizability by minimizing overfitting[120]. To construct an unbiased feature space to train the
model, we decided *a priori* that the model's inputs would be anatomical location, extracellular
waveform, and firing statistics. We elected not to use summary statistics because they provide an
impoverished set of information compared to the inputs we selected. We optimized the model's
architecture fully independently from our ground truth dataset by leveraging n=3,090 curated but
unlabeled units that were recorded in the experiments used to create the ground-truth library but
were not activated optogenetically. We trained variational autoencoders to reconstruct the
waveforms and 3D-ACGs of the unlabeled units, and optimized the architecture of the
autoencoders based on the quality of the reconstruction, independently from the ground truth
dataset. In the final classifier, we used encoder networks of the two autoencoders to reduce the
dimensionality of the waveforms and 3D-ACGs of the ground-truth library. The output of the
autoencoders, along with the layer of each neuron, served as inputs used to train the final
classifier on the ground truth dataset. Thus, no aspect of the model's feature space or architecture
was chosen based on the model's performance on the ground truth dataset.

Our classifier is a "semi-supervised" model because the variational autoencoders were tuned and
trained with unsupervised learning on a set of unlabeled neurons while the complete classifier
was trained with supervised learning on a separate set of ground-truth identified neurons. We
derived our strategy from the "M1" model[121].

Variational autoencoder pre-training on unlabeled data
We trained two separate autoencoders to reconstruct the waveforms and log-scaled 3D-ACGs of
our unlabeled units. Ultimately, the encoder networks of the autoencoders, trained on our
unlabeled data, compressed the input data into two 10-dimensional 'latent spaces' for the 2 input
features: 3D-ACG and waveform. The training objective of the variational autoencoders was the
'Evidence Lower Bound' loss[70] modified to include a β term to encourage disentanglement of the
latent space (Higgins et al. 2016). During training, we employed a Kullback–Leibler divergence
annealing procedure to enhance model stability and convergence[122]. Both variational
autoencoders were trained through gradient descent with the Adam optimizer, complemented by
a cosine-annealing learning rate strategy with periodic warm restarts[123].

To both facilitate model convergence and yield high-quality reconstructions, we manually
adjusted variational autoencoder parameters to adapt the model to our specific data
characteristics and improve its performance in subsequent tasks. This procedure did not rely on
the ground truth dataset, so we could adjust hyperparameters freely without overfitting to the
classification task.

● The final *waveform variational autoencoder* consisted of a 2-layer multilayer perceptron
(MLP) encoder with Gaussian Error Linear Units (GeLU) non-linearities[124] and a 2-layer
MLP decoder also with GeLU non-linearities. It was trained for 60 epochs with η=1e-4, β=5
and a mini-batch size of 128.

● The final *3D-ACG variational autoencoder* consisted of a 2-layer convolutional neural
network (CNN) encoder with average pooling after convolutions, batch normalization, and
rectified linear unit (ReLU) non-linearities, and a 2-layer MLP decoder with ReLU non-
linearities. It was trained for 60 epochs with η=5e-4, β=5 and a mini-batch size of 32.

The analysis described in Supplementary Figure 7 ensures that our trained variational
autoencoders accurately captured the variance in our data.

<u>Semi-supervised classifier</u>
The final classifier model consisted serially of: (1) the waveform and temporal feature
"variational autoencoders" pretrained on unlabeled data to reduce the dimensionality of the input
features; (2) a multi-headed input layer that accepted the latent spaces of the waveform and 3D-
ACG variational autoencoders, along with a "one-hot-encoded" 3-bit binary code of the unit's
cerebellar layer; (3) a single fully-connected hidden layer with 100 units that processed the 3
normalized inputs; (4) an output layer with one output unit for each of the 5 cell types (Figure
6C). The value of the output units sums to 1 via a softmax function so that the output of the
classifier is the probability that a given set of inputs is from each of the 5 cell types. Between the
input (2) and fully-connected (3) steps, we applied batch normalization[87] to equate the
contributions of waveform, discharge statistics, and layer. The fully-connected hidden layer had
a Gaussian prior that encouraged each network unit to have activation values across the training
set with zero mean and unit variance. We trained the weights of the complete classifier on the
data in the ground-truth library using gradient descent with a leave-one-out cross-validation
strategy. We trained the models until convergence or for 20 epochs, whichever came first, with
η=1e-3, a mini-batch size of 128 and the AdamW optimizer[123]. We allowed the weights in the
pre-trained variational autoencoders to change during optimization to allow fine-tuning that
caused a small improvement in performance on the downstream classification task.

Finally, we took several steps to ensure that our models are robustly trained and capable of
generalizing well across datasets:

1. To account for "class imbalance" created by the different number of neurons in each cell
type, we performed random oversampling of the under-represented cell types for every
model after splitting into testing and validation data[125].

2. We assessed the performance of all models through leave-one-out cross-validation, which
has a lower bias and comparable variance to other cross-validation methods[126,127] and has
been used in the past to model small datasets such as ours[20]. Thus, leave-one-out cross-
validation is better than other cross-validation methods but worse at estimating the
generalization error than having an independent test set. We note that our models seemed to
perform well on independent "expert-classified" test sets.

3. We did not tune the hyperparameters of the final semi-supervised classifier or of its training
procedure. We used only predefined heuristic values and trained until convergence.

4. We adopted a strategy to prevent *confidence miscalibration*, the tendency of deep neural
networks to exhibit over-confidence in their predictions[128]. We corrected the overconfidence
of each model instance by applying a last-layer Laplace approximation to the output
layer[83,84]. Further, for each leave-one-out sample, we created a "deep ensemble"[129] by
training an ensemble of 10 models with random initial conditions. We then averaged the
probability for each cell type across model instantiations. Each model generated an average
prediction probability for each cell type, yielding a set of 5 values (for the 5 classes) that
summed to 1. To quantify classifier confidence, we averaged the predicted probability for
each cell type across the 10 instantiations of the model and computed the *confidence ratio* as
the ratio of the highest- to second-highest predicted cell-type for the input features from each
cell in our samples. We chose a confidence ratio of 2 as the *confidence threshold* here, but
higher thresholds could be applied to increase confidence in each prediction of cell type.

Generalization of prediction to unlabeled mouse and macaque cerebellar neuron cell type
We predicted the cell type of mouse (*Medina*) and macaque (*Lisberger*) cerebellar neurons that
were not involved in the classifier training procedures using an ensemble classifier that utilized
all ground truth neurons and initial conditions (202 x 10 = 2020 models in total). Each of the
2020 models was slightly different from the others due to the combination of the 10 initial
conditions and the 'leave-one-out' procedure used to train them and produced slightly different
results. The predicted cell-type of each neuron in the unlabeled sample was chosen as that with
the maximum average prediction across the 2020 models. We applied the *confidence ratio* and
confidence threshold as we had for the ground-truth library.

1047

Author contributions

Contribution	Maxime Beau*	David J. Herzfeld*	Francisco Naveros Arrabal*	Marie E. Hemelt*	Federico D'Agostino*	Marlies Oostland*	Alvaro Sánchez-López*	Young Yoon Chung	Michael Maibach	Hannah N. Stabb	M. Gabriela Martínez-Lopera	Agoston Lajko	Marie Zedler	Shogo Ohmae	Nathan J. Hall	Beverley A. Clark†	Dana Cohen†	Stephen G. Lisberger†	Dimitar Kostadinov†	Court Hull*	Michael Häusser†	Javier F. Medina†
Conceptualization	•	•														•	•	•	•	•	•	•
Mouse data collection	•	•		•		•	•	•	•	•		•	•							•		
Mouse data analysis	•		•	•	•	•	•				•											
Monkey data collection		•																				
Monkey data analysis		•												•								
Classifier design	•	•		•																		
Software development	•	•	•		•							•	•									
Manuscript, led writing																•						
Manuscript, major contribution to figures	•	•		•																•		•
Manuscript, major contribution to writing	•	•																•	•	•	•	•
Manuscript, contribution to figures and/or writing	•	•	•	•	•	•	•									•	•	•	•	•	•	•
Supervision	•												•		•	•	•	•	•	•	•	•
Funding															•		•		•		•	•

* co-first author
† co-last author

Data availability
The data that support the findings of our study will be made publicly available at the time of publication.

Code availability
The custom analysis code used in our study will be publicly available on Github at the time of publication for major packages. Other code will be available from the corresponding author upon request.

References

1. Ramón y Cajal, S. *Histologie Du Système Nerveux de l'homme & Des Vertébrés*. (Maloine, Paris, 1909). doi:10.5962/bhl.title.48637.
2. Fishell, G. & Heintz, N. The Neuron Identity Problem: Form Meets Function. *Neuron* **80**, 602–612 (2013).
3. Masland, R. H. Neuronal cell types. *Curr. Biol.* **14**, R497–R500 (2004).
4. Migliore, M. & Shepherd, G. M. An integrated approach to classifying neuronal phenotypes. *Nat. Rev. Neurosci.* **6**, 810–818 (2005).
5. Zeng, H. & Sanes, J. R. Neuronal cell-type classification: challenges, opportunities and the path forward. *Nat. Rev. Neurosci.* **18**, 530–546 (2017).
6. Luo, L., Callaway, E. M. & Svoboda, K. Genetic Dissection of Neural Circuits. *Neuron* **57**, 634–660 (2008).
7. Siletti, K. *et al.* Transcriptomic diversity of cell types across the adult human brain. *Science* **382**, eadd7046 (2023).
8. Hodge, R. D. *et al.* Conserved cell types with divergent features in human versus mouse cortex. *Nature* **573**, 61–68 (2019).
9. Poulin, J.-F., Tasic, B., Hjerling-Leffler, J., Trimarchi, J. M. & Awatramani, R. Disentangling neural cell diversity using single-cell transcriptomics. *Nat. Neurosci.* **19**, 1131–1141 (2016).
10. Sugino, K. *et al.* Molecular taxonomy of major neuronal classes in the adult mouse forebrain. *Nat. Neurosci.* **9**, 99–107 (2006).
11. Doyle, J. P. *et al.* Application of a Translational Profiling Approach for the Comparative Analysis of CNS Cell Types. *Cell* **135**, 749–762 (2008).
12. Josh Huang, Z. & Zeng, H. Genetic Approaches to Neural Circuits in the Mouse. *Annu. Rev. Neurosci.* **36**, 183–215 (2013).
13. Fujita, H., Kodama, T. & du Lac, S. Modular output circuits of the fastigial nucleus for diverse motor and nonmotor functions of the cerebellar vermis. *eLife* **9**, e58613 (2020).
14. Ecker, J. R. *et al.* The BRAIN Initiative Cell Census Consortium: Lessons Learned toward Generating a Comprehensive Brain Cell Atlas. *Neuron* **96**, 542–557 (2017).
15. Bota, M. & Swanson, L. W. The neuron classification problem. *Brain Res. Rev.* **56**, 79–88 (2007).
16. Jun, J. J. *et al.* Fully integrated silicon probes for high-density recording of neural activity. *Nature* **551**, 232–236 (2017).
17. Steinmetz, N. A. *et al.* Neuropixels 2.0: A miniaturized high-density probe for stable, long-term brain recordings. *Science* **372**, eabf4588 (2021).
18. Gouwens, N. W. *et al.* Classification of electrophysiological and morphological neuron types in the mouse visual cortex. *Nat. Neurosci.* **22**, 1182–1195 (2019).
19. Haar, S., Givon-Mayo, R., Barmack, N. H., Yakhnitsa, V. & Donchin, O. Spontaneous Activity Does Not Predict Morphological Type in Cerebellar Interneurons. *J. Neurosci.* **35**, 1432–1442 (2015).
20. Van Dijck, G. *et al.* Probabilistic Identification of Cerebellar Cortical Neurones across Species. *PLOS ONE* **8**, e57669 (2013).
21. Lima, S., Hromádka, T., Znamenskiy, P. & Zador, A. PINP: a new method of tagging neuronal populations for identification during in vivo electrophysiological recording. *PloS One* **4**, (2009).

22. Tye, K. M. & Deisseroth, K. Optogenetic investigation of neural circuits underlying brain disease in animal models. *Nat. Rev. Neurosci.* **13**, 251–266 (2012).

23. Petreanu, L., Huber, D., Sobczyk, A. & Svoboda, K. Channelrhodopsin-2-assisted circuit mapping of long-range callosal projections. *Nat. Neurosci.* **10**, 663–668 (2007).

24. Petreanu, L., Mao, T., Sternson, S. M. & Svoboda, K. The subcellular organization of neocortical excitatory connections. *Nature* **457**, 1142–1145 (2009).

25. Sjulson, L., Cassataro, D., DasGupta, S. & Miesenböck, G. Cell-Specific Targeting of Genetically Encoded Tools for Neuroscience. *Annu. Rev. Genet.* **50**, 571–594 (2016).

26. Lee, C., Lavoie, A., Liu, J., Chen, S. X. & Liu, B. Light Up the Brain: The Application of Optogenetics in Cell-Type Specific Dissection of Mouse Brain Circuits. *Front. Neural Circuits* **14**, 18 (2020).

27. Liu, X. *et al.* Near-infrared manipulation of multiple neuronal populations via trichromatic upconversion. *Nat. Commun.* **12**, 5662 (2021).

28. Hull, C. & Regehr, W. G. The Cerebellar Cortex. *Annu. Rev. Neurosci.* **45**, 151–175 (2022).

29. Striedter, G. F., Bullock, T. H., Preuss, T. M., Rubenstein, J. & Krubitzer, L. A. *Evolution of Nervous Systems*. (Academic Press, 2016).

30. Jacobs, B. *et al.* Comparative neuronal morphology of the cerebellar cortex in afrotherians, carnivores, cetartiodactyls, and primates. *Front. Neuroanat.* **8**, (2014).

31. Häusser, M. & Clark, B. A. Tonic Synaptic Inhibition Modulates Neuronal Output Pattern and Spatiotemporal Synaptic Integration. *Neuron* **19**, 665–678 (1997).

32. Khaliq, Z. M., Gouwens, N. W. & Raman, I. M. The contribution of resurgent sodium current to high-frequency firing in Purkinje neurons: an experimental and modeling study. *J. Neurosci. Off. J. Soc. Neurosci.* **23**, 4899–4912 (2003).

33. Raman, I. M. & Bean, B. P. Ionic currents underlying spontaneous action potentials in isolated cerebellar Purkinje neurons. *J. Neurosci. Off. J. Soc. Neurosci.* **19**, 1663–1674 (1999).

34. Amat, S. B. *et al.* Using c-kit to genetically target cerebellar molecular layer interneurons in adult mice. *PLOS ONE* **12**, e0179347 (2017).

35. Gurnani, H. & Silver, R. A. Multidimensional population activity in an electrically coupled inhibitory circuit in the cerebellar cortex. *Neuron* **109**, 1739-1753.e8 (2021).

36. Zhang, X.-M. *et al.* Highly restricted expression of Cre recombinase in cerebellar Purkinje cells. *genesis* **40**, 45–51 (2004).

37. Hull, C. & Regehr, W. G. Identification of an Inhibitory Circuit that Regulates Cerebellar Golgi Cell Activity. *Neuron* **73**, 149–158 (2012).

38. Chow, L. M. L. *et al.* Inducible Cre recombinase activity in mouse cerebellar granule cell precursors and inner ear hair cells. *Dev. Dyn. Off. Publ. Am. Assoc. Anat.* **235**, 2991–2998 (2006).

39. Eccles, J. C., Ito, M. & Szentágothai, J. *The Cerebellum as a Neuronal Machine*. (Springer, Berlin, Heidelberg, 1967). doi:10.1007/978-3-662-13147-3.

40. Eccles, J. C., Llinás, R. & Sasaki, K. The excitatory synaptic action of climbing fibres on the Purkinje cells of the cerebellum. *J. Physiol.* **182**, 268–296 (1966).

41. Thach, W. T. Discharge of cerebellar neurons related to two maintained postures and two prompt movements. II. Purkinje cell output and input. *J. Neurophysiol.* **33**, 537–547 (1970).

42. Kostadinov, D., Beau, M., Blanco-Pozo, M. & Häusser, M. Predictive and reactive reward signals conveyed by climbing fiber inputs to cerebellar Purkinje cells. *Nat. Neurosci.* **22**, 950–962 (2019).

43.   Mathy, A. *et al.* Encoding of Oscillations by Axonal Bursts in Inferior Olive Neurons.
Neuron **62**, 388–399 (2009).

44.   Davie, J. T., Clark, B. A. & Häusser, M. The origin of the complex spike in cerebellar
Purkinje cells. *J. Neurosci. Off. J. Soc. Neurosci.* **28**, 7599–7609 (2008).

45.   Llinás, R. & Sugimori, M. Electrophysiological properties of in vitro Purkinje cell dendrites
in mammalian cerebellar slices. *J. Physiol.* **305**, 197–213 (1980).

46.   Hill, D. N., Mehta, S. B. & Kleinfeld, D. Quality Metrics to Accompany Spike Sorting of
Extracellular Signals. *J. Neurosci.* **31**, 8699–8705 (2011).

47.   Laboy-Juárez, K. J., Ahn, S. & Feldman, D. E. A normalized template matching method for
improving spike detection in extracellular voltage recordings. *Sci. Rep.* **9**, 12087 (2019).

48.   Fabre, J. M. J., Beest, E. H. van, Peters, A. J., Carandini, M. & Harris, K. D. Bombcell:
automated curation and cell classification of spike-sorted electrophysiology data. Zenodo
https://doi.org/10.5281/zenodo.8172822 (2023).

49.   Cohen, J. Y., Haesler, S., Vong, L., Lowell, B. B. & Uchida, N. Neuron-type-specific
signals for reward and punishment in the ventral tegmental area. *Nature* **482**, 85–88 (2012).

50.   Walsh, J. V., Houk, J. C., Atluri, R. L. & Mugnaini, E. Synaptic Transmission at Single
Glomeruli in the Turtle Cerebellum. *Science* **178**, 881–883 (1972).

51.   Taylor, A., Elias, S. A. & Somjen, G. Focal synaptic potentials due to discrete mossy-fibre
arrival volleys in the cerebellar cortex. *Proc. R. Soc. Lond. B Biol. Sci.* **231**, 217–230
(1987).

52.   Sibille, J. *et al.* High-density electrode recordings reveal strong and specific connections
between retinal ganglion cells and midbrain neurons. *Nat. Commun.* **13**, 5218 (2022).

53.   Arenkiel, B. R. *et al.* In Vivo Light-Induced Activation of Neural Circuitry in Transgenic
Mice Expressing Channelrhodopsin-2. *Neuron* **54**, 205–218 (2007).

54.   Niell, C. M. & Stryker, M. P. Highly selective receptive fields in mouse visual cortex. *J.*
Neurosci. Off. J. Soc. Neurosci. **28**, 7520–7536 (2008).

55.   Petersen, P. C., Siegle, J. H., Steinmetz, N. A., Mahallati, S. & Buzsáki, G. CellExplorer: A
framework for visualizing and characterizing single neurons. *Neuron* **109**, 3594-3608.e2
(2021).

56.   Badura, A. & De Zeeuw, C. I. Cerebellar Granule Cells: Dense, Rich and Evolving
Representations. *Curr. Biol.* **27**, R415–R418 (2017).

57.   D'Angelo, E. *et al.* The cerebellar Golgi cell and spatiotemporal organization of granular
layer activity. *Front. Neural Circuits* **7**, 93 (2013).

58.   Barmack, N. H. & Yakhnitsa, V. Functions of Interneurons in Mouse Cerebellum. *J.*
Neurosci. **28**, 1140–1152 (2008).

59.   Ophir, O., Shefi, O. & Lindenbaum, O. Neuronal Cell Type Classification using Deep
Learning. Preprint at https://doi.org/10.48550/arXiv.2306.00528 (2023).

60.   Lee, E. K. *et al.* Non-linear dimensionality reduction on extracellular waveforms reveals
cell type diversity in premotor cortex. *eLife* **10**, e67490 (2021).

61.   Lee, K., Carr, N., Perliss, A. & Chandrasekaran, C. WaveMAP for identifying putative cell
types from in vivo electrophysiology. *STAR Protoc.* **4**, 102320 (2023).

62.   Ruigrok, T. J. H., Hensbroek, R. A. & Simpson, J. I. Spontaneous Activity Signatures of
Morphologically Identified Interneurons in the Vestibulocerebellum. *J. Neurosci.* **31**, 712–
724 (2011).

63.   Hensbroek, R. A. *et al.* Identifying Purkinje cells using only their spontaneous simple spike
activity. *J. Neurosci. Methods* **232**, 173–180 (2014).

64. Prsa, M., Dash, S., Catz, N., Dicke, P. W. & Thier, P. Characteristics of Responses of Golgi
Cells and Mossy Fibers to Eye Saccades and Saccadic Adaptation Recorded from the
Posterior Vermis of the Cerebellum. *J. Neurosci.* **29**, 250–262 (2009).

65. Mountcastle, V. B. Modality and topographic properties of single neurons of cat's somatic
sensory cortex. *J. Neurophysiol.* **20**, 408–434 (1957).

66. Gold, C., Henze, D. A., Koch, C. & Buzsáki, G. On the Origin of the Extracellular Action
Potential Waveform: A Modeling Study. *J. Neurophysiol.* **95**, 3113–3128 (2006).

67. Katai, S. *et al.* Classification of extracellularly recorded neurons by their discharge patterns
and their correlates with intracellularly identified neuronal types in the frontal cortex of
behaving monkeys. *Eur. J. Neurosci.* **31**, 1322–1338 (2010).

68. Trainito, C., von Nicolai, C., Miller, E. K. & Siegel, M. Extracellular Spike Waveform
Dissociates Four Functionally Distinct Cell Classes in Primate Cortex. *Curr. Biol.* **29**, 2973-
2982.e5 (2019).

69. Shin, S.-L. *et al.* Regular Patterns in Cerebellar Purkinje Cell Simple Spike Trains. *PLOS*
ONE **2**, e485 (2007).

70. Kingma, D. P. & Welling, M. Auto-Encoding Variational Bayes. Preprint at
https://doi.org/10.48550/arXiv.1312.6114 (2022).

71. Lackey, E. P. *et al.* Cerebellar circuits for disinhibition and synchronous inhibition. *BioRxiv*
Prepr. Serv. Biol. 2023.09.15.557934 (2023) doi:10.1101/2023.09.15.557934.

72. Senzai, Y. & Buzsáki, G. Physiological Properties and Behavioral Correlates of
Hippocampal Granule Cells and Mossy Cells. *Neuron* **93**, 691-704.e5 (2017).

73. Lemon, R. N., Baker, S. N. & Kraskov, A. Classification of Cortical Neurons by Spike
Shape and the Identification of Pyramidal Neurons. *Cereb. Cortex* **31**, 5131–5138 (2021).

74. Ardid, S. *et al.* Mapping of Functionally Characterized Cell Classes onto Canonical Circuit
Operations in Primate Prefrontal Cortex. *J. Neurosci.* **35**, 2975–2991 (2015).

75. Mitchell, J. F., Sundberg, K. A. & Reynolds, J. H. Differential attention-dependent response
modulation across cell classes in macaque visual area V4. *Neuron* **55**, 131–141 (2007).

76. Simpson, J. I., Hulscher, H. C., Sabel-Goedknegt, E. & Ruigrok, T. J. H. Between in and
out: linking morphology and physiology of cerebellar cortical interneurons. in *Progress in*
Brain Research vol. 148 329–340 (Elsevier, 2005).

77. Holtzman, T., Rajapaksa, T., Mostofi, A. & Edgley, S. A. Different responses of rat
cerebellar Purkinje cells and Golgi cells evoked by widespread convergent sensory inputs.
J. Physiol. **574**, 491–507 (2006).

78. Hady, M. F. A. & Schwenker, F. Semi-supervised Learning. in *Handbook on Neural*
Information Processing (eds. Bianchini, M., Maggini, M. & Jain, L. C.) 215–239 (Springer,
Berlin, Heidelberg, 2013). doi:10.1007/978-3-642-36657-4_7.

79. Peikari, M., Salama, S., Nofech-Mozes, S. & Martel, A. L. A Cluster-then-label Semi-
supervised Learning Approach for Pathology Image Classification. *Sci. Rep.* **8**, 7193
(2018).

80. van Engelen, J. E. & Hoos, H. H. A survey on semi-supervised learning. *Mach. Learn.* **109**,
373–440 (2020).

81. Mahmud, M. S. & Fu, X. Unsupervised classification of high-dimension and low-sample
data with variational autoencoder based dimensionality reduction. in *2019 IEEE 4th*
International Conference on Advanced Robotics and Mechatronics (ICARM) 498–503
(2019). doi:10.1109/ICARM.2019.8834333.

82. Ganaie, M. A., Hu, M., Malik, A. K., Tanveer, M. & Suganthan, P. N. Ensemble deep learning: A review. *Eng. Appl. Artif. Intell.* **115**, 105151 (2022).

83. Kristiadi, A., Hein, M. & Hennig, P. Being Bayesian, even just a bit, fixes overconfidence in ReLU networks. in *Proceedings of the 37th International Conference on Machine Learning* vol. 119 5436–5446 (JMLR.org, 2020).

84. Daxberger, E. *et al.* Laplace Redux - Effortless Bayesian Deep Learning. in *Advances in Neural Information Processing Systems* vol. 34 20089–20103 (Curran Associates, Inc., 2021).

85. Hendrycks, D. & Gimpel, K. Bridging Nonlinearities and Stochastic Regularizers with Gaussian Error Linear Units. (2016).

86. Taha, A. A., Hennig, L. & Knoth, P. Confidence estimation of classification based on the distribution of the neural network output layer. *arXiv.org* https://arxiv.org/abs/2210.07745v2 (2022).

87. Ioffe, S. & Szegedy, C. Batch normalization: accelerating deep network training by reducing internal covariate shift. in *Proceedings of the 32nd International Conference on International Conference on Machine Learning - Volume 37* 448–456 (JMLR.org, Lille, France, 2015).

88. Buzsáki, G., Anastassiou, C. A. & Koch, C. The origin of extracellular fields and currents — EEG, ECoG, LFP and spikes. *Nat. Rev. Neurosci.* **13**, 407–420 (2012).

89. Senzai, Y., Fernandez-Ruiz, A. & Buzsáki, G. Layer-Specific Physiological Features and Interlaminar Interactions in the Primary Visual Cortex of the Mouse. *Neuron* **101**, 500-513.e5 (2019).

90. Strominger, N. L., Demarest, R. J. & Laemle, L. B. Cerebral Cortex. in *Noback's Human Nervous System, Seventh Edition: Structure and Function* (eds. Strominger, N. L., Demarest, R. J. & Laemle, L. B.) 429–451 (Humana Press, Totowa, NJ, 2012). doi:10.1007/978-1-61779-779-8_25.

91. Andersen, P., Morris, R., Amaral, D., Bliss, T. & O'Keefe, J. *The Hippocampus Book.* xx, 832 (Oxford University Press, New York, NY, US, 2007).

92. Basso, M. A. & May, P. J. Circuits for Action and Cognition: A View from the Superior Colliculus. *Annu. Rev. Vis. Sci.* **3**, 197–226 (2017).

93. Tanaka, I. & Ezure, K. Overall distribution of GLYT2 mRNA-containing versus GAD67 mRNA-containing neurons and colocalization of both mRNAs in midbrain, pons, and cerebellum in rats. *Neurosci. Res.* **49**, 165–178 (2004).

94. Ishihara, N., Armsen, W., Papadopoulos, T., Betz, H. & Eulenburg, V. Generation of a mouse line expressing Cre recombinase in glycinergic interneurons. *genesis* **48**, 437–445 (2010).

95. Wagner, M. J., Kim, T. H., Savall, J., Schnitzer, M. J. & Luo, L. Cerebellar granule cells encode the expectation of reward. *Nature* **544**, 96–100 (2017).

96. Altman, J. & Bayer, S. A. Time of origin and distribution of a new cell type in the rat cerebellar cortex. *Exp. Brain Res.* **29**, 265–274 (1977).

97. Lainé, J. & Axelrad, H. The candelabrum cell: A new interneuron in the cerebellar cortex. *J. Comp. Neurol.* **339**, 159–173 (1994).

98. Osorno, T. *et al.* Candelabrum cells are ubiquitous cerebellar cortex interneurons with specialized circuit properties. *Nat. Neurosci.* **25**, 702–713 (2022).

99. Lainé, J. & Axelrad, H. Extending the cerebellar Lugaro cell class. *Neuroscience* **115**, 363–374 (2002).

1287   100. Schilling, K., Oberdick, J., Rossi, F. & Baader, S. L. Besides Purkinje cells and granule
1288   neurons: an appraisal of the cell biology of the interneurons of the cerebellar cortex.
1289   *Histochem. Cell Biol.* **130**, 601–615 (2008).
1290   101. Lipovsek, M. *et al.* Patch-seq: Past, Present, and Future. *J. Neurosci.* **41**, 937–946 (2021).
1291   102. Cadwell, C. R. *et al.* Electrophysiological, transcriptomic and morphologic profiling of
1292   single neurons using Patch-seq. *Nat. Biotechnol.* **34**, 199–203 (2016).
1293   103. Liu, J. *et al.* Integrative analysis of in vivo recording with single-cell RNA-seq data reveals
1294   molecular properties of light-sensitive neurons in mouse V1. *Protein Cell* **11**, 417–432
1295   (2020).
1296   104. Turner, N. L. *et al.* Reconstruction of neocortex: Organelles, compartments, cells, circuits,
1297   and activity. *Cell* **185**, 1082-1100.e24 (2022).
1298   105. Madisen, L. *et al.* A toolbox of Cre-dependent optogenetic transgenic mice for light-
1299   induced activation and silencing. *Nat. Neurosci.* **15**, 793–802 (2012).
1300   106. Jelitai, M., Puggioni, P., Ishikawa, T., Rinaldi, A. & Duguid, I. Dendritic excitation–
1301   inhibition balance shapes cerebellar output during motor behaviour. *Nat. Commun.* **7**,
1302   13722 (2016).
1303   107. Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization system
1304   for the whole mouse brain. *Nat. Neurosci.* **13**, 133–140 (2010).
1305   108. Aller, M. i. *et al.* Cerebellar granule cell Cre recombinase expression. *genesis* **36**, 97–103
1306   (2003).
1307   109. Daigle, T. L. *et al.* A Suite of Transgenic Driver and Reporter Mouse Lines with Enhanced
1308   Brain-Cell-Type Targeting and Functionality. *Cell* **174**, 465-480.e22 (2018).
1309   110. Pachitariu, M. *et al.* MouseLand/Kilosort2: 2.0 final. Zenodo
1310   https://doi.org/10.5281/zenodo.4147288 (2020).
1311   111. Mayrhofer, J. M. *et al.* Distinct Contributions of Whisker Sensory Cortex and Tongue-Jaw
1312   Motor Cortex in a Goal-Directed Sensorimotor Transformation. *Neuron* **103**, 1034-1043.e5
1313   (2019).
1314   112. Bastianelli, E. Distribution of calcium-binding proteins in the cerebellum. *The Cerebellum*
1315   **2**, 242–262 (2003).
1316   113. Herzfeld, D. J., Joshua, M. & Lisberger, S. G. Rate versus synchrony codes for cerebellar
1317   control of motor behavior. *Neuron* **111**, 2448-2460.e6 (2023).
1318   114. Robinson, D. A. A Method of Measuring Eye Movement Using a Scleral Search Coil in a
1319   Magnetic Field. *IEEE Trans. Bio-Med. Electron.* **10**, 137–145 (1963).
1320   115. Hall, N. J., Herzfeld, D. J. & Lisberger, S. G. Evaluation and resolution of many challenges
1321   of neural spike sorting: a new sorter. *J. Neurophysiol.* **126**, 2065–2090 (2021).
1322   116. Gold, C., Girardin, C. C., Martin, K. A. C. & Koch, C. High-Amplitude Positive Spikes
1323   Recorded Extracellularly in Cat Visual Cortex. *J. Neurophysiol.* **102**, 3340–3351 (2009).
1324   117. Beau, M. *et al.* NeuroPyxels: loading, processing and plotting Neuropixels data in python.
1325   Zenodo https://doi.org/10.5281/zenodo.5509776 (2021).
1326   118. Lisberger, S. G. & Pavelko, T. A. Vestibular signals carried by pathways subserving
1327   plasticity of the vestibulo-ocular reflex in monkeys. *J. Neurosci.* **6**, 346–354 (1986).
1328   119. van Welie, I., Roth, A., Ho, S. S. N., Komai, S. & Häusser, M. Conditional Spike
1329   Transmission Mediated by Electrical Coupling Ensures Millisecond Precision-Correlated
1330   Activity among Interneurons In Vivo. *Neuron* **90**, 810–823 (2016).
1331   120. Verstynen, T. & Kording, K. P. Overfitting to 'predict' suicidal ideation. *Nat. Hum. Behav.*
1332   **7**, 680–681 (2023).

121. Kingma, D. P., Rezende, D. J., Mohamed, S. & Welling, M. Semi-supervised learning with
deep generative models. in *Proceedings of the 27th International Conference on Neural*
Information Processing Systems - Volume 2 3581–3589 (MIT Press, Cambridge, MA, USA,
2014). doi:https://doi.org/10.48550/arXiv.1406.5298.
122. Bowman, S. R. *et al.* Generating Sentences from a Continuous Space. in *Proceedings of the*
20th SIGNLL Conference on Computational Natural Language Learning (eds. Riezler, S. &
Goldberg, Y.) 10–21 (Association for Computational Linguistics, Berlin, Germany, 2016).
doi:10.18653/v1/K16-1002.
123. Loshchilov, I. & Hutter, F. SGDR: Stochastic Gradient Descent with Warm Restarts.
Preprint at https://doi.org/10.48550/arXiv.1608.03983 (2017).
124. Hendrycks, D. & Gimpel, K. Gaussian Error Linear Units (GELUs). Preprint at
https://doi.org/10.48550/arXiv.1606.08415 (2023).
125. Lemaître, G., Nogueira, F. & Aridas, C. K. Imbalanced-learn: A Python Toolbox to Tackle
the Curse of Imbalanced Datasets in Machine Learning. *J. Mach. Learn. Res.* **18**, 1–5
(2017).
126. Wang, B. & Zou, H. Honest leave-one-out cross-validation for estimating post-tuning
generalization error. *Stat* **10**, e413 (2021).
127. Zhang, Y. & Yang, Y. Cross-validation for selecting a model selection procedure. *J.*
Econom. **187**, 95–112 (2015).
128. Abdar, M. *et al.* A review of uncertainty quantification in deep learning: Techniques,
applications and challenges. *Inf. Fusion* **76**, 243–297 (2021).
129. Lakshminarayanan, B., Pritzel, A. & Blundell, C. Simple and Scalable Predictive
Uncertainty Estimation using Deep Ensembles. in *Advances in Neural Information*
Processing Systems vol. 30 (Curran Associates, Inc., 2017).
1357
1358