Upregulated cholesterol biosynthesis facilitates the survival of methylation-retaining AML cells following decitabine treatment.

Danielle R Bond[1,2^*], Sean M Burnard[1,2^], Kumar Uddipto[1,2], Kooper V Hunt[1,2], Brooke M Harvey[1,2], Luiza Steffens Reinhardt[1,3], Charley Lawlor-O'Neill[1], Ellise A Roper[1], Sam Humphries[1,2], Heather C. Murray[1,2], Nikola A Bowden[4,5], Anoop K Enjeti[2,4,6,7], Nicole M Verrills[1,2], Carlos Riveros[8], Kim-Anh Lê Cao[9], Heather J Lee[1,2*]

[1]*School of Biomedical Science and Pharmacy, The University of Newcastle, Australia.*

[2]*Precision Medicine Research Program, Hunter Medical Research Institute, Australia.*

[3]*Cancer Detection and Therapy Research Program, Hunter Medical Research Institute, Australia.*

[4]*School of Medicine and Public Health, The University of Newcastle, Australia.*

[5]*Drug Repurposing and Medicines Program, Hunter Medical Research Institute, Australia.*

[6]*Department of Haematology, Calvary Mater Hospital, Australia.*

[7]*New South Wales Health Pathology, John Hunter Hospital, Australia.*

[8]*Data Science, Hunter Medical Research Institute, Australia.*

[9]*Melbourne Integrative Genomics, School of Mathematics and Statistics, The University of Melbourne, Australia.*

^ Joint lead authors

* Corresponding authors

Abstract

DNA hypomethylating agents (HMAs) are used to treat acute myeloid leukaemia (AML) and myelodysplasia patients who are unsuitable for intensive chemotherapy, but low response rates and therapy-resistant relapse remain significant challenges. To optimise HMA efficacy, we must understand how resistance and relapse arise from cells that survive treatment. Here we combine single-cell multi-omic analysis with parallel colony-forming assays to link HMA-induced molecular heterogeneity with functional consequences in AML cells. HMAs, azacytidine (AZA) and decitabine (DAC), induced global epigenetic heterogeneity associated with upregulation of inflammatory responses and cell death pathways in a subset of hypomethylated cells. Some AML cells maintained high DNA methylation during treatment, and these methylation-retaining cells had increased self-renewal capacity following DAC, but not AZA. Molecular profiling of individual colonies revealed upregulated cholesterol biosynthesis as an adaptation to HMA treatment, and inhibition by rosuvastatin enhanced DAC effects *in vitro* and *in vivo*. Thus, HMA-induced heterogeneity has important implications for AML cell growth and statins are a candidate co-treatment strategy to delay or prevent HMA-resistant relapse.

Introduction

Hypomethylating agents (HMAs) are DNA methyltransferase (DNMT) inhibitors that are used to treat patients with Acute Myeloid Leukaemia (AML) and the pre-leukaemic condition Myelodysplastic Syndrome (MDS). Two commonly prescribed HMAs are the cytidine analogues, azacytidine (AZA, 5-azacytidine) and decitabine (DAC, 2'-deoxy-5-azacytidine), which are incorporated into DNA during replication[1,2]. This leads to degradation of DNMTs[3] and loss of DNA methylation in subsequent cell divisions[2,4,5].

The relatively limited side effects of these epigenetic therapies make them useful alternatives to standard intensive chemotherapies, and they are routinely administered to older, or otherwise unfit, AML patients. While single agent HMA treatment extends survival in many patients[6-12], only 20% have a complete response to therapy[13]. Responses are also short-lived (e.g., 8-15 months[14]), with acquired resistance leading to relapse in most patients. To address these limitations, many clinical trials are testing co-treatment strategies with some studies delivering improved outcomes[15]. For example, the pro-apoptotic therapy, venetoclax, has increased response rates in elderly AML patients undergoing HMA treatment[16]. However, relapse remains a significant problem with the median duration of response being 11-18 months for patients treated with both venetoclax and HMA therapy[16,17]. To improve the long-term benefits of HMA therapy, we must understand the

molecular mechanisms responsible for HMA efficacy, as well as the processes by which relapse

arises from cells that survive treatment.

While initial studies demonstrated that HMA treatment can eliminate cancer cells with self-renewal

capacity[18], some leukaemic stem cell (LSC) populations were shown to survive AZA treatment in AML

and MDS patients[19]. This may be facilitated by integrin signaling in the bone marrow

microenvironment, which induces quiescence in blasts and predicts AZA response in MDS patients[20].

Another study demonstrated that LSCs increase CD70 expression following HMA treatment, and

CD70 blockade by cusatuzumab was shown to reduce the self-renewal capacity of AML patient

blasts[21]. A phase I/II clinical trial of this combination recently reported a marginal improvement in

survival compared to historical data from a similar patient cohort[22,23].

Altered pyrimidine metabolism can also facilitate HMA resistance. Prior to DNA incorporation, both

DAC and AZA must be converted into a deoxycytidine triphosphate analogue, Aza-dCTP[24]. Unlike

DAC, AZA can also be converted to AZA-CTP and incorporated into RNA, where it influences

transcript stability and translation[25,26]. Distinct enzymes catalyse these reactions and are

dysregulated at relapse[24]. For example, Deoxycytidine Kinase (DCK) is required for the metabolism of

DAC, but not AZA. Loss of *DCK* promotes resistance to DAC[27], and its expression is downregulated in

patients who develop relapse on DAC, but not AZA[24].

In this study, we further characterise the cellular processes that facilitate AML cell survival and

proliferation following HMA treatment. Our single-cell multi-omic analysis reveals global HMA-

induced DNA methylation heterogeneity and methylation-retaining cells that appear to evade

treatment. In parallel colony-forming assays, we show that methylation-retaining cells have a growth

advantage following treatment with DAC, but not AZA, and reveal upregulation of cholesterol

biosynthesis in cells surviving treatment. Co-treatment with statins enhances the effects of DAC in

colony assays and extends the survival of leukaemia-bearing mice. Together, our work suggests that

relapse may arise from metabolic adaptation in therapy-evading cells and identifies a candidate

treatment strategy to enhance the long-term efficacy of HMA therapy.

80 Results

81 DNA methylation heterogeneity induced by HMA treatment

To characterise the responses of individual AML cells to HMA treatment, we performed single-cell

analysis of DNA methylation. AML cell lines (HL-60, MOLM-13, MV-4-11) were treated with low

doses of decitabine (DAC) or azacytidine (AZA) to induce maximal demethylation with minimal

effects on cell growth and viability (Supplementary Fig. 1). After 3 days of HMA treatment, striking

heterogeneity in DNA methylation levels was observed (Fig. 1A, Supplementary Table 1). While

untreated cells had homogeneously high levels of DNA methylation (e.g., HL-60: 65-73%), the extent

of hypomethylation varied substantially among cells treated with DAC (e.g., HL-60: 17-69%) or AZA

(e.g., HL-60: 20-69%). Interestingly, a small proportion (1-5%) of methylation-retaining cells (Fig. 1A,

dashed boxes) displayed no evidence of HMA-induced hypomethylation, with DNA methylation

levels comparable to untreated cells. The extent of HMA-induced hypomethylation was related to

cell division rate, as indicated by positive correlations between DNA methylation and CellTrace

fluorescence (Fig. 1B). This is consistent with HMA incorporation during replication[2] and confirms

that slowly dividing cells can avoid the effects of HMA treatment.

We next examined DNA methylation in different genomic contexts to test whether heterogeneity is

observed across the genome. Contexts with high levels of DNA methylation in untreated HL-60 cells

(i.e., exons, introns, intergenic regions) showed the greatest reductions upon HMA treatment (Fig.

1C), and this loss of methylation was accompanied by increased cell-to-cell heterogeneity (Fig. 1D).

Active promoters marked by H3K4me3 were the only genomic features without increased DNA

methylation heterogeneity following HMA treatment.

To explore the consequences of DNA methylation heterogeneity on other layers of genetic

regulation, we used the multi-omic data collected from HL-60 cells. The scNMT-seq method profiles

DNA methylation, DNA accessibility and gene expression in parallel[28], allowing these molecular

modalities to be correlated with each other across the genome of individual cells (Fig. 1E). Untreated

cells showed the expected trends, with DNA methylation being negatively correlated to accessibility

and gene expression, while DNA accessibility and transcription were positively correlated. In HMA-

treated cells, the relationship between DNA methylation and accessibility was weakened in all

genomic contexts. HMA treatment also weakened the correlation between DNA methylation (in CpG

islands, promoters, exons, and introns) and expression of associated genes. In contrast, HMA

treatment had minimal impact on associations between DNA accessibility and gene expression.

These observations imply that loss of DNA methylation is not always accompanied by increased

accessibility and transcription, but may introduce epigenetic noise in an otherwise well-regulated

system.

Figure 1. HMA treatment induces DNA methylation heterogeneity in AML cells. HL-60, MOLM-13
and MV-4-11 cells were labelled with CellTrace and treated with decitabine (DAC; 100nM) or
azacytidine (AZA; HL-60: 2000nM, MOLM-13 and MV-4-11: 500nM) every 24h for 72h. Single cells
collected by indexed FACS were subjected to scNMT-seq (HL-60) or scTEM-seq (MOLM-13, MV-4-
11). **A)** Violin plots of DNA methylation levels in single HL-60 (left), MOLM-13 (centre) and MV-4-11
(right) cells. Superimposed points show single-cell values from untreated (UNT, orange), DAC (cyan)
and AZA (purple) groups. Dashed boxes surround DAC and AZA cells with methylation levels within
the range of UNT samples. Data are shown for 185-222 cells collected from 2-3 replicate
experiments in each cell line (UNT, *n = 27-38*; DAC *n = 63-93*; AZA *n = 68-91*). **B)** Scatter plots
comparing CellTrace fluorescence and DNA methylation in single cells, with linear regression analysis
and F-test *p*-values. **C)** Violin plots of DNA methylation levels in different genomic contexts from HL-
60 scNMT-seq data. **D)** Violin plots of DNA methylation heterogeneity, as determined by pairwise
dissimilarity analysis, within different genomic contexts from HL-60 scNMT-seq data. **E)** Box and
whisker plots of Pearson correlation coefficients computed within single cells from HL-60 scNMT-seq
data. DNA methylation and DNA accessibility were considered in different genomic contexts, and
individual loci were matched based on genomic co-ordinates. Correlations were performed between
DNA methylation and DNA accessibility (top), DNA methylation and gene expression (middle), and
DNA accessibility and gene expression (bottom). Boxes depict the interquartile range (IQR) with
median. Whiskers extend to the highest and lowest data points within 1.5 x IQR of the first and third
quartile. Outlying data points are not shown.

Transcriptional programs linked to HMA-induced epigenetic heterogeneity

135
HMA mechanisms of action have been difficult to precisely define because genome-wide loss of DNA

methylation is associated with pleiotropic transcriptional changes[15]. HMA-induced promoter

hypomethylation is thought to allow re-expression of tumour suppressor genes[29], re-activation of

DNA repair pathways[30] and upregulation of differentiation markers[31,32]. HMA treatment can also

increase expression of cancer testis antigens[33] and enhance antigen presentation on cancer cells[34].

Furthermore, genome-wide de-repression of transposable elements (TEs) has been shown to trigger

a viral mimicry response in which dsRNA stimulate interferon signalling and apoptosis[35-37]. To clarify

how HMA-induced epigenetic heterogeneity influences transcriptional responses we performed a

multivariate analysis to integrate the three molecular layers: DNA methylation, DNA accessibility,

and transcription. We excluded untreated cells to focus on variability among HMA-treated cells and

applied an unsupervised sparse Partial Least Squares (sPLS) method[38]. This method performed

feature selection to identify variably expressed transcripts that are highly correlated to changes in

DNA methylation and accessibility in regulatory regions (CpG islands, promoters, H3K4me3 sites and

H3K27ac sites) and 3kb genomic windows.

The 200 transcript features selected by sPLS were further examined in both treated and untreated

cells, revealing 4 groups of cells with distinct transcriptional profiles (Fig. 2A, B). Consistent with the

induction of viral mimicry, a gradient of TE expression was observed across component 1. For

example, cell group 3 had high expression of LINE: L2a and low global DNA methylation levels (Fig.

2A, C and D). Gene ontology (GO) over-representation analysis (ORA) was performed for each of the

3 clusters of transcript features (Fig. 2B; Supplementary Table 2). Cell group 3 had low expression of

genes in cluster 3, which were related to translation and inhibition of cell death (Fig. 2E,

Supplementary Table 3). Simultaneously, cell group 3 had high expression of genes in cluster 1,

which were enriched in terms related to immune inflammatory response and positive regulation of

cell death (Fig. 2F, Supplementary Table 3). This transcriptional profile is consistent with the

expected effects of HMA treatment[1], and 29 of the 78 genes in expression cluster 1 were

significantly upregulated by DAC and/or AZA in matched bulk RNA sequencing (RNA-seq) data

(Supplementary Table 2). In contrast, cell group 1 displayed an inverted gene expression pattern

which was shared with many untreated cells (observed:expected ratio = 1.48). This suggests that cell

group 1 did not activate transcriptional pathways commonly associated with HMA treatment,

despite low methylation levels in most cells (Fig. 2A, D). The expression profile of cell group 2 was

intermediate between groups 1 and 3, whereas cell group 4 showed distinctive upregulation of gene

expression cluster 2. No ontology terms from gene cluster 2 retained significance after multiple

testing correction.

Figure 2. HMA-induced epigenetic heterogeneity influences transcriptional programs in AML cells. HMA treated (AZA and DAC only) HL-60 scNMT-seq data underwent multivariate feature selection by sparse least squares (sPLS) using an unsupervised model. **A)** sPLS projection of cells based on transcript features coloured by cell group 1-4 (from B). **B)** Heatmap of transcript features selected by sPLS displaying all samples (treated and untreated) as columns, are split by k-means clustering and grouped by treatment. Individual gene and TE expression levels (rows) are z-score normalised and split by k-means clustering with internal hierarchical clustering. **C-D)** sPLS projections of cells based on transcript features are coloured by **C)** LINE:L2:L2a expression, and **D)** global methylation level. **E-F)** Tree plots of the gene ontology (GO) over-representation analysis (ORA) for gene expression clusters 3 and 1. **G)** Heatmaps summarising the DNA methylation and DNA accessibility features selected by sPLS. Samples (columns) are ordered according to the heatmap in B. The average methylation or accessibility of all sPLS selected features across all samples (treated and untreated) is displayed on the left for each genomic context. The two heatmaps show z-score normalised averages of DNA methylation and accessibility for all sPLS selected features in each genomic context. **H)** Pearson correlations were computed between gene expression and DNA methylation (top) or accessibility (bottom) of associated promoters. Bar graphs show the percentage of correlations ($p <$ 0.05) with negative (left) and positive (right) coefficients for all genes and filtered by gene expression cluster (identified in D). **I)** Correlations between *RHEX* expression and DNA methylation of adjacent loci (top) are displayed together with annotated short and long interspersed nuclear elements (SINE, LINE), long terminal repeat (LTR), DNA, and Simple repeat sequences (bottom). The blue shading highlights a promoter-proximal region of intron 1 that included 6 regions with negative correlations (cor < -0.3, $p <$ 0.05) between DNA methylation and *RHEX* expression.

When considering the epigenetic features selected by sPLS (Supplementary Tables 4 and 5), we noted that cell group 3 was the only group to have relatively low DNA methylation and high chromatin accessibility across several genomic contexts (Fig. 2G). This suggests that the transcriptional response to HMA treatment depends on both reductions in DNA methylation and gains in accessibility. Consistently, transcript features from expression cluster 1 had predominantly positive correlations with accessibility features, and many negative correlations to methylation features, especially in CpG islands and 3kb genomic windows (Supplementary Fig. 2).

To test whether epigenetic alterations in cis-regulatory elements could influence transcriptional responses to HMA treatment, we next correlated gene expression to DNA methylation and accessibility in nearby genomic loci. Genes from expression cluster 1 showed a significant shift toward negative correlations with promoter methylation ($p = 2.15 \times 10^{-11}$, χ^2 test) and positive correlations with promoter accessibility ($p = 2.26 \times 10^{-8}$, χ^2 test) (Fig. 2H, Supplementary Table 6). An interesting example is *RHEX* (regulator of hemoglobinization and erythroid cell expansion), which is highly expressed in AML[39]. Negative correlations between *RHEX* expression and DNA methylation were concentrated in a promoter-proximal region of intron 1 that contained several conserved long terminal repeat (LTR) TEs (Fig. 2I). A previous study has identified an AML-specific *RHEX* transcript resulting from onco-exaptation of an upstream LTR2B element[39], so HMA-induced hypomethylation could induce other non-canonical transcripts by activating additional TEs.

In summary, our single-cell multi-omic analysis has identified transcriptional changes linked to the

patterns of epigenetic heterogeneity in HMA-treated AML cells. Importantly, activation of genes

involved in inflammatory response and cell death pathways (expression cluster 1) with simultaneous

down-regulation of inhibitory genes involved in these pathways (expression cluster 3) is observed in

only a sub-set of hypomethylated cells (cell group 3) (Fig. 2A, C, D).

Functional consequences of HMA-induced heterogeneity

To determine the functional consequences of epigenetic and transcriptional heterogeneity in HMA-

treated cells, we next performed colony-forming assays. DAC and AZA significantly decreased colony

counts (Fig. 3A), confirming that HMA treatment decreases the self-renewal capacity of AML cells[18].

DAC treatment also significantly increased the size of MOLM-13 and MV-4-11 colonies, indicative of

increased proliferation during colony formation (Supplementary Fig. 3A). To characterise the

molecular profiles of colonies formed after HMA treatment, we performed single-colony analysis of

DNA methylation (Fig. 3B, C). Following DAC treatment (Fig. 3B, solid fill), we observed a high

percentage of colonies (34-65%) with DNA methylation levels comparable to untreated colonies

(≥75%). This was in stark contrast to the low percentage of methylation-retaining cells (1-5%)

observed in single-cell data after 72h DAC treatment (Fig. 1A, dashed box; Fig. 3B, dashed line).

Surprisingly, this shift towards higher DNA methylation levels was far less pronounced in colonies

established following AZA treatment (Fig. 3C).

To assess recovery of DNA methylation following HMA treatment, a time-course analysis was

performed during the colony-forming assay (Fig. 3D). DNA methylation levels in HMA-treated HL-60

colonies were low throughout the time-course (Fig. 3D, top), mirroring an analysis performed in

suspension culture (Supplementary Fig. 1E). Similar results were obtained for MOLM-13 and MV-4-

11 colonies derived after AZA treatment (Fig. 3D, centre and bottom). In contrast, high levels of DNA

methylation were observed at early stages of colony formation (experiment day 6) following DAC

treatment of MOLM-13 and MV-4-11 cells (Fig. 3D, centre and bottom). This suggests that the shift

toward high DNA methylation observed in these colonies (Fig. 3B) is not due to a gradual recovery of

methylation. Rather, our data is consistent with the selection of highly methylated cells in colony-

forming assays performed after DAC treatment. We deduce that methylation-retaining cells have

increased self-renewal and proliferative capacity relative to hypomethylated cells, after treatment

with DAC, but not AZA.

Figure 3. Highly methylated AML cells display a growth and survival advantage following
treatment with DAC, but not AZA. A) HL-60, MOLM-13, and MV-4-11 colony counts following
treatment with DAC (cyan) or AZA (purple) vs. untreated cells (UNT, orange). **B-C)** Density plots show
the average DNA methylation levels for single cells collected on experiment day 3 (dashed line) and
individual colonies collected on experiment day 17 (solid fill) following treatment with DAC or AZA.
HL-60 scNMT-seq data were filtered for cytosines within SINE Alu sites for direct comparison to
scTEM-seq data from colonies. Data are shown for 288 colonies collected from triplicate
experiments in each cell line ($n = 96$ per treatment). **D)** Time-course experiment showing changes in
average DNA methylation of cells collected at different time points throughout the colony-forming
assay (experiment days 6, 10 and 17). Values for experiment day 3 were obtained from single-cell
data (Fig. 1A). Data in A and D are expressed as mean +/- standard error of the mean (SEM).
Statistical analysis of colony counts (A) was performed using ordinary one-way ANOVA with
Dunnett's multiple comparisons test with a $p < 0.05$ cut-off for significance ($p < 0.03$*, $p < 0.006$**, p
< 0.0002***).

Transcriptional programs associated with recovery following HMA exposure

To identify cellular processes active during recovery from HMA exposure, we next generated

matched transcriptomes from the same set of colony samples. Single-colony RNA-seq and principal

component analysis (PCA) showed that DAC and AZA samples were generally distinct from untreated

samples, regardless of their global DNA methylation levels, in all cell lines (Fig. 4A). This implies that

HMA exposure has substantial effects on the transcriptome, even in highly methylated cells.

Figure 4. HMA treatment has sustained transcriptional effects in both methylation-retaining and hypomethylated cells. A) Principal Component Analysis (PCA) plots of single-colony RNA-seq data from AML cell lines, highlighting treatment groups (UNT = orange; DAC = cyan; AZA = purple) and matched mean DNA methylation levels (circle: high > 75%; triangle: low < 75%). Data shown for 119 - 220 colonies collected from 3 replicate experiments in each cell line (UNT, *n = 14-73*; DAC, *n = 46-78*; AZA, *n = 56-73*). **B)** Heatmap of the top 2000 highly variable genes from colony RNA-seq data. Samples are ordered by decreasing global methylation levels (green gradient) within each treatment group. Rows are grouped by K-means clusters based on gene expression, with hierarchical clustering by Euclidean distance within each cluster. **C)** GO analysis of the clusters from the top 2000 highly variable genes from B. The size of the circles indicates the gene ratio (number of genes from the input list annotated to the GO term divided by the total number of genes in the input list), and the colour represents the significance of the adjusted *p*-value. Gene clusters are colour-coded on the y-axis and GO processes are shown on the x-axis.

Of the 2000 most variably expressed genes among HL-60 samples, only 215 had increased

expression specific to hypomethylated colonies (Fig. 4B, cluster 3; Supplementary Table 8). Many of

these genes (42.3%) were upregulated following 72h treatment with either DAC or AZA in bulk HL-60

RNA-seq data, and several were associated with activation of inflammatory responses within the

sPLS model (e.g., *S100A8, S100A9*; Supplementary Table 8; Fig. 2B, cluster 1). In contrast, genes in

cluster 5 were upregulated following HMA exposure in both hypomethylated and highly methylated

colonies (Fig. 4B). Only 3.5% of these genes were induced by HMA treatment in bulk RNA-seq data,

but several were included in gene expression cluster 3 from the sPLS model (e.g., *MYC, MPO*;

Supplementary Table 8; Fig. 2B). Thus, some transcriptional changes that occur immediately after

HMA treatment are maintained in hypomethylated colonies, while other genes are upregulated

during colony formation and are independent of HMA-induced global hypomethylation. GO ORA

revealed an enrichment of anti-microbial and immune-related processes among both

hypomethylation-dependent and -independent gene sets, whereas 'small molecule biosynthetic

process' was uniquely over-represented among the hypomethylation-independent cluster 5 genes

(Fig. 4C; Supplementary Table 9).

To identify cellular processes that could favour the growth of methylation-retaining cells (Fig. 3), we

next focused on the HL-60 colonies derived following DAC treatment. Among these samples, the

broad range of DNA methylation levels (Fig. 3B) allowed us to compute correlations to gene and TE

expression (Fig. 5A). We found an enrichment for negative correlations with TE expression,

consistent with their upregulation in hypomethylated cells[35-37]. We also identified transcripts with

both significant negative (cor < -0.4, adjusted *p*-value < 0.05, *n* = 722) and positive (cor > 0.4,

adjusted *p*-value < 0.05, *n* = 345) correlations to global DNA methylation levels and divided these

genes into 4 clusters based on their expression patterns across treatment groups (Fig. 5B, C;

Supplementary Table 10). Genes with increased expression in hypomethylated colonies (Fig. 5B,

cluster 2) were enriched for GO terms related to defense responses, cell motility and chemotaxis

(Supplementary Table 11). Interestingly, we also identified 233 genes with specific upregulation in

highly methylated HL-60 colonies derived following DAC treatment (Fig. 5B, cluster 1). These genes

displayed significant enrichment of cholesterol-related ontologies, which included many enzymes

required for *de novo* cholesterol biosynthesis downstream of mevalonate[40] (Figure 5C,

Supplementary Table 11; Supplementary Fig. 4). Other genes involved in the mevalonate pathway

(e.g., *MVD, MVK,* and *PMVK*) were increased by HMA treatment in most HL-60 colonies, regardless

of their global DNA methylation level. MOLM-13 and MV-4-11 colonies also displayed increased

expression of cholesterol biosynthesis genes after HMA treatment (Supplementary Fig. 5).

Interestingly, these changes were not observed immediately following HMA treatment, nor after

long-term culture in suspension (i.e., in day 3 and day 21 HL-60 bulk RNA-seq data). On the contrary,

several members of the cholesterol biosynthesis pathway were significantly decreased (FDR < 0.05,

Log2(fold change) < -1) following 72h DAC and/or AZA treatment of HL-60 cells (*LSS, DHCR7,*

Supplementary Table 8; *HMGCS1, HMGCR,* data not shown). These observations suggest that

increased cholesterol production is a delayed response to HMA treatment that occurs independently

of methylation changes and facilitates the self-renewal and proliferation of AML cells.

Inhibition of cholesterol biosynthesis following HMA exposure

To test whether inhibition of cholesterol biosynthesis could enhance HMA efficacy, we performed

co-treatments with rosuvastatin in colony-forming assays. Rosuvastatin is a potent inhibitor of the

rate limiting enzyme of the cholesterol biosynthetic pathway, HMG-CoA reductase (HMGCR), and is

frequently prescribed to reduce the risk of cardiovascular disease associated with

hypercholesterolaemia[41]. In all cell lines, rosuvastatin caused a trend towards decreased colony

numbers following DAC treatment, with a significant decrease in MOLM-13 colonies, whereas co-

treatment with AZA had varied effects (Fig. 5D). Following DAC and rosuvastatin co-treatment,

colonies also displayed significantly reduced size (Supplementary Fig. 3B) and DNA methylation

levels (Fig. 5E) in MOLM-13 and MV-4-11 cell lines, where strong selection for highly methylated

cells was previously noted (Fig. 3B). This suggests that inhibition of cholesterol biosynthesis

specifically inhibits the self-renewal capacity of methylation-retaining cells and identifies statins as a

candidate co-treatment strategy to target HMA-evading AML blasts.

The efficacy and potential clinical utility of DAC and rosuvastatin co-treatment was then tested in

immunocompromised mice engrafted with luciferase-tagged MOLM-13 cells (Fig. 5F). In a dose

optimisation experiment, combining DAC and rosuvastatin led to a significant increase in median

survival for all doses tested (Fig. 5F, left). The 1mg/kg/day dose of rosuvastatin was chosen for

further validation in this model, as it is a low dose that is commonly prescribed for preventing

cardiovascular disease (equivalent to approximately 5-10mg/day oral dose in humans). DAC and

rosuvastatin co-treatment once again led to a significant increase in median survival compared to

DAC treatment alone (24 vs. 21 days) in this aggressive *in vivo* AML model (Fig. 5F, right).

Rosuvastatin alone provided no benefit, with survival times comparable to vehicle treated mice. DAC

and rosuvastatin co-treated mice also exhibited a trend towards decreased tumour burden, as

measured by BLI, between days 17 – 24 of the experiment, compared to mice treated with DAC

alone (Supplementary Fig. 6). These *in vivo* results provide evidence that HMAs combined with

statins have the potential to decrease tumour burden and improve AML survival.

Figure 5. Upregulation of cholesterol biosynthesis in cells surviving HMA therapy can be targeted
to enhance treatment efficacy. A) Volcano plot showing the Pearson correlation coefficient and
adjusted p-value for correlations between gene or transposable element (TE) expression and global
DNA methylation levels from HL-60 colonies derived following DAC treatment. The upper density
plot shows a bias toward negative correlations, especially between global DNA methylation level and
TE expression (red). **B)** Simplified heatmap of k-means clustering for the 1,067 genes with significant
correlations to global DNA methylation level (adjusted p-value ≤ 0.05 and 0.4 ≤ correlation estimate
≤ 0.4, from A). **C)** Summarised tree plots displaying GO terms with significant (adjusted p-value <
0.05) over-representation in clusters 1 and 2 (from B). **D)** Colony counts for HL-60 (top), MOLM-13
(middle) and MV-4-11 (bottom) cell lines obtained following HMA and rosuvastatin co-treatments.

Data from No HMA (orange), DAC (cyan) and AZA (purple) groups are normalised to the
corresponding 0 µM rosuvastatin control. Means ± SEM for $n = 3$ experiments. Significance
determined by two-way ordinary ANOVA with Dunnett's multiple comparisons test, $p < 0.05$* vs.
corresponding 0 µM rosuvastatin control. **E)** DNA methylation of colonies formed following DAC and
rosuvastatin co-treatments. Means ± SEM for $n = 3$ experiments. Significance determined by one-
way ANOVA with Tukey's multiple comparisons test, $p < 0.05$* vs. corresponding 0 µM rosuvastatin
control. **F)** Left: rosuvastatin dose optimisation experiment showing median survival of mice
engrafted with MOLM-13-luc cells following treatment with DAC (0.2mg/kg/day) +/- rosuvastatin (1,
10, 40mg/kg/day) on a treatment schedule of '5 days on, 2 days off' for 3 cycles. Right: Validation of
survival benefit when DAC (0.2mg/kg/day) is combined with rosuvastatin (1mg/kg/day) in mice
engrafted with MOLM-13-luc AML cells. Survival analyses were performed using Kaplan-Meier
analysis followed by the Log-rank (Mantel-Cox) test and an adjusted p-value of < 0.05 was
considered statistically significant. Left: $p < 0.002$**, $p < 0.0001$****; Right: $p < 0.05$*, $p < 0.005$**.

Discussion

The clinical benefits of HMA therapy are limited by the rapid development of treatment-resistant

relapse. We have characterised the heterogeneous responses of AML cells to HMA treatment,

revealing new insights into how cells survive and adapt to treatment.

Our single-cell multi-omic analyses revealed global DNA methylation heterogeneity induced by

treatment, and transcriptional responses linked to epigenetic changes. We observed activation of

inflammatory response and cell death pathways in only a minor subset of hypomethylated cells (Fig.

2B), consistent with our previous report of heterogeneous TE expression following DAC treatment[42]

and scRNA-seq data from a colorectal cancer cell line[43]. These observations suggest that additional

epigenetic modifications, transcription factors, or other mechanisms, can suppress transcriptional

responses in hypomethylated cells (e.g., H3K9me3[44,45]). Alternatively, loss of methylation at specific

loci may be required for HMAs to exert their effects. Our observations are consistent with the lack of

correlation between HMA-induced hypomethylation and clinical response[46-48], and support the use

of locus-specific methylation changes to build a predictor of patient response[49].

In contrast to our single-cell analyses, Li *et al.* reported reduced epigenetic and transcriptional

variance in AML cells collected after 12 weeks of AZA treatment in a transgenic mouse model[50]. We

suggest that HMAs initially increase epigenetic and transcriptional diversity, allowing some cells to

gain a relative growth or survival advantage. Subsequent expansion of those clones would lead to

the reduced heterogeneity reported by Li *et al.*[50]. Consistently, we found that some AML cells retain

high levels of DNA methylation during HMA treatment (Fig. 1A) and have a relative growth

advantage following drug withdrawal (Fig. 3). Methylation-retaining cells tended to divide less

frequently during treatment (Fig. 1B) but had higher self-renewal and proliferative capacity than

hypomethylated cells after DAC treatment (Fig. 3). Interestingly, this selection for methylation-

retaining cells was not observed following AZA treatment, reflecting previous reports of distinct

effects of these HMAs[51-53]. Since AZA is incorporated into both DNA and RNA[54], while DAC is

restricted to DNA, we speculate that RNA-mediated toxicities (such as translational inhibition[25,26])

could prevent the growth of highly methylated cells following AZA treatment. Together, our results

suggest that methylation-retaining cells are a likely source of AML relapse, especially following DAC

therapy.

We observed upregulated cholesterol biosynthesis, particularly in colonies that had high DNA

methylation after HMA treatment (Fig. 5 and Supplementary Fig. 4). This indicates that HMA

exposure causes transcriptional changes and metabolic alterations, even in highly methylated cells.

Previous studies have shown that HMAs perturb the homeostasis of pyrimidine metabolism

independently of DNA methylation changes[24], and similar effects have been linked to altered

cholesterol and lipid metabolism in AZA-treated liver cell lines[55]. Therefore, we speculate that

upregulation of cholesterol biosynthesis is a delayed response to HMA treatment that occurs in

highly methylated cells, potentially via altered pyrimidine metabolism.

Increased cholesterol demand is an established feature of AML cells[56], and upregulation of

cholesterol biosynthesis genes has shown prognostic value in AML[57-59]. While the precise

mechanisms by which cholesterol confers a survival advantage remain unclear, inhibiting cholesterol

biosynthesis has been shown to sensitize AML cells to radiation and chemotherapy[60,61]. In addition,

we have demonstrated that inhibition of cholesterol biosynthesis by rosuvastatin can decrease the

self-renewal capacity and global DNA methylation levels of MOLM-13 colonies when combined with

DAC treatment (Fig. 5D, E). This suggests that the upregulation of cholesterol biosynthesis facilitates

the self-renewal and proliferation of cells that retain DNA methylation during HMA treatment.

Recently, cholesterol metabolism was also linked to DAC resistance in AML cell lines, with statin co-

treatment showing synergistic inhibition of *in vitro* AML cell growth[62]. *In vivo*, we observed

significantly improved survival of leukaemia-bearing mice treated with DAC and rosuvastatin (Fig.

5F), suggesting that co-treatment may increase the duration of HMA response in some AML and

MDS patients.

Encouragingly, a recent retrospective analysis of MDS patients (including some HMA-treated high-

risk cases) reported improved survival and reduced progression to AML for patients who

commenced statin treatments three months before or after MDS diagnosis[63]. Current clinical trials

are testing the safety of a statin (pitavastatin) in combination with AZA and venetoclax in AML

patients[64] and our results imply that DAC and statin co-treatments should also be assessed. This

therapeutic avenue is of particular interest since statins are commonly prescribed, well-tolerated,

oral medications, which could be rapidly repositioned for use in AML and MDS patients receiving
HMA therapy.

Methods

Cell lines and culture

AML cell lines, HL-60 (ATCC #CCL-240), MOLM-13 (DSMZ #ACC-554), and MV-4-11 (ATCC #CRL-9591)
were maintained in tissue culture flasks (Greiner Bio-One) at 37°C, 5% CO_2, and sub-cultured at
500,000 cells/mL every 2-3 days with fresh medium. HL-60 cells were maintained in Iscove's
Modified Dulbecco's media (IMDM; Sigma-Aldrich) supplemented with 4mM GlutaMAX (Thermo
Fisher Scientific) and 10% Fetal Bovine Serum (FBS; Sigma-Aldrich). MOLM-13 and MV-4-11 cells
were maintained in Roswell Park Memorial Institute 1640 media (RPMI; Sigma-Aldrich)
supplemented with 2mM GlutaMAX and 10% FBS. All cell lines were mycoplasma negative based on
routine testing using MycoAlert Mycoplasma Detection Kit (Lonza). Cell line authentication was
routinely performed by the Australian Genome Research Facility (AGRF).

CellTrace staining

Prior to drug treatment (Day 0), AML cells ($2x10^6$ cells/mL) were stained with 1µM (MOLM-13 and
MV-4-11) or 3µM (HL-60) CellTrace Far Red (Thermo Fisher Scientific) to monitor cell divisions,
according to manufacturer's instructions. Cells with uniformly high CellTrace fluorescence
underwent fluorescence activated cell sorting (FACS) prior to treatment with hypomethylating
agents (HMAs) in MOLM-13 and MV-4-11 cell lines, whereas all cells were used for HL-60
treatments.

HMA treatments

AML cell lines were treated with HMAs at various doses in suspension culture, every 24h for 72h
total. All cell lines were treated with 100nM decitabine (DAC; Selleckchem #S1200), HL-60 cells were
treated with 2000nM azacytidine (AZA; Selleckchem # S1782), and MOLM-13 and MV-4-11 cells
were treated with 500nM AZA. Untreated cells (UNT) were given an equal volume of 0.1% DMSO in
UltraPure™ DNase/RNase-Free Distilled Water (Thermo Fisher Scientific) and served as a negative
control. After HMA treatment, cells were prepared for FACS or colony forming assays, as described
below.

Fluorescence activated cell sorting (FACS)

455 HMA treated cells were stained with propidium iodide (PI, 1.5µg/mL) and prepared for FACS. Viable
456 (PI⁻) single cells were sorted into 2.5µL of RLT PLUS buffer (Qiagen) containing 2.5U SUPERas-In
457 (Thermo Fisher Scientific) in LoBind 96-well full skirted plates (Eppendorf) using indexed sorting on a

458   FACS Aria II (BD Biosciences). Plates were sealed and briefly centrifuged before storage at -80°C for

459   sequencing analyses.

Colony-forming assays

HMA treated cells were seeded in MethoCult Optimum (H4034; STEMCELL Technologies Inc.) at 500

cells/well in 6-well plates, with rosuvastatin (Selleckchem # S2169) added to the MethoCult at

various doses (0, 1, 10, 30µM). Cells were then cultured at 37°C, 5% CO_2 for 14 days to allow colony

formation. Wells containing colonies were imaged using Cytation3 (Biotek). Colony counts and sizes

were analysed using ImageJ software. Individual colonies were manually plucked using a 20µL

pipette tip into 100µL of media, centrifuged at 200xg for 5 mins, and then resuspended in 20µL of

RLT PLUS buffer before storage at -80°C. Alternatively, all colonies in each well were collected by

resuspending the MethoCult Optimum media (and colonies) in 3mL of standard culture media

(IMDM or RPMI), centrifuging at 200xg for 5 mins, and resuspending the cell pellet in 20-50µL RLT

PLUS buffer, before storage at -80°C.

Library preparation and sequencing

scNMT-seq library preparation and sequencing

For scNMT-seq, matched scNOMe-seq and scRNA-seq libraries were prepared from sorted HL-60

single cells, as previously described[28]. Minor modifications to the published protocol are in the

Supplementary Methods.

For scNOMe-seq libraries, paired-end 150bp sequencing was performed on the NovaSeq (Illumina)

platform. For scRNA-seq libraries, paired-end 75bp sequencing was performed on the NovaSeq or

NextSeq (Illumina) platform.

scTEM-seq library preparation and sequencing

480   For scTEM-seq analysis of global DNA methylation levels in single MOLM-13 and MV-4-11 cells,

481   library preparation was performed as previously described[65].

482   Paired-end 150bp sequencing was performed on the MiSeq (Illumina) platform.

Colony TEM-seq and RNA-seq library preparation and sequencing

Single colony TEM-seq (Fig. 3B, C) and parallel RNA-seq analysis was performed as described[65,66] with

minor modifications. Lysates from single colonies (HL-60: 7.5µL; MOLM-13 and MV-4-11: 2.5µL)

were used to separate genomic DNA and mRNA. During single colony TEM-seq library preparation,

the number of SINE Alu amplification cycles was reduced to 29. For RNA-seq analysis, 15 cycles of

cDNA amplification were used.

TEM-seq analysis of pooled colonies (Fig. 3D, 5E) was performed as described[65,66] using 5-10μL of cell

lysate as input for bisulphite conversion, and 29 cycles for SINE Alu amplification.

All TEM-seq libraries were sequenced using 150bp paired-end sequencing on the MiSeq (Illumina)

platform. For RNA-seq libraries, paired-end 75bp sequencing was performed on the NextSeq or

NovaSeq platform.

Data processing

495 Sequencing data were processed and aligned as described in the Supplementary Methods.

scNMT-seq data analysis

Quality control

For scNMT-seq data, cells were required to pass both scNOMe-seq and scRNA-seq quality control

(QC). Cells that had less than 500,000 CpG sites covered, less than 5,000,000 GpC sites covered,

greater than 15% CHH methylation rate, or less than 2% GpC methylation, failed scNOMe-seq QC.

For scRNA-seq, QC was performed using bam files from hisat2, and the SeqMonk[67] (v1.47.1) 'RNA-

seq QC Plot'. Cells that had less than 70% reads in exons, or less than 15% genes measured, failed

scRNA-seq QC. In total, 222 scNMT-seq samples passed QC (Supplementary Table 1).

scNOMe-seq normalisation and batch correction

scNOMe-seq libraries provide information on both DNA methylation (CpG sites) and DNA

accessibility (GpC sites). For both CpG (methylation) and GpC (accessibility) datasets, several

genomic annotation contexts were considered: introns, exons, intergenic regions, CpG islands,

promoters (-1500 to +500 bp of transcription start sites), H3K4me3 sites (ENCODE[68-70] accession ID:

ENCFF021JBH, experiment: ENCSR000DUO) and H3K27ac sites (ENCODE[68-70] accession ID:

ENCFF763UAG, experiment: ENCSR919WLM). In addition, unbiased 3kb windows of the whole

genome were generated with a step size of 1.5kb.

For DNA methylation, the CpG methylation rate was estimated within each annotation window using

the Bayes binomial approximation as in Smallwood et. al.[71].

The GpC methylation, which marks accessible DNA in scNOMe-seq libraries, is introduced in vitro

using a bacterial GpC methyltransferase enzyme (Supplementary Methods). To remove batch effects

resulting from differences in enzymatic activity, data normalisation and batch correction were

performed as follows. GpC data for the whole genome was aggregated in windows of 500kb in

length with 250kb overlap separately for methylated and unmethylated GpC counts for each cell.

Per-cell pooled size factors were computed from these 500kb windows using the method of Lun et.

al.[72] scaling by total library size, as implemented in the single-cell R package "scuttle" (v1.8.4)[73].

Batch scaling factors were estimated from corrected methylated and unmethylated window log

counts using the rescaleBatches method from the R package "batchelor" (v1.14.1)[74]. Per cell

methylated and unmethylated cell scaling factors were calculated as the ratio of the batch-corrected

sum of counts to the mean sum of counts across cells. Finally, unscaled methylated and

unmethylated counts in each cell were independently scaled by the product of the cell pooled size

factor and the methylated/unmethylated count batch correction factor, respectively. GpC

methylation rate for each annotation window was then computed using the normalised batch

corrected counts by the Bayesian binomial approximation.

From the overall distribution of counts across in the annotation layer for CpG and GpC methylation

data, minimum total count thresholds per window of 5 counts (CpG) and 20 counts (GpC) were

established and applied to discard windows with unreliable methylation rate estimation.

Pairwise distance analysis of DNA methylation heterogeneity

To assess the DNA methylation heterogeneity per treatment group and genomic context (Fig. 1D),

pairwise CpG methylation distance analysis was performed. The mean absolute methylation

difference was computed for each cell pair (A, B) as the mean of the absolute difference in

methylation rate at each common cytosine position in the relevant genomic annotation. To make

the comparison of methylation patterns meaningful, only cytosine loci with data in both cells in the

pair were used. These mean absolute methylation differences were grouped by the treatment

combination of the cell pair. The global summaries shown in Fig. 1D corresponds to the groups

where both cells in the pair had the same treatment. Higher values indicate a more heterogeneous

methylation pattern when cells in the same treatment group are compared vis-a-vis.

Cell-wise correlation analysis

To assess the relationships between DNA methylation, DNA accessibility and gene expression within

individual cells (Fig. 1E), Pearson correlations were computed using HL-60 scNMT-seq data. For this

analysis, RNA-seq data was normalised and log transformed per batch using

'scuttle::logNormCounts()' (v1.6.2)[73] without batch correction or prior count filtering. DNA

methylation was correlated to DNA accessibility at matched loci based on genomic coordinates. For

correlations with gene expression, methylation and accessibility measurements at promoters,

introns and exons were matched to the corresponding transcript. For CpG islands, H3K27ac sites and

H3K4me3 sites, methylation and accessibility measurements were matched to all transcripts within

10kb. For intergenic regions and 3kb genomic windows, methylation and accessibility measurements

were matched to every transcript within 1bp to assess the expression of immediately adjacent

genes. For each cell, Pearson correlation estimates were then computed using all matched values

and the cor.test function in R.

scRNAseq normalisation and batch correction

scRNA-seq libraries from HL-60 scNMT-seq data were filtered to remove lowly expressed genes,

requiring at least 5 counts in 10% of cells. Normalisation and variance stabilisation were performed

by scTransform[75] and batch correction by anchor-based integration using the R package "Seurat"

(v4.2.0)[76]. First, batches were independently normalised by scTransform. The top 5,000 most

variable features that were in common across batches were identified to determine anchors for

integration and batch correction of the data (using default parameter and k.weigh=50), applied to

and retaining those 5,000 commonly variable features. Finally, a sparse RNA-seq matrix was utilised,

whereby gene imputation calculations were ignored and removed by reintroducing 'NAs' in place of

genes with originally 'missing data' (zeros). Downstream analyses considered only autosomal genes

(Chr1-22).

Integrative sparse partial least squares (sPLS) analysis

Mixomics[38] (v6.20.0) was used to perform a multivariate integrative analysis of HL-60 scNMT-seq

data (Fig. 2A-D). Feature selection was performed to identify variably expressed transcripts that are

highly correlated to changes in DNA Methylation and accessibility following HMA treatment. We

performed an unsupervised sparse Partial Least Squares (sPLS) analysis using the function

'mixOmics::mint.block.spls()' which combines a multivariate integrative (MINT) method and a

multiblock sPLS integrative analysis. MINT[77] accounts for multiple batches (Supplementary Table 1)

measured on the same variables, while the multiblock sPLS seeks for correlated patterns between

DNA methylation and DNA accessibility rates that are split into multiple genomic regions ('blocks')

and explain (correlated to) the predictor (transcriptome).

To focus on transcriptomic and epigenetic changes resulting from HMA treatment only treated cells

(AZA and DAC) were included in the sPLS model. The genomic regions included in this analysis for

both DNA methylation and DNA accessibility were CpG islands, promoters, H3K27ac sites, H3K4me3

sites and 3kb windows. The rates from these genomic regions were filtered to retain only those

detected in greater than 10% of cells. The sPLS model was implemented assessing 2

components, selecting 50 features per component and per block (genomic region) in the

DNA methylation and DNA accessibility data sets, and 100 genes per component in the

transcriptome data set. More details are provided in our GitHub page.

Heatmap visualisation and identification of cell and expression clusters from sPLS selected features

was performed using ComplexHeatmap[78] (v2.12.1). sPLS selected features for components 1 and 2

were extracted using the function 'mixOmics::selectVar()'. Heatmap visualisation was performed on

sPLS selected features using transcriptomic (converted to z-score) and epigenetic rates (mean of

features in genomic regions and converted to z-score) that were entered into the model and

included both treated and untreated cells. K-means clustering was performed on sPLS selected

transcriptomic features, first on Gene Expression (row_km=3) followed by Cell Group

(column_km=4).

sPLS sample projections (Fig. 2A, C and D) were plotted using ggplot2[79] (v3.3.6) by extracting the

sPLS components 1 and 2 for a given block (RNA or epigenetic genomic region) and overlayed with

relevant information i.e. cell group identified from k-means clustering and treated cell type or

average DNA methylation.

Gene Ontology (GO) Over Representation Analysis (ORA) was calculated using clusterProfiler[80]

(v4.4.4) for 'biological process' and displayed using enrichplot[81] V1.16.1 (Fig. 2E, F). Gene Expression

k-means clusters (Fig. 2D) and sPLS selected features per component (1-2) were assessed by

'enrichGO(p.adj=0.05, p.adj.method = "fdr", q.val.threshold = 0.4)' and the list of genes from the

batch corrected transcriptome dataset (entered into the sPLS model) as the background. Results

were displayed as treeplots using default settings for pairwise 'termsim()' and 'treeplot(nCluster=5,

showCategory = 10)'.

The correlation of sPLS features (Supplementary Fig. 2) was calculated as a similarity matrix using

'mixOmics::circosPlot()' on the sPLS model. The results were displayed using ComplexHeatmap

showing DNA methylation and DNA accessibility features related to transcript features split by the

previously identified Gene Expression k-means clusters.

Locus-specific correlation analysis

To compare gene expression to adjacent epigenetic features, locus-specific correlations were

performed using HL-60 scNMT-seq data (Fig. 2H and I). DNA methylation and DNA accessibility

measurements were paired to genes based on annotation (Promoters) or by strand-aware position

within 10kbp of the gene transcription start site (CpG islands, H3K27ac sites, H3K4me3 sites and 3kb

windows). For paired sites, Pearson correlations were computed between CpG or GpC methylation

rate and log gene expression values. All cells with data (i.e. UNT, DAC and AZA groups) were

combined in these correlations, and a minimum of 22 cells with paired data (i.e. both gene

expression and DNA methylation/accessibility measurements) were required for the correlation to

be performed.

Colony sequencing analysis

RNA-seq quality control, normalisation and batch correction

Samples were excluded if they had less than 35% genes measured, or less than 70% reads in exons for HL-60 and MOLM-13 samples, or less than 65% reads in exons for MV-4-11 samples. RNA-seq data from single colonies were filtered to remove lowly expressed genes, requiring at least 5 counts in 3 samples. For each cell line, normalisation was performed by 'scuttle::logNormCounts()' (v1.6.2) and batch corrected using mutual nearest neighbours method by 'batchelor::mnnCorrect()' (v1.12.3) with default parameters. Downstream analyses considered only autosomal genes (Chr1-22).

Highly variable gene analysis

For Figure 4, highly variable genes (HVGs) were identified from colony RNA-seq data and PCA was performed using 'scater::calculatePCA(ntop = 2,000)'. K-means clusters of HVGs was determined using the r stats package (v4.2.1) with 'kmeans(centers = 8, iter.max = kmeans.iter, nstart = 50)'. Heatmapping of HVGs and k-means cluster was performed using 'ComplexHeatmap::pheatmap()' with z-scored values and Euclidean distance hierarchical clustering within row clusters (k-means groups) and columns (samples) ordered by treatment and descending average global methylation level.

GO ORA of k-means clusters was compared using clusterProfiler for 'biological process' by 'compareCluster(pAdjustMethod = "fdr", p.adj.threshold = 0.05, qvalueCutoff=0.4)' and the full list of genes from the batch corrected dataset (for each cell type) as the background list. Plots were created using 'clusterProfiler::dotplot(showCategory = 3) + coord_flip()'.

Correlation analysis

Pearson correlations comparing gene expression to mean global methylation in DAC HL-60 colonies (Fig. 5A) were performed using 'cor.test()' and underwent Benjamini–Hochberg false discovery rate adjustment using 'p.adjust(method="BH")'. Gene clustering and heatmap visualisation was performed on significantly correlated genes (p.adj ≤ 0.05 & cor.value.estimate ≤ -0.4 or cor.value.estimate ≥ 0.4). The average expression of each gene was calculated per treatment group with DAC split into high (≥75%) and low (<75%) global methylation groups. The R package "pheatmap"[82] (v1.0.12) was used to plot the mean centred treatment group average expression levels with rows aggregated into 4 'kmeans_k' clusters. The genes from each 'Kmeans_k' cluster was extracted and underwent GO ORA for biological process individually using 'enrichGO()' with fdr adjustment and results displayed as treeplots.

AML cell-line xenograft model

All experimental procedures were reviewed, approved, and carried out according to the Animal Care and Ethics Committee of the University of Newcastle (approval number: A-2023-303), and with consideration of the ARRIVE guidelines (Supplementary Methods).

MOLM-13 cells transduced with firefly luciferase (MOLM-13-luc) were a kind gift from Dr Charles de Bock (Children's Cancer Institute, UNSW Sydney). Five-week-old female NOD.Cg-Prkdc scid Il2rg tm1Wjl /SzJ (NSG) mice were obtained from the Australian Bioresources (ABR, Moss Vale, NSW, Australia) and were acclimatised for one week prior to any experimental procedure. The NSG mice were inoculated with MOLM-13-luc cells ($5x10^5$ cells suspended in 100 µL of PBS) by injection into the lateral tail vein. Tumour burden was assessed by bioluminescence imaging (BLI) twice a week using an IVIS Spectrum *in vivo* imaging system (PerkinElmer, Waltham, MA, USA), following intraperitoneal injections of 150 mg/kg D-luciferin (Promega, Alexandria, NSW, Australia) and under anaesthesia with isoflurane. Treatments commenced on day 6 post-engraftment, when a positive luminescence signal was detected.

In a pilot experiment, three different doses of rosuvastatin were co-administered with DAC. Mice (*n* = 5 per group) were treated by intraperitoneal injection of either vehicle (2% DMSO, 30% PEG300 in water), DAC (0.2 mg/Kg in saline), or rosuvastatin (1 mg/kg, 10 mg/kg, or 40 mg/kg in 30% PEG300 in water) combined with DAC (0.2 mg/Kg) once a day (5 days on, 2 days off) for up to 3 weeks. The animals were monitored until they reached ethical endpoint.

In a second experiment, mice (*n*= 15 per group) received vehicle (2% DMSO, 30% PEG300 in water), DAC (0.2 mg/Kg), rosuvastatin (1 mg/Kg), or DAC + rosuvastatin treatments via intraperitoneal injections once a day (5 days on, 2 days off) for up to 3 weeks, and mice were monitored until ethical endpoint.

Survival analyses were performed using Kaplan-Meier analysis followed by the Log-rank (Mantel-Cox) test. All statistical analyses were performed using GraphPad Prism v. 9.0 (GraphPad Software, La Jolla, CA, USA). An adjusted *p*-value of < 0.05 was considered statistically significant.

Data and code availability

Sequencing data and analysis code will be made available upon reasonable request and published following peer review.

References

1　Stomper, J., Rotondo, J. C., Greve, G. & Lübbert, M. Hypomethylating agents (HMA) for the
treatment of acute myeloid leukemia and myelodysplastic syndromes: mechanisms of
resistance and novel HMA-based therapies. *Leukemia* (2021).
https://doi.org:10.1038/s41375-021-01218-0

2　Jones, P. A. & Taylor, S. M. Cellular differentiation, cytidine analogs and DNA methylation.
Cell **20**, 85-93 (1980). https://doi.org:http://dx.doi.org/10.1016/0092-8674(80)90237-8

3　Patel, K. *et al.* Targeting of 5-aza-2′-deoxycytidine residues by chromatin-associated DNMT1
induces proteasomal degradation of the free enzyme. *Nucleic Acids Research* **38**, 4313-4324
(2010). https://doi.org:10.1093/nar/gkq187

4　Mayyas, I. M. *et al.* Hairpin-bisulfite sequencing of cells exposed to decitabine documents
the process of DNA demethylation. *Epigenetics*, 1-9 (2020).
https://doi.org:10.1080/15592294.2020.1861169

5　Issa, J. P. *et al.* Safety and tolerability of guadecitabine (SGI-110) in patients with
myelodysplastic syndrome and acute myeloid leukaemia: a multicentre, randomised, dose-
escalation phase 1 study. *Lancet Oncol* **16**, 1099-1110 (2015).
https://doi.org:10.1016/S1470-2045(15)00038-8

6　Fenaux, P. *et al.* Efficacy of azacitidine compared with that of conventional care regimens in
the treatment of higher-risk myelodysplastic syndromes: a randomised, open-label, phase III
study. *Lancet Oncol* **10**, 223-232 (2009). https://doi.org:10.1016/s1470-2045(09)70003-8

7　Jabbour, E. *et al.* Randomized phase 2 study of low-dose decitabine vs low-dose azacitidine
in lower-risk MDS and MDS/MPN. *Blood* **130**, 1514-1522 (2017).
https://doi.org:10.1182/blood-2017-06-788497

8　Fenaux, P. *et al.* Azacitidine prolongs overall survival compared with conventional care
regimens in elderly patients with low bone marrow blast count acute myeloid leukemia. *J*
Clin Oncol **28**, 562-569 (2010). https://doi.org:10.1200/jco.2009.23.8329

9　Kantarjian, H. M. *et al.* Survival advantage with decitabine versus intensive chemotherapy in
patients with higher risk myelodysplastic syndrome: comparison with historical experience.
Cancer **109**, 1133-1137 (2007). https://doi.org:10.1002/cncr.22508

10　Kantarjian, H. M. *et al.* Multicenter, randomized, open-label, phase III trial of decitabine
versus patient choice, with physician advice, of either supportive care or low-dose
cytarabine for the treatment of older patients with newly diagnosed acute myeloid
leukemia. *J Clin Oncol* **30**, 2670-2677 (2012). https://doi.org:10.1200/jco.2011.38.9429

11　Lee, J. H. *et al.* Comparison of 7-day azacitidine and 5-day decitabine for treating
myelodysplastic syndrome. *Ann Hematol* **92**, 889-897 (2013).
https://doi.org:10.1007/s00277-013-1702-8

12　Dombret, H. *et al.* International phase 3 study of azacitidine vs conventional care regimens in
older patients with newly diagnosed AML with >30% blasts. *Blood* **126**, 291-299 (2015).
https://doi.org:10.1182/blood-2015-01-621664

13　Helbig, G. *et al.* Real Life Data on Efficacy and Safety of Azacitidine Therapy for
Myelodysplastic Syndrome, Chronic Myelomonocytic Leukemia and Acute Myeloid
Leukemia. *Pathology & Oncology Research* **25**, 1175-1180 (2019).
https://doi.org:10.1007/s12253-018-00574-0

14　Stahl, M. *et al.* Hypomethylating agents in relapsed and refractory AML: outcomes and their
predictors in a large international patient cohort. *Blood Adv* **2**, 923-932 (2018).
https://doi.org:10.1182/bloodadvances.2018016121

15　Fennell, K. A., Bell, C. C. & Dawson, M. A. Epigenetic therapies in acute myeloid leukemia:
where to from here? *Blood* **134**, 1891-1901 (2019).
https://doi.org:10.1182/blood.2019003262

16 DiNardo, C. D. *et al.* Venetoclax combined with decitabine or azacitidine in treatment-naive,
elderly patients with acute myeloid leukemia. *Blood* **133**, 7-17 (2019).
https://doi.org:10.1182/blood-2018-08-868752
17 DiNardo, C. D. *et al.* Azacitidine and Venetoclax in Previously Untreated Acute Myeloid
Leukemia. *New England Journal of Medicine* **383**, 617-629 (2020).
https://doi.org:10.1056/NEJMoa2012971
18 Tsai, H. C. *et al.* Transient low doses of DNA-demethylating agents exert durable antitumor
effects on hematological and epithelial tumor cells. *Cancer Cell* **21**, 430-446 (2012).
https://doi.org:10.1016/j.ccr.2011.12.029
19 Craddock, C. *et al.* Azacitidine fails to eradicate leukemic stem/progenitor cell populations in
patients with acute myeloid leukemia and myelodysplasia. *Leukemia* **27**, 1028-1036 (2013).
https://doi.org:10.1038/leu.2012.312
20 Unnikrishnan, A. *et al.* Integrative Genomics Identifies the Molecular Basis of Resistance to
Azacitidine Therapy in Myelodysplastic Syndromes. *Cell Rep* **20**, 572-585 (2017).
https://doi.org:10.1016/j.celrep.2017.06.067
21 Riether, C. *et al.* Targeting CD70 with cusatuzumab eliminates acute myeloid leukemia stem
cells in patients treated with hypomethylating agents. *Nature Medicine* **26**, 1459-1467
(2020). https://doi.org:10.1038/s41591-020-0910-8
22 Pabst, T. *et al.* Results from a phase I/II trial of cusatuzumab combined with azacitidine in
patients with newly diagnosed acute myeloid leukemia who are ineligible for intensive
chemotherapy. *Haematologica* **108**, 1793-1802 (2023).
https://doi.org:10.3324/haematol.2022.281563
23 Zeidan, A. M. *et al.* Clinical outcomes of older patients with AML receiving hypomethylating
agents: a large population-based study in the United States. *Blood Adv* **4**, 2192-2201 (2020).
https://doi.org:10.1182/bloodadvances.2020001779
24 Gu, X. *et al.* Decitabine- and 5-azacytidine resistance emerges from adaptive responses of
the pyrimidine metabolism network. *Leukemia* **35**, 1023-1036 (2021).
https://doi.org:10.1038/s41375-020-1003-x
25 Aimiuwu, J. *et al.* RNA-dependent inhibition of ribonucleotide reductase is a major pathway
for 5-azacytidine activity in acute myeloid leukemia. *Blood* **119**, 5229-5238 (2012).
https://doi.org:10.1182/blood-2011-11-382226
26 Diesch, J. *et al.* Inhibition of CBP synergizes with the RNA-dependent mechanisms of
Azacitidine by limiting protein synthesis. *Nature Communications* **12**, 6060 (2021).
https://doi.org:10.1038/s41467-021-26258-z
27 Gruber, E., Franich, R. L., Shortt, J., Johnstone, R. W. & Kats, L. M. Distinct and overlapping
mechanisms of resistance to azacytidine and guadecitabine in acute myeloid leukemia.
Leukemia **34**, 3388-3392 (2020). https://doi.org:10.1038/s41375-020-0973-z
28 Clark, S. J. *et al.* scNMT-seq enables joint profiling of chromatin accessibility DNA
methylation and transcription in single cells. *Nature Communications* **9**, 781 (2018).
https://doi.org:10.1038/s41467-018-03149-4
29 Daskalakis, M. *et al.* Demethylation of a hypermethylated P15/INK4B gene in patients with
myelodysplastic syndrome by 5-Aza-2'-deoxycytidine (decitabine) treatment. *Blood* **100**,
2957-2964 (2002). https://doi.org:10.1182/blood.V100.8.2957
30 Kiziltepe, T. *et al.* 5-Azacytidine, a DNA methyltransferase inhibitor, induces ATR-mediated
DNA double-strand break responses, apoptosis, and synergistic cytotoxicity with doxorubicin
and bortezomib against multiple myeloma cells. *Molecular Cancer Therapeutics* **6**, 1718-
1727 (2007). https://doi.org:10.1158/1535-7163.Mct-07-0010
31 Dunshee, D. R. *et al.* CC-486 Mechanism Imparted By Extended Exposure of Azacitidine
Upregulates Myeloid Differentiation Markers and Induces Cell Death in AML Cells. *Blood*
136, 33-34 (2020). https://doi.org:https://doi.org:10.1182/blood-2020-137606

32   Momparler, R. L., Bouchard, J. & Samson, J. Induction of differentiation and inhibition of
777   DNA methylation in HL-60 myeloid leukemic cells by 5-AZA-2▯-deoxycytidine. *Leukemia*
778   *Research* **9**, 1361-1366 (1985). https://doi.org:https://doi.org/10.1016/0145-
779   2126(85)90123-7
780   33   Goodyear, O. *et al.* Induction of a CD8+ T-cell response to the MAGE cancer testis antigen by
781   combined treatment with azacitidine and sodium valproate in patients with acute myeloid
782   leukemia and myelodysplasia. *Blood* **116**, 1908-1918 (2010). https://doi.org:10.1182/blood-
783   2009-11-249474
784   34   Nahas, M. R. *et al.* Hypomethylating agent alters the immune microenvironment in acute
785   myeloid leukaemia (AML) and enhances the immunogenicity of a dendritic cell/AML vaccine.
786   *Br J Haematol* **185**, 679-690 (2019). https://doi.org:10.1111/bjh.15818
787   35   Roulois, D. *et al.* DNA-Demethylating Agents Target Colorectal Cancer Cells by Inducing Viral
788   Mimicry by Endogenous Transcripts. *Cell* **162**, 961-973 (2015).
789   https://doi.org:10.1016/j.cell.2015.07.056
790   36   Liu, M. *et al.* Vitamin C increases viral mimicry induced by 5-aza-2'-deoxycytidine. *Proc Natl*
791   *Acad Sci U S A* **113**, 10238-10244 (2016). https://doi.org:10.1073/pnas.1612262113
792   37   Chiappinelli, Katherine B. *et al.* Inhibiting DNA Methylation Causes an Interferon Response in
793   Cancer via dsRNA Including Endogenous Retroviruses. *Cell* **162**, 974-986 (2015).
794   https://doi.org:10.1016/j.cell.2015.07.011
795   38   Rohart, F., Gautier, B., Singh, A. & Le Cao, K. A. mixOmics: An R package for 'omics feature
796   selection and multiple data integration. *PLoS Comput Biol* **13**, e1005752 (2017).
797   https://doi.org:10.1371/journal.pcbi.1005752
798   39   Deniz, Ö. *et al.* Endogenous retroviruses are a source of enhancers with oncogenic potential
799   in acute myeloid leukaemia. *Nature Communications* **11**, 3506 (2020).
800   https://doi.org:10.1038/s41467-020-17206-4
801   40   Huang, B., Song, B.-l. & Xu, C. Cholesterol metabolism in cancer: mechanisms and
802   therapeutic opportunities. *Nature Metabolism* **2**, 132-141 (2020).
803   https://doi.org:10.1038/s42255-020-0174-0
804   41   Teramoto, T. & Watkins, C. Review of efficacy of rosuvastatin 5 mg. *Int J Clin Pract* **59**, 92-
805   101 (2005). https://doi.org:10.1111/j.1742-1241.2005.00346.x
806   42   Hunt, K. V. *et al.* scTEM-seq: Single-cell analysis of transposable element methylation to link
807   global epigenetic heterogeneity with transcriptional programs. *Scientific Reports* **12**, 5776
808   (2022). https://doi.org:10.1038/s41598-022-09765-x
809   43   Takeshima, H. *et al.* Low-dose DNA demethylating therapy induces reprogramming of
810   diverse cancer-related pathways at the single-cell level. *Clinical Epigenetics* **12**, 142 (2020).
811   https://doi.org:10.1186/s13148-020-00937-y
812   44   Griffin, G. K. *et al.* Epigenetic silencing by SETDB1 suppresses tumour intrinsic
813   immunogenicity. *Nature* (2021). https://doi.org:10.1038/s41586-021-03520-4
814   45   Irwin, R. *et al.* UHRF1 suppresses viral mimicry through both DNA methylation-dependent
815   and -independent mechanisms. *bioRxiv*, 2020.2008.2031.274894 (2020).
816   https://doi.org:10.1101/2020.08.31.274894
817   46   Maria Teresa, V. *et al.* Why methylation is not a marker predictive of response to
818   hypomethylating agents. *Haematologica* **99**, 613-619 (2014).
819   https://doi.org:10.3324/haematol.2013.099549
820   47   Unnikrishnan, A. *et al.* AZA-MS: a novel multiparameter mass spectrometry method to
821   determine the intracellular dynamics of azacitidine therapy in vivo. *Leukemia* **32**, 900-910
822   (2018). https://doi.org:10.1038/leu.2017.340
823   48   Yang, A. S. *et al.* DNA methylation changes after 5-aza-2'-deoxycytidine therapy in patients
824   with leukemia. *Cancer Res* **66**, 5495-5503 (2006). https://doi.org:10.1158/0008-5472.CAN-
825   05-2385

826   49   Greve, G. *et al.* In vivo kinetics of early, non-random methylome and transcriptome changes
827   induced by DNA-hypomethylating treatment in primary AML blasts. *Leukemia* **37**, 1018-1027
828   (2023). https://doi.org:10.1038/s41375-023-01876-2

829   50   Li, S. *et al.* Somatic Mutations Drive Specific, but Reversible, Epigenetic Heterogeneity States
830   in AML. *Cancer Discovery* **10**, 1934-1949 (2020). https://doi.org:10.1158/2159-8290.Cd-19-
831   0897

832   51   Hollenbach, P. W. *et al.* A Comparison of Azacitidine and Decitabine Activities in Acute
833   Myeloid Leukemia Cell Lines. *PLOS ONE* **5**, e9001 (2010).
834   https://doi.org:10.1371/journal.pone.0009001

835   52   Aumer, T. *et al.* Comprehensive comparison between azacytidine and decitabine treatment
836   in an acute myeloid leukemia cell line. *Clinical Epigenetics* **14**, 113 (2022).
837   https://doi.org:10.1186/s13148-022-01329-0

838   53   Flotho, C. *et al.* The DNA methyltransferase inhibitors azacitidine, decitabine and zebularine
839   exert differential effects on cancer gene expression in acute myeloid leukemia cells.
840   *Leukemia* **23**, 1019-1028 (2009). https://doi.org:10.1038/leu.2008.397

841   54   Li, L. H., Olin, E. J., Buskirk, H. H. & Reineke, L. M. Cytotoxicity and mode of action of 5-
842   azacytidine on L1210 leukemia. *Cancer Res* **30**, 2760-2769 (1970).

843   55   Poirier, S. *et al.* The epigenetic drug 5-azacytidine interferes with cholesterol and lipid
844   metabolism. *J Biol Chem* **289**, 18736-18751 (2014).
845   https://doi.org:10.1074/jbc.M114.563650

846   56   Vitols, S., Norgren, S., Juliusson, G., Tatidis, L. & Luthman, H. Multilevel regulation of low-
847   density lipoprotein receptor and 3-hydroxy- 3-methylglutaryl coenzyme A reductase gene
848   expression in normal and leukemic cells. *Blood* **84**, 2689-2698 (1994).
849   https://doi.org:10.1182/blood.v84.8.2689.2689

850   57   Banker, D. E. *et al.* Cholesterol synthesis and import contribute to protective cholesterol
851   increments in acute myeloid leukemia cells. *Blood* **104**, 1816-1824 (2004).
852   https://doi.org:10.1182/blood-2004-01-0395

853   58   Bai, S. *et al.* Lipid profile as a novel prognostic predictor for patients with acute myeloid
854   leukemia. *Frontiers in Oncology* **13** (2023). https://doi.org:10.3389/fonc.2023.950732

855   59   Kuliszkiewicz-Janus, M., Małecki, R. & Mohamed, A. Lipid changes occuring in the course of
856   hematological cancers. *Cellular and Molecular Biology Letters* **13** (2008).
857   https://doi.org:10.2478/s11658-008-0014-9

858   60   Dimitroulakos, J. *et al.* Increased Sensitivity of Acute Myeloid Leukemias to Lovastatin-
859   Induced Apoptosis: A Potential Therapeutic Approach. *Blood* **93**, 1308-1318 (1999).
860   https://doi.org:10.1182/BLOOD.V93.4.1308

861   61   Li, H. Y., Appelbaum, F. R., Willman, C. L., Zager, R. A. & Banker, D. E. Cholesterol-modulating
862   agents kill acute myeloid leukemia cells and sensitize them to therapeutics by blocking
863   adaptive cholesterol responses. *Blood* **101**, 3628-3634 (2003).
864   https://doi.org:10.1182/blood-2002-07-2283

865   62   Yabushita, T. *et al.* Mitotic perturbation is a key mechanism of action of decitabine in
866   myeloid tumor treatment. *Cell Reports* **42**, 113098 (2023).
867   https://doi.org:https://doi.org/10.1016/j.celrep.2023.113098

868   63   Afzal, A., Fiala, M. A., Jacoby, M. A. & Walter, M. J. Statin use in myelodysplastic syndromes
869   is associated with a better survival and decreased progression to leukemia. *Blood Advances*
870   **7**, 3838-3841 (2023). https://doi.org:10.1182/bloodadvances.2023009818

871   64   Brem, E. A. *et al.* A Phase 1 Study of Adding Pitavastatin to Venetoclax-Based Therapy in AML
872   and CLL/SLL. *Blood* **142**, 5940-5940 (2023). https://doi.org:10.1182/blood-2023-188086

873   65   Hunt, K. V. *et al.* scTEM-seq: Single-cell analysis of transposable element methylation to link
874   global epigenetic heterogeneity with transcriptional programs. *Sci Rep* **12**, 5776 (2022).
875   https://doi.org:10.1038/s41598-022-09765-x

66 Hunt KV, B. S., Bond DR, Lee HJ. Protocol for targeted analysis of transposable element
methylation levels and transcriptome in single cells using scTEM-seq. *ProtocolExchange*
(2022). https://doi.org:10.21203/rs.3.pex-2075/v1
67 SeqMonk (Babraham Bioinformatics).
68 Benjamin, C. H. *et al.* The ENCODE Uniform Analysis Pipelines. *bioRxiv*,
2023.2004.2004.535623 (2023). https://doi.org:10.1101/2023.04.04.535623
69 An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**, 57-74
(2012). https://doi.org:10.1038/nature11247
70 Luo, Y. *et al.* New developments on the Encyclopedia of DNA Elements (ENCODE) data
portal. *Nucleic Acids Res* **48**, D882-d889 (2020). https://doi.org:10.1093/nar/gkz1062
71 Smallwood, S. A. *et al.* Single-cell genome-wide bisulfite sequencing for assessing epigenetic
heterogeneity. *Nature Methods* **11**, 817-820 (2014). https://doi.org:10.1038/nmeth.3035
72 L. Lun, A. T., Bach, K. & Marioni, J. C. Pooling across cells to normalize single-cell RNA
sequencing data with many zero counts. *Genome Biology* **17**, 75 (2016).
https://doi.org:10.1186/s13059-016-0947-7
73 McCarthy, D. J., Campbell, K. R., Lun, A. T. L. & Wills, Q. F. Scater: pre-processing, quality
control, normalization and visualization of single-cell RNA-seq data in R. *Bioinformatics* **33**,
1179-1186 (2017). https://doi.org:10.1093/bioinformatics/btw777
74 Haghverdi, L., Lun, A. T. L., Morgan, M. D. & Marioni, J. C. Batch effects in single-cell RNA-
sequencing data are corrected by matching mutual nearest neighbors. *Nature Biotechnology*
36, 421-427 (2018). https://doi.org:10.1038/nbt.4091
75 Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq
data using regularized negative binomial regression. *Genome Biology* **20**, 296 (2019).
https://doi.org:10.1186/s13059-019-1874-1
76 Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e3529
(2021). https://doi.org:10.1016/j.cell.2021.04.048
77 Rohart, F., Eslami, A., Matigian, N., Bougeard, S. & Lê Cao, K.-A. MINT: a multivariate
integrative method to identify reproducible molecular signatures across independent
experiments and platforms. *BMC Bioinformatics* **18**, 128 (2017).
https://doi.org:10.1186/s12859-017-1553-8
78 Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847-2849 (2016).
https://doi.org:10.1093/bioinformatics/btw313
79 Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New York, 2016).
80 Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *The*
Innovation **2**, 100141 (2021). https://doi.org:https://doi.org/10.1016/j.xinn.2021.100141
81 Yu, G. enrichplot: Visualization of Functional Enrichment Result. R package. (2022).
82 Kolde, R. & Kolde, M. R. Package 'pheatmap'. *R package* **1**, 790 (2015).

Acknowledgements

Nicole Cole (University of Newcastle) provided technical support for fluorescence activated cell

sorting. Al J. Abadi (Melbourne Integrative Genomics, University of Melbourne) assisted with the

implementation of sPLS analysis. The authors thank Professor Tri Phan (Garvan Institute) and

Professor Xu Dong Zhang (University of Newcastle) for critical review of the manuscript.

H.J.L. and C.R. received funding from the National Health and Medical Research Council of Australia
(NHMRC; GNT1143614, GNT1173892 and GNT2016283). K.A.L.C. and H.J.L. received funding from
the Australian Research Council (ARC; DP200102903). H.J.L. and H.C.M. received funding from the
Cancer Institute NSW (ECF171145 and ECF1299). D.R.B received funding from Cure Cancer Australia
Foundation (CCAF2023-Bond) and NSW Health Pathology North. N.A.B received funding from the
McGuigan family through the Hunter Medical Research Institute (HMRI778). The contents of the
published material are solely the responsibility of the research institutions involved or individual
authors and do not reflect the views of funding agencies.

Author Contributions

H.J.L. conceived and oversaw the project. D.R.B., A.K.E., C.R., K.A.L.C. and H.J.L. acquired funding.
D.R.B., K.U., K.V.H., B.H., C.L.O., E.A.R., S.H., L.S.R., H.M. and H.J.L. performed experiments. D.R.B.,
S.M.B., K.U., K.V.H., B.H., C.L.O., L.S.R., C.R. and H.J.L. performed data analysis. S.M.B. and C.R.
developed analytical approaches. S.M.B. managed sequencing data and analysis code. D.R.B., N.A.B.,
A.K.E, N.M.V., K.A.L.C. and H.J.L. supervised work. D.R.B., S.M.B., K.U., B.H., L.S.R., C.R. and H.J.L.
prepared figures and wrote the manuscript. All authors read and approved the manuscript.

Competing Interests

936 A.K.E. declares the following competing interests: Advisory board/ Honoraria from AbbVie, Astellas,
937 Gilead, Servier, Jazz, Otsuka/Astex and RACE Oncology; Speakers fees from AbbVie, Otsuka/Astex,
938 Astellas and Jazz; Research Funding from RACE oncology and Otsuka/Astex.

Correspondence

940 Correspondence should be directed to Dr Heather Lee and Dr Danielle Bond.