Hypomethylating agents induce epigenetic and transcriptional heterogeneity with implications for AML cell self-renewal

Danielle R Bond[1,2^*], Sean M Burnard[1,2^], Kumar Uddipto[1,2], Kooper V Hunt[1,2], Brooke M Harvey[1,2],

Luiza Steffens Reinhardt[1,3,9], Charley Lawlor-O'Neill[1], Ellise A Roper[1], Sam Humphries[1,2], Heather C.

Murray[1,2], Abdul Mannan[1,2], Matthew D Dun[1,2], Charles E de Bock[4,5], Nikola A Bowden[6,7], Anoop K

Enjeti[2,6,8,9], Nicole M Verrills[1,2], Carlos Riveros[6], Kim-Anh Lê Cao[10], Heather J Lee[1,2*]

[1]*School of Biomedical Science and Pharmacy, The University of Newcastle, Australia.*

[2]*Precision Medicine Research Program, Hunter Medical Research Institute, Australia.*

[3]*Cancer Detection and Therapy Research Program, Hunter Medical Research Institute, Australia.*

[4]*Children's Cancer Institute, Lowy Cancer Research Centre, Randwick, NSW 2031, Australia*

[5]*School of Clinical Medicine, Faculty of Medicine, UNSW Sydney, NSW 2052, Australia*

[6]*School of Medicine and Public Health, The University of Newcastle, Australia.*

[7]*Drug Repurposing and Medicines Program, Hunter Medical Research Institute, Australia.*

[8]*Department of Haematology, Calvary Mater Hospital, Australia.*

[9]*New South Wales Health Pathology, John Hunter Hospital, Australia.*

[10]*Melbourne Integrative Genomics, School of Mathematics and Statistics, The University of Melbourne, Australia.*

^ These authors contributed equally to this work; * Corresponding authors

Correspondence should be directed to A/Prof Heather J Lee: Heather.lee@newcastle.edu.au

Competing Interests

A.K.E. declares the following competing interests: advisory board and honoraria from AbbVie,

Astellas, Gilead, Servier, Jazz, Otsuka/Astex, and RACE Oncology; speaker fees from AbbVie,

25   Otsuka/Astex, Astellas and Jazz; Research Funding from RACE oncology and Otsuka/Astex. N.A.B.

26   receives research funding from Merck KGaA and Bristol Myers Squibb.

Abstract

DNA hypomethylating agents (HMAs) are used to treat acute myeloid leukemia (AML) and

myelodysplasia patients who are unsuitable for intensive chemotherapy. However, low response

rates and therapy-resistant relapse remain significant challenges. To improve outcomes, we must

understand how AML cells survive HMA treatment and continue to proliferate following therapy. We

combine single-cell multiomics with parallel colony-forming assays to link HMA-induced

heterogeneity with functional consequences in AML cellss. Azacytidine (AZA) and decitabine (DAC)

induced global epigenetic heterogeneity, associated with upregulation of inflammatory responses

and cell death pathways in a subset of hypomethylated cells. Some cells maintained high DNA

methylation levels during treatment, and these methylation-retaining cells had increased self-

renewal capacity following DAC treatment in two *FLT3-ITD* AML cell lines. Transcriptional profiling of

colonies formed after HMA treatment revealed many genes with altered expression in both

methylation-retaining and hypomethylated cells, with increased expression of cholesterol-related

genes observed in all cell lines. Inhibition of the cholesterol biosynthesis pathway by rosuvastatin

enhanced HMA effects on colony formation *in vitro* and extended survival in two *in vivo* models of

AML. Our study demonstrates that HMA-induced epigenetic heterogeneity has implications for AML

cell growth and identifies statins as a candidate co-treatment strategy to improve HMA efficacy.

Introduction

Hypomethylating agents (HMAs), such as azacytidine (AZA, 5-azacytidine) and decitabine (DAC, 2'-

deoxy-5-azacytidine), are used to treat patients with acute myeloid leukemia (AML) and

myelodysplastic syndrome (MDS) (1). AZA and DAC are cytidine analogs, which are incorporated into

DNA during replication (1, 2). This leads to the degradation of DNA methyltransferase (DNMT)

enzymes (3) and loss of DNA methylation in subsequent cell divisions (2, 4). HMA mechanisms of

action have been difficult to precisely define because genome-wide loss of methylation is associated

with pleiotropic transcriptional changes (5). HMA-induced promoter hypomethylation permits the

re-expression of tumor suppressor genes (6), re-activation of DNA repair pathways (7) and

upregulation of differentiation markers in AML cells (8). HMA treatment can also increase the

expression of cancer testis antigens (9) and enhance antigen presentation in cancer cells (10).

Furthermore, genome-wide de-repression of transposable elements (TEs) has been shown to trigger

a viral mimicry response, in which dsRNA stimulates interferon signaling and apoptosis (11-13).

Unlike DAC, AZA can also be incorporated into RNA, which influences transcript stability and

translation (14, 15).

While single-agent HMA treatment extends survival in many patients (16-18), only 20% show a

complete response (CR) to therapy (19). Responses are also short-lived (e.g., 8-15 months (20)), with

most patients developing HMA-resistant relapse. To address these limitations, clinical trials have

tested co-treatment strategies, with some delivering improved outcomes (5). For example, the pro-

apoptotic therapy venetoclax has increased response rates in elderly AML patients undergoing HMA

treatment (21). However, relapse remains a significant problem, with the median duration of

response being 11-18 months for patients treated with both venetoclax and HMA therapy (21, 22).

To improve the long-term benefits of HMA therapy, it is essential to understand how AML cells

survive treatment and proliferate to cause relapse.

In this study, we characterize HMA-induced molecular heterogeneity with implications for the self-

renewal capacity of AML cells. We perform single-cell multiomic analysis to reveal global HMA-

induced DNA methylation heterogeneity and methylation-retaining cells that appear to evade

treatment. In parallel colony-forming assays, we show that methylation-retaining cells have a growth

and self-renewal advantage following treatment with DAC, but not AZA, in two *FLT3-ITD* cell lines.

We also identify the upregulation of cholesterol-related genes in colonies formed after HMA

treatment, and identify statins as a candidate treatment strategy to enhance the long-term efficacy

of HMAs.

Results

DNA methylation heterogeneity induced by HMA treatment

We performed single-cell analysis of DNA methylation in AML cell lines (HL-60, MOLM-13, and MV-4-

11; Supplementary Table 1) treated with low doses of decitabine (DAC) or azacytidine (AZA) to

induce maximal demethylation with minimal effects on cell growth and viability (Supplementary Fig.

S1). After three days of HMA treatment, a striking level of DNA methylation heterogeneity was

observed (Fig. 1A; Supplementary Table 2). While untreated cells showed homogeneously high levels

of DNA methylation (e.g., HL-60: 65-73%), the extent of hypomethylation varied substantially among

cells treated with DAC (e.g., HL-60: 17-69%) or AZA (e.g., HL-60: 20-69%). Interestingly, a small

proportion (1-5%) of methylation-retaining cells (Fig. 1A, dashed boxes) displayed no evidence of

HMA-induced hypomethylation. The extent of HMA-induced hypomethylation was related to cell

division rate, as indicated by positive correlations between DNA methylation and CellTrace

fluorescence (Fig. 1B; Supplementary Fig. S2). This is consistent with HMA incorporation during

replication (2) and confirms that slowly dividing cells can avoid the effects of HMA treatment.

Genomic contexts with high levels of DNA methylation in untreated HL-60 cells (such as exons,

introns, and intergenic regions) showed the most pronounced reduction following HMA treatment

(Fig. 1C), with this loss of methylation accompanied by increased cell-to-cell heterogeneity (Fig. 1D).

In contrast, active promoters marked by H3K4me3 were the only genomic features to show a

significantly decreased DNA methylation heterogeneity after HMA treatment, likely due to their low

levels of DNA methylation in untreated cells.

To explore links between DNA methylation heterogeneity and other layers of genetic regulation, we

used multi-omic data collected from HL-60 cells. The scNMT-seq method profiles DNA methylation,

DNA accessibility, and gene expression in parallel (23), allowing these molecular modalities to be

correlated with each other across the genome of individual cells (Fig. 1E). Untreated cells showed

the expected trends, with DNA methylation being negatively correlated with accessibility and gene

expression, whereas DNA accessibility and transcription were positively correlated. In HMA-treated

cells, the relationship between DNA methylation and accessibility was significantly weakened in all

genomic contexts. HMA treatment also significantly weakened the correlation between DNA

methylation (in promoters, exons, and introns) and expression of associated genes. In contrast, HMA

treatment had a minimal effect on the association between DNA accessibility and gene expression.

These observations imply that loss of DNA methylation is not always accompanied by increased

accessibility and transcription.

Transcriptional programs linked to HMA-induced epigenetic heterogeneity

To clarify how HMA-induced epigenetic heterogeneity influences transcriptional responses, we

performed a multivariate analysis to integrate the three molecular layers: DNA methylation, DNA

accessibility, and transcription. We excluded untreated cells to focus on variability among HMA-

treated cells and applied an unsupervised sparse Partial Least Squares (sPLS) method (27). This

approach was used to identify variably expressed transcripts that are highly correlated to changes in

DNA methylation and accessibility in regulatory regions (CpG islands, promoters, H3K4me3 sites, and

H3K27ac sites) and 3kb genomic windows.

The 200 transcript features selected by sPLS were further examined in both treated and untreated

cells, revealing 4 groups of cells with distinct transcriptional profiles (Fig. 2A, B). A significant and

negative correlation ($r = -0.37$, $p < 1 \times 10^{-7}$) between TE LINE:L2a expression and component 1 was

observed (Fig. 2C; Supplementary Fig. S3A), consistent with activation of viral mimicry in cell group

3. However, there was no significant relationship between global DNA methylation and component

1, and cell group 3 had similar methylation levels to other HMA-treated cells (Fig. 2D; Supplementary

Fig. S3D). This indicates that global methylation levels are insufficient to explain the transcriptional

responses of HMA-treated AML cells.

Gene ontology (GO) over-representation analysis (ORA) was performed for each of the three clusters

of transcript features (Fig. 2B; Supplementary Table 3). Cell group 3 had low expression of genes in

cluster 3, which were related to translation and inhibition of cell death (Fig. 2E; Supplementary Table

4). Cell group 3 also had high expression of genes in cluster 1, which were enriched in terms related

to immune inflammatory response and positive regulation of cell death (Fig. 2F; Supplementary

Table 4). This transcriptional profile is consistent with the expected effects of HMA treatment (1),

and 29 of the 78 genes in cluster 1 were significantly upregulated by DAC and/or AZA in matched

bulk RNA sequencing (RNA-seq) data (Supplementary Table 3). In contrast, cell group 1 displayed an

inverted gene expression pattern and included many untreated cells (observed:expected ratio =

1.48). This suggests that cell group 1 did not activate transcriptional pathways commonly associated

with HMA treatment, despite low methylation levels in most cells (Fig. 2A, D). The expression profile

of cell group 2 was intermediate between groups 1 and 3, whereas cell group 4 showed distinctive

upregulation of genes expressed in cluster 2. No ontology terms from cluster 2 retained significance

after multiple testing correction.

The sPLS model also selected epigenetic features that were correlated to the 200 transcript features,

without considering the direction of correlation nor the proximity of features (Tables S5, S6). For cell

group 3, the selected DNA methylation features have low methylation in many cells, whereas the

DNA accessibility features tend to have high accessibility (Fig. 2B, G). This was in contrast to other

cell groups, suggesting that the transcriptional response to HMA treatment is associated with

reductions in DNA methylation and gains in accessibility. Consistently, transcript features from

expression cluster 1 had predominantly positive correlations with accessibility features and many

negative correlations with methylation features, especially in CpG islands and 3kb genomic windows

(Supplementary Fig. S4).

To test whether epigenetic alterations in *cis*-regulatory elements could influence transcriptional

responses to HMA treatment, we next correlated gene expression with DNA methylation and

accessibility in nearby genomic loci. Genes from expression cluster 1 showed a significant shift

toward negative correlations with promoter methylation (p = 2.15x10^{-11}, χ^2 test) and positive

correlations with promoter accessibility (p = 2.26x10^{-8}, χ^2 test), suggesting that these genes are

particularly sensitive to loss of DNA methylation in *cis* (Fig. 2H, I; Supplementary Table 7). An

interesting example is *RHEX* (regulator of hemoglobinization and erythroid cell expansion), which is

highly expressed in AML (32). Negative correlations between *RHEX* expression and DNA methylation

were concentrated in a promoter-proximal region of intron 1, which contains several conserved long

terminal repeat (LTR) TEs (Fig. 2I).

In summary, our single-cell multiomic analysis identified transcriptional changes linked to patterns of

epigenetic heterogeneity in HMA-treated AML cells. Importantly, activation of genes involved in

inflammatory response and cell death pathways (expression cluster 1) was observed in only a subset

of hypomethylated cells (cell group 3) (Fig. 2A, C, D).

Effects of HMA-induced heterogeneity on the self-renewal capacity of AML

cells

To determine the consequences of epigenetic and transcriptional heterogeneity on AML cell self-

renewal, colony-forming assays were performed. DAC and AZA significantly decreased colony counts

(Fig. 3A), as previously reported (33). DAC treatment also significantly increased the size of colonies

in the two *FLT3-ITD* cell lines, MOLM-13 and MV-4-11 (Supplementary Table 1, Fig. S5A). To

characterize the molecular profiles of colonies formed after HMA treatment, we performed single-

colony analysis of DNA methylation (Fig. 3B-C; Supplementary Table 8). Following DAC treatment

(Fig. 3B, solid fill) many colonies (34-65%) had DNA methylation levels comparable to untreated

colonies (≥75%). This was in stark contrast to the low percentage of methylation-retaining cells (1-

5%) observed in single-cell data after 72h of DAC treatment (Fig. 1A, dashed box; Fig. 3B, dashed

line). Surprisingly, this shift toward higher DNA methylation levels was far less pronounced in

colonies established following AZA treatment (Fig. 3C).

To assess the recovery of DNA methylation following HMA treatment, time-course analysis was

performed during colony formation (Fig. 3D). DNA methylation levels in HMA-treated HL-60 colonies

were low throughout the time course (Fig. 3D, top), mirroring an analysis performed in suspension

culture (Supplementary Fig. S1E). Similar results were obtained for MOLM-13 and MV-4-11 colonies

derived after AZA treatment (Fig. 3D, center and bottom). In contrast, high levels of DNA

methylation were observed in the early stages of colony formation (experiment day 6) following DAC

treatment of MOLM-13 and MV-4-11 cells (Fig. 3D, center and bottom). This suggests that the shift

toward high DNA methylation observed in these colonies (Fig. 3B) was not due to gradual recovery

of methylation. Rather, our data are consistent with the selection of highly methylated cells in

colony-forming assays performed after DAC treatment. We deduce that methylation-retaining cells

can have increased self-renewal and proliferative capacity relative to hypomethylated cells after

HMA treatment, but this may depend on the mutational profile of the AML cells and the HMA used.

Transcriptional changes in hypomethylated and methylation-retaining colonies

Next, we generated matched transcriptomes from the same set of colony samples (Supplementary

Table 8). Single-colony RNA-seq and principal component analysis (PCA) showed that DAC and AZA

samples were generally distinct from untreated samples, regardless of their global DNA methylation

levels, in all cell lines (Fig. 4A). This implies that HMA exposure has substantial effects on the

transcriptome, even in highly methylated cells.

Of the 2000 most variably expressed genes among the HL-60 samples, only 215 had increased

expression specific to hypomethylated colonies (Fig. 4B, cluster 3; Supplementary Table 9). Many of

these genes (42.3%) were upregulated after 72h of treatment with either DAC or AZA in bulk HL-60

RNA-seq data, and several were associated with activation of inflammatory responses within the

sPLS model (e.g., *S100A8* and *S100A9*; Supplementary Table 9; Fig. 2B, cluster 1). In contrast, the

genes in cluster 5 were upregulated following HMA exposure in both hypomethylated and highly

methylated colonies (Fig. 4B), and only 3.5% of these genes were induced by HMA treatment in bulk

RNA-seq data (Supplementary Table 9). Thus, some transcriptional changes that occur immediately

after HMA treatment are maintained in hypomethylated colonies, whereas other genes are

upregulated during colony formation and are independent of HMA-induced global hypomethylation.

GO ORA revealed an enrichment of anti-microbial and immune-related processes among both

hypomethylation-dependent and -independent gene sets, whereas 'small molecule biosynthetic

process' and 'cholesterol biosynthetic process' were over-represented among the hypomethylation-

independent cluster 5 genes (Fig. 4C; Supplementary Table 10). HL-60 colonies that retained DNA

methylation following DAC treatment also had particularly high expression of cholesterol-related

genes, including many enzymes required for *de novo* cholesterol biosynthesis downstream of

mevalonate (34) (Supplementary Fig. S6-7; Tables S11, S12).

We next performed a focused analysis of genes annotated to 'cholesterol biosynthetic process'

(GO:0006695) across all cell lines. Like HL-60 colonies, MOLM-13 and MV-4-11 samples displayed

increased expression of many of these genes after HMA treatment (Supplementary Fig. S8).

Furthermore, a statistical analysis revealed that 5 genes (*SREBF1*, *SCAP*, *PMVK*, *MVD* and *LSS*) were

significantly increased (Pairwise Wilcoxon test with Benjamini-Hochberg correction, FDR = < 0.05) by

both DAC and AZA in colonies derived from all 3 cell lines (Fig. 5A). These genes form part of a

regulatory feedback loop that maintains cholesterol homeostasis by up-regulating biosynthesis in

response to low cellular cholesterol (35) (Supplementary Fig. S6). Since cholesterol regulation is

altered in both methylation-retaining and hypomethylated colonies, targeting this metabolic

pathway could overcome the initial heterogeneity induced by HMA treatment (Fig. 1-3).

Inhibition of cholesterol pathways following HMA exposure

To further investigate the relationship between cholesterol regulation and self-renewal capacity

following HMA treatment, we performed colony-forming assays in the presence of rosuvastatin.

Rosuvastatin is a potent inhibitor of the rate-limiting enzyme of the cholesterol biosynthetic

pathway, HMG-CoA reductase (HMGCR), and is frequently prescribed to reduce the risk of

cardiovascular diseases associated with hypercholesterolaemia (36). AZA and rosuvastatin co-

treatment had varied effects in AML cell lines (Fig. 5B, C), with synergistic effects on colony-

formation observed in MV-4-11 cells (Supplementary Fig. S5C). Meanwhile, DAC and rosuvastatin

had synergistic effects on colony formation in all cell lines (Supplementary Fig. S5C), with

significantly fewer colonies formed upon addition of 10 or 30 μM rosuvastatin in MOLM-13 cells (Fig.

5B). Colonies also displayed significantly reduced size (Supplementary Fig. S5B) and DNA methylation

levels (Fig. 5C) following DAC and rosuvastatin treatment of MOLM-13 and MV-4-11 cell lines, where

strong selection for highly methylated cells was previously noted (Fig. 3B). This suggests that

inhibition of cholesterol biosynthesis is effective even against methylation-retaining cells that

maintain self-renewal capacity following HMA treatment.

The efficacy and potential clinical utility of DAC and rosuvastatin co-treatment was then tested in

immunocompromised mice engrafted with luciferase-tagged MOLM-13 cells. In a dose optimization

experiment, combining DAC and rosuvastatin led to a significant increase in median survival for all

doses tested (Fig. 6A; Supplementary Table 13). The 1 mg/kg/day dose of rosuvastatin was chosen

for further validation in this model, as it is a low dose commonly prescribed for preventing

cardiovascular disease (equivalent to approximately 5mg/day oral dose in humans). Co-treatment

with DAC and rosuvastatin once again led to a significant increase in median survival compared to

DAC treatment alone (24 vs. 21 days) in this aggressive *in vivo* AML model (Fig. 6B; Supplementary

Table 14). In contrast, rosuvastatin alone significantly reduced the median survival time of vehicle-

treated mice (18 vs. 19 days; Fig. 6B; Supplementary Table 14). Finally, we assessed AZA and DAC in

combination with rosuvastatin in a patient-derived xenograft model of AML (AML-16). Rosuvastatin

(1mg/kg/day) co-treatment significantly extended the median survival of AZA-treated mice by 8.5

days (Fig. 6C; Supplementary Table 15) and decreased leukemia burden (Fig. 6D). Meanwhile,

combining rosuvastatin with DAC (0.2mg/kg/day) had no significant effect on survival or leukemia

burden in this model (Supplementary Fig. S9). These *in vivo* results provide evidence that HMAs

combined with statins have the potential to decrease leukemia burden and improve AML survival,

with further research required to optimize drug selection and dosing in patient cohorts.

Discussion

A thorough understanding of the ways in which AML cells survive HMA therapy and initiate

recurrence is essential to overcome the clinical challenge of treatment-resistant relapse. Our unique

study has revealed HMA-induced heterogeneity and methylation-retaining cells that evade the

epigenetic effects of treatment. We also identify metabolic pathways associated with the self-

renewal of HMA-treated AML cells, which can be targeted to improve HMA effects.

Our single-cell multiomic analyses revealed global DNA methylation heterogeneity induced by

treatment and transcriptional responses linked to epigenetic changes. We observed activation of

inflammatory response and cell death pathways in only a minor subset of hypomethylated cells (Fig.

2B), consistent with our previous report of heterogeneous TE expression following DAC treatment

(24) and scRNA-seq data from a colorectal cancer cell line (37). These observations suggest that

additional epigenetic modifications, transcription factors, or other mechanisms can suppress

transcriptional responses in hypomethylated cells (e.g., H3K9me3 (38, 39)). Alternatively, loss of

methylation at specific loci may be required for HMAs to exert their effects. Our observations are

consistent with the lack of correlation between HMA-induced global hypomethylation and clinical

response (40, 41) and support the use of locus-specific methylation changes to build a predictor of

patient response (42).

In contrast to our single-cell analyses, Li *et al.* reported reduced epigenetic and transcriptional

variance in AML cells collected after 12 weeks of AZA treatment in a transgenic mouse model (43).

We suggest that HMAs initially increase epigenetic and transcriptional diversity, allowing some cells

to gain a relative growth or survival advantage. Subsequent expansion of these clones would lead to

reduced heterogeneity, as reported by Li *et al.* (43). Consistently, we found that some AML cells

retained high levels of DNA methylation during HMA treatment (Fig. 1A) and had a relative growth

advantage following drug withdrawal (Fig. 3). Methylation-retaining cells tended to divide less

frequently during treatment (Fig. 1B) but had higher self-renewal and proliferative capacity than

hypomethylated cells after DAC treatment (Fig. 3). Interestingly, this selection for methylation-

retaining cells was less pronounced following AZA treatment, reflecting previous reports of distinct

effects of these HMAs (44, 45). Since AZA is incorporated into both DNA and RNA (46), whereas DAC

is restricted to DNA, we speculate that RNA-mediated toxicities (such as translational inhibition (14,

15)) could prevent the growth of highly methylated cells following AZA treatment. Taken together,

our results show that heterogeneous responses to HMA treatment can influence AML cell self-

renewal with potential implications for relapse.

We observed upregulated expression of cholesterol biosynthesis genes in colonies formed after

HMA treatment (Fig. 5; Supplementary Fig. S6, 8). These changes were observed in both

hypomethylated and highly methylated colonies, suggesting that HMA exposure has persistent

transcriptional effects that are independent of global methylation loss. Rapidly proliferating AML

cells are known to have high cholesterol demands (47), with cholesterol and related metabolites

required for cell membrane synthesis, lipid rafts, cell cycle progression and activation of oncogenic

signaling molecules (via prenylation of RAS, Rho/Rac) (Supplementary Fig. S7) (34, 48). In colony-

forming assays rosuvastatin synergistically enhanced the effects of DAC (Fig. 5B; Supplementary Fig.

S5C), suggesting that rapid consumption of cholesterol supports AML cell self-renewal after HMA

treatment (35). Since this metabolic adaptation occurs even in highly methylated cells, statin co-

treatment could have broad impact to inhibit AML cell growth despite initial heterogeneity in HMA

responses.

Altered cholesterol regulation is an established feature of AML with relevance to prognosis and

therapy (47, 49, 50). Recently, cholesterol metabolism was also linked to DAC resistance in AML cell

lines, with statin co-treatment showing synergistic inhibition of *in vitro* AML cell growth (51). Others

have shown that inhibition of cholesterol biosynthesis can sensitize AML cells to radiation and

chemotherapy (52, 53). *In vivo*, we observed significantly improved survival of leukemia-bearing

mice treated with HMA and rosuvastatin (Fig. 6) suggesting that co-treatment may increase the

duration of response in some AML and MDS patients. Encouragingly, a recent retrospective analysis

of MDS patients (including some HMA-treated high-risk cases) reported improved survival and

reduced progression to AML in patients who commenced statin treatments three months before or

after MDS diagnosis (54). Current clinical trials are testing the safety of a statin (pitavastatin) in

combination with AZA and venetoclax in AML patients (55). This therapeutic avenue is of particular

interest since statins are commonly prescribed, well-tolerated, oral medications, which could be

rapidly repositioned for use in AML and MDS patients receiving HMA therapy. Our results support

further investigation of the long-term benefits of statin and HMA co-treatments in different AML

subtypes.

Methods

Cell lines and culture

AML cell lines (Supplementary Table 1), HL-60 (ATCC #CCL-240), MOLM-13 (DSMZ #ACC-554), and

MV-4-11 (ATCC #CRL-9591), were maintained in tissue culture flasks (Greiner Bio-One) at 37°C and

5% CO_2, and sub-cultured at 500,000 cells/mL every 2-3 days with fresh medium. HL-60 cells were

maintained in Iscove's Modified Dulbecco's medium (IMDM; Sigma-Aldrich) supplemented with

4mM GlutaMAX (Thermo Fisher Scientific) and 10% Fetal Bovine Serum (FBS; Sigma-Aldrich). MOLM-

13 and MV-4-11 cells were maintained in Roswell Park Memorial Institute 1640 medium (RPMI;

Sigma-Aldrich) supplemented with 2mM GlutaMAX and 10% FBS.

CellTrace staining

320   AML cells (2-2.5x10^6 cells/mL) were stained with 1μM (MOLM-13 and MV-4-11) or 3μM (HL-60)

321   CellTrace Far Red (Thermo Fisher Scientific), according to the manufacturer's instructions. MOLM-13

322   and MV-4-11 cells with uniformly high CellTrace fluorescence were purified by fluorescence-

323 activated cell sorting (FACS) prior to treatment with hypomethylating agents (HMAs), whereas all

324 cells were used for HL-60 treatments.

HMA treatments

326 HMA treatments were performed in suspension culture, with drugs added every 24h for 72h total.

327 All cell lines were treated with 100nM decitabine (DAC; Selleckchem #S1200), HL-60 cells were

328 treated with 2000nM azacytidine (AZA; Selleckchem # S1782), while MOLM-13 and MV-4-11 cells

329 were treated with 500nM AZA. Untreated cells (UNT) received 0.1% DMSO (vehicle control).

Fluorescence activated cell sorting (FACS)

331 HMA-treated cells were stained with propidium iodide (PI, 1.5µg/mL) and prepared for FACS. Viable

332 (PI⁻) single cells were sorted into 2.5µL of RLT PLUS buffer (Qiagen) containing 2.5U SUPERas-In

333 (Thermo Fisher Scientific) in 96-well plates using indexed sorting on a FACS Aria II (BD Biosciences).

334 Plates were sealed and briefly centrifuged before storage at -80°C for sequencing analysis.

Colony-forming assays

HMA-treated cells were seeded in MethoCult Optimum (H4034; STEMCELL Technologies Inc.) at 500

cells/well in 6-well plates with rosuvastatin (Selleckchem # S2169) added to the MethoCult at

various doses (0, 1, 10, 30µM). Cells were cultured at 37°C and 5% CO_2 for 14 days, and colonies

were imaged using Cytation3 (Biotek). Colony counts and sizes were analyzed using ImageJ software.

Individual colonies were manually plucked using a 20µL pipette tip into 100µL of media, centrifuged

at 200xg for 5 min, and then resuspended in 20µL of RLT PLUS buffer before storage at -80°C.

Alternatively, all colonies in each well were collected by resuspending the MethoCult Optimum

media (and colonies) in 3mL of standard culture media (IMDM or RPMI), centrifuging at 200xg for 5

min, and resuspending the cell pellet in 20-50µL of RLT PLUS buffer before storage at -80°C.

Library preparation and sequencing

scNMT-seq library preparation and sequencing

For scNMT-seq, matched scNOMe-seq and scRNA-seq libraries were prepared from sorted HL-60

single cells as previously described (23). Minor modifications to the published protocol were as

follows: 1) Amplified cDNA was purified using a 0.6:1 volumetric ratio of AMPure XP Beads (Beckman

Coulter) and eluted in 15µL water. 2) During the fifth repeat of first-strand synthesis reaction, the

samples were held at 37°C for 90min. 3) The oligo used in the second-strand synthesis reaction (5'-

CAGACGTGTGCTCTTCCGATCTNNNNNNN-3') was modified to be compatible with the NEBNext oligos

(multiplex oligos for Illumina, dual index sets, New England Biolabs), which were used to amplify

scNOMe-seq libraries. 4) A 0.65:1 volumetric ratio of AMPure XP Beads was used to purify products

of both the first- and second-strand synthesis reactions, as well as the amplified scNOMe-seq

libraries.

For scNOMe-seq libraries, paired-end 150bp sequencing was performed using the NovaSeq

(Illumina) platform. For scRNA-seq libraries, paired-end 75bp sequencing was performed on the

NovaSeq or NextSeq (Illumina) platform.

scTEM-seq library preparation and sequencing

For scTEM-seq analysis of global DNA methylation levels in single MOLM-13 and MV-4-11 cells,

library preparation was performed as described (24). Paired-end 150bp sequencing was performed

on the MiSeq (Illumina) platform.

Colony TEM-seq and RNA-seq library preparation and sequencing

Single-colony TEM-seq (Fig. 3B, C) and parallel RNA-seq analyses were performed as described (24,

25) with minor modifications. Lysates from single colonies (HL-60: 7.5µL; MOLM-13 and MV-4-11:

2.5µL) were used to separate genomic DNA and mRNA. During single-colony TEM-seq library

preparation, the number of SINE Alu amplification cycles was reduced to 29. For RNA-seq analysis,

15 cycles of cDNA amplification were performed.

TEM-seq analysis of pooled colonies (Fig. 3D, 5E) was performed as described (25) using 5-10µL of

cell lysate as input for bisulfite conversion and 29 cycles for SINE Alu amplification.

TEM-seq libraries were sequenced using 150bp paired-end sequencing on the MiSeq platform. For

RNA-seq libraries, paired-end 75bp sequencing was performed on the NextSeq or NovaSeq platform.

Data processing

Sequencing data were processed and aligned as described in the Supplementary Information.

scNMT-seq data analysis

Quality control, data normalization and batch correction were performed as described in the

Supplementary Information. All analyses were performed in R v4.2.1, unless otherwise stated.

Pairwise distance analysis of DNA methylation heterogeneity

To assess DNA methylation heterogeneity per treatment group and genomic context (Fig. 1D),

pairwise CpG methylation distance analysis was performed. The mean absolute methylation

difference was computed for each cell pair (A, B) as the mean of the absolute difference in

methylation rate at each common cytosine position in the relevant genomic annotation. To make

the comparison of methylation patterns meaningful, only cytosine loci with data from both cells in

the pair were used. The mean absolute methylation differences were grouped according to the

treatment combination of the cell pairs. The global summaries shown in Fig. 1D correspond to

groups in which both cells in the pair had the same treatment. Higher values indicate a more

heterogeneous methylation pattern when cells in the same treatment group were compared vis-à-

vis.

Cell-wise correlation analysis

To assess the relationships between DNA methylation, DNA accessibility, and gene expression within

individual cells (Fig. 1E), Pearson correlations were computed using HL-60 scNMT-seq data. For this

analysis, RNA-seq data was normalized and log transformed per batch using

'scuttle::logNormCounts()' (v1.6.2) (26) without batch correction or prior count filtering. DNA

methylation was correlated to DNA accessibility at matched loci, based on genomic co-ordinates. For

correlations with gene expression, methylation and accessibility measurements at promoters,

introns, and exons were matched to the corresponding transcript. For CpG islands, H3K27ac sites

and H3K4me3 sites, methylation and accessibility measurements were matched to all transcripts

within 10kb. For intergenic regions and 3kb genomic windows, methylation and accessibility

measurements were matched to every transcript within 1bp to assess the expression of immediately

adjacent genes. For each cell, Pearson's correlation estimates were then computed using all

matched values and the 'cor.test()' function in R.

Integrative sparse partial least squares (sPLS) analysis

Mixomics (27) (v6.20.0) was used to perform a multivariate integrative analysis of the HL-60 scNMT-

seq data (Fig. 2A-D). Feature selection was performed to identify variably expressed transcripts that

were highly correlated with changes in DNA Methylation and accessibility after HMA treatment. We

performed an unsupervised sparse Partial Least Squares (sPLS) analysis using the function

'mixOmics::mint.block.spls()' which combines a multivariate integrative (MINT) method and a

multiblock sPLS integrative analysis. MINT (28) accounts for multiple batches (Supplementary Table

2) measured on the same variables, while the multiblock sPLS seeks for correlated patterns between

DNA methylation and DNA accessibility rates that are split into multiple genomic regions ('blocks')

and explain (correlated to) the predictor (transcriptome).

To focus on transcriptomic and epigenetic changes resulting from HMA treatment, only treated cells

(AZA and DAC) were included in the sPLS model. The genomic regions included in this analysis for

both DNA methylation and DNA accessibility were CpG islands, promoters, H3K27ac sites, H3K4me3

sites and 3kb windows. The rates from these genomic regions were filtered to retain only those

detected in greater than 10% of cells. The sPLS model was implemented assessing two components,

selecting 50 features per component and per block (genomic region) in the DNA methylation and

DNA accessibility datasets, and 100 genes per component in the transcriptome dataset. Further

details are provided on GitHub.

Heatmap visualization and identification of cell and expression clusters from sPLS-selected features

was performed using the ComplexHeatmap package (v2.12.1). sPLS selected features for

components 1 and 2 were extracted using the function 'mixOmics::selectVar()'. Heatmap

visualization was performed on sPLS-selected features using transcriptomic (converted to z-score)

and epigenetic rates (mean of features in genomic regions and converted to z-score) that were

entered into the model and included both treated and untreated cells. K-means clustering was

performed on sPLS-selected transcriptomic features, first on Gene Expression (row_km=3), followed

by Cell Group (column_km=4).

sPLS sample projections (Fig. 2A, C, and D) were plotted using ggplot2 (29) (v3.3.6) by extracting the

sPLS components 1 and 2 for a given block (RNA or epigenetic genomic region) and overlaid with

relevant information, that is, the cell group identified from k-means clustering and treated cell type

or average DNA methylation.

Gene Ontology (GO) Over Representation Analysis (ORA) was calculated using clusterProfiler (30)

(v4.4.4) for 'biological process' and displayed using enrichplot (v1.16.1) (Fig. 2E, F). Gene Expression

k-means clusters (Fig. 2B) and sPLS selected features per component (1-2) were assessed by

'enrichGO(p.adj=0.05, p.adj.method = "fdr", q.val.threshold = 0.4)' and the list of genes from the

batch corrected transcriptome dataset (entered into the sPLS model) as the background. Results

were displayed as treeplots using default settings for pairwise 'termsim()' and 'treeplot(nCluster=5,

showCategory = 10)'.

The correlation of sPLS features (Supplementary Fig. S3) was calculated as a similarity matrix using

'mixOmics::circosPlot()' on the sPLS model. The results were displayed using ComplexHeatmap,

showing DNA methylation and DNA accessibility features related to transcript features split by the

previously identified Gene Expression k-means clusters.

Locus-specific correlation analysis

To compare gene expression to adjacent epigenetic features, locus-specific correlations were

performed using HL-60 scNMT-seq data (Fig. 2H, I). DNA methylation and DNA accessibility

measurements were paired to genes based on annotation (promoters) or by strand-aware position

within 10kbp of the gene transcription start site (CpG islands, H3K27ac sites, H3K4me3 sites and 3kb

windows). For paired sites, Pearson's correlations were computed between CpG or GpC methylation

rate and log gene expression values. All cells with data (i.e., the UNT, DAC, and AZA groups) were

combined in these correlations, and a minimum of 22 cells with paired data (i.e., both gene

expression and DNA methylation/accessibility measurements) were required for the correlation to

be performed.

Colony sequencing analysis

Quality control, data normalization and batch correction were performed as described in the

Supplementary Information. All analyses were performed in R v4.2.1.

Highly variable gene analysis

For Figure 4, highly variable genes (HVGs) were identified from colony RNA-seq data and principal

component analysis (PCA) was performed using 'scater::calculatePCA(ntop = 2,000)'. K-means

clusters of HVGs were determined using the R stats package (v4.2.1) with 'kmeans(centers = 8,

iter.max = kmeans.iter, nstart = 50)'. Heatmapping of HVGs and k-means cluster was performed

using 'ComplexHeatmap::pheatmap()' with z-scored values and Euclidean distance hierarchical

clustering within row clusters (k-means groups) and columns (samples) ordered by treatment and

descending average global methylation level.

GO ORA of k-means clusters was compared using clusterProfiler for 'biological process' by

'compareCluster(pAdjustMethod = "fdr", p.adj.threshold = 0.05, qvalueCutoff=0.4)' and the full list

of genes from the batch corrected dataset (for each cell type) as the background list. Plots were

created using 'clusterProfiler::dotplot(showCategory = 3) + coord_flip()'.

Analysis of cholesterol biosynthesis gene expression

For Figure 5A, genes from the cholesterol biosynthesis pathway (GO:0006695) were analyzed from

batch corrected expression data for each cell line (HL-60, MOLM-13 and MV-4-11). After batch

correction, 40 genes out of 58 genes (from GO:0006695) were in common across all three cell lines

datasets. In total, for each cell line there were 41 (HL-60), 41 (MOLM-13) and 43 (MV-4-11) genes

analyzed from GO:0006695. The R package rstatix (v0.7.2) was used to perform pairwise Wilcoxon

test between treatment groups with UNT samples as the reference group for each gene. Benjamini-

Hochberg false discovery rate (FDR) correction was performed on all p-values, per cell line. Boxplots

were generated using ggpubr (v0.6.0) displaying the batch corrected relative expression with

'pointrange' error.

AML xenograft models

Five-week-old female NOD.Cg-Prkdc scid Il2rg tm1Wjl /SzJ (NSG) mice were obtained from the

Australian Bioresources (ABR, Moss Vale, NSW, Australia) and acclimatized for one week prior to any

experimental procedure.

MOLM-13 cells transduced with firefly luciferase (MOLM-13-luc; 5×10^5 cells suspended in 100 μL of

PBS) were injected into the lateral tail vein of NSG mice. Leukemia burden was assessed by

bioluminescence imaging (BLI) twice a week using an IVIS Spectrum *in vivo* imaging system

(PerkinElmer, Waltham, MA, USA), following intraperitoneal injections of 150 mg/kg D-luciferin

(Promega, Alexandria, NSW, Australia) under anesthesia with isoflurane. Treatments commenced on

day 6 post-engraftment when a positive luminescence signal was detected. Mice (n = 5-15/group)

were treated by intraperitoneal injection of either vehicle (2% DMSO, 30% PEG300 in water), DAC

(0.2 mg/Kg in saline), rosuvastatin (1 mg/kg), or combination rosuvastatin (1 mg/kg, 10 mg/kg, or 40

mg/kg in 30% PEG300 in water) and DAC (0.2 mg/Kg) once a day (5 days on, 2 days off) for up to 3

weeks. The animals were monitored until they reached the ethical endpoint.

For patient derived xenograft (PDX) experiments, NSG mice were inoculated with 2×10^6 AML-16 cells

(*FLT3-ITD*, *NPM1*, *IDH2*, and *WT1* mutant) (31) in 100μL of PBS, by injection into the lateral tail vein.

Leukemia burden (% hCD45+ cells in peripheral blood) was assessed 1-2 times weekly via flow

cytometry (FACSCanto II). Treatment commenced once an average leukemia burden of 1% CD45$^+$

cells was achieved (~3.5 weeks). Mice (n= 6-11 per group) received vehicle (2% DMSO, 30% PEG300

in water), DAC (0.2 mg/Kg), AZA (1mg/Kg), rosuvastatin (1 mg/kg), DAC + rosuvastatin, or AZA +

rosuvastatin, via intraperitoneal injections once a day (5 days on, 2 days off) for 1 week (cycle 1),

followed by twice per week for an additional 2 weeks (2 cycles). Mice were monitored until the

ethical endpoint (>25% huCD45$^+$ cells).

Survival Analysis

Survival analyses were performed using Kaplan-Meier analysis followed by the log-rank (Mantel-Cox)

test. All statistical analyses were performed using GraphPad Prism v. 9.0 (GraphPad Software, La

Jolla, CA, USA). Differences were considered statistically significant at an adjusted *p*-value < 0.05.

Data Availability

507   Sequencing data generated in this study has been deposited in the GEO database (GSE255339).

508   Relevant code for this manuscript is available on GitHub via:

509   www.github.com/canepi/HMA_heterogeneity

References

1.   Stomper J, Rotondo JC, Greve G, Lübbert M. Hypomethylating agents (HMA) for the

treatment of acute myeloid leukemia and myelodysplastic syndromes: mechanisms of resistance and

novel HMA-based therapies. Leukemia. 2021.

2.   Jones PA, Taylor SM. Cellular differentiation, cytidine analogs and DNA methylation. Cell.

1980;20(1):85-93.

3.   Patel K, Dickson J, Din S, Macleod K, Jodrell D, Ramsahoye B. Targeting of 5-aza-2'-

deoxycytidine residues by chromatin-associated DNMT1 induces proteasomal degradation of the

free enzyme. Nucleic Acids Research. 2010;38(13):4313-24.

4.   Issa JP, Roboz G, Rizzieri D, Jabbour E, Stock W, O'Connell C, et al. Safety and tolerability of

guadecitabine (SGI-110) in patients with myelodysplastic syndrome and acute myeloid leukaemia: a

multicentre, randomised, dose-escalation phase 1 study. Lancet Oncol. 2015;16(9):1099-110.

5.   Fennell KA, Bell CC, Dawson MA. Epigenetic therapies in acute myeloid leukemia: where to

from here? Blood. 2019;134(22):1891-901.

6.   Daskalakis M, Nguyen TT, Nguyen C, Guldberg P, Kohler G, Wijermans P, et al.

Demethylation of a hypermethylated P15/INK4B gene in patients with myelodysplastic syndrome by

5-Aza-2'-deoxycytidine (decitabine) treatment. Blood. 2002;100(8):2957-64.

7.   Kiziltepe T, Hideshima T, Catley L, Raje N, Yasui H, Shiraishi N, et al. 5-Azacytidine, a DNA

methyltransferase inhibitor, induces ATR-mediated DNA double-strand break responses, apoptosis,

and synergistic cytotoxicity with doxorubicin and bortezomib against multiple myeloma cells.

Molecular Cancer Therapeutics. 2007;6(6):1718-27.

8.   Dunshee DR, Dai Y, Jang JC, Risueño A, Jeyaraju DV, Hagner P, et al. CC-486 Mechanism

Imparted By Extended Exposure of Azacitidine Upregulates Myeloid Differentiation Markers and

Induces Cell Death in AML Cells. Blood. 2020;136:33-4.

9.   Goodyear O, Agathanggelou A, Novitzky-Basso I, Siddique S, McSkeane T, Ryan G, et al.

Induction of a CD8+ T-cell response to the MAGE cancer testis antigen by combined treatment with

azacitidine and sodium valproate in patients with acute myeloid leukemia and myelodysplasia.

Blood. 2010;116(11):1908-18.

10.   Nahas MR, Stroopinsky D, Rosenblatt J, Cole L, Pyzer AR, Anastasiadou E, et al.

Hypomethylating agent alters the immune microenvironment in acute myeloid leukaemia (AML) and

enhances the immunogenicity of a dendritic cell/AML vaccine. Br J Haematol. 2019;185(4):679-90.

11.   Roulois D, Loo Yau H, Singhania R, Wang Y, Danesh A, Shen SY, et al. DNA-Demethylating

Agents Target Colorectal Cancer Cells by Inducing Viral Mimicry by Endogenous Transcripts. Cell.

2015;162(5):961-73.

12.   Liu M, Ohtani H, Zhou W, Orskov AD, Charlet J, Zhang YW, et al. Vitamin C increases viral

mimicry induced by 5-aza-2'-deoxycytidine. Proc Natl Acad Sci U S A. 2016;113(37):10238-44.

13.   Chiappinelli Katherine B, Strissel Pamela L, Desrichard A, Li H, Henke C, Akman B, et al.

Inhibiting DNA Methylation Causes an Interferon Response in Cancer via dsRNA Including

Endogenous Retroviruses. Cell. 2015;162(5):974-86.

14.   Aimiuwu J, Wang H, Chen P, Xie Z, Wang J, Liu S, et al. RNA-dependent inhibition of

ribonucleotide reductase is a major pathway for 5-azacytidine activity in acute myeloid leukemia.

Blood. 2012;119(22):5229-38.

15.   Diesch J, Le Pannérer M-M, Winkler R, Casquero R, Muhar M, van der Garde M, et al.

Inhibition of CBP synergizes with the RNA-dependent mechanisms of Azacitidine by limiting protein

synthesis. Nature Communications. 2021;12(1):6060.

16.   Fenaux P, Mufti GJ, Hellstrom-Lindberg E, Santini V, Finelli C, Giagounidis A, et al. Efficacy of

azacitidine compared with that of conventional care regimens in the treatment of higher-risk

myelodysplastic syndromes: a randomised, open-label, phase III study. Lancet Oncol.

2009;10(3):223-32.

17.   Fenaux P, Mufti GJ, Hellström-Lindberg E, Santini V, Gattermann N, Germing U, et al.

Azacitidine prolongs overall survival compared with conventional care regimens in elderly patients

with low bone marrow blast count acute myeloid leukemia. J Clin Oncol. 2010;28(4):562-9.

18.   Kantarjian HM, O'Brien S, Huang X, Garcia-Manero G, Ravandi F, Cortes J, et al. Survival

advantage with decitabine versus intensive chemotherapy in patients with higher risk

myelodysplastic syndrome: comparison with historical experience. Cancer. 2007;109(6):1133-7.

19.   Helbig G, Chromik K, Woźniczka K, Kopińska AJ, Boral K, Dworaczek M, et al. Real Life Data

on Efficacy and Safety of Azacitidine Therapy for Myelodysplastic Syndrome, Chronic

Myelomonocytic Leukemia and Acute Myeloid Leukemia. Pathology & Oncology Research.

2019;25(3):1175-80.

20.   Stahl M, DeVeaux M, Montesinos P, Itzykson R, Ritchie EK, Sekeres MA, et al.

Hypomethylating agents in relapsed and refractory AML: outcomes and their predictors in a large

international patient cohort. Blood Adv. 2018;2(8):923-32.

21.   DiNardo CD, Pratz K, Pullarkat V, Jonas BA, Arellano M, Becker PS, et al. Venetoclax

combined with decitabine or azacitidine in treatment-naive, elderly patients with acute myeloid

leukemia. Blood. 2019;133(1):7-17.

22.   DiNardo CD, Jonas BA, Pullarkat V, Thirman MJ, Garcia JS, Wei AH, et al. Azacitidine and

Venetoclax in Previously Untreated Acute Myeloid Leukemia. New England Journal of Medicine.

2020;383(7):617-29.

23.   Clark SJ, Argelaguet R, Kapourani CA, Stubbs TM, Lee HJ, Alda-Catalinas C, et al. scNMT-seq

enables joint profiling of chromatin accessibility DNA methylation and transcription in single cells.

Nature Communications. 2018;9(1):781.

24.   Hunt KV, Burnard SM, Roper EA, Bond DR, Dun MD, Verrills NM, et al. scTEM-seq: Single-cell

analysis of transposable element methylation to link global epigenetic heterogeneity with

transcriptional programs. Scientific Reports. 2022;12(1).

25.   Hunt KV BS, Bond DR, Lee HJ. Protocol for targeted analysis of transposable element

methylation levels and transcriptome in single cells using scTEM-seq. ProtocolExchange. 2022.

26.   McCarthy DJ, Campbell KR, Lun ATL, Wills QF. Scater: pre-processing, quality control,

normalization and visualization of single-cell RNA-seq data in R. Bioinformatics. 2017;33(8):1179-86.

27.   Rohart F, Gautier B, Singh A, Le Cao KA. mixOmics: An R package for 'omics feature selection

and multiple data integration. PLoS Comput Biol. 2017;13(11):e1005752.

28.   Rohart F, Eslami A, Matigian N, Bougeard S, Lê Cao K-A. MINT: a multivariate integrative

method to identify reproducible molecular signatures across independent experiments and

platforms. BMC Bioinformatics. 2017;18(1):128.

29.   Wickham H. ggplot2: Elegant Graphics for Data Analysis: Springer-Verlag New York; 2016.

30.   Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrichment tool

for interpreting omics data. The Innovation. 2021;2(3):100141.

31.   Lee EM, Yee D, Busfield SJ, McManus JF, Cummings N, Vairo G, et al. Efficacy of an Fc-

modified anti-CD123 antibody (CSL362) combined with chemotherapy in xenograft models of acute

myelogenous leukemia in immunodeficient mice. Haematologica. 2015;100(7):914-26.

32.   Deniz Ö, Ahmed M, Todd CD, Rio-Machin A, Dawson MA, Branco MR. Endogenous

retroviruses are a source of enhancers with oncogenic potential in acute myeloid leukaemia. Nature

Communications. 2020;11(1):3506.

33.   Tsai HC, Li H, Van Neste L, Cai Y, Robert C, Rassool FV, et al. Transient low doses of DNA-

demethylating agents exert durable antitumor effects on hematological and epithelial tumor cells.

Cancer Cell. 2012;21(3):430-46.

34.   Huang B, Song B-l, Xu C. Cholesterol metabolism in cancer: mechanisms and therapeutic

opportunities. Nature Metabolism. 2020;2(2):132-41.

35.   Duan Y, Gong K, Xu S, Zhang F, Meng X, Han J. Regulation of cholesterol homeostasis in

health and diseases: from mechanisms to targeted therapeutics. Signal Transduction and Targeted

Therapy. 2022;7(1):265.

36.   Teramoto T, Watkins C. Review of efficacy of rosuvastatin 5 mg. Int J Clin Pract.

2005;59(1):92-101.

37.   Takeshima H, Yoda Y, Wakabayashi M, Hattori N, Yamashita S, Ushijima T. Low-dose DNA

demethylating therapy induces reprogramming of diverse cancer-related pathways at the single-cell

level. Clinical Epigenetics. 2020;12(1):142.

38.   Griffin GK, Wu J, Iracheta-Vellve A, Patti JC, Hsu J, Davis T, et al. Epigenetic silencing by

SETDB1 suppresses tumour intrinsic immunogenicity. Nature. 2021.

39.   Irwin R, Scullion C, Thursby S, Sun M, Thakur A, Rothbart S, et al. UHRF1 suppresses viral

mimicry through both DNA methylation-dependent and -independent mechanisms. bioRxiv.

2020:2020.08.31.274894.

40.   Unnikrishnan A, Vo ANQ, Pickford R, Raftery MJ, Nunez AC, Verma A, et al. AZA-MS: a novel

multiparameter mass spectrometry method to determine the intracellular dynamics of azacitidine

therapy in vivo. Leukemia. 2018;32(4):900-10.

41.   Yang AS, Doshi KD, Choi SW, Mason JB, Mannari RK, Gharybian V, et al. DNA methylation

changes after 5-aza-2'-deoxycytidine therapy in patients with leukemia. Cancer Res.

2006;66(10):5495-503.

42.   Greve G, Andrieux G, Schlosser P, Blagitko-Dorfs N, Rehman U-U, Ma T, et al. In vivo kinetics

of early, non-random methylome and transcriptome changes induced by DNA-hypomethylating

treatment in primary AML blasts. Leukemia. 2023;37(5):1018-27.

43.   Li S, Chen X, Wang J, Meydan C, Glass JL, Shih AH, et al. Somatic Mutations Drive Specific, but

Reversible, Epigenetic Heterogeneity States in AML. Cancer Discovery. 2020;10(12):1934-49.

44.   Aumer T, Gremmelmaier CB, Runtsch LS, Pforr JC, Yeşiltaç GN, Kaiser S, et al. Comprehensive

comparison between azacytidine and decitabine treatment in an acute myeloid leukemia cell line.

Clinical Epigenetics. 2022;14(1):113.

45.   Flotho C, Claus R, Batz C, Schneider M, Sandrock I, Ihde S, et al. The DNA methyltransferase

inhibitors azacitidine, decitabine and zebularine exert differential effects on cancer gene expression

in acute myeloid leukemia cells. Leukemia. 2009;23(6):1019-28.

46.   Li LH, Olin EJ, Buskirk HH, Reineke LM. Cytotoxicity and mode of action of 5-azacytidine on

L1210 leukemia. Cancer Res. 1970;30(11):2760-9.

47.   Vitols S, Norgren S, Juliusson G, Tatidis L, Luthman H. Multilevel regulation of low-density

lipoprotein receptor and 3-hydroxy- 3-methylglutaryl coenzyme A reductase gene expression in

normal and leukemic cells. Blood. 1994;84(8):2689-98.

48.   Chiarella E, Nisticò C, Di Vito A, Morrone HL, Mesuraca M. Targeting of Mevalonate-

Isoprenoid Pathway in Acute Myeloid Leukemia Cells by Bisphosphonate Drugs. Biomedicines.

2022;10(5).

49.   Banker DE, Mayer SJ, Li HY, Willman CL, Appelbaum FR, Zager RA. Cholesterol synthesis and

import contribute to protective cholesterol increments in acute myeloid leukemia cells. Blood.

2004;104(6):1816-24.

50.   Bai S, Wang H, Shao R, Fu B, Lu S, Wang J, et al. Lipid profile as a novel prognostic predictor

for patients with acute myeloid leukemia. Frontiers in Oncology. 2023;13.

51.   Yabushita T, Chinen T, Nishiyama A, Asada S, Shimura R, Isobe T, et al. Mitotic perturbation

is a key mechanism of action of decitabine in myeloid tumor treatment. Cell Reports.

2023;42(9):113098.

52.   Dimitroulakos J, Nohynek D, Backway KL, Medley DW, Yeger H, Freedman MH, et al.

Increased Sensitivity of Acute Myeloid Leukemias to Lovastatin-Induced Apoptosis: A Potential

Therapeutic Approach. Blood. 1999;93(4):1308-18.

53.   Li HY, Appelbaum FR, Willman CL, Zager RA, Banker DE. Cholesterol-modulating agents kill

acute myeloid leukemia cells and sensitize them to therapeutics by blocking adaptive cholesterol

responses. Blood. 2003;101(9):3628-34.

54.   Afzal A, Fiala MA, Jacoby MA, Walter MJ. Statin use in myelodysplastic syndromes is

associated with a better survival and decreased progression to leukemia. Blood Advances.

2023;7(15):3838-41.

55.   Brem EA, O'Brien SM, Becker PS, Jeyakumar D, Pinter-Brown LC, Kirk C, et al. A Phase 1 Study

of Adding Pitavastatin to Venetoclax-Based Therapy in AML and CLL/SLL. Blood.

2023;142(Supplement 1):5940-.

Acknowledgements

Nicole Cole (University of Newcastle) provided technical support for fluorescence activated cell sorting. Al J. Abadi (Melbourne Integrative Genomics, University of Melbourne) assisted with the implementation of sPLS analysis. The authors thank Professor Tri Phan (Garvan Institute) and Professor Xu Dong Zhang (University of Newcastle) for critically reviewing this manuscript.

H.J.L., C.R. and M.D.D. received funding from the National Health and Medical Research Council of Australia (NHMRC) (GNT1143614, GNT1180782, GNT1173892, and GNT2016283). K.A.L.C. and H.J.L. received funding from the Australian Research Council (ARC) (DP200102903). H.J.L. and H.C.M. received funding from the Cancer Institute NSW (2018/ECF001 and ECF1299, respectively). D.R.B received funding from Cure Cancer Australia Foundation (CCAF2023-Bond) and NSW Health Pathology North. N.A.B. received funding from the McGuigan family through the Hunter Medical Research Institute (HMRI778). The contents of the published material are solely the responsibility of the research institutions involved or individual authors, and do not reflect the views of funding agencies.

Author Contributions

H.J.L. conceived and oversaw the project. D.R.B., A.K.E., C.R., K.A.L.C., and H.J.L. acquired funding. D.R.B., K.U., K.V.H., B.H., C.L.O., E.A.R., S.H., L.S.R., H.M., and H.J.L. performed experiments. D.R.B., S.M.B., K.U., K.V.H., B.H., C.L.O., L.S.R., C.R., and H.J.L. performed data analysis. S.M.B. and C.R. developed analytical approaches. S.M.B. managed sequencing data and analysis code. D.R.B., N.A.B., A.K.E, N.M.V., K.A.L.C., and H.J.L. supervised work. A.M., M.D.D. and C.E.d.B. provided critical resources. D.R.B., S.M.B., K.U., B.H., L.S.R., C.R., and H.J.L. prepared figures and wrote the manuscript. All authors have read and approved the manuscript.

Ethics Declarations

All experimental procedures using animals were reviewed, approved, and performed according to the Animal Care and Ethics Committee of the University of Newcastle (approval number: A-2023-303), and with consideration of the ARRIVE guidelines (Supplementary Information).

Figure Legends

Figure 1. HMA treatment induces DNA methylation heterogeneity in AML cells. HL-60, MOLM-13 and MV-4-11 cells were labelled with CellTrace and treated with decitabine (DAC; 100nM) or azacytidine (AZA; HL-60: 2000nM, MOLM-13 and MV-4-11: 500nM) every 24h for 72h. Single cells collected by indexed FACS were subjected to scNMT-seq (HL-60) or scTEM-seq (MOLM-13, MV-4-11). **A)** Violin plots of DNA methylation levels in single HL-60 (left), MOLM-13 (middle) and MV-4-11 (right) cells. Superimposed points show single-cell values from untreated (UNT, orange), DAC (cyan) and AZA (purple) groups. Dashed boxes surround DAC and AZA cells with methylation levels within the range of UNT samples. Data are shown for 185-222 cells collected from 2-3 replicate experiments in each cell line (UNT, $n = 27$-38; DAC $n = 63$-93; AZA $n = 68$-91). **B)** Scatter plots comparing CellTrace fluorescence and DNA methylation in single cells, with linear regressions, F-test p-values, and Pearson correlation coefficients (r). **C)** Violin plots of DNA methylation levels in different genomic contexts from HL-60 scNMT-seq data. **D)** Violin plots of DNA methylation heterogeneity, as determined by pairwise dissimilarity analysis, within different genomic contexts from HL-60 scNMT-seq data. **E)** Box and whisker plots of Pearson correlation coefficients computed within single cells from HL-60 scNMT-seq data. DNA methylation and DNA accessibility were considered in different genomic contexts, and individual loci were matched based on genomic co-ordinates. Correlations were performed between DNA methylation and DNA accessibility (top), DNA methylation and gene expression (middle), and DNA accessibility and gene expression (bottom). Boxes depict the interquartile range (IQR) with median. Whiskers extend to the highest and lowest data points within 1.5 x IQR of the first and third quartile. Outlying data points are not shown.

Statistical analysis was performed using ordinary one-way ANOVA with Dunnett's (A) or Šídák's (C-E)

multiple comparisons test: * $p < 0.05$, ** $p < 0.0005$ vs. UNT.

Figure 2. HMA-induced epigenetic heterogeneity influences transcriptional programs in AML cells.

HMA treated (AZA and DAC only) HL-60 scNMT-seq data underwent multivariate feature selection by

sparse least squares (sPLS) using an unsupervised model. **A)** sPLS projection of cells based on

transcript features coloured by cell group 1-4 (from B). **B)** Heatmap of transcript features selected by

sPLS displaying all samples (treated and untreated) as columns, are split by k-means clustering and

grouped by treatment. Individual gene and TE expression levels (rows) are z-score normalised and

split by k-means clustering with internal hierarchical clustering. **C-D)** sPLS projections of cells based

on transcript features are coloured by **C)** LINE:L2:L2a expression, and **D)** global methylation level. **E-**

F) Tree plots of the gene ontology (GO) over-representation analysis (ORA) for gene expression

clusters 3 and 1. **G)** Heatmaps summarising the DNA methylation and DNA accessibility features

selected by sPLS. Samples (columns) are ordered according to the heatmap in B. The average

methylation or accessibility of all sPLS selected features across all samples (treated and untreated) is

displayed on the left for each genomic context. The two heatmaps show z-score normalised

averages of DNA methylation and accessibility for all sPLS selected features in each genomic context.

H) Pearson correlations were computed between gene expression and DNA methylation (top) or

accessibility (bottom) of associated promoters. Bar graphs show the percentage of correlations ($p <$

0.05) with negative (left) and positive (right) coefficients for all genes and filtered by gene expression

cluster (identified in D). **I)** Correlations between *RHEX* expression and DNA methylation of adjacent

loci (top) are displayed together with annotated short and long interspersed nuclear elements (SINE,

LINE), long terminal repeat (LTR), DNA, and Simple repeat sequences (bottom). The blue shading

highlights a promoter-proximal region of intron 1 that included 6 regions with negative correlations

(cor < -0.3, $p < 0.05$) between DNA methylation and *RHEX* expression.

Figure 3. Highly methylated AML cells display a growth and survival advantage following

treatment with DAC, but not AZA. A) HL-60, MOLM-13, and MV-4-11 colony counts following

treatment with DAC (cyan) or AZA (purple) vs. untreated cells (UNT, orange). **B-C)** Density plots show

the average DNA methylation levels for single cells collected on experiment day 3 (dashed line) and

individual colonies collected on experiment day 17 (solid fill) following treatment with DAC or AZA.

HL-60 scNMT-seq data were filtered for cytosines within SINE Alu sites for direct comparison to

scTEM-seq data from colonies. Data are shown for 288 colonies collected from triplicate

experiments in each cell line ($n = 96$ per treatment). **D)** Time-course experiment showing changes in

average DNA methylation of cells collected at different time points throughout the colony-forming

assay (experiment days 6, 10 and 17). Values for experiment day 3 were obtained from single-cell

data (Fig. 1A). Data in A and D are expressed as mean +/- standard error of the mean (SEM).

Statistical analysis of colony counts (A) was performed using ordinary one-way ANOVA with

Dunnett's multiple comparisons test with a $p < 0.05$ cut-off for significance ($p < 0.03$*, $p < 0.006$**, p

< 0.0002***).

Figure 4. HMA treatment has sustained transcriptional effects in both methylation-retaining and

hypomethylated cells. A) Principal Component Analysis (PCA) plots of single-colony RNA-seq data

from AML cell lines, highlighting treatment groups (UNT = orange; DAC = cyan; AZA = purple) and

matched mean DNA methylation levels (circle: high > 75%; triangle: low < 75%). Data shown for 119 -

220 colonies collected from 3 replicate experiments in each cell line (UNT, $n = 14-73$; DAC, $n = 46-78$;

AZA, $n = 56-73$). **B)** Heatmap of the top 2000 highly variable genes from colony RNA-seq data.

Samples are ordered by decreasing global methylation levels (green gradient) within each treatment

group. Rows are grouped by K-means clusters based on gene expression, with hierarchical clustering

by Euclidean distance within each cluster. **C)** GO analysis of the clusters from the top 2000 highly

variable genes from B. The size of the circles indicates the gene ratio (number of genes from the

input list annotated to the GO term divided by the total number of genes in the input list), and the

color represents the significance of the adjusted *p*-value. Gene clusters are color-coded on the y-axis

and GO processes are shown on the x-axis.

Figure 5. Altered cholesterol regulation in cells surviving HMA therapy can be targeted to enhance

treatment efficacy. A) Box and whisker plots of relative expression for *SREBF1*, *PMVK*, *MVD* and *LSS*

in HL-60 (left), MOLM-13 (middle) and MV-4-11 (right) colony data from Figure 5. Pairwise Wilcoxon

test between treatment groups was performed with UNT samples as the reference group, for genes

present within 'cholesterol biosynthetic process' (GO:0006695). Boxes depict the interquartile range

(IQR) with median. Whiskers extend to the highest and lowest data points within 1.5 x IQR of the

first and third quartile. Benjamini-Hochberg false discover rate (FDR) correction was performed on

all p-values, per cell line: FDR < 0.05*, FDR < 0.01**, FDR < 0.001***. **B)** Colony counts for HL-60

(top), MOLM-13 (middle) and MV-4-11 (bottom) cell lines obtained following HMA and rosuvastatin

co-treatments. Data from No HMA (orange), DAC (cyan) and AZA (purple) groups are normalised to

the corresponding 0 µM rosuvastatin control. Means ± SEM for *n = 3* experiments. Significance

determined by two-way ordinary ANOVA with Dunnett's multiple comparisons test, *p* < 0.05* vs.

corresponding 0 µM rosuvastatin control. **C)** DNA methylation of colonies formed following DAC and

rosuvastatin co-treatments. Means ± SEM for *n = 3* experiments. Significance determined by one-

way ANOVA with Tukey's multiple comparisons test, *p* < 0.05* vs. corresponding 0 µM rosuvastatin

control.

Figure 6. HMA and statin co-treatment increases survival *in vivo*. A) Rosuvastatin dose optimisation

experiment showing median survival of mice engrafted with MOLM-13-luc cells following treatment

with DAC (0.2mg/kg/day) +/- rosuvastatin (1, 10, 40mg/kg/day) on a treatment schedule of '5 days

on, 2 days off' for 3 cycles. **B)** Validation of survival benefit when DAC (0.2mg/kg/day) is combined

with rosuvastatin (1mg/kg/day) in mice engrafted with MOLM-13-luc AML cells. **C)** AZA and

rosuvastatin co-treatment significantly extended median survival of NSG mice engrafted with a

patient-derived xenograft (AML-16 model). Mice were treated with AZA (1mg/kg/day) +/-

rosuvastatin (1mg/kg/day) for 5 days on, 2 days off in cycle 1, followed by twice per week for an

additional 2 cycles. **D)** Leukemia burden (% human CD45$^+$ cells) post-engraftment and following the

cessation of treatment. Mann-Whitney test (unpaired, non-parametric, two-tailed t-test) was used

to test for statistical significance. Survival analyses (A-C) were performed using Kaplan-Meier

analysis followed by the Log-rank (Mantel-Cox) test and an adjusted p-value of < 0.05 was

considered statistically significant. $p < 0.05$*, $p < 0.005$**, $p < 0.0001$***.

Figure 2

Figure 3

Figure 4

Figure 6

A)

	Median Survival (days)
Vehicle (n = 4)	18
DAC (n = 5)	20
DAC + 1 mg/kg ROS (n = 5)	26
DAC + 10 mg/kg ROS (n = 5)	26
DAC + 40 mg/kg ROS (n = 5)	22

B)

	Median Survival (days)
Vehicle (n = 10)	19
1 mg/kg ROS (n = 15)	18
DAC (n = 8)	21
DAC + 1 mg/kg ROS (n = 12)	24

C)

	Median Survival (days)
Vehicle (n = 11)	48
1 mg/kg ROS (n = 11)	48
AZA (n = 6)	69.5
AZA + 1 mg/kg ROS (n = 7)	78

D)