Orc6 dissociation from chromatin prevents premature loading of MCM at G2 and

tetraploid production

Yoko Hayashi-Takanaka[1,2,*], Ichiro Hiratani[2], Tokuko Haraguchi[1] and Yasushi Hiraoka[1*]

[1] Graduate School of Frontier Biosciences, Osaka University, 1-3 Yamadaoka, Suita

565-0871, Japan

[2] RIKEN Center for Biosystems Dynamics Research, 2-2-3 Minatojima-minamimachi,

Chuo-ku, Kobe 650-0047 Japan

[*]Correspondence should be addressed to:

Yoko Hayashi-Takanaka

RIKEN Center for Biosystems Dynamics Research, 2-2-3 Minatojima-minamimachi,

Chuo-ku, Kobe 650-0047 Japan

E-mail: yoko.hayashi.yr@riken.jp

or

Yasushi Hiraoka

Graduate School of Frontier Biosciences, Osaka University, 1-3, Yamadaoka, Suita,

565-0871, Japan

E-mail: hiraoka@fbs.osaka-u.ac.jp

Abstract

DNA replication is tightly regulated to occur only once per cell cycle. The untimely

initiation of DNA replication can result in genome instability, leading to aneuploidy,

which has been associated with early senescence and cancer. The pre-replication

complex, comprising the origin recognition complex (ORC; Orc1-6), Cdc6, Cdt1, and

MCM, is required for initiating DNA replication, although the function of Orc6 is yet to

be elucidated. Here, we show that Orc6 dissociates from chromatin upon entering the

S-phase and that the Orc6 dissociation depends on proteasome activity. Treatment that

inhibits proteasome activity, which declines with aging, increases the senescence marker

p21 levels, and promotes cell cycle arrest in human immortalized hTERT-RPE1 cells.

This treatment induced large nuclei with high levels of chromatin-bound Orc6 and

MCM without undergoing mitosis. When the proteasome activity recovered, those cells

with high levels of chromatin-bound Orc6 and MCM proceeded to whole-genome DNA

replication, confirming that they were tetraploid G1 cells. We propose that

proteasome-dependent dissociation of Orc6 from chromatin after S-phase is essential for

preventing MCM reloading and the subsequent development of tetraploid cells.

Keywords

Aging/DNA replication/ORC/ pre-replication complex(pre-RC)/ tetraploid cells

Genomic DNA replication is essential for the proliferation of living organisms. In eukaryotes, this process is highly regulated to ensure that all chromosomes replicate once per cell cycle. An impairment of this process results in abnormal DNA replication, which can lead to genomic instability and promote senescence or oncogenesis (Thakur *et al*, 2022).

DNA replication begins with binding of pre-replication complexes (pre-RCs) to replication origins in the G1 phase (Hu & Stillman, 2023). The pre-RC comprises four types of proteins: the origin recognition complex (ORC), minichromosome maintenance (MCM), Cdc6, and Cdt1. MCM is a replicative helicase consisting of Mcm2–7 subunits (Bell & Botchan, 2013; Deegan & Diffley, 2016; Ishimi, 2018). Cdc6 and Cdt1 assist the recruitment of the MCM complex to chromatin. Upon entering the S-phase, the pre-RC disassembles. Abnormal pre-RC loading onto chromatin after the S-phase can initiate DNA re-replication before mitosis (Vassilev & DePamphilis, 2017; Limas & Cook, 2019). Therefore, after the onset of S-phase, Cdt1 and Cdc6 protein levels are highly regulated through proteasomal degradation mediated by E3 ubiquitin ligases (Nishitani *et al*, 2006; Walter *et al*, 2016).

ORC consists of the Orc1–6 hexamer and serves as a scaffold for assembling the other pre-RC components at the DNA replication origins during the G1 phase (Bell & Labib, 2016). In metazoans, replication origins do not share a distinct consensus sequence; instead, they are defined by their association with ORC binding. Metazoan ORC exists as a complete hexamer only during the G1 phase. Orc2–5 bind to chromatin throughout the cell cycle, whereas Orc1 selectively dissociates from chromatin via ubiquitin-mediated proteolysis after entering the S-phase (Li & DePamphilis, 2002; Méndez *et al*, 2002; Tatsumi *et al*, 2003). Orc1–5 exhibit structural similarities, whereas

Orc6 shows no structural similarity to other ORC subunits (Duncker *et al*, 2009),

implying functional dissimilarity. Orc6 has the potential to independently bind to DNA

regardless of Orc1-5 (Xu *et al*, 2020), participate in DNA repair (Lin *et al*, 2022), and

contribute to mitosis (Bernal & Venkitaraman, 2011; Semple *et al*, 2006; Balasov *et al*,

2009) in various living organisms. However, the function of human Orc6 during the cell

cycle remains largely unknown.

Proteasomes serve as the major cellular proteolytic machinery. Low levels of

proteasome activity are observed in replicative senescent cells (Sabath *et al*, 2020), and

its inhibition induces cellular senescence in normal human fibroblasts (Chondrogianni

et al, 2003; Chondrogianni & Gonos, 2004; Ukekawa *et al*, 2004; Torres *et al*, 2006).

Senescent cells are produced by evading mitosis and exhibit chromatin-bound pre-RCs,

indicating their presence in the tetraploid G1 phase (Panopoulos *et al*, 2014; Johmura *et*

al, 2014; Zeng *et al*, 2023). However, the role of proteasome activity in pre-RC

formation during cellular senescence remains elusive.

In this study, we investigated changes in Orc6 levels during the cell cycle in

human cells using multicolor immunofluorescence-based single-cell plot analysis

(hereafter referred to as "single-cell plot analysis"). Single-cell plot analysis enables

quantitative measurements of Orc6 levels in individual cells using fluorescence

microscopy images (Hayashi-Takanaka *et al*, 2020, 2021). Therefore, it provides the

advantage of not requiring cell synchronization when examining target protein levels

during the cell cycle. In this study, we found that the continued presence of Orc6 on

chromatin after the S phase entry resulted in the formation of tetraploid cells.

Results

Dissociation of Orc6 from chromatin during the S phase

To quantify chromatin-bound Orc6 in individual human cells, we performed single-cell

plot analysis using fluorescence microscopy to measure multiple intracellular

components within each cell (Hayashi-Takanaka *et al*, 2020, 2021). An asynchronous

population of hTERT-RPE1cells was fixed using two different methods ("direct fixation"

or "pre-extraction," Fig. 1A). The "direct fixation" method detects whole nuclear

proteins, whereas the "pre-extraction" method only detects chromatin-bound fractions

(Hayashi-Takanaka *et al*, 2021). The fixed cells were stained with Hoechst 33342 (DNA

amount indicator, referred to as "Hoechst"), EdU (S-phase indicator), and an anti-Orc6

antibody (Fig. EV1). The same cells were stained with an anti-Orc2 antibody to detect

the behavior of Orc2–5 subunits on chromatin throughout the cell cycle (Vashee *et al*,

2001; Dhar *et al*, 2001). The fluorescence images indicated that Orc6 staining was

relatively uniform in the cell population prepared by the direct fixation method, whereas

it was variable among the cells prepared by the pre-extraction method (compare the

upper and lower panels for Orc6; Fig. 1A). In contrast, Orc2 staining was relatively

uniform in both methods (right panels, Fig. 1A). Single-cell plot analysis was performed

based on the images shown in Fig. 1A, and the normalized intensities of EdU, Orc6, and

Orc2 were plotted against the DNA content determined by the Hoechst intensity (each

dot represents one cell, Fig. 1B). EdU-positive S-phase cells (>-2 on the $\log_2$ scale) are

denoted as orange dots in the EdU panels (left panels, Fig. 1B), and the same cells are

denoted as orange dots in both Orc6 (middle panels, Fig. 1B) and Orc2 panels (right

panels, Fig. 1B). In the direct fixation method, the whole fraction of Orc6 levels were

relatively constant, ranging from -1 to 1 on the $\log_2$ scale throughout the cell cycle

(upper middle panel, Fig. 1B). In contrast, when employing the pre-extraction method,

Orc6 levels in the chromatin-bound fraction were high in the G1 phase, decreased

immediately after S-phase entry, and remained low in the S and G2 phases (lower

middle panel, Fig. 1B). The Orc2 levels remained relatively constant at the G1/S

transition and changed little during the S phase in both direct and pre-extraction

methods (right panels, Fig. 1B). These results indicate that Orc6, but not Orc2,

dissociates from chromatin during the S phase in hTERT-RPE1 cells.

To understand whether such Orc6 dissociation was also detected in other cell

lines, we examined changes in chromatin-bound Orc6 levels during the cell cycle in cell

lines derived from cancer cells (HeLa and U2OS cells) and normal fibroblast cells

(IMR90 cells) using the pre-extraction method (Fig. EV2A, B). The decrease in Orc6

levels at the transition from G1 to S phase was smaller in cancer cell lines (HeLa and

U2OS cells, Fig. EV2B) than in normal cell lines (IMR90 and hTERT-RPE1 cells, Fig.

EV2B), suggesting that chromatin-bound Orc6 levels and their regulation during the

cell cycle may vary between normal and cancer cells or between cell types.

Involvement of proteasome for dissociation of Orc6 from chromatin

As the levels of Cdt1, a component of the pre-RC, are regulated by proteasome activity,

we investigated whether Orc6 was also a target for proteasome activity using MG132, a

proteasome inhibitor. The hTERT-RPE1 cells were treated with 20 μM MG132 for

varying durations (1, 2, and 4 h), fixed using the two different fixation methods, and

stained with Hoechst and anti-Orc6 antibody. The same cells were co-stained with

anti-Cdt1 and anti-Orc2 antibodies as controls for proteasomal degradation (Fig. 2A).

Single-cell plot analysis showed that Orc6 levels in the whole fraction (obtained by the

direct-fixation method) remained almost unchanged for up to 2 h of MG132 treatment

(blue dots, left panels in Fig. 2A), whereas those in the chromatin-bound fraction

(obtained by the pre-extraction method) increased with MG132 treatment over time

after the S phase (blue dots, right panels in Fig. 2A). Cdt1 levels (a positive control)

increased in cells progressing through the S and G2 phases after 1 h of treatment and

increased further with extended treatment periods in both fixation methods (orange dots,

Fig. 2A), indicating that proteasomal degradation occurred normally in cells under these

experimental conditions. Orc2 levels were almost unchanged for up to 4 h in the direct

fixation method (green dots, left panels in Fig. 2A), and they slightly increased for 4 h

treatment in the pre-extraction method (green dots, right panels in Fig. 2A). These

results suggest that Orc6, and to a lesser extent Orc2, was not degraded by the

proteasome during the S and G2 phases of the cell cycle but dissociated from chromatin

via proteasome activity.

To compare the increase in Orc6 levels during the S phase due to MG132

treatment with that in Cdt1 and Orc2 levels, S-phase cells were selected based on the

Hoechst value (1.1< Hoechst <1.75) from the single-cell plot analysis (Fig. 2A), and the

fluorescence intensity of Orc6 in each cell was plotted against the Hoechst value (each

dot represents a single cell, Fig. 2B). Plot analysis indicated that there were no or slight

increases in the Orc6 and Orc2 levels in the whole fraction for up to 4 h treatment,

whereas a large increase was observed in the total Cdt1 level with duration of MG132

treatment (whole fraction, Fig. 2B). Large increase was observed in the

chromatin-bound fraction of Orc6 in S-phase cells with duration of MG132 treatment

(Orc6, chromatin-bound fraction, Fig. 2B). The increase in chromatin-bound Cdt1 and

Orc2 levels was lower than that in Orc6 (chromatin-bound fraction, Fig. 2B). Orc6

chromatin binding was not affected by depletion of Orc1 and Orc2 (Fig. EV3),

suggesting that the increase in chromatin-bound Orc6 levels in MG132 treatment was

not attributable to the presence of other ORC components on chromatin.

To further confirm this result, we examined Orc6 protein levels in whole-cell

lysates and chromatin fractions obtained by sucrose cushion fractionation (Fig. 2C).

Cells were treated with 20 μM MG132 for varying durations (1, 2, and 4 h), and Orc6

levels were analyzed in whole cell lysates and chromatin fractions using western

blotting (Fig. 2D). The results indicated that the Orc6 levels in whole-cell lysates

remained unchanged throughout MG132 treatment (whole, top row, Fig. 2D). In

contrast, those in the chromatin fractions increased approximately 3-fold (top row, Fig.

2D). As a control, Cdt1 levels increased over time in the whole-cell lysates

(approximately 7-fold) and chromatin fractions (approximately 3-fold) after MG132

treatment. Orc2 levels remained almost unchanged but slightly increased after the

treatment (1.2–1.5-fold). These results are consistent with those of the single-plot

analysis (Fig. 2A), suggesting that Orc6 dissociates from chromatin during the S phase

via proteasome function without undergoing degradation.

Aged cells have high levels of chromatin-bound Orc6 after the S phase

To understand the relationship between Orc6 dissociation from chromatin and aging, we

examined the dynamics of chromatin-bound Orc6 in human diploid fibroblast cells of

different ages (Fig. 3A). IMR90 cells at passages 18 (younger cells) and 32 (older cells

containing senescent cells) were fixed using the pre-extraction method and stained with

Hoechst and anti-Orc6 antibody. The cells were co-stained with an anti-Mcm2 antibody

as a marker for the cell cycle phases. Levels of chromatin-bound Mcm2 increase during

the G1 phase, peak at the G1/S transition, decrease during the S phase, and become

lowest at the G2 phase (Mcm2, passage 18, Fig. 3B) (Zhou *et al*, 2020;

Hayashi-Takanaka *et al*, 2021; Zeng *et al*, 2023). Immunostaining of passage 32 cells

with an anti-Orc6 antibody occasionally showed enlarged nuclei with high levels of

chromatin-bound Orc6 (arrows in Fig. 3A), whereas passage 18 cells did not (Fig. 3A).

Single-cell plot analysis revealed the presence of cells with high Orc6 signals (Orc6

>0.5 on the $\log_2$ scale) and high DNA content (Hoechst >1.8) among passage 32 cells

(magenta dots in the lower left panel, Fig. 3B). In contrast, a limited number of cells

exhibiting these characteristics were observed among passage 18 cells (upper left panel

in Fig. 3B). The cells with high Orc6 levels (magenta dots in the Orc6 panel) are plotted

as magenta dots in the Mcm2 panel. Some of these magenta-marked cells (with high

DNA content >1.8) showed low Mcm2 signals (blue-boxed region, Fig. 3B), indicating

that these cells were in the G2 phase. These results suggest that Orc6 remained bound to

chromatin until the G2 phase in a subset of aged cells. Additionally, some of the

magenta-marked cells showed high Mcm2 signals (red-boxed region, Fig. 3B),

suggesting that these cells were in a non-G2-phase state.

Induction of senescence by MG132 treatment causes aberrant loading of

chromatin-bound Orc6 after the S phase

Senescent cells exhibit low proteasomal activity and impaired proteasome induces

cellular senescence (Chondrogianni *et al*, 2003; Sabath *et al*, 2020). We first confirmed

whether MG132 treatment induced cellular senescence in hTERT-RPE1 cells. We

examined levels of p21 (a marker of cellular senescence) in cells treated with 312 nM

MG132 for 18 h. Under these conditions, the incidence of cell death was lower than that

in treatment with higher concentrations (> 625 nM). Immunostaining using an anti-p21

antibody showed an increase in the number of the p21-positive cells by MG132

treatment compared with the DMSO control (Fig. 4A). Western blotting analysis also

showed an increase in p21 protein levels by MG132 treatment compared with the

DMSO control (Fig. 4B). These results suggest that inhibition of proteasome activity by

MG132 induces the state similar to cellular senescence.

Next, we induced cellular senescence in immortalized hTERT-RPE1 cells by

inhibiting proteasome activity with MG132 treatment and examined the Orc6 dynamics

during the cell cycle. hTERT-RPE1 cells were treated with relatively low MG132

concentrations (78, 156, and 312 nM) for 18 h. Cells were fixed using the pre-extraction

method, stained with anti-Orc6, anti-Mcm3, and anti-Cdt1 antibodies, and subjected to

single-cell plot analysis (Fig. 4C). MG132 treatment produced cells with high levels of

chromatin-bound Orc6 with high DNA content, which were denoted as magenta dots in

Orc6 panels (Orc6 >0.5 on the $\log_2$ scale, Hoechst >1.8, Fig. 4C). The number of

magenta cells increased with increasing MG132 concentrations. In contrast, few such

cells appeared in the control DMSO cells (Fig. 4C). The cells labeled with magenta dots

in Orc6 panels were also denoted as magenta dots in the Mcm3 and Ctd1 panels (middle

and lower panels, Fig. 4C). Some of the magenta-marked cells had low Mcm3 signals

(blue-boxed region, Fig. 4C; blue arrows, Fig. EV4A), indicating that these cells were

in the G2 phase. Notably, other magenta-marked cells had high Mcm3 levels (red-boxed

region, Fig. 4C; red arrows, Fig. EV4A), suggesting that these cells are in a

non-G2-phase state. Co-staining with Cdt1 showed that most of the magenta-marked

cells exhibited relatively high levels of Cdt1 compared with those of G2 phase cells in

the DMSO control. Based on these observations, we hypothesized that these red-boxed

cells with high Mcm3 levels represent G1-phase cells.

Defects in the proteasome cause the formation of tetraploid cells

To elucidate the emergence of high-Orc6 cells with high DNA content (magenta dots in

Fig. 4C) by MG132 treatment, we performed a time-course analysis of MG132

treatment (Fig. 4D) and analyzed cell cycle stages of the cells by staining with

anti-Mcm2 (G1-phase marker) and anti-histone H3 phosphorylated at the 10^{th} serine

residue (H3S10ph; M-phase marker) antibodies (Hayashi-Takanaka *et al*, 2009). When

cells were treated with DMSO as a control, cells with high histone H3S10ph signals

were detected (magenta dots, Fig. 4D; Fig. EV4B), indicating normal cell cycle

progression under these conditions (leftmost, lower panel, Fig. 4D). In contrast, upon

treatment with MG132, cells with high histone H3S10ph signals gradually disappeared

by 8 h. The number of cells with high DNA content (Hoechst ~2) and low Mcm2 levels

gradually increased for 4–12 h (blue-circled regions, Fig. 4D), suggesting that the first

4–12 h of treatment resulted in the accumulation of G2-phase cells. Longer MG132

treatments (16–24 h) showed an increase in the number of cells with high DNA content

(Hoechst ~2) and high Mcm2 levels (red-boxed regions, Fig. 4D). During the 16–24 h

period, no M-phase cells were observed. These results suggest that the cells with high

DNA content (Hoechst, ~2) and high Mcm2 levels represent tetraploid G1 cells

produced by skipping the M phase. We also confirmed that 24 h treatment with MG132

produced large nuclei with high levels of cyclin B1 (Fig. EV4C), which is expressed in

the late G2 and mitosis phases but less in the G1 phase (Behbehani *et al*, 2012). This

result suggests that the aberrant loading of MCM onto chromatin begins in the G2

phase.

To clarify the involvement of proteasomes in the formation of tetraploid G1

cells, we examined the depletion effects of PSMD7, a 26S proteasome subunit, required

for proteolytic activity (Shi *et al*, 2018), using siRNA treatment. Cells were subjected to

a 2-day treatment with siRNA targeting PSMD7. Western blotting analysis showed that

the amount of PSMD7 was reduced to approximately 20% of that of the control siRNA

(Fig. 4E). Under these conditions, siRNA-treated cells were fixed using the

pre-extraction method, stained with anti-Orc6 and anti-Mcm2 antibodies, and subjected

to single-cell plot analysis (Fig. 4F). The results showed that PSMD7 depletion

produced cells with high chromatin-bound Orc6 levels and high DNA content (magenta

dots, Orc6 >0.5 on the $\log_2$ scale, Hoechst >1.8, Fig. 4F). Some of these cells exhibited

low Mcm2 levels signifying the G2 phase (blue-boxed region, Fig. 4F), whereas others

exhibited high Mcm2 levels signifying the G1 phase (red-boxed region, Fig. 4F),

suggesting that PSMD7 plays a role in dissociation of Orc6 from chromatin at the onset

of S phase. These results also suggest that proteasome defects are responsible for the

abnormal loading of Orc6 onto chromatin after the S phase, leading to the production of

tetraploid G1 cells.

Tetraploid cells formed by proteasome inhibition progress to the S phase

We hypothesized that cells with abnormal MCM loading onto chromatin, with a

Hoechst value of approximately 2, would undergo DNA replication without mitosis. To

investigate this hypothesis, cells were treated with 312 nM MG132 for 24 h (day 1),

washed, and cultured in a fresh MG132-free medium for 24 h (day 2), and DNA

synthesis was monitored (Fig. 5A). To label cells undergoing DNA replication, EdU

was added to the culture medium 30 min prior to fixation at the end of day 1 or 2.

Subsequently, the cells were fixed using the pre-extraction method, stained with EdU,

anti-Mcm2, and anti-Orc6 antibodies, and subjected to single-cell plot analysis (Fig.

5B). EdU-positive cells were denoted as orange dots, and the same cells were also

denoted as orange dots in the Mcm2 and Orc6 panels (Fig. 5B). The results showed that

on day 1 of cell culture with MG132, the number of EdU-positive cells was limited,

with the majority of the cells being arrested at Hoechst levels of 1 or 2. Among the cells

with Hoechst levels of approximately 2, some cells had high levels of Mcm2 (red-boxed

regions, Fig. 5B). One day after MG132 removal (day 2), EdU-positive cells were

detected with Hoechst ranges of 1–2 and 2–4 (orange dots, day-2 panels, Fig. 5B).

These results indicate that cells with Hoechst levels of 2–4 were produced by an

additional round of DNA replication, implying the occurrence of whole-genome

duplication in these cells (EdU panels, day 2, Fig. 5B). Mcm2 patterns exhibited

similarities between cell populations with Hoechst levels 1–2 and 2–4 (Mcm2 panel,

day 2, Fig. 5B): Mcm2 levels increased during the G1 phase and decreased during the S

phase, reaching their lowest levels in the G2 phase. These results suggest that cells with

high Mcm2 levels and DNA content of Hoechst ~2 (red-boxed region, Fig. 5B)

proceeded to another round of DNA replication.

To understand the cell cycle phases of cells with high Orc6 levels and DNA

content of Hoechst ~2, these cells were denoted as magenta dots in the Orc6 panels

(Orc6 > 0.5 on the $\log_2$ scale, Fig. 5B'). The same cells were denoted as magenta dots

also in the Mcm2 panels (Fig. 5B'). As indicated in Fig. 4C, high Orc6 cell populations

with DNA content of Hoechst ~2 were divided into two groups on day 1: one group

exhibited high Mcm2 levels, whereas the other exhibited relatively low Mcm2 levels

(red- and blue-boxed regions, respectively, Fig. 5B'; red and blue arrows, respectively,

Fig.5C). In contrast, most of the high Orc6 cell population exhibited high Mcm2 levels

on day 2, indicating that these cells were in the G1 phase. These results suggest that

some of the high-Orc6 cells with DNA content of Hoechst ~2 are tetraploid G1 cells,

whereas others are diploid G2 cells. These results also suggest that tetraploid cells

produced by proteasome inhibitors can undergo DNA replication.

Involvement of Orc6 in MCM loading during tetraploid formation

To investigate whether the presence of Orc6 on chromatin after S phase entry

contributes to MCM loading, we performed proteasome inhibitor treatment following

siRNA treatment to deplete the Orc6 protein. Initially, cells were treated with Orc6

siRNA (siOrc6) for 42 h to assess the effect of Orc6 depletion alone on MCM loading

(Fig. 6A). Western blotting analysis confirmed that siRNA targeting Orc6 efficiently

reduced the total amount of Orc6 protein (Fig. 6B), and the chromatin-bound Orc6 (Fig.

EV3B). Following a 42-h siRNA treatment, the cells were fixed using the pre-extraction

method and stained with Hoechst and anti-Mcm2 antibody. Single-cell plot analysis

showed that the distribution pattern of siOrc6-treated cells resembled that of the

negative control (Fig. 6C, Fig. EV5A). As a positive control, we examined the effect of

Orc1 depletion, as Orc1 is a subunit essential for MCM loading on chromatin (Ogawa *et*

al, 1999; Samson *et al*, 2016; Coulombe *et al*, 2019) and is degraded in a

proteasome-dependent manner after the S-phase entry (Li & DePamphilis, 2002;

Méndez *et al*, 2002; Tatsumi *et al*, 2003). Upon treatment with Orc1 siRNA (siOrc1),

chromatin-bound Mcm2 levels were reduced compared to that in the negative control

(Fig. 6C), and the number of G1-phase cells increased while the number of S-phase

cells decreased (Fig. EV5A). These results confirmed the significance of Orc1 for

Mcm2 loading onto chromatin during the S phase and indicated that Orc1 depletion

moderately arrested the cell cycle at the G1 phase in hTERT-RPE1 cells. In contrast,

Orc6 depletion with siRNA had a weaker effect on Mcm2 loading onto chromatin when

compared to that in Orc1 (Fig. 6C and Fig. EV5A).

335   Next, we examined the effect of Orc6 depletion on the loading of Mcm2 onto

chromatin induced by proteasome-inhibitor treatment. Cells were treated with siRNA

for 18 h, followed by MG132 treatment for an additional 24 h under siRNA treatment

(Fig. 6D, Fig. EV5B). Following the treatment, cells were fixed using the pre-extraction

method, stained with Hoechst and anti-Mcm2 antibody, and then analyzed using

single-cell plot analysis. The results showed that cells with high Mcm2 levels and DNA

content of Hoechst ~2, that is, cells with MCM loading on chromatin after the S phase,

were observed in the negative control (siControl, Fig. 6E). These cells were recognized

as tetraploid G1 cells formed by skipping the M phase due to proteasome-inhibitor

treatment, as mentioned above (Fig. 4E). These tetraploid G1 cells with high MCM

levels (Hoechst >1.8 and Mcm2 >‒2 on the $\log_2$ scale) were selected (magenta dots, Fig.

6E) and quantified as a percentage of the total number of cells with Hoechst signals

>1.8 (Fig. 6F). The results showed that the percentage of tetraploid cells with high

MCM levels was $35.2 \pm 6.7\%$ in the siControl. In contrast, in Orc1-depleted cells, it was

significantly reduced to $5.7 \pm 2.4\%$ (Fig. 6F), indicating the significance of Orc1 in

MCM loading in tetraploid G1 cells. Similarly, the percentage of high-MCM cells was

reduced to $13.3 \pm 8.4\%$ in Orc6-depleted cells compared to that in the siControl ($p =$

0.0006, Fig. 6F). These results suggest that Orc6 is involved in the reloading of MCM

in tetraploid G1 cells, albeit to a lesser extent than Orc1.

We also examined the involvement of Cdc6, another pre-RC component, in

MCM loading following siRNA treatment. Western blotting analysis confirmed the

effective reduction of Cdc6 protein levels using siRNA (Fig. 6G). Cells were treated

with siRNA targeting Cdc6 for 18 h and then treated with MG132 for an additional 24 h

(Fig. 6D). The cells were fixed using the pre-extraction method and stained with

Hoechst and anti-Mcm2 antibody. Single-cell plot analysis showed a reduction in the

number of tetraploid G1 cells exhibiting high MCM levels (magenta dots, Hoechst >1.8,

and Mcm2 >−2 on the $\log_2$ scale) upon depletion of Cdc6 siRNA (Fig. 6H), similar to

Orc6 siRNA effects. Therefore, Orc6 and another pre-RC factor, Cdc6, are involved in

the chromatin loading of MCM during the tetraploid G1 phase.

In summary, during the G1 phase, Orc6 binds to chromatin in conjunction with

other pre-RC components. Upon entering the S phase, Cdt1 and Orc1 proteins are

degraded by proteasome activity, while Orc6 is dissociated from the chromatin (Fig.

7A). Orc2–5 remain on chromatin throughout the cell cycle. These cells progress to

mitosis, producing normal diploid cells (Fig. 7A). When cells lose proteasome activity,

Orc6, Orc1, Cdt1, and MCMs are prematurely reloaded onto chromatin without

progressing to mitosis, thereby producing tetraploid cells (Fig. 7B).

Discussion

In this study, we demonstrated the dynamic changes in chromatin-bound Orc6 levels

during the cell cycle in hTERT-RPE1 cells. The chromatin-bound Orc6 levels increased

until the G1/S transition, decreased upon entering the S phase, and remained low until

the next G1 phase, although the total amount of Orc6 was almost unchanged during the

cell cycle. Contradictory to our results, previous western blotting analysis using

synchronized HeLa cells in different cell cycle phases showed that the levels of

chromatin-bound Orc6 were nearly constant throughout the cell cycle (Méndez *et al*,

2002). Our single-cell plot analysis showed that the difference in the Orc6 levels

between the G1 and S/G2 phases in two cancer cell lines (HeLa and U2OS cells) was

smaller than that in normal cell lines (IMR90, and hTERT-RPE1 cells) (Fig. 1, Fig.

EV2). This suggests that regulating chromatin-bound Orc6 levels during the cell cycle

could potentially serve as a distinguishing factor between cancer and normal cells.

We demonstrated that Orc6 removal from chromatin after the S phase by a

proteasome-dependent mechanism plays a significant role in inhibiting abnormal MCM

loading. Aberrant MCM loading onto chromatin after the S phase can induce an

additional round of DNA replication without mitotic division, leading to tetraploidy and

consequent cancer development. To prevent this, the ubiquitin-proteasome system

regulates the protein levels of pre-RC factors such as Cdt1, Cdc6, and Orc1 during the

cell cycle (Nishitani *et al*, 2006; Walter *et al*, 2016; Li & DePamphilis, 2002; Méndez *et*

al, 2002; Tatsumi *et al*, 2003). Orc6 is not degraded at the protein level and is

dissociated from chromatin by proteasome activity. Therefore, proteasome activity is

presumed to differently regulate pre-RC protein levels on chromatin, thereby preventing

abnormal MCM loading. Given that Orc6 is the most diverse ORC subunit protein

(Dhar & Dutta, 2000), its dissociation from chromatin during the S phase may have

evolved as a new mechanism to prevent aberrant MCM loading onto chromatin.

The molecular mechanism underlying Orc6 dissociation from chromatin

remains uncertain. Orc6 has no known ubiquitinated amino acid residues for subsequent

proteasomal degradation (Coulombe *et al*, 2019). A possible explanation is the

involvement of post-translational modifications, such as phosphorylation, as observed in

S. cerevisiae Orc6 (Chen *et al*, 2007; Chen & Bell, 2011). Recently, the significance of

S-phase-specific Orc6 phosphorylation for genome stability through DNA damage

checkpoints has been reported (Lin *et al*, 2023). These protein modifications could be

responsible for Orc6 dissociation from chromatin, mediated by proteasome activity

through the degradation of modifying enzymes. Moreover, proteasomes may have

nonproteolytic functions. Multiple ATPases within the proteasome enable DNA

unwinding (Makino *et al*, 1996, 1999). Consequently, a proteasome lacking proteolytic

activity can facilitate DNA unwinding, allowing chromatin binding to ORC, including

Orc6. Furthermore, proteasome activity inhibition can induce nucleosome

hyperacetylation at transcription start sites (Kinyamu *et al*, 2020), which coincides with

replication origins (Sugimoto *et al*, 2018), potentially facilitating the chromatin binding

of Orc6.

Reduced proteasome activity impairs various cellular activities, including

senescence (Vilchez *et al*, 2014; Huang *et al*, 2022). Recent studies have shown that

cellular senescence corresponds to a tetraploid state in the G1 phase, which appears

when cells skip the M phase (Panopoulos *et al*, 2014; Johmura *et al*, 2014; Zeng *et al*,

2023). Consistently, we observed that inhibition of proteasome activity by MG132

resulted in tetraploid G1 cells, and that subsequent removal of the inhibitor restored

proteasome activity, promoting cell progression to the S phase (Figs. 4, 5). Therefore,

reduction in proteasome activity during aging may result in the production of tetraploid

G1 cells, thereby increasing the risk of cancer development.

Our results indicate the involvement of pre-RC factors Orc6, Orc1, and Cdc6

in tetraploid cell production. Although the deletion of either Orc1 or Orc2 has a

profound effect on cell cycle progression (Okano-Uchida *et al*, 2018; Chou *et al*, 2021),

Orc6 siRNA had a minimal effect on cell cycle progression in hTERT-RPE1 (Fig. 6,

this study) and U2OS cells (Lin *et al*, 2022). These results suggest that incomplete ORC

hexamers could recruit pre-RC to chromatin after the S phase in differentiated cells,

although all subunits of the Orc1–6 hexamer are presumably essential in

undifferentiated cells such as those in early embryogenesis (Prasanth *et al*, 2002; Ortega

et al, 2016). Therefore, it may be important to eliminate both Orc1 and Orc6 from the

chromatin at the onset of the S phase. Certain carcinomas with elevated Orc6 levels

(Pan *et al*, 2022; Tang *et al*, 2023) indicate that the presence of excessive amounts of

Orc6 might hinder Orc6 dissociation from chromatin, potentially resulting in the

reloading of pre-RC onto chromatin. Our findings indicate that normal cells can

progress to the tetraploid G1 phase when the proteasome activity is reduced, a

significant step associated with whole-genome duplication related to carcinogenesis

(Bielski *et al*, 2018; López *et al*, 2020; Gemble *et al*, 2022). Therefore, our findings

advance comprehension of the emerging replication factor dynamics through

proteasome activity and correlate with the process of tetraploid cell formation, which is

associated with senescence and cancer development.

Materials and Methods

Antibodies for immunostaining

The primary antibodies used were anti-Orc6 (3A4, sc-32735; Santa Cruz

Biotechnology), anti-Orc2 (M055-3, MBL), anti-Cdt1 (D10F11; Cell Signaling

Technology), anti-Mcm3 (sc-390480; Lot#J1118, Santa Cruz Biotechnology;

Anti-BM28; 610700, BD transduction), anti-Mcm2 (D7G11, Cell Signaling

Technology), anti-Orc1 (F-10, sc-398734, Santa Cruz Biotechnology), anti-GAPDH

(14C10; Cell Signaling Technology), anti-histone H3 (1G1) (Nozawa *et al*, 2010),

anti-PSMD7 (EPR13517, ab181072; Abcam), anti-Cdc6 (sc-9964, Santa Cruz

Biotechnology), anti-p21 (12D1; Cell Signaling Technology), anti-histone H3S10ph

(CMA311, (Hayashi-Takanaka *et al*, 2009)) and cyclin B1 (AB_397617, BD

biosciences). The specificity of these antibodies used for immunostaining is shown in

Fig. EV1. We previously confirmed by single-cell plot analysis based on

immunostaining that chromatin-bound Mcm2 and Mcm3 showed similar dynamics

during the cell cycle (Hayashi-Takanaka *et al*, 2021).

For fluorescent staining, the secondary antibodies were as follows: Alexa 488

donkey anti-mouse IgG (715-545-151; Jackson ImmunoResearch), Cy5 donkey

anti-mouse IgG (715-175-151; Jackson ImmunoResearch), Cy3 donkey anti-Rat IgG

(712-165-153; Jackson ImmunoResearch), Alexa 488 donkey anti-rabbit IgG

(711-545-152; Jackson ImmunoResearch) and Cy3 donkey anti-rabbit IgG

(711-005-152; Jackson ImmunoResearch).

Cell culture and chemicals

hTERT-RPE1 (ATCC, CRL-4000), IMR90 (ATCC, CCL-186), U2OS (ATCC, HTB-96),

and HeLa (RCB0007) cell lines were grown in a culture medium containing

high-glucose Dulbecco's modified Eagle's medium (Sigma-Aldrich) supplemented with

penicillin/streptomycin (100 units/mL penicillin, 100 μg/mL streptomycin; Fujifilm)

and 10% fetal calf serum (Thermo Fisher Scientific), as previously described

(Hayashi-Takanaka *et al*, 2021).

siRNA transfection was performed using Lipofectamine RNAiMAX (Thermo

Fisher Scientific) with siRNAs (SASI_Hs01_00069342 for Orc6,

SASI_Hs01_00038888 for Orc1, SASI_Hs02_00341067 for Orc2, and Universal

Negative Control #1, Sigma-Aldrich; s11403 for PSMD7, and s2745 for Cdc6, Ambion,

Thermo Fisher Scientific) in accordance with the manufacturer's protocol.

MG132 (Fujifilm) was dissolved in DMSO.

Fluorescence staining for single-cell plot analysis

Immunostaining was performed as previously described (Hayashi-Takanaka *et al*, 2011,

2021, 2015). Briefly, the cells ($\sim 2 \times 10^4$ cells/cm^2) were plated on a 12-well plate with a

coverslip (15 mm diameter; No. 1S; Matsunami) at the bottom and cultured for more

than 1 day.

In the direct fixation method, cells were first fixed with 1 mL of the fixative

(2% paraformaldehyde [Electron Microscopy Sciences] dissolved in 250 mM HEPES

[pH 7.4]) for 10 min at room temperature (approximately 25°C), followed by

permeabilization for 15 min using 1 % Triton X-100 in PBS, and subsequently blocked

with Blocking One (Nacalai Tesque, Inc.).

In the pre-extraction method, prior to fixation, cells were first treated with 1

mL of 0.2% Triton X-100 solution containing 20 mM HEPES (pH 7.4), 100 mM NaCl,

and 300 mM sucrose for 3 min on ice. After removing the Triton X-100 solution, the

cells were then fixed using the fixative described above. This fixation method removes

the nuclear membranes and loosely bound chromatin proteins.

The fixed cells were incubated with specific primary antibodies (0.2–1 μg/mL)

for 2 h at room temperature (approximately 25°C) and washed three times (10 min

each) using PBS. When cells were stained with multiple antibodies, the cells were

simultaneously incubated with these antibodies. The cells were treated with secondary

antibodies (1 μg/mL) and Hoechst (0.1 μg/mL) for 2 h at room temperature

(approximately 25°C) and subsequently washed three times with PBS. The cells were

mounted using the ProLong Glass Antifade Mountant (Thermo Fisher Scientific).

The replication foci were labeled by culturing the cells in a medium containing

10 μM 5-ethynyl-2′-deoxyuridine (EdU) for 30 min. Subsequently, the cells were fixed

using the direct fixation or pre-extraction method, and the resulting EdU signals were

detected with Alexa Fluor 647 using a Click-iT EdU Imaging Kit (Thermo Fisher

Scientific).

Microscopy

Fluorescence images were obtained using a DeltaVision Elite system (GE Healthcare

Inc.) equipped with a pco.edge 4.2 sCMOS camera (PCO) and an Olympus 40×

UApo/340 oil immersion objective lens (NA 0.65 with iris) with DeltaVision Elite Filter

Sets (DAPI-FITC-TRITC-Cy5).

Single-cell plot analysis

Single-cell plot analysis was used to measure the levels of multiple intracellular

components based on fluorescence microscopic images and plot their correlation in

individual cells (Hayashi-Takanaka *et al*, 2020, 2021). The fluorescence intensity of

each nucleus was measured using NIS Elements software (version 3.0; Nikon). The

background intensity outside the cells was subtracted. The nuclear areas of individual

cells were determined by automatic thresholding using Hoechst signals. The sum of the

intensities (i.e., the average intensity × nuclear area) in each nucleus was measured

across all the fluorescence channels. Cells in the M phase were excluded from analysis,

except for those shown in Fig. 4D. In Fig. 4D, M-phase cells were used for the analysis.

To plot the Hoechst intensity distribution of 450–600 nuclei, the signal intensity in each

nucleus was normalized relative to those with the 20th lowest and 20th highest

intensities, designated as 1 and 2, respectively, because nuclei with the lowest and

highest intensities were sometimes outliers. Hoechst intensity was plotted on a linear

scale. The average intensities of Orc6, EdU, Orc2, Cdt1, Mcm2, and Mcm3 were set to

1 (0 on the $\log_2$ scale) and plotted on the $\log_2$ scale. The fluorescence intensities were

normalized to time "0 h" (Figs. 2A, 2B), DMSO treatment (Figs. 4D, 4E, EV4C, EV4D),

the negative control siRNA (Figs. 6C, 6E, 6H, EV3A, EV4F, EV4D) and MG132

treatment for day 1 (Fig. 5). The values relative to the average were plotted using

Mathematica (version 13.0) (Wolfram Research).

Preparation of chromatin fractions and whole-cell lysates

Native chromatin fractions (hereafter, referred to as "chromatin fractions") were

prepared by the method previously described (Sheu & Stillman, 2006; Quan *et al*, 2015)

with some modifications (Hayashi-Takanaka *et al*, 2021). The cells were washed with

cold PBS, suspended in buffer A (50 mM HEPES [pH 7.4], 100 mM KCl, 2.5 mM

$MgCl_2$, 0.1 mM $ZnSO_4$, 1 mM ATP, 1 mM dithiothreitol [DTT], and protease inhibitor

cocktail [EDTA-free, Nacalai Tesque, Inc.]), and collected via centrifugation (1,300 *g*, 3

min, 4°C). The cells were subsequently treated with 200 μL of 0.2% Triton X-100 in

buffer A, layered with 200 μL of 30% sucrose cushion in buffer A containing 0.2%

Triton X-100, and centrifuged at $20,000 \times g$ for 10 min at 4°C. The pellets were

collected as chromatin fractions and dissolved in $1\times$ SDS loading buffer. The

suspension was then subjected to SDS-PAGE.

To prepare whole-cell lysates, the cells were washed with PBS and suspended

in an equal volume of $2\times$ SDS loading buffer. The suspension was subjected to

SDS-PAGE.

Western blotting analysis

Chromatin fractions and whole-cell lysates were suspended in $1\times$ SDS loading buffer

and subjected to 10%, 12% or 15% SDS-PAGE to separate proteins. Proteins were

transferred onto Immobilon-P PVDF membranes (Merck) using a semi-dry blotting

system (ATTO). After blocking with 5% skim milk (Nacalai Tesque, Inc.), the

membranes were incubated with the primary antibodies. After incubation with

peroxidase-conjugated secondary antibodies (GE Healthcare), the signals were detected

using ImmunoStar LD (Fujifilm Wako).

Acknowledgements

We thank Ms. Chizuru Ohtsuki and Yoshino Kubota for their technical assistance, and

Drs. Hisao Masai, Haruhiko Asakawa, and Yasuhiro Hirano for insightful discussion.

We also thank Dr. Hiroshi Kimura for histone H3 antibodies and Dr. Hisao Masukata for

critically reading the manuscript.

This work was supported by JSPS KAKENHI grants: JP18H05530,

JP20K20582 and JP20K08698 (to IH); JP18H05533, JP20H00454 and JP23K05636 (to

YH); and JP18H05528 (to TH). This work was also supported by the Naito Foundation,

Data Availability

571   Source data are available from the corresponding authors upon reasonable request.

Conflict of interest

574   The authors declare no competing interests.

576   **References**

577   Balasov M, Huijbregts RPH & Chesnokov I (2009) Functional analysis of an Orc6
578   mutant in Drosophila. *Proc Natl Acad Sci U S A* 106: 10672–10677

579   Behbehani GK, Bendall SC, Clutter MR, Fantl WJ & Nolan GP (2012) Single-cell mass
580   cytometry adapted to measurements of the cell cycle. *Cytometry A* 81: 552–566

581   Bell SD & Botchan MR (2013) The minichromosome maintenance replicative helicase.
582   *Cold Spring Harb Perspect Biol* 5: a012807

583   Bell SP & Labib K (2016) Chromosome Duplication in Saccharomyces cerevisiae.
584   *Genetics* 203: 1027–1067

585   Bernal JA & Venkitaraman AR (2011) A vertebrate N-end rule degron reveals that Orc6
586   is required in mitosis for daughter cell abscission. *J Cell Biol* 192: 969–978

587   Bielski CM, Zehir A, Penson AV, Donoghue MTA, Chatila W, Armenia J, Chang MT,
588   Schram AM, Jonsson P, Bandlamudi C, *et al* (2018) Genome doubling shapes the
589   evolution and prognosis of advanced cancers. *Nat Genet* 50: 1189–1195

590   Chen S & Bell SP (2011) CDK prevents Mcm2–7 helicase loading by inhibiting Cdt1
591   interaction with Orc6. *Genes Dev* 25: 363–372

592   Chen S, De Vries MA & Bell SP (2007) Orc6 is required for dynamic recruitment of Cdt1
593   during repeated Mcm2–7 loading. *Genes Dev* 21: 2897–2907

594   Chondrogianni N & Gonos ES (2004) Proteasome inhibition induces a senescence-like
595   phenotype in primary human fibroblasts cultures. *Biogerontology* 5: 55–61

596   Chondrogianni N, Stratford FLL, Trougakos IP, Friguet B, Rivett AJ & Gonos ES (2003)
597   Central role of the proteasome in senescence and survival of human fibroblasts:
598   induction of a senescence-like phenotype upon its inhibition and resistance to
599   stress upon its activation. *J Biol Chem* 278: 28026–28037

600   Chou H-C, Bhalla K, Demerdesh OE, Klingbeil O, Hanington K, Aganezov S, Andrews P,
601   Alsudani H, Chang K, Vakoc CR, *et al* (2021) The human origin recognition
602   complex is essential for pre-RC assembly, mitosis, and maintenance of nuclear
603   structure. *Elife* 10: e61797

Coulombe P, Nassar J, Peiffer I, Stanojcic S, Sterkers Y, Delamarre A, Bocquet S &
Méchali M (2019) The ORC ubiquitin ligase OBI1 promotes DNA replication
origin firing. *Nat Commun* 10: 2426

Deegan TD & Diffley JF (2016) MCM: one ring to rule them all. *Current Opinion in*
Structural Biology 37: 145–151

Dhar SK, Delmolino L & Dutta A (2001) Architecture of the human origin recognition
complex. *J Biol Chem* 276: 29067–29071

Dhar SK & Dutta A (2000) Identification and Characterization of the Human ORC6
Homolog. *Journal of Biological Chemistry* 275: 34983–34988

Duncker BP, Chesnokov IN & McConkey BJ (2009) The origin recognition complex
protein family. *Genome Biol* 10: 214

Gemble S, Wardenaar R, Keuper K, Srivastava N, Nano M, Macé A-S, Tijhuis AE,
Bernhard SV, Spierings DCJ, Simon A, *et al* (2022) Genetic instability from a
single S phase after whole-genome duplication. *Nature* 604: 146–151

Hayashi-Takanaka Y, Hayashi Y, Hirano Y, Miyawaki-Kuwakado A, Ohkawa Y, Obuse
C, Kimura H, Haraguchi T & Hiraoka Y (2021) Chromatin loading of MCM
hexamers is associated with di-/tri-methylation of histone H4K20 toward
S phase entry. *Nucleic Acids Res* 49: 12152–12166

Hayashi-Takanaka Y, Kina Y, Nakamura F, Becking LE, Nakao Y, Nagase T, Nozaki N
& Kimura H (2020) Histone modification dynamics as revealed by multicolor
immunofluorescence-based single-cell analysis. *J Cell Sci* 133: jcs243444

Hayashi-Takanaka Y, Maehara K, Harada A, Umehara T, Yokoyama S, Obuse C,
Ohkawa Y, Nozaki N & Kimura H (2015) Distribution of histone H4
modifications as revealed by a panel of specific monoclonal antibodies.
Chromosome Res 23: 753–766

Hayashi-Takanaka Y, Yamagata K, Nozaki N & Kimura H (2009) Visualizing histone
modifications in living cells: spatiotemporal dynamics of H3 phosphorylation
during interphase. *J Cell Biol* 187: 781–790

Hayashi-Takanaka Y, Yamagata K, Wakayama T, Stasevich TJ, Kainuma T, Tsurimoto
T, Tachibana M, Shinkai Y, Kurumizaka H, Nozaki N, *et al* (2011) Tracking

634 epigenetic histone modifications in single cells using Fab-based live endogenous
635 modification labeling. *Nucleic Acids Research* 39: 6475–6488

636 Hu Y & Stillman B (2023) Origins of DNA replication in eukaryotes. *Mol Cell* 83: 352–
637 372

638 Huang W, Hickson LJ, Eirin A, Kirkland JL & Lerman LO (2022) Cellular senescence:
639 the good, the bad and the unknown. *Nat Rev Nephrol* 18: 611–627

640 Ishimi Y (2018) Regulation of MCM2-7 function. *Genes Genet Syst* 93: 125–133

641 Johmura Y, Shimada M, Misaki T, Naiki-Ito A, Miyoshi H, Motoyama N, Ohtani N,
642 Hara E, Nakamura M, Morita A, *et al* (2014) Necessary and Sufficient Role for a
643 Mitosis Skip in Senescence Induction. *Molecular Cell* 55: 73–84

644 Kinyamu HK, Bennett BD, Bushel PR & Archer TK (2020) Proteasome inhibition
645 creates a chromatin landscape favorable to RNA Pol II processivity. *J Biol Chem*
646 295: 1271–1287

647 Li C-J & DePamphilis ML (2002) Mammalian Orc1 Protein Is Selectively Released from
648 Chromatin and Ubiquitinated during the S-to-M Transition in the Cell Division
649 Cycle. *Molecular and Cellular Biology* 22: 105–116

650 Limas JC & Cook JG (2019) Preparation for DNA replication: the key to a successful S
651 phase. *FEBS Lett* 593: 2853–2867

652 Lin Y-C, Liu D, Chakraborty A, Kadyrova LY, Song YJ, Hao Q, Mitra J, Hsu RYC, Arif
653 MK, Adusumilli S, *et al* (2022) Orc6 is a component of the replication fork and
654 enables efficient mismatch repair. *Proc Natl Acad Sci U S A* 119: e2121406119

655 Lin Y-C, Liu D, Chakraborty A, Macias V, Brister E, Sonalkar J, Shen L, Mitra J, Ha T,
656 Kajdacsy-Balla A, *et al* (2023) DNA Damage-Induced, S-Phase Specific
657 Phosphorylation of Orc6 is Critical for the Maintenance of Genome Stability. *Mol*
658 *Cell Biol* 43: 143–156

659 López S, Lim EL, Horswell S, Haase K, Huebner A, Dietzen M, Mourikis TP, Watkins
660 TBK, Rowan A, Dewhurst SM, *et al* (2020) Interplay between whole-genome
661 doubling and the accumulation of deleterious alterations in cancer evolution.
662 *Nat Genet* 52: 283–293

Makino Y, Yogosawa S, Kanemaki M, Yoshida T, Yamano K, Kishimoto T, Moncollin V,
Egly JM, Muramatsu M & Tamura T (1996) Structures of the rat proteasomal
ATPases: determination of highly conserved structural motifs and rules for their
spacing. *Biochem Biophys Res Commun* 220: 1049–1054

Makino Y, Yoshida T, Yogosawa S, Tanaka K, Muramatsu M & Tamura TA (1999)
Multiple mammalian proteasomal ATPases, but not proteasome itself, are
associated with TATA-binding protein and a novel transcriptional activator,
TIP120. *Genes Cells* 4: 529–539

Méndez J, Zou-Yang XH, Kim S-Y, Hidaka M, Tansey WP & Stillman B (2002) Human
Origin Recognition Complex Large Subunit Is Degraded by Ubiquitin-Mediated
Proteolysis after Initiation of DNA Replication. *Molecular Cell* 9: 481–491

Nishitani H, Sugimoto N, Roukos V, Nakanishi Y, Saijo M, Obuse C, Tsurimoto T,
Nakayama KI, Nakayama K, Fujita M, *et al* (2006) Two E3 ubiquitin ligases,
SCF-Skp2 and DDB1-Cul4, target human Cdt1 for proteolysis. *EMBO J* 25:
1126–1136

Nozawa R-S, Nagao K, Masuda H-T, Iwasaki O, Hirota T, Nozaki N, Kimura H & Obuse
C (2010) Human POGZ modulates dissociation of HP1alpha from mitotic
chromosome arms through Aurora B activation. *Nat Cell Biol* 12: 719–727

Ogawa Y, Takahashi T & Masukata H (1999) Association of fission yeast Orp1 and
Mcm6 proteins with chromosomal replication origins. *Mol Cell Biol* 19: 7228–
7236

Okano-Uchida T, Kent LN, Ouseph MM, McCarty B, Frank JJ, Kladney R, Cuitino MC,
Thompson JC, Coppola V, Asano M, *et al* (2018) Endoreduplication of the mouse
genome in the absence of ORC1. *Genes Dev* 32: 978–990

Ortega MA, Nguyen H & Ward WS (2016) ORC proteins in the mammalian zygote. *Cell*
Tissue Res 363: 195–200

Pan Q, Li F, Ding Y, Huang H & Guo J (2022) ORC6 acts as a biomarker and reflects
poor outcome in clear cell renal cell carcinoma. *J Cancer* 13: 2504–2514

Panopoulos A, Pacios-Bras C, Choi J, Yenjerla M, Sussman MA, Fotedar R & Margolis
RL (2014) Failure of cell cleavage induces senescence in tetraploid primary cells.

Mol Biol Cell 25: 3105–3118

Prasanth SG, Prasanth KV & Stillman B (2002) Orc6 Involved in DNA Replication,
Chromosome Segregation, and Cytokinesis. *Science* 297: 1026–1031

Quan Y, Xia Y, Liu L, Cui J, Li Z, Cao Q, Chen XS, Campbell JL & Lou H (2015)
Cell-Cycle-Regulated Interaction between Mcm10 and Double Hexameric
Mcm2-7 Is Required for Helicase Splitting and Activation during S Phase. *Cell*
Reports 13: 2576–2586

Sabath N, Levy-Adam F, Younis A, Rozales K, Meller A, Hadar S, Soueid-Baumgarten S
& Shalgi R (2020) Cellular proteostasis decline in human senescence. *Proc Natl*
Acad Sci U S A 117: 31902–31913

Samson RY, Abeyrathne PD & Bell SD (2016) Mechanism of Archaeal MCM Helicase
Recruitment to DNA Replication Origins. *Mol Cell* 61: 287–296

Semple JW, Da-Silva LF, Jervis EJ, Ah-Kee J, Al-Attar H, Kummer L, Heikkila JJ,
Pasero P & Duncker BP (2006) An essential role for Orc6 in DNA replication
through maintenance of pre-replicative complexes. *EMBO J* 25: 5150–5158

Sheu Y-J & Stillman B (2006) Cdc7-Dbf4 Phosphorylates MCM Proteins via a Docking
Site-Mediated Mechanism to Promote S Phase Progression. *Molecular Cell* 24:
101–113

Shi K, Zhang J-Z, Zhao R-L, Yang L & Guo D (2018) PSMD7 downregulation induces
apoptosis and suppresses tumorigenesis of esophageal squamous cell carcinoma
via the mTOR/p70S6K pathway. *FEBS Open Bio* 8: 533–543

Sugimoto N, Maehara K, Yoshida K, Ohkawa Y & Fujita M (2018) Genome-wide
analysis of the spatiotemporal regulation of firing and dormant replication
origins in human cells. *Nucleic Acids Res* 46: 6683–6696

Tang M, Chen J, Zeng T, Ye D-M, Li Y-K, Zou J & Zhang Y-P (2023) Systemic analysis of
the DNA replication regulator origin recognition complex in lung
adenocarcinomas identifies prognostic and expression significance. *Cancer Med*
12: 5035–5054

Tatsumi Y, Ohta S, Kimura H, Tsurimoto T & Obuse C (2003) The ORC1 Cycle in
Human Cells. *Journal of Biological Chemistry* 278: 41528–41534

Thakur BL, Ray A, Redon CE & Aladjem MI (2022) Preventing excess replication origin
activation to ensure genome stability. *Trends Genet* 38: 169–181

Torres C, Lewis L & Cristofalo VJ (2006) Proteasome inhibitors shorten replicative life
span and induce a senescent-like phenotype of human fibroblasts. *J Cell Physiol*
207: 845–853

Ukekawa R, Maegawa N, Mizutani E, Fujii M & Ayusawa D (2004) Proteasome
inhibitors induce changes in chromatin structure characteristic of senescent
human fibroblasts. *Biosci Biotechnol Biochem* 68: 2395–2397

Vashee S, Simancek P, Challberg MD & Kelly TJ (2001) Assembly of the human origin
recognition complex. *J Biol Chem* 276: 26666–26673

Vassilev A & DePamphilis M (2017) Links between DNA Replication, Stem Cells and
Cancer. *Genes* 8: 45

Vilchez D, Saez I & Dillin A (2014) The role of protein clearance mechanisms in
organismal ageing and age-related diseases. *Nat Commun* 5: 5659

Walter D, Hoffmann S, Komseli E-S, Rappsilber J, Gorgoulis V & Sørensen CS (2016)
SCFCyclin F-dependent degradation of CDC6 suppresses DNA re-replication.
Nat Commun 7: 10530

Xu N, You Y, Liu C, Balasov M, Lun LT, Geng Y, Fung CP, Miao H, Tian H, Choy TT, *et*
al (2020) Structural basis of DNA replication origin recognition by human Orc6
protein binding with DNA. *Nucleic Acids Research* 48: 11146–11161

Zeng J, Hills SA, Ozono E & Diffley JFX (2023) Cyclin E-induced replicative stress
drives p53-dependent whole-genome duplication. *Cell* 186: 528-542.e14

Zhou Y, Pozo PN, Oh S, Stone HM & Cook JG (2020) Distinct and sequential
re-replication barriers ensure precise genome duplication. *PLoS Genet* 16:
e1008988

748
749

 Figure Legends

Figure 1. Detection of chromatin-bound Orc6 by single-cell plot analysis

(**A**) Representative fluorescence images of Hoechst, EdU, Orc6, and Orc2 in

hTERT-RPE1 cells prepared using the direct fixation method (upper panels) and the

pre-extraction method (lower panels). Scale bars, 10 μm. (**B**) Single-cell plot analysis

based on the images in (**A**). Each dot represents the intensities of EdU, Orc6, and Orc2

in a single individual cell against the Hoechst intensities. The orange dots represent

S-phase cells based on EdU intensities ($\log_2$[EdU] >−2). n = 450 (number of cells

examined in each panel).

Figure 2. Effects of proteasome inhibitors on the dynamics of Orc6 during the cell

cycle

(**A**) Single-cell plot analysis of Orc6 (top), Cdt1 (middle), and Orc2 (bottom) in

hTERT-RPE1 cells. Cells were treated with 20 μM MG132 for 0, 1, 2, or 4 h (timescale

on the top), fixed using the direct fixation method (left panels) and the pre-extraction

method (right panels), and stained simultaneously with anti-Orc6, anti-Cdt1, and

anti-Orc2 antibodies. The cells were counterstained with Hoechst. Each dot represents

the intensity of Orc6, Cdt1, and Orc2 in a single individual cell plotted against the

Hoechst intensity. n = 450 (number of cells examined in each panel). (**B**) Orc6, Cdt1,

and Orc2 levels in the S-phase cells. Cells corresponding to the S phase (1.1< Hoechst

<1.75) are selected from (**A**), and their fluorescence intensities were plotted. The

average intensity in MG132-untreated cells was defined as 1. (**C**) Schematic diagram of

the biochemical fractionation method for whole-cell lysates and chromatin fractions.

(**D**) Western blotting analysis using antibodies against Orc6, Cdt1, Orc2, histone H3,

and GAPDH. hTERT-RPE1 cells were treated with 20 μM MG132 for the indicated

time periods (h). Whole-cell lysates (left) and chromatin fractions (right) were prepared

as described above and analyzed using western blotting. Histone H3 and GAPDH are

loading controls. The positions of molecular mass markers (kDa) are indicated to the

right.

Figure 3. Aberrantly high levels of chromatin-bound Orc6 and Mcm2 in aged

fibroblast cells

(**A**) IMR90 cells (passage numbers 18 and 32) were fixed using the pre-extraction

method and stained using Hoechst, anti-Orc6, and anti-Mcm2 antibodies.

Representative fluorescence images are shown. Arrows indicate the relatively larger
nucleus with high Orc6 signals. Scale bars, 10 μm. (**B**) Single-cell plot analysis based
on the images in (**A**). Each dot represents the intensities of Orc6 and Mcm2 in an
individual cell plotted against the Hoechst intensities. The magenta dots in the Orc6
panels represent cells with fluorescent signals of Hoechst >1.8 and $\log_2$(Orc6) >0.5. The
same cells were marked by magenta in the Mcm2 panels. The red-boxed regions
represent cells with high MCM levels ($\log_2$[Mcm2] >-1) and high DNA content
(Hoechst >1.8), and the blue-boxed regions represent cells with low MCM levels
($\log_2$[Mcm2] <-1) and high DNA content (Hoechst >1.8). n = 450 (number of cells
examined in each panel).
795
Figure 4. Inhibition of proteasome activity causes cellular senescence and
aberrantly high levels of chromatin-bound Orc6 and MCM
(**A**) Representative fluorescence images of Hoechst and p21 staining in hTERT-RPE1
cells treated with DMSO or 312 nM MG132 for 18 h. Cells were fixed using the direct
fixation method. Scale bars, 10 μm. (**B**) Western blotting analysis of whole-cell lysates
using antibodies against p21 and GAPDH. hTERT-RPE1 cells were treated with DMSO
as a control or 312 nM MG132 for 1 day before harvesting. The positions of molecular
mass markers (kDa) are indicated to the right. GAPDH is a loading control. (**C**)
Single-cell plot analysis of Orc6 (top), Mcm3 (middle), and Cdt1 (bottom) in
hTERT-RPE1 cells. Cells were treated with DMSO as a control or the indicated
concentrations of MG132 for 18 h, fixed using the pre-extraction method, and stained
with Hoechst, anti-Orc6, anti-Mcm3, and anti-Cdt1 antibodies. The cells showing high
Orc6 levels ($\log_2$[Orc6] >0.5) and high DNA content (Hoechst >1.8) are marked
magenta in the Orc6 panels. The same cells are marked magenta also in the Mcm3 and
Cdt1 panels. n = 450 (cells examined in each panel). The red-boxed regions represent
cells with high Mcm3 levels and high DNA content, and the blue-boxed region
represents cells with low Mcm3 levels and high DNA content. (**D**) Single-cell plot
analysis of Mcm2 (top) and histone H3S10ph (bottom) in hTERT-RPE1 cells. Cells
were treated with DMSO as a control or 312 nM MG132 for the indicated time periods.
n = 450 (number of cells examined in each panel). The cells showing high H3S10ph
levels ($\log_2$[H3S10ph] >1) are marked magenta in the H3S10ph panels. The blue-circled
regions represent cells with low Mcm2 levels and high DNA content. The red-boxed
regions represent cells with high Mcm2 and high DNA content. (**E**) Efficiency of siRNA
knockdown of PSMD7 expression. hTERT-RPE1 cells were treated with siRNA for the
negative control or PSMD7 for 2 days before harvesting. Western blotting analysis of

whole-cell lysates was performed using antibodies against PSMD7 and GAPDH. The
positions of molecular mass markers (kDa) are indicated to the right. GAPDH is a
loading control. (**F**) Single-cell plot analysis of Orc6 and Mcm2 in hTERT-RPE1 cells
treated with negative control-siRNA (siControl) or PSMD7-siRNA (siPSMD7). Cells
were treated with the indicated siRNA for 2 days, fixed using the pre-extraction method,
and stained with Hoechst, anti-Orc6, and anti-Mcm2 antibodies. The cells showing high
Orc6 levels ($\log_2$[Orc6] >0.5) and high DNA content (Hoechst >1.8) are marked
magenta in the Orc6 panels. The same cells are marked magenta also in the Mcm2
panels. The red-boxed region represents cells with high Mcm2 levels ($\log_2$[Mcm2] >⁻2)
and high DNA content (Hoechst >1.8), and the blue-boxed region represents cells with
low Mcm2 levels ($\log_2$[Mcm2] <⁻2) and high DNA content (Hoechst >1.8). n = 500
(number of cells examined in each panel).
833
Figure 5. Detection of cell cycle progression in tetraploid cells and Orc6 behavior
(**A**) Timing diagram of MG132 treatment and EdU labeling used in (**B**). hTERT-RPE1
cells were incubated with 312 nM MG132 for a day (day 1) (left panels in **B**), washed
with fresh medium without MG132, and incubated for another day (day 2) (right panels
in **B**). The cells were labeled with EdU for 30 min before the sample collection on day 1
and day 2. (**B**) Single-cell plot analysis of EdU (top), Mcm2 (middle), and Orc6
(bottom). Cells were fixed using the pre-extraction method and stained with Hoechst,
anti-EdU, anti-Mcm2, and anti-Orc6 antibodies. The orange dots represent S-phase cells
based on EdU intensities ($\log_2$[EdU] >2). The red-boxed regions represent cells with
high Mcm2 levels ($\log_2$[Mcm2] >⁻1) and high DNA contents (1.8< Hoechst <2.1). n =
600 (cells examined in each panel). (**B'**) The same data from (**B**) is reused, although it
was marked differently. The cells showing high Orc6 levels ($\log_2$[Orc6] >0.5) and high
DNA content (Hoechst >1.8) are marked magenta in the Orc6 panels. The same cells are
marked magenta also in the Mcm2 panels. The red-boxed region represents cells with
high Mcm2 levels ($\log_2$[Mcm2] >⁻0.5) and high DNA content (1.8< Hoechst <2.1), and
the blue-boxed region represents cells with low Mcm2 levels ($\log_2$[Mcm2] <0.5) and
high DNA content (1.8< Hoechst <2.1). (**C**) Representative fluorescence images of
Hoechst, Orc6, and Mcm2 staining used in single cell plot analysis in (**B'**). Red and
blue arrows indicated typical cells in the red and blue boxes, respectively, in Figure (**B'**).
Scale bars, 10 μm.
854
Figure 6. Effects of Orc6 depletion on the formation of tetraploid cells

(**A**) Timing diagram of siRNA treatment. Cells were treated with siRNA against a

negative control (siControl), Orc1 as a positive control (siOrc1), or Orc6 (siOrc6) for 42

h. (**B**) Western blotting analysis using antibodies against Orc6, Orc1, and histone H3 for

the whole-cell fraction from hTERT-RPE1 cells treated with siRNA against a negative

control (siControl), Orc1 (siOrc1), and Orc6 (siOrc6). The positions of molecular mass

markers (kDa) are indicated to the right. (**C**) Single-cell plot analysis of Mcm2 in

hTERT-RPE1 cells treated with siRNA (siControl, siOrc1, or siOrc6). Cells were

treated with siRNA, fixed using the pre-extraction method, and stained with Hoechst

and anti-Mcm2 antibody. n = 450 (number of cells examined in each panel). (**D**) Timing

diagram of siRNA treatment and addition of 312 nM MG132. hTERT-RPE1 cells were

treated with siRNA. After 18 h, MG132 was added to the medium at a final

concentration of 312 nM, and the cells were incubated for an additional 24 h in the

presence of siRNA. (**E**) Single-cell plot analysis of Mcm2 in the cells treated with

siRNA (siControl, siOrc1, or siOrc6). Cells showing high Mcm2 levels ($\log_2$[Mcm2]

>−1.5) and high DNA content (Hoechst > 1.8) are marked magenta. n = 450 (number of

cells examined in each panel). (**F**) Percentages of the magenta-marked cells

($\log_2$[Mcm2] >−1.5 and Hoechst >1.8) among the total G2 phase cells (Hoechst >1.8)

calculated based on (**E**). Columns and error bars represent the mean and standard

deviation of four independent experimental results. Statistical differences ($p = 0.0006$)

were determined using a one-sided t-test. (**G**) Western blotting analysis using antibodies

against Cdc6 and GAPDH for whole-cell lysates from cells treated with siRNA against

negative control (siControl) or Cdc6 (siCdc6), as described in (**D**). The positions of

molecular mass markers (kDa) are indicated to the right. (**H**) Single-cell plot analysis of

chromatin-bound Mcm2 in hTERT-RPE1 cells treated with siRNA (siControl or siCdc6)

as described in (**D**). Cells were treated with siRNA against the negative control

(siControl) or Cdc6 (siCdc6). After incubation with MG132, the cells were fixed using

the pre-extraction method and stained with Hoechst and anti-Mcm2 antibody. Cells

showing high MCM levels ($\log_2$[Mcm2] >−2) and high DNA content (Hoechst >1.8) are

marked magenta. n = 450 (number of cells examined in each panel).

Figure 8. Schematic diagram of Orc6 behavior

(**A**) Process with normal proteasome activity. After DNA replication, Cdt1 and Orc1

proteins are degraded, and Orc6 is removed from the chromatin, preventing MCM

reloading. These cells enter mitosis and produce normal diploid cells. Orc2–5 represents

a complex of Orc2, Orc3, Orc4, and Orc5. (**B**) When cells lose proteasome activity,

Orc6, Orc1, Cdt1, and MCM are reloaded onto chromatin, producing tetraploid cells by
skipping mitosis.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A. **Normal proteasome activity**

Mitosis

Diploid

B. **Low proteasome activity**

Mitosis
Skip

Tetraploid

Figure 7