1

Proteasome-dependent Orc6 removal from chromatin upon S-phase entry safeguards against MCM reloading and tetraploidy

Yoko Hayashi-Takanaka[1,2,*], Ichiro Hiratani[2], Tokuko Haraguchi[1] and Yasushi Hiraoka[1*]

[1] Graduate School of Frontier Biosciences, Osaka University, 1-3 Yamadaoka, Suita 565-0871, Japan

[2] RIKEN Center for Biosystems Dynamics Research, 2-2-3 Minatojima-minamimachi, Chuo-ku, Kobe 650-0047 Japan

[*]Correspondence should be addressed to:

Yoko Hayashi-Takanaka

RIKEN Center for Biosystems Dynamics Research, 2-2-3 Minatojima-minamimachi, Chuo-ku, Kobe 650-0047 Japan

E-mail: yoko.hayashi.yr@riken.jp

or

Yasushi Hiraoka

Graduate School of Frontier Biosciences, Osaka University, 1-3, Yamadaoka, Suita, 565-0871, Japan

E-mail: hiraoka@fbs.osaka-u.ac.jp

Abstract

DNA replication is tightly regulated to occur only once per cell cycle, as untimely re-initiation can lead to aneuploidy, which is associated with early senescence and cancer. The pre-replication complex (comprising Orc1-6, Cdc6, Cdt1, and MCM) is essential for the initiation of DNA replication, but the dynamics and function of Orc6 during the cell cycle remain elusive. Here, we demonstrate that Orc6 associates with chromatin during G1-phase and dissociates upon S-phase entry. The dissociation of Orc6 from chromatin is dependent on proteasome activity, and inhibition of the proteasome leads to the accumulation of chromatin-bound Orc6, which promotes abnormal MCM loading after S-phase entry without undergoing mitosis in human immortalized hTERT-RPE1 cells. Following release from proteasome inhibition, cells with elevated levels of chromatin-bound Orc6 and MCM proceed to the next replication phase as tetraploid cells. Our findings suggest that the proteasome-dependent dissociation of Orc6 after DNA replication is critical for preventing inappropriate MCM reloading and tetraploid formation.

Keywords (3-6 words)

Orc6; DNA replication; MCM; proteasome; tetraploid; single-cell plot analysis

Introduction

Genomic DNA replication is fundamental to the proliferation of living organisms. In eukaryotes, this process is tightly regulated to ensure that all chromosomes replicate only once per cell cycle. Disruptions in this regulation can result in abnormal DNA replication, leading to genomic instability, which may promote senescence or oncogenesis (Thakur et al., 2022).

DNA replication begins with the assembly of pre-replication complexes (pre-RCs) at replication origins during G1 phase (Hu and Stillman, 2023). The pre-RC consists of four key components: the origin recognition complex (ORC), minichromosome maintenance (MCM) complex, Cdc6, and Cdt1. MCM, a replicative helicase, is composed of Mcm2–7 subunits (Bell and Botchan, 2013; Deegan and Diffley, 2016; Ishimi, 2018), while Cdc6 and Cdt1 facilitate the recruitment of MCM to chromatin. Upon entry into S-phase, the pre-RC disassembles. Aberrant loading of pre-RCs onto chromatin after the S-phase initiation can trigger DNA re-replication without mitosis (Vassilev and DePamphilis, 2017; Limas and Cook, 2019), underscoring the need for tight regulation of Cdt1 and Cdc6 levels by proteasome degradation through E3 ubiquitin ligases to prevent pre-RC formation beyond G1 phase (Nishitani et al., 2006; Walter et al., 2016).

The ORC is a six-subunit complex (Orc1–6), which binds chromatin to mark replication origins for pre-RC assembly during G1 phase (Bell and Labib, 2016; Ganier et al., 2019; Hu and Stillman, 2023). While Orc2–5 remain chromatin-bound throughout the cell cycle, Orc1 dissociates from chromatin after S-phase entry via ubiquitin-mediated proteolysis (Li and DePamphilis, 2002; Méndez et al., 2002; Tatsumi et al., 2003; Coulombe et al., 2019). Orc1–5 share structural similarities, whereas Orc6 is more similar

to TFIIB (Duncker et al., 2009; Liu et al., 2011; Feng et al., 2021), distinguishing it functionally from other ORCs. Orc6 can bind DNA independently of Orc1-5 (Xu et al., 2020) and has roles in chromosome segregation and cytokinesis (Prasanth et al., 2002; Chesnokov et al., 2003; Semple et al., 2006; Balasov et al., 2009; Bernal and Venkitaraman, 2011). Recent studies suggest additional functions for Orc6, including its involvement in replication fork progression, ATR activation, and mismatch repair (Lin et al., 2022). However, the specific role of human Orc6 in the cell cycle remains largely unexplored.

Proteasomes are the primary machinery for protein degradation in cells. Reduced proteasome activity is observed in replicative senescent cells (Sabath et al., 2020) and inhibiting proteasomes induces cellular senescence in normal human fibroblasts (Chondrogianni & Gonos, 2004; Chondrogianni et al., 2003; Torres et al., 2006; Ukekawa et al., 2004). Senescent cells, which bypass mitosis, exhibit chromatin-bound pre-RCs, suggesting a predisposition to enter S-phase as tetraploid cells (Johmura et al., 2014; Panopoulos et al., 2014; Zeng et al., 2023). However, it remains unclear whether reduced proteasome activity in cellular senescence is involved in pre-RC formation.

In this study, we explored the dynamics of Orc6 during the cell cycle in human cells using multicolor immunofluorescence-based single-cell plot analysis (hereafter referred to as "single-cell plot analysis"). This method allows for the quantitative measurements of Orc6 levels in individual cells from fluorescence microscopy images, eliminating the need for cell synchronization (Hayashi-Takanaka et al., 2020; Hayashi-Takanaka et al., 2021). Our findings suggest that the regulation of Orc6 localization throughout the cell cycle is crucial for proper pre-RC formation, and its dysregulation can lead to replication as tetraploid cells.

Results

Dissociation of Orc6 from chromatin during S-phase

To quantify chromatin-bound Orc6 in individual human cells, we performed single-cell

plot analysis using fluorescence microscopy to measure multiple intracellular

components in each cell (Hayashi-Takanaka et al., 2020; Hayashi-Takanaka et al.,

2021). An asynchronous population of hTERT-RPE1 cells was fixed using two different

methods: "direct fixation" and "pre-extraction" (Fig. 1A). The direct fixation method

detects whole nuclear proteins, while the pre-extraction method detects only the

chromatin-bound fraction (Hayashi-Takanaka et al., 2021). The fixed cells were stained

with Hoechst 33342 (a DNA indicator, referred to as "Hoechst"), EdU (an S-phase

marker), and an anti-Orc6 antibody (Fig. 1A). The same cells were also stained with an

anti-Orc2 antibody to detect the behavior of Orc2–5 subunits on chromatin throughout

the cell cycle. The specificity of these antibodies was validated by immunostaining as

shown in Fig. S1.

Fluorescence imaging revealed that Orc6 staining was relatively uniform in

the cell population when using the direct fixation method but showed variability among

the cells prepared by the pre-extraction method (compare upper and lower panels for

Orc6 in Fig. 1A). In contrast, Orc2 staining appeared punctate and remained consistent

across all cells in both methods (right panels, Fig. 1A). Single-cell plot analysis was

performed based on these images: the DNA content was determined by Hoechst

intensity normalized from 1 to 2, and the normalized intensities of EdU, Orc6, and Orc2

were plotted against the DNA content, where each dot represents a single cell (Fig. 1B).

EdU-positive S-phase cells (>-2 on the $\log_2$ scale), marked as orange dots in the EdU

panels (left panels, Fig. 1B), are also shown as orange dots in both the Orc6 and Orc2

panels (middle and right panels, Fig. 1B). In the direct fixation method, the whole

fraction of Orc6 levels remained relatively constant, ranging from −1 to 1 on the $\log_2$

scale throughout the cell cycle (upper middle panel, Fig. 1B). However, using the pre-

extraction method, chromatin-bound Orc6 levels were high during G1-phase, dropped

immediately after S-phase entry, and stayed low through S- and G2-phases (lower

middle panel, Fig. 1B). Orc2 levels remained relatively constant during the G1/S

transition and showed minimal changes during the S-phase in both fixation methods

(right panels, Fig. 1B). These results indicate that Orc6, but not Orc2, dissociates from

chromatin during S-phase in hTERT-RPE1 cells.

To determine if Orc6 dissociation occurs in other cell types, we analyzed

changes in chromatin-bound Orc6 levels during the cell cycle in cancer cell lines (HeLa

and U2OS cells) and normal fibroblast cells (IMR90 cells) using the pre-extraction

method (Fig. S2A). The reduction in Orc6 levels during the G1- to S-phase transition

was less pronounced in cancer cell lines (HeLa and U2OS cells, Fig. S2B) compared to

normal cell lines (IMR90 and hTERT-RPE1 cells, Fig. S2B). This suggests that the

regulation of chromatin-bound Orc6 levels during the cell cycle may differ between

normal and cancer cells, or among different cell types.

Involvement of the proteasome in Orc6 dissociation from chromatin

Since the levels of Cdt1, a component of the pre-RC, are regulated by proteasome

activity, we examined whether Orc6 is also a target for proteasome regulation using the

proteasome inhibitor MG132. hTERT-RPE1 cells were treated with 20 μM MG132 for

varying durations (1, 2, and 4 h), fixed using the two different fixation methods, and

stained with Hoechst and an anti-Orc6 antibody. The same cells were co-stained with

anti-Cdt1 and anti-Orc2 antibodies as controls for proteasome degradation (Fig. 2A).

Single-cell plot analysis revealed that Orc6 levels in the whole fraction (obtained by the

direct-fixation method) remained relatively unchanged for up to 2 h of MG132

treatment (blue dots, left panels in Fig. 2A). However, in the chromatin-bound fraction

(obtained by the pre-extraction method), Orc6 was predominantly bound to chromatin

during G1-phase and dissociated after S-phase entry under control conditions (DMSO

treatment). Upon the treatment with MG132 for 1 h, the decrease in Orc6 levels at the

G1- to S-phase transition was less pronounced (blue dots, right panels in Fig. 2A). After

2 h of MG132 treatment, Orc6 remained bound to chromatin after the S-phase, similar

to its levels in G1-phase.

Cdt1 levels increased in cells progressing through S and G2 phases after 1 h

of treatment and continued to increase with extended treatment in both fixation methods

(orange dots, Fig. 2A), indicating that proteasome degradation was functioning

normally under these experimental conditions. Orc2 levels remained relatively

unchanged for up to 4 h in the direct fixation method (green dots, left panels in Fig.

2A), and slightly increased after 4 h in the pre-extraction method (green dots, right

panels in Fig. 2A). These results suggest that Orc6, and to a lesser extent Orc2, is not

degraded by the proteasome during the S and G2 phases but instead dissociates from

chromatin via proteasome activity.

To compare the increase in Orc6 levels during the S-phase due to MG132

treatment with that of Cdt1 and Orc2, S-phase cells were selected based on Hoechst

intensity (1.1< Hoechst <1.75) from the single-cell plot analysis (Fig. 2A), and then the

fluorescence intensity was plotted during the time course of MG132 treatment (each dot

represents a single cell, Fig. 2B and S3D, E). The results showed little to no increase in

Orc6 and Orc2 levels in the whole fraction after up to 4 h of treatment (1–2-fold; whole

fraction, Fig. 2B), while a large increase in Cdt1 levels was observed with increasing

duration of MG132 treatment (up to 8-fold; whole fraction, Fig. 2B). In the chromatin-

bound fraction, Orc6 levels in S-phase cells showed a substantial increase with

prolonged MG132 treatment (up to 8-fold; chromatin-bound fraction, Fig. 2B), while

those in G1- and G2-phase cells increased mildly (up to 4-fold; chromatin-bound

fraction, Fig. S3D, E). The increase in chromatin-bound Cdt1 and Orc2 levels was

smaller compared to Orc6 (up to 5-fold and 3-fold, respectively; chromatin-bound

fraction, Fig. 2B). Orc6 binding to chromatin was not affected by the depletion of either

Orc1 or Orc2 (Fig. S3A-C), suggesting that the MG132-induced increase in chromatin-

bound Orc6 was independent of the presence of Orc1 or Orc2. Similar analyses for G1

and G2 phases showed that the changes in chromatin-bound Orc6 due to MG132 were

less pronounced than in S-phase (Fig. S3D, E).

To further confirm these results, we examined Orc6 protein levels in whole-cell

lysates and chromatin fractions obtained via sucrose cushion fractionation (Fig. 2C).

Cells were treated with 20 μM MG132 for varying durations (1, 2, and 4 h), and Orc6

levels were analyzed in whole-cell lysates and chromatin fractions by western blotting

(Fig. 2D). Orc6 levels in whole-cell lysates remained unchanged throughout MG132

treatment (whole, top row, Fig. 2D). In contrast, in the chromatin fractions, Orc6 levels

increased approximately 3-fold (top row, Fig. 2D). When treated with MG132 for 2–4

hours, the chromatin-bound fraction of Orc6 was almost equivalent to the total fraction

(Supplementary information "Blot transparency"), indicating that most Orc6 binds to

chromatin upon inhibition of proteasome activity. The amount of Cdt1 was also

examined as a control to confirm the effect of the proteasome inhibitor. Cdt1 levels

increased over time in both whole-cell lysates (approximately 7-fold) and chromatin

fractions (approximately 3-fold) after MG132 treatment. Orc2 levels remained relatively

unchanged, with only a slight increase (1.2–1.5-fold). These results are consistent with

the single-plot analysis (Fig. 2A, B, S3D, E), suggesting that Orc6 dissociates from

chromatin during the S-phase via proteasome activity, without undergoing degradation.

IMR90 cells with repeated passages exhibit high levels of chromatin-bound Orc6

after S-phase completion

In our examination of Orc6 protein levels across various cell types, we observed that

IMR90 cells (human diploid fibroblasts) undergoing repeated passages displayed both

high levels of chromatin-bound Orc6 and enlarged nuclei (Fig. 3A). To explore the

relationship between cellular aging and pre-RC formation, we analyzed the dynamics of

chromatin-bound Orc6 and MCM levels during the cell cycle in IMR90 cells at different

passage numbers (Fig. 3A). Cells from passage 18 (younger cells) and passage 32 (older

cells, containing cells with large nuclei) were fixed using the pre-extraction method and

stained with Hoechst, anti-Orc6, and anti-Mcm2 antibodies. Single-cell plot analysis

showed that in passage 18 cells, the dynamics of chromatin-bound Orc6 and Mcm2

throughout the cell cycle were similar to those seen in hTERT-RPE1 cells (Fig. S4A).

Specifically, chromatin-bound Mcm2 levels increased during G1-phase, peaked at the

G1/S transition, and then decreased during the S-phase, reaching their lowest levels in

G2-phase (upper right panel in Fig. 3B) (Håland et al., 2015; Hayashi-Takanaka et al.,

2021; Zeng et al., 2023; Zhou et al., 2020). Chromatin-bound Orc6 levels were high in

G1-phase and decreased during S/G2-phase, similar to those seen in hTERT-RPE1 cells

(upper left panel in Fig.3B, second right in Fig. S2B). However, in passage 32 cells, a

subset of cells exhibited abnormally high levels of Orc6 (Orc6 >0.85 on the $\log_2$ scale)

after DNA replication was completed (1.8<Hoechst <2.1, magenta dots in the Orc6

panel, lower left panel in Fig. 3B). When these cells with high Orc6 levels (magenta

dots) were plotted on the Mcm2 panel, most of them displayed elevated Mcm2 levels

(red-boxed region, Fig. 3B) instead of the expected lower levels (blue-boxed region,

Fig. 3B). This suggests that in passage 32 IMR90 cells, pre-RC formation, which

normally occurs in G1 phase, was abnormally retained in G2 phase. These cells

progressed further through the cell cycle, resulting in the formation of cells with large

nuclei (2<Hoechst<4 in lower panels, Fig. 3B). Such cells were rarely observed in

passage 18 (upper left panel in Fig. 3B). In passage 18 (young cells), low levels of SA-

βGal and p21 protein, markers of cellular senescence, were observed (Fig. S4A). In

contrast, in passage 32 (aged cells, often containing large nuclei), some cells exhibited

high levels of SA-βGal and/or p21 protein (Fig. S4A). These results indicate that

IMR90 cells undergo cellular senescence between 18 and 32 passages. Given that

cellular senescence is known to be induced by impaired proteasome function

(Chondrogianni et al., 2003; Sabath et al., 2020), we hypothesized that the emergence of

cells with abnormal loading of Orc6 and MCM is related to the reduction of proteasome

activity.

Low proteasome activity causes aberrant loading of chromatin-bound Orc6 after

the S-phase

To test this hypothesis, we examined whether the reduction of proteasome activity in

immortalized cells (hTERT-RPE1 cells) leads to the emergence of cells with high Orc6

and MCM loading on chromatin after replication, similar to passage 32 IMR90 cells. To

determine the concentrations of MG132 that reproduce cellular senescence, hTERT-

RPE1 cells were treated with various concentrations of MG132 for longer periods (18

hours). Whereas most cells died at high concentrations used in Figure 2 (> 625 nM), cell

death was less frequent at a lower concentration of 312 nM MG132. Thus, we applied

MG132 at 312 nM or lower concentrations in the following experiments.

As a marker of cellular senescence, we first examined the levels of p21, a cell

cycle inhibitor that increases upon MG132 treatment with p53 stabilization (Kwon et

al., 2002; Zhu et al., 2007). Western blot analysis and immunostaining revealed elevated

p21 protein levels in MG132-treated cells compared to the DMSO control (Fig. 4A, Fig.

S4C), suggesting that proteasome inhibition by MG132 induces cell cycle arrest.

Similarly, cells positive for SA-βGal (senescence-associated β-galactosidase), a

hallmark of cellular senescence, were not detected on day 2 of treatment but appeared

on day 6 (Fig. S4D). Conversely, Ki67, a protein expressed in all phases of the cell

cycle except G0 phase (Scholzen and Gerdes, 2000), was widespread in MG132-treated

cells on day 1 but had disappeared in some cells on day 4 (Fig. S4E). Therefore,

treatment with low concentrations of MG132 for 18-24 h primarily led to cell cycle

arrest, while continuous treatment over several days induced cellular senescence.

Next, we examined the effect of low MG132 concentrations on Orc6

dynamics throughout the cell cycle. hTERT-RPE1 cells were treated with 78, 156, and

312 nM MG132 for 18 h. Cells were fixed using the pre-extraction method and stained

with anti-Orc6, anti-Mcm3, and anti-Cdt1 antibodies (Fig. 4C), followed by single-cell

plot analysis (Fig. 4C). We used anti-Mcm3 instead of anti-Mcm2 due to compatibility

with immunostaining; previous single-cell plot analysis confirmed that chromatin-

bound Mcm2 and Mcm3 exhibit similar cell cycle dynamics (Fig. S4F) (Hayashi-

Takanaka et al., 2021). MG132 treatment produced cells with high levels of chromatin-

bound Orc6 and high DNA content, marked as magenta dots in Orc6 panels (Orc6 >0.5

on the $\log_2$ scale, Hoechst >1.8, Fig. 4C). The number of these magenta cells increased

with increasing MG132 concentrations, whereas few were observed in the control

DMSO cells (Fig. 4C). These cells marked in Orc6 panels were also magenta-marked in

the Mcm3 and Ctd1 panels (middle and lower panels, Fig. 4C). Some of these cells

exhibited low Mcm3 signals (blue arrows in Fig. 4B; the blue-boxed region in Fig. 4C),

while others exhibited high Mcm3 levels (red arrows in lower panels in Fig. 4B; red-

boxed region in Fig. 4C), suggesting that these cells are in an abnormal G2-phase state.

Co-staining with Cdt1 revealed that most magenta-marked cells also had relatively high

levels of Cdt1 (Fig. 4C). Based on these observations, we hypothesized that the red-

boxed cells with high Mcm3 levels represent G1-phase cells, where pre-RC formation

had occurred, and the cells were ready to enter the S-phase.

To confirm the involvement of proteasomes in the abnormal pre-RC

formation after the S-phase, we examined the effects of depleting PSMD7, a subunit of

the 26S proteasome necessary for proteolytic activity (Shi et al., 2018), using siRNA

treatment. Cells were treated with PSMD7-targeting siRNA for 2 days, and Western

blotting showed that PSMD7 levels were reduced to approximately 20% of those in

control siRNA-treated cells (Fig. 4D). Under these conditions, cells were fixed using the

pre-extraction method, stained with anti-Orc6 and anti-Mcm2 antibodies, and subjected

to single-cell plot analysis (Fig. 4E, Fig. S4F). The results showed that PSMD7

depletion led to cells with high chromatin-bound Orc6 levels and high DNA content

(magenta dots, Orc6 >0.5 on the $\log_2$ scale, Hoechst >1.8, Fig. 4E). Some of these cells

had low Mcm2 levels (blue-boxed region, Fig. 4E), whereas others had high Mcm2

levels (red-boxed region, Fig. 4E), suggesting that proteasomal defects contribute to the

abnormal loading of Orc6 onto chromatin after the S-phase.

To investigate whether Orc6 binding to chromatin depends on Cdt1, we

examined the turnover of Cdt1. As the degradation of Cdt1 is regulated by enzymes

with NEDD8-activating enzyme activity (Lin et al., 2010), we suppressed Cdt1

degradation by treating cells with MLN4924, an inhibitor of the NEDD8-activating

enzyme. Treatment with 250 nM MLN4924 for 24 h significantly increased chromatin-

bound Cdt1 levels after the S phase; however, Orc6 levels did not increase beyond those

observed in the G1 phase, and MCM did not show abnormal loading (Fig. 4F). These

results suggest that the chromatin-bound Orc6 is regulated by the proteasome activity

and is not directly dependent on the presence of Cdt1 on chromatin.

Inhibition of proteasome activity causes pre-RC formation without mitosis

To examine whether cells with high levels of Orc6 and MCM, accompanied by high

DNA content (red box in Fig. 4C), appear without entering mitosis due to MG132

treatment, we performed a time-course analysis to determine the timing of their

appearance. We examined cell cycle progression using anti-Mcm2 antibody and anti-

histone H3 serine 10 phosphorylation (H3S10ph; an M-phase marker) antibodies

(Hayashi-Takanaka et al., 2009). In the case of control treated with DMSO, cells with

high histone H3S10ph signals were observed (Fig. 5A; magenta dots, bottom left in Fig.

5B), indicating normal cell cycle progression. The percentage of histone H3S10ph-

positive cells (magenta dots, ~5% of total) is consistent with the duration of mitosis in

hTERT-RPE1 cells from our previous study (~1 h, Hayashi-Takanaka et al., 2009;

Hayashi-Takanaka et al., 2021). In contrast, after MG132 treatment, cells with high

histone H3S10ph signals gradually disappeared by 8 h. During the first 4–12 h of

treatment, cells with high DNA content (Hoechst ~2) and low Mcm2 levels gradually

increased (blue-circled regions, Fig. 5B), suggesting accumulation of G2-phase cells.

Prolonged MG132 treatments (16–24 h) led to an increase in cells with high DNA

content (Hoechst ~2) and high Mcm2 levels (red-boxed regions, Fig. 5B), while no M-

phase cells were detected. These results suggest that cells with high DNA content

(Hoechst, ~2) and high Mcm2 levels represent tetraploid G1 cells produced by skipping

the M phase.

Using the same conditions, we analyzed the dynamics of chromatin-bound

Orc6 and Cdt1. Chromatin-bound Orc6 levels increased in S-phase cells by 4 h after

MG132 addition (upper panel in Fig. 5C), similar to those observed with higher

concentrations of MG132 (20 μM, Fig. 2). Chromatin-bound Cdt1 levels showed little

change 4 h after treatment (Fig. 5C) but increased by 8 h in cells with Hoechst value

around 2. The number of cells with high Orc6 and Cdt1 levels continued to increase by

12 h (orange-boxed regions, Fig. 5C), while cells with high levels of MCM appeared

later at 16-20 h (red-boxed regions, Fig. 5B). These results suggest that MCM does not

immediately bind to chromatin following the binding of Orc6 and Cdt1. We also

performed experiments combining MCM and Orc6 and obtained essentially the same

results (Fig. S5).

Cells forming pre-RC via proteasome inhibition can progress through the cell cycle

as tetraploids

In assessing cells with high DNA content and elevated levels of Orc6 and MCM on

chromatin, we hypothesized that these cells are in the tetraploid G1 phase and may

subsequently progress through the cell cycle. To test this hypothesis, cells were treated

with 312 nM MG132 for 24 h (day 1), washed, and then cultured in MG132-free

medium for an additional 24 h (day 2) (Fig. 6A). EdU was added to the culture medium

30 min before fixation at the end of both days to label cells undergoing DNA

replication. Cells were then fixed using the pre-extraction method and stained with anti-

Mcm2, and anti-Orc6 antibodies (Fig. 6B), and subjected to single-cell plot analysis

(Fig. 6C). EdU-positive cells were marked as orange dots, and the same cells were also

identified in the Mcm2 and Orc6 panels (Fig. 6C). On day 1, the number of EdU-

positive cells was limited, with most cells arrested at Hoechst levels of 1 or 2. Among

the cells with Hoechst levels around 2, a subset exhibited high levels of Mcm2 (red-

boxed regions: Hoechst >1.8 and Mcm2 >−1.5 on the $\log_2$ scale), accounting for 33.3

±3.7% of the total (mean and standard error of the mean in three independent

experiments, Fig. 6B). On day 2, after MG132 removal, EdU-positive cells appeared in

the Hoechst range of 2–4 (orange arrows in Fig.6B; orange dots in day 2 panels),

comprising 7.68 ±0.40 % of the total (mean and standard error of the mean in three

independent experiments, Fig. 6B). These results indicate that cells with Hoechst levels

of 2–4 underwent another round of DNA replication, suggesting whole-genome

duplication (EdU panels, day 2, Fig. 6C). The Mcm2 patterns were similar between

cells with Hoechst levels 1–2 and 2–4 (Mcm2 panel, day 2, Fig. 6C), with Mcm2 levels

increasing in G1-phase and decreasing during S-phase, reaching their lowest levels in

G2-phase. These results suggest that the cells with high MCM levels after S-phase

induced by proteasome inhibition (red-boxed region, Fig. 6C) were in G1-phase of

tetraploidy, capable of undergoing an additional round of DNA replication.

To further clarify the cell cycle phases of cells with high Orc6 levels and a

Hoechst value around 2, these cells were marked as magenta dots in the Orc6 panels

(Orc6 > 0.5 on the $\log_2$ scale, Fig. 6C'). The same cells were also marked as magenta

dots in the Mcm2 panels (Fig. 6C'). As observed in Fig. 4D, the population of cells with

high Orc6 and Hoechst ~2 on day 1 split into two groups (Fig. 6C'): one with high

Mcm2 levels and the other with relatively low Mcm2 levels (red and blue arrows,

respectively, Fig.6B; red- and blue-boxed regions, respectively, Fig. 6C'). On day 2,

however, most of the high Orc6 population exhibited high Mcm2 levels. These results

suggest that Orc6 binds to chromatin before MCM loading, followed by the next

replication phase.

Involvement of Orc6 in MCM loading in mitosis-skipped cells

To determine whether the presence of Orc6 on chromatin after S-phase entry contributes

to MCM loading, we investigated MCM levels on chromatin following Orc6 depletion

via siRNA in hTERT-RPE1 cells (Fig. 7). First, we examined the effect of Orc6 siRNA

(siOrc6) on MCM loading during G1-phase, together with Orc1, which is a subunit

essential for MCM loading (Coulombe et al., 2019; Ogawa et al., 1999; Samson et al.,

2016) and is known to be degraded in a proteasome-dependent manner after the S-phase

entry (Li and DePamphilis, 2002; Méndez et al., 2002; Tatsumi et al., 2003). Western

blot analysis confirmed the efficient depletion of both Orc1 and Orc6 after 42 h of

siRNA treatment (Fig. 7A, B). Then, cells were fixed using the pre-extraction method,

stained with Hoechst and anti-Mcm2 antibody, and subjected to single-cell plot

analysis. Single-cell plot analysis showed that cells treated with siOrc1 or siOrc6

accumulated at G1 phase (81.0% and 71.3%, middle and right panels, respectively in

Fig. 7C) compared with control siRNA (66.2%, left panel in Fig. 7C). To quantify

MCM loading, cells with high MCM levels (cyan dots, half of the maximum value of

Mcm2 in Hoechst <1.125, Fig. 7C) were expressed as a percentage of the cells with

Hoechst signals >1.125 (Fig. 7D). The percentage of cells with high MCM levels was

significantly reduced in Orc1-depleted cells ($3.8 \pm 0.8\%$ compared to $26.4 \pm 6.8\%$ in

siControl; Fig. 7D). Orc6 depletion caused a relatively small reduction in Mcm2 loading

($9.6 \pm 2.3\%$ in Fig. 7D). These results suggest that Orc6 plays a supporting role, while

Orc1 plays a critical role in MCM loading.

We next examined the effect of Orc6 depletion on Mcm2 loading induced by

proteasome inhibition. Cells were treated with siRNA for 18 h, followed by MG132

treatment for 24 h while maintaining siRNA treatment (Fig. 7E). After treatment, cells

were fixed using the pre-extraction method, stained with Hoechst and anti-Mcm2

antibody, and analyzed using single-cell plot analysis (Fig. 7F; Fig. S6A). Cells with

doubled DNA content (Hoechst ~2) accumulated by MG132 treatment in all siControl,

siOrc1 and siOrc6 (Fig. 7F). Cells with high MCM levels and Hoechst ~2 (magenta

dots, Hoechst >1.8 and Mcm2 >−1.5 on the $\log_2$ scale, Fig. 7F) were quantified as a

percentage of the total cells with Hoechst >1.8 (Fig. 7G). Orc1 depletion reduced the

percentage of high-MCM cells to $5.7 \pm 1.2\%$ (Fig. 7G), underscoring the importance of

Orc1 in MCM loading in tetraploid G1 cells. Orc6 depletion modestly reduced the

percentage of high-MCM cells ($13.3 \pm 4.2\%$) compared to the siControl ($35.2 \pm 3.4\%$).

These results suggest that while Orc6 contributes to MCM reloading in tetraploid G1

cells, its role is less prominent than that of Orc1.

We also examined a role of Cdc6, another pre-RC component, in MCM

loading after siRNA treatment. Western blot analysis confirmed effective Cdc6

reduction (Fig. 7H). Cells were treated with siCdc6 for 18 h, followed by MG132 for 24

h (Fig. 7E). The cells were then fixed using the pre-extraction method and stained with

Hoechst and anti-Mcm2 antibody. Single-cell plot analysis revealed a reduction in

tetraploid G1 cells high MCM levels (magenta dots, Hoechst >1.8, and Mcm2 >−1.5 on

the $\log_2$ scale) after Cdc6 depletion (Fig. 7I, J; Fig. S6B), similar to the effect of Orc6

depletion. This suggests that Orc6 contributes as much as Cdc6 to MCM loading during

the tetraploid G1-phase in hTERT-RPE1 cells.

In summary, during G1-phase, Orc6 binds to chromatin along with other pre-

RC components. Upon S-phase entry, Cdt1 and Orc1 are degraded by proteasome

activity, while Orc6 dissociates from chromatin (Fig. 8A). Orc2–5 remain bound to

chromatin throughout the cell cycle. These cells proceed through mitosis, producing

normal diploid cells (Fig. 8A). However, in cells with reduced proteasome activity,

Orc6, Orc1, Cdt1, and MCM are reloaded onto chromatin. These cells bypass mitosis

and proceed into replication, leading to tetraploid formation (Fig. 8B).

Discussion

In this study, we demonstrated dynamic changes in chromatin-bound Orc6 levels during

the cell cycle in hTERT-RPE1 cells. The levels of chromatin-bound Orc6 increased

until the G1/S transition, decreased upon entry into the S-phase, and remained low until

the next G1-phase, despite the total amount of Orc6 remaining almost unchanged

throughout the cell cycle. Contrary to our results, previous western blot analysis using

synchronized HeLa cells showed that chromatin-bound Orc6 levels remained nearly

constant across different cell cycle phases (Méndez et al., 2002). However, our single-

cell plot analysis revealed a more pronounced difference in Orc6 levels between G1 and

S/G2 phases in normal cell lines (IMR90, and hTERT-RPE1 cells) compared to cancer

cell lines (HeLa and U2OS cells) (Fig. 1, Fig. S2). This suggests that the regulation of

chromatin-bound Orc6 during the cell cycle could serve as a distinguishing factor

between normal and cancerous cells.

We further demonstrated that the removal of Orc6 from chromatin after the S-

phase, through a proteasome-dependent mechanism, plays a crucial role in preventing

abnormal MCM loading. Aberrant MCM loading beyond the S-phase can initiate an

additional round of DNA replication without mitotic division, leading to tetraploidy,

which may contribute to cancer development. To prevent such occurrences, the

ubiquitin-proteasome system regulates the levels of several pre-RC factors, such as

Cdt1, Cdc6, and Orc1, at specific cell cycle phases (Li and DePamphilis, 2002; Méndez

et al., 2002; Nishitani et al., 2006; Tatsumi et al., 2003; Walter et al., 2016).

Interestingly, unlike other pre-RC factors, Orc6 is not degraded at the protein level but

is instead dissociated from chromatin via proteasome activity. This indicates that the

proteasome regulates pre-RC protein levels on chromatin in a distinct manner, thereby

preventing abnormal MCM loading. Certain carcinomas exhibit elevated Orc6 levels

(Pan et al., 2022; Tang et al., 2023), suggesting that excess Orc6 might interfere with its

chromatin dissociation, potentially leading to the reloading of pre-RC onto chromatin.

Our results also suggest that eliminating Orc6 from chromatin at the onset of the S-

phase is critical. Given that Orc6 is the most evolutionarily diverse ORC subunit (Dhar

and Dutta, 2000), its dissociation from chromatin during the S-phase may represent an

auxiliary mechanism uniquely evolved to prevent aberrant MCM loading.

The precise molecular mechanism driving Orc6 dissociation from chromatin

remains unclear. Orc6 lacks known ubiquitinated amino acid residues for subsequent

proteasome degradation (Coulombe et al., 2019). One possibility is that the chromatin-

bound Orc6 levels are regulated by factors degraded through proteasome activity, such

as Orc1 or Cdt1. However, our experiments showed that reducing Orc1 and Cdt1 levels

did not affect the chromatin-bound Orc6 levels. Another plausible explanation involves

post-translational modifications, such as phosphorylation, as observed with *S. cerevisiae*

Orc6 (Chen and Bell, 2011; Chen et al., 2007). Notably, S-phase-specific

phosphorylation of Orc6 has been shown to contribute to genome stability via DNA

damage checkpoints (Lin et al., 2023). These modifications may facilitate Orc6

dissociation from chromatin, possibly mediated by the degradation of modifying

enzymes through proteasome activity. In addition, proteasomes may have non-

proteolytic functions. For example, the ATPase activity within the proteasome can

unwind DNA (Makino et al., 1996; Makino et al., 1999), suggesting that a proteasome

lacking proteolytic activity might still facilitate chromatin binding by ORC, including

Orc6. Furthermore, inhibition of proteasome activity has been associated with

nucleosome hyperacetylation at transcription start sites (Kinyamu et al., 2020), which

overlaps with replication origins (Sugimoto et al., 2018). This hyperacetylation may

promote Orc6 binding to chromatin.

A decline in proteasome activity affects a variety of cellular processes,

including senescence (Huang et al., 2022; Vilchez et al., 2014). Recent studies show

that cellular senescence is associated with a tetraploid state in G1-phase, which occurs

when cells bypass M-phase (Johmura et al., 2014; Panopoulos et al., 2014; Zeng et al.,

2023). In agreement, we observed that proteasome inhibition by MG132 led to

tetraploid G1 cells, and subsequent removal of the inhibitor restored proteasome

activity, promoting cell progression into S-phase (Figs. 4, 5). This implies that age-

related declines in proteasome activity may contribute to the production of tetraploid G1

cells, increasing the risk of cancer development.

478   In conclusion, our results show that normal cells can progress to a tetraploid

G1-phase when proteasome activity is reduced, a key step linked to whole-genome

duplication and carcinogenesis (Bielski et al., 2018; Gemble et al., 2022; López et al.,

2020). This study contributes to our understanding of the interplay between replication

factor dynamics and proteasome activity, further connecting these processes to

tetraploid cell formation, senescence, and cancer development.

Materials and Methods

Antibodies for immunofluorescence staining and Western blotting

The primary antibodies used were anti-Orc6 (1:300 for immunofluorescence staining

(IF) and 1:1000 for Western blot (WB); RRID:AB_670295, 3A4, sc-32735; Santa Cruz

Biotechnology), anti-Orc2 (1:300 for IF and 1:1000 for WB; RRID:AB_592395, M055-

3, MBL), anti-Cdt1 (1:300 for IF and 1:1000 for WB; RRID:AB_10896851, D10F11;

Cell Signaling Technology), anti-Mcm3 (1:1000 for IF; sc-390480; Lot#J1118, Santa

Cruz Biotechnology; 1:300 (IF); RRID:AB_398024, Anti-BM28; 610701, BD

transduction), anti-Mcm2 (1:1000 for IF and WB; RRID:AB_2142137, D7G11, Cell

Signaling Technology), anti-Orc1 (1:100 for IF and 1:300 for WB; F-10, sc-398734,

Santa Cruz Biotechnology), anti-GAPDH (1:1000 for WB; RRID:AB_1903993, 14C10;

Cell Signaling Technology), anti-histone H3 (1G1) (1:1000 for WB; Nozawa et al.,

2010), anti-PSMD7 (1:1000 for IF and WB; EPR13517, ab181072; Abcam), anti-Cdc6

(1:300 for IF and 1:1000 for WB,RRID:AB_627236, sc-9964, Santa Cruz

Biotechnology), anti-p21 (1:1000 for IF and WB; RRID:AB_823586, 12D1; Cell

Signaling Technology), and anti-histone H3S10ph (1:2000 for IF; CMA311, (Hayashi-

Takanaka et al., 2009)). The specificity of these antibodies used for immunostaining is

shown in Fig. S1. We previously confirmed by single-cell plot analysis based on

immunostaining that chromatin-bound Mcm2 and Mcm3 showed similar dynamics

during the cell cycle (Hayashi-Takanaka et al., 2021).

For fluorescent staining, the secondary antibodies were as follows: Alexa 488

donkey anti-mouse IgG (RRID:AB_2341099, 715-545-151; Jackson ImmunoResearch),

Cy5 donkey anti-mouse IgG (RRID:AB_2340820, 715-175-151; Jackson

ImmunoResearch), Cy3 donkey anti-Rat IgG (RRID:AB_2340667, 712-165-153;

Jackson ImmunoResearch), Alexa 488 donkey anti-rabbit IgG (RRID:AB_2313584,

711-545-152; Jackson ImmunoResearch) and Cy3 donkey anti-rabbit IgG

(RRID:AB_2340585, 711-005-152; Jackson ImmunoResearch).

Cell culture and chemicals

hTERT-RPE1 (ATCC, CRL-4000), IMR90 (ATCC, CCL-186), U2OS (ATCC, HTB-

96), and HeLa (RCB0007) cell lines were grown in a culture medium containing high-

glucose Dulbecco's modified Eagle's medium (Sigma-Aldrich) supplemented with

penicillin/streptomycin (100 units/mL penicillin, 100 μg/mL streptomycin; Fujifilm)

and 10% fetal calf serum (Thermo Fisher Scientific), as previously described. Unless

otherwise noted, the experiments were performed using hTERT-RPE1 cells.

siRNA transfection was performed using Lipofectamine RNAiMAX (Thermo

Fisher Scientific) with siRNAs (SASI_Hs01_00069342 for Orc6,

SASI_Hs01_00069343 for Orc6#2, SASI_Hs01_00038888 for Orc1,

SASI_Hs02_00341067 for Orc2, and Universal Negative Control #1, Sigma-Aldrich;

s11403 for PSMD7, s11401 for PSMD7#2, s2745 for Cdc6, and s2744 for Cdc6#2,

Ambion, Thermo Fisher Scientific) following the manufacturer's protocol.

MG132 (Fujifilm) and MLN4924 (MedChemExpress) were dissolved in

DMSO.

Fluorescence staining for single-cell plot analysis

Immunostaining was performed as previously described (Hayashi-Takanaka et al.,

2011; Hayashi-Takanaka et al., 2015; Hayashi-Takanaka et al., 2021). Briefly, the cells

($\sim 2 \times 10^4$ cells/cm^2) were plated on a 12-well plate with a coverslip (15 mm diameter;

No. 1S; Matsunami) at the bottom and cultured for over 1 day.

In the direct fixation method, cells were first fixed with 1 mL of the fixative

(2% paraformaldehyde [Electron Microscopy Sciences] dissolved in 250 mM HEPES

[pH 7.4]) for 10 min at room temperature (approximately 25°C), followed by

permeabilization for 15 min using 1 % Triton X-100 in PBS, and subsequently blocked

with Blocking One (Nacalai Tesque, Inc.).

In the pre-extraction method, before fixation, cells were first treated with 1 mL

of 0.2% Triton X-100 solution containing 20 mM HEPES (pH 7.4), 100 mM NaCl, and

300 mM sucrose for 3 min on ice. After removing the Triton X-100 solution, the cells

were fixed using the fixative described above. This fixation method removes the nuclear

membranes and loosely bound chromatin proteins.

The fixed cells were incubated with specific primary antibodies (0.2–1 μg/mL)

for 2 h at room temperature (approximately 25°C) and washed three times (10 min

each) using PBS. When cells were stained with multiple antibodies, the cells were

simultaneously incubated with these antibodies. The cells were treated with secondary

antibodies (1 μg/mL) and Hoechst (0.1 μg/mL) for 2 h at room temperature

(approximately 25°C) and subsequently washed three times with PBS. The cells were

mounted using the ProLong Glass Antifade Mountant (Thermo Fisher Scientific).

The replication foci were labeled by culturing the cells in a medium containing

10 μM 5-ethynyl-2′-deoxyuridine (EdU) for 30 min. Subsequently, the cells were fixed

using the direct fixation or pre-extraction method, and the resulting EdU signals were

detected with Alexa Fluor 647 using a Click-iT EdU Imaging Kit (Thermo Fisher

Scientific).

For staining of SA-βGal (senescence-associated β-galactosidase), live cells

were treated with the reagents according to commercial protocols (Cellular Senescence

Detection Kit, SPiDER-βGal. Dojindo).

Microscopy

Fluorescence images were obtained using a DeltaVision Elite system (GE Healthcare

Inc.) equipped with a pco.edge 4.2 sCMOS camera (PCO) and an Olympus 40×

UApo/340 oil immersion objective lens (NA 0.65 with iris) with DeltaVision Elite Filter

Sets (DAPI-FITC-TRITC-Cy5).

Single-cell plot analysis

Single-cell plot analysis was used to measure the levels of multiple intracellular

components based on fluorescence microscopic images and plot their correlation in

individual cells (Hayashi-Takanaka et al., 2020; Hayashi-Takanaka et al., 2021). The

fluorescence intensity of each nucleus was measured using NIS Elements software

(version 3.0; Nikon). The background intensity outside the cells was subtracted. The

nuclear areas of individual cells were determined by automatic thresholding using

Hoechst signals. The sum of the intensities (i.e., the average intensity × nuclear area) in

each nucleus was measured across all the fluorescence channels. Cells in the M-phase

were excluded from analysis except those shown in Fig. 5B, 5C. In Fig. 5B, 5C, M-

phase cells were used for the analysis. To plot the Hoechst intensity distribution of 450–

600 nuclei, the signal intensity in each nucleus was normalized relative to those with the

20th lowest and 20th highest intensities, designated as 1 and 2, respectively, because

nuclei with the lowest and highest intensities were sometimes outliers. Hoechst intensity

was plotted on a linear scale. The average intensities of Orc6, EdU, Orc2, Cdt1, Mcm2,

and Mcm3 were set to 1 (0 on the $\log_2$ scale) and plotted on the $\log_2$ scale. The

fluorescence intensities were normalized to time "0 h" (Figs. 2A, 2B), DMSO treatment

(Figs. 4C, 4F), the negative control siRNA (Figs. 4E, 7C, 7F, 7I, S3A, S3C, S4G, S5C,

S5D) and MG132 treatment for day 1 (Fig. 6C). The values relative to the average were

plotted using Mathematica (version 13.0) (Wolfram Research).

Preparation of chromatin fractions and whole-cell lysates

Native chromatin fractions (hereafter, referred to as "chromatin fractions") were

prepared by the method previously described (Quan et al., 2015; Sheu and Stillman,

2006) with some modifications (Hayashi-Takanaka et al., 2021). The cells were washed

with cold PBS, suspended in buffer A (50 mM HEPES [pH 7.4], 100 mM KCl, 2.5 mM

$MgCl_2$, 0.1 mM $ZnSO_4$, 1 mM ATP, 1 mM dithiothreitol [DTT], and protease inhibitor

cocktail [EDTA-free, Nacalai Tesque, Inc.]), and collected via centrifugation (1,300 g, 3

min, 4°C). The cells were subsequently treated with 200 µL of 0.2% Triton X-100 in

buffer A, layered with 200 µL of 30% sucrose cushion in buffer A containing 0.2%

Triton X-100, and centrifuged at 20,000 × *g* for 10 min at 4°C. The pellets were

collected as chromatin fractions and dissolved in 1× SDS loading buffer. The

suspension was then subjected to SDS-PAGE.

To prepare whole-cell lysates, the cells were washed with PBS and suspended

in an equal volume of 2× SDS loading buffer. The suspension was subjected to SDS-

PAGE.

Western blot analysis

Chromatin fractions and whole-cell lysates were suspended in 1× SDS loading buffer

and subjected to 10%, 12%, or 15% SDS-PAGE to separate proteins. Proteins were

transferred onto Immobilon-P PVDF membranes (Merck) using a semi-dry blotting

system (ATTO). After blocking with 5% skim milk (Nacalai Tesque, Inc.), the

membranes were incubated with the primary antibodies. After incubation with

peroxidase-conjugated secondary antibodies (GE Healthcare), the signals were detected

using ImmunoStar LD (Fujifilm Wako) and the ChemiDoc image system (BioRad). The

intensity of the signals was quantified using Image Lab software (BioRad).

Acknowledgements

We thank Ms. Chizuru Ohtsuki and Yoshino Kubota for their technical assistance, and

Drs. Hisao Masai, Haruhiko Asakawa, and Yasuhiro Hirano for insightful discussion.

We also thank Dr. Hiroshi Kimura for histone H3 antibodies and Dr. Hisao Masukata for

critically reading the manuscript.

This work was supported by JSPS KAKENHI grants: JP18H05530,

JP20K20582 and JP20K08698 (to IH); JP18H05533, JP20H00454 and JP23K05636 (to

YH); and JP18H05528 (to TH). This work was also supported by the Naito Foundation,

Urakami Foundation for Food and Food Culture Promotion, and the Foundation for

Promotion of Cancer Research in Japan (to YHT).

Data Availability

Source data are available from the corresponding authors upon reasonable request.

Conflict of interest

The authors declare no competing interests.

References

Balasov, M., Huijbregts, R. P. H. and Chesnokov, I. (2009). Functional analysis of an
Orc6 mutant in Drosophila. *Proc Natl Acad Sci U S A* **106**, 10672–10677.

Bell, S. D. and Botchan, M. R. (2013). The minichromosome maintenance replicative
helicase. *Cold Spring Harb Perspect Biol* **5**, a012807.

Bell, S. P. and Labib, K. (2016). Chromosome Duplication in Saccharomyces cerevisiae.
Genetics **203**, 1027–1067.

Bernal, J. A. and Venkitaraman, A. R. (2011). A vertebrate N-end rule degron reveals
that Orc6 is required in mitosis for daughter cell abscission. *J Cell Biol* **192**, 969–
978.

Bielski, C. M., Zehir, A., Penson, A. V., Donoghue, M. T. A., Chatila, W., Armenia, J.,
Chang, M. T., Schram, A. M., Jonsson, P., Bandlamudi, C., et al. (2018).
Genome doubling shapes the evolution and prognosis of advanced cancers. *Nat*
Genet **50**, 1189–1195.

Chen, S. and Bell, S. P. (2011). CDK prevents Mcm2–7 helicase loading by inhibiting
Cdt1 interaction with Orc6. *Genes Dev.* **25**, 363–372.

Chen, S., De Vries, M. A. and Bell, S. P. (2007). Orc6 is required for dynamic recruitment
of Cdt1 during repeated Mcm2–7 loading. *Genes Dev.* **21**, 2897–2907.

Chesnokov, I. N., Chesnokova, O. N. and Botchan, M. (2003). A cytokinetic function of
Drosophila ORC6 protein resides in a domain distinct from its replication activity.
Proc. Natl. Acad. Sci. U.S.A. **100**, 9150–9155.

Chondrogianni, N. and Gonos, E. S. (2004). Proteasome inhibition induces a senescence-
like phenotype in primary human fibroblasts cultures. *Biogerontology* **5**, 55–61.

Chondrogianni, N., Stratford, F. L. L., Trougakos, I. P., Friguet, B., Rivett, A. J. and
Gonos, E. S. (2003). Central role of the proteasome in senescence and survival of
human fibroblasts: induction of a senescence-like phenotype upon its inhibition
and resistance to stress upon its activation. *J Biol Chem* **278**, 28026–28037.

Coulombe, P., Nassar, J., Peiffer, I., Stanojcic, S., Sterkers, Y., Delamarre, A., Bocquet,
S. and Méchali, M. (2019). The ORC ubiquitin ligase OBI1 promotes DNA

659   replication origin firing. *Nat Commun* **10**, 2426.

660   **Deegan, T. D. and Diffley, J. F.** (2016). MCM: one ring to rule them all. *Current Opinion*
661   *in Structural Biology* **37**, 145–151.

662   **Dhar, S. K. and Dutta, A.** (2000). Identification and Characterization of the Human
663   ORC6 Homolog. *Journal of Biological Chemistry* **275**, 34983–34988.

664   **Duncker, B. P., Chesnokov, I. N. and McConkey, B. J.** (2009). The origin recognition
665   complex protein family. *Genome Biol* **10**, 214.

666   **Feng, X., Noguchi, Y., Barbon, M., Stillman, B., Speck, C. and Li, H.** (2021). The
667   structure of ORC-Cdc6 on an origin DNA reveals the mechanism of ORC
668   activation by the replication initiator Cdc6. *Nat Commun* **12**, 3883.

669   **Ganier, O., Prorok, P., Akerman, I. and Méchali, M.** (2019). Metazoan DNA replication
670   origins. *Curr Opin Cell Biol* **58**, 134–141.

671   **Gemble, S., Wardenaar, R., Keuper, K., Srivastava, N., Nano, M., Macé, A.-S., Tijhuis,**
672   **A. E., Bernhard, S. V., Spierings, D. C. J., Simon, A., et al.** (2022). Genetic
673   instability from a single S phase after whole-genome duplication. *Nature* **604**,
674   146–151.

675   **Håland, T. W., Boye, E., Stokke, T., Grallert, B. and Syljuåsen, R. G.** (2015).
676   Simultaneous measurement of passage through the restriction point and MCM
677   loading in single cells. *Nucleic Acids Res* **43**, e150–e150.

678   **Hayashi-Takanaka, Y., Yamagata, K., Nozaki, N. and Kimura, H.** (2009). Visualizing
679   histone modifications in living cells: spatiotemporal dynamics of H3
680   phosphorylation during interphase. *J Cell Biol* **187**, 781–790.

681   **Hayashi-Takanaka, Y., Yamagata, K., Wakayama, T., Stasevich, T. J., Kainuma, T.,**
682   **Tsurimoto, T., Tachibana, M., Shinkai, Y., Kurumizaka, H., Nozaki, N., et al.**
683   (2011). Tracking epigenetic histone modifications in single cells using Fab-based
684   live endogenous modification labeling. *Nucleic Acids Research* **39**, 6475–6488.

685   **Hayashi-Takanaka, Y., Maehara, K., Harada, A., Umehara, T., Yokoyama, S., Obuse,**
686   **C., Ohkawa, Y., Nozaki, N. and Kimura, H.** (2015). Distribution of histone H4
687   modifications as revealed by a panel of specific monoclonal antibodies.
688   *Chromosome Res* **23**, 753–766.

Hayashi-Takanaka, Y., Kina, Y., Nakamura, F., Becking, L. E., Nakao, Y., Nagase, T.,
Nozaki, N. and Kimura, H. (2020). Histone modification dynamics as revealed
by multicolor immunofluorescence-based single-cell analysis. *J Cell Sci* **133**,
jcs243444.

Hayashi-Takanaka, Y., Hayashi, Y., Hirano, Y., Miyawaki-Kuwakado, A., Ohkawa, Y.,
Obuse, C., Kimura, H., Haraguchi, T. and Hiraoka, Y. (2021). Chromatin
loading of MCM hexamers is associated with di-/tri-methylation of histone
H4K20 toward S phase entry. *Nucleic Acids Res* **49**, 12152–12166.

Hu, Y. and Stillman, B. (2023). Origins of DNA replication in eukaryotes. *Mol Cell* **83**,
352–372.

Huang, W., Hickson, L. J., Eirin, A., Kirkland, J. L. and Lerman, L. O. (2022). Cellular
senescence: the good, the bad and the unknown. *Nat Rev Nephrol* **18**, 611–627.

Ishimi, Y. (2018). Regulation of MCM2-7 function. *Genes Genet Syst* **93**, 125–133.

Johmura, Y., Shimada, M., Misaki, T., Naiki-Ito, A., Miyoshi, H., Motoyama, N.,
Ohtani, N., Hara, E., Nakamura, M., Morita, A., et al. (2014). Necessary and
Sufficient Role for a Mitosis Skip in Senescence Induction. *Molecular Cell* **55**, 73–
84.

Kinyamu, H. K., Bennett, B. D., Bushel, P. R. and Archer, T. K. (2020). Proteasome
inhibition creates a chromatin landscape favorable to RNA Pol II processivity. *J*
Biol Chem **295**, 1271–1287.

Kwon, Y. H., Jovanovic, A., Serfas, M. S., Kiyokawa, H. and Tyner, A. L. (2002). P21
functions to maintain quiescence of p27-deficient hepatocytes. *J Biol Chem* **277**,
41417–41422.

Li, C.-J. and DePamphilis, M. L. (2002). Mammalian Orc1 Protein Is Selectively
Released from Chromatin and Ubiquitinated during the S-to-M Transition in the
Cell Division Cycle. *Molecular and Cellular Biology* **22**, 105–116.

Limas, J. C. and Cook, J. G. (2019). Preparation for DNA replication: the key to a
successful S phase. *FEBS Lett* **593**, 2853–2867.

Lin, J. J., Milhollen, M. A., Smith, P. G., Narayanan, U. and Dutta, A. (2010). NEDD8-
targeting drug MLN4924 elicits DNA rereplication by stabilizing Cdt1 in S phase,

triggering checkpoint activation, apoptosis, and senescence in cancer cells. *Cancer Res* **70**, 10310–10320.

Lin, Y.-C., Liu, D., Chakraborty, A., Kadyrova, L. Y., Song, Y. J., Hao, Q., Mitra, J., Hsu, R. Y. C., Arif, M. K., Adusumilli, S., et al. (2022). Orc6 is a component of the replication fork and enables efficient mismatch repair. *Proc Natl Acad Sci U S A* **119**, e2121406119.

Lin, Y.-C., Liu, D., Chakraborty, A., Macias, V., Brister, E., Sonalkar, J., Shen, L., Mitra, J., Ha, T., Kajdacsy-Balla, A., et al. (2023). DNA Damage-Induced, S-Phase Specific Phosphorylation of Orc6 is Critical for the Maintenance of Genome Stability. *Mol Cell Biol* **43**, 143–156.

Liu, S., Balasov, M., Wang, H., Wu, L., Chesnokov, I. N. and Liu, Y. (2011). Structural analysis of human Orc6 protein reveals a homology with transcription factor TFIIB. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 7373–7378.

López, S., Lim, E. L., Horswell, S., Haase, K., Huebner, A., Dietzen, M., Mourikis, T. P., Watkins, T. B. K., Rowan, A., Dewhurst, S. M., et al. (2020). Interplay between whole-genome doubling and the accumulation of deleterious alterations in cancer evolution. *Nat Genet* **52**, 283–293.

Makino, Y., Yogosawa, S., Kanemaki, M., Yoshida, T., Yamano, K., Kishimoto, T., Moncollin, V., Egly, J. M., Muramatsu, M. and Tamura, T. (1996). Structures of the rat proteasomal ATPases: determination of highly conserved structural motifs and rules for their spacing. *Biochem Biophys Res Commun* **220**, 1049–1054.

Makino, Y., Yoshida, T., Yogosawa, S., Tanaka, K., Muramatsu, M. and Tamura, T. A. (1999). Multiple mammalian proteasomal ATPases, but not proteasome itself, are associated with TATA-binding protein and a novel transcriptional activator, TIP120. *Genes Cells* **4**, 529–539.

Méndez, J., Zou-Yang, X. H., Kim, S.-Y., Hidaka, M., Tansey, W. P. and Stillman, B. (2002). Human Origin Recognition Complex Large Subunit Is Degraded by Ubiquitin-Mediated Proteolysis after Initiation of DNA Replication. *Molecular Cell* **9**, 481–491.

Nishitani, H., Sugimoto, N., Roukos, V., Nakanishi, Y., Saijo, M., Obuse, C., Tsurimoto,

T., Nakayama, K. I., Nakayama, K., Fujita, M., et al. (2006). Two E3 ubiquitin ligases, SCF-Skp2 and DDB1-Cul4, target human Cdt1 for proteolysis. *EMBO J* **25**, 1126–1136.

Nozawa, R.-S., Nagao, K., Masuda, H.-T., Iwasaki, O., Hirota, T., Nozaki, N., Kimura, H. and Obuse, C. (2010). Human POGZ modulates dissociation of HP1alpha from mitotic chromosome arms through Aurora B activation. *Nat Cell Biol* **12**, 719–727.

Ogawa, Y., Takahashi, T. and Masukata, H. (1999). Association of fission yeast Orp1 and Mcm6 proteins with chromosomal replication origins. *Mol Cell Biol* **19**, 7228–7236.

Pan, Q., Li, F., Ding, Y., Huang, H. and Guo, J. (2022). ORC6 acts as a biomarker and reflects poor outcome in clear cell renal cell carcinoma. *J Cancer* **13**, 2504–2514.

Panopoulos, A., Pacios-Bras, C., Choi, J., Yenjerla, M., Sussman, M. A., Fotedar, R. and Margolis, R. L. (2014). Failure of cell cleavage induces senescence in tetraploid primary cells. *Mol Biol Cell* **25**, 3105–3118.

Prasanth, S. G., Prasanth, K. V. and Stillman, B. (2002). Orc6 Involved in DNA Replication, Chromosome Segregation, and Cytokinesis. *Science* **297**, 1026–1031.

Quan, Y., Xia, Y., Liu, L., Cui, J., Li, Z., Cao, Q., Chen, X. S., Campbell, J. L. and Lou, H. (2015). Cell-Cycle-Regulated Interaction between Mcm10 and Double Hexameric Mcm2-7 Is Required for Helicase Splitting and Activation during S Phase. *Cell Reports* **13**, 2576–2586.

Sabath, N., Levy-Adam, F., Younis, A., Rozales, K., Meller, A., Hadar, S., Soueid-Baumgarten, S. and Shalgi, R. (2020). Cellular proteostasis decline in human senescence. *Proc Natl Acad Sci U S A* **117**, 31902–31913.

Samson, R. Y., Abeyrathne, P. D. and Bell, S. D. (2016). Mechanism of Archaeal MCM Helicase Recruitment to DNA Replication Origins. *Mol Cell* **61**, 287–296.

Scholzen, T. and Gerdes, J. (2000). The Ki-67 protein: from the known and the unknown. *J Cell Physiol* **182**, 311–322.

Semple, J. W., Da-Silva, L. F., Jervis, E. J., Ah-Kee, J., Al-Attar, H., Kummer, L., Heikkila, J. J., Pasero, P. and Duncker, B. P. (2006). An essential role for Orc6

in DNA replication through maintenance of pre-replicative complexes. *EMBO J*
25, 5150–5158.

Sheu, Y.-J. and Stillman, B. (2006). Cdc7-Dbf4 Phosphorylates MCM Proteins via a
Docking Site-Mediated Mechanism to Promote S Phase Progression. *Molecular*
Cell **24**, 101–113.

Shi, K., Zhang, J.-Z., Zhao, R.-L., Yang, L. and Guo, D. (2018). PSMD7 downregulation
induces apoptosis and suppresses tumorigenesis of esophageal squamous cell
carcinoma via the mTOR/p70S6K pathway. *FEBS Open Bio* **8**, 533–543.

Sugimoto, N., Maehara, K., Yoshida, K., Ohkawa, Y. and Fujita, M. (2018). Genome-
wide analysis of the spatiotemporal regulation of firing and dormant replication
origins in human cells. *Nucleic Acids Res* **46**, 6683–6696.

Tang, M., Chen, J., Zeng, T., Ye, D.-M., Li, Y.-K., Zou, J. and Zhang, Y.-P. (2023).
Systemic analysis of the DNA replication regulator origin recognition complex in
lung adenocarcinomas identifies prognostic and expression significance. *Cancer*
Med **12**, 5035–5054.

Tatsumi, Y., Ohta, S., Kimura, H., Tsurimoto, T. and Obuse, C. (2003). The ORC1 Cycle
in Human Cells. *Journal of Biological Chemistry* **278**, 41528–41534.

Thakur, B. L., Ray, A., Redon, C. E. and Aladjem, M. I. (2022). Preventing excess
replication origin activation to ensure genome stability. *Trends Genet* **38**, 169–
181.

Torres, C., Lewis, L. and Cristofalo, V. J. (2006). Proteasome inhibitors shorten
replicative life span and induce a senescent-like phenotype of human fibroblasts.
J Cell Physiol **207**, 845–853.

Ukekawa, R., Maegawa, N., Mizutani, E., Fujii, M. and Ayusawa, D. (2004).
Proteasome inhibitors induce changes in chromatin structure characteristic of
senescent human fibroblasts. *Biosci Biotechnol Biochem* **68**, 2395–2397.

Vassilev, A. and DePamphilis, M. (2017). Links between DNA Replication, Stem Cells
and Cancer. *Genes* **8**, 45.

Vilchez, D., Saez, I. and Dillin, A. (2014). The role of protein clearance mechanisms in
organismal ageing and age-related diseases. *Nat Commun* **5**, 5659.

Walter, D., Hoffmann, S., Komseli, E.-S., Rappsilber, J., Gorgoulis, V. and Sørensen,
C. S. (2016). SCFCyclin F-dependent degradation of CDC6 suppresses DNA re-
replication. *Nat Commun* **7**, 10530.

Xu, N., You, Y., Liu, C., Balasov, M., Lun, L. T., Geng, Y., Fung, C. P., Miao, H., Tian,
H., Choy, T. T., et al. (2020). Structural basis of DNA replication origin
recognition by human Orc6 protein binding with DNA. *Nucleic Acids Research*
48, 11146–11161.

Zeng, J., Hills, S. A., Ozono, E. and Diffley, J. F. X. (2023). Cyclin E-induced replicative
stress drives p53-dependent whole-genome duplication. *Cell* **186**, 528-542.e14.

Zhou, Y., Pozo, P. N., Oh, S., Stone, H. M. and Cook, J. G. (2020). Distinct and
sequential re-replication barriers ensure precise genome duplication. *PLoS*
Genet **16**, e1008988.

Zhu, Q., Wani, G., Yao, J., Patnaik, S., Wang, Q.-E., El-Mahdy, M. A., Praetorius-Ibba,
M. and Wani, A. A. (2007). The ubiquitin-proteasome system regulates p53-
mediated transcription at p21waf1 promoter. *Oncogene* **26**, 4199–4208.

 Figure Legends

Figure 1. Detection of chromatin-bound Orc6 by single-cell plot analysis

(**A**) Representative fluorescence images of Hoechst, EdU, Orc6, Orc2, and merged

(Hoechst/EdU/Orc6) in hTERT-RPE1 cells prepared using the direct fixation method

(upper panels) and the pre-extraction method (lower panels). Scale bars, 10 μm. (**B**)

Single-cell plot analysis based on the images in (**A**). Each dot represents the EdU, Orc6,

and Orc2 intensities in a single individual cell against the Hoechst intensities. The

orange dots represent S-phase cells based on EdU intensities ($\log_2$[EdU] > −2). n = 450

(number of cells examined in each panel).

Figure 2. Effects of proteasome inhibitors on the dynamics of Orc6 during the cell

cycle

Single-cell plot analysis of Orc6 (top), Cdt1 (middle), and Orc2 (bottom) in hTERT-

RPE1 cells. Cells were treated with 20 μM MG132 for 0, 1, 2, or 4 h (timescale on the

top), fixed using the direct fixation method (left panels) and the pre-extraction method

(right panels). Each dot represents the intensity of Orc6, Cdt1, and Orc2 in a single

individual cell plotted against the Hoechst intensity. n = 450 (number of cells examined

in each panel). (**B**) Orc6, Cdt1, and Orc2 levels in the S-phase cells. Cells

corresponding to the S-phase (1.1 < Hoechst < 1.75) were selected from (**A**), and their

fluorescence intensities were plotted. The average intensity in MG132-untreated cells (0

h) was defined as 1. G1- and G2-phase cells are shown in Fig. S3D, E. (**C**) Schematic

diagram of the biochemical fractionation method for whole-cell lysates and chromatin

fractions. (**D**) Western blot analysis using antibodies against Orc6, Cdt1, Orc2, histone

H3, and GAPDH. hTERT-RPE1 cells were treated with 20 μM MG132 for the indicated

periods (h). Whole-cell lysates (left) and chromatin fractions (right) were prepared and

analyzed using western blotting. Histone H3 and GAPDH are loading controls.

Molecular weight markers (kDa) are indicated on the right.

Figure 3. Aberrantly high levels of chromatin-bound Orc6 and Mcm2 in aged

fibroblast cells

(**A**) IMR90 cells (passage numbers 18 and 32) were fixed using the pre-extraction

method and stained using Hoechst, anti-Orc6, and anti-Mcm2 antibodies.

Representative fluorescence images are shown. Arrows indicate relatively larger nuclei

with high Orc6 signals. Scale bars, 10 μm. (**B**) Single-cell plot analysis based on the

images in (**A**). Each dot represents the intensity of Orc6 and Mcm2 in an individual cell

plotted against Hoechst intensity. In the range of $1.8 <$ Hoechst < 2.1, the magenta dots

in the Orc6 panels represent cells with fluorescent signals of $\log_2(\text{Orc6}) > 0.85$. The

same cells are also marked magenta in the Mcm2 panels. The red-boxed regions

represent cells with high MCM levels ($\log_2[\text{Mcm2}] > -1.5$), and the blue-boxed regions

represent cells with low MCM levels ($\log_2[\text{Mcm2}] < -1.5$) in the range of $1.8 <$ Hoechst

< 2.1. n $= 450$ (number of cells examined in each panel).

Figure 4. MG132 treatment increases the cells with high chromatin-bound Orc6

after the S-phase

(**A**) Western blot analysis of whole-cell lysates using antibodies against p21 (cell cycle

marker) and GAPDH (loading control). hTERT-RPE1 cells were treated with DMSO or

312 nM MG132 for 1 day before harvesting. Molecular weight markers (kDa) are

indicated on the right. (**B**) Representative fluorescence images of Hoechst, Orc6, Mcm3,

and Cdt1 staining in hTERT-RPE1 cells treated with DMSO or 312 nM MG132 for 18

h. Cells were fixed using the pre-fixation method and stained with Hoechst, anti-Orc6,

anti-Mcm3, and anti-Cdt1 antibodies. G1 (yellow arrows) and G2 (blue arrows) were

labeled based on single-cell plot analysis in (**C**). Red arrows indicate a representative

cell categorized in the red-boxed region in (**C**). Scale bars, 10 μm. (**C**) Single-cell plot

analysis based on images of hTERT-RPE1 cells treated with DMSO or the indicated

concentration of MG132, fixed and stained as described in (B). The cells with high Orc6

levels ($\log_2[\text{Orc6}] > 0.5$) and high DNA content (Hoechst > 1.8) are marked magenta in

the Orc6 panels. The same cells are also marked magenta in the Mcm3 and Cdt1 panels.

The red-boxed region represents cells with high Mcm3 levels and high DNA content,

and the blue-boxed region represents cells with low Mcm3 levels and high DNA

content. n $= 450$ (cells examined in each panel). (**D**) Efficiency of siRNA knockdown of

PSMD7. Western blot analysis of hTERT-RPE1 cell lysates was performed with

antibodies against PSMD7 and GAPDH (loading control) after 2 days of siRNA

treatment. Molecular weight markers (kDa) are indicated on the right. (**E**) Single-cell

plot analysis of Orc6 and Mcm2 in hTERT-RPE1 cells treated with siControl or

siPSMD7. Cells were treated with siRNA for 2 days, fixed, and stained with Hoechst,

anti-Orc6, and anti-Mcm2 antibodies. Cells with high Orc6 levels ($\log_2[\text{Orc6}] > 0.5$)

and high DNA content (Hoechst > 1.8) are marked magenta in the Orc6 panels. The

same cells are also marked magenta in the Mcm2 panels. The red-boxed region

represents cells with high Mcm2 levels ($\log_2[\text{Mcm2}] > -1.5$) and high DNA content

(Hoechst > 1.8), and the blue-boxed region represents cells with low Mcm2 levels

($\log_2$[Mcm2] < ‑1.5) and high DNA content (Hoechst > 1.8). n = 500 (number of cells

examined in each panel). Another experiment with a different siRNA is shown in Fig.

S6A. (**F**) Single-cell plot analysis of hTERT-RPE1 cells treated with DMSO or 250 nM

MLN4924 for 24 h. The cells were fixed using the pre-extraction method, and stained

with Hoechst, anti-Orc6, anti-Mcm3, and anti-Cdt1 antibodies. n = 450 (number of cells

examined in each panel).

Figure 5. Time course of MG132-induced changes in Orc6, Mcm2 and Cdt1

indicate abnormal Pre-RC formation without mitosis

(**A**) Images of Hoechst, Mcm2, histone H3S10ph, and EdU in the DMSO-treated cells

fixed using the pre-extraction method. Combination of Mcm2, EdU and histone

H3S10ph signals identify cells in G1, S, G2, and M phases. The merged image shows

Hoechst (blue), Mcm2 (red), and histone H3S10ph (green). Scale bar, 10 µm. (**B**)

Single-cell plot analysis of Mcm2 (top), histone H3S10ph (bottom) in cells treated with

DMSO or 312 nM MG132 for the indicated durations. Cells with high H3S10ph levels

($\log_2$[H3S10ph] > 1) are marked magenta in the H3S10ph panels. Blue-circled regions

represent cells with low Mcm2 levels and high DNA content. Red-boxed regions

represent cells with high Mcm2 and high DNA content. n = 425 (number of cells

examined in each panel). (**C**) Single-cell plot analysis of Orc6 (top) and Cdt1 (bottom)

in hTERT-RPE1 cells treated with DMSO or 312 nM MG132 for the indicated

durations. Orange-boxed regions represent cells with high Orc6, Cdt1, and high DNA

content in the 12-h treatment.

Figure 6. Cells forming pre-RC via proteasome inhibition can progress through the

cell cycle as tetraploids

(**A**) Timing diagram for MG132 treatment and EdU labeling used in (**B**) and (**C**).

hTERT-RPE1 cells were incubated with 312 nM MG132 for a day (day 1) (left panels in

B), washed with fresh medium without MG132, and incubated for another day (day 2)

(right panels in **B**). The cells were labeled with EdU for 30 min before the sample

collection on day 1 and day 2. (**B**) Representative fluorescence images of Hoechst,

Orc6, and Mcm2 staining used in single-cell plot analysis in (**C** and **C'**). Red and blue

arrows indicate typical cells in the red and blue boxes in (**C** and **C'**). Orange arrows

indicate a tetraploid S-phase cell (orange dots of day 2 in (**C**)). Scale bars, 10 µm. (**C**)

Single-cell plot analysis of EdU (top), Mcm2 (middle), and Orc6 (bottom). The orange

dots represent S-phase cells based on EdU intensities ($\log_2$[EdU] > 2) with high DNA

contents (Hoechst > 1.8). The red-boxed regions represent cells with high Mcm2 levels

($\log_2$[Mcm2] > −1.5) and high DNA contents (1.8 < Hoechst < 2.1). n = 600 (cells

examined in each panel). (**C'**) The same data from (**C**) is reused but marked differently.

Cells with high Orc6 levels ($\log_2$[Orc6] > 0.5) and high DNA content (Hoechst > 1.8)

are marked magenta in the Orc6 panels. The same cells are also marked magenta in the

Mcm2 panels. The red-boxed region represents cells with high Mcm2 levels

($\log_2$[Mcm2] > −1.5) and high DNA content (1.8 < Hoechst < 2.1). The blue-boxed

region represents cells with low Mcm2 levels ($\log_2$[Mcm2] < -1.5) and high DNA

content (1.8 < Hoechst < 2.1).

Figure 7. Effects of Orc6 depletion on the formation of tetraploid cells

(**A**) Schematic diagram of siRNA treatment timing. hTERT-RPE1 cells were treated

with siRNA targeting a negative control (siControl), Orc1 as a positive control (siOrc1),

or Orc6 (siOrc6) for 42 h. (**B**) Western blot analysis of whole-cell fractions from cells

treated with siRNA against a negative control (siControl), Orc1 (siOrc1), and Orc6

(siOrc6), using antibodies against Orc6, Orc1, and histone H3. Molecular weight

markers (kDa) are indicated on the right. (**C**) Single-cell plot analysis of Mcm2 in

hTERT-RPE1 cells treated with siRNA (siControl, siOrc1, or siOrc6). siRNA-treated

cells were fixed using the pre-extraction method and stained with Hoechst and anti-

Mcm2 antibody. Cells with high Mcm2 levels (> half the maximum Mcm2 value,

excluding the 4 highest values among 450 cells) and low DNA content (Hoechst <

1.125) are marked cyan. n = 450 (number of cells examined in each panel). (**D**)

Percentages of cyan-marked cells (high MCM) among G1-phase cells (Hoechst <

1.125) based on (**C**). Columns and error bars represent the mean and standard error of

the mean of three independent experiments. Statistical difference (p = 0.004) was

determined using a one-sided t-test. (**E**) Timing diagram for siRNA treatment and

MG132 addition. hTERT-RPE1 cells were treated with siRNA. After 18 h of siRNA

treatment, MG132 was added at 312 nM for an additional 24 h, with continued siRNA

treatment. (**F**) Single-cell plot analysis of Mcm2 in hTERT-RPE1 cells treated with

siRNA (siControl, siOrc1, or siOrc6), with MG132 addition. Cells with high Mcm2

levels ($\log_2$[Mcm2] > −1.5) and high DNA content (Hoechst > 1.8) are marked magenta.

n = 450 (number of cells examined in each panel). Another experiment with different

siRNA is shown in Fig. S6A. (**G**) Percentage of magenta-marked cells ($\log_2$[Mcm2] >

−1.5 and Hoechst > 1.8) among G2-phase cells (Hoechst >1.8) based on (**F**). Columns

and error bars represent the mean and standard error of the mean of four independent

experimental results. Statistical differences (p = 0.0006) were determined using a one-

sided t-test. (**H**) Western blot analysis of whole-cell lysates from hTERT-RPE1 cells

treated with siControl or siCdc6), using antibodies against Cdc6 and GAPDH.

Molecular weight markers (kDa) are indicated on the right. (**I**) Single-cell plot analysis

of chromatin-bound Mcm2 in hTERT-RPE1 cells treated with siControl or siCdc6. After

MG132 treatment, cells were fixed using the pre-extraction method and stained with

Hoechst and anti-Mcm2 antibody. Cells with high MCM levels ($\log_2[\text{Mcm2}] > -1.5$)

and high DNA content (Hoechst > 1.8) are marked magenta. n = 450 (number of cells

examined in each panel). Another experiment with a different siRNA is shown in Fig.

S6B. (**J**) Percentage of magenta-marked cells ($\log_2[\text{Mcm2}] > -1.5$ and Hoechst > 1.8)

among G2-phase cells (Hoechst > 1.8) based on (**I**). Columns and error bars represent

the mean and standard error of the mean of three independent experiments. Statistical

difference ($p = 0.001$) was determined using a one-sided *t*-test.

Figure 8. Schematic diagram of Orc6 behavior

(**A**) Normal proteasome activity: After DNA replication, Cdt1 and Orc1 are degraded by

proteasome activity, and Orc6 dissociates from chromatin, preventing MCM reloading.

Cells enter mitosis and produce normal diploid cells. Orc2–5 represents a complex of

Orc2, Orc3, Orc4, and Orc5. (**B**) Loss of proteasome activity: Orc6, Orc1, Cdt1, and

MCM are reloaded onto chromatin without mitosis, allowing cells to re-enter the

replication phase as tetraploid cells.

Figure 1

Figure 2

A

IMR90

	Hoechst	Orc6	Mcm2
passage18			
passage 32			

B

● 1.8<Hoechst < 2.1 & log$_2$[Orc6] > 0.85

Orc6

passage 18
passage 32

log$_2$(Orc6)

Hoechst

Mcm2

log$_2$(Mcm2)

Hoechst

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A. **Normal proteasome activity**

Diploid

B. **Low proteasome activity**

Tetraploid

Figure 8

Supplemental Figures

Figure S1. Validation of antibodies used in this study

Western blot analysis of whole-cell lysates from hTERT-RPE1 cells using antibodies against Orc6, Orc1, Orc2, Cdt1, GAPDH, histone H3, Mcm2, Mcm3, PSMD7, Cdc6, and p21. Molecular weight markers (kDa) are indicated on the right of each blot. The percentage of polyacrylamide gel used is shown below each blot. Asterisks (*) indicate the proteins of interest detected at their expected molecular weights.

Figure S2. Behaviors of chromatin-bound Orc6 during the cell cycle in HeLa, U2OS, IMR90, and hTERT-RPE1 cells

(**A**) Representative fluorescence images of Hoechst and Orc6 staining in HeLa (left), U2OS (second left), IMR90 (second right), and hTERT-RPE1 (right) cells. Cells were fixed using the pre-extraction method and stained with Hoechst and anti-Orc6 antibody. The method of pre-extraction reproducibly makes the nuclei of HeLa cells smaller, but not in U2OS cells, for unknown reasons. Scale bars, 10 μm. (**B**) Single-cell plot analysis based on the images in (**A**). Each dot represents the intensities of Orc6 in an individual cell plotted against the Hoechst intensities. Magenta arrows show the difference in Orc6 levels from the 10[th] highest intensity in G1 (Hoechst < 1.1) to the Orc6 average intensity in early S-phase (1.15 < Hoechst < 1.4). n = 450 (number of cells examined in each panel).

Figure S3. Chromatin binding of Orc6 is independent of the presence of Orc1 and Orc2

(**A**) Single-cell plot analysis of Orc6 in hTERT-RPE1 cells treated for 24 h with siRNA against a negative control (siControl), Orc1 (siOrc1), Orc2 (siOrc2), and Orc6 (siOrc6), and fixed using the pre-extraction method to obtain the chromatin-bound fraction. The cells were stained with Hoechst and anti-Orc6 antibody. n = 450 (number of cells examined in each panel). (**B**) Western blot analysis using antibodies against Orc6, Orc1, Orc2, and histone H3. hTERT-RPE1 cells were treated for 24 h with siRNA (siControl, siOrc1, siOrc2, or siOrc6). Chromatin fractions were prepared from these cells using the sucrose cushion fractionation as described in Methods and subjected to western blot analyses. Histone H3 is a loading control. Molecular weight markers (kDa) are indicated on the right. (**C**) Single-cell plot analysis of Orc6 in hTERT-RPE1 cells treated for 24 h

with siRNA against a negative control (siControl), Orc1 (siOrc1), Orc2 (siOrc2), and Orc6 (siOrc6), and directly fixed to obtain the chromatin-bound fraction. The cells were stained with Hoechst and anti-Orc6 antibody. n = 450 (number of cells examined in each panel). (**D, E**) Orc6, Cdt1, and Orc2 levels in G1- **(D)** and G2-phases cells **(E)**. Cells corresponding to G1-phase (Hoechst < 1.1) or G2-phase (1.75 < Hoechst) are selected from (Fig. 2A), and their fluorescence intensities were plotted. The average intensity in MG132-untreated cells (0 h) was defined as 1.

Figure S4. Effect of MG132 treatment on hTERT-RPE1 cells

(**A**) Representative fluorescence images of Hoechst, SA-βGal and p21 staining in IMR90 cells (passage numbers 18 and 32). Cells were fixed using the direct fixation

method. **(B)** Single-cell plot analysis of IMR90 cells (passage 18) co-immunostained with EdU and Mcm3. Cells were treated with EdU for 30 min prior to collection and then fixed using the pre-extraction method. Each dot represents the intensities of EdU and Mcm3 in an individual cell, plotted against Hoechst 33342 intensities. Orange dots represent S-phase cells based on EdU intensities (log2(EdU) > –2). The number of cells examined in each panel is 450. **(C)** Representative fluorescence images of Hoechst and p21 staining in hTERT-RPE1 cells treated with DMSO or 312 nM MG132 on day 1. Cells were fixed using the direct fixation method. **(D)** Representative images of Hoechst and SA-βGal staining in cells treated with DMSO or 312 MG132 on day 2 and on day 6. Before the direct fixation method for Hoechst staining, cells were incubated with the reagent according to the manual. **(E)** Representative images of Hoechst and Ki67 staining in cells treated with DMSO or 312 nM MG132 on day 1 and on day 4. Cells were fixed using the direct fixation method. Red arrows indicate Ki67 negative cells. **(F)** Single-cell plot analysis of hTERT-RPE1 cells co-immunostained with Mcm2 and Mcm3. The number of cells examined in each panel is 450. **(G)** Single-cell plot analysis of Orc6 and Mcm2 in hTERT-RPE1 cells treated with negative control-siRNA (siControl) or two types of PSMD7-siRNA (siPSMD7). Cell conditions and immunostaining methods were the same as in Fig.4E. Scale bars, 10 μm.

Figure S5. Time course from MG132 treatment to the appearance of cells with high chromatin-bound MCM and Orc6

Single-cell plot analysis of Mcm2 (top) and Orc6 (bottom) in cells treated with DMSO or 312 nM MG132 for the indicated durations. The same conditions but different combinations are shown in Fig. 5B, 5C. n = 450 (number of cells examined in each panel).

Figure S6. Single-cell plot analysis in siRNA-treated cells

(A, B) Single-cell plot analysis of Mcm2 in cells treated with siOrc6 and siOrc6#2 (**A**), and siCdc6 and siCdc6#2 (**B**). Cells with high Mcm2 levels (log$_2$[Mcm2] > ‑1.5) and high DNA content (Hoechst > 1.8) are marked magenta. n = 450 (number of cells examined in each panel).

Western blot transparency_1

Fig. 7B **Fig. 7H**

Western blot transparency_2