Event-Related Potential markers of Subjective Cognitive Decline and Mild Cognitive Impairment during a sustained visuo-attentive task

A. A. Vergani[1,2], S. Mazzeo[4,6,7], V. Moschini[4], R. Burali[5], M. Lassi[1,2], L. G. Amato[1,2], J. Carpaneto[1,2], G. Salve-strini[5], C. Fabbiani[3,5], G. Giacomucci[3,4], C. Morinelli[4], F. Emiliani[3], M. Scarpino[5], S. Bagnoli[3], A. Ingannato[3], B. Nacmias[3,5], S. Padiglioni[4], S. Sorbi[3,4,5], V. Bessi[3,4,*], A. Grippo[5], A. Mazzoni[1,2]

[1] The BioRobotics Institute, Sant'Anna School of Advanced Studies, viale Rinaldo Piaggio 34, 56025 Pontedera-Pisa, Italy
[2] Department of Excellence in Robotics and AI, Sant'Anna School of Advanced Studies, viale Rinaldo Piaggio 34, 56025 Pontedera-Pisa, Italy
[3] Department of Neuroscience, Psychology, Drug Research and Child Health, Azienda Ospedaliero-Universitaria Careggi, Largo Brambilla 3, Florence, 50134, Italy.
[4] Research and Innovation Centre for Dementia-CRIDEM, Azienda Ospedaliero-Universitaria Careggi, Largo Brambilla 3, Florence, 50134, Italy.
[5] IRCCS Fondazione Don Carlo Gnocchi, via di Scandicci, 269, 50143 Florence, Italy
[6] Vita-Salute San Raffaele University, Via Olgettina, 58, 20132, Milano, Italy
[7] IRCCS Policlinico San Donato, Piazza Edmondo Malan, 2, 20097, San Donato Milanese, Italy

* Corresponding author:
Dr. Valentina Bessi
Department of Neuroscience, Psychology, Drug Research and Child Health,
University of Florence, AOU Careggi, Largo Brambilla 3, Florence, 50134, Italy.
Research and Innovation Centre for Dementia-CRIDEM, AOU Careggi, Florence, Italy.
Email: valentina.bessi@unifi.it.
Voice : (office) Tel +39 055-7948660

Abstract

Subjective cognitive decline (SCD), mild cognitive impairment (MCI), or severe Alzheimer's disease stages are still lacking clear electrophysiological correlates. In 178 individuals (119 SCD, 40 MCI, and 19 healthy subjects (HS)), we analysed event-related potentials recorded during a sustained visual attention task, aiming to distinguish biomarkers associated with clinical conditions and task performance. We observed condition-specific anomalies in event-related potentials (ERPs) during visual encoding (P1/N1/P2) and decision-making (P300/P600/P900): SCD individuals showed attenuated dynamics compared to HS, while MCI individuals showed amplified dynamics, except for P300, which matched clinical severity. ERP features confirmed a non-monotonic trend, with MCI showing higher neural resource recruitment. Moreover, task performance correlated with condition-specific ERP gain and latencies across early and late ERP components. These findings enhanced the understanding of the neural mechanisms underlying cognitive decline in SCD and MCI and suggested potential biomarkers for early diagnosis and intervention.

Highlights

- In encoding (P1/N1/P2) and decision (P600/P900) ERPs, SCD individuals showed attenuated dynamics compared to HS, while MCI individuals exhibited amplified dynamics compared to SCD.

- P300 dynamics matched clinical severity.

- MCI individuals demonstrated higher recruitment of neural resources, indicating a non-monotonic trend in ERP features between clinical conditions.

- Task performance correlated with condition-specific gain and latencies across multiple ERP components.

Key Words

Subjective Cognitive Decline (SCD); Mild Cognitive Impairment (MCI); EEG; Event-Related Potentials;

1. Introduction

Neurocognitive disorders affect 6-50 million people worldwide, with prevalence doubling every five years, particularly among those aged 50-80. This trend poses a significant societal burden, with various factors contributing to dementia, including neurological, systemic, and psychiatric conditions. Alzheimer's disease (AD) is the most prevalent cause of neurocognitive decline (1). AD involves the accumulation of beta-amyloid plaques and neurofibrillary tangles, leading to neurodegeneration and cognitive decline, eventually resulting in dementia. This process unfolds over decades, with amyloid buildup occurring years before symptoms. Stages range from subtle cognitive changes to full-blown dementia. The initial stage, Subjective Cognitive Decline (SCD), involves self-reported cognitive decline while performance on standardized tests remains within the normal range when adjusted for age, sex, and education (2,3). Mild Cognitive Impairment (MCI) occurs when pathological scores on neuropsychological tests are present without a significant impact on daily life activities. It serves as a transitional stage between normal aging and the more severe cognitive decline seen in dementia (4). In the realm of dementia research, SCD and MCI hold paramount significance as they fall within the spectrum of AD. Patients affected by these conditions present an opportunity for intervention with recently developed Disease-Modifying Therapies (DMTs) approved for AD (5,6). Indeed, it is widely acknowledged that DMTs should be administered during the early stages of the disease, prior to the onset of neurodegeneration (7).

Seeking reliable biomarkers for early AD diagnosis is crucial. Common biomarkers like MRI, FDG-PET, and CSF are invasive and not widely available. Hence, researchers explore accessible options, with EEG showing promise (8). Nevertheless, despite these efforts, only a limited number of studies have delved into this promising avenue (e.g., biomarking conditions as SCD and MCI (9–14), MCI against AD (15), across CSF (16) and ApoE ε-4 allele (17)).

Additionally, in dementia EEG studies, sensory event-related potentials are examined (e.g., auditory (18) and visual (19–22)). Specifically, visual event-related potentials suggest a compelling hypothesis about brain alterations in the visual system that could help detect early structural changes linked to anomalies in ERPs (23–25). For example, by recording EEG during a visuo-memory task, Waninger et al (26) found amplitude suppression of late positive potentials (∼400ms) in MCI against healthy subjects over right occipital and temporal channels. Other studies enquired early phase of visual processing as the encoding of stimulus: Krasodomska et al (27) found N95 wave dynamics alterations in AD, as other colleagues in last decays detect visual evoked potential anomalies in dementia patients (28,29). Hence, an unresolved critical aspect is how visual alterations manifest across various stages of cognitive decline.

Along with the early visual alterations associated with cognitive decline, abnormalities of the late post-stimulus ERP components related to the quality of decision-making are known. Among the best studied is the P300, which is altered in gain or latency in pathological conditions such as AD dementia (30–35), as well as for the P600, which has been observed that its abnormalities are associated with increased risk of conversion to dementia in MCI patients (36–42).

Following the hypothesis that the domain of visual faculties is important to highlight electrophysiological abnormalities, which can be traced back to cognitive decline, we performed an exploratory study by implementing a visual-attentive experimental paradigm and recorded cortical EEG activity during task execution. To this end, we first monitored the behavioural outcomes of SCD and MCI patients, and compared with Healthy Subjects (HS), then investigated the presence of ERP correlates specific to clinical condition and their performance, thereby enhancing our understanding of electrophysiology along the dementia continuum.

2. Methods

2.1 Clinical protocol

The clinical protocol of the PREVIEW project (ClinicalTrials.gov Identifier: NCT05569083) has been published previously (43). In brief, PREVIEW is a longitudinal study on Subjective Cognitive Decline started in October 2020 with the aim to identify features derived from easily accessible, cost-effective and non-invasive assessment to accurately detect SCD patients who will progress to AD dementia. All participants were collected in agree with the Declaration of Helsinki and with the ethical standards of the Committee

134 on Human Experimentation of Careggi University Hospital (Florence, Italy). The study was approved by
135 the local Institutional Review Board (reference 15691oss).

2.2 Participants
We enrolled 178 individuals (117F), including 119 SCD patients (85F), 40 MCI patients (24F), and 19
healthy individuals (8F). All participants underwent thorough family and clinical history evaluations,
neurological examinations, extensive neuropsychological assessments, premorbid intelligence estimation,
and depression evaluations.

The following inclusion criteria were adopted: satisfied criteria for SCD (44) or MCI (45); Mini Mental
State Examination (MMSE) score >24, corrected for age and education; normal functioning on the Activi-
ties of Daily Living (ADL) and the Instrumental Activities of Daily Living (IADL) scales unsatisfied criteria
for AD diagnosis according to National Institute on Aging-Alzheimer's Association (NIA-AA) criteria (46).
Exclusion criteria were history of head injury, current neurological and/or systemic disease, symptoms of
psychosis, major depression, substance use disorder; complete data loss of patients' follow-up; use of any
medication with known effects on EEG oscillations, such as benzodiazepines or antiepileptic drugs.

A subset of 58 patients underwent CSF collection for assessment of $A\beta_{42}$, $A\beta_{42}/A\beta_{40}$, total-tau (t-tau) and
phosphorylated-tau (p-tau). Among these, 31 patients also underwent cerebral amyloid-PET. Normal val-
ues for CSF biomarkers were: $A\beta_{42}$>670 pg/ml, $A\beta_{42}/A\beta_{40}$ ratio>0.062, t-tau<400 pg/ml and p-tau<60
pg/ml (47). Methods used CSF collection, biomarker analysis, and amyloid-PET acquisition and rating are
described in further detail elsewhere (43,48). Patients who underwent AD biomarker assessment, were
classified as A+ if at least one of the amyloid biomarkers (CSF $A\beta_{42}$, $A\beta_{42}/A\beta_{40}$ or amyloid PET) indicated
the presence of $A\beta$ pathology, and as A- if none of the biomarkers indicated the presence of $A\beta$ pathology.
In cases where there were conflicting results between CSF and Amyloid PET, only the pathological result
was considered. Patients were classified as T+ or T- based on whether their CSF p-tau concentrations
were higher or lower than the cut-off value, respectively. Similarly, patients were classified as N+ or N-
depending on whether their t-tau concentrations were higher or lower than the cut-off value. Using this
initial classification, we applied the NIA-AA Research Framework (49) to define the following groups:
ATN Negative (46 of 58; 29 SCD of 34 + 17 MCI of 24): normal AD biomarkers (A-/T-/N-) and ATN Posi-
tive (12 of 58; 5 SCD + 7 MCI): pathological AD biomarkers (if at least A or T or N were +).

2.3 Visuo-attentive task
163 The visuo-attentive experimental paradigm chosen is the 3-Choice Vigilance Test (3-CVT) requires identi-
164 fying a target shape (upward triangle) among two distractor shapes (downward triangle and diamond)
165 (29). Shapes are shown for 0.2 seconds with varied interstimulus intervals in the 20-minute task. Partici-
166 pants press left for targets (70%) and right for distractors (30%). Performance is evaluated using reac-
167 tion time, accuracy, and F-Measure (50).

2.4 EEG devices
169 EEG data were collected from eligible subjects at IRCCS Don Gnocchi (Florence, Italy) using the 64-
170 channel Galileo-NT system (E.B. Neuro S.p.a.). Sensor placement followed the extended 10/20 system
171 (51). Signals were recorded unipolarly at 512 Hz. Electrode impedances were maintained between 7 and
172 10 KOhm; if exceeded, electrodes were readjusted, and affected segments were removed.

2.5 EEG preprocessing and ERP component definition
EEG processing included band-pass filtering (1-45 Hz), noisy channel interpolation, average re-
referencing, and artefactual component exclusion via ICA (52). Trials lasted 1000 ms, with 200 ms for
stimulus presentation and 800 ms for response (participants had in average 421.32 trials (std=77.57)).
ERPs were epoch-aligned with correct responses to the target stimulus, segmented from 0 to 1000 ms
with a -100 ms baseline. Average EEG signals from occipital (PO7, PO8, O1, Oz, O2) and central channels
(FC1, FCz, FC2, C1, Cz, C2) were computed for encoding and decision-making analysis, respectively. In the
stimulus encoding phase, particular interest was shown in the P1, N1 and P2 components, while in the
decision phase, the components of greatest interest were P300, P600 and P900.

2.6 Neural features computations

We extracted neural features from defined ERP components, including voltage peaks, latencies, and integrals (Simpson method) from each channel of occipital and central scalp parcellations. The computation of the integration time window was performed once the peak of the canonical components was identified, with the width of the window proportional to the extent of the component. We also introduced an occipital-seed based functional connectivity metrics with the aim to evaluate the amount of similarity between seed channels and other channels of the scalp. The similarity measure adopted is the Spearman rank-order correlation coefficient (53–55) computed within encoding (0-200ms) timeframe. We then counted the channels that had correlations with (p < 0.05) and (r > 0.90) and computed the percentage relative to the total number of channels. This measurement allowed us to assess the extent of extra-seed recruitment of channels (the more the channels are similar to the occipital seed, the more they are engaged with the occipital seed, the more are the neural sources used during encoding process). A value close to 0% indicated that no channel was highly correlated with the occipital seed, while a value greater than 0% indicated the extent of extra-seed recruitment.

2.7 Patients' descriptors

Patients underwent an extensive neuropsychological examination (see specific references in (43)), including global measurements (MMSE), attention (Trial Making Test A/B and BA, visual search), and premorbid intelligence estimation (TIB). Personality traits (Big Five Factors Questionnaire - BFFQ), and leisure activities evaluation (structured interviewed regarding participation in intellectual, sporting and social activities)

2.8 Computational notes

Non-parametric analysis was employed, with statistics presented as mean values and std (where possible). Pair and group analysis were done with Kruskal-Wallis H test. Statistical differences in ERP voltage dynamics were assessed instant-by-instant by computing H statistics on voltage values at considered groups: i.e., for each instant t and group g we first compute the matrices $\left[v_g\right]_t^{c \times n}$, where v_g is the set of voltage values of a given group g, $c \times n$ is the matrix dimension of c (number of channels) and n (number of subjects within the group g); then, we statistically compared each group matrix as an empirical distribution of voltage values. The multiple comparison correction used for ERP dynamics was the FDR Benjamini-Hochberg, while for all the other test was Bonferroni. Data preprocessing utilized EEGLAB (52), while postprocessing and visualization were performed using Python libraries (numpy, scipy, pandas). Scripts of pre and postprocessing are available at https://github.com/albertoarturovergani/PREVIEWTCVT. Due to the sensitivity of personal information in this research, the data that support the findings of this study are available from the corresponding author, upon reasonable request.

2.9 Outliers management

All 178 participants performed the 3-CVT task. Poor behavioural performance was not grounds for exclusion, as we aimed to reflect the variability encountered in real-world applications. Instead, we applied an exclusion criterion based on neural features extracted from ERP dynamics: participants who exceeded the 3-sigma cut-off (56) for at least two neural features were excluded. This led to the exclusion of 2 MCI and 4 SCD patients.

2.10 Visual summary

A summary of the results and methodologies implemented can be seen in Figure 1. Cognitive decline followed a monotonically decreasing course with age and the speed of the loss of mental faculties depended on the presence of a cognitive pathology such as SCD, MCI or severe forms of dementia such as AD (Fig. 1A). Task performance also followed a monotonically decreasing course in line with the severity of the clinical condition (Fig. 1B). Simultaneous recording of the EEG signal during task performance (Fig. 1C) captured the dynamics of occipital and central ERP components distinct for clinical conditions (Fig. 1D). The components distinguished significantly for their dynamics across clinical conditions were the occipital P1 and N1 components and the central components (P300, P600, P900). Feature extraction from the identified components revealed a non-monotonic trend in their values, showing MCIs to have a greater recruitment of neural resources (Fig. 1E).

Figure 1. Visual abstract. (A) Monotonic decreasing relationship between age and cognitive decline, modu-
lated in its course by the presence of SCD or MCI pathology (or eventually AD). Dashed line traces the patho-
logical deviation from healthy path line. (B) Performance on the visuo-attentive task 3-CVT following a mon-
otonic decreasing course according to the severity of the pathology. (C) Experimental EEG signal recording
setting while participants were performing the 3-CVT task. (D) ERP dynamics extracted from occipital and
central channels separated by clinical condition, where occipital channels probed the encoding phase of the
stimulus, and the central channels probed the decision-making phase of the stimulus. (E) Non-monotonic (V-
shape) trend of ERP features reflecting increased recruitment of neural resources by MCI patients. Dashed
line traces the hypothetical monotonic trend which has been altered by the features values.

3. Results

3.1 Participant profiling

The multidimensional profile indicated that individuals with SCD and MCI differed significantly on several
clinical, psychological and behavioural scales (see group statistics in Table 1). Patients with MCI had a
higher age, lower educational level and lower performance on cognitive and visuomotor assessments
than subjects with SCD and HS. Subjects with SCD demonstrated higher levels of intellectual and social
activity than those with MCI and were more likely to have a family history of Alzheimer's disease. HS sub-
jects, on the other hand, were only examined on a few scales, showed the highest scores on cognitive tests
(e.g. MMSE), were younger and had the highest levels of education compared to both SCD and MCI groups.
Overall, the HS subjects showed better cognitive functioning and better education levels, highlighting
their status as a reference or control group in this study.

 Table 1. Demographic, clinical characteristics and task performance of study participants.

Class	Scale	HS (n=19)	SCD (n=119)	MCI (n=40)	p(H)
Task	F-Measure [a.u.]	0.91 (0.03)	0.900 (0.05)	0.869 (0.05)	*0.02 (14.22)
Task	Accuracy [%]	95.37 (3.22)	94.235 (4.84)	90.025 (7.62)	*0.01 (14.62)
Task	Reaction time [s]	0.45 (0.08)	0.464 (0.09)	0.485 (0.09)	1.00 (3.01)
Clinical	MMSE [a.u.]	29.15 (1.09)	27.514 (2.14)	26.392 (2.40)	**0.00 (18.73)
Clinical	Education [years]	14.95 (3.32)	13.412 (3.65)	10.325 (3.88)	***0.00 (23.73)
Clinical	Age [years]	62.53 (5.08)	65.403 (9.23)	72.750 (8.37)	***0.00 (23.79)
Clinical	TIB [a.u]	- (-)	113.326 (3.60)	110.618 (6.38)	0.65 (4.66)
Clinical	Age at onset [years]	- (-)	56.983 (9.03)	63.300 (10.06)	**0.01 (12.85)
Clinical	Family history of AD [count]	- (-)	74 (-)	25 (-)	0.15 (12.85)
Visuo-attentive	Trail Making Test B-A [a.u.]	- (-)	33.353 (35.47)	115.213 (177.39)	*0.02 (10.69)
Visuo-attentive	Trail Making Test A [a.u.]	- (-)	29.624 (14.37)	40.475 (39.89)	1.00 (2.67)
Visuo-attentive	Trail Making Test A [a.u.]	- (-)	63.084 (43.91)	155.725 (190.79)	*0.03 (10.37)
Visuo-attentive	Attentive matrices [a.u.]	- (-)	51.002 (6.45)	47.812 (7.93)	0.19 (6.85)
Leasure time	Intellectual activity [a.u.]	- (-)	18.866 (3.38)	16.994 (5.32)	0.23 (6.51)
Leasure time	Social activity [a.u.]	- (-)	9.039 (2.80)	8.082 (2.69)	1.00 (3.92)
Leasure time	Sport activity [a.u.]	- (-)	6.534 (2.48)	6.027 (2.27)	1.00 (0.54)
Psychological	Extraversion [a.u.]	- (-)	45.756 (5.51)	44.375 (5.97)	1.00 (3.78)
Psychological	Agreableness [a.u.]	- (-)	50.832 (7.59)	48.400 (6.38)	0.07 (8.63)
Psychological	Coscientiousness [a.u.]	- (-)	49.887 (7.67)	49.000 (7.01)	1.00 (0.44)
Psychological	Emotive stability [a.u.]	- (-)	50.202 (7.08)	49.250 (6.91)	1.00 (2.01)
Psychological	Openness of mind [a.u.]	- (-)	46.109 (6.25)	42.725 (6.07)	0.10 (7.95)

Note. The table is organized into scales divided by category: clinical, leisure time, psychological (Big Five), visual-attentive, and task performance. The values represent the mean and standard deviation (SD) for each scale across three groups: Subjective Cognitive Decline (SCD, N=119), Mild Cognitive Impairment (MCI, N=40), and Healthy Subjects (HS, N=19). Pairwise comparisons use the Kruskal-Wallis test, with p-values (p(H)) associated with the H-statistic. Significant p-values are marked with asterisks: * (p < 0.05), ** (p < 0.01), *** (p < 0.001), **** (p < 0.0001).

3.2 Task performance follows the severity of cognitive decline

The participants showed behaviour in line with the severity of their condition (Fig. 2). Although reaction time showed no significant differences between conditions (H=3.01, p=1.00; Tab1), there was an observable trend towards slower responses (Fig. 2A). Accuracy differed significantly between group conditions (H=14.62, p=0.01; Tab1), as well as for the F-Measure (H=14.22, p=0.02; Tab1). The post-hoc analyses (Fig. 2B) showed that significant differences were attributable to SCD and MCI (H=12.63 p=0.001) and between HS and MCI (H=8.06, p=0.0135), whereas they were not significant between HS and SCD (H=0.81, p=1.00). Post-hoc F-Measure analysis (Fig. 2C) showed a significant difference between SCD and MCI (H=10.86, p=0.0023) and between HS and MCI (H=9.87, p=0.00503), but not between HS and SCD (H=1.563, p=0.6337). The dichotomy of the F-measure based on the median resulted in two distinct cate-

gories: low performance and high performance (Fig. 2C), highlighting that there were MCI and SCD patients capable of achieving performance equal to healthy subjects.

Figure 2. Behavioural outcomes across conditions. (A) Reaction time [s]. (B) Accuracy [%]. (C) F-measure [a.u.]. To the right of (C) is the Cumulative Density Function of F-measure dichotomised into 'low' and 'high' levels with reference to the median of the aggregated conditions. Pairwise statistics based on Kruskal-Wallis H test with p-value corrected by Bonferroni's method (alpha=0.05). P-value annotation legend: ns: 0.05 < p <= 1, *: 0.01 < p <= 0.05, **: 0.001 < p <= 0.01, ***: 0.0001 < p <= 0.001. Colour code: HS (blue), SCD (orange), MCI (red), Low-performance subjects (green) and High-performance subjects (cyan).

3.3 Clinical conditions have specific ERP dynamics

The study of ERP dynamics in the occipital and central channels revealed condition-specific potential patterns. In the occipital channels, statistically significant temporal sequences ($p<0.05$) were observed in the amplitudes between clinical conditions (Fig. 3A), particularly for the P1 (54-82 [ms]; HS>SCD for 57.14% of time and MCI>SCD for 100%) and N1 (93-195 [ms]; HS<MCI for 75.51% of time, HS<SCD for 100% of time, MCI<SCD for 59.18% of time) potentials. Interestingly, the temporal dynamics of the amplitudes of the P1 and N1 profiles are not monotonically modulated by condition severity. In fact, a greater attenuation of the amplitude profiles for both P1 and N1 is evident in patients with SCD than in MCI and HS. In the central channels, significant differences in the instantaneous amplitudes were observed between clinical conditions (Fig. 3B) for the P300/600/900 components ($p<0.05$; >335 [ms]; HS>MCI for 64.41% of time, HS>SCD for 90.59% of time, MCI≠SCD for 41.47% of time), which were overall attenuated in patients compared to HS, but MCI, particularly for P600 showed an amplified gain compared to SCD.

Figure 3. ERP dynamics across conditions. (A) . ERP computed as average of signals in the cluster of occipital channels (PO7, PO8, O1, Oz, O2) representative of the encoding phase of the stimulus. (B) ERP computed as average of signals in the cluster of central channels (FC1, FCz, FC2, C1, Cz, C2) representative of the decision-making phase regarding the stimulus. Both panels (A) and (B): Bold representation is the overall mean within each group and shading is the standard deviation. The measures on top are the instantaneous H-statistic of the Kruskal-Wallis test and the associated p-value corrected by Bonferroni's method (alpha<0.05). Temporal instants associated with a p<0.05 are highlighted with a vertical violet bar. P1/N1/P2 and P300/600/900 labels stand for the name of the canonical event-related potentials relative to the encoding phase and decision-making phase, respectively. Colour code: HS (blue), SCD (orange), MCI (red).

Table 2. ERP features (latency, peak, integral) extracted from ERP dynamics across group conditions.

Type	ERP	SCD (N=115)	MCI (N=38)	HS (N=19)	SCD vs MCI p(H)	SCD vs HS p(H)	MCI vs HS p(H)	Age p(r)
Lat	P1	73.57 (21.95)	73.79 (15.08)	70.02 (16.78)	1 (0.44)	* 0.0486 (5.78)	0.167 (3.66)	1 (0.04)
Lat	P2	203.11 (28.76)	208.94 (28.81)	216.96 (27.75)	0.0617 (5.36)	*** 0.0001 (17.51)	0.1043 (4.46)	**0.007 (0.23)
Lat	N1	123.02 (37.47)	129.79 (25.25)	128.33 (15.29)	0.0891 (4.73)	* 0.0455 (5.9)	1 (0.34)	**0.0019 (0.26)
Lat	P300	410.1 (50.35)	421.41 (55.33)	418.38 (48.41)	** 0.0034 (10.58)	0.3485 (2.47)	1 (0.78)	0.441 (0.11)
Lat	P600	605.45 (84.83)	610.14 (81.84)	599.35 (85.84)	1 (0.8)	1 (0.46)	0.5388 (1.8)	0.2889 (0.13)
Lat	P900	882.86 (64.27)	860.4 (64.57)	878.17 (68.41)	**** 0.0 (22.87)	1 (0.73)	0.0574 (5.49)	0.5076 (-0.11)
Peak	P1	2.46 (2.08)	2.61 (1.45)	2.66 (2.06)	** 0.0094 (8.73)	1 (0.85)	1 (0.87)	1 (-0.04)
Peak	P2	2.47 (2.46)	2.38 (3.02)	2.18 (2.6)	1 (0.92)	1 (0.57)	1 (0.03)	**0.0066 (-0.23)
Peak	N1	-4.54 (3.29)	-5.2 (2.45)	-7.56 (4.29)	*** 0.0006 (13.7)	**** 0.0 (43.85)	*** 0.0001 (17.4)	*0.0273 (-0.2)
Peak	P300	1.96 (1.46)	1.53 (1.45)	2.12 (1.48)	*** 0.0009 (13.04)	0.5702 (1.72)	*** 0.001 (12.84)	*0.0216 (-0.2)
Peak	P600	1.53 (1.18)	1.72 (1.23)	1.92 (0.96)	* 0.0117 (8.34)	*** 0.0002 (15.75)	0.6603 (1.5)	1 (-0.02)
Peak	P900	0.94 (0.97)	1.04 (0.92)	1.46 (0.99)	0.1908 (3.44)	**** 0.0 (31.96)	*** 0.001 (12.79)	0.9337 (0.08)
Int	P1	155.24 (103.99)	159.13 (73.05)	183.7 (115.81)	0.1013 (4.51)	0.0686 (5.18)	1 (0.76)	1 (0.03)
Int	P2	732.62 (333.01)	826.6 (321.07)	976.17 (424.23)	** 0.0013 (12.36)	**** 0.0 (30.87)	* 0.0116 (8.34)	1 (0.05)
Int	N1	402.47 (214.06)	462.44 (203.13)	561.25 (251.41)	*** 0.0005 (14.33)	**** 0.0 (33.34)	** 0.0065 (9.39)	0.3366 (0.12)
Int	P300	278.46 (175.54)	278.18 (159.95)	333.91 (170.8)	1 (0.17)	*** 0.0009 (13.06)	** 0.0051 (9.85)	0.7511 (-0.09)
Int	P600	254.53 (177.68)	285.65 (168.7)	316.67 (168.95)	* 0.019 (7.45)	*** 0.0002 (16.02)	0.3085 (2.66)	1 (0.01)
Int	P900	150.38 (140.86)	181.09 (136.33)	206.24 (146.99)	*** 0.0005 (14.09)	*** 0.0001 (16.93)	0.4285 (2.15)	0.1853 (0.14)

Note. The table presents the mean and standard deviation (SD) for various Event-Related Potential (ERP) components measured in latency (Lat [ms]), peak amplitude (Peak [µV]), and integral amplitude (Int [µV×mS]) across three groups: Subjective Cognitive Decline (SCD, N=115), Mild Cognitive Impairment (MCI, N=38), and Healthy Subjects (HS, N=19). Pairwise comparisons between groups (SCD vs. MCI, SCD vs. HS, MCI vs. HS) are shown with p-values (p(H)) from the Kruskal-Wallis test. Features correlations with age are indicated by p-values (p(r)) from the Spearman correlation test. Significant p-values are marked with asterisks: * (p < 0.05), ** (p < 0.01), *** (p < 0.001), **** (p < 0.0001).

Analysis of ERP features (peak, latency, and integral) revealed significant differences (p<0.05) between the group conditions (statistics detailed in Table 2). The observed latency differences included: P1 (SCD>HS), P2 (SCD < HS), N1 (SCD < HS), P300 (SCD < MCI), and P900 (SCD > MCI). Peak differences were noted as follows: P1 (SCD < MCI), N1 (HS < MCI < SCD), P300 (SCD < MCI; MCI < HS), P600 (SCD < MCI; SCD < HS), and P900 (SCD < HS; MCI < HS). Integrals differed between clinical conditions for P1 (SCD < HS), P2 (SCD < HS; MCI < HS), N1 (HS < MCI < SCD), P300 (SCD < HS; MCI < HS), P600 (SCD < MCI; SCD < HS), and P900 (SCD < MCI; SCD < HS). Interestingly, these characteristics mirrored the observed ERP dynamics, suggesting a non-monotonic (V-shaped) trend across participant conditions, indicating that MCI participants exhibited similar feature values to HS participants. Moreover, we found correlated with age latency of P2 and N1 and the peak of P300, while anticorrelated with peak of P2, N1, and P300. Instead, correlation study between latencies and peaks (S-Fig. 1) showed a significant association for patients, specifically for P1 (SCD and MCI), P2 (MCI), N1 (SCD), and P300.

3.4 ERP features follow a non-monotonic ordering showing attenuation for SCD and amplification for MCI

Previous analyses have suggested that electrophysiological correlates of clinical conditions might follow a non-monotonic pattern, i.e., ERP features do not follow a structure of the observational values ordered with the severity of the cognitive pathology. To study the specific contribution of the feature type (latency, peak, integral) on the non-monotonic ordering, we aggregated and normalized the feature values of a given type with respect to the HS values of the participants, indicating them as percentage deviations (Fig. 6). The normalized latency showed no significant difference between clinical conditions (Fig. 6A) (HSvsSCD, H=1.42 p=0.069; SCDvsMCI, H=1.74, p=0.83; HSvsMCI, H=0.089, p=1.00). The normalized peak showed a significant difference between HS and SCD (Fig. 6B) (HSvsSCD, H=8.55, p=0.010), but not between other pairs (HSvsMCI, H=4.43, p=0.10; SCDvsMCI, H=0.69, p=1.000). The normalized integral showed significant differences between the clinical conditions (HSvsSCD, H=2.60, p=<0.0001; HSvsMCI, H=7.446, p=0.019; SCDvsMCI, H=9.226, p=0.0071), therefore the values follow a statistical non-monotonic ordering (HS>SCD<MCI)(Fig. 6C). We than selected the normalized integral to probe the robustness across features of the non-monotonicity trend.

*Figure 6. Aggregated feature types normalized by HS values. The feature types (latency, peak, integral) aggregated the canonical potentials (P1/N1/P2/P300/P600/P900) and are normalized by HS mean (0% variations means the values are closed to HS average value; if % is >0/<0 means positive/negative percentage deviation from HS mean). Each panel showed bar and line plots to highlight mono/non-monotonic trend. (A) ERP latency. (B) ERP peak. (C) ERP integral. Pairwise statistics based on Kruskal H test with p-value corrected by Bonferroni's method (alpha=0.05). P-value annotation legend: ns: 0.05 < p <= 1, *: 0.01 < p <= 0.05, **: 0.001 < p <= 0.01, ***: 0.0001 < p <= 0.001. Colour code: SCD (orange), MCI (red).*

An overall mean decrease in feature values of 22.29% was observed in patients with SCD compared to HS (Fig. 7A). When looking at specific features, the largest significant percentage deviations (p<0.05) between SCD and HS concerned all integrals of P1/N1/P2/P300/P600/P900. In patients with MCI, a mean

percentage deviation of -14.35% was observed from HS (Fig. 7B). When looking at specific features, the
largest significant percentage deviations (p<0.05) from MCI to HS concerned N1 and P300.

Moreover, we normalized the integral of SCD as percentage of deviation from MCI, to investigate the dif-
ference across patients (Fig. 7C). The overall attenuation in percentage terms was -10.98% for the SCDs
compared to the MCI, showing significant statistics particularly for the P1, N1, P2, P600, and P900 com-
ponents. The Spearman association analysis of the characteristics with age showed that P900 was weakly
statistically linked with age (r=0.14, p=0.061), while the other normalized integral features were not as-
sociated with age: N1 (r=0.12, p=0.11), P1 (r=0.02, p=0.72), P2 (r=0.04, p=0.52), P300 (r=-0.08, p=0.25)
and P600 (r=0.014, p=0.84).

Figure 7. Radar plots of normalized integral features. The graphs show the percentage deviations of each
ERP integral feature from the HS case of the SCD (A) and MCI (B) conditions, and the percentage deviation of
SCD from MCI (C). The hexagonal box on each radar plot represents the percentage limit of 0% (HS for A-B
and MCI for C), the positive or negative deviations of which show the variations from that reference. The
numbers on the corner points of the polygon indicate the deviation in percentage terms for each integral
characteristic. The legend shows the average of the percentage changes of the integrals (indicated by ○).
The pairwise statistic based on the Kruskal H test with p-value corrected by Bonferroni's method (al-
*pha=0.05) is close to each percentage pair. Annotation legend P-value: ns: 0.01 < p <= 1, ~: 0.05<p<0.1, *:*
*0.01 < p <= 0.05, **: 0.001 < p <= 0.01, ***: 0.0001 < p <= 0.001. Colour code: HS (blue), SCD (orange), MCI*
(red).

3.5 Performance is modulated by distinct ERP dynamics in encoding and decision-making phases
After identifying electrophysiological correlates associated with the clinical conditions, we investigated
the ERP correlates of task performance. HS showed no instantaneous statistical differences in potential
amplitude during the encoding phase (S-Fig. 2A), whereas they did show statistical differences (p<0.05)
during the decision-making phase, highlighting an increase in P600 (507.81-728.51 [ms]) and P900
(>742.18 [ms]) gain in the case of high performers compared to low performers (S-Fig. 2B). In SCD pa-

tients, statistically significant variations (p<0.05) were observed during the encoding phase (Fig4. A) in the intervals related to P1 (35.15-56.64 [ms]) and P2 (72.26-107.42 [ms]), and during the decision phase (Fig4. B) in the intervals related to P300 and P600 (300-615.23 [ms]), and, to a lesser extent, P900 (significant spots over 835 ms). ERP-related feature extraction (statistics in Table 3) revealed significant differences (p<0.05) between low and high-performance conditions. Latencies differed for P1 (high < low), N1 (high < low), P300 (high < low), and P600 (high < low). Peak amplitudes and integrals showed differences for P300 (high > low) and P600 (high > low), as well as their integrals. Age correlated with latency of P2 and N1, but anticorrelated with peak of P2.

Figure 4. ERP dynamics across performance in SCD. (A) . ERP computed in the cluster of occipital channels (PO7, PO8, O1, Oz, O2) representative of the encoding phase of the stimulus. (B) ERP computed in cluster of central channels (FC1, FCz, FC2, C1, Cz, C2) representative of the decision-making phase regarding the stimulus. Both panels (A) and (B): Bold representation is the overall mean within each group and shading is the

399 *Table 3. ERP features (latency, peak, integral) extracted from ERP dynamics across task performance in SCD*

Type	ERP	SCD High (N=60)	SCD Low (N=55)	High vs Low p(H)	Age p(r)
Lat	P1	71.97 (23.29)	75.31 (20.24)	** 0.0018 (9.69)	1 (0.01)
Lat	P2	202.32 (29.6)	203.98 (27.79)	0.5351 (0.38)	** 0.0027 (0.31)
Lat	N1	117.45 (34.93)	129.1 (39.15)	**** 0.0 (16.9)	*** 0.0001 (0.38)
Lat	P300	406.63 (50.92)	413.89 (49.45)	** 0.0092 (6.78)	0.1618 (0.18)
Lat	P600	593.35 (79.57)	618.64 (88.35)	*** 0.0009 (11.01)	0.4813 (0.13)
Lat	P900	882.16 (63.41)	883.62 (65.2)	0.7374 (0.11)	1 (0.03)
Peak	P1	2.49 (2.28)	2.43 (1.85)	0.4064 (0.69)	0.452 (-0.13)
Peak	P2	2.57 (2.25)	2.36 (2.66)	0.0943 (2.8)	** 0.0093 (-0.27)
Peak	N1	-4.71 (3.54)	-4.37 (2.99)	0.7063 (0.14)	0.1519 (-0.18)
Peak	P300	2.23 (1.38)	1.66 (1.48)	**** 0.0 (24.93)	0.0795 (-0.21)
Peak	P600	1.66 (1.07)	1.39 (1.28)	**** 0.0 (19.66)	1 (-0.06)
Peak	P900	0.96 (0.89)	0.92 (1.05)	0.1061 (2.61)	0.6103 (0.12)
Int	P1	163.29 (111.0)	146.46 (94.99)	0.2274 (1.46)	0.8254 (-0.1)
Int	P2	726.72 (315.02)	739.04 (351.48)	0.8905 (0.02)	1 (-0.01)
Int	N1	404.16 (210.76)	400.62 (217.59)	0.8616 (0.03)	1 (0.06)
Int	P300	296.39 (173.74)	258.89 (175.41)	*** 0.0007 (11.46)	1 (-0.08)
Int	P600	271.01 (156.36)	236.56 (196.77)	**** 0.0 (17.06)	1 (0.0)
Int	P900	153.12 (126.79)	147.4 (154.69)	0.2821 (1.16)	0.3802 (0.14)

Note. The table presents the mean and standard deviation (SD) for various Event-Related Potential (ERP)
components measured in latency (Lat [ms]), peak amplitude (Peak [μV]), and integral amplitude (Int
[μV×mS]) across two SCD groups: low performance (N=59) and high performance (N=55). Pairwise compar-
isons between groups are shown with p-values (p(H)) from the Kruskal-Wallis test. Features correlations
with age are indicated by p-values (p(r)) from the Spearman correlation test. Significant p-values are
*marked with asterisks: * (p < 0.05), ** (p < 0.01), *** (p < 0.001), **** (p < 0.0001).*

The study of electrophysiological correlates of performance in MCI revealed significant differences
(p<0.05) during both the encoding and decision phases. In the encoding phase (Fig. 5A), significant time
instants were associated with P1 (44.92-56.64 ms), N1 (115.23-126.95 ms), and P2 (218.75-316.40 [ms]).
In the decision phase (Fig. 5B), significant time instants were observed for P300 (300-408 ms), P600
(423.82-541.01 [ms]), and spots of P900 (957.03-978.51 [ms]). Feature extraction of peak amplitude,
peak latency, and integral (statistics detailed in Table 4) showed significant differences (p<0.05) for P2
latencies (high < low), N1 peaks (high > low), and peaks of P300 and P600 (both high > low). The inte-
grals showed statistical differences for P1 (high < low), N1 (high < low), and P300, P600, and P900 (all
high > low. Age correlated only with the integral of P1.

Figure 5. ERP dynamics across performance in SCD. (A) . ERP computed in the cluster of occipital channels
(PO7, PO8, O1, Oz, O2) representative of the encoding phase of the stimulus. (B) ERP computed in cluster of
central channels (FC1, FCz, FC2, C1, Cz, C2) representative of the decision-making phase regarding the stimu-
lus. Both panels (A) and (B): Bold representation is the overall mean within each group and shading is the
standard deviation. The measures on top are the instantaneous H-statistic of the Kruskal-Wallis test and the
associated p-value corrected by Bonferroni's method (alpha<0.05). Temporal instants associated with a
p<0.05 are highlighted with a vertical violet bar. P1/N1/P2 and P300/600/900 labels stand for the name of
the canonical event-related potentials relative to the encoding phase and decision-making phase respective-
ly. Colour code: low performance subjects (green), high performance subjects (cyan).

 Table 4. ERP features (latency, peak, integral) extracted from ERP dynamics across task performance in MCI

Type	ERP	MCI Low (N=26)	MCI High (N=12)	Low vs High p(H)	Age p(r)
Lat	P1	73.48 (15.15)	74.45 (14.9)	0.8375 (0.04)	1 (0.1)
Lat	P2	213.1 (29.63)	199.93 (24.66)	** 0.0039 (8.33)	1 (0.13)
Lat	N1	127.48 (22.75)	134.8 (29.35)	0.1094 (2.56)	1 (-0.04)
Lat	P300	420.3 (52.29)	423.83 (61.33)	0.6527 (0.2)	1 (-0.1)
Lat	P600	612.66 (86.68)	604.68 (69.92)	0.5731 (0.32)	1 (0.12)
Lat	P900	861.48 (68.1)	858.07 (56.08)	0.9801 (0.0)	1 (-0.16)
Peak	P1	2.71 (1.62)	2.42 (0.95)	0.3164 (1.0)	0.3609 (0.26)
Peak	P2	2.64 (3.09)	1.82 (2.79)	0.1804 (1.79)	1 (0.02)
Peak	N1	-5.49 (2.42)	-4.57 (2.38)	** 0.0093 (6.76)	0.0925 (-0.35)
Peak	P300	1.38 (1.45)	1.84 (1.41)	** 0.0043 (8.17)	1 (-0.13)
Peak	P600	1.62 (1.22)	1.95 (1.23)	* 0.0261 (4.95)	1 (0.03)
Peak	P900	1.04 (0.95)	1.04 (0.87)	0.5937 (0.28)	1 (0.1)
Int	P1	171.52 (78.62)	132.28 (49.51)	*** 0.001 (10.86)	* 0.0336 (0.41)
Int	P2	853.19 (335.23)	768.97 (279.46)	0.1146 (2.49)	0.6592 (0.2)
Int	N1	483.26 (206.22)	417.32 (188.54)	* 0.0162 (5.78)	0.2671 (0.28)
Int	P300	260.38 (155.77)	316.74 (162.1)	** 0.0048 (7.97)	1 (-0.08)
Int	P600	269.14 (168.77)	321.45 (162.92)	* 0.0173 (5.67)	1 (0.0)
Int	P900	175.38 (145.16)	193.45 (113.93)	* 0.0489 (3.88)	1 (0.06)

*Note. The table presents the mean and standard deviation (SD) for various Event-Related Potential (ERP) components measured in latency (Lat [ms]), peak amplitude (Peak [μV]), and integral amplitude (Int [μV×mS]) across two MCI groups: low performance (N=26) and high performance (N=12). Pairwise comparisons between groups are shown with p-values (p(H)) from the Kruskal-Wallis test. Features correlations with age are indicated by p-values (p(r)) from the Spearman correlation test. Significant p-values are marked with asterisks: * (p < 0.05), ** (p < 0.01), *** (p < 0.001), **** (p < 0.0001).*

3.6 ATN classification is associated with a specific ERP dynamic in SCD and MCI patients

After studying the ERP dynamics between clinical conditions and performance levels, we investigated the correlates of ATN classification. Our results showed that, aggregating SCD and MCI patients undergoing CSF examination (N=58), no significant differences were observed between positive and negative ATN status along the temporal dynamics of encoding in occipital channels and decision-making in central channels (S-Fig. 3). However, when stratifying the ATN classification according to clinical condition, significant results emerged for both patients with SCD (N=34) and MCI (N=24). In SCDs (S-Fig. 4) a significant time window was found corresponding to N1 (123-144 [ms]), which represented an anticipatory time offset in case of a positive ATN. Stratifying the ATN by MCI condition (see S-Fig. 5), no significant differences were observed within the encoding window. However, differences were noted during the decision phase, in the attenuation of the amplitude peaks of P600 (476-591 [ms]) and P900 (750-945 [ms], and spots over >986[ms]) in the case of ATN positive MCI.

3.7 Encoding FC reflects the non-monotonic ordering of ERP features and

The observation that MCI patients showed a higher recruitment of neural resources prompted us to investigate the topographic state of the scalp, with particular attention to the visual encoding phase. To this end, the use of the Encoding FC (see Methods) highlighted the non-monotonic ordering of activation between clinical conditions, showing higher values of extra-occipital channel recruitment for HS (11.05%) and MCI (11.84%), compared to SCD (7.50%) (Fig6A). A significant difference was observed between SCD and MCI (H=11.86, p=0.0017), whereas not between HS and SCD (H=4.563, p=0.098) and between HS and MCI (H=2.028, p=1.000). Therefore, the similarity of the features values between HS and MCI is supported by the increased general recruitment of the scalp in relation to the occipital seed. This similarity in the increase in recruitment between HS and MCI appears to be specific to low-performing HS (H=6.70, p=0.028;

Fig. 6A), as encoding FC above the overall median, although this effect of performance on FC is less significant on SCD (H=0.13, p=1.000) and MCI (H=0.070, p=1.000). Moreover, stratifying the patients with respect to ATN classification, we found no significant differences of Encoding FC across biological severity between SCD (H=0.0654, p=1.000) and MCI (H=0.0053, p=1.000) (Fig. 6B). The association study according to Spearman's metric showed a significant correlation of Encoding FC with age (r=0.23, p=0.001) and anti-correlation with education (r=-0.15, p=0.04).

Figure 6. Encoding FC computed across clinical conditions and ATN classification. (A) Boxplot of clinical conditions. (B) Boxplot of clinical conditions stratified by ATN status. Topographic maps left to (A) are indicative of the scalp potentials spread during the encoding process (0-200ms) as the FC is over/under the FC median. Colormaps are in voltage range -1/+1 [μV]. Higher FC means higher similarity between the occipital seed and other channels, indicative of increased extra-occipital scalp recruitment. Pairwise comparisons between groups are shown with p-values from the Kruskal-Wallis test. Significant p-values are marked with asterisks: * (p < 0.05), ** (p < 0.01), *** (p < 0.001), **** (p < 0.0001).

4. Discussion

The study analysed electrophysiological correlates of sustained visual attention in SCD and MCI. We identified condition-specific anomalies in event-related potentials (ERPs) during visual encoding (P1/N1/P2) and decision-making (P300/P600/P900). Individuals with SCD displayed reduced ERP dynamics compared to HS, whereas those with MCI exhibited heightened dynamics, except for the P300, which corresponded with clinical severity. By looking the ERP features, in particular peaks, integrals and normalized integrals, the trend of the features values followed a non-monotonic order, indicating higher values for MCI if compared with SCD. Task performance correlated with encoding anomalies across conditions and decision-making potential gain, revealing an amplification of the P300 and P600 potentials in both SCD and MCI individuals.

4.1 Visuo-attentive impairment along the cognitive decline

These findings support the hypothesis that visual sensory abnormalities characterize SCD and MCI patients to varying degrees (42). For example, occipital P1 and N1 potentials are thought to represent aspects of visual-attentive processes, including their cost (P1) and benefit (N1) (57–59). Open hypotheses suggest that P1 and N1 may not solely originate from the primary visual cortex, with N1 potentially linked to occipital-parietal/temporal/frontal generators (60), whereas P1 from extra V1 regions (V2,V3, dorsal V4) (61,62). Therefore, the recorded abnormalities in early visual components between SCD and MCI may indicate a broader impairment of the early attentional mechanism in visual processing (63).

Moreover, the increase in P1 gain observed in the case of MCI is consistent with the observation of other
colleagues, mainly, however, oriented towards auditory stimuli (64–66). What we added in our study is
that SCD patients showed a dramatic reduction in P1 compared to controls. This fact indicated that not
only P1 gain is relevant for the transition from SCD to MCI, but also its attenuation in the case of the tran-
sition between HS and SCD. The characteristic of observing an attenuation of ERP components in SCDs,
compared to HSs, and a gain in MCIs compared to SCDs, was also valid for other early visual components,
such as N1 and P2. Thus, the candidate marker of the evolution of cognitive decline along the predemen-
tia stages must consider the asymmetrical dynamics between SCDs and MCIs, where the former show
mainly an attenuation of potential compared to HSs and the latter a gain compared to SCDs.

Patients showed cognitive decline in task performance. During the encoding phase, high and low perfor-
mance reflected the anomalies observed among the clinical conditions. During the decision phase, howev-
er, more regularity is also present between conditions, highlighting that higher performance correlates
with increased central scalp recruitment, mainly P300 and P600. These two components are ERP known
in the literature as a correlate of decision quality (38,67). Similar dynamics, but in the case of late positive
potential (LPP) was found by Waninger et al which detected performance correlations with LPP recorded
on parietal channels, but during a working load visual memory test that is the Standardized Image Recog-
nition test (SIR) (25).

In general, across patients, P300, P600 and P900 were observed in our study to be attenuated in gain or
latencies in the case of patients compared to HS. This results are in line with previous observations pf
P300 dynamics in relation to cognitive decline (30–32), or, regarding P600, which has been observed
that its abnormalities are associated with increased risk of conversion to dementia in MCI patients (36–
38). P900, on the other hand, is a component recently discovered by Rosenfeld et al (68) that could poten-
tially serve as an index of the use of countermeasures and associated to P300 (69). To our knowledge,
modulation of the P900 component as a function of the severity of cognitive pathology has not been re-
ported before, but it has been observed in the sleep (70,71) and in active problem solving. (72)

The reduction in P300 amplitudes associated with cognitive decline has been topographically linked to
sources mainly in the medial frontal cortex, right dorsolateral prefrontal cortex, right inferior parietal
lobe (73), and in general in a non-pathological condition at the temporo-parietal junction (TPJ), supple-
mentary motor cortex (SMA), anterior cingulate cortex (ACC), superior temporal gyrus (STG), insula, and
dorsolateral prefrontal cortex (74). Since these anatomically highly interconnected brain regions are part
of a network associated with sustained attention, explain why P300 gain is reduced in the case of the
most severe cognitive pathology. Interestingly, the P300 is the only component that appears to show mo-
notony with the clinical condition, i.e. its attenuation follows the severity of the clinical condition.

On the other hand, modulation of P600 appears to be associated with cognitive alterations of a linguistic
nature (75,76), and the generating sources may also involve the basal ganglia (65,66). Although most
studies on P600 are linguistic, it appears to be related to the systemic release of norepinephrine by the
locus coeruleus (77), which influences the quality of decision making, memory, and attention (78), the
cells of which are known to be profoundly reduced in the case of AD (78–83).

4.2 Cognitive reserve, Encoding FC and non-monotonic ERP trend
Cognitive reserve (CR) is recognized for its role in influencing cognitive decline, potentially shielding
against dementia symptoms despite existing brain alterations (84). SCD patients exhibited higher proxy
scores of CR compared to MCI patients, as evidenced by measures of leisure activities and clinical scales.
This suggests a potentially greater capacity for brain resilience in supporting cognitive functions among
SCD patients. The CR is related to the concept of brain efficiency (85–87), i.e., lower utilization of cortical
activity for getting performance, in fact its contrary - neural inefficiency - has been associated with low CR
(88,89). The overuse of neural resources during the performance of a task is known to increase in relation
to age, particularly the extra-engagement of prefrontal cortex, and modulated by factors influencing CR
such as IQ (90) and sport practice (91,92).

In our study, we observed that MCI patients exhibited higher Encoding FC compared to SCD patients, sug-
gesting a higher utilization of cognitive resources to perform the task. Additionally, the non-monotonicity
of the observed features aligns with the general amplification of scalp activity in MCI patients compared

to SCD patients, which may also reflect deficits in brain efficiency. Furthermore, the fact that low-performing HS have a higher Encoding FC is indicative of excessive neural resource utilization associated with low-level visual-attentive performance. Instead, across participants, we found that the use of higher Encoding FC was associated directly with age and inversely with education, and this latter is known to be linked with intelligence (93). Thus, our results are in line with studies showing that intelligence and age modulate brain efficiency in terms of adaptive control of neural resource consumption (94,95), in particular in AD and related conditions (96–98)

Hence, the non-monotonic ordering of features could be explained by the brain inefficiency that some patients have, particularly MCI patients who showed the greatest extra-occipital recruitment. In this sense, modelling neural inefficiency could add causal contributions with respect to cognitive reserve and the underlying biophysical factors (99). A cause-effect paradigm as a modelling framework (100,101) of pathological AD-type neural degeneration could explain the mechanisms underlying the observed non-monotonicity in scalp potentials, and thus explaining why the electrophysiological correlate does not follow a monotonic change in line with the continuous gradient of cognitive decline (Fig. 1 in (102)).

Of note, SCD patients, showing less scalp similarity with the occipital seed compared to other channels, reflected a reduced dipole effect on the scalp, whereas HS and MCIs exhibited a dipole topography characterized by occipital negativity and frontal positivity when using higher neural resources. This occipito-frontal dipole pattern alterations resembles EEG microstate classes C and D (103), which have been associated with AD and non-AD conditions in recent research (104).

4.3 Strengths and weakness

Strengths include large sample size, multimodal data (EEG and patient descriptors), and inclusion of CSF markers in a subset. Weaknesses: limited robustness of CSF markers' statistical significance, low healthy subject number (focused on SCD vs. MCI), monocentric study without follow-ups (ongoing in PREVIEW study).

4.4 Future works

A direct application of the ERP neural features identified in this study is training machine learning algorithms to classify patients based on learned diagnostic categories (8,105–113). It is important to note the possible role of misclassifications, i.e., the assignment of a clinical category of worse or lesser severity. The consequence of a misclassification towards a less severe category, e.g. for an MCI to be classified as SCD, might indicate that the neural correlate of that MCI patient is more akin to that of an SCD and thus might have a less severe biological pathology or greater protection underlying it. Conversely, an SCD classified as MCI might indicate a neural correlate akin to a greater neurophysiological and biological alteration, which could be a risk factor for cognitive decline. Therefore, in the approach to machine learning of features extracted from neural correlates it will be important to evaluate the degree of technical error of the algorithm (false misclassification) from actual misclassifications that could give fundamental indications for the patient's clinical course.

4.5 Conclusion

Current cognitive decline biomarkers (114), such as PET neuroimaging (115) or CSF biomarkers (116), are costly, invasive, and impractical for large-scale use. Our study aims to overcome these limitations by exploring features obtainable through clinical assessments, neuropsychological evaluations, and non-invasive methods like EEG and blood tests (117). Validating multiple neural features from EEG (114,118,119) is crucial to establish their preventive and diagnostic potential for Alzheimer's disease and, in general, along the cognitive decline continuum, in order to improve and support the clinical decision-making (120).

Acknowledgements

We thank Ahmet Kaymak, Niccolo Meneghetti, Fabio Taddeini, Giovanni Vecchiato, Giacomo Privato, Salvatore Falciglia, Michela Rocchetti and Greta Carnevali for their relevant comments which have improved the quality of the research work.

Data availability
Scripts of pre and postprocessing are available at
https://github.com/albertoarturovergani/PREVIEWTCVT. Due to the sensitivity of personal information
in this research, the data that support the findings of this study are available from the corresponding au-
thor, upon reasonable request.

Conflicts
The authors declared no conflict of interest.

Funding Sources
This project was funded by Tuscany Region - PRedicting the EVolution of SubjectIvE Cognitive Decline to
Alzheimer's Disease With machine learning - PREVIEW - CUP. D18D20001300002 and by Project funded
under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.3 - Call for
tender No. 341 of 15/03/2022 of Italian Ministry of University and Research funded by the European Un-
ion – NextGenerationEU Award Number: Project code PE0000006, Concession Decree No. 1553 of
11/10/2022 adopted by the Italian Ministry of University and Research, CUP D93C22000930002, "A mul-
tiscale integrated approach to the study of the nervous system in healt and disease" (MNESYS).

Bibliography
1. Livingston G, Huntley J, Sommerlad A, Ames D, Ballard C, Banerjee S, et al. Dementia prevention,
   intervention, and care: 2020 report of the Lancet Commission. The Lancet. 2020 Aug
   8;396(10248):413–46.

2. Jessen F, Amariglio RE, van Boxtel M, Breteler M, Ceccaldi M, Chételat G, et al. A conceptual
   framework for research on subjective cognitive decline in preclinical Alzheimer's disease. Alz-
   heimers Dement. 2014 Nov;10(6):844–52.

3. Jessen F, Amariglio RE, Buckley RF, Flier WM van der, Han Y, Molinuevo JL, et al. The characterisa-
   tion of subjective cognitive decline. The Lancet Neurology. 2020 Mar 1;19(3):271–8.

4. Gauthier S, Reisberg B, Zaudig M, Petersen RC, Ritchie K, Broich K, et al. Mild cognitive impair-
   ment. The Lancet. 2006 Apr 15;367(9518):1262–70.

5. van Dyck CH, Swanson CJ, Aisen P, Bateman RJ, Chen C, Gee M, et al. Lecanemab in Early Alz-
   heimer's Disease. New England Journal of Medicine. 2023 Jan 5;388(1):9–21.

6. Budd Haeberlein S, Aisen PS, Barkhof F, Chalkias S, Chen T, Cohen S, et al. Two Randomized Phase
   3 Studies of Aducanumab in Early Alzheimer's Disease. J Prev Alzheimers Dis. 2022;9(2):197–210.

7. Guest FL, Rahmoune H, Guest PC. Early Diagnosis and Targeted Treatment Strategy for Improved
   Therapeutic Outcomes in Alzheimer's Disease. In: Guest PC, editor. Reviews on New Drug Targets
   in Age-Related Disorders [Internet]. Cham: Springer International Publishing; 2020 [cited 2022
   Nov 27]. p. 175–91. (Advances in Experimental Medicine and Biology). Available from:
   https://doi.org/10.1007/978-3-030-42667-5_8

8. Rossini PM, Miraglia F, Vecchio F. Early dementia diagnosis, MCI-to-dementia risk prediction, and
   the role of machine learning methods for feature extraction from integrated biomarkers, in par-
   ticular for EEG signal analysis. Alzheimer's & Dementia. 2022;18(12):2699–706.

9. Alexander DM, Arns MW, Paul RH, Rowe DL, Cooper N, Esser AH, et al. Eeg markers for cognitive
   decline in elderly subjects with subjective memory complaints. J Integr Neurosci. 2006
   Mar;05(01):49–74.

10.   Babiloni C, Visser PJ, Frisoni G, De Deyn PP, Bresciani L, Jelic V, et al. Cortical sources of rest-
ing EEG rhythms in mild cognitive impairment and subjective memory complaint. Neurobiology of
Aging. 2010 Oct 1;31(10):1787–98.

11.   Ganapathi AS, Glatt RM, Bookheimer TH, Popa ES, Ingemanson ML, Richards CJ, et al. Differ-
entiation of Subjective Cognitive Decline, Mild Cognitive Impairment, and Dementia Using
qEEG/ERP-Based Cognitive Testing and Volumetric MRI in an Outpatient Specialty Memory Clinic.
Journal of Alzheimer's Disease. 2022 Jan 1;90(4):1761–9.

12.   Che J, Cheng N, Jiang B, Liu Y, Liu H, Li Y, et al. Executive function measures of participants
with mild cognitive impairment: Systematic review and meta-analysis of event-related potential
studies. International Journal of Psychophysiology. 2024 Mar 1;197:112295.

13.   Wang J, Li C, Yu X, Zhao Y, Shan E, Xing Y, et al. Effect of emotional stimulus on response in-
hibition in people with mild cognitive impairment: an event-related potential study. Front Neuro-
sci [Internet]. 2024 Apr 30 [cited 2024 Jul 10];18. Available from:
https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2024.1357435/full

14.   Xiao S, Li Y, Liu M, Li Y. Electrophysiological evidence of impaired cognitive reappraisal in
amnestic mild cognitive impairment: An event-related potential study. Behavioural Brain Re-
search. 2022 Jun 3;427:113800.

15.   Paitel ER, Samii MR, Nielson KA. A systematic review of cognitive event-related potentials in
mild cognitive impairment and Alzheimer's disease. Behavioural Brain Research. 2021 Jan
1;396:112904.

16.   Smailovic U, Koenig T, Kåreholt I, Andersson T, Kramberger MG, Winblad B, et al. Quantita-
tive EEG power and synchronization correlate with Alzheimer's disease CSF biomarkers. Neurobiol
Aging. 2018 Mar;63:88–95.

17.   Cintra MTG, Ávila RT, Soares TO, Cunha LCM, Silveira KD, de Moraes EN, et al. Increased
N200 and P300 latencies in cognitively impaired elderly carrying ApoE ε-4 allele. International
Journal of Geriatric Psychiatry. 2018;33(2):e221–7.

18.   Morrison C, Rabipour S, Knoefel F, Sheppard C, Taler V. Auditory Event-related Potentials in
Mild Cognitive Impairment and Alzheimer's Disease. Current Alzheimer Research. 2018 Jul
1;15(8):702–15.

19.   Harding GFA, Wright CE, Orwin A. Primary Presenile Dementia: The Use of the Visual Evoked
Potential as a Diagnostic Indicator. The British Journal of Psychiatry. 1985 Nov;147(5):532–9.

20.   Kolev V, Yordanova J, Basar-Eroglu C, Basar E. Age effects on visual EEG responses reveal dis-
tinct frontal alpha networks. Clinical Neurophysiology. 2002 Jun 1;113(6):901–10.

21.   Karamacoska D, Barry RJ, De Blasio FM, Steiner GZ. EEG-ERP dynamics in a visual Continuous
Performance Test. International Journal of Psychophysiology. 2019 Dec 1;146:249–60.

22.   Saito H, Yamazaki H, Matsuoka H, Matsumoto K, Numachi Y, Yoshida S, et al. Visual event-
related potential in mild dementia of the Alzheimer's type. Psychiatry and Clinical Neurosciences.
2001;55(4):365–71.

23.   Chapman RM, Bragdon HR. Evoked Responses to Numerical and Non-Numerical Visual Stim-
uli while Problem Solving. Nature. 1964 Sep;203(4950):1155–7.

24.   Armstrong RA. Alzheimer's Disease and the Eye. Journal of Optometry. 2009 Jan 1;2(3):103–
11.

25.   Javitt DC, Martinez A, Sehatpour P, Beloborodova A, Habeck C, Gazes Y, et al. Disruption of
early visual processing in amyloid-positive healthy individuals and mild cognitive impairment. Alz-
heimers Res Ther. 2023 Feb 28;15(1):42.

26.   Waninger S, Berka C, Meghdadi A, Karic MS, Stevens K, Aguero C, et al. Event-related poten-
tials during sustained attention and memory tasks: Utility as biomarkers for mild cognitive im-
pairment. Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring. 2018 Jan
1;10:452–60.

27.   Krasodomska K, Lubiński W, Potemkowski A, Honczarenko K. Pattern electroretinogram
(PERG) and pattern visual evoked potential (PVEP) in the early stages of Alzheimer's disease. Doc
Ophthalmol. 2010 Oct 1;121(2):111–21.

28.   Pollock VE, Schneider LS, Chui HC, Henderson V, Zemansky M, Sloane RB. Visual evoked po-
tentials in dementia: A meta-analysis and empirical study of Alzheimer's disease patients. Biologi-
cal Psychiatry. 1989 Apr 15;25(8):1003–13.

29.   Mangun GR. Neural mechanisms of visual selective attention. Psychophysiology.
1995;32(1):4–18.

30.   Hedges D, Janis R, Mickelson S, Keith C, Bennett D, Brown BL. P300 Amplitude in Alzheimer's
Disease: A Meta-Analysis and Meta-Regression. Clin EEG Neurosci. 2016 Jan 1;47(1):48–55.

31.   Polich J, Ehlers CL, Otis S, Mandell AJ, Bloom FE. P300 latency reflects the degree of cognitive
decline in dementing illness. Electroencephalography and Clinical Neurophysiology. 1986 Feb
1;63(2):138–44.

32.   Braverman ER, Chen TJH, Schoolfield J, Martinez-Pons M, Arcuri V, Varshavskiy M, et al. De-
layed P300 latency correlates with abnormal Test of Variables of Attention (TOVA) in adults and
predicts early cognitive decline in a clinical setting. Adv Therapy. 2006 Jul 1;23(4):582–600.

33.   van Deursen JA, Vuurman EFPM, Smits LL, Verhey FRJ, Riedel WJ. Response speed, contin-
gent negative variation and P300 in Alzheimer's disease and MCI. Brain and Cognition. 2009 Apr
1;69(3):592–9.

34.   Jiang S, Qu C, Wang F, Liu Y, Qiao Z, Qiu X, et al. Using event-related potential P300 as an
electrophysiological marker for differential diagnosis and to predict the progression of mild cogni-
tive impairment: a meta-analysis. Neurol Sci. 2015 Jul 1;36(7):1105–12.

35.   Parra M, Ascencio L, Urquina H, Manes F, Ibanez A. P300 and Neuropsychological Assess-
ment in Mild Cognitive Impairment and Alzheimer Dementia. Front Neurol [Internet]. 2012 Dec 5
[cited 2024 Jul 13];3. Available from:
https://www.frontiersin.org/journals/neurology/articles/10.3389/fneur.2012.00172/full

36.   Olichney JM, Taylor JR, Gatherwright J, Salmon DP, Bressler AJ, Kutas M, et al. Patients with
MCI and N400 or P600 abnormalities are at very high risk for conversion to dementia. Neurology.
2008 May 6;70(19_part_2):1763–70.

37.   Olichney JM, Pak J, Salmon DP, Yang JC, Gahagan T, Nowacki R, et al. Abnormal P600 word
repetition effect in elderly persons with preclinical Alzheimer's disease. Cognitive Neuroscience.
2013 Dec 1;4(3–4):143–51.

38.   Dröge A, Fleischer J, Schlesewsky M, Bornkessel-Schlesewsky I. Neural mechanisms of sen-
tence comprehension based on predictive processes and decision certainty: Electrophysiological
evidence from non-canonical linearizations in a flexible word order language. Brain Research.
2016 Feb 15;1633:149–66.

39.   Paitel ER, Samii MR, Nielson KA. A systematic review of cognitive event-related potentials in
mild cognitive impairment and Alzheimer's disease. Behavioural Brain Research. 2021 Jan
1;396:112904.

40.   Kimiskidis VK, Papaliagkas VT. Event-related potentials for the diagnosis of mild cognitive
impairment and Alzheimer's disease. Expert Opinion on Medical Diagnostics. 2012 Jan 1;6(1):15–
26.

41.   Gu L, Zhang Z. Exploring Potential Electrophysiological Biomarkers in Mild Cognitive Impair-
ment: A Systematic Review and Meta-Analysis of Event-Related Potential Studies. Journal of Alz-
heimer's Disease. 2017 Jan 1;58(4):1283–92.

42.   Morrison C, Rabipour S, Taler V, Sheppard C, Knoefel F. Visual Event-Related Potentials in
Mild Cognitive Impairment and Alzheimer's Disease: A Literature Review. Current Alzheimer Re-
search. 2019 Jan 1;16(1):67–89.

43.   Mazzeo S, Lassi M, Padiglioni S, Vergani AA, Moschini V, Scarpino M, et al. PRedicting the
EVolution of SubjectIvE Cognitive Decline to Alzheimer's Disease With machine learning: the PRE-
VIEW study protocol. BMC Neurology. 2023 Aug 12;23(1):300.

44.   Jessen F, Amariglio RE, van Boxtel M, Breteler M, Ceccaldi M, Chételat G, et al. A conceptual
framework for research on subjective cognitive decline in preclinical Alzheimer's disease. Alz-
heimer's & Dementia. 2014;10(6):844–52.

45.   Albert MS, DeKosky ST, Dickson D, Dubois B, Feldman HH, Fox NC, et al. The diagnosis of
mild cognitive impairment due to Alzheimer's disease: recommendations from the National Insti-
tute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's dis-
ease. Alzheimers Dement. 2011 May;7(3):270–9.

46.   McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR, Kawas CH, et al. The diagnosis
of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging-
Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimer's
& Dementia. 2011 May 1;7(3):263–9.

47.   Alcolea D, Pegueroles J, Muñoz L, Camacho V, López-Mora D, Fernández-León A, et al.
Agreement of amyloid PET and CSF biomarkers for Alzheimer's disease on Lumipulse. Annals of
Clinical and Translational Neurology. 2019;6(9):1815–24.

48.   Giacomucci G, Mazzeo S, Bagnoli S, Casini M, Padiglioni S, Polito C, et al. Matching Clinical
Diagnosis and Amyloid Biomarkers in Alzheimer's Disease and Frontotemporal Dementia. Journal
of Personalized Medicine. 2021 Jan;11(1):47.

49.   Jack CR, Bennett DA, Blennow K, Carrillo MC, Feldman HH, Frisoni GB, et al. A/T/N: An unbi-
ased descriptive classification scheme for Alzheimer disease biomarkers. Neurology. 2016 Aug
2;87(5):539–47.

50.   Stikic M, Johnson RR, Levendowski DJ, Popovic DP, Olmstead RE, Berka C. EEG-derived esti-
mators of present and future cognitive performance. Front Hum Neurosci. 2011;5:70.

51.   Danjou P, Viardot G, Maurice D, Garcés P, Wams EJ, Phillips KG, et al. Electrophysiological
assessment methodology of sensory processing dysfunction in schizophrenia and dementia of the
Alzheimer type. Neuroscience & Biobehavioral Reviews. 2019 Feb 1;97:70–84.

52.   Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-trial EEG dynam-
ics including independent component analysis. J Neurosci Methods. 2004 Mar 15;134(1):9–21.

53.   Siems M, Pape AA, Hipp JF, Siegel M. Measuring the cortical correlation structure of sponta-
neous oscillatory activity with EEG and MEG. NeuroImage. 2016 Apr 1;129:345–55.

54.   Spearman C. The Proof and Measurement of Association Between Two Things. East Norwalk,
CT, US: Appleton-Century-Crofts; 1961. 45 p. (Studies in individual differences: The search for in-
telligence).

55.   Tan A, Hu L, Tu Y, Chen R, Hung YS, Zhang Z. N1 Magnitude of Auditory Evoked Potentials
and Spontaneous Functional Connectivity Between Bilateral Heschl's Gyrus Are Coupled at Inter-
individual Level. Brain Connectivity. 2016 Jul;6(6):496–504.

56.   Pukelsheim F. The Three Sigma Rule. The American Statistician. 1994 May 1;48(2):88–91.

57.   Luck SJ, Hillyard SA, Mouloua M, Woldorff MG, Clark VP, Hawkins HL. Effects of spatial cuing
on luminance detectability: Psychophysical and electrophysiological evidence for early selection.
Journal of Experimental Psychology: Human Perception and Performance. 1994;20:887–904.

58.   Luck SJ, Heinze HJ, Mangun GR, Hillyard SA. Visual event-related potentials index focused at-
tention within bilateral stimulus arrays. II. Functional dissociation of P1 and N1 components. Elec-
troencephalography and Clinical Neurophysiology. 1990 Jun 1;75(6):528–42.

59.   Mangun GR, Hillyard SA. Modulations of sensory-evoked brain potentials indicate changes in
perceptual processing during visual-spatial priming. J Exp Psychol Hum Percept Perform. 1991
Nov;17(4):1057–74.

60.   Clark VP, Fan S, Hillyard SA. Identification of early visual evoked potential generators by reti-
notopic and topographic analyses. Human Brain Mapping. 1994;2(3):170–87.

61.   Woldorff M g., Fox P t., Matzke M, Lancaster J l., Veeraswamy S, Zamarripa F, et al. Retino-
topic organization of early visual spatial attention effects as revealed by PET and ERPs. Human
Brain Mapping. 1997;5(4):280–6.

62.   Di Russo F, Martínez A, Sereno MI, Pitzalis S, Hillyard SA. Cortical sources of the early com-
ponents of the visual evoked potential. Human Brain Mapping. 2002 Feb;15(2):95–111.

63.   Di Russo F, Spinelli D. Electrophysiological evidence for an early attentional mechanism in
visual processing in humans. Vision Research. 1999 Sep 1;39(18):2975–85.

64.   Green DL, Payne L, Polikar R, Moberg PJ, Wolk DA, Kounios J. P50: A candidate ERP bi-
omarker of prodromal Alzheimer's disease. Brain Research. 2015 Oct 22;1624:390–7.

65.   Golob EJ, Johnson JK, Starr A. Auditory event-related potentials during target detection are
abnormal in mild cognitive impairment. Clinical Neurophysiology. 2002 Jan 1;113(1):151–61.

66.   Golob EJ, Starr A. Effects of stimulus sequence on event-related potentials and reaction time
during target detection in Alzheimer's disease. Clinical Neurophysiology. 2000 Aug 1;111(8):1438–
49.

67.   Polich J. Updating P300: An integrative theory of P3a and P3b. Clinical Neurophysiology.
2007 Oct 1;118(10):2128–48.

68.   Rosenfeld JP, Labkovsky E. New P300-based protocol to detect concealed information: Re-
sistance to mental countermeasures against only half the irrelevant stimuli and a possible ERP in-
dicator of countermeasures. Psychophysiology. 2010;47(6):1002–10.

69.   Meixner JB, Labkovsky E, Peter Rosenfeld J, Winograd M, Sokolovsky A, Weishaar J, et al.
P900: A Putative Novel ERP Component that Indexes Countermeasure Use in the P300-Based
Concealed Information Test. Appl Psychophysiol Biofeedback. 2013 Jun 1;38(2):121–32.

70.   Hull J, Harsh J. P300 and sleep-related positive waveforms (P220, P450, and P900) have dif-
ferent determinants. Journal of Sleep Research. 2001;10(1):9–17.

71.   Weitzman ED, Kremen H. Auditory evoked responses during different stages of sleep in man.
Electroencephalography and Clinical Neurophysiology. 1965 Jan 1;18(1):65–70.

72.   Zhao Y, Tu S, Lei M, Qiu J, Ybarra O, Zhang Q. The neural basis of breaking mental set: an
event-related potential study. Exp Brain Res. 2011 Jan 1;208(2):181–7.

73.   Anderer P, Saletu B, Semlitsch HV, Pascual-Marqui RD. Non-invasive localization of P300
sources in normal aging and age-associated memory impairment. Neurobiology of Aging. 2003
May 1;24(3):463–79.

74.   Mulert C, Pogarell O, Juckel G, Rujescu D, Giegling I, Rupp D, et al. The neural basis of the
P300 potential. European Archives of Psychiatry and Clinical Neurosciences. 2004 Jun
1;254(3):190–8.

75.   Leckey M, Federmeier KD. The P3b and P600(s): Positive contributions to language compre-
hension. Psychophysiology. 2020;57(7):e13351.

76.   Gouvea AC, Phillips C, Kazanina N, Poeppel D. The linguistic processes underlying the P600.
Language and Cognitive Processes. 2010 Feb 1;25(2):149–88.

77.   Sassenhagen J, Schlesewsky M, Bornkessel-Schlesewsky I. The P600-as-P3 hypothesis revisit-
ed: Single-trial analyses reveal that the late EEG positivity following linguistically deviant material
is reaction time aligned. Brain and Language. 2014 Oct 1;137:29–39.

78.   Theofilas P, Ehrenberg AJ, Dunlop S, Di Lorenzo Alho AT, Nguy A, Leite REP, et al. Locus co-
eruleus volume and cell population changes during Alzheimer's disease progression: A stereologi-
cal study in human postmortem brains with potential implication for early-stage biomarker dis-
covery. Alzheimer's & Dementia. 2017 Mar 1;13(3):236–46.

79.   Tomlinson BE, Irving D, Blessed G. Cell loss in the locus coeruleus in senile dementia of Alz-
heimer type. Journal of the Neurological Sciences. 1981 Mar 1;49(3):419–28.

80.   Mather M, Harley CW. The Locus Coeruleus: Essential for Maintaining Cognitive Function
and the Aging Brain. Trends in Cognitive Sciences. 2016 Mar 1;20(3):214–26.

81.   Matchett BJ, Grinberg LT, Theofilas P, Murray ME. The mechanistic link between selective
vulnerability of the locus coeruleus and neurodegeneration in Alzheimer's disease. Acta Neuropa-
thol. 2021 May 1;141(5):631–50.

82.   Bartus RT, Dean RL, Beer B, Lippa AS. The Cholinergic Hypothesis of Geriatric Memory Dys-
function. Science. 1982 Jul 30;217(4558):408–14.

83.   Betts MJ, Cardenas-Blanco A, Kanowski M, Spottke A, Teipel SJ, Kilimann I, et al. Locus co-
eruleus MRI contrast is reduced in Alzheimer's disease dementia and correlates with CSF Aβ lev-
els. Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring. 2019 Dec 1;11:281–5.

84.   Stern Y. Cognitive reserve. Neuropsychologia. 2009 Aug 1;47(10):2015–28.

85.   Cabeza R, Anderson N, Locantore J, Mcintosh a. Aging Gracefully: Compensatory Brain Activ-
ity in High-Performing Older Adults. NeuroImage. 2002 Nov;17(3):1394–402.

86.   Cabeza R, Albert M, Belleville S, Craik FIM, Duarte A, Grady CL, et al. Maintenance, reserve
and compensation: the cognitive neuroscience of healthy ageing. Nat Rev Neurosci. 2018
Nov;19(11):701–10.

87.   Barulli D, Stern Y. Efficiency, capacity, compensation, maintenance, plasticity: emerging con-
cepts in cognitive reserve. Trends in Cognitive Sciences. 2013 Oct 1;17(10):502–9.

88.   Gu L, Chen J, Gao L, Shu H, Wang Z, Liu D, et al. Cognitive reserve modulates attention pro-
cesses in healthy elderly and amnestic mild cognitive impairment: An event-related potential
study. Clinical Neurophysiology. 2018 Jan 1;129(1):198–207.

89.   Speer ME, Soldan A. Cognitive reserve modulates ERPs associated with verbal working
memory in healthy younger and older adults. Neurobiology of Aging. 2015 Mar 1;36(3):1424–34.

90.   Gajewski PD, Falkenstein M, Thönes S, Wascher E. Stroop task performance across the
lifespan: High cognitive reserve in older age is associated with enhanced proactive and reactive
interference control. NeuroImage. 2020 Feb 15;207:116430.

91.   Berchicci M, Lucci G, Pesce C, Spinelli D, Di Russo F. Prefrontal hyperactivity in older people
during motor planning. NeuroImage. 2012 Sep 1;62(3):1750–60.

92.   Berchicci M, Lucci G, Di Russo F. Benefits of Physical Exercise on the Aging Brain: The Role of
the Prefrontal Cortex. The Journals of Gerontology: Series A. 2013 Nov 1;68(11):1337–41.

93.   Ritchie SJ, Tucker-Drob EM. How Much Does Education Improve Intelligence? A Meta-
Analysis. Psychol Sci. 2018 Aug 1;29(8):1358–69.

94.   Ramchandran K, Zeien E, Andreasen NC. Distributed neural efficiency: Intelligence and age
modulate adaptive allocation of resources in the brain. Trends in Neuroscience and Education.
2019 Jun 1;15:48–61.

95.   Lipp I, Benedek M, Fink A, Koschutnig K, Reishofer G, Bergner S, et al. Investigating Neural
860   Efficiency in the Visuo-Spatial Domain: An fmri Study. PLOS ONE. 2012 Dec 12;7(12):e51316.

861   96.   Fischer FU, Wolf D, Tüscher O, Fellgiebel A, on behalf of Alzheimer's Disease Neuroimaging
862   Initiative. Structural Network Efficiency Predicts Resilience to Cognitive Decline in Elderly at Risk
863   for Alzheimer's Disease. Front Aging Neurosci [Internet]. 2021 Feb 22 [cited 2024 Jul 12];13.
864   Available from: https://www.frontiersin.org/journals/aging-
865   neuroscience/articles/10.3389/fnagi.2021.637002/full

866   97.   Mevel K, Chételat G, Eustache F, Desgranges B. The Default Mode Network in Healthy Aging
867   and Alzheimer's Disease. International Journal of Alzheimer's Disease. 2011;2011(1):535816.

868   98.   Stern Y. Cognitive Reserve and Alzheimer Disease. Alzheimer Disease & Associated Disor-
869   ders. 2006 Jun;20(2):112.

870   99.   Fischer FU, Gerber S, Tüscher O, Initiative the ADN. A mathematical model of the Alz-
871   heimer's Disease biomarker cascade demonstrates statistical pitfalls in identifying neurobiological
872   surrogates of cognitive reserve [Internet]. bioRxiv; 2023 [cited 2024 Jul 11]. p.
873   2023.10.26.563793. Available from:
874   https://www.biorxiv.org/content/10.1101/2023.10.26.563793v1

875   100.   Lavanga M, Stumme J, Yalcinkaya BH, Fousek J, Jockwitz C, Sheheitli H, et al. The virtual aging
876   brain: a model-driven explanation for cognitive decline in older subjects [Internet]. bioRxiv; 2022
877   [cited 2023 Jul 13]. p. 2022.02.17.480902. Available from:
878   https://www.biorxiv.org/content/10.1101/2022.02.17.480902v2

879   101.   Amato LG, Vergani AA, Lassi M, Fabbiani C, Mazzeo S, Burali R, et al. Personalized modeling
880   of Alzheimer's disease progression estimates neurodegeneration severity from EEG recordings.
881   Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring. 2024;16(1):e12526.

882   102.   Sperling RA, Aisen PS, Beckett LA, Bennett DA, Craft S, Fagan AM, et al. Toward defining the
883   preclinical stages of Alzheimer's disease: Recommendations from the National Institute on Aging-
884   Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimer's
885   & Dementia. 2011;7(3):280–92.

886   103.   Koenig T, Prichep L, Lehmann D, Sosa PV, Braeker E, Kleinlogel H, et al. Millisecond by Milli-
887   second, Year by Year: Normative EEG Microstates and Developmental Stages. NeuroImage. 2002
888   May 1;16(1):41–8.

889   104.   Lassi M, Fabbiani C, Mazzeo S, Burali R, Vergani AA, Giacomucci G, et al. Degradation of EEG
890   microstates patterns in subjective cognitive decline and mild cognitive impairment: Early bi-
891   omarkers along the Alzheimer's Disease continuum? Neuroimage Clin. 2023;38:103407.

892   105.   Doan DNT, Ku B, Choi J, Oh M, Kim K, Cha W, et al. Predicting Dementia With Prefrontal Elec-
893   troencephalography and Event-Related Potential. Front Aging Neurosci [Internet]. 2021 Apr 13
894   [cited 2024 Jul 11];13. Available from: https://www.frontiersin.org/journals/aging-
895   neuroscience/articles/10.3389/fnagi.2021.659817/full

896   106.   Zhang J, Xia J, Liu X, Olichney J. Machine Learning on Visibility Graph Features Discriminates
897   the Cognitive Event-Related Potentials of Patients with Early Alzheimer's Disease from Healthy
898   Aging. Brain Sciences. 2023 May;13(5):770.

107.   Kim NH, Yang DW, Choi SH, Kang SW. Machine Learning to Predict Brain Amyloid Pathology in Pre-dementia Alzheimer's Disease Using QEEG Features and Genetic Algorithm Heuristic. Front Comput Neurosci [Internet]. 2021 Nov 11 [cited 2024 Jul 11];15. Available from: https://www.frontiersin.org/journals/computational-neuroscience/articles/10.3389/fncom.2021.755499/full

108.   Rutkowski TM, Abe MS, Komendzinski T, Sugimoto H, Narebski S, Otake-Matsuura M. Machine learning approach for early onset dementia neurobiomarker using EEG network topology features. Front Hum Neurosci [Internet]. 2023 Jun 16 [cited 2024 Jul 11];17. Available from: https://www.frontiersin.org/journals/human-neuroscience/articles/10.3389/fnhum.2023.1155194/full

109.   Chedid N, Tabbal J, Kabbara A, Allouch S, Hassan M. The development of an automated machine learning pipeline for the detection of Alzheimer's Disease. Sci Rep. 2022 Oct 28;12(1):18137.

110.   Ieracitano C, Mammone N, Hussain A, Morabito FC. A novel multi-modal machine learning based approach for automatic classification of EEG recordings in dementia. Neural Networks. 2020 Mar 1;123:176–90.

111.   Kim M jae, Youn YC, Paik J. Deep learning-based EEG analysis to classify normal, mild cognitive impairment, and dementia: Algorithms and dataset. NeuroImage. 2023 May 15;272:120054.

112.   Sibilano E, Buongiorno D, Lassi M, Grippo A, Bessi V, Sorbi S, et al. Understanding the Role of Self-Attention in a Transformer Model for the Discrimination of SCD From MCI Using Resting-State EEG. IEEE Journal of Biomedical and Health Informatics. 2024 Jun;28(6):3422–33.

113.   Sibilano E, Brunetti A, Buongiorno D, Lassi M, Grippo A, Bessi V, et al. An attention-based deep learning approach for the classification of subjective cognitive decline and mild cognitive impairment using resting-state EEG. J Neural Eng. 2023 Feb;20(1):016048.

114.   Frisoni GB, Festari C, Massa F, Ramusino MC, Orini S, Aarsland D, et al. European intersocietal recommendations for the biomarker-based diagnosis of neurocognitive disorders. The Lancet Neurology. 2024 Mar 1;23(3):302–12.

115.   Sabri O, Sabbagh MN, Seibyl J, Barthel H, Akatsu H, Ouchi Y, et al. Florbetaben PET imaging to detect amyloid beta plaques in Alzheimer's disease: phase 3 study. Alzheimers Dement. 2015 Aug;11(8):964–74.

116.   Buerger K, Ewers M, Pirttilä T, Zinkowski R, Alafuzoff I, Teipel SJ, et al. CSF phosphorylated tau protein correlates with neocortical neurofibrillary pathology in Alzheimer's disease. Brain. 2006 Nov;129(Pt 11):3035–41.

117.   Mengel D, Soter E, Ott JM, Wacker M, Leyva A, Peters O, et al. Blood biomarkers confirm subjective cognitive decline (SCD) as a distinct molecular and clinical stage within the NIA-AA framework of Alzheimer's disease [Internet]. medRxiv; 2024 [cited 2024 Jul 16]. p. 2024.07.10.24310205. Available from: https://www.medrxiv.org/content/10.1101/2024.07.10.24310205v1

118.   Babiloni C, Blinowska K, Bonanni L, Cichocki A, De Haan W, Del Percio C, et al. What electrophysiology tells us about Alzheimer's disease: a window into the synchronization and connectivity of brain neurons. Neurobiology of Aging. 2020 Jan 1;85:58–73.

940   119.   Horvath A, Szucs A, Csukly G, Sakovics A, Stefanics G, Kamondi A. EEG and ERP biomarkers of
941   Alzheimer's disease: a critical review. Frontiers in bioscience (Landmark edition). 2018 Jan
942   1;23:183–220.

943   120.   Knopman DS. For a dementia diagnosis, clinical acumen must precede biomarkers. The Lan-
944   cet Neurology. 2024 Mar 1;23(3):225–6.

945
946

Supporting Materials

S-Figure 1. Scatterplots between peak and latency of ERP canonical components. (A) P1. (B) P2. (C) N1. (D) P300. (E) P600. (F). P900. Legend contains the Spearman correlation measure and p-value between latency and peak for each clinical condition. Color code: HS (blu), SCD (orange), MCI (red).

S-Figure 2. ERP dynamics across performance in HS. (A) . ERP computed in the cluster of occipital channels
(PO7, PO8, O1, Oz, O2) representative of the encoding phase of the stimulus. (B) ERP computed in cluster of
central channels (FC1, FCz, FC2, C1, Cz, C2) representative of the decision-making phase regarding the stimu-
lus. Both panels (A) and (B): Bold representation is the overall mean within each group and shading is the
standard deviation. The measures on top are the instantaneous H-statistic of the Kruskal-Wallis test and the
associated p-value corrected by Bonferroni's method (alpha<0.05). Temporal instants associated with a
p<0.05 are highlighted with a vertical violet bar. P1/N1/P2 and P300/600/900 labels stand for the name of
the canonical event-related potentials relative to the encoding phase and decision-making phase respective-
ly. Colour code: low performance subjects (green), high performance subjects (cyan).

S-Figure 3. ERP dynamics across ATN conditions. (A) . ERP computed in the cluster of occipital channels (PO7, PO8, O1, Oz, O2) representative of the encoding phase of the stimulus. (B) ERP computed in the cluster of central channels (FC1, FCz, FC2, C1, Cz, C2) representative of the decision-making phase regarding the stimulus. Both panels (A) and (B): Bold representation is the overall mean within each group and shading is the standard deviation. The measures on top are the instantaneous H-statistic of the Kruskal-Wallis test and the associated p-value corrected by Bonferroni's method (alpha<0.05). Temporal instants associated with a p<0.05 are highlighted with a vertical violet bar. P1/N1/P2 and P300/600/900 labels stand for the name of the canonical event-related potentials relative to the encoding phase and decision-making phase, respectively. Colour code: ATN positive (brown), ATN negative (purple), ATN unknown (black).

S-Figure 4. ERP dynamics across ATN conditions in SCD patients. (A) . ERP computed in the cluster of occipi-
tal channels (PO7, PO8, O1, Oz, O2) representative of the encoding phase of the stimulus. (B) ERP computed
in the cluster of central channels (FC1, FCz, FC2, C1, Cz, C2) representative of the decision-making phase re-
garding the stimulus. Both panels (A) and (B): Bold representation is the overall mean within each group
and shading is the standard deviation. The measures on top are the instantaneous H-statistic of the Kruskal-
Wallis test and the associated p-value corrected by Bonferroni's method (alpha<0.05). Temporal instants as-
sociated with a p<0.05 are highlighted with a vertical violet bar. P1/N1/P2 and P300/600/900 labels stand
for the name of the canonical event-related potentials relative to the encoding phase and decision-making
phase respectively. Colour code: ATN positive (brown), ATN-SCD negative (purple), ATN-SCD unknown
(black).

S-Figure 5. ERP dynamics across ATN conditions in MCI patients. (A) . ERP computed in the cluster of occipi-
tal channels (PO7, PO8, O1, Oz, O2) representative of the encoding phase of the stimulus. (B) ERP computed
in the cluster of central channels (FC1, FCz, FC2, C1, Cz, C2) representative of the decision-making phase re-
garding the stimulus. Both panels (A) and (B): Bold representation is the overall mean within each group
and shading is the standard deviation. The measures on top are the instantaneous H-statistic of the Kruskal-
Wallis test and the associated p-value corrected by Bonferroni's method (alpha<0.05). Temporal instants as-
sociated with a p<0.05 are highlighted with a vertical violet bar. P1/N1/P2 and P300/600/900 labels stand
for the name of the canonical event-related potentials relative to the encoding phase and decision-making
phase, respectively. Colour code: ATN-MCI positive (brown), ATN-MCI negative (purple), ATN-MCI unknown
(black).