A Tale of Twins: The Distinct Roles of Intracellular and Extracellular Formate in Regulating Flagellar and Pathogenicity Island-1 Genes in *Salmonella* Typhimurium

Debapriya Mukherjee[1], Dipshikha Chakravortty[1,2*]

[1]Department of Microbiology and Cell Biology, Division of Biological Sciences, Indian Institute of Science, Bangalore, India.

[2]School of Biology, Indian Institute of Science Education and Research, Thiruvananthapuram, 695551.

[*]Corresponding author: dipa@iisc.ac.in

Tel: 0091 80 2293 2842

Fax: 0091 80 2360 2697

Abstract

Host-derived short-chain fatty acids (SCFAs) are extensively being studied for their role in the virulence and pathogenesis of *Salmonella* Typhimurium (STM). Formate, an SCFA present in the ileum, functions as a signalling molecule to enhance STM invasion. However, the role of intracellular formate in *Salmonella* virulence remains poorly understood. To investigate this, we generated knockouts of *pflB* (pyruvate-formate lyase) and *focA* (formate transporter). Disruption of formate production through *pflB* deletion led to reduced flagellation and increased *hilA* and *prgH* expression, attributed to elevated intracellular pH and membrane damage. This suppression of flagellar machinery drives a shift from adhesion to invasion, regulated by RpoE via the CsrA/*csrB* pathway. Additionally, we demonstrate that upon compensation for intracellular deficiency of formate, STM *ΔpflB* starts to utilize formate as a signalling molecule to regulate downstream processes. To the best of our knowledge, this study is the first to establish the critical role of the *pflB* gene in maintaining intracellular pH and controlling virulence gene expression in STM. Lastly, our findings emphasize the importance of modulating *pflB* expression across intestinal regions to optimize STM invasion.

Keywords: Flagella/ Formate/ Intracellular pH/ Membrane depolarization/ Short Chain Fatty Acids (SCFAs)

Introduction

The genus *Salmonella* consists of Gram-negative gammaproteobacteria that infect a wide range

of human and animal hosts [1]. Non-typhoidal serovars, such as *Salmonella* Typhimurium, are

commonly associated with self-limiting diarrheal illness with relatively low fatality. According

to the Global Burden of Diseases, Injuries, and Risk Factors Study (GBD) 2017, there were an

estimated 95.1 million cases of *Salmonella* enterocolitis worldwide, resulting in 50,771 deaths

[2-4]. Although non-typhoidal infections are typically confined to the intestines, these serovars

can occasionally invade sterile tissues, leading to bacteraemia, meningitis, and other focal

infections, which significantly increase mortality rates.[5].

Salmonella Typhimurium enters the host via contaminated food and water. One of the defence

mechanisms in the host is the presence of short-chain fatty acids (SCFAs) in the small intestine.

SCFAs are produced through the anaerobic fermentation of non-digestible polysaccharides,

such as resistant starches and dietary fibres. The SCFAs found in the intestine—formate (C1),

acetate (C2), propionate (C3), and butyrate (C4), with concentrations reaching up to 10–100

mM. Recent studies have extensively explored the role of SCFAs ((like butyrate and

propionate) in *Salmonella* infections [6-10]. For example, butyrate downregulates the

expression of pathogenicity island-1 (SPI-1) genes in *S.* Typhimurium, thereby reducing

bacterial invasion and translocation from the intestines to the bloodstream [11-13].

Formate present in the intestine has been previously shown to act as a diffusible signal

promoting *Salmonella* Typhimurium invasion. [14]. Imported, unmetabolized formic acid

functions as a cytoplasmic signal by binding to HilD, the master transcriptional regulator of

Salmonella invasion, thereby preventing inhibitory fatty acids from binding and extending the

activity of HilD, which leads to the derepression of invasion genes [15]. Moreover, formate

utilization by respiratory enzymes such as formate dehydrogenase-N (FdnGHI) and formate

dehydrogenase-O (FdoGHI) plays a key role in *Salmonella* colonization of the gut [16]. *S.*

Typhimurium can source formate from both its own pyruvate-formate lyase (PflB) and from

the intestinal environment [17, 18]. However, the role of endogenously produced formate in

regulating *Salmonella* virulence and pathogenesis have seldom been reported [19].

In this study, we examined the role of endogenously produced formate in the pathogenic

lifestyle of *S.* Typhimurium. To the best of our knowledge, we are the first to identify the role

of the *pflB* gene in regulating cytosolic pH of STM, which is instrumental in maintaining the

flagellar operon. Quite interestingly, in the wild-type strain with an intact formate pool,

extracellular formate functions as a signalling molecule via the BarA/SirA two-component

system, highlighting the distinct roles of intracellular and extracellular formate pools in

regulating *Salmonella* virulence. We demonstrate that regulating the *pflB* gene in various

regions of the intestine may be a strategy utilized by *Salmonella* to optimize its invasion of the

distal ileum, its primary target site.

Materials and Methods

Bacterial strains and plasmids

The wild-type *Salmonella enterica* subspecies *enterica* serovar Typhimurium strain 14028S

used in this study was generously provided by Professor Michael Hensel from the Max von

Pettenkofer Institute for Hygiene and Medical Microbiology, Germany. Knockout and

complement strains were generated specifically for this study. Bacterial strains were stored as

glycerol stocks at -80°C and revived on fresh Luria-Bertani (LB) agar plates (HiMedia) with

appropriate antibiotics as needed (50 µg/ml Kanamycin, 50 µg/ml Ampicillin, 25 µg/ml

89   Chloramphenicol). A single colony was inoculated into a fresh LB tube with antibiotics when

90   required and incubated at 37°C with shaking (170 rpm) to obtain an overnight culture. Strains

91   containing the pKD46 plasmid were incubated overnight at 30°C with shaking (170 rpm). A

92   complete list of bacterial strains and plasmids used in this study is provided in Table 1.

Construction of the knockout strains of *Salmonella*

The knockout strains of *Salmonella* were generated using the one-step chromosomal gene

inactivation technique developed by Datsenko and Wanner [20]. In brief, STM (WT) was

transformed with the pKD46 plasmid, which expresses the λ-Red recombinase system under

the control of an arabinose-inducible promoter. Transformed cells were selected on LB plates

containing 50 μg/ml Ampicillin. To generate the knockout strains, a single colony of STM

(WT)/pKD46 was inoculated into fresh LB medium with 50 μg/ml Ampicillin and 50 mM

arabinose, then incubated overnight at 30°C with shaking (170 rpm). The overnight culture was

then subcultured into fresh LB medium and incubated under the same conditions for 2.5 hours

to reach an OD_{600} of 0.35-0.4. Electrocompetent STM (WT)/pKD46 cells were prepared by

washing the bacterial pellet with chilled, autoclaved MilliQ water and 10% glycerol.

Kanamycin (KanR, 1.5 kb) and Chloramphenicol (ChlR, 1.1 kb) resistance cassettes were

amplified from pKD4 and pKD3 plasmids, respectively, using knockout-specific primers. The

amplified resistance cassettes were then electroporated into STM (WT)/pKD46, and

transformed cells were selected on LB agar plates with the appropriate antibiotics (Kanamycin

or Chloramphenicol). Knockout colonies were confirmed using gene-specific intragenic

primers, knockout confirmatory primers, and antibiotic-cassette-specific primers.

For generating double-knockout strains, the one-step chromosomal gene inactivation method

was slightly modified. Briefly, STM Δ*pflB* was transformed with the pKD46 plasmid. An

overnight culture of the transformed strain was subcultured into fresh LB medium supplemented with 50 mM arabinose and 50 µg/ml Ampicillin, then incubated at 30°C with shaking (170 rpm) for 2.5 hours until it reached an OD600 of 0.3-0.4. The chloramphenicol resistance cassette (1.1 kb), flanked by regions homologous to *focA*, was electroporated into electrocompetent STM Δ*pflB*/pKD46 cells to generate STM Δ*focAΔpflB*. The double-knockout strains were selected on LB agar plates containing Kanamycin (50 µg/ml) and Chloramphenicol (25 µg/ml). These double-knockout colonies were further confirmed with knockout confirmatory and gene-specific intragenic primers. The primers used for generating knockouts are listed in Table 2.

Construction of complemented and overexpression strains of *Salmonella*

The *pflB* and *csrA* genes were amplified from the genomic DNA of wild-type *Salmonella* using gene-specific cloning primers. Both the colony PCR products and the empty pQE60 cloning vector were digested with BamHI-HF and HindIII-HF (NEB) at 37°C for 1 hour. After double digestion, the insert and vector were purified and ligated using T4 DNA Ligase (NEB) at 16°C overnight. The ligation product was then transformed into *Escherichia coli* TG1 using the heat shock method. Successful cloning was confirmed by double digestion of the recombinant plasmid. The confirmed recombinant plasmid was subsequently transformed into *Salmonella* strains to create complemented and overexpression strains. Primers used for generating complement strains are listed in Table 2.

RNA Isolation and RT-qPCR

Logarithmic-phase cultures of bacterial cells was centrifuged at 6,000 rpm at 4°C, and the

resulting pellet was resuspended in TRIzol reagent. The suspension was stored at -80°C until

further use. RNA was isolated from the lysed supernatants using the chloroform-isopropanol

method. The RNA pellet was washed with 70% ethanol and resuspended in 30 µl of DEPC-

treated MilliQ water. RNA concentration was measured with a NanoDrop spectrophotometer

(Thermo Fisher Scientific), and quality was assessed on a 2% agarose gel. Following this, 2 µg

of RNA was treated with DNase using DNase TURBO (Thermo Fisher Scientific) at 37°C for

30 minutes. DNase was inactivated by adding 5 mM Na_2EDTA (Thermo Fisher Scientific) and

incubating at 65°C for 10 minutes. cDNA synthesis was carried out using the PrimeScript RT

reagent kit (Takara, CAT RR037A) according to the manufacturer's instructions. Quantitative

real-time PCR (qRT-PCR) was performed with a TB Green RT-qPCR kit (Takara) on the

QuantStudio 5 Real-Time PCR system. Intragenic primers were used to examine gene

expression, and expression levels were normalized to 16S rRNA. Primer sequences are listed

in Table 2.

Eukaryotic cell lines and growth conditions

The HeLa, Caco-2, HepG2, and RAW 264.7 cell lines were cultured in Dulbecco's Modified

Eagle's Medium (DMEM, Sigma-Aldrich) supplemented with 10% Fetal Calf Serum (FCS,

Gibco), at 37°C with 5% CO_2.

Gentamicin Protection Assay and Intracellular Cell Survival Assay

Caco-2 and RAW 264.7 cells were seeded into tissue culture-treated 24-well plates. For

infection of the epithelial Caco-2 cells, overnight cultures of STM WT, STM Δ*pflB*, STM

Δ*pflB*/pQE60:*pflB*, and STM Δ*focA*Δ*pflB* were subcultured in fresh LB medium, with or

without supplementation of 40 mM sodium formate (Sigma). For infection of the macrophage

cell line RAW 264.7, the overnight cultures were used directly. Both cell types were infected

at a multiplicity of infection (MOI) of 10. The plates were centrifuged at 1000 rpm for 10

minutes, after which the cells were incubated at 37°C with 5% CO_2 for 25 minutes. Following

incubation, cells were washed with 1X phosphate-buffered saline (PBS) to remove

extracellular, non-adherent, and loosely adherent bacteria, and fresh media containing 100

µg/ml gentamicin was added. After a 1-hour incubation, cells were washed again with 1X PBS

and exposed to 25 µg/ml gentamicin until lysis. Cell lysis was performed at 2 hours and 16

hours post-infection using 0.1% Triton X-100, and the lysates were plated onto *Salmonella-*

Shigella (SS) agar alongside the pre-inoculum. The percent invasion and fold proliferation were

determined using the following formulas:

Percent invasion/phagocytosis= [C.F.U at 2h/ C.F.U of Pre-Inoculum] *100

Fold proliferation= [C.F.U at 16h/ C.F.U of 2h]

Adhesion assay

The adhesion assay was performed by modifying the previously reported protocol [21].

Overnight cultures (12 hours old) of STM WT, STM Δ*pflB*, and STM Δ*fliC* were subcultured

into fresh LB medium, with or without supplementation of 40 mM sodium formate (Sigma).

The logarithmic phase of these bacterial strains was used to infect the cells at a multiplicity of

infection (MOI) of 10. The plates were centrifuged at 1000 rpm for 10 minutes, followed by

incubation at 37°C with 5% CO_2 for 10 minutes. To remove extracellular, unadhered, and

loosely adhered bacteria, the cells were washed twice with 1X phosphate-buffered saline

(PBS). Subsequently, the cells were lysed with 0.1% Triton X-100, and the lysates were plated

on Salmonella-Shigella (SS) agar alongside the pre-inoculum. The percent adhesion was determined using the following formula:

Percent adhesion= [C.F.U at 10 mins Post Infection/ C.F.U of Pre-Inoculum] *100

TEM

Transmission Electron Microscopy was performed by modifying the protocol by Marathe *et al* [22]. In brief, overnight cultures of STM WT, STM *ΔpflB*, STM *ΔfocA*, STM *ΔfocAΔpflB*, STM WT/*csrA*, STM *ΔpflB*/*csrA*, STM *ΔrpoE*, and STM *ΔpflBΔrpoE* were subcultured (1:100) in fresh LB medium, with or without supplementation (+/-F), and incubated at 37°C for 2.5–3 hours under shaking conditions. The bacterial samples were then washed twice with 1X phosphate-buffered saline (PBS) and resuspended in 50 µl of 1X PBS. A 5 µl aliquot of the suspension was applied to a glow-discharged copper grid and allowed to air-dry. The bacteria were subsequently negatively stained using 1% uranyl acetate. After air-drying, the samples were imaged using Transmission Electron Microscopy (JEM-100 CX II; JEOL).

Confocal Microscopy

Confocal Microscopy was performed by modifying the protocol by Marathe *et al* [22]. In summary, 12-hour-old stationary-phase cultures of STM WT, STM *ΔpflB*, STM *ΔfocAΔpflB*, STM WT/*csrA*, STM *ΔpflB*/*csrA*, STM *ΔrpoE*, and STM *ΔpflBΔrpoE*, with or without supplementation (+/-F), were subcultured (1:100) into fresh LB medium and incubated at 37°C for 2.5–3 hours under shaking conditions until reaching the logarithmic phase. A 100 µl aliquot of the log-phase culture was spotted onto sterile coverslips and air-dried overnight. Once completely dried, the samples were fixed with 3.5% paraformaldehyde (PFA). The PFA was washed off with 1X PBS, and the samples were stained with a rabbit-raised anti-Fli primary

antibody (1:100 dilution in 2.5% BSA solution, Difco) for 3 hours at room temperature. After

washing off the primary antibody with 1X PBS, the samples were incubated with Alexa-488-

conjugated anti-rabbit secondary antibody (Sigma) for 45 minutes at room temperature. Before

mounting the coverslips onto clean glass slides, the samples were stained with DAPI (4',6-

diamidino-2-phenylindole) at 100 µg/ml for 5 minutes. The coverslips were sealed with

transparent nail polish. All images were captured using a Zeiss 880 microscope and analyzed

with Zeiss ZEN Black 2012 software.

Gas Chromatography-Mass Spectrometry

GC-MS was performed by modifying the protocol by Hughes *et al* [23]. Briefly, overnight

cultures of STM WT, STM *ΔpflB*, STM *ΔfocA*, STM *ΔfocAΔpflB* were subcultured (1:100) into

100 ml of LB Broth. After 2.5-3h of incubation at 37˚C under shaking conditions, 50 ml of the

individual cultures were centrifuged at 6,000 rpm for 10 mins. The pellet was washed twice

with 1X PBS to remove the remnants of the media. Finally, the pellet was resuspended in 1 ml

of 1X PBS, and the suspension was sonicated to lyse the bacterial cells. The cell debris was

removed by centrifugation. Before sonication, the CFU in the pellet had been recorded by

plating.

Sodium formate (Sigma) was used as an external standard. External standards and biological

samples were derivatized before the Mass Spectrometry. 2 M hydrochloric acid was added in

a 1:1 ratio to the samples to protonate formate. Liquid-liquid extraction was done which

extracted formic acid into ethyl acetate (Sigma-Aldrich). To remove any remaining water, the

ethyl acetate extract was passed through a column of anhydrous sodium sulfate (Sigma-

Aldrich). The ethyl acetate extract of formic acid was then incubated at 80 ºC for 1 h with the

derivatization reagent N, O-Bis(trimethylsilyl)trifluoro acetamide (Sigma Aldrich) with 1%

Chlorotrimethylsilane (Fluka). The derivatized samples were then transferred to autosampler

vials for analysis by Gas Chromatography Mass Spectrometry (Agilent technologies, 7890A GC, 5975C MS). The injection temperature was set to 200 °C and the injection split ratio was 1:40 with an injection volume of 1 µL. The oven temperature was set to start at 40 °C for 4 min, with an increase to 120 °C at 5 °C per min and to 200 °C at 25 °C per min with a final hold at this temperature for 3 min. Helium carrier gas had a flow rate of 1.2mL/min with constant flow mode. A 30 m × 0.25 mm × 0.40 µm DB-624 column from Agilent was used. The interface temperature was set to 220 °C. The electron impact ion source temperature was at 230 °C, with an ionization energy of 70 eV and 1235 EM volts. For qualitative experiments, selected ion monitoring (single quadrupole mode) of 103 m/z scan had been performed. The retention time for formate and deuterated formate was 5.9 min. The target and reference (qualifier) ions for formate were m/z = 103 and m/z = 75, respectively. The formate concentration of each sample was interpolated from a standard curve with the known standards. It was normalized with the CFU values of each sample to obtain the intracellular formate concentration/CFU.

To determine the formate transportation into the cell, we subcultured overnight cultures of STM WT, STM *ΔpflB*, STM *ΔfocA*, and STM *ΔfocAΔpflB* into fresh LB tubes. During the subculture, Sodium Formate was supplemented into each tube to maintain a concentration of 1M. Post 2.5-3h of incubation at 37˚C under shaking conditions, the bacterial cells were removed from the media by centrifuging the culture twice at 6,000 rpm for 10 mins. A 2 µm filter was used to remove any other bacterial cells left in the spent media. Following this, a similar protocol as mentioned above was used to quantify the residual formate in the spent media. It was compared to the media where bacterial cells had not been subcultured.

Bisbenzimide Assay

The bisbenzimide assay was performed by modifying the protocol by Roy Chowdhury *et al*

[24]. Briefly, overnight cultures of STM WT, STM *ΔpflB*, and STM *ΔfliC* were subcultured

into fresh LB tubes(+/-F). Logarithmic phase cultures of each condition were taken and the

OD_{600} was adjusted to 0.1. To 180 µl of this bacterial suspension, 20 µl of Bisbenzimide dye

(10µg/ml) was added. It was incubated at 37˚C for 10 mins. The intracellular DNA-bound

bisbenzimide was quantified in a TECAN 96 well microplate reader using a 346 nm excitation

and 460 nm emission filter.

Staining with DiBAC$_4$

The DiBAC$_4$ staining was performed by modifying the protocol by Roy Chowdhury *et al* [25].

Membrane potential or membrane porosity was quantified by using a membrane potential

sensitive fluorescent dye bis-(1,3-dibutyl barbituric acid)- trimethylene oxonol (Invitrogen).

For this purpose, overnight culture of STM WT, STM *ΔpflB*, and STM *ΔfliC* were subcultured

with or without the supplementation of 40 mM Sodium Formate (Sigma). Around $4.5*10^7$ cells

of the logarithmic phase culture were incubated with 1 µg/ml of DiBAC$_4$ dye at 37˚C for 30

mins. The percentage of DiBAC$_4$-positive cells and the Median Fluorescence Intensity (MFI)

of DiBAC$_4$ was determined by flow cytometry (BD FACSVerse by BD Biosciences-US).

BCECF-AM Staining

273 The BCECF-AM staining was performed by modifying the protocol by Roy Chowdhury *et al*

274 [24]. The acidification of the cytosol with the change in pH of extracellular conditions was

275 determined using a cell-permeable dual excitation ratiometric dye 2',7'-Bis-(2-Carboxyethyl)-

276 5- (and-6)-Carboxyfluorescein, Acetoxymethyl Ester (BCECF-AM). Briefly, the logarithmic

phase cultures of STM WT, STM *ΔpflB* (+/-F) were resuspended in buffers of different pH

(5,5.5,6,7). They were incubated for 2 hours at 37˚C before the addition of 1 mg/ml BCECF-

AM to make a final concentration of 20 μM. The cells were incubated with the dye for 30

minutes. Post-incubation, the cells were analysed by flow cytometry (BD FACSVerse by BD

Biosciences, US) with 488 nm and 405 nm excitation and 535 nm excitation channels. The

MFI of the bacterial cells at 488 nm and 405 nm was obtained from BD FACSuite software

and the ratio of 488/405 was utilized to determine the cytosolic acidification in response to

changing pH conditions.

Intracellular pH measurement by pHuji plasmid

STM WT and STM *ΔpflB* were transformed with the plasmid pBAD:pHuji (Plasmid #61555)

that encodes a pH-sensitive red fluorescent protein pHuji [26, 27]. The logarithmic cultures of

STM WT/pBAD:pHuji and STM *ΔpflB*/pBAD:pHuji (+/-F) were incubated in phosphate

buffers of pH 5,6,7,8,9 in the presence of 40 μM Sodium Benzoate to equilibrate the

intracellular pH with the extracellular one. The resultant fluorescent intensity was plotted as a

function of pH and was fitted to obtain a standard curve. The fluorescence values obtained from

LB-grown cultures of the bacterial cells (+/- F) were interpolated in the standard curve to obtain

the intracellular pH of the bacterial cells *in-vitro*.

Ethics Statement for Animal Experiments

All the animal experiments have been approved by the Institutional Animal Ethics Clearance

Committee (IAEC), Indian Institute of Science, Bangalore. The registration number is

48/1999/CPCSEA. The guidelines formulated by the Committee for the Purpose of Control

and Supervision of Experiments on Animals (CPCSEA) were strictly followed. The ethical

clearance number for this study is CAF/Ethics/853/2021.

Mice Survival Assay

6-8 weeks old male C57BL/6 mice and Balb/c mice were acquired from the specific-pathogen

free conditions of Central Animal Facility, IISc. To draw a comparison between the mortality

rate of WT and mutant strains, the mice (n=10) were orally gavaged with 10^8 cells from the

stationary phase culture of each. After infection, mice were monitored every day for survival,

and the data was represented as percent survival.

Determination of bacterial burden in different organs

To understand the organ burden of the WT and mutant strain post-infection in mice, 10^7 cells

from the overnight culture of the bacterial strains were orally gavaged into 6-8 weeks old

C57BL/6 mice. 5 days post-infection, the mice were ethically euthanized, and the liver, spleen,

and brain were acquired. The organs were homogenized, and appropriate dilutions of the

homogenate were plated on *Salmonella-Shigella* (SS Agar). The data was represented was

log_{10}CFU/gm-wt.

Invasion Assay in mice

Invasion assay in mice was performed by modifying the protocol by Chandra *et al* [28]. Briefly,

10^7 cells from the overnight culture of the bacterial cells were orally gavaged into 6-8 weeks

old male C57BL/6 mice. 6 hours post-infection, the mice were ethically euthanized, and the

322   intestine was acquired. It was homogenized, and appropriate dilutions of the homogenate were

323   plated on *Salmonella-Shigella* (SS Agar). Corresponding fecal matter was also collected, and

324   it was also plated post homogenization. The data was represented was $\log_{10}$CFU/gm-wt.

326   Intraperitoneal infection of mice

327   Intraperitoneal infection of mice was performed with 10^3 CFU from the overnight cultures of

328   WT and mutant strains. 2 days post-infection, the mice were ethically euthanized, and the liver

329   and the spleen were obtained. The organ burden of the bacterial strains was determined by the

330   same protocol as above.

332   Statistical Analysis

Each experiment has been repeated 1 to 8 times independently as mentioned in the figure

legends. The statistical analyses were performed by a unpaired Student's t-test or two-way

ANOVA multiple comparison test as denoted in the figure legends. The data obtained from the

mice experiments were analyzed using the Mann-Whitney *U* test. The *p*-values below 0.05

were considered significant. The results are indicated as Mean $\pm$ SD or Mean $\pm$ SEM as

mentioned in the figure legends. All the data were plotted and analysed using GraphPad Prism

8.4.2.

342   Results

The deletion of the *pflB* gene led to decreased invasion efficiency in the Caco-2 cell line, which corresponded to a reduced organ burden in C57BL/6 mice five days after infection through oral gavaging

To investigate the role of endogenous formate levels in *Salmonella* virulence and pathogenesis, we created knockout strains of *pflB* and *focA* using the one-step gene inactivation technique outlined by Datsenko and Wanner [20]. FocA is the formate transporter in *Salmonella* Typhimurium that functions as a passive export channel at high external pH and switches to an active importer of formate at low pH [29]. The deletion of either gene did not impact the *in vitro* growth of *Salmonella* in either complex LB media or M9 Minimal Media (**Figure S1 A-B**). Since the *focA* and *pflB* genes are part of the same operon, we investigated whether deleting one gene affected the expression of the other [30]. Our study revealed that deleting the upstream *focA* gene resulted in decreased expression of the downstream *pflB* gene (**Figure S1C**). However, deletion of *pflB* increased the expression of *focA* (**Figure S1D**).

To confirm that deleting *pflB* and *focA* indeed depleted the native formate pool of the cell, we collected lysates from logarithmic phase cultures of all strains and quantified the formate concentration using Gas Chromatography-Mass Spectrometry (GC-MS). After normalizing by CFU, we observed that there was a significant reduction in the relative intracellular formate concentrations upon deletion of *pflB*, whereas deletion of *focA* did not cause a notable change. There was a significantly reduced intracellular formate level in STM *ΔfocAΔpflB* (**Figure 1A**). This finding suggested the possibility of an alternative formate transporter. To investigate further, we incubated secondary cultures of all strains in formate-supplemented media and measured formate concentration in the spent media after 3 hours by GC-MS. We found a significant reduction in formate concentration in the spent media of STM *ΔfocA* compared to the control, indicating the presence of an alternative formate importer (**Figure 1B**). Since STM

Δ*focA* retained intracellular formate levels similar to STM WT and our data suggested the

presence of an alternate formate transporter, we focused our study on STM Δ*pflB*, where we

could successfully deplete the intracellular formate pool.

Intracellular Cell Survival Assay (ICSA) using the human colorectal carcinoma cell line Caco-

2 showed that STM *ΔpflB* exhibited reduced invasion efficiency compared to STM WT, but it

had similar fold proliferation to the latter (**Figure 1C**). The invasion deficiency in STM *ΔpflB*

was also rescued by complementing the gene with an exogenous pQE60 plasmid (**Figure S1E**).

In the murine macrophage cell line RAW 264.7, STM *ΔpflB* showed a decreased phagocytosis

and hyperproliferation compared to STM WT (**Figure 1E, F**).

To determine if the invasion deficiency observed in STM *ΔpflB* was consistent in an animal

model, we orally gavaged 10^7 CFU of STM WT and STM *ΔpflB* into 6–8-week-old male

C57BL/6 mice. Adhesion-deficient STM *ΔfliC* served as a control. Six hours post-infection,

we observed a reduced bacterial load in the mouse intestine for STM Δ*pflB*, which was

associated with increased faecal shedding compared to STM WT (**Figure 1G, H, I**). These

findings were consistent with the results for the control strain STM *ΔfliC*, which is known to

have reduced adhesion and is cleared more rapidly from the host [31].

To further understand the role of *pflB* deletion in overall *Salmonella* pathogenesis in *in-vivo*

mouse model, we orally gavaged 10^7 CFU of bacteria in 6-8 weeks old male C57BL/6 mice

and euthanized the mice 5 days post-infection (dpi) to determine the bacterial burden in the

liver, the spleen, and the brain (**Figure 1J, K, L**). We found a significantly reduced bacterial

burden in the liver and the spleen when infected with STM Δ*pflB* as compared to infection with

STM WT (**Figure 1K, L**). We also found that C57BL/6 mice infected with STM Δ*pflB* had a

better survival rate as compared to mice infected with STM WT (**Figure 1M**). We subjected

C57BL/6 mice to an intra-peritoneal infection with 10^3 CFU of both the WT and the knockout

strains. Under these conditions, we observed no significant change in the organ burden of STM WT and STM *ΔpflB* 3dpi (**Figure S2A-D**). All these data suggested that the reduced organ burden of STM *ΔpflB* upon oral gavaging is due to its attenuated adhesion in the gastrointestinal tract, which promoted a faster clearance from the host body.

STM *ΔpflB* was deficient in flagella and the deficiency was partially recovered upon supplementation of 40 mM sodium formate

Salmonella Typhimurium uses adhesins, such as flagella, to attach to host cells during infection [32-34]. Following successful adhesion, it employs the *Salmonella* Pathogenicity Island-1 (SPI-1) machinery to invade non-phagocytic epithelial cells [35-37]. Our previous findings indicated that STM Δ*pflB* has reduced invasion efficiency in Caco-2 cells. We opted to proceed by supplementing exogenous formate to STM Δ*pflB* to determine whether the effects observed from depleting intracellular formate could indeed be reversed through the addition of extracellular formate. In our further studies, we used a concentration of 40 mM (indicated as F henceforth), as it yielded maximal expression of the SPI-1 regulatory gene *hilA* in STM WT in a concentration-dependent analysis ranging from 0 to 50 mM (**Figure S3A**). Interestingly, we observed that STM Δ*pflB* exhibited increased expression of *prgH*, which encodes a structural protein of the SPI-1 apparatus (**Figure 2A**). This expression was further elevated in STM Δ*pflB*+F. Additionally, STM WT+F showed significantly higher *prgH* expression compared to STM WT, consistent with the findings of Huang *et al* [17].

We also found that STM Δ*pflB* had lower *fliC* expression, which was partially restored under +F conditions. Formate supplementation reduced *fliC* expression in STM WT as well, compared to the untreated control (**Figure 2B**). We confirmed this via Transmission Electron Microscopy (TEM) as well, where we found that STM WT+F had lesser flagella compared to

untreated control. STM *ΔpflB* had a reduced flagellar number, which was partially restored

upon supplementation of F (**Figure 2C**). Quantitative analysis of normalized MFI in confocal

microscopy and adhesion assays in the Caco-2 cell line further validated the findings from qRT-

PCR and TEM (**Figure 2 D-F**). STM WT+F also showed lesser motility on swim agar plate

compared to STM WT, however, we did not see a recovery in motility of STM *ΔpflB* upon F

supplementation (**Figure 2 G**). This indicates that while F supplementation in STM *ΔpflB* may

restore the transcription level of *fliC* and its protein expression, it does not lead to a recovery

in flagella-mediated motility. The partially restored adhesion observed in STM *ΔpflB*+F in the

Caco-2 cell line could be attributed to the increased surface hydrophobicity of the flagellar

filaments, resulting from the recovery in protein levels. This enhanced hydrophobicity

promotes stronger interactions for effective adhesion to eukaryotic host cells [38]. F

supplementation in STM *ΔpflB* was also associated with an improved invasion in Caco-2 cell

line. Complementation of the *pflB* via an extragenous plasmid in STM *ΔpflB* also led to an

improved percent invasion compared to the knockout strain. In line with our previous data, we

also found that STM WT+F had a higher percent invasion compared to STM WT (**Figure 2H**).

Further experimentation involved incubating secondary cultures of STM WT and STM *ΔpflB*

in varying concentrations of formate. Notably, formate significantly inhibited *fliC* expression

in STM WT starting from a concentration of 1 mM, while a concentration as low as 0.125 mM

was sufficient to restore reduced *fliC* expression in STM *ΔpflB* (**Figure S3B**). Additionally, 10

mM formate significantly upregulated *hilA*—the master regulator of SPI-1—in both strains

(**Figure S3C**).

The double knockout strain STM *ΔfocAΔpflB* exhibited similar growth kinetics as STM WT in

LB broth and M9 Minimal Media (**Figure S4A, B**). However, due to its depleted formate pool,

it was also deficient in flagella, with partial recovery in anti-Fli MFI and adhesion in Caco-2

cell line when incubated with formate (**Figure S4 C, D, E**). In line with our previous findings,

we observed that oral gavaging of STM *ΔfocA* led to a similar bacterial burden in the liver and

the spleen as STM WT 5 dpi. STM *ΔfocAΔpflB* showed lesser organ burden in the liver and

spleen of the C57 mice model (**Figure S4 F**).

STM *ΔpflB* had a higher intracellular pH and higher membrane depolarization, both of which were recovered upon F supplementation

Previous studies in *Escherichia coli* indicated that formate translocation through the FocA

channel and its metabolism via the FHL-1 (Formate Hydrogen Lyase-1) complex are crucial

for maintaining cellular pH homeostasis during anaerobic growth or fermentation [39-45]. This

raised the question of whether formate synthesis during aerobic growth in *Salmonella*

Typhimurium could also contribute to maintaining intracellular pH (ipH). Additionally, weak

acids like acetate and benzoate can cross the cell membrane in their neutral forms and

subsequently dissociate, thereby reducing ipH [46]. Based on this, we also hypothesized that

the endogenous formate pool might play a role in maintaining intracellular pH (ipH). To

investigate how changes in the surrounding media's pH affect intracellular pH, we incubated

STM WT and STM *ΔpflB* (+/- F) in phosphate buffers at pH levels 5.5, 6, and 7. After

incubation, we stained the cells with the pH-sensitive dye BCECF-AM. A higher 488/405 nm

fluorescence ratio for STM *ΔpflB* under all pH conditions indicated a higher ipH in the

knockout strain compared to the WT (**Figure 3A**).

To further confirm these results, we transformed STM WT and STM *ΔpflB* with the plasmid

pBAD-pHuji, which encodes a pH-sensitive red fluorescent protein. The higher ipH observed

in STM *ΔpflB*/pBAD-pHuji confirmed that *pflB* plays a significant role in maintaining

cytosolic pH in *Salmonella* Typhimurium. Consistent with the BCECF-AM staining results, formate did not affect the ipH of STM WT/pBAD-pHuji (**Figure 3B**). These findings suggest the possibility that extracellular and intracellular formate pools might independently regulate the expression of *fliC* and *hilA* through distinct mechanisms.

One of the causes of membrane depolarization in biological systems is disrupted pH homeostasis [47]. To investigate membrane potential in all strains with and without formate supplementation, we analysed the exponential cultures of STM *ΔpflB* in LB. These cultures showed a higher population of DiBAC$_4$-positive cells, indicating increased membrane depolarization. Supplementation with formate successfully reduced membrane depolarization in STM *ΔpflB*, but it had no effect on membrane polarization in STM WT (**Figure 3D, E**). Similar results were observed when cells were stained with the DNA-binding fluorescent dye bisbenzimide (**Figure 3C**).

Furthermore, the extent of membrane depolarization in STM *ΔpflB* was reduced when stationary-phase bacteria were sub cultured into acidic F-media (pH 5) (**Figure 3E**). The H$^+$ ions in the acidic F-media may help counteract the disrupted cytosolic acidification in the *pflB* knockout strain, thereby restoring membrane polarization.

These findings suggest that the intracellular formate pool is crucial for maintaining intracellular pH (ipH) in *Salmonella* Typhimurium, which in turn supports cell membrane polarization. We also explored whether the elevated ipH in the *pflB* knockout strain contributes to disrupted flagellar apparatus synthesis. Incubation of STM *ΔpflB* in acidic LB medium (pH 6) led to an increase in the *fliC* transcript level (**Figure S5 A**), indicating that higher ipH is responsible for these downstream effects.

STM *ΔpflB* had a higher transcript level of *csrB*, raising the possibility of flagella inhibition via the *csr* sRNA system

To investigate how formate regulates the flagellar operon, we measured the transcript levels of several genes across different classes, including class I (*flhD*), class II (*flgA*, *fliA*), and class III (*flgK*, *motA*) (**Figure 4A**) [48-56]. We discovered that in the formate-deficient *pflB* knockout strain, inhibition of the flagellar operon begins at the level of class I genes or the genes that regulate class I genes. The transcription of the *flhDC* promoter is controlled by various factors, including OmpR, Crp, HdfR, H-NS, QseBC, FliA, LrhA, and others [50, 57-59]. One of the most extensively studied mechanisms of post-transcriptional regulation of *flhDC* involves the Csr system. The Csr system consists of the RNA-binding protein CsrA and the small regulatory RNAs (sRNAs) *csrB* and *csrC* [60]. CsrA is a 61-amino-acid protein that binds to the mRNA of target genes, thereby either inhibiting or enhancing their translation [61]. CsrA can bind to the *flhD* transcript and stabilize it by preventing RNase E cleavage at the 5′ end [62]. The activity of the CsrA protein is antagonized by small regulatory RNAs *csrB* and *csrC*. A single *csrB* RNA molecule can bind up to 18 CsrA molecules, whereas a single *csrC* molecule can bind up to 9 CsrA molecules [63-65]. In *S.* Typhimurium, CsrA has been seen to regulate motility, virulence, carbon storage, and secondary metabolism [66, 67]. The transcription of *csrB* and *csrC* is regulated by the BarA/SirA two-component systems (TCS) orthologues in *E. coli* and *S.* Typhimurium[68-71].

We observed an increase in *csrB* transcript levels in STM *ΔpflB*, which decreased in STM *ΔpflB*+F (**Figure 4B**). We also observed a reduction in *csrA* transcript levels in STM *ΔpflB,* which was recovered upon incubation in F supplemented LB or LB with acidic pH (pH 6) (**Figure S5B**). Since *csrB* directly binds to CsrA and inhibits its activity, we decided to clone *csrA* into the extragenous plasmid pQE60 and transform this recombinant plasmid into STM

ΔpflB. Overexpressing *csrA* through IPTG induction partially restored the reduced expression

of *fliC* in STM *ΔpflB* (**Figure 4C**). CsrA is known to bind and stabilize the *flhD* transcript.

However, we speculate that an excess of CsrA molecules in the cell could potentially stabilize

flhD to the point of inhibiting its translation. This may explain why *csrA* overexpression in

STM WT led to a decrease in *fliC* expression. Overexpression of *csrA* mediated downregulation

of motility has been demonstrated in *Bacillus subtilis* and *Clostridium difficile* as well [72, 73].

The increase in flagellar numbers upon *csrA* overexpression in STM *ΔpflB* was visualized and

quantified using TEM and confocal microscopy (**Figure 4D**, **E, G**). This finding was

corroborated by an adhesion assay with the Caco-2 cell line, where STM *ΔpflB*/p: *csrA* showed

a higher adhesion percentage compared to STM *ΔpflB* (**Figure 4F**).

522   The extracytoplasmic sigma factor RpoE enhances the expression of csrB,

523   which in turn limits the expression of *fliC*

In *E. coli*, RpoE (σ^E) functions as an extracytoplasmic stress factor essential for cell viability

and regulating the response to membrane stress[74]. The membrane-bound anti-sigma factor

RseA binds to σ^E, aided by the periplasmic protein RseB [75, 76]. In the absence of envelope

stress, RseA binding inhibits σ^E from interacting with RNA polymerase [76-79]. When

envelope stress occurs, the proteolytically unstable RseA is quickly degraded, releasing free σ^E

molecules to carry out downstream functions. These functions include activating genes

involved in the biogenesis, transport, and assembly of lipopolysaccharides (LPS),

phospholipids, and outer membrane proteins (OMPs), as well as proteases and chaperones that

maintain or repair outer membrane (OM) integrity [76]. Recent studies have demonstrated that

σ^E also indirectly activates the transcription of *csrB/C* sRNAs from σ^{70} promoters [80, 81].

These findings have been established in *E. coli*. Deletion of *rpoE* in STM showed a reduction

in *csrB* transcript levels (~0.588), suggesting that a similar mechanism may operate in STM as well **(Figure 5A)**.

Our previous data demonstrated that formate produced by PflB helps maintain the intracellular pH (ipH) of the cell. Consequently, deletion of the *pflB* gene disrupts ipH balance, leading to membrane depolarization. We hypothesized that this membrane stress in *STM ΔpflB* could cause RseA degradation, thereby freeing σ^E to promote *csrB* transcription. Our results showed that the elevated *csrB* expression observed in *STM ΔpflB* was significantly reduced in *STM ΔpflBΔrpoE* (**Figure 5A**). Additionally, the suppressed expression of *flhD* and *fliC* in *STM ΔpflB* was restored in *STM ΔpflBΔrpoE* (**Figure 5B**). Confocal microscopy and TEM provided further confirmation of this finding (**Figure 5C-E**). However, we did not observe any recovery in the swim halo upon deletion of *rpoE* from STM *ΔpflB*, which shows that the recovery in translational level and protein level is not getting reflected in the motility of the attenuated strain (**Figure 5F**). However, there was an improved adhesion of *STM ΔpflBΔrpoE* to Caco-2 cells compared to *STM ΔpflB* (**Figure 5G**). This improved adhesion might be due to the enhanced hydrophobicity due to increased FliC protein in STM *ΔpflBΔrpoE* compared to STM *ΔpflB* [38]. These results suggest that the reduction in flagella synthesis in STM *ΔpflB* could be mediated by the extra cytoplasmic stress factor σ^E.

Formate regulates the balance between flagellar expression and SPI-1 gene expression in the intestine of C57BL/6 mice

Our previous data showed that *STM ΔpflB* exhibited a higher abundance of the *prgH* transcript. To determine if this increased expression of the SPI-1 gene was controlled by the same regulatory loop, we analysed the transcript levels of the SPI-1 genes *hilA* and *prgH*. We found that the elevated expression of *hilA* and *prgH* in *STM ΔpflB* was reduced in *STM ΔpflBΔrpoE*

(**Figure 6A**). This confirmed that intracellular formate regulates the shift from flagellar

expression to SPI-1 gene expression.

Our previous results showed that formate supplementation enhances *prgH* expression in both

STM WT and STM *ΔpflB* (**Figure 2A**). Flagellar motility and epithelial cell invasion are

regulated by the BarA/SirA two-component system (TCS) [70]. As expected, STM *ΔsirA*

exhibited low levels of *hilA* and *prgH* expression, and formate supplementation in STM *ΔsirA*

did not significantly alter the expression of these genes. Additionally, deletion of *sirA* in STM

ΔpflB eliminated any significant upregulation of *hilA* and *prgH* in response to formate

supplementation (**Figure 6B**). This indicates that extracellular formate influences the virulence

and pathogenicity of *Salmonella* via the BarA/SirA TCS.

Interestingly, we observed that STM *ΔpflBΔsirA* continued to show recovery in *fliC* transcript

levels under formate supplementation. This suggests that intracellular formate regulates

Salmonella pathogenesis independently of the BarA/SirA TCS (**Figure 6C**). Therefore, when

the intracellular formate pool is maintained by the *pflB* gene, extracellular formate modulates

SPI-1 gene expression through the BarA/SirA TCS. However, if the intracellular formate pool

is disrupted, the resulting membrane stress triggers RpoE-mediated overexpression of *csrB*,

which is associated with reduced *fliC* transcript levels and increased *hilA* and *prgH* expression.

As expected, STM *ΔsirA* exhibited low expression levels of *hilA* and *prgH*, and formate (F)

supplementation in STM *ΔsirA* did not result in any significant change in the expression of

these genes. Furthermore, deletion of *sirA* in STM *ΔpflB* eliminated any significant

upregulation of *hilA* and *prgH* in response to formate supplementation (**Figure 6B**). This

demonstrates that extracellular formate influences the virulence and pathogenesis of

Salmonella through the BarA/SirA TCS. Interestingly, we observed that STM *ΔpflBΔsirA*

continued to show recovery in *fliC* transcript levels with formate supplementation. This

indicates that intracellular formate regulates *Salmonella* pathogenesis independently of the

BarA/SirA TCS (**Figure 6C**). Thus, when the intracellular formate pool is maintained by the

pflB gene, extracellular formate modulates SPI-1 gene expression via the BarA/SirA TCS.

However, if the intracellular formate pool is disrupted, the resulting membrane stress triggers

RpoE-mediated overexpression of *csrB*, which is associated with reduced *fliC* transcript levels

and increased *hilA* and *prgH* expression. We proceeded to understand whether achieving

intracellular formate concentrations of STM Δ*pflB* to similar levels as STM WT and further

supplementing it with formate can replicate the behaviour of wild-type strain in presence of

formate. We subcultured the stationary phase cultures of STM WT and STM Δ*pflB* and

supplemented them higher concentrations of formate (70, 90, 110, 150 mM). We found that

there was a substantial decrease in the in the *fliC* transcript level in STM WT upon

supplementation of higher concentrations of formate. STM Δ*pflB* showed a remarkable

recovery in the *fliC* transcript level at 90 mM concentration, following which the expression

levels declined in the presence of 110 mM and 150 mM formate (**Figure 6D**). GC-MS mediated

quantification of intracellular formate concentration revealed that a concentration of 90 mM

was enough to replenish the depleted intracellular formate pool in STM Δ*pflB,* which explains

the highest expression of *fliC* at this concentration (**Figure 6E**). Beyond this, supplementation

of formate reduces the expression of *fliC* as seen in **Figure 6D**. This mimics the scenario of

supplementing F to STM WT with an intact formate pool. This was also visualized under TEM

and quantified by confocal microscopy (**Figure S6A, B, C**).

To explore the physiological relevance of this phenomenon, we examined *pflB* expression

levels in different sections of the intestine, since flagella and SPI-1 play critical roles in the

intestinal phase of *Salmonella* infection. We observed that *pflB* expression decreased

progressively from the duodenum to the ileum (**Figure 6F**). This finding led us to hypothesize

that *Salmonella* downregulates its *pflB* gene as it approaches its invasion site—the distal ileum.

This downregulation could induce a partial formate deficiency within the bacterial cell, causing membrane depolarization, which in turn activates σ^E, increases *csrB* abundance, downregulates flagellar operon transcription, and upregulates the SPI-1 gene cascade, ultimately help in bacterial invasion at its target site of ileum.

To test this hypothesis, we collected duodenum and ileum sections from mice infected with STM WT and STM *ΔpflB* and stained them with an anti-SipC antibody to measure levels of the SPI-1 effector protein (**Figure 6G**). We normalized the fluorescence from the secondary antibody, specific to the rabbit anti-SipC antibody, against mCherry fluorescence from bacteria transformed with the pFPV plasmid. In mice infected with STM WT, we found that the normalized mean fluorescence intensity (MFI) increased from the duodenum to the ileum. In contrast, STM *ΔpflB*-infected mice showed a decrease in normalized MFI from the duodenum to the ileum (**Figure 6H**). These results indicate that the formate pool maintained by the PflB enzyme regulates SPI-1 gene expression, and its depletion may be a strategy used by *Salmonella* Typhimurium to maximize SPI-1 expression, and thereby invasion, in the ileum.

Discussion

Salmonella possesses a network of genes and regulatory factors that manages the balance between its extracellular and intracellular modes of existence [82]. Research on *Salmonella* Typhimurium has demonstrated overlapping regulatory mechanisms between the flagellar secretion system and the SPI-1. The flagellar protein FliZ is known to activate the primary regulator of flagella synthesis, FlhD, through post-translational modification. Kage *et al*. have shown that FliZ can also regulate HilD, a key regulator of SPI-1, at the transcriptional level [83, 84]. HilD has been found to bind directly to the promoter region of *flhDC*, upstream of the P5 transcriptional start site, thus enhancing its expression [85]. Other studies indicate that SPI-

1 expression can suppress flagellar gene expression, aiding in pathogenicity and virulence of *Salmonella*. A mutation in *fliZ* reduces *hilA* expression by half, while a knockout of *sirA* decreases *hilA* expression tenfold and increases flagella expression a hundredfold [86-89]. Furthermore, recent research by Saleh *et al*. has shown that HilD activation can trigger a SPI-1 dependent induction of the stringent response and significantly reduce the Proton Motive Force (PMF). Although flagellation remains unaffected, HilD activation results in a motility defect in *S.* Typhimurium [90]. Collectively, these studies reveal that *Salmonella* can toggle between flagella synthesis and SPI-1 gene expression to optimize adhesion and the delivery of SPI-1 effectors. There are several potential benefits to downregulating flagella while upregulating SPI-1 genes. First, reducing flagella expression prevents the bacteria from moving away from a target cell primed for invasion. Second, because the flagellar apparatus is immunogenic, downregulating it may help the bacteria evade the host immune response [91-93]. Lastly, simultaneous production of both systems could lead to interference, potentially causing the improper secretion of flagellin and SPI-1 effectors [94].

In our study, we knocked out the gene encoding the pyruvate-formate lyase enzyme, effectively depleting the cell of its native formate pool. Although the knockout strain displayed increased expression of the SPI-1 gene *prgH*, it showed reduced transcript levels of *fliC*. This was reflected in decreased invasion and adhesion capabilities in a human epithelial cell line. These deficiencies were partially restored with the addition of 40 mM formate, which also explains the strain's hyperproliferation in the RAW 264.7 macrophage cell line [95]. Additionally, we found that the STM Δ*pflB* mutant was unable to maintain its intracellular pH to the same extent as the STM wild type, resulting in a membrane depolarization phenotype. Membrane stress triggered the sigma factor RpoE to increase the expression of *csrB*. Since *csrB* is known to bind to and inhibit the CsrA protein, we hypothesized that the elevated *csrB* transcript levels led to reduced *fliC* expression. Overexpressing *csrA* via an extragenous plasmid partially restored

fliC expression, supporting our hypothesis. Furthermore, deletion of *rpoE* in the STM Δ*pflB*

strain successfully reversed the flagella-deficient phenotype, providing further confirmation.

This demonstrated that the extra-cytoplasmic sigma factor RpoE mediates downstream effects

in response to cell membrane stress.

In all our experiments, we also observed that formate supplementation led to reduced flagellar

expression in STM WT. Interestingly, adding formate did not affect the intracellular pH or

membrane potential of STM WT, which led us to hypothesize that extracellular formate impacts

STM pathogenicity differently than intracellular formate synthesis. Huang *et al*. demonstrated

that extracellular formate functions as a diffusible signal to enhance invasion, but this effect

was independent of the BarA/SirA Two-Component System and was also unrelated to *csrB*,

which is directly regulated by the response regulator SirA. In our study, we found that STM

WT with formate supplementation (STM WT+F) showed higher expression of SPI-1 genes

[96]. Contrary to previous findings, we observed that *csrB* transcript levels were higher in STM

WT+F compared to STM WT, suggesting that elevated *csrB* levels might be linked to reduced

fliC expression [97]. This indicates that formate is a crucial metabolite for *Salmonella*, with a

critical concentration determining whether it acts through the TCS pathway or by regulating

the cell's intracellular pH. Additionally, we found that supplementing extracellular formate to

restore the intracellular formate pool caused STM Δ*pflB* to mimic the wild-type strain,

exhibiting reduced *fliC* levels in the presence of formate.

Salmonella Typhimurium has been extensively studied for its ability to adapt its metabolism to

conditions within the host. Previous research has shown that spatially distinct *Salmonella*

populations in the mucous layer and intestinal lumen differ in their energy metabolism.

Transcript levels of pyruvate-formate lyase and formate dehydrogenases were found to be

higher in the mucous layer than in the lumen population [16]. This formate metabolism

provided a fitness advantage in the mucous layer. In our study, we observed that *pflB*

transcription was highest in the large intestine and gradually decreased in the small intestine,

the primary site of *Salmonella* infection. This suggests that *Salmonella* modulates *pflB*

expression in response to local gut microenvironment signals. We propose a model in which

Salmonella downregulates the *pflB* gene upon reaching its infection site, disrupting its

intracellular formate pool. This can result in partial membrane depolarization, which

downregulates the flagellar apparatus and promotes invasion into epithelial cells through

upregulated expression of the SPI-1 system. The specific signals and genetic mechanisms

controlling this modulation remain unclear.

Overall, our study contributes to understanding how intracellular metabolism in *Salmonella*

influences its ability to adhere to and invade non-phagocytic intestinal epithelial cells, and it

highlights the potential for uncovering additional host-pathogen metabolic interactions in

future research.

Acknowledgements

We acknowledge the Departmental Real-Time Facility, Departmental Confocal Microscopy

Facility, Divisional Electron Microscopy Facility, Divisional Mass Spectrometry Facility,

Divisional Flow Cytometry Facility, and Central Animal Facility at IISc for providing us

opportunity for experimentation. Mr. Sumitlal K and Ms. Navya have helped in image

acquisition in Confocal Microscopy. Dr. Sushma and Keerthana helped with image acquisition

in Transmission Electron Microscopy. Dr. Muralidhar and Dr. Sunitha helped in formate

quantitation studies using Gas Chromatography Mass Spectrometry. Ms. Lipika, Ms. Dharani,

and Mr. Munish helped in data acquisition from Flow Cytometry. Dr. Akshay Datey is acknowledged for his initial supervision in the study. The anti-SipC antibody was a kind gift from Prof. Dr. Ayub Qadri, National Institute of Immunology, Delhi, India.

Funding information

We express our heartfelt gratitude to the financial support from the Department of Biotechnology (DBT), Ministry of Science and Technology; Department of Science and Technology (DST), Ministry of Science and Technology. DC acknowledges DAE for the SRC Outstanding Investigator Award and funds, ASTRA Chair Professorship, and TATA Innovation fellowship funds. The authors jointly acknowledge the DBT-IISc Partnership Program. Infrastructure support from ICMR (Center for Advanced Study in Molecular Medicine), DST (FIST), and UGC-CAS (special assistance) is acknowledged. DM acknowledges the IISc fellowship. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

DM and DC have conceived the study and designed the experiments. DM has performed all the experiments and participated in acquiring the data. DM has analysed the data, constructed the figures, and wrote the original draft of the manuscript. DM and DC have participated in the proofreading and editing of the manuscript. DC has supervised the study and helped in fund acquisition. All the authors have read and approved the manuscript.

Declaration of interest

The authors declare no conflict of interest.

Data availability

Data can be available from the authors upon reasonable request.

Data deposition

Link for pre-print: https://doi.org/10.1101/2024.01.30.577997

Thesis link

This research has not yet been included as part of any PhD thesis till current date.

References

1. Galanis, E., et al., *Web-based surveillance and global Salmonella distribution, 2000-2002.* Emerg Infect Dis, 2006. **12**(3): p. 381-8.
2. Collaborators, G.B.D.C.o.D., *Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980-2017: a systematic analysis for the Global Burden of Disease Study 2017.* Lancet, 2018. **392**(10159): p. 1736-1788.
3. DALYs, G.B.D. and H. Collaborators, *Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017.* Lancet, 2018. **392**(10159): p. 1859-1922.
4. Disease, G.B.D., I. Injury, and C. Prevalence, *Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017.* Lancet, 2018. **392**(10159): p. 1789-1858.

5. Crump, J.A., et al., *Epidemiology, Clinical Presentation, Laboratory Diagnosis, Antimicrobial*
Resistance, and Antimicrobial Management of Invasive Salmonella Infections. Clin Microbiol
Rev, 2015. **28**(4): p. 901-37.
6. Maslowski, K.M., et al., *Regulation of inflammatory responses by gut microbiota and*
chemoattractant receptor GPR43. Nature, 2009. **461**(7268): p. 1282-6.
7. Tan, J., et al., *The role of short-chain fatty acids in health and disease.* Advances in
immunology, 2014. **121**: p. 91-119.
8. Thorburn, A.N., L. Macia, and C.R. Mackay, *Diet, metabolites, and "western-lifestyle"*
inflammatory diseases. Immunity, 2014. **40**(6): p. 833-42.
9. Huffnagle, G.B., *Increase in dietary fiber dampens allergic responses in the lung.* Nat Med,
2014. **20**(2): p. 120-1.
10. Machado, M.G., V. Sencio, and F. Trottein, *Short-Chain Fatty Acids as a Potential Treatment*
for Infections: a Closer Look at the Lungs. Infect Immun, 2021. **89**(9): p. e0018821.
11. Boyen, F., et al., *Coated fatty acids alter virulence properties of Salmonella Typhimurium and*
decrease intestinal colonization of pigs. Veterinary microbiology, 2008. **132**(3-4): p. 319-327.
12. Gantois, I., et al., *Butyrate specifically down-regulates Salmonella pathogenicity island 1 gene*
expression. Applied and environmental microbiology, 2006. **72**(1): p. 946-949.
13. Lawhon, S.D., et al., *Intestinal short-chain fatty acids alter Salmonella typhimurium invasion*
gene expression and virulence through BarA/SirA. Molecular microbiology, 2002. **46**(5): p.
1451-1464.
14. Chowdhury, R., et al., *Salmonella Invasion Is Controlled by Competition among Intestinal*
Chemical Signals. mBio, 2023. **14**(2): p. e0001223.
15. Chowdhury, R., et al., *Salmonella Invasion Is Controlled by Competition among Intestinal*
Chemical Signals. mBio, 2023. **14**(2): p. e00012-23.
16. Winter, M.G., et al., *Formate oxidation in the intestinal mucus layer enhances fitness of*
Salmonella enterica serovar Typhimurium. mBio, 2023. **14**(4): p. e0092123.
17. Huang, Y., et al., *Formate Acts as a Diffusible Signal To Induce <i>Salmonella</i> Invasion.*
Journal of Bacteriology, 2008. **190**(12): p. 4233-4241.
18. Winter, M.G., et al., *Formate oxidation in the intestinal mucus layer enhances fitness of*
Salmonella enterica serovar Typhimurium. Mbio, 2023. **14**(4): p. e00921-23.
19. Abernathy, J., et al., *Deletions in the pyruvate pathway of Salmonella Typhimurium alter SPI1-*
mediated gene expression and infectivity. J Anim Sci Biotechnol, 2013. **4**(1): p. 5.
20. Datsenko, K.A. and B.L. Wanner, *One-step inactivation of chromosomal genes in Escherichia*
coli K-12 using PCR products. Proc Natl Acad Sci U S A, 2000. **97**(12): p. 6640-5.
21. Gagnon, M., et al., *Comparison of the Caco-2, HT-29 and the mucus-secreting HT29-MTX*
intestinal cell models to investigate Salmonella adhesion and invasion. J Microbiol Methods,
2013. **94**(3): p. 274-9.
22. Marathe, S.A., et al., *Curcumin Reduces the Motility of Salmonella enterica Serovar*
Typhimurium by Binding to the Flagella, Thereby Leading to Flagellar Fragility and Shedding.
J Bacteriol, 2016. **198**(13): p. 1798-1811.
23. Hughes, E.R., et al., *Microbial Respiration and Formate Oxidation as Metabolic Signatures of*
Inflammation-Associated Dysbiosis. Cell Host Microbe, 2017. **21**(2): p. 208-219.
24. Roy Chowdhury, A., et al., *Salmonella Typhimurium outer membrane protein A (OmpA)*
renders protection from nitrosative stress of macrophages by maintaining the stability of
bacterial outer membrane. PLoS Pathog, 2022. **18**(8): p. e1010708.
25. Chowdhury, A.R., et al., *Loss of outer membrane protein A (OmpA) impairs the survival of*
Salmonella Typhimurium by inducing membrane damage in the presence of ceftazidime and
meropenem. J Antimicrob Chemother, 2022. **77**(12): p. 3376-3389.
26. Shen, Y., et al., *pHuji, a pH-sensitive red fluorescent protein for imaging of exo- and*
endocytosis. J Cell Biol, 2014. **207**(3): p. 419-32.

27. Hajra, D., et al., *SIRT1 and SIRT3 impact host mitochondrial function and host-Salmonella pH balance during infection.* bioRxiv, 2023: p. 2023.09.11.557159.

28. Chandra, K., et al., *GH18 family glycoside hydrolase Chitinase A of Salmonella enhances virulence by facilitating invasion and modulating host immune responses.* PLoS Pathog, 2022. **18**(4): p. e1010407.

29. Lu, W., et al., *pH-dependent gating in a FocA formate channel.* Science, 2011. **332**(6027): p. 352-4.

30. Dong, R., et al., *DsrA Modulates Central Carbon Metabolism and Redox Balance by Directly Repressing pflB Expression in Salmonella Typhimurium.* Microbiol Spectr, 2022. **10**(1): p. e0152221.

31. Parker, C.T. and J. Guard-Petter, *Contribution of flagella and invasion proteins to pathogenesis of Salmonella enterica serovar enteritidis in chicks.* FEMS microbiology letters, 2001. **204**(2): p. 287-291.

32. Haiko, J. and B. Westerlund-Wikstrom, *The role of the bacterial flagellum in adhesion and virulence.* Biology (Basel), 2013. **2**(4): p. 1242-67.

33. Rossez, Y., et al., *Bacterial flagella: twist and stick, or dodge across the kingdoms.* PLoS Pathog, 2015. **11**(1): p. e1004483.

34. Gut, A.M., et al., *Salmonella infection - prevention and treatment by antibiotics and probiotic yeasts: a review.* Microbiology (Reading), 2018. **164**(11): p. 1327-1344.

35. Agbor, T.A. and B.A. McCormick, *Salmonella effectors: important players modulating host cell function during infection.* Cell Microbiol, 2011. **13**(12): p. 1858-69.

36. Finlay, B.B., S. Ruschkowski, and S. Dedhar, *Cytoskeletal rearrangements accompanying salmonella entry into epithelial cells.* J Cell Sci, 1991. **99 (Pt 2)**: p. 283-96.

37. Zhou, D. and J. Galan, *Salmonella entry into host cells: the work in concert of type III secreted effector proteins.* Microbes Infect, 2001. **3**(14-15): p. 1293-8.

38. Horstmann, J.A., et al., *Methylation of Salmonella Typhimurium flagella promotes bacterial adhesion and host cell invasion.* Nature communications, 2020. **11**(1): p. 2013.

39. Kammel, M., C. Pinske, and R.G. Sawers, *FocA and its central role in fine-tuning pH homeostasis of enterobacterial formate metabolism.* Microbiology (Reading), 2022. **168**(10).

40. Suppmann, B. and G. Sawers, *Isolation and characterization of hypophosphite-resistant mutants of Escherichia coli: identification of the FocA protein, encoded by the pfl operon, as a putative formate transporter.* Molecular Microbiology, 1994. **11**(5): p. 965-982.

41. Kammel, M. and R.G. Sawers, *The FocA channel functions to maintain intracellular formate homeostasis during Escherichia coli fermentation.* Microbiology, 2022. **168**(4).

42. Beyer, L., et al., *Coordination of FocA and pyruvate formate-lyase synthesis in Escherichia coli demonstrates preferential translocation of formate over other mixed-acid fermentation products.* J Bacteriol, 2013. **195**(7): p. 1428-35.

43. Wiechert, M. and E. Beitz, *Mechanism of formate-nitrite transporters by dielectric shift of substrate acidity.* EMBO J, 2017. **36**(7): p. 949-958.

44. Lyu, M., et al., *Structural basis of transport and inhibition of the Plasmodium falciparum transporter PfFNT.* EMBO reports, 2021. **22**(3): p. e51628.

45. Helmstetter, F., et al., *Formate-nitrite transporters carrying nonprotonatable amide amino acids instead of a central histidine maintain pH-dependent transport.* J Biol Chem, 2019. **294**(2): p. 623-631.

46. Minamino, T., et al., *Effect of intracellular pH on rotational speed of bacterial flagellar motors.* J Bacteriol, 2003. **185**(4): p. 1190-4.

47. Putnam, R.W., *Chapter 17 - Intracellular pH Regulation*, in *Cell Physiology Source Book (Fourth Edition)*, N. Sperelakis, Editor. 2012, Academic Press: San Diego. p. 303-321.

48. Kutsukake, K. and N. Ide, *Transcriptional analysis of the flgK and fliD operons of Salmonella typhimurium which encode flagellar hook-associated proteins.* Mol Gen Genet, 1995. **247**(3): p. 275-81.

49. Gillen, K.L. and K.T. Hughes, *Transcription from two promoters and autoregulation contribute*
to the control of expression of the Salmonella typhimurium flagellar regulatory gene flgM. J
Bacteriol, 1993. **175**(21): p. 7006-15.
50. Ikebe, T., S. Iyoda, and K. Kutsukake, *Structure and expression of the fliA operon of Salmonella*
typhimurium. Microbiology (Reading), 1999. **145 (Pt 6)**: p. 1389-1396.
51. Ohnishi, K., et al., *Gene fliA encodes an alternative sigma factor specific for flagellar operons*
in Salmonella typhimurium. Mol Gen Genet, 1990. **221**(2): p. 139-47.
52. Karlinsey, J.E., et al., *Completion of the hook-basal body complex of the Salmonella*
typhimurium flagellum is coupled to FlgM secretion and fliC transcription. Mol Microbiol,
2000. **37**(5): p. 1220-31.
53. Ohnishi, K., et al., *A novel transcriptional regulation mechanism in the flagellar regulon of*
Salmonella typhimurium: an antisigma factor inhibits the activity of the flagellum-specific
sigma factor, sigma F. Mol Microbiol, 1992. **6**(21): p. 3149-57.
54. Fraser, G.M., et al., *Substrate specificity of type III flagellar protein export in Salmonella is*
controlled by subdomain interactions in FlhB. Mol Microbiol, 2003. **48**(4): p. 1043-57.
55. Hirano, T., et al., *Substrate specificity classes and the recognition signal for Salmonella type III*
flagellar export. J Bacteriol, 2003. **185**(8): p. 2485-92.
56. Frye, J., et al., *Identification of new flagellar genes of Salmonella enterica serovar*
Typhimurium. J Bacteriol, 2006. **188**(6): p. 2233-43.
57. Kutsukake, K., *Autogenous and global control of the flagellar master operon, flhD, in*
Salmonella typhimurium. Mol Gen Genet, 1997. **254**(4): p. 440-8.
58. Tanabe, Y., et al., *The transcript from the sigma(28)-dependent promoter is translationally*
inert in the expression of the sigma(28)-encoding gene fliA in the fliAZ operon of Salmonella
enterica serovar Typhimurium. J Bacteriol, 2011. **193**(22): p. 6132-41.
59. Ko, M. and C. Park, *H-NS-Dependent regulation of flagellar synthesis is mediated by a LysR*
family protein. J Bacteriol, 2000. **182**(16): p. 4670-2.
60. Vakulskas, C.A., et al., *Regulation of bacterial virulence by Csr (Rsm) systems.* Microbiol Mol
Biol Rev, 2015. **79**(2): p. 193-224.
61. Romeo, T., *Global regulation by the small RNA-binding protein CsrA and the non-coding RNA*
molecule CsrB. Mol Microbiol, 1998. **29**(6): p. 1321-30.
62. Yakhnin, A.V., et al., *CsrA activates flhDC expression by protecting flhDC mRNA from RNase E-*
mediated cleavage. Mol Microbiol, 2013. **87**(4): p. 851-66.
63. Liu, M.Y., et al., *The RNA molecule CsrB binds to the global regulatory protein CsrA and*
antagonizes its activity in Escherichia coli. J Biol Chem, 1997. **272**(28): p. 17502-10.
64. Weilbacher, T., et al., *A novel sRNA component of the carbon storage regulatory system of*
Escherichia coli. Mol Microbiol, 2003. **48**(3): p. 657-70.
65. Dubey, A.K., et al., *RNA sequence and secondary structure participate in high-affinity CsrA-*
RNA interaction. RNA, 2005. **11**(10): p. 1579-87.
66. Altier, C., M. Suyemoto, and S.D. Lawhon, *Regulation of Salmonella enterica serovar*
typhimurium invasion genes by csrA. Infect Immun, 2000. **68**(12): p. 6790-7.
67. Lawhon, S.D., et al., *Global regulation by CsrA in Salmonella typhimurium.* Mol Microbiol,
2003. **48**(6): p. 1633-45.
68. Suzuki, K., et al., *Regulatory circuitry of the CsrA/CsrB and BarA/UvrY systems of Escherichia*
coli. J Bacteriol, 2002. **184**(18): p. 5130-40.
69. Suzuki, K., et al., *Regulatory circuitry of the CsrA/CsrB and BarA/UvrY systems of Escherichia*
coli. Journal of bacteriology, 2002. **184**(18): p. 5130-5140.
70. Teplitski, M., R.I. Goodier, and B.M. Ahmer, *Pathways Leading from BarA/SirA to Motility*
andVirulence Gene Expression in Salmonella. Journal of bacteriology, 2003. **185**(24): p. 7257-
7265.
71. Weilbacher, T., et al., *A novel sRNA component of the carbon storage regulatory system of*
Escherichia coli. Molecular microbiology, 2003. **48**(3): p. 657-670.

72. Gu, H., et al., *Carbon storage regulator CsrA plays important roles in multiple virulence-*
associated processes of Clostridium difficile. Microbial pathogenesis, 2018. **121**: p. 303-309.
73. Yakhnin, H., et al., *CsrA of Bacillus subtilis regulates translation initiation of the gene*
encoding the flagellin protein (hag) by blocking ribosome binding. Molecular microbiology,
2007. **64**(6): p. 1605-1620.
74. Hayden, J.D. and S.E. Ades, *The extracytoplasmic stress factor, sigmaE, is required to*
maintain cell envelope integrity in Escherichia coli. PLoS One, 2008. **3**(2): p. e1573.
75. Ahuja, N., et al., *Analyzing the interaction of RseA and RseB, the two negative regulators of*
the sigmaE envelope stress response, using a combined bioinformatic and experimental
strategy. J Biol Chem, 2009. **284**(8): p. 5403-13.
76. Lima, S., et al., *Dual molecular signals mediate the bacterial response to outer-membrane*
stress. Science, 2013. **340**(6134): p. 837-41.
77. De Las Penas, A., L. Connolly, and C.A. Gross, *The sigmaE-mediated response to*
extracytoplasmic stress in Escherichia coli is transduced by RseA and RseB, two negative
regulators of sigmaE. Mol Microbiol, 1997. **24**(2): p. 373-85.
78. Missiakas, D., et al., *Modulation of the Escherichia coli sigmaE (RpoE) heat-shock*
transcription-factor activity by the RseA, RseB and RseC proteins. Mol Microbiol, 1997. **24**(2):
p. 355-71.
79. Campbell, E.A., et al., *Crystal structure of Escherichia coli sigmaE with the cytoplasmic*
domain of its anti-sigma RseA. Mol Cell, 2003. **11**(4): p. 1067-78.
80. Pannuri, A., et al., *Circuitry Linking the Catabolite Repression and Csr Global Regulatory*
Systems of Escherichia coli. J Bacteriol, 2016. **198**(21): p. 3000-3015.
81. Yakhnin, H., et al., *Circuitry Linking the Global Csr- and sigma(E)-Dependent Cell Envelope*
Stress Response Systems. J Bacteriol, 2017. **199**(23).
82. Fabrega, A. and J. Vila, *Salmonella enterica serovar Typhimurium skills to succeed in the host:*
virulence and regulation. Clin Microbiol Rev, 2013. **26**(2): p. 308-41.
83. Kage, H., et al., *Coordinated regulation of expression of Salmonella pathogenicity island 1*
and flagellar type III secretion systems by ATP-dependent ClpXP protease. J Bacteriol, 2008.
190(7): p. 2470-8.
84. Saini, S., et al., *FliZ Is a posttranslational activator of FlhD4C2-dependent flagellar gene*
expression. J Bacteriol, 2008. **190**(14): p. 4979-88.
85. Singer, H.M., et al., *The Salmonella Spi1 virulence regulatory protein HilD directly activates*
transcription of the flagellar master operon flhDC. J Bacteriol, 2014. **196**(7): p. 1448-57.
86. Eichelberg, K. and J.E. Galan, *The flagellar sigma factor FliA (sigma(28)) regulates the*
expression of Salmonella genes associated with the centisome 63 type III secretion system.
Infect Immun, 2000. **68**(5): p. 2735-43.
87. Iyoda, S., et al., *A flagellar gene fliZ regulates the expression of invasion genes and virulence*
phenotype in Salmonella enterica serovar Typhimurium. Microb Pathog, 2001. **30**(2): p. 81-
90.
88. Lucas, R.L., et al., *Multiple factors independently regulate hilA and invasion gene expression*
in Salmonella enterica serovar typhimurium. J Bacteriol, 2000. **182**(7): p. 1872-82.
89. Goodier, R.I. and B.M. Ahmer, *SirA orthologs affect both motility and virulence.* J Bacteriol,
2001. **183**(7): p. 2249-58.
90. Saleh, D.O., et al., *SPI-1 virulence gene expression modulates motility of Salmonella*
Typhimurium in a proton motive force- and adhesins-dependent manner. PLoS Pathog, 2023.
19(6): p. e1011451.
91. Gewirtz, A.T., et al., *Salmonella typhimurium induces epithelial IL-8 expression via Ca(2+)-*
mediated activation of the NF-kappaB pathway. J Clin Invest, 2000. **105**(1): p. 79-92.
92. McSorley, S.J., B.T. Cookson, and M.K. Jenkins, *Characterization of CD4+ T cell responses*
during natural infection with Salmonella typhimurium. J Immunol, 2000. **164**(2): p. 986-93.

93. Takahashi, A., et al., *Production of beta-defensin-2 by human colonic epithelial cells induced*
by Salmonella enteritidis flagella filament structural protein. FEBS Lett, 2001. **508**(3): p. 484-
8.
94. Ellermeier, C.D. and J.M. Slauch, *RtsA and RtsB coordinately regulate expression of the*
invasion and flagellar genes in Salmonella enterica serovar Typhimurium. J Bacteriol, 2003.
185(17): p. 5096-108.
95. Olsen, J.E., et al., *The role of flagella and chemotaxis genes in host pathogen interaction of*
the host adapted Salmonella enterica serovar Dublin compared to the broad host range
serovar S. Typhimurium. BMC Microbiol, 2013. **13**: p. 67.
96. Huang, Y., et al., *Formate acts as a diffusible signal to induce Salmonella invasion.* J Bacteriol,
2008. **190**(12): p. 4233-41.
97. Teplitski, M., R.I. Goodier, and B.M. Ahmer, *Pathways leading from BarA/SirA to motility and*
virulence gene expression in Salmonella. J Bacteriol, 2003. **185**(24): p. 7257-65.

Tables:

Table 1: List of strains

Strains/ plasmids	Characteristics	Source/ Information
Salmonella enterica subspecies *enterica* serovar Typhimurium strain 14028S	Wild type (WT)	Gift from Prof. M. Hensel
S. Typhimurium *ΔfocA*	Chl^R	This study
S. Typhimurium *ΔpflB*	Kan^R	This study
S. Typhimurium *ΔpflB*/pQE60	Kan^R, Amp^R	This study
S. Typhimurium *ΔpflB*/pQE60: *pflB*	Kan^R, Amp^R	This study
S. Typhimurium *ΔfocAΔpflB*	Kan^R, Chl^R	This study
S. Typhimurium *ΔfliC*	Kan^R	Laboratory stock
S. Typhimurium/pQE60: *csrA*	Amp^R	This study
S. Typhimurium *ΔpflB*/pQE60: *csrA*	Kan^R, Amp^R	This study

pKD4	Plasmid with FRT flanked Kanamycin-resistant cassette	
pKD3	Plasmid with FRT flanked Chloramphenicol-resistant cassette	
pKD46	Plasmid expressing λ-red recombinase system, Amp^R	
pQE60 vector	Low copy number plasmid, Amp^R	Laboratory stock
pHuji		Addgene

 Table 2: List of primers:

PRIMER NAME	SEQUENCE (5'-3')
16S rRNA RT FP	GTGAGGTAACGGCTCACCAA
16S rRNA RT RP	TAACCGCAACACCTTCCTCC
hilA RT FP	GCCGGTGACCATTACGAAGA
hilA RT RP	AAGAGAGAAGCGGGTTGGTG
prgH RT FP	CGTTTTCAGGTGTTGCCAGG
prgH RT RP	TACGCGGCTCATCGAAATGA
flhD RT FP	CGCCTCGGTATCAACGAAGA
flhD RT RP	CACTTCATTGAGCAGACGCG
fliC RT FP	CTAAACAAACTGGGTGGCGC

fliC RT RP	GCACCCAGGTCAGAACGTAA
flgA RT FP	ATTATGTCGCCGTTGCC
flgA RT RP	TGACCTGTACTCGTTGC
flgK RT FP	AACCACCGATCAGTATC
flgK RT RP	TCGTCAGGTTATAGGTG
fliA RT FP	GCTGGATGAATTACGCA
fliA RT RP	CCACTCATCGTAAGAGA
motA RT FP	TTGCTGGCGTTGCTCTA
motA RT RP	GATGATCAGGCGCAGAT
csrB RT FP	CGTACAACGAAGCGAACGTC
csrB RT RP	TGTGACCTTACGGCCTGTTC
focA KO FP	ACACCGTAATTGCATAAAAGCCATGCGACTTACGGGCCTACATATGAATATCCTCCTTAG

focA KO RP	ATGCTCGTTACCACGCAGGTAAATGACCCAGTATGTCAACGTGTAGGCTGGAGCTGCTTC
pflB KO FP	ATGTCCGAGCTTAATGAAAAGTTAGCCACAGCCTGGGAACATATGAATATCCTCCTTAG
pflB KO RP	TACATGGTCTGCGTGAAGGTACGAGTAATAACGTCCTGCTGTGTAGGCTGGAGCTGCTTC
focA RT FP	CGGAAAGCCACATCAGTAAC
focA RT RP	ATCGGGGGTATCTGTTTCTC
pflB RT FP	ATAACGTCCTGCTGCTGTTC
pflB RT RP	ATGGTTACTTCCACCACGAA
pflB cloning FP	CGCGGATCCATGTCCGAGCTTAATGA
pflB cloning RP	CCCAAGCTTCAGTCAAACCCATTACA
rpoE KO FP	GGTTTGGGGAGACATTACCTCGGATGAGCGAGCAGTTACATATGAATATCCTCCTTAG
rpoE KO RP	CCTAATACCTTTTCCAGTATCCCGCTATCGTCAACGGTGTAGGCTGGAGCTGCTTC

sirA KO FP	CTATCAGTAGCGTTATCCCTATTCTGGAGATATTCCTTTGCATATGAATATCCTCCTTAG
sirA KO RP	ATACGATAGCGATAGCTGTTCACCGTTTTAGGACTGAGATGTGTAGGCTGGAGCTCTTC
csrA cloning FP	CGCGGATCCATGCTGATTCTGACTCG
csrA cloning RP	CCCAAGCTTTTAGTAACTGGACTGCTGG

984 **Figures**

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

991 **Figure 6** N=4, n=2

Graphical abstract

Supplementary Figure 1

A.

Intraperitoneal
administration of 10³ CFU
of bacterial strains

Euthanization of mice 3dpi,
organs collected for enumeration
of bacterial burden by plating

6-8 weeks old,
male C57BL/6
mice

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

999   **Supplementary Figure 6**

Figure legends:

Figure 1: The deletion of the *pflB* gene led to decreased invasion efficiency in the Caco-2 cell line, which corresponded to a reduced organ burden in C57BL/6 mice five days after infection through oral gavaging

A. GC-MS mediated quantification of relative intracellular formate in STM WT, STM *ΔpflB*, STM *ΔfocA*, STM *ΔfocAΔpflB*. Data is represented as Mean+/-SEM of N=3, n=6.

B. GC-MS mediated quantification of residual formate in the spent media post incubation by STM WT, STM *ΔpflB*, STM *ΔfocA*, STM *ΔfocAΔpflB*. Data is represented as Mean+/-SEM of N=3, n=3.

C. Percent invasion of STM WT and STM *ΔpflB* in Caco-2 cell line. Data is represented as Mean+/-SEM of N=3, n=3.

D. Fold proliferation of STM WT and STM *ΔpflB* in Caco-2 cell line. Data is represented as Mean+/-SEM of N=3, n=3.

E. Percent phagocytosis of STM WT and STM *ΔpflB* in RAW 264.7 cell line. Data is represented as Mean+/-SEM of N=3, n=3.

F. Fold proliferation of STM WT and STM *ΔpflB* in RAW 264.7 cell line. Data is represented as Mean+/-SEM of N=3, n=3.

G. Schematic showing the protocol followed for invasion assay in C57BL/6 mice

H. Bacterial burden in the intestine post 6h of infecting the mice with STM WT and STM *ΔpflB* via oral gavaging. Data is represented as Mean+/-SEM of N=2, n=5.

I. Bacterial burden in the feces of mice post 6h of infection with STM WT and STM *ΔpflB* via oral gavaging. Data is represented as Mean+/-SEM of N=2, n=5.

J. Schematic showing the protocol followed for determining the organ burden of STM WT and STM *ΔpflB* 5 days post oral gavaging

K. Organ burden of STM WT and STM *ΔpflB* in liver 5 days post oral gavaging. Data is represented as Mean+/-SEM of N=3, n=5.

L. Organ burden of STM WT and STM *ΔpflB* in spleen 5 days post oral gavaging. Data is represented as Mean+/-SEM of N=3, n=5.

M. Survival rates of C57BL/6 mice post infection with STM WT and STM *ΔpflB* through oral route.

(Unpaired Student's t-test for column graphs, Two-way Anova for grouped data, Mann-Whitney U-test for animal experiment data (**** $p < 0.0001$, *** $p < 0.001$, ** $p<0.01$, * $p<0.05$))

Figure 2: STM *ΔpflB* was deficient in flagella and the deficiency was recovered upon supplementation of formate

A. RT-qPCR mediated study of expression profile of *prgH* in the logarithmic phase cultures of STM WT, STM *ΔpflB* (+/-F). Data is represented as Mean+/-SEM of N=3, n=3.

B. RT-qPCR mediated study of expression profile of *fliC* in the logarithmic phase cultures of STM WT, STM *ΔpflB* (+/-F). Data is represented as Mean+/-SEM of N=2, n=3.

C. TEM assisted visualization of the flagellar structures in STM WT, STM *ΔpflB* (+/-F)

D. Confocal microscopy assisted visualization of the flagellar structures in STM WT, STM *ΔpflB* (+/-F). Images are representative of N=2, n=5.

E. Normalized Median Fluorescence Intensity (MFI), i.e.., MFI of Secondary antibody for Rabbit generated Anti-fli antibody/ MFI for DAPI in confocal microscopy assisted visualization of the flagellar structures in STM WT, STM *ΔpflB* (+/-F). Data is represented as Mean+/-SEM of N=2, n=5.

F. Swim motility of STM WT and STM *ΔpflB* on 0.25% Agar containing LB plates

G. Percent adhesion of STM WT, STM *ΔpflB*, STM *ΔfliC*(+/-F) in Caco-2 cell line. Data is represented as Mean+/-SEM of N=5, n=3.

H. Percent invasion of STM WT, STM *ΔpflB*, STM *ΔpflB*/pQE60:*pflB* (+/-F) in Caco-2 cell line. Caco-2 cell line. Data is represented as Mean+/-SEM of N=2, n=2

(Unpaired Student's t-test for column graphs, Two-way Anova for grouped data, Mann-Whitney U-test for animal experiment data (**** $p < 0.0001$, *** $p < 0.001$, ** $p<0.01$, * $p<0.05$))

Figure 3: STM *ΔpflB* had a higher intracellular pH and higher membrane depolarization, both of which were recovered upon formate supplementation

A. BCECF-AM dye mediated elucidation of the relative ipH in the logarithmic cultures of STM WT, STM *ΔpflB* (+/-F) in phosphate buffers of pH 5.5, 6, and 7. Data is represented as Mean+/-SEM of N=4, n=4.

B. Elucidation of the ipH in the logarithmic cultures of STM WT/pBAD-pHuji and STM *ΔpflB*/pBAD-pHuji (+/-F). Data is represented as Mean+/-SEM of N=4, n=3.

C. Bisbenzimide assay to quantify the membrane damage in the logarithmic cultures of STM WT, STM *ΔpflB*, STM *ΔfliC* (+/-F). Data is represented as Mean+/-SEM of N=6, n=5.

D. Representative FACS percent positive and histogram profiles of $DiBAC_4$ staining in the logarithmic cultures of STM WT, STM *ΔpflB*, STM *ΔfliC* (+/-F). Data is representative of N=4, n=3.

E. Flow cytometry obtained quantification of percent- $DiBAC_4$ positive cells in the logarithmic cultures of STM WT, STM *ΔpflB*, STM *ΔfliC* (+/-F) incubated in LB media (pH 7.2) and F-media (pH 5.5). Data is represented as Mean+/-SEM of N=4, n=3.

(Unpaired Student's t-test for column graphs, Two-way Anova for grouped data, Mann-Whitney U-test for animal experiment data (**** $p < 0.0001$, *** $p < 0.001$, ** $p<0.01$, * $p<0.05$))

Figure 4: STM *ΔpflB* had a higher transcript level of *csrB*, raising the possibility of flagella inhibition via the *csr* sRNA system

A. RT-qPCR mediated expression profile of the flagellar operon genes *flhD*, *flgA*, *flgK*, *fliA*, *motA* in the logarithmic cultures of STM WT, STM *ΔpflB (+/-F)*. Data is represented as Mean+/-SEM of N=3, n=3.

B. RT-qPCR mediated expression profile of the sRNA *csrB* in the logarithmic cultures of STM WT, STM *ΔpflB (+/-F)*. Data is represented as Mean+/-SEM of N=6, n=3.

C. RT-qPCR mediated expression profile of the *fliC* in the logarithmic cultures of STM WT and STM *ΔpflB* transformed with an pQE60*: csrA* under an IPTG-inducible system. Data is represented as Mean+/-SEM of N=3, n=3.

D. Confocal microscopy assisted visualization of the flagellar structures in STM WT/pQE60: *csrA*, STM *ΔpflB*/pQE60: *csrA*. Images are representative of N=2, n=10.

E. Normalized Median Fluorescence Intensity (MFI), i.e.., MFI of Secondary antibody for Rabbit generated Anti-fli antibody/ MFI for DAPI in confocal microscopy assisted visualization of the flagellar structures in STM WT/pQE60: *csrA*, STM *ΔpflB*/pQE60: *csrA*. Data is represented as Mean+/-SEM of N=2, n=10

F. Percent adhesion of STM WT/pQE60: *csrA* and STM *ΔpflB*/pQE60: *csrA* in Caco-2 cell line. Data is represented as Mean+/-SEM of N=3, n=3.

G. TEM assisted visualization of the flagellar structures in STM WT/pQE60: *csrA*, STM *ΔpflB*/pQE60: *csrA*. Data is representative of N=2, n=10.

(Unpaired Student's t-test for column graphs, Two-way Anova for grouped data, Mann-Whitney U-test for animal experiment data (**** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$))

Figure 5: The extra cytoplasmic sigma factor RpoE increases the expression of *csrB* and therefore restricts the expression of *fliC*

A. RT-qPCR mediated expression profile of the sRNA *csrB* in the logarithmic cultures of STM WT, STM *ΔpflB*, STM *ΔrpoE*, STM *ΔpflBΔrpoE*. Data is represented as Mean+/-SEM of N=4, n=3.

B. RT-qPCR mediated expression profile of the flagellar operon genes *flhD* and *fliC* in the logarithmic cultures of STM WT, STM *ΔpflB*, STM *ΔrpoE*, STM *ΔpflBΔrpoE*. Data is represented as Mean+/-SEM of N=2, n=3.

C. Confocal microscopy assisted visualization of the flagellar structures in STM WT, STM *ΔpflB*, STM *ΔrpoE*, STM *ΔpflBΔrpoE*. Images are representative of N=2, n=10.

D. Normalized Median Fluorescence Intensity (MFI), i.e.., MFI of Secondary antibody for Rabbit generated Anti-fli antibody/ MFI for DAPI in confocal microscopy assisted visualization of the flagellar structures in STM WT, STM *ΔpflB*, STM *ΔrpoE*, STM *ΔpflBΔrpoE*. Data is represented as Mean+/-SEM of N=2, n=10.

E. TEM assisted visualization of the flagellar structures in STM WT, STM *ΔpflB*, STM *ΔrpoE*, STM *ΔpflBΔrpoE*. Data is representative of N=2, n=10.

F. Swim motility of STM WT, STM *ΔpflB*, STM *ΔrpoE*, STM *ΔpflBΔrpoE* on 0.25% Agar containing LB plates

G. Percent adhesion of STM WT, STM *ΔpflB*, STM *ΔrpoE*, STM *ΔpflBΔrpoE* in Caco-2 cell line. Data is represented as Mean+/-SEM of N=2, n=3.

(Unpaired Student's t-test for column graphs, Two-way Anova for grouped data, Mann-Whitney U-test for animal experiment data (**** $p < 0.0001$, *** $p < 0.001$, ** $p<0.01$, * $p<0.05$))

Figure 6: PflB maintains the switch between the flagellar expression and SPI-1 gene expression in intestine of the C57BL/6

A. RT-qPCR mediated expression profile of the SPI-1 genes *hilA* and *prgH* in the logarithmic cultures of STM WT, STM *ΔpflB*, STM *ΔrpoE*, STM *ΔpflBΔrpoE*. Data is represented as Mean+/-SEM of N=4, n=3.

B. RT-qPCR mediated expression profile of the SPI-1 genes *hilA* and *prgH* in the logarithmic cultures of STM WT, STM *ΔpflB*, STM *ΔsirA*, STM *ΔpflBΔsirA*. Data is represented as Mean+/-SEM of N=4, n=3.

C. RT-qPCR mediated expression profile of *fliC* gene in the logarithmic cultures of STM WT, STM *ΔpflB*, STM *ΔrpoE*, STM *ΔpflBΔrpoE*. Data is represented as Mean+/-SEM of N=3, n=3.

D. RT-qPCR mediated expression profile of *fliC* gene in the logarithmic cultures of STM WT and STM *ΔpflB* supplemented with 70, 90, 110, 150 mM of sodium formate. Data is represented as Mean+/-SEM of N=3, n=3.

E. GC-MS mediated quantification of relative intracellular formate in STM WT, STM *ΔpflB*, supplemented with 40, 90, 110 mM Sodium formate. Data is represented as Mean+/-SEM of N=4, n=2

F. RT-qPCR mediated expression profile of *pflB* gene in the duodenum and ileum of the intestine of C57BL/6 mice. Data is represented as Mean+/-SEM of N=3, n=3.

G. Confocal microscopy assisted visualization of the SPI-1 effector SipC protein in the duodenum and ileum of mice infected with STM WT and STM *ΔpflB*. Images are representative of n=7.

H. Confocal microscopy assisted visualization of the SPI-1 effector SipC protein in the duodenum and ileum of mice infected with STM WT and STM *ΔpflB*. Images are representative of n=7.

1155

1156 (Unpaired Student's t-test for column graphs, Two-way Anova for grouped data, Mann-
1157 Whitney U-test for animal experiment data (**** p < 0.0001, *** p < 0.001, ** p<0.01, *
1158 p<0.05))

1159

Supplementary Figure 1: STM *ΔpflB*, and STM *ΔfocA* are not growth deficient in LB
Broth, M9 Minimal Media. Deletion of pflB caused an enhanced expression of *focA*

A. Growth curves of STM WT, STM *ΔpflB* and STM *ΔfocA* in LB broth, M9 Minimal
Media. Data is represented as Mean+/-SEM of N=2, n=2.
B. Growth curves of STM WT, STM *ΔpflB* and STM *ΔfocA* in M9 Minimal Media. Data
is represented as Mean+/-SEM of N=2, n=2.
C. RT-qPCR mediated expression profile of *pflB* gene in the logarithmic cultures of STM
ΔfocA. Data is represented as Mean+/-SD of n=3.
D. RT-qPCR mediated expression profile of *focA* gene in the logarithmic cultures of STM
ΔpflB. Data is represented as Mean+/-SD of n=3.

(Unpaired Student's t-test for column graphs, Two-way Anova for grouped data, Mann-
Whitney U-test for animal experiment data (**** p < 0.0001, *** p < 0.001, ** p<0.01, *
p<0.05))

1173

1174 **Supplementary Figure 2: Intraperitoneal infection of STM WT and STM *ΔpflB* lead to**
1175 **no significant change in the bacterial burden in blood, liver, and spleen 3 dpi**

1176 A. Schematic showing the protocol followed for determining the organ burden of STM WT
1177 and STM *ΔpflB* 3 days post intraperitoneal infection

B-D. Bacterial burden of STM WT and STM *ΔpflB* in blood (B), liver (C), and spleen. Data is represented as Mean+/-SEM of N=2, n=5.

(Unpaired Student's t-test for column graphs, Two-way Anova for grouped data, Mann-Whitney U-test for animal experiment data (**** $p < 0.0001$, *** $p < 0.001$, ** $p<0.01$, * $p<0.05$))

Supplementary Figure 3: Formate modulates the expression of *fliC* and *hilA* in STM WT and STM *ΔpflB* in a concentration dependent manner.

A. Expression of *hilA* in STM WT in formate supplemented at concentrations of 10, 20, 30, 40, and 50 mM. Data is representative of N=2, n=3 and expressed as Mean+/- SD.

B. Expression of *fliC* in STM WT and STM *ΔpflB* in formate supplemented at concentrations of 0.067, 0.125, 0.5, 1, 10 mM. Data is represented as Mean+/-SEM of N=4, n=3.

C. Expression of *hilA* in STM WT and STM *ΔpflB* in formate supplemented at concentrations of 0.067, 0.125, 0.5, 1, 10 mM. Data is represented as Mean+/-SEM of N=4, n=3.

(Unpaired Student's t-test for column graphs, Two-way Anova for grouped data, Mann-Whitney U-test for animal experiment data (**** $p < 0.0001$, *** $p < 0.001$, ** $p<0.01$, * $p<0.05$))

Supplementary Figure 4: STM *ΔfocAΔpflB* had an attenuated flagellar apparatus, with partial recovery upon supplementation of extragenous F

A. Growth curves of STM WT, STM $\Delta focA\Delta pflB$ in LB broth and M9 Minimal Media. Data is represented as Mean+/-SEM of N=2, n=2.

B. Growth curves of STM WT, STM $\Delta pflB$ and STM $\Delta focA$ in M9 Minimal Media. Data is represented as Mean+/-SEM of N=2, n=2.

C. Confocal microscopy assisted visualization of the flagellar structures in STM WT, STM $\Delta focA\Delta pflB$ (+/-F). Images are representative of N=2, n=5.

D. Normalized Median Fluorescence Intensity (MFI), i.e.., MFI of Secondary antibody for Rabbit generated Anti-fli antibody/ MFI for DAPI in confocal microscopy assisted visualization of the flagellar structures in STM WT, STM $\Delta focA\Delta pflB$ (+/-F). Data is represented as Mean+/-SEM of N=2, n=5.

E. Percent adhesion of STM WT and STM $\Delta focA\Delta pflB$ in Caco-2 cell line. Data is represented as Mean+/-SEM of N=5, n=3.

F. Organ burden of STM WT and STM $\Delta focA\Delta pflB$ in liver and spleen of C57BL/6 mice 5 days post oral gavaging. Data is represented as Mean+/-SEM of N=2, n=5.

(Unpaired Student's t-test for column graphs, Two-way Anova for grouped data, Mann-Whitney U-test for animal experiment data (**** $p < 0.0001$, *** $p < 0.001$, ** $p<0.01$, * $p<0.05$))

Supplementary Figure 5: Decreased expression of *fliC* in STM $\Delta pflB$ can be altered upon incubating the bacterial cultures in acidic media (pH of 6).

A. RT-qPCR mediated expression profile of *fliC* in the logarithmic cultures of STM WT, STM $\Delta pflB$ incubated in acidic LB media (pH 6). Data is represented as Mean+/-SEM of N=3, n=3.

B. RT-qPCR mediated expression profile of *csrA* in the logarithmic cultures of STM WT, STM *ΔpflB* incubated in acidic LB media (pH 6). Data is represented as Mean+/-SEM of N=3, n=3.

(Unpaired Student's t-test for column graphs, Two-way Anova for grouped data, Mann-Whitney U-test for animal experiment data (**** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$))

Supplementary Figure 6: Compensation of intracellular formate pool in STM *ΔpflB* makes it mimic the behaviour of STM WT

A. Confocal microscopy assisted visualization of the flagellar structures in STM WT, STM *ΔpflB* supplemented with 40, 90, 110 mM of sodium formate. Images are representative of N=3, n≥5.

B. Normalized Median Fluorescence Intensity (MFI), i.e.., MFI of Secondary antibody for Rabbit generated Anti-fli antibody/ MFI for DAPI in confocal microscopy assisted visualization of the flagellar structures in STM WT, STM *ΔpflB* supplemented with 40, 90, 110 mM of sodium formate. Data is represented as Mean+/-SEM of N=3, n≥5.

C. TEM assisted visualization of the flagellar structures in STM WT and STM *ΔpflB* supplemented with 40, 90, 110 mM of sodium formate. Data is representative of N=2, n=10.