Identifying a Distractor Produces Object-Based Inhibition in an Allocentric Reference Frame for Saccade Planning

Coleman E. Olenick[1*], Heather Jordan[1], Mazyar Fallah[1,2]

1. Department of Human Health and Nutritional Sciences, College of Biological Science, University of Guelph, Guelph, Ontario N1G 2W1, Canada
2. Canadian Action and Perception Network, Canada

Abstract:

We investigated whether distractor inhibition occurs relative to the target or fixation in a

perceptual decision-making task using a purely saccadic response. Previous research has shown

that during the process of discriminating a target from distractor, saccades made to a target

deviate towards the distractor. Once discriminated, the distractor is inhibited, and trajectories

deviate away from the distractor. Saccade deviation magnitudes provide a sensitive measure of

target-distractor competition dependent on the distance between them. While saccades are

planned in an egocentric reference frame (locations represented relative to fixation), object-based

inhibition has been shown to occur in an allocentric reference frame (objects represented relative

to each other independent of fixation). By varying the egocentric and allocentric distances of the

target and distractor, we found that only egocentric distances contributed to saccade trajectories

shifts towards the distractor during active decision-making. When the perceptual decision-

making process was complete, and the distractor was inhibited, both ego- and allocentric

distances independently contributed to saccade trajectory shifts away from the distractor. This is

consistent with independent spatial and object-based inhibitory mechanisms. Therefore, we

suggest that distractor inhibition is maintained in cortical visual areas with allocentric maps

which then feeds into oculomotor areas for saccade planning.

Introduction

When I view a tabletop full of coffee cups, I can locate my cup relative to myself in a viewpoint-dependent, egocentric reference frame. When I want to reach for my cup, there is considerable evidence that both my eye- and hand-movements are programmed using an ego-centric spatial map of the scene. However, if I want to avoid knocking over my colleagues' cups, I need to direct my actions towards my cup while considering the locations of the non-target cups. This requires an allocentric (object-to-object) spatial map of the scene, where objects are represented in a viewpoint-independent reference frame. There is converging evidence from behavioral, electrophysiological, and imaging studies that we use both viewpoint-dependent egocentric, and viewpoint-independent allocentric reference frames in spatial cognitive tasks using reaching[1][2][3][4].

When there are multiple objects in the visual scene, attending to and acting on, relevant stimuli may require the suppression of irrelevant stimuli competing for attentional selection. Yet there is no clear delineation about the reference frames supporting this suppression. In models of attention based on visual processing for example, the biased competition and selective tuning models, the suppression of irrelevant stimuli changes in strength based on the proximity to an attended stimulus[5][6]. However, oculomotor models of spatial attention invoke regions encoding relevant and irrelevant object locations in retinotopic coordinates, notably the superior colliculus (SC) and frontal eye fields (FEF)[7][8][9].

The receptive fields of oculomotor regions also receive inputs in non-retinotopic reference frames, for example eye position-based gain modulations of the FEF[10][11]. There is also extensive work on the study of gain fields in parietal regions representing multiple reference frames[12][13] that feed into the FEF and SC. While these inputs provide a source of non-

retinotopic information, their effects are largely modelled as multiplicative gain factors on an underlying retinotopic map giving rise to the oculomotor plans.

Recently, saccadic eye movement trajectories have been shown to reflect the continuous competition resolution between objects[14]. Specifically, at short object processing times, before the target object has been differentiated from a distractor and decision making is still actively taking place, saccades deviate towards the distracting object. However, once the target and distractor are differentiated after greater processing time, and decision making has completed, the saccades deviate away from the distractor. The directional deviation of a saccade can be used to differentiate between a distractor or location that has been inhibited and one that has not, for a given single saccade[15-26]. Saccades are planned in an egocentric reference frame on a retinotopic map of space[27] and activity on this map has been shown to cause saccade trajectory deviations[28-30]. The deviation towards a distractor is due to increased activity at the location of the distractor, whereas suppression at the distractor location leads to deviation away[25][29-30]. Because saccade planning is egocentric[31-33], it is assumed that the suppression of distractors would also be egocentric.

However, behavioral studies in humans have shown that object-based suppression can also occur in allocentric reference frames. For example, object-based inhibition of return (IOR) has been shown to occur in an allocentric reference frame[34-37]. Both trajectory deviations and IOR can occur under the same conditions but do so on different timescales[38]. The complex objects used by Giuricich, and colleagues[14] required late-stage visual processing to discriminate target from distractor[39-40]. These late-stage ventral stream areas (e.g. TE, TEO, IT cortex) support object representations in allocentric reference frames[41], likely resulting from large, bilateral receptive fields with position and size invariance[41-43]. So, if saccade planning occurs in an egocentric

reference frame, but the resolution of object competition is dependent on regions with allocentric

representations and both involve inhibition of a distractor, what is the reference frame of the

resulting distractor suppression?

In this study we independently varied the egocentric and allocentric distance of a target and

distractor to investigate the reference frame of distractor suppression. The egocentric distance of

the target and distractor is their eccentricity, while the allocentric distance describes the distance

between the target and distractor or target-distractor distance (T-D distance), which is

independent of the viewpoint or eye position. Varying the eccentricity of the target and distractor

while keeping the angle between them constant, leads to a range of T-D distances. Using saccade

trajectories to infer the state of decision-making, we identified when the distractor has been

suppressed through the resulting deviation away from the distractor. The magnitude of the

saccade deviation is associated with the strength of the distractor-related activity as has been

demonstrated through both recording and stimulation studies of the SC[28-29] and FEF[30]. The

amount of deviation at different egocentric and allocentric distances will reflect the reference

frame of distractor activation and suppression. As the oculomotor system encodes saccades to a

target in an egocentric reference frame, we expect to find egocentric reference frame effects on

saccade trajectories both during active competition before the decision is made and after, when

the distractor is suppressed. However, if distractor suppression is dependent on object-based

inhibition developing in cortical visual areas, once the competition is resolved, saccade

trajectories should also be shifted based on T-D distance in an allocentric reference frame.

Methods

Participants

Forty naïve participants (8 males; age: 18-27, mean = 19.6 years old) were recruited for the

current experiment. All participants had normal or corrected to normal vision, gave written

informed consent, and were compensated with $15 CAD for their time. Ethics approval of the

experiment was obtained from the Research Ethics Board at the University of Guelph.

Figure 1: Experimental Overview A) The stimuli were selected from 2 sets of 6 objects, each

2° X 2°. The target and distractor on a trial were drawn from the same set, with each object

varying by one line addition or removal, relative to adjacently numbered objects in the set. The

objects were designed to avoid resemblance to any English characters or numbers to ensure their

novelty to participants. B) The objects were presented at 15 possible locations at 3 eccentricities

(depicted in color for clarity in the figure only: 7 [red], 10 [green], and 13 [blue] dva from

fixation), and 5 branches separated by 45° in the upper half of the display. The target and

distractor were always objects from the same set and presented along adjacent branches. C)

Trials began with participants fixating a central cross, which was replaced with a preview of the

target for that trial. After participants pressed a button, they were required to again fixate a cross

for 200ms before the target and distractor appeared simultaneously. The objects remained on

screen until either a saccade was made to the target or distractor or 750ms had elapsed. Trials

without a response in 750ms were presented again at a later point in the block, and there were

some trials where only the target was presented to establish a saccade baseline. D) The sum

curvature of a saccade was calculated as the area under the curve between the actual saccade

trajectory and a path directly between its start and end points. The endpoint angle of a saccade

was calculated as the polar angle between the saccade endpoint and the center of the target

object. This example saccade is showing deviation away from the distractor direction and would

be negative. See methods for more detail.

Stimuli and Apparatus

Participants were seated in a dimly lit room 57cm away from a 21" CRT monitor (60Hz,

1024×768), with movement reduced by use of a chin and head rest. Stimulus presentation and

data collection was controlled by Presentation software (Neurobehavioral Systems, Inc.,

Berkeley, CA), and participant responses were made using a response box (RB-540, Cedrus

Corp., San Pedro, CA) and eye movements. The eye movements of participants were recorded

from their right eye using an Eyelink II infrared eye tracker (500Hz; SR Research, ON, Canada).

The target and distractor stimuli were white (CIExy = [.29, 0.30]; luminance = 126.02cd/m^2) on

a black (CIExy = [.27, 0.26]; luminance = 0.20cd/m^2) background and were randomly selected

from a set previously used[14][25][39] (Figure 1a). Each stimulus consisted of 6- or 7-line segments

(1° × 0.08°). The stimuli resemble novel alphanumeric characters, requiring similar processing in

late-stage ventral stream areas (e.g. TE, TEO, IT cortex)[40][44].

Procedure

Participants completed a delayed match to sample task, making saccades to a previously shown

target presented with a distractor (Figure 1c). Eye position was calibrated at the beginning of the

experiment, at the start of each block, and additionally if necessary. Each trial began with a

fixation cross (0.4° X 0.4°; CIExy = [.x=29, y=0.30]; 126.02cd/m^2) at the center of the screen.

Following 200 ms fixation, the target object was presented at the centre of the display and

remained until the participant pressed a designated button on a response box. The target object

was replaced by a fixation cross and the participant had to maintain fixation within a 2X2 dva

square window centered on the cross for 200 ms or the trial was terminated. The fixation marker

remained on the display while the target and a distractor object appeared simultaneously for 750

ms or until the participant made a saccade to one of the objects. Participants received a written

error message and audible tone if they failed to make a saccade or made a saccade to the

distractor object; error trials were presented later in within the same block.

There were 15 potential positions for the objects, organized into 5 different branches separated

by 45°, and placed at egocentric eccentricities of 7, 10, or 13 dva (Figure 1b). The various

eccentricities for the target and distractor yield a range of allocentric distances between them (T-

D distance), ranging from 5.36 dva to 9.95 dva (5.36, 7.07, 7.65, 9.22, 9.45, and 9.95 dva). The

target and distractor were always 45° apart, an angular distance previously investigated that has

been shown to provide reliable saccade trajectory deviations[14]. Within each block, each

combination of target and distractor eccentricities appeared with equal probability (3x3

combinations) and were evenly distributed across the array of possible positions (Figure 1b),

resulting in 72 distinct target-distractor pairings (trials), pseudorandomly interleaved across the

block. Note that all objects were presented on or above the horizontal meridian and thus when

the target or distractor appeared on a horizontal branch, the other object necessarily appeared 45°

above the meridian. However, this was not predictive as an object appearing on the horizontal

meridian had equal probability of being the target or the distractor across the block of trials.

Additionally, each block contained target-only trials at each of the 15 object positions, acting as

a baseline, producing a total block size of 87 trials. The objects used as target and distractor were

always from the same set of the two sets used in the experiment (Figure 1a) but varied in their

similarity to one another. This range of stimuli were used to increase the complexity of the task

minimizing perceptual learning. Each participant completed 5 experimental blocks (435 total

trials) and a 10-trial practice block presented at the beginning of the experiment that was not

included in analysis.

Data Analysis

The investigation of the effects of eccentricity made use of multiple regression across target and

distractor eccentricities. To differentiate an allocentric reference frame from an egocentric

reference frame, we a priori determined to investigate both trials with equal target and distractor

eccentricity, and trials with equal T-D distance but flipped target and distractor eccentricities

(e.g. target at 7 dva/distractor at 13 dva vs target at 13 dva/distractor at 7dva). All multiple

comparisons tests and regression analyses have been subject to Benjamini-Hochberg corrections

for multiple comparisons

Saccade Detection: Saccades were detected using in-house MATLAB (Mathworks, Natick, MA)

scripts following pre-determined criteria; an eye movement with a velocity greater than 20°/s for

at least 8 ms with a peak velocity greater than 50°/s. Saccades were evaluated manually to

remove trials where the first saccade was not made to the target or distractor (9.59%). The

saccades not made to the target or distractor were a mix of different outcomes: short saccades

where the intended target of a saccade could not be determined, saccades that were initially made

in the direction of one of the two objects but not reaching it before making a second saccade (a

double-step saccade), and saccades that were partially occluded by an eye blink. Trials with short

(amplitude < 1° [0.65%]) or express saccades (Saccade Reaction Time (SRT) < 100 ms [1.97%])

or where no eye position data was recorded (3.64%) were excluded from further analysis.

Overall, 84.15% of recorded trials were included for analysis.

Accuracy: Accuracy was calculated for each participant as the proportion of trials that a saccade

was made to the target rather than the distractor. A 4X4 dva box around the target or distractor

was used to determine which object was selected for a given saccade. All participants'

performance exceeded chance (binomial distribution with p = 0.5 [50%]) with 65.2 ± 8.9%

correct. *Saccade Trajectory Metrics*: To quantify the deviation of a saccade from a straight path,

the sum curvature and endpoint angle for each saccade was calculated. The sum curvature

provides a measure of the strength of target and distractor representations in the cortical saccade

planning process[17-18][30][45]. Sum curvature was calculated as the total deviation of a saccade

from a straight line between its start point and endpoint (reference line). That is, the magnitude

of deviation from the reference line and direction (sign) of each eye position sample for a

saccade were summed to generate a single metric for each saccade. The endpoint angle of a

saccade is suggested to result from the outcome of a subcortical vector average of saccade plans

to both the target and distractor[44-45]. The endpoint angle was calculated as the polar angle (in

degrees) between the reference line and a straight path from the saccade start point to the center

of the target. The reference line was translated prior to computation such that the start of each

saccade was on the cartesian origin to account for small variations in initial fixation location.

Both metrics were normalized by subtracting the mean values from target only trials and

corrected such that positive values reflect deviations towards the distractor and negative values

reflect deviations away from the distractor.

Processing Time Division: Following Giuricich and colleagues[14], trajectory metrics were used

to separate saccadic responses made between actively competing and inhibited distractor stages

as these different processes may not rely on the same reference frame(s). When two items are

potential saccade targets, a saccade made to one has its trajectory shifted towards the other. As

an actively competing distractor in a perceptual discrimination is a potential saccade target, this

results in saccade trajectories made to the target shifting towards the distractor[15-18][21-24][29-30]. In

contrast, with enough time the discrimination decision-making is completed, wherein the target

is identified and the distractor is inhibited. An inhibited distractor results in saccade trajectories

shifting away from it[19-20][25-26][29-30]. The change in saccade deviation direction over time has

been used to separate saccades executed at different stages of visual processing [14-26][29-30][45]. To

determine the range of saccade latencies falling within each stage (actively competing vs

inhibited distractor), sum curvature was organized by saccade latency and a 10 ms width sliding

window was used to create a series of binned saccades. Within each of these bins, a two-tailed t-

test against 0 was performed to determine whether there was significant overall curvature

towards or away from the distractor at each saccade latency (Figure 2b).

Figure 2 Raw Saccade Trajectories and Deviation Direction Over Processing Time. A) 80
random saccade traces to targets at 7, 10, and 13 dva (from left to right, respectively), rectified to
a distractor appearing on the right, but uncorrected for saccade start position, are plotted. Black
squares represent the center of the intended targets (note that the actual targets had a width of 2
dva), and the traces are color-coded by object processing time; red and blue reflecting near
100ms and 350ms respectively. B) The average sum curvature of saccades across all participants
has been plotted over saccade latency. A 10 ms sliding window over saccade latency was used to
group saccades and the average sum curvature within each bin is represented by the shaded grey
points, with the solid line reflecting a 10 ms moving average over the shaded points. Positive

values indicate deviation towards the distractor, where the distractor is still actively competing

with the target, while negative values indicate deviations away after the distractor has been

inhibited. The red points along the bottom represent the center of saccade latency bins where

saccades showed significant curvature towards or away from the distractor (different from 0).

The blue shaded region represents a transition between an active and inhibited distractor that has

been omitted from further analysis.

Results

Prior to analysis of the task performance and saccade trajectory metrics, the saccades were

divided into two categories: those in which the distractor was actively competing with the target

and those after it had been inhibited. This was accomplished by examining the saccadic sum

curvature over object processing time (Figure 2b). By using a sliding 10 ms window to group

saccades by object processing time and performing t-tests against 0 on sum curvature within each

window, we identified time ranges with consistent significant deviation towards the actively

competing distractor (100-196 ms) or away from the inhibited distractor (252-750 ms). In total,

there were 10407 saccades outside the transition zone; 62.5% at short processing time and 37.5%

at long processing times. These ranges are consistent with prior studies[14][25][39][46]. Saccades

whose latencies fell in the transition zone between these 2 ranges were excluded from further

analysis. It is important to note that not all participants had data in all combinations of

experimental conditions and saccadic response times, making the use of repeated measures

ANOVA impractical. Our analysis instead uses mixed-effects regression models as a well-

established alternative[47-49] for the purposes of the current study.

Accuracy

In an egocentric reference frame, increasing the distance between fixation and the target would

be expected to have the opposite effect as increasing the distance between fixation and the

distractor, due to the central bias in attention (where attention and visual processing is stronger

closer to fixation and drops off with distance) as shown by other visual search paradigms[50-52].

The central bias predicts that as the distance between the target and fixation increases, accuracy

decreases, but as the distance between distractor and fixation increases, accuracy improves. To

investigate the relationship between an egocentric reference frame and accuracy over processing

times, a multiple linear regression with target and distractor eccentricity, and processing time

(short and long) as factors was performed. The overall regression was found to be highly

significant ($R^2 = 0.641$, $F(7,680) = 173.2$, $p < 6.4\text{X}10^{-15}$). Increasing target eccentricity was

found to decrease accuracy, while increasing distractor eccentricity increased accuracy (Figure

3). The effect of both target and distractor eccentricity significantly decreased from short to long

processing time. There was a significant interaction between target and distractor eccentricity

that did not change with processing time, suggesting a relationship between object positions

beyond their individual eccentricities, that will be explored in the next section. Finally, there was

no change in accuracy across processing times independent from the other relationships

mentioned above. However, a Wilcoxon signed rank test across each participant revealed a

significant increase in accuracy at longer processing times ($p = 3.19\text{X}10^{-15}$).

Figure 3: Accuracy by Target and Distractor Eccentricity. The average accuracy plotted over

target and distractor eccentricity. A) Accuracy at short processing times while competition

between target and distractor is still active. B) Accuracy at long processing times when

competition has been resolved and the distractor had been inhibited. Error bars represent mean ±

95% confidence intervals. The bars and lines are colored according to the target eccentricity of 7

dva (red), 10 dva (green), and 13 dva (blue).

To identify any effects of an allocentric reference frame on accuracy, a multiple linear regression

with processing time and target-distractor distance (T-D distance) was performed on the subset

of trials in which the target eccentricity and distractor eccentricity were equal, i.e. equal

egocentric distances (Table 1; Figure 4 B, C). When the target and distractor were both presented

at an eccentricity of 7, 10, or 13 dva, the T-D distances were 5.36, 7.65, and 9.95 dva,

respectively. The regression across T-D distance was significant overall, and robust across

participants (R^2: 0.332, $F(3,224) = 37.14$, $p < 2.13\text{X}10^{-15}$). While overall accuracy increased

significantly from short to long processing time (from regression: $p = 0.0156$), T-D distance had

no influence on accuracy at short processing times. Critically, accuracy varied by T-D distance at

long processing times, when the distractor is inhibited, decreasing significantly as the T-D

distance increased.

Figure 4: Accuracy by Target-Distractor (T-D) Distance Pairs. A) Example of target and

distractor positions that have an equal distance between them (T-D distance) but either the target

is closer to fixation (orange), or the distractor is closer to fixation (purple). Accuracy for targets

and distractors presented at the same eccentricity but varying T-D distances plotted for B) short

processing times and C) long processing times. The average accuracy plotted over T-D distance

pairs for D) short processing times, and E) long processing times. Error bars represent mean ±

95% confidence intervals. Note that the pairs of orange and purple dots in panels B & C are

offset slightly for visibility and that the actual T-D distances for each target closer and distractor

closer pair are: 7.07, 9.22, and 9.45 dva from left to right. Pairwise comparisons between the

allocentric distance pairs are indicated by n.s: not significant, *: $p < 0.05$ ***: $p < 0.001$.

Table 1: Regression Coefficients, and p-values for Accuracy and Trajectory Metrics Over

Processing Time. Each cell consists of the regression coefficient and its p-value: *: $p < 0.05$, **:

$p < 0.01$, ***: $p < 0.001$, n.s.: not significant. Note: the values reported for accuracy regressions

were determined through separate linear regressions for short and long processing time after

differences had been established. ŧ is used to indicate coefficients that were not significantly

different between processing times. The T-D distance regression was performed only on the

points where target and distractor eccentricities were equal. The eccentricity interaction reflects a

changing effect for the increase of target or distractor eccentricity as the eccentricity of the other

object increases.

	Regression Predictor	Accuracy	Sum Curvature	Endpoint Angle
Short Processing Time	Target Eccentricity	-0.139 $< 2\text{x}10^{-16}$ ***	+1.040 $1.1\text{x}10^{-4}$ ***	-1.392 $7.2\text{x}10^{-5}$***
	Distractor Eccentricity	+0.035 $5.5\text{x}10^{-3}$ **	+0.039 0.8494 n.s.	-2.234 $< 2\text{x}10^{-16}$***
	Eccentricity Interaction	+0.005 $3.11\text{x}10^{-5}$ ***	-0.036 0.1282 n.s.	0.153 $5.5\text{x}10^{-7}$ ***
	T-D Distance	+0.004 0.521 n.s.	+0.446 $1.6\text{x}10^{-4}$ ***	-0.915 $5.3\text{x}10^{-12}$ ***

Long Processing Time	Target Eccentricity	-0.076 4.2x10⁻⁴ ***	+0.168 0.4668 n.s.	-0.316 0.1061 n.s.
	Distractor Eccentricity	-0.015 0.4843 n.s.	+0.261⁺ 0.2407 n.s.	-0.442 0.0206 *
	Eccentricity Interaction	0.004⁺ 0.0671 n.s.	-0.047⁺ 0.0313 n.s.	+0.040 0.0338 *
	T-D Distance	-0.023 0.0104 *	-0.669 2.3x10⁻⁸ ***	+0.074 0.511 n.s.

The other method we employed to differentiate egocentric and allocentric coding was comparing

conditions where the T-D distance between the target and distractor was identical, but their

eccentricities varied Using paired t-tests (Figure 4 D, E), we see significant differences between

eccentricities with the target or distractor closer to fixation at both processing times (all $p <$

1.933×10^{-3}). However, the differences between the target closer and distractor closer conditions

decreased substantially, indicating a reduction in the influence of an egocentric reference frame.

Taken together, the reduction in center bias and variation in accuracy across T-D distance shown

in the significant regression in the previous section at long processing times indicate that

accuracy is driven by object distance in both egocentric and allocentric reference frames.

Saccade Trajectory Deviations

To better understand the reference frames of target and distractor representations, we next

investigated their effects on saccade trajectory metrics, which provide a continuous and

separable measure of distractor activity and suppression. Separate multiple linear regressions

were performed on the saccade curvature and endpoint angle for short and long processing times

with target and distractor eccentricity as predictors (see Table 1; Figure 5).

At *short processing* time, both regressions were significant overall (Sum Curvature: $R^2 = 0.0449$,

$F(3,3658) = 57.39$, $p < 9.11 \times 10^{-16}$ | Endpoint Angle: $R^2 = 0.0598$, $F(3,3658) = 77.53$, $p <$

7.98×10^{-16}). At short processing time, where saccades deviate towards the distractor, only

increasing target eccentricity led to greater saccade curvatures, consistent with an egocentric

target representation whose strength diminishes with eccentricity. In contrast, the endpoint angle

decreased as the eccentricity of the target and the eccentricity of the distractor increased, again

consistent with an egocentric representation. A significant interaction effect between target and

distractor eccentricity on endpoint angle indicates that the relative effect of the eccentricity of an

object changes with the eccentricity of the other object, and may be suggestive of an allocentric

effect and will be followed-up on in later analyses.

Figure 5: Saccade Trajectory Metrics by Target and Distractor Eccentricity. The average

A) sum curvature and B) endpoint angle plotted over target and distractor eccentricity. Error bars

represent mean ± 95% confidence intervals. Values above 0 reflect saccade deviations towards a

distractor while the decision-making process is active. Values below 0 represent deviations away

from the distractor, after object competition has been resolved and the distractor has been

suppressed. The bars and lines are colored according to the target eccentricity of 7 dva (red), 10

dva (green), and 13 dva (blue).

At *long processing* times, regressions for sum curvature and endpoint angle were both small but

significant (Sum Curvature: $R^2 = 0.018$, $F(3,3176) = 19.43$, $p = 5.8 \times 10^{-12}$ | Endpoint Angle: $R^2 =$

0.0025, $F(3,2492) = 3.045$, $p = 0.0051$). Changes to the target eccentricity had no effect on the

sum curvature or endpoint angle. The distractor eccentricity was correlated with the endpoint

angle, but not the sum curvature. Interestingly, the interaction between target and distractor

eccentricity was significantly correlated with both trajectory metrics, again suggesting target-

distractor distance effects in an allocentric reference frame. Together, these results show a

decreased strength of egocentric object representations feeding target-distractor competition in

saccade planning after the decision-making process has been resolved.

To investigate the influence of an allocentric reference frame, linear regressions were performed

on T-D distance for both saccade metrics when the target and distractor eccentricity were equal

(Figure 6 A, B). In addition, pairwise t-tests were used to compare conditions where the T-D

distance of the objects was equal, but their eccentricities varied (Figure 6 C-F). At *short*

processing times, T-D distance had a significant effect on both the sum curvature ($R^2 = 0.0115$,

$F(1,1231) = 14.37$, $p = 2.9 \times 10^{-4}$), and endpoint angle ($R^2 = 0.0379$, $F(1,1231) = 48.53$, $p =$

1.54×10^{-11}). However, pairwise comparisons across equal T-D distance pairs also found

significant differences for both metrics (all $p < 0.0432$). Together these findings suggest that at

short processing times saccade trajectories are driven by egocentric rather than allocentric

representations.

Conversely, at *long processing* times T-D distance was only correlated with the sum curvature

($R^2 = 0.0277$, $F(1,1110) = 31.65$, $p = 6.06 \times 10^{-8}$), and not endpoint angle ($R^2 = 0.0005$, $F(1,880)$

$= 0.432$, $p = 0.5491$). Pairwise comparisons revealed no differences in sum curvature when target

and distractor eccentricities varied while the T-D distance was equal (all $p > 0.2441$), supporting

allocentric rather than egocentric effects. There were no effects for saccade endpoint angles

when T-D distance was equal across varying target and distractor eccentricities.

Figure 6: Target-Distractor (T-D) Distance. A & B: The average sum curvature and endpoint

angle plotted over T-D distance when the target and distractor have equal eccentricity. Values

above 0 reflect saccade deviations towards a distractor while the decision-making process is

active. Values below 0 represent deviations away from the distractor, after object competition

has been resolved and the distractor has been suppressed. C-F: The saccade metrics plotted over

T-D distance pairs with the same conventions as in Figure 3, where orange reflects the conditions

with the target presented closer to fixation and purple reflects trials with the distractor closer to

fixation. Error bars represent mean ± 95% confidence intervals. Note that the pairs are offset

slightly for visibility and that the actual T-D distance is between the 2 points. Significant

regression effects across T-D distance and comparisons within eccentricity pairs are indicated

with ***: $p < 0.001$.

Discussion

In this study, we varied the distance of target and distractor objects relative to a central fixation

point, and one another, in a saccadic delayed match to sample task to investigate the reference

frame of object-based suppression in the oculomotor system. We found that for short object

processing times, while the competition between target and distractor is active[14], both target and

distractor are represented in an egocentric reference frame. However, the completion of decision

making led to the emergence of distractor suppression, which occurred in an allocentric reference

frame.

Saccades executed at short processing times displayed a strong bias for selecting objects closer to

fixation (less eccentric). Additionally, the trajectory shifts of these saccades towards the

distractor were consistent with weighted vector averaging, as has been previously reported for

saccade deviations towards distracting objects[14-18][21-24][29-30][45]. Our results for both the saccade

trajectories and accuracy suggest that the weight of each object decreases linearly as its

eccentricity increases. The strength of this spatial bias alters saccade target selection independent

of the identity of the stimuli. Our results lend support to the idea that early saccades are based on

bottom-up salience rather than top-down relevance[53] which affects both accuracy and saccade

curvature[54]. The eccentricity gradient may result from either an inherent attentional bias[50-52], or

reflect a visually-based salience difference (i.e. cortical magnification). Both of these

possibilities reflect an egocentric reference frame early in the formation of object

representations.

In contrast, long processing times prior to saccade execution resulted in deviations away from

the distractor indicating that the distractor had been suppressed[19-20][25-26][29-30][45]. When the

stimuli are processed for longer prior to initiating the saccade, the target is identified and we find

that the distractor is suppressed in an allocentric reference frame. The magnitude of the curvature

is no longer affected by the eccentricity of the distractor. Instead, the magnitude of the curvature

increases as the distance between the target and distractor increases, reflecting a shift from an

egocentric to allocentric reference frame. However, there continues to be an effect of *target*

eccentricity at long processing times, indicating that an egocentric reference frame is still being

utilized. This arises from saccades to targets being programmed in egocentric coordinates[31-30].

Object-based suppression in an allocentric reference frame has been found in object-based

inhibition of return (IOR)[34-35], and to previous saccade locations[36-37][55-56]. The current work

extends these findings, identifying resolution of target/distractor competition as the initiator of

distractor suppression that occurs in an allocentric frame of reference.

The stimuli used in our study are similar to alphanumeric characters[40], which have previously

been suggested to require late-stage ventral stream visual processing in IT cortex[39-40] where

objects are represented allocentrically[41-43]. The transformation from a gaze-centered reference

frame to hand-centered to guide reaching has previously been shown to occur in the posterior

parietal cortex (PPC)[57-58], making it the most probable site for task-relevant reference frame

transitions(Andersen and Zipser, 1988)[12][59]. Therefore, this would suggest that object-based

distractor suppression arising in area IT[60] would be carried by projections to PPC[61]

subsequently feeding into the oculomotor system (i.e. SC, FEF). As PPC is associated with

multiple reference frames[12][59][62-64], it would be able to map allocentric object suppression onto

the gaze-centered egocentric reference frame used in saccade planning.

The simultaneous presence of multiple items necessitates discrimination of the target from a

distractor. The suppression of the distractor would provide an effective mechanism to improve

target selection and motor planning. Our finding that this suppression is allocentric highlights the

involvement of object-based distractor inhibition outside of the motor system influencing the

saccade planning process. We find that we can measure this allocentric object-based attentional

mechanisms in a temporally precise manner manifested through saccade curvatures.

In summary, object-based suppression occurs in an allocentric reference frame affecting target

selection and the subsequent motor action. The early representation of potential saccade targets is

egocentric and strongly weighted by eccentricity as they compete for selection, with a bias for

the object closest to the center of gaze. With longer processing time, the target is identified, and

the distractor is suppressed. The target continues to be represented in an egocentric reference

frame, while the strength of distractor suppression is dependent on target-distractor distance in an

allocentric reference frame. Object-based suppression likely is a general mechanism of visual

processing that affects motor planning, as seen here, as well as perceptual decision making, as

seen in prior studies.

References

1. Ball, K., Smith, D., Ellison, A. & Schenk, T. Both egocentric and allocentric cues support
spatial priming in visual search. *Neuropsychologia* **47**, 1585–1591 (2009).

2. Galati, G., Pelle, G., Berthoz, A. & Committeri, G. Multiple reference frames used by the
human brain for spatial perception and memory. *Experimental brain research* **206**, 109–
20 (2010).

3. Ruotolo, F. *et al.* Neural correlates of egocentric and allocentric frames of reference
combined with metric and non-metric spatial relations. *Neuroscience* **409**, 235–252
(2019).

4. Burgess, N. Spatial memory: how egocentric and allocentric combine. *Trends in*
Cognitive Sciences **10**, 551–557 (2006).

5. Tsotsos, J. K. *et al.* Modeling visual attention via selective tuning. *Artificial Intelligence*
78, 507–545 (1995).

6. Desimone, R. & Duncan, J. Neural Mechanisms of Selective Visual Attention. *Annu. Rev.*
Neurosci. **18**, 193–222 (1995).

7. Moore, T., Armstrong, K. M. & Fallah, M. Visuomotor Origins of Covert Spatial
Attention. *Neuron* **40**, 671–683 (2003).

8. Moore, T. & Fallah, M. Control of eye movements and spatial attention. *Proceedings of*
the National Academy of Sciences **98**, 1273–1276 (2001).

9. Moore, T. & Fallah, M. Microstimulation of the Frontal Eye Field and Its Effects on
Covert Spatial Attention. *Journal of Neurophysiology* **91**, 152–162 (2004).

10. Balan, P. F. & Ferrera, V. P. Effects of Gaze Shifts on Maintenance of Spatial Memory in
Macaque Frontal Eye Field. *J. Neurosci.* **23**, 5446–5454 (2003).

11. Cassanello, C. R. & Ferrera, V. P. Computing vector differences using a gain field-like

mechanism in monkey frontal eye field. *The Journal of Physiology* **582**, 647–664 (2007).

12. Andersen, R. A. & Zipser, D. The role of the posterior parietal cortex in coordinate

transformations for visual–motor integration. *Can. J. Physiol. Pharmacol.* **66**, 488–501

(1988).

13. Stricanne, B., Andersen, R. A. & Mazzoni, P. Eye-centered, head-centered, and

intermediate coding of remembered sound locations in area LIP. *Journal of*

Neurophysiology **76**, 2071–2076 (1996).

14. Giuricich, C., Green, R. J., Jordan, H. & Fallah, M. Target–Distractor Competition

Modulates Saccade Trajectories in Space and Object Space. *eNeuro* **10**, (2023).

15. Sheliga, B. M., Riggio, L. & Rizzolatti, G. Orienting of attention and eye movements.

Exp Brain Res **98**, 507–522 (1994).

16. Sheliga, B. M., Riggio, L. & Rizzolatti, G. Spatial attention and eye movements. *Exp*

Brain Res **105**, 261–275 (1995).

17. Van der Stigchel, S., Meeter, M. & Theeuwes, J. Eye movement trajectories and what

they tell us. *Neuroscience & Biobehavioral Reviews* **30**, 666–679 (2006).

18. Van der Stigchel, S. Recent advances in the study of saccade trajectory deviations.

Vision Research **50**, 1619–1627 (2010).

19. McSorley, E., Haggard, P. & Walker, R. Time Course of Oculomotor Inhibition Revealed

by Saccade Trajectory Modulation. *Journal of Neurophysiology* **96**, 1420–1424 (2006).

20. McSorley, E., Haggard, P. & Walker, R. The spatial and temporal shape of oculomotor

inhibition. *Vision Research* **49**, 608–614 (2009).

21. McSorley, E. & McCloy, R. Saccadic eye movements as an index of perceptual decision-making. *Exp Brain Res* **198**, 513–520 (2009).

22. van Zoest, W., Donk, M. & Van der Stigchel, S. Stimulus-salience and the time-course of saccade trajectory deviations. *Journal of Vision* **12**, 16 (2012).

23. Wang, Z. & Theeuwes, J. Distractor Evoked Deviations of Saccade Trajectory Are Modulated by Fixation Activity in the Superior Colliculus: Computational and Behavioral Evidence. *PLOS ONE* **9**, e116382 (2014).

24. Kehoe, D. H., Rahimi, M. & Fallah, M. Perceptual color space representations in the oculomotor system are modulated by surround suppression and biased selection. *Frontiers in Systems Neuroscience* **12**, 1 (2018).

25. Kehoe, D. H. & Fallah, M. Rapid accumulation of inhibition accounts for saccades curved away from distractors. *Journal of Neurophysiology* **118**, 832–844 (2017).

26. Castellotti, S., Szinte, M., Del Viva, M. M. & Montagnini, A. Saccadic trajectories deviate toward or away from optimally informative visual features. *iScience* **26**, 107282 (2023).

27. Robinson, D. A. Eye movements evoked by collicular stimulation in the alert monkey. *Vision Research* **12**, 1795–1808 (1972).

28. Aizawa, H. & Wurtz, R. H. Reversible Inactivation of Monkey Superior Colliculus. I. Curvature of Saccadic Trajectory. *Journal of Neurophysiology* **79**, 2082–2096 (1998).

29. McPeek, R. M., Han, J. H. & Keller, E. L. Competition Between Saccade Goals in the Superior Colliculus Produces Saccade Curvature. *Journal of Neurophysiology* **89**, 2577–2590 (2003).

30. McPeek, R. M. Incomplete Suppression of Distractor-Related Activity in the Frontal Eye Field Results in Curved Saccades. *Journal of Neurophysiology* **96**, 2699–2711 (2006).

31. Ottes, F. P., Van Gisbergen, J. A. M. & Eggermont, J. J. Visuomotor fields of the superior colliculus: A quantitative model. *Vision Research* **26**, 857–873 (1986).

32. Sommer, M. A. & Wurtz, R. H. Composition and Topographic Organization of Signals Sent From the Frontal Eye Field to the Superior Colliculus. *Journal of Neurophysiology* **83**, 1979–2001 (2000).

33. Ben Hamed, S., Duhamel, J.-R., Bremmer, F. & Graf, W. Representation of the visual field in the lateral intraparietal area of macaque monkeys: a quantitative receptive field analysis. *Exp Brain Res* **140**, 127–144 (2001).

34. Jordan, H. & Tipper, S. P. Object-based inhibition of return in static displays. *Psychonomic Bulletin & Review* **5**, 504–509 (1998).

35. Tipper, S. P., Jordan, H. & Weaver, B. Scene-based and object-centered inhibition of return: Evidence for dual orienting mechanisms. *Perception & Psychophysics* **61**, 50–60 (1999).

36. Sogo, H. & Takeda, Y. Effect of previously fixated locations on saccade trajectory during free visual search. *Vision Research* **46**, 3831–3844 (2006).

37. Pertzov, Y., Zohary, E. & Avidan, G. Rapid Formation of Spatiotopic Representations As Revealed by Inhibition of Return. *J. Neurosci.* **30**, 8882–8887 (2010).

38. Godijn, R., Theeuwes, J. & Godijn, R. The relationship between inhibition of return and saccade trajectory deviations. *J Exp Psychol Hum Percept Perform* **30**, 538–554 (2004).

39. Kehoe, D. H., Lewis, J. & Fallah, M. Oculomotor target selection is mediated by complex objects. *Journal of neurophysiology* **126**, 845–863 (2021).

40. Nobre, A. C., Allison, T. & McCarthy, G. Word recognition in the human inferior

temporal lobe. *Nature* **372**, 260–263 (1994).

41. Ito, M., Tamura, H., Fujita, I. & Tanaka, K. Size and position invariance of neuronal

responses in monkey inferotemporal cortex. *Journal of Neurophysiology* **73**, 218–226

(1995).

42. Desimone, R. & Gross, C. G. Visual areas in the temporal cortex of the macaque. *Brain*

Research **178**, 363–380 (1979).

43. Boussaoud, D., Desimone, R. & Ungerleider, L. G. Visual topography of area TEO in the

macaque. *Journal of Comparative Neurology* **306**, 554–575 (1991).

44. Kehoe, D. H., Aybulut, S. & Fallah, M. Higher order, multifeatural object encoding by the

oculomotor system. *Journal of Neurophysiology* **120**, 3042–3062 (2018).

45. Kehoe, D. H. & Fallah, M. Oculomotor feature discrimination is cortically mediated.

Front Syst Neurosci **17**, 1251933 (2023).

46. van Leeuwen, J., Smeets, J. B. J. & Belopolsky, A. V. Forget binning and get SMART:

Getting more out of the time-course of response data. *Atten Percept Psychophys* **81**,

2956–2967 (2019).

47. Cnaan, A., Laird, N. M. & Slasor, P. Using the general linear mixed model to analyse

unbalanced repeated measures and longitudinal data. *Statistics in Medicine* **16**, 2349–

2380 (1997).

48. Wallace, D. & Green, S. B. Analysis of Repeated Measures Designs with Linear Mixed

Models. in *Modeling Intraindividual Variability With Repeated Measures Data*

(Psychology Press, 2001).

49. Krueger, C. & Tian, L. A Comparison of the General Linear Mixed Model and Repeated

Measures ANOVA Using a Dataset with Multiple Missing Data Points. *Biological*

Research For Nursing **6**, 151–157 (2004).

50. Wolfe, J. M., O'Neill, P. & Bennett, S. C. Why are there eccentricity effects in visual

search? Visual and attentional hypotheses. *Perception & Psychophysics* **60**, 140–156

(1998).

51. Van Heusden, E., Olivers, C. N. L. & Donk, M. The effects of eccentricity on attentional

capture. *Atten Percept Psychophys* **86**, 422–438 (2024).

52. Yoo, S.-A., Tsotsos, J. K. & Fallah, M. The Attentional Suppressive Surround:

Eccentricity, Location-Based and Feature-Based Effects and Interactions. *Frontiers in*

Neuroscience **0**, 710 (2018).

53. Fecteau, J. H. & Munoz, D. P. Salience, relevance, and firing: a priority map for target

selection. *Trends in Cognitive Sciences* **10**, 382–390 (2006).

54. White, B. J., Theeuwes, J. & Munoz, D. P. Interaction between Visual- and Goal-related

Neuronal Signals on the Trajectories of Saccadic Eye Movements. *Journal of Cognitive*

Neuroscience **24**, 707–717 (2012).

55. Jonikaitis, D. & Belopolsky, A. V. Target–Distractor Competition in the Oculomotor

System Is Spatiotopic. *J. Neurosci.* **34**, 6687–6691 (2014).

56. van Leeuwen, J. & Belopolsky, A. V. Distractor displacements during saccades are

reflected in the time-course of saccade curvature. *Sci Rep* **8**, 2469 (2018).

57. Buneo, C. A., Batista, A. P., Jarvis, M. R. & Andersen, R. A. Time-invariant reference

frames for parietal reach activity. *Exp Brain Res* **188**, 77–89 (2008).

58. Bremner, L. R. & Andersen, R. A. Temporal Analysis of Reference Frames in Parietal

Cortex Area 5d during Reach Planning. *Journal of Neuroscience* **34**, 5273–5284 (2014).

59. Cohen, Y. E. & Andersen, R. A. A common reference frame for movement plans in the

posterior parietal cortex. *Nat Rev Neurosci* **3**, 553–562 (2002).

60. Ramezanpour, H. & Fallah, M. The role of temporal cortex in the control of attention.

Current Research in Neurobiology **3**, 100038 (2022).

61. Choi, S. H., Jeong, G., Kim, Y. B. & Cho, Z. H. Proposal for human visual pathway in the

extrastriate cortex by fiber tracking method using diffusion-weighted MRI. *NeuroImage*

220, 117145 (2020).

62. Zaehle, T. *et al.* The neural basis of the egocentric and allocentric spatial frame of

reference. *Brain Research* **1137**, 92–103 (2007).

63. Chen, Z. Object-based attention: A tutorial review. *Atten Percept Psychophys* **74**, 784–

802 (2012).

64. Derbie, A. Y. *et al.* Common and distinct neural trends of allocentric and egocentric

spatial coding: An ALE meta-analysis. *European Journal of Neuroscience* **53**, 3672–3687

(2021).

Acknowledgements: Funding was provided by NSERC Discovery (RGPIN-2016-05296) and

CIHR Operating (102482) grants to MF, and a University of Guelph Undergraduate Summer

Research Assistantship to CEO.

Author contributions: All authors designed the study. CEO collected and analyzed the data. All

authors interpreted the results. CEO wrote the draft manuscript. HJ and MF revised the draft. All

authors reviewed and approved the final manuscript.

Data Availability: The data used in this study are being analysed for a second manuscript but

can be made available from the authors upon reasonable request. Requests should be directed to

the corresponding author.

Additional Information: The authors declare no competing interests.