1

A "torn bag mechanism" of small extracellular vesicle release via limiting membrane rupture of e*n bloc* released amphisomes (amphiectosomes)

Tamás Visnovitz[1, 2, *, #], Dorina Lenzinger[1,*], Anna Koncz[1,3 *], Péter M Vizi[1], Tünde Bárkai[1], Krisztina V Vukman[1], Alicia Galinsoga[1], Krisztina Németh[1,3], Kelsey Fletcher[1], Zsolt I Komlósi[1], Péter Lőrincz[4], Gábor Valcz[1,3,5], Edit I Buzás[1,3,6#]

[1]Semmelweis University, Department of Genetics, Cell- and Immunobiology, Nagyvárad tér 4. 1089 Budapest, Hungary

[2]ELTE Eötvös Loránd University, Department of Plant Physiology and Molecular Plant Biology, Pázmány Péter sétány 1/c, 1117 Budapest, Hungary

[3]HUN-REN-SU Translational Extracellular Vesicle Research Group, Nagyvárad tér 4. 1089 Budapest, Hungary

[4]ELTE Eötvös Loránd University, Department of Anatomy, Cell and Developmental Biology, Pázmány Péter sétány 1/c, 1117 Budapest, Hungary

[5]Department of Image Analysis, 3DHISTECH Ltd, Budapest, Hungary

[6]HCEMM-SU Extracellular Vesicle Research Group, Hungary, Nagyvárad tér 4. 1089 Budapest, Hungary

\#   Correspondence: buzas.edit@med.semmelweis-univ.hu and visnovitz.tamas@med.semmelweis-univ.hu

*   These authors contributed equally

Keywords:

exosomes, amphisome, extracellular vesicles, limiting membrane, secretion, biogenesis

Abstract

Recent studies showed an unexpected complexity of extracellular vesicle (EV) biogenesis pathways.
We previously found evidence that human colorectal cancer cells *in vivo* release large multivesicular
body-like structures *en bloc*. Here, we tested whether this large extracellular vesicle type is unique to
colorectal cancer cells. We found that all cell types we studied (including different cell lines and cells
in their original tissue environment) released multivesicular large EVs. We also demonstrated that
upon spontaneous rupture of the limiting membrane of the multivesicular large EVs, their intraluminal
vesicles (ILVs) escaped to the extracellular environment by a "torn bag mechanism". We proved that
the multivesicular large EVs were released by ectocytosis of amphisomes (hence, we termed them
amphiectosomes). Both ILVs of amphiectosomes and small EVs separated from conditioned media
were either exclusively CD63 or LC3B positive. According to our model, upon fusion of
multivesicular bodies with autophagosomes, fragments of the autophagosomal inner membrane curl up
to form LC3B positive ILVs of amphisomes, while CD63 positive small EVs are of multivesicular
body origin. Our data suggest a novel common release mechanism for small EVs, distinct from the
exocytosis of multivesicular bodies or amphisomes, as well as the small ectosome release pathway.

Introduction

Extracellular vesicles (EVs) are phospholipid bilayer enclosed structures[1-3], which have important roles in cellular homeostasis and intercellular communication. Exosomes have been defined as small (~ 50-200 nm) EVs (sEVs) of endosomal origin[1,3]. Although autophagy is a major cellular homeostatic mechanism, and is implicated in a broad spectrum of human diseases, the intersection of autophagy and exosome secretion remains poorly understood. Recently, regulatory interactions have been shown between autophagy-related molecules and EV biogenesis[4,5]. Furthermore, the LC3-conjugation machinery was demonstrated to specify the cargo packaged into EVs[6]. Importantly, both others and we reported the secretion of LC3-carrying exosomes[6,7]. Particularly relevant to the findings presented here, is the implication of amphisomes (hybrid organelles formed by the fusion of late endosomes/multivesicular bodies (MVBs) with autophagosomes[8,9]) in EV biogenesis. It was suggested that fusion of the limiting membrane of amphisomes with the plasma membrane of cells results in a subsequent release of exosomes by exocytosis[1,3,10]. The current study was prompted by our recent data showing the *in vivo en bloc* release of large, MVB-like sEV clusters by human colorectal cancer cells[11]. Here we investigated if this was a colorectal cancer cell-specific phenomenon. Unexpectedly, we found that it was a general, autophagy-related mechanism of sEV release that we designated as "torn bag mechanism".

Results and Discussion

In this study, we analyzed *in situ* fixed, cultured cells with the released EVs preserved in their original microenvironment on a gelatin/fibronectin coated surface, and we detected large multivesicular EVs (MV-lEVs) in sections of different immersion fixed organs. We tested tumorous HT29, HepG2, and non-tumorous HEK293, HEK293T-PalmGFP, HL1 cell lines, as well as primary suspension type bone marrow derived mast cells (BMMCs). In addition, we studied ultrathin sections of mouse kidney and liver.

By the analysis of transmission electron micrographs of all tested cell types, we identified budding (Fig.1A-G) and secretion (Fig.1H-N) of MV-lEVs carrying intraluminal vesicles (ILVs). Importantly, in all cases we found evidence for the extracellular rupture of the limiting membrane of MV-lEVs and the release of ILVs (Fig.1O-U). For this novel type of sEV release, we suggest the designation "torn bag mechanism" which is distinct from the exocytosis of MVBs and amphisomes[1,3,10] and from the release of plasma membrane derived small EV (sEVs) by ectocytosis[12].

Most relevant to the *in vivo* conditions, we also observed the same phenomenon within the blood vessels of ultrathin sections of both murine kidney (Fig.1V) and liver (Fig.1W,X). In these cases, both

the intact MV-lEVs (Fig.1V-X) and the "torn bag release" of sEVs (Fig.1V) were detected. Fig.1X

shows that a circulating leukocyte releases MV-lEVs by ectocytosis.

Next, we decided to further investigate the subcellular origin of the ILVs within the secreted MV-

lEVs. We analyzed the microenvironment of *in situ* fixed HEK293T-PalmGFP cells by confocal

microscopy. The PalmGFP signal of HEK293T-PalmGFP cells principally associates with the plasma

membrane[13] (Fig.2_S1C-E). The green fluorescence helped us to identify the plasma membrane-

derived limiting membrane of MV-lEVs. In agreement with our previous findings on HT29 colorectal

cancer cells, within the MV-lEVs, we found CD63/ALIX (Fig.2A,G) CD81/ALIX (Fig.2B,H),

CD63/TSG101 (Fig.2.C,I) and CD81/TSG101 (Fig.2D,J) double positive ILVs or ILV clusters. In line

with a recent finding[12], CD81 was present in the plasma membrane-derived limiting membrane

(Fig.2B,D,F), while CD63 was only found inside the MV-lEVs (Fig.2A,C,E).

Next we studied the possible autophagy related aspects of the secreted MV-lEVs. ILVs were tested for

the autophagy marker LC3B in parallel with CD63 and CD81. Although LC3B, CD63 and CD81 were

all present in association with the ILVs (Fig.2E,F), the LC3B and CD63 (Fig.2K) and the LC3B and

CD81 (Fig.2L) signals did not overlap. Fig.2M shows that while the known sEV markers (CD63,

CD81, TSG101 and ALIX) strongly co-localized with each other, LC3B positivity hardly showed co-

localization with CD63 or CD81. Immunocytochemistry results of HT29 and HepG2 cells further

validated the findings obtained with the HEK293T-PalmGFP cells (Fig.2_S2). The ILVs of

HEK293T-PalmGFP and HepG2 cell lines were Rab7 positive (Fig.2_S1A,B) suggesting a late

endosomal origin. Western blot validation of the applied antibodies is summarized in Fig.2_S3. The

sEV markers were also tested on separated sEVs of HEK293T-PalmGFP cells by transmission

electron microscopy (TEM). Fig.2P confirms the typical sEV morphology. With TEM double

immunogold labelling, using anti-LC3B and anti-CD63 antibodies simultaneously, we found distinct

LC3B positive (Fig.2Q, Fig.2_S5O) and CD63 positive (Fig.2R, Fig.2_S5O) sEVs. Based on the

analysis of TEM images, the diameters of unlabeled, and LC3B positive and negative sEVs were

determined (Fig.2S). The LC3B positive sEVs had a significantly larger diameter as compared to the

LC3B negative ones. The simultaneous presence of CD63, CD81, TSG101, ALIX and the

autophagosome marker LC3B within the MV-lEVs (rather than on their limiting membrane) identified

the extracellular MV-lEVs as *en bloc* released amphisomes.

The "torn bag mechanism" was also monitored by live cell SIM[2] super-resolution microscopy analysis

of HEK293T-PalmGFP-LC3RFP cells (Fig.2N,O). The release of the LC3 positive red fluorescent

signal was detected within a relatively short period of time (260 sec) (Fig.2O). We could rule out the

possibility that rupture of the limiting membrane detected by TEM (Fig.1O-T,V) was a fixation

artefact by showing the release of LC3 positive sEVs from amphiectosomes with live-cell imaging.

Characterization of the in-house developed HEK293T-PalmGFP-LC3RFP cell line is shown in

Fig.2_S4.

In the following step, we addressed the question whether LC3, associated with the ILVs of MV-lEVs,

indeed reflected autophagy origin. We tested MVBs (Fig.2_S5A,F,K), autophagosomes

(Fig.2_S5B,G,L), amphisomes (Fig.2_S5C,H,M), amphiectosomes (Fig.2_S5D,I,N) and isolated sEV

fractions of the same cells (Fig.2_S5E,J,O). Using immune electron microscopy, as expected, we

found CD63 single positivity in MVBs (Fig.2_S5K). In autophagosomes, the limiting phagophore

membrane was LC3B positive, and CD63 positivity was also present (Fig.2_S5L). The limiting

membrane of amphisomes was LC3B negative, and the internal membranous structures were either

LC3B or CD63 positive (Fig.2_S5M). The same immunostaining was also observed in the ILVs of the

released amphiectosomes (Fig.2_S5N). Importantly, the separated sEVs of HEK293T-PalmGFP cells

were either LC3B or CD63 positive (Fig.2_S5O). Thus, we confirmed our confocal microscopy results

at the ultrastructural level. Using immunogold TEM, we provided further evidence for the

budding/ectocytosis mechanism of amphiectosome release (Fig.2_S5P). The diameters of ILVs within

MVBs, amphisomes and amphiectosomes were compared (Fig.2_S5Q), and the differences were

likely due to the different pH and osmotic conditions within these organelles. In agreement with our

observations with separated sEVs, LC3B positive ILVs had a significantly larger diameter than the

LC3B negative ones (Fig.2_S5R) possibly indicating their different intracellular origin. Based on all

the above findings, we propose the following model (Fig.3A): autophagosomes and MVBs fuse to

form amphisomes, and the inner, LC3 positive membrane of autophagosomes disintegrates[14]. It curls

up and forms LC3 positive ILVs. Therefore, amphisomes contain both MVB-derived and

autophagosome-derived, LC3 negative and positive ILVs, respectively. The amphisome is next

released from the cell by ectocytosis. Finally, the plasma membrane-derived limiting membrane

ruptures enabling the ILVs escape to the extracellular space by a "torn bag mechanism".

To investigate the process of amphiectosome release, we exposed the MV-lEV releasing cells to

different *in vitro* treatments (Fig.3B). The release of MV-lEVs was monitored by confocal microscopy

of *in situ* fixed cell cultures. Optimal test conditions were determined (Fig.3_S1A-F) and the results

are summarized in Fig.3C. Cytochalasin B did not have any effect on the discharge of MV-lEVs

suggesting that the release did not involve a major actin-dependent mechanism. In contrast, there was

a significant reduction of the MV-lEV secretion upon exposure of the cells to Colchicine indicating a

role of microtubules in the release of the MV-lEVs. While Rapamycin significantly reduced the

discharge of MV-lEVs, Chloroquine and Bafilomycin induced an enhanced MV-lEV secretion.

Rapamycin activates autophagic degradation[15], therefore, it induces a shift towards degradation as

opposed to secretion. The lysosomotropic agents Chloroquine and Bafilomycin are known to interfere

with the acidification of lysosomes[16,17]. By blocking the degradation pathway of MVBs/amphisomes

(Fig.3B), an enhanced sEV secretion is observed. This effect is well known for exosome secretion

from MVBs[18,19]. The diameters of the released MV-lEVs were determined based on confocal images

(Fig.3_S1G). Metabolic activity of the cells was determined by a Resazurin assay, and a significant

reduction was detected upon exposure of the cells to Rapamycin (Fig.3_S1D) in line with previously

published data[20]. LC3II is the membrane-associated, lipidated autophagic form of LC3[21] and it is the

hallmark of autophagy related membranes[14]. Importantly, by Western blot, we not only showed the

presence of the membrane-bound LC3II in serum-free, lEV-depleted (sEV containing) conditioned

medium of both HEK293T-PalmGFP and HEK293T-PalmGFP-LC3RFP cells, but the amount of

LC3II substantially increased upon Chloroquine treatment (Fig.3D). Raw data of Western blots are

available in Fig.3_S2.

Recent advances in the EV field shed light on migrasomes, a special type of MV-lEVs[22,23]. With their

pomegranate-like ultrastructure, migrasomes resemble amphiectosomes. Therefore, we tested the

presence of TSPAN4, a migrasome limiting membrane marker[23], in amphiectosomes. Fig.4A,B,G,H

show that although TSPAN4 was present intraluminally in the HEK293T-PalmGFP-derived MV-

lEVs, it was clearly absent from their limiting membrane. Surprisingly, we identified two different

HT29 cell-derived MV-lEV populations: one with intraluminal TSPAN4 only (Fig.4C,E,I,K), and

another one with a TSPAN4 positive limiting membrane (Fig.4D,F,J,L). This raised the possibility that

the latter population corresponded to migrasomes. Our co-localization analysis also confirmed the

existence of two distinct MV-lEV populations (Fig.4M). Next, we carried out live-cell imaging on

HEK293T-PalmGFP-LC3RFP cells. The released MV-lEVs were either LC3 positive or negative

intraluminally (Fig.4N,O). Our TEM images confirmed that certain cell types can release both

migrasome-like structures and amphiectosomes. MV-lEVs with typical migrasome-associated

retraction fiber(s) were detected in the case of HL1 (Fig.4P), HEK293T-PalmGFP (Fig.4Q) and

BMMC cells (Fig.4R). Importantly, the same cells also released amphiectosomes by budding from the

cell surface (Fig.4S-U). Taken together, based on the absence of TSPAN4 in their limiting membrane

and their lack of association with retraction fibers, amphiectosomes appear to be distinct from

migrasomes. Besides migrasomes, another MV-lEV type was described in the case of gastrointestinal

tumors and low-grade glioblastoma cells referred to as spheresome[24,25]. However, there is no data on a

relationship of spheresome release and autophagy. Recently, endothelial cell derived,

multicompartmented microvesicles (MCMVs) were shown to protrude and pinch-off from the cell

surface releasing ILVs by a mechanism similar to exocytosis[26]. The absence of protrusion clusters

described for MCMV[26] distinguishes amphiectosomes from MCMVs.

Our approach, involving *in situ* fixation of cultures and tissues, made it possible to recognize sEV

release from amphiectosomes by the "torn bag mechanism". We propose that this mechanism can be

easily missed if conditioned medium is subjected to centrifugation, SEC purification or even to simple

pipetting, which may rupture of the limiting membrane of amphiectosomes. This aligns with our

observation that the spontaneous escape of ILVs from untouched amphiectosomes can occur as early

as 5 minutes after amphiectosome release. Our data presented here suggest that amphiectosome
secretion and the "torn bag mechanism" may have a significant, however, previously unrecognized
role in sEV biogenesis.

Figures

Figure 1. Transmission electron microscopic detection of the release and
extracellular fate of large, multivesicular EVs secreted by different cell lines
and cells in mouse organs

Major steps of the release of large, multivesicular EVs were detected in the case of all tested cell lines including the immortal, non-tumorous HEK293T-PalmGFP (A,H,O), HEK293 (B,I,P), the tumorous cell lines HepG2 (C,J,Q) and HT29 (D,K,R), the beating cardiomyocyte cell line HL1 (E,L,S) and the primary suspension of bone marrow derived mast cells (BMMCs) (F,M,T). The different phases of EV secretion were also captured in the circulation of mouse kidney (V) and liver (W,X). According to the electron micrographs, we found evidence for the budding (A-G,X) and secretion (H-N,V,W) of the multivesicular large EVs (MV-lEVs). We also detected the extracellular rupture of the limiting membrane of the released MV-lEVs with the escape of the intraluminal vesicles (ILVs) by a "torn bag mechanism" (O-U,V). Although it is not always clear whether the secreted MV-

201   lEVs have a single or double limiting membrane, several micrographs suggest the presence of the double

202   membrane (Y-AF) in the secreted MV-lEVs. In the case of BMMCs (Y), the release phase of a multivesicular

203   structure is captured. The bottom portion of this structure embedded in the cytoplasm is surrounded by a single

204   membrane (white arrow head) while the upper (budding) portion is covered by double membrane (asterisk).

Figure 2. Detection of conventional sEV markers and the LC3 protein in HEK293T-PalmGFP cell-derived EVs

Widely used sEV markers (CD63, CD81, ALIX and TSG101) and LC3B were tested in MV-lEVs found in the microenvironment of the releasing cells by confocal microscopy after *in situ* fixation (A-F). Normalized fluorescence intensities were calculated in order to determine the relative localization of the limiting membrane (PalmGFP), the conventional sEV markers and the LC3B signal (G-L). Fluorescence intensity peaks of sEV markers were largely overlapping with each other, while the LC3B signal and the sEV markers showed separation. Co-localization rates were also calculated (M). The sEV markers co-localized with one another as no significant difference was found among them. In contrast, low co-localization rates were detected between the "classical" sEV markers and LC3B (one-way ANOVA, p<0.0001, n=8-26 confocal images). Real time release of LC3 positive sEVs by the "torn bag mechanism" was studied in the case of HEK293T-PalmGFP-LC3RFP cells by Elyra7 SIM² super-resolution live cell imaging (N,O). Images were recorded continuously, and selected serial time points are shown. LC3 positive, red fluorescent small particles were released within a 5 min timeframe (O) and are indicated by white arrows. Presence of CD63 and LC3B were detected in the case of a separated sEV fraction using immunogold TEM. HEK293T-PalmGFP derived sEV fraction is shown by negative-positive contrast without immune labelling (P). In double-labelled immunogold TEM images (Q,R), distinct LC3B

positive (Q) and CD63 positive (R) sEVs were found. However, CD63-LC3B double positive EVs were not

detected. Black arrowheads indicate 10 nm gold particles identifying LC3B, while white arrowheads show 5 nm

gold particles corresponding to the presence of CD63. Quantitative analysis of TEM images was performed (S),

and the diameters of different EV populations were determined. The LC3B negative population was significantly

smaller than the LC3B positive one ($p < 0.0001$, t-test; n=79-100). No difference was detected when the

immunogold labelled sEV fraction (either LC3B positive or negative, LC3B+/-) and the unlabeled sEV fraction

(sEV) were compared ($p < 0.05$, t-test, n=112-179).

Fig.2_S1. Confocal microscopic analysis of HEK293T-PalmGFP and
HepG2 cells derived lEVs; HEK293T-PalmGFP cells

The MV-lEVs released by HEK293T-PalmGFP (A) and HepG2 cells (B) were tested for CD63 sEV and Rab7
late endosomal markers. Both CD63 and Rab7 were present in the ILVs. The GFP positivity of the HEK293T-
PalmGFP cells was controlled by immunofluorescence microscopy on control (C) and Chloroquine-treated (D)
cells. Co-localization of Palm-GFP and the anti-GFP signal was calculated in the plasma membrane and in the
cytoplasm (E) and Pearson's correlations were visualized. The green fluorescence in the plasma membrane was
clearly GFP-dependent in the case of control (Ctrl Membrane) and Chloroquine-treated (Chloro Membrane)
samples, while in the cytoplasm, the correlation was significantly weaker (Ctrl Cytoplasm and Chloro
Cytoplasm). Within the cytoplasm, the correlation was stronger in Chloroquine-treated cells, suggesting that the
phagophore membrane may accumulate Palm-GFP ($p < 0.0001$, t-test; $n = 45$).

Fig.2_S2 Confocal microscopic images of amphiectosome release by HT29 and HepG2 cells

The intraluminal EVs of amphiectosomes were found to be positive for LC3B and CD81 in HT29 cultures (A) and HepG2 cells (H). In the case of HT29 cells, phases of the "torn bag mechanism" were captured, including a

249   secreted intact amphiectosome (B), MV-lEV with ruptured limiting membrane releasing internal vesicles (C,

250   white arrow), an inside-out secreted amphiectosome (D, white arrow), and an amphiectosome with a fully

251   disintegrated limiting membrane and released sEVs (E). The presence of TSG101/CD63 (F), ALIX/CD81 (G),

252   and LC3B/CD81 (H) was also examined in the released amphiectosomes of the HepG2 cell line.

 Fig.2_S3. Western blot validation of antibodies used in immune-
fluorescence detection

Lysates from three distinct human cell lines (HEK293, HepG2 and HT29) were employed in the validation process to reduce the cell line-specific variations in the study. Protein bands lacking posttranslational modifications are noted as "w/o PTM." Numerous posttranslational modifications of CD63 and CD81 are widely recognized.

Fig.2_S4. Characterization of the HEK293T-PalmGFP-LC3RFP cell line.

Confocal microscopy (A) and Western blot analysis (B) were employed to characterize the HEK293T-PalmGFP-
LC3RFP cell line. For overnight Chloroquine treatments, 30 μM Chloroquine was applied. Punctate LC3
fluorescence observed in (A) corresponds to autophagosomes. In panel (B), whole-cell lysates from HEK293T-
PalmGFP (lane i), HEK293T-PalmGFP-LC3RFP (lane ii) and Chloroquine-treated HEK293T-PalmGFP-
LC3RFP (lane iii) samples were examined. Gray arrows indicate LC3I, while black arrows highlight LC3II. Full
Western Blot images can be find in Fig.3_S2 supplementary figure.

Fig.2_S5. Structures involved in amphiectosome release

Multivesicular body (MVB, A), autophagosome (B), amphisome (C), amphiectosome (D) and secreted sEVs (E) were identified by TEM with and without immunogold labelling of HEK293T-PalmGFP cell cultures. White arrowheads (5 nm gold particles) indicate CD63 and black arrowheads (10 nm gold particles) show LC3B. While MVBs (F) were LC3B negative (K), we detected CD63 positivity on the surface of the ILVs (K). In an autophagosome (G), the limiting membrane layers were double positive for CD63 and LC3B (L). In contrast, the internal membranes of autophagosome were CD63 single positive (L). In the case of an amphisome (H), heterogeneous membrane structures were visible with variable size and morphology. The ILVs were either CD63 or LC3B positive (M). The amphiectosomes were located in the extracellular space and contained ILVs of different size and shape (I). The ILVs of amphiectosome (as in case of amphisome), were either CD63 or LC3B positive (N). Secreted sEVs (J) with immunogold labelling (O), were also found to be either CD63 or LC3B positive. Release of an amphiectosome is shown (P) with CD63 and LC3B immunogold signals. Higher magnification of the insert is indicated by the black rectangle. It shows either CD63 or LC3B positive ILVs. Size distributions of ILVs of MVBs, amphisomes and amphiectosomes were determined on Epon-embedded ultrathin sections (Q). Although the ILV sizes differed significantly (one-way ANOVA, ****: p<0.0001, n=73, 138 and 595, respectively), the majority of ILVs had a diameter between 40-100 nm. The diameter of LC3B positive and negative ILVs of amphiectosomes was assessed on TEM images of immunogold labelled ultrathin sections (R).

289   LC3B negative ILVs were significantly smaller than the LC3B positive ones, while the ILVs in the Epon-
290   embedded sections did not differ from the LC3B positive ones (one-way ANOVA, p <0.001, n= 595, 101 and
291   70, respectively).

Figure 3. Amphiectosome release and its modulation

Based on our data, a model of amphiectosome release was generated (A). According to this model, the fusion of MVBs and autophagosomes forms amphisomes. The LC3B positive membrane layer (indicated in orange) undergoes disintegration and forms LC3B positive ILVs inside the amphisome. Later, the amphisome is released into the extracellular space by ectocytosis and can be identified extracellularly as an amphiectosome. Finally, the limiting membrane(s) of the amphiectosome is ruptured and the ILVs are released as sEVs into the extracellular space by a "torn bag mechanism". In order to further support our model on amphiectosome release and "torn bag" EV secretion different *in vitro* treatments were applied. Cytochalasin B, Colchicine, Chloroquine, Bafilomycin A1 and Rapamycin were used to modulate amphisome release. Points of actions are summarized (B). While Cytochalasin B inhibits actin-dependent membrane budding and cell migration, Colchicine blocks the microtubule-dependent intracellular trafficking. Chloroquine and Bafilomycin have similar, Rapamycin have opposite effects on lysosome-autophagosome or lysosome-amphisome fusion. Chloroquine and Bafilomycin inhibits lysosomal degradation while Rapamycin accelerates it. Based on confocal microscopy, Cytochalasin B (CytoB) did not alter the dynamics of amphiectosome release (C). In contrast, both Colchicine (Colch) and Rapamycin (Rapa) significantly inhibited the release of amphiectosomes, while Chloroquine (Chloro) and Bafilomycin (Bafilo) increased the release frequency. Between Chloroquine and Bafilomycin effect significant difference was not detected (C). Results are shown as mean ± SD of 3-4 independent biological replicates,

analyzed by one-way ANOVA, *: $p < 0.05$, **: $p < 0.01$, ns: non-significant. Presence of membrane-bound

(lipidated) LC3II was tested by Western blotting. The total protein content of serum-, cell- and large EV-

depleted conditioned medium of HEK293T-PalmGFP (PalmGFP) and HEK293T-PalmGFP-LC3RFP

(PalmGFP-LC3RFP) cells was purified by precipitation and 20 µg of the protein samples were loaded on the gel

(D). The lipidated LC3II band was detected in all cases. Relative expression of control (Ctrl) and Chloro treated

samples were determined by densitometry. Chloroquine treatment increased the LC3II level by approximately by

two fold. Results are shown as mean ± SD of n=6 biological replicates.

Fig.3_S1. Dose determination for treatments and size distribution of MV-lEVs

Relative metabolic activity was assessed through a Resazurin assay (A-E) during treatment optimization. The red dashed line indicates 100 % metabolic activity, representing control cells. Results are presented as mean ± SD values of n=3-4 independent biological replicates. Student's unpaired t-test was performed to compare control and treated cells (*: p<0.05, **: p<0.01, ****: p<0.0001). For Colchicine treatment, alterations in the microtubular network were observed through immunocytochemistry (F) and documented using an epifluorescent microscope. Changes in the diameter of released amphiectosomes under various treatments were determined on confocal microscopy images. A significant reduction in size was identified only in the case of Rapamycin treatment (G, one-way ANOVA test *: p<0.05, n= 95-101).

331 **Fig.3_S2. Unedited Western blots of Fig.2_S3 and Fig.3**

1: HEK 293T-PalmGFP
2: HEK 293T-PalmGFP-LC3RFP
3: Chloroquine treated HEK 293T-PalmGFP-LC3RFP

Subfigure A displays the unedited Western blots utilized for the quantification in Fig.3D. Black arrows indicate
the location of LC3I, while red squares represent the quantified LC3II. In Subfigure B, the unedited blots from
Fig.2_S3B are presented. Black arrows denote the LC3I-RFP fusion protein, while red arrows indicate the
LC3II-RFP fusion protein. The blue arrow points to the native LC3I, while the green arrow represents the native
LC3II protein.

Figure 4. Comparison of amphiectosomes and migrasomes

Commonly used sEV markers (CD63, CD81) and TSPAN4, a suggested migrasome marker, were tested in *in situ* fixed intact MV-lEVs of HEK293T-PalmGFP (A,B) and HT29 (C-F) cells by confocal microscopy. Normalized fluorescence intensities were calculated in order to determine the relative localization of the limiting membrane (with PalmGFP or lactadherin labelling) and the CD63/TSPAN4 and CD81/TSPAN4 markers (G-L). In the case of HEK293T-PalmGFP-derived EVs, there were no "classical migrasomes" with TSPAN4 in their limiting membrane were found. The TSPAN4 signal was only detected intraluminally in the MV-lEVs. The limiting membranes of HT29-derived MV-lEVs were either TSPAN4 positive and negative. The co-localization

rate between the limiting membrane and TSPAN4 was low in case of HEK293T-PalmGFP-derived EVs. In the

case of HT29 cells, two MV-lEV populations were identified: one with low and one with high co-localization

rates (M). Live cell imaging of HEK293T-PalmGFP-LC3RFP cells showed migrasomes-like (retraction fiber-

associated) MV-lEVs with or without intraluminal LC3 positivity (N,O). Using TEM, we could identify

structures with retraction fiber-associated migrasome morphology in the case of HL1 (P), HEK293T-PalmGFP

(Q) and BMMC (R) cells. For comparison, budding of amphiectosomes of the same HL1 (S), HEK293T-

PalmGFP (T) and BMMC cells (U) are shown (without being associated with long retractions fibers).

Material and methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Cell line (*Homo sapiens*)	HEK293 human kidney(embryonic)	ECACC (Sigma)	#85120602	Batch No: 18E026
Cell line (*Homo sapiens*)	HT29 Caucasian colon adenocarcinoma grade II	ECACC (Sigma)	#91072201	Batch No 09K003
Cell line (*Homo sapiens*)	HepG2 human hepatocyte carcinoma	ECACC (Sigma)	#85011430	Batch No: 19B009
Cell line (Mus musculus)	HL1 mouse cardiomyocyte cell line, atrial	Merck	# SCC065	Batch No: RD1601001
Cell line (*Homo sapiens*)	HEK293T-PalmGFP human kidney(embryonic) expressing palmitoylated GFP	Kind gift of Charles Lai https://doi.org:10.1038/ncomms8029		Resorted before MCB preparation
Cell line (*Homo sapiens*)	HEK293T-PalmGFP-LC3RFP human kidney(embryonic) expressing palmitoylatedGFP and RFP tagged LC3	This paper		See material and methods
Cell line (Mus musculus)	BMMC bone marrow derived mast cells	Primary cell culture https://doi.org/10.1002/jev2.12023		
Lentiviral particles	LentiBrite RFP-LC3 Lentiviral Biosensor	Merck	17-10143	Batch No: 3530171

Biological sample (*Mus musculus*)	own animal house		C57BL/6	male, 12 weeks old
Antibody	rabbit polyclonal anti-CD63 (C-terminal)	Sigma/Merck	SAB2109138	IF (1:200) WB (1:500)
Antibody	mouse monoclonal anti-CD63	SantaCruz Biotechnology	MX-49.129.5 clone: sc-5275	IF (1:200) TEM (1:50) WB (1:1,000)
Antibody	rabbit polyclonal anti-CD81	Sigma/Merck	SAB3500454	IF (1:200) WB (1:2,500)
Antibody	mouse monoclonal anti-CD81	Invitrogen	MA5-13548 clone: 1.3.3.22	IF (1:100) WB (1:100)
Antibody	rabbit polyclonal anti-TSG101	Sigma/Merck	HPA006161	IF (1:200) WB (1:1,000)
Antibody	rabbit polyclonal anti-ALIX (C-terminal)	Sigma/Merck	SAB420047	IF (1:200) WB (1:1,000)
Antibody	rabbit monoclonal anti-LC3B	Sigma/Merck	ZRB100 clone: 12K5	IF (1:200) TEM (1:50) WB (1:1,000)
Antibody	rabbit monoclonal anti-LC3A	Sigma/Merck	ZRB1125 clone: 3J12	IF (1:200) WB (1:1,000)
Antibody	rabbit polyclonal anti-TSPAN4	Bioss	BS-9413R	IF (1:200)
Antibody	mouse monoclonal anti-Rab7	Sigma/Merck	R8779 clone: Rab7-117	IF (1:200) WB (1:1,000)
Antibody	mouse monoclonal anti-α-tubulin	Sigma/Merck	T9026 clone: DM1A	IF (1:200)
Antibody	mouse monoclonal anti-GFP	Sigma/Merck	G6539 clone: GFP-20	IF (1:200) WB (1:1,000)
Antibody	mouse monoclonal anti-RFP	Invitrogen	MA5-15257 clone: RF5R	WB (1:1,000)
Antibody	goat anti-mouse IgG-ATTO550	Sigma/Merck	43394	IF (1:1,000)

Antibody	goat anti-rabbit IgG-ATTO647N	Sigma/Merck	40839	IF (1:1,000)
Antibody	goat polyclonal anti-rabbit IgG Fc (HRP)	abcam	ab97200	WB (1:10,000)
Antibody	goat polyclonal anti-mouse IgG Fc (HRP)	abcam	ab97265	WB (1:10,000)
Antibody	goat polyclonal anti-rabbit IgG (whole molecule) 10 nm gold pre-adsorbed	abcam	ab27234	TEM (1:50)
Antibody	goat polyclonal anti-mouse IgG (whole molecule) 5 nm gold pre-adsorbed	Sigma/Merck	G7527	TEM (1:50)
Chemical compound, drug	Bafilomycin A1	Sigma/Merck	B1793	Lot Number: 0000190389
Chemical compound, drug	Colchicine	Serva	77120.02	Lot Number: 190300
Chemical compound, drug	Chloroquine-diphosphate	Invitrogen	P36236 C	Lot Number: 2441325
Chemical compound, drug	Rapamycin	Sigma/Merck	R0395	Lot Number: 0000084976
Chemical compound, drug	Cytochalasin B	Sigma	C2743	Lot Number: 037M4083V
Chemical compound, drug	FBS	Biosera	FB-1090/500	Lot Number: 015BS575
Software, algorithm	LASX	Leica	Leica Application Suite X 3.5.5.19976	
Software, algorithm	ZEN Blue	Zeiss	ZEN 2.3 lite	
Software, algorithm	iTEM	Olympus	iTEM 5.1	

Software, algorithm	Image J	https://imagej.net/ij/	v1.54g	
Software, algorithm	Prism9	GraphPad	GraphPad Prism 9.4.1	
Software, algorithm	Biorender	https://www.biorender.com/		

Cell lines

The HEK293 human embryonic kidney, the HepG2 human hepatocyte carcinoma cell line and the HT29 human colon adenocarcinoma cell lines were purchased from the European Collection of Authenticated Cell Cultures (ECACC) through their distributor (Sigma). The HL1 cell line was purchased from Millipore. The HEK293T-PalmGFP human embryonic kidney cells was kindly provided by Charles P. Lai[27]. Mouse bone marrow-derived mast cells (BMMCs) were differentiated and expanded as we described previously[28]. The HEK293, HEK293T-PalmGFP and HepG2 cell lines were grown in DMEM (Gibco)[29,30], the HT29 cells were cultured in RPMI 1640 (Gibco)[11], while the HL1 cells were grown in Claycomb medium[31]. All cells were cultured with 10 % fetal bovine serum (FBS, BioSera) in the presence of 100 U/mL of Penicillin and 100 μg/mL Streptomycin (Sigma).

Before analysis by confocal microscopy, the cells were cultured on the surface of gelatin-fibronectin coated glass coverslips (VWR). The coating solution contained 0.02 % gelatin (Sigma) and 5 mg/mL fibronectin (Invitrogen). Coverslips were coated overnight (O/N) at 37 °C.

For transmission electron microscopy (TEM), the adherent cells (HEK293, HEK293T-PalmGFP, HepG2, HT29 and HL1) were grown in gelatin-fibronectin coated 8-well Flux Cell Culture Slides (SPL).

Cell cultures were tested regularly for Mycoplasma infection by PCR, with the following PCR primers: GAAGAWATGCCWTATTTAGAAGATGG and CCRTTTTGACTYTTWCCAC-CMAGTGGTTGTTG [31].

Generation of HEK293T-PalmGFP-LC3RFP cell line

For generation of a stable HEK293T-PalmGFP-LC3RFP cell line, HEK293T-PalmGFP cells were transfected by LentiBrite RFP-LC3 Lentiviral particles (Merck) according to the instructions of the manufacturer. The GFP-RFP double positive cells were sorted by a HS800 Cell Sorter (SONY) and cell banks (master and working cell banks) were prepared. The success of the stable transfection was analyzed by immunocytochemistry and Western blotting. Results are shown in Fig.2_S3.

Confocal microscopy

Confocal microscopy was carried out as we described earlier[28,31] with some modifications. Briefly, unlike the majority of the studies in the field of EVs, here we analyzed untouched, *in situ* fixed and cultured cells and their

microenvironment. Since centrifugation may disrupt the limiting membrane of amphiectosomes, the *in situ*
fixation made it possible to observe them in their intact form. The culture medium was gently removed by
pipetting from above the cells leaving a thin medium layer only (approximately 150 μL of liquid on the cells).
Without any further washing, cells were *in situ* fixed by 4 % paraformaldehyde (PFA) in phosphate buffer saline
(PBS) for 20 min at room temperature (RT). The released lEVs were either fixed and captured during the release
or were preserved on the gelatin/fibronectin surface coating. After fixation, 3x 5 min washes with 50 mM
glycine in PBS were carried out. In the case of the non-fluorescent HepG2 and HT29 cells, a lactadherin-based
plasma membrane staining was performed[13,28,31]. Lactadherin (Haematologic Technologies) was conjugated to
ATTO488 fluorophore (abcam) according to the instructions of the manufacturer. The lactadherin-ATTO488
conjugate was added to the cells in 1:100 dilution in PBS (for 1 h, RT). The unbound lactadherin was removed
by washing with PBS (3 times, 5 min, RT) and post-fixation was carried out by 4 % PFA (20 min, RT). PFA was
removed by washes with 50 mM glycine in PBS (3 times, 5 min, RT). Blocking and permeabilization of the cells
were performed by 10 % FBS with 0.1 % TritonX-100 (Sigma) in PBS (1 h, RT). In general, primary antibodies
were applied in 1:200 dilution O/N at 4 °C in the above blocking and permeabilization solution. Excess primary
antibodies were eliminated by washing with the blocking and permeabilization solution (3 times, 5 min RT). The
secondary antibodies were applied in 1:1000 dilution in 1 % FBS in PBS (1 h, RT). Unbound secondary
antibodies were eliminated by washing (1 % FBS, in PBS, 2 times, 5 min; PBS 2 times, 5 min; water 2 times, 5
min) and the samples were mounted in ProLong Diamond with DAPI (Invitrogen). Slides were examined by
Leica SP8 Lightning confocal microscope with adaptive lightning mode using a HC PL APO CS2 63x/1.40 OIL
objective with hybrid detector. The applied lookup tables (LUT) were linear during this study. For image
analysis and co-localization studies, we applied Leica LASX software using the unprocessed raw images. In case
of co-localization studies, 20 % threshold and 10 % background settings were applied.

Transmission electron microscopy:

Adherent cells (HEK293, HEK293T-PalmGFP, HepG2, HT29 and HL1), as well as mouse (C57BL/6) kidney
and liver tissues pieces (approx. 1.5 mm x 1.5 mm) were immersed in and fixed by 4 % glutaraldehyde (48 h/4
°C), post-fixed by 1% osmium tetroxide (2 h, RT) and were embedded into Epon resin (Electron Microscopy
Sciences) as described previously[32]. In the case of BMMCs, 920 μL cell suspension was complemented with 80
μL 50% glutaraldehyde to reach the final 4 % glutaraldehyde concentration. Cells were fixed for 48h at 4°C and
were post-fixed by 1% osmium tetroxide (2 h, RT). During sample preparation, BMMCs were collected by
gravity-based sedimentation. Due to the high viscosity of Epon resin, BMMCs were embedded in LR White low
viscosity resin (SPI Supplies) according to the instructions of the manufacturer. Ultrathin sections (60 nm) were
contrasted by uranyl acetate (3.75 %, 10 min, RT) and lead citrate (12 min, RT).

For immunogold TEM studies, cells and tissues were fixed by 4 % PFA with 0.1% glutaraldehyde (48 h, 4 °C)
and were post-fixed by 0.5 % osmium tetroxide (30 min, RT). Samples were embedded into LR White and were
immunogold labelled as described previously[11]. The contrast was enhanced by uranyl acetate (3.75 %, 1 min,
RT) and lead citrate (2 min, RT). Purified HEK293T-PalmGFP-derived sEV-s were detected by negative-
positive contrasting and immunogold labelling as described previously[31]. Antibodies were used in 1:50 dilution.
Detailed list of the used antibodies is available in the Key Resources Table.

For all electron microscopic studies, a JEOL 1011 transmission electron microscope was used. Images were

captured with the help of Olympus iTEM software and for image analysis, Image J software was used.

Live cell imaging

The HEK293T-PalmGFP-LC3RFP stable cell line was cultured the same way as HEK293T-PalmGFP cells.

Before the experiments, gelatin-fibronectin coated 10-well coverslip bottom chamber slide (Greiner-BioOne)

was seeded and treated by 30 µM Chloroquine O/N. Release of migrasomes, amphiectosomes and sEVs were

followed by the Leica SP8 Lightning confocal microscope equipped with an Okolab environmental chamber and

a Zeiss ELYRA 7 with Lattice SIM² super-resolution fluorescent microscope with the help of 63x/1.4 plan-

apochromat Oil objective. For image analysis, we applied Leica LASX, Zeiss ZEN Blue and Image J software.

Modulation of amphiectosome release

In order to test the release mechanism of amphiectosomes and to distinguish them from migrasomes, different

treatments were applied overnight (O/N) in fresh, serum containing cell culture medium except for Colchicine,

where 1 h treatment was selected. Maturation and fusion of endosomes and lysosomes were inhibited by 15.5

µg/mL (30 µM) Chloroquine (Invitrogen) and 10 nM BafilomycinA1 (Sigma). Actin polymerization was

inhibited by 125 ng/mL Cytochalasin B (Sigma). Tubulin polymerization and function were inhibited by 250

pg/mL Colchicine, while an autophagy-related degradation was induced by 50 ng/mL Rapamycin. The selected

concentrations were determined based both on literature data and our preliminary experiments (Fig.3_S1).

Cellular metabolic activity was determined by a metabolic activity-based Resazurin assay[31]. Fresh cell culture

medium was added to control cultures a day before the *in situ* fixation. Reagents were diluted in fresh cell

culture medium. Leica TCS SP8 Lightning confocal microscope was used for detection of amphiectosome

release. A few hundred µm² sized area with 15-20 µm in height was tile-scanned with a few hundred cells. The

MV-lEVs were recognized as CD63 positive EVs surrounded by GFP positive membrane. They were counted

and were normalized to the number of nuclei.

Western blotting

Presence of proteins and specificity of the used primary antibodies were confirmed by Western blotting as

described previously[31]. For accurate quantification (free of variations potentially caused by EV purification), we

analyzed cell-, serum- and lEV- free conditioned medium. The cells were cultured O/N in a serum-free culture

medium. After harvesting, the cells were eliminated by centrifugation (300 g, 10 min at 4 °C) followed by a

2.000g centrifugation (30 min at 4 °C) to eliminate lEVs. Total protein content of the conditioned, serum-, cell-

and lEV-free medium was precipitated by trichloroacetic acid as described previously[31,33]. The protein pellets

were suspended in in cOmplete Protease Inhibitor Cocktail (Roche) containing radio-immunoprecipitation assay

(RIPA) buffer.

When whole cell lysate was tested for validation of antibodies and the HEK293T-PalmGFP-LC3RFP cell line,

cells were lysed in cOmplete Protease Inhibitor Cocktail (Roche) containing RIPA buffer.

Polyacrylamide gel electrophoresis was carried out using 10 % gels (acrylamide/bis-acrylamide ratio 37.5:1) or

any kDa precast gels (Biorad) and a MiniProtean (BioRad) gel running system. For better solubilization of

membrane proteins, equal volumes of 0.1 % TritonX-100, Laemmli buffer and samples were mixed as described

previously[34]. Approximately 10-30 µg protein were loaded into each well. Following electrophoretic separation,

proteins were transferred to PVDF membranes (Serva). Membranes were blocked with 5 % skimmed milk

powder or 5 % BSA in washing buffer for 1 h. Primary antibodies were applied in 1:1000 dilution with the

exception of the anti-CD63 (rabbit), anti-CD81 (rabbit) and anti-CD81 (mouse) antibodies where 1:500, 1:2500

and 1:100 dilutions were used, respectively. Peroxidase-labelled secondary antibodies were applied in 1:10 000

dilution. The signals were detected by ECL Western Blotting Substrate (Thermo Scientific) with an Imager

CHEMI Premium (VWR) image analyzer system. In case of quantification, equal protein amounts were loaded

to the gels. Within a biological replicate, the control and Chloroquine-treated samples were run on the same gels.

To enable comparison, the relative expression of control and Chloroquine-treated samples were determined and

compared.

Software and statistical analysis

For image capturing, analysis and co-localization studies, Leica LAS X, Zeiss ZEN Blue, Olympus iTEM and

Image J software were used. Figures and graphs were generated using GraphPad Prism 9.4.1 and Biorender

(BioRender.com). For statistical analysis, standard deviation was calculated. Unpaired two tailed student t-tests

and one-way ANOVA were used (* $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$).

Acknowledgements

This research was funded by the NVKP_16-1-2016-0004 grant of the Hungarian National Research,

Development and Innovation Office (NKFIH), ÚNKP-23-3-I-SE-2, the Semmelweis Innovation Fund (STIA

2020 KFI), Hungarian Scientific Research Fund (OTKA K120237 and FK 138851), Eötvös Loránd University

Excellence Fund (EKA 2022/045-P101), Hungary Academy of Sciences, LP2022-13/2022, VEKOP-2.3.2-16-

2016-00002, VEKOP-2.3.3-15-2017-00016, the Higher Education Excellence Program (FIKP) and the

Therapeutic Thematic Programme TKP2021-EGA-23. This study was also supported by the grants RRF-2.3.1-

21-2022-00003 (National Cardiovascular Laboratory Program) and 2019-2.1.7-ERA-NET-2021-00015. The

project has received funding from the EU's Horizon 2020 Research and Innovation Programme under grant

agreement No. 739593.

The authors would like to thank Gábor Valcz for for his invaluable support throughout the research process, for

critically reviewing the manuscript and for his insightful comments. The authors would like to thank the ZEISS

Microscopy Customer Center Team for the collaboration, and Dr. Abel Pereira da Graça for the possibility to use

the Elyra7 SIM[2] super-resolution microscope. The authors are grateful to Györgyné Vidra, Györgyi Balogh and

Andrea Orbán for their technical help and advises.

Conflict of interest

References

Buzas, E. I. The roles of extracellular vesicles in the immune system. *Nat Rev Immunol* **23**, 236-250 (2023). https://doi.org:https://doi.org/10.1038/s41577-022-00763-8

György, B. *et al.* Membrane vesicles, current state-of-the-art: emerging role of extracellular vesicles. *Cellular and Molecular Life Sciences* **68**, 2667-2688 (2011). https://doi.org:10.1007/s00018-011-0689-3

Théry, C. *et al.* Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *Journal of Extracellular Vesicles* **7**, 1535750 (2018). https://doi.org:10.1080/20013078.2018.1535750

Murrow, L., Malhotra, R. & Debnath, J. ATG12–ATG3 interacts with Alix to promote basal autophagic flux and late endosome function. *Nature Cell Biology* **17**, 300-310 (2015). https://doi.org:10.1038/ncb3112

Guo, H. *et al.* Atg5 Disassociates the V1V0-ATPase to Promote Exosome Production and Tumor Metastasis Independent of Canonical Macroautophagy. *Developmental Cell* **43**, 716-730.e717 (2017). https://doi.org:https://doi.org/10.1016/j.devcel.2017.11.018

Leidal, A. M. *et al.* The LC3-conjugation machinery specifies the loading of RNA-binding proteins into extracellular vesicles. *Nature Cell Biology* **22**, 187-199 (2020). https://doi.org:10.1038/s41556-019-0450-y

Minakaki, G. *et al.* Autophagy inhibition promotes SNCA/alpha-synuclein release and transfer via extracellular vesicles with a hybrid autophagosome-exosome-like phenotype. *Autophagy* **14**, 98-119 (2018). https://doi.org:10.1080/15548627.2017.1395992

Berg, T. O., Fengsrud, M., Strømhaug, P. E., Berg, T. & Seglen, P. O. Isolation and Characterization of Rat Liver Amphisomes: EVIDENCE FOR FUSION OF AUTOPHAGOSOMES WITH BOTH EARLY AND LATE ENDOSOMES*. *Journal of Biological Chemistry* **273**, 21883-21892 (1998). https://doi.org:https://doi.org/10.1074/jbc.273.34.21883

Fader, C. M., Sánchez, D., Furlán, M. & Colombo, M. I. Induction of Autophagy Promotes Fusion of Multivesicular Bodies with Autophagic Vacuoles in K562 Cells. *Traffic* **9**, 230-250 (2008). https://doi.org:https://doi.org/10.1111/j.1600-0854.2007.00677.x

Jeppesen, D. K. *et al.* Reassessment of Exosome Composition. *Cell* **177**, 428-445.e418 (2019). https://doi.org:https://doi.org/10.1016/j.cell.2019.02.029

Valcz, G. *et al.* En bloc release of MVB-like small extracellular vesicle clusters by colorectal carcinoma cells. *Journal of Extracellular Vesicles* **8**, 1596668 (2019). https://doi.org:10.1080/20013078.2019.1596668

Mathieu, M. *et al.* Specificities of exosome versus small ectosome secretion revealed by live intracellular tracking of CD63 and CD9. *Nature Communications* **12**, 4389 (2021). https://doi.org:10.1038/s41467-021-24384-2

Kovács, K. D. *et al.* Nanoinjection of extracellular vesicles to single live cells by robotic fluidic force microscopy. *Journal of Extracellular Vesicles* **12**, 12388 (2023). https://doi.org:https://doi.org/10.1002/jev2.12388

Klionsky, D. J. *et al.* Guidelines for the use and interpretation of assays for monitoring autophagy (4th edition)1. *Autophagy* **17**, 1-382 (2021). https://doi.org:10.1080/15548627.2020.1797280

Xie, D. *et al.* Autophagy Contributes to the Rapamycin-Induced Improvement of Otitis Media. *Frontiers in Cellular Neuroscience* **15** (2022).

Chen, P. M., Gombart, Z. J. & Chen, J. W. Chloroquine treatment of ARPE-19 cells leads to lysosome dilation and intracellular lipid accumulation: possible implications of lysosomal dysfunction in macular degeneration. *Cell Biosci* **1**, 10 (2011). https://doi.org/10.1186/2045-3701-1-10

Wang, R. *et al.* Molecular basis of V-ATPase inhibition by bafilomycin A1. *Nature Communications* **12**, 1782 (2021). https://doi.org:10.1038/s41467-021-22111-5

557   18   Edgar, J. R., Manna, P. T., Nishimura, S., Banting, G. & Robinson, M. S. Tetherin is an
558   exosomal tether. *eLife* **5**, e17180 (2016). https://doi.org:10.7554/eLife.17180
559   19   Ortega, F. G. *et al.* Interfering with endolysosomal trafficking enhances release of bioactive
560   exosomes. *Nanomedicine: Nanotechnology, Biology and Medicine* **20**, 102014 (2019).
561   https://doi.org:https://doi.org/10.1016/j.nano.2019.102014
562   20   Zhang, Z. *et al.* Rapamycin maintains NAD(+)/NADH redox homeostasis in muscle cells. *Aging
563   (Albany NY)* **12**, 17786-17799 (2020).
564   21   Tanida, I., Ueno, T. & Kominami, E. LC3 and Autophagy. *Methods Mol Biol* **445**, 77-88 (2008).
565   https://doi.org:10.1007/978-1-59745-157-4_4
566   22   Liang, H. *et al.* The formation of migrasomes is initiated by the assembly of sphingomyelin
567   synthase 2 foci at the leading edge of migrating cells.
568   23   Ma, L. *et al.* Discovery of the migrasome, an organelle mediating release of cytoplasmic
569   contents   during   cell   migration.   *Cell   Research*   **25**,   24-38   (2015).
570   https://doi.org:10.1038/cr.2014.135
571   24   Baselga, M. *et al.* Spheresomes are the main extracellular vesicles in low-grade gliomas.
572   *Scientific Reports* **13**, 11180 (2023). https://doi.org:10.1038/s41598-023-38084-y
573   25   Junquera, C. *et al.* Biogenesis of a new type of extracellular vesicles in gastrointestinal
574   stromal tumors: ultrastructural profiles of spheresomes. *Histochem Cell Biol* **146**, 557-567
575   (2016). https://doi.org:10.1007/s00418-016-1460-5
576   26   Petersen, J. D., Mekhedov, E., Kaur, S., Roberts, D. D. & Zimmerberg, J. Endothelial cells
577   release microvesicles that harbour multivesicular bodies and secrete exosomes. *Journal of
578   Extracellular Biology* **2**, e79 (2023). https://doi.org:https://doi.org/10.1002/jex2.79
579   27   Lai, C. P. *et al.* Visualization and tracking of tumour extracellular vesicle delivery and RNA
580   translation   using   multiplexed   reporters.   *Nature   Communications*   **6**,   7029   (2015).
581   https://doi.org:10.1038/ncomms8029
582   28   Vukman, K. V. *et al.* An implanted device enables in vivo monitoring of extracellular vesicle-
583   mediated spread of pro-inflammatory mast cell response in mice. *Journal of Extracellular
584   Vesicles* **10**, e12023 (2020). https://doi.org:https://doi.org/10.1002/jev2.12023
585   29   Németh, K. *et al.* High fat diet and PCSK9 knockout modulates lipid profile of the liver and
586   changes the expression of lipid homeostasis related genes. *Nutrition & Metabolism* **20**, 19
587   (2023). https://doi.org:10.1186/s12986-023-00738-z
588   30   Németh, K. *et al.* Extracellular vesicle release and uptake by the liver under normo- and
589   hyperlipidemia.   *Cellular   and   Molecular   Life   Sciences*   **78**,   7589-7604   (2021).
590   https://doi.org:10.1007/s00018-021-03969-6
591   31   Koncz, A. *et al.* Endoplasmin Is a Hypoxia-Inducible Endoplasmic Reticulum-Derived Cargo of
592   Extracellular Vesicles Released by Cardiac Cell Lines. *Membranes* **13** (2023).
593   32   Olah, I., Kendall, C. & Glick, B. Differentiation of bursal secretory-dendritic cells studied with
594   anti-vimentin   monoclonal   antibody.   *The   Anatomical   Record*   **233**,   111-120   (1992).
595   https://doi.org:https://doi.org/10.1002/ar.1092330115
596   33   Koontz,   L.   TCA   precipitation.   *Methods   Enzymol*   **541**,   3-10   (2014).
597   https://doi.org:10.1016/b978-0-12-420119-4.00001-x
598   34   Visnovitz, T., Solti, Á., Csikós, G. & Fricke, W. Plasma membrane H+-ATPase gene expression,
599   protein level and activity in growing and non-growing regions of barley (Hordeum vulgare)
600   leaves. *Physiologia Plantarum* **144**, 382-393 (2012). https://doi.org:10.1111/j.1399-
601   3054.2012.01578.x

602