1 **Title**

2 Buckling instability underlies vertebral segmentation during axolotl tail regeneration
3
Authors

Wouter Masselink[1,2,*], Tobias Gerber[3,4], Vijayishwer Singh Jamwal[5,6], Francisco Falcon[1,2,7], Tom

Deshayes[1,8], Riley Grindle[5], Ryan P. Seaman[5], Marco Röcklinger[9], Sofia-Christina

Papadopoulos[5,10], Gisela Deneke[1], Adrija Adhikary[1], Orestis G. Andriotis[9], Marko Pende[5], Yuka

Taniguchi-Sugiura[1], Tzi-Yang Lin[1,7], Thomas Kurth[11], Jingkui Wang[1,2], Detlev Arendt[4], Ji-Feng

Fei[12], Barbara Treutlein[3, 13], Fred W. Kolling[14], Philipp J. Thurner[9], Joel H. Graber[5], Edouard

Hannezo[15], Elly M. Tanaka[1,2,*], Prayag Murawala[1,5,10,*]

11
1. Research Institute of Molecular Pathology (IMP), Vienna BioCenter (VBC), 1030 Vienna Austria.

2. Institute of Molecular Biotechnology of the Austrian Academy of Sciences (IMBA), Vienna

BioCenter (VBC), 1030 Vienna.

3. Max Planck Institute for Evolutionary Anthropology, 04103 Leipzig, Germany.

4. European Molecular Biology Laboratory (EMBL), Developmental Biology Unit, 69117

Heidelberg, Germany.

5. MDI Biological Laboratory (MDIBL), Bar Harbor, ME 04609, USA.

6. Department of Molecular & Biomedical Sciences, University of Maine, Orono, ME 04469, USA.

7. Vienna BioCenter PhD Program, Doctoral School of the University of Vienna and Medical

University of Vienna, 1030 Vienna, Austria.

8. École Normale Supérieure de Lyon, Université Claude Bernard Lyon 1, Université de Lyon, 69342

Lyon, France.

9. Institute of Lightweight Design and Structural Biomechanics, TU Wien, 1060, Vienna, Austria.

10. Department of Nephrology and Hypertension, Hannover Medical School, 30625 Hannover,
Germany.

11. Core Facility Electron Microscopy and Histology. Center for Molecular and Cellular
Bioengineering (CMCB), Technology Platform, Technische Universität Dresden, 01307, Dresden,
Germany.

12. Department of Pathology, Guangdong Provincial People's Hospital (Guangdong Academy of
Medical Sciences), Southern Medical University, 510080 Guangzhou, China

13. Department of Biosystems Science and Engineering, ETH Zurich, Basel 4058, Switzerland.

14. Dartmouth Cancer Center, Dartmouth Geisel School of Medicine, Lebanon, New Hampshire, USA

15. Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria

* Correspondence: wouter.masselink@imba.oeaw.ac.at, elly.tanaka@imba.oeaw.ac.at,
pmurawala@mdibl.org

Abstract

Primary body-axis development is a highly conserved process that proceeds through somitogenesis and subsequent subdivision into dermatome, myotome, and sclerotome. Defects in somitic-clock genes such as *Hes7* lead to vertebral-segmentation defects in mice and fish. Here we show that in the axolotl, although *Hes7* is necessary for proper embryonic vertebral segmentation, it is—surprisingly—dispensable during tail regeneration. We investigated the mechanism of vertebral segmentation during regeneration which initially occurs through extension of a cartilage rod ventral to the spinal cord. We find that the regenerating cartilage rod undergoes a periodic wrinkling that provides a template for vertebral segmentation. Via direct mechanical measurements and biophysical perturbations, we show that a model of compression-induced buckling instability can predict vertebral segmentation. The cartilage rod and other somitic derivatives (muscle, cartilage, tendon, fibroblasts) arise from tendon-like, *Lfng*⁺ multi-potent mesenchymal progenitors, which display a gene regulatory state distinct from somitic progenitors. In summary, we uncover a mechanism of vertebral segmentation during axolotl tail regeneration that is distinct from the somite-based developmental mechanism.

Main text

Regeneration of the axolotl tail, which is composed of a spinal cord surrounded by segmented vertebrae and muscle, is a rare example of primary body-axis regeneration among vertebrates. While the cellular and molecular basis of spinal-cord regeneration in axolotl has been characterized[1,2], very little is known about how somitic derivatives such as vertebrae are regenerated in axolotl. In vertebrate animals, embryonic development and regeneration have similar outcomes, but their starting points differ, and the question of when the pathways underlying regeneration converge with those underlying development has fascinated generations of researchers[3]. Beyond the differences in starting points, patterning during regeneration of post-embryonic structures occurs at much larger spatial scales compared to embryonic patterning, which suggests that different mechanisms could be at play. During vertebrate embryonic development, patterning of the primary body axis is dependent on somitogenesis. Somites are epithelial balls of cells on either side of the notochord and are specified through the clock-and-wavefront model[4,5]. Somites contain multipotent progenitors that respond to cues from the notochord and then lay out the architecture of the tail; these progenitors are the source of vertebrae and myomeres in addition to other connective-tissue cell types[6,7]. In the axolotl, regeneration takes place in the absence of the notochord and somites; instead, blastema cells condense into a cartilage rod, after which vertebrae emerge[8]. In this study we asked: what is the cell source and mechanism of vertebral patterning during regeneration?

Hallmarks of somitogenesis are absent during tail regeneration

We investigated two hallmarks of vertebral segmentation during embryogenesis and regeneration, i.e., the formation of epithelial somites, and their subsequent resegmentation. During

embryogenesis in axolotls and other vertebrates, the tail contains histologically distinct somites

that give rise to vertebrae, muscles, and other types of connective-tissue cells. We found that during

regeneration of the axolotl tail, the tail blastema does not contain histologically distinct somites,

in contrast to embryogenesis (Fig. 1a, b, Extended Data Fig. 1a). Instead, the earliest visibly

segmented structures during regeneration are myomeres which appear around 14 days post

amputation (dpa) (Fig.1b, Extended Data Fig. 1a). Nevertheless, despite the absence of somites,

both the number of myomeres and vertebrae are correctly re-established in the axolotl tail (Fig.

1c-e) (uninjured: 20.10 +/- 1.45 vs regenerated 19.20 +/- 2.10) [9]. Axolotl embryos, similar to many

other vertebrates, undergo resegmentation, where rostro-caudal specification of somites into

Uncx[+] and *Tbx18*[+] domains results in a half-segment off-set between myomeres and vertebrae

(Fig. 1f-h)[10–12]. Consistent with our observation of a histological absence of somites during

regeneration (Fig. 1b, Extended Data Fig. 1a), we found that somite-dependent resegmentation is

also absent during regeneration (Fig. 1i-l).

To understand the processes underlying re-establishment of periodic segments during

axolotl tail regeneration, we performed Xenium spatial transcriptomics (Fig. 2a) on stage-28

embryos and on tail blastemas at 3, 7, 10, and 14 dpa (Fig. 2b-e, Extended Data Fig. 1b-i, 2). We

selected 100 target genes to visualize not only cell and lineage identity but also pathways known

to participate in tissue patterning and signaling, including distal HOX genes, and the FGF, Notch,

retinoic acid (RA), Wnt, BMP, and SHH signaling pathways (Extended Data Table 1). Using

Xenium clustering analysis, we identified eight cell types in embryo and eight in tail blastema (Fig.

2c, e, Extended Data Fig. 2a, h, o).We found that both embryos and blastemas express distal HOX

genes, *Fgfr1*, *Wnt5a,* and the Wnt/B-cat downstream target *Aldh1a2*, as well as the cell identity

markers *Scx, Lfng,* and *Meox1*(Extended Data Fig. 1b-i, 2c-e, j-l, q-s). While *Hes7*, *Tbx6*, and

Mesp2 expression was present in the embryonic tail bud, it was absent in the tail blastema (Fig. 2f,

g, Extended Data Fig. 2f, m, t).

In axolotl embryos, we observed polarized somitic expression of resegmentation markers

Tbx18 and *Uncx4.1*, but neither was expressed in the tail blastema (Extended Data Fig. 1d-e, 2g,

n, u). Hematoxylin and eosin (H&E) staining and the absence of *Tbx18/Uncx4.1* rostro-caudal

polarity strongly argue against classic epithelial somites at blastema stages. Together, these results

show that while the tail bud and tail blastema exhibit some similarities in their expression of genes

associated with *Fgf*, *Wnt*, and RA signaling, we did not detect somite like epithelial units in the

blastema via H&E staining.

Hes7 is dispensable for vertebral segmentation during regeneration

To determine directly whether somitogenesis-dependent patterning takes place during

regeneration, we generated a somitic-clock mutant axolotl through targeted mutagenesis of *Hes7*.

The Hairy and enhancer of split (Hes/Her) family of bHLH transcription factors is a family of

highly conserved oscillatory genes involved in somitogenesis, including *Hes7* (Extended Data Fig.

3a, b), which shows variable expression in the axolotl tail bud consistent with a role as a somitic

oscillator (Fig. 2f)[13]. *Hes7* mutant mice display aberrant somite-boundary formation, resulting in

vertebral-segmentation defects[14,15]. To generate *Hes7* mutant axolotls, we injected eggs with

Cas9/gRNA complexes representing a mixture of four gRNAs targeting all three exons of the *Hes7*

gene sequence (Extended Data Fig. 3b). We were unable to recover homozygous *Hes7* mutant

axolotls, as they are apparently embryonic lethal. We could, however, analyze mosaic F0 crispants,

which escaped embryonic lethality. We imaged vertebrae of F0 crispants using Alizarin Red S in

live, 4-5-cm-long animals. As in mice, these mutant axolotls displayed irregular vertebral

segmentation and fusions (Fig. 2h, i) consistent with a role for *Hes7* in somitogenesis during

embryonic development. Surprisingly these mutant tails recover a wildtype phenotype following

amputation, at which time the tails display regular vertebral segmentation and the normal number

of tail vertebrae (Fig. 2j-l, Extended Data Fig. 3c-e). To ensure that this observation did not result

from positive selection of wildtype and heterozygotic cells in the regenerate, we analyzed editing

efficiencies at each exonic target site in the unamputated tail close to the amputation plane and in

the regenerated tails. We found no significant difference in editing efficiency among the target

sites (Fig. 2m), indicating that, vertebral segmentation during axolotl tail regeneration proceeds

without a detectable requirement for *Hes7* under mosaic loss-of-function conditions. In summary,

we show that while vertebral segmentation takes place during tail regeneration, it does so through

a non-canonical somite-independent program.

Somite-like lineage potential in the regenerating tail

We next set out to identify the source of the newly regenerated musculo-skeletal tissues in

the axolotl tail. To survey the potency of cells that regenerate the tail, we performed clonal barcode

lineage tracing in the regenerated axolotl tail using a CellTag-based strategy adapted to a third-

generation foamy virus vector system[16–18]. We generated three unique libraries, each containing a

9-nt barcode and a defined 6-nt tag (V1, V2, V3) (Fig. 3a, Extended Data Fig. 4a-c) associated

with the 5' end of a GFP-encoding insert. High-titer viral libraries (Extended Data Fig. 4d) were

used to infect each tail four days prior to amputation (V1), as well as four (V2) and seven (V3)

dpa (Fig. 3b, c). We recovered V1, V2, and V3 barcodes from single-cell RNA-sequencing

(scRNA-seq) data of flow-sorted GFP$^+$ cells at 21 dpa (Fig. 3d-e, Extended Data Fig. 4e-h). To

identify putative clones, we calculated the Jaccard similarity index of barcodes detected in cells

with somite-derivative signatures (tenocytes, intermediate fibroblasts, dermis, fin mesenchyme,

chondrocytes, and myogenic cells) because of fate-mapping studies presented in the next section

(Fig. 3f). This analysis identified clones ranging in size from 2-6 cells, which suggests an

approximately 6-19% recovery rate when compared to an expected average clone size of 32.89

(V1), 16.91 (V2), and 10.27 (V3) cells, assuming a blastema cell-cycle length of 100 hours[9]. Of

the 48 clones analyzed, 39 contained diverse combinations of 2-4 cell types (Fig. 3f). Notably,

myogenic cells showed a clonal relationship with each of the other five cell types. Moreover,

computational mixing of the two replicates and subsequent clone-calling resulted in correct

annotation of 47 of 48 clones, indicating a sufficient diversity of barcodes (Extended Data Fig. 4h-

i). Although we did not recover any one clone containing all six cell types, these data are

compatible with a model in which a multipotent progenitor in the tail that has potential similar to

that of pre-somitic mesoderm, and that may give rise to Dermo-Myo-Sclerogenic (DMS) lineages

during regeneration (Extended Data Fig. 4j).

Intermyotomal cells are the source of multipotent musculo-skeletal progenitors

We next aimed to identify and characterize the origin and nature of these progenitors,

hypothesizing that they may arise at least in part from connective tissue. We first grafted vertebrae

from a constitutively expressing *CAGGs:GFP* transgenic axolotl into an unlabeled host axolotl

and observed minimal contribution from vertebrae cells (Extended Data Fig. 5a-d), eliminating

vertebrae as a cell source. We then performed Cre/*lox*P genetic fate mapping of tail regeneration

using two connective-tissue-specific *Cre-ERT2*-expressing drivers (*Col1A2* and *Twist3*) in

combination with a *CAGGs:LoxP-STOP-LoxP-Cherry* reporter, and inducing recombination 10

days prior to tail amputation. Prior to amputation, the *Col1A2* driver broadly labeled the tail

connective-tissue population (fin mesenchyme, dermal fibroblasts, tenocytes, and periskeletal and

skeletal cells) (Fig. 4a). The *Twist3* driver labeled a smaller subset of tail connective tissue

including fin mesenchyme, dermal fibroblasts, and skeletal lineage, but showed very limited

labeling of connective-tissue cells residing at the intermyotomal boundaries (Fig. 4b). After tail

amputation, the *Twist3* driver showed limited participation of the labeled cells to the regenerate

(Fig. 4d, f, Extended Data Fig. 5e-h). In contrast, the *Col1A2* driver contributed labeled cells to

regenerated fin mesenchyme, tenocytes, and vertebrae as well as muscle (Fig. 4c, e, Extended Data

Fig. 5e-h), consistent with the notion of a common source cell for connective-tissue and muscle

lineages. This further excludes dermal fibroblasts and fin mesenchymal cells as the likely cell

source and suggests tenocytes as the likely cells source. To assess whether muscle labeling is due

to non-specific labeling of the myogenic lineage, such as satellite cells, we performed sectioning

and immunostaining with the muscle-progenitor marker PAX7. In converted, unamputated tails

we observed no colocalization between Cherry$^+$ cells and *PAX7$^+$* muscle progenitors (Extended

Data Fig. 5i). However, regenerated tails showed the presence of Cherry$^+$/PAX7$^-$, Cherry$^-$/PAX7$^+$,

and Cherry$^+$/PAX7$^+$ populations in addition to Cherry$^+$ muscle fibers, suggesting that *Col1A2*-

labeled Cherry+ cells give rise to at least some PAX7+ muscle progenitors (Extended Data Fig.

5j).

To molecularly pinpoint the potential source cells, we generated scRNAseq data of flow-

sorted cells from both the *Col1A2* and *Twist3* driver lines (Extended Data Fig. 6a) and annotated

six connective-tissue cell types (tenocytes, fibroblasts, chondrocytes, fin mesenchyme, dermis, and

periskeleton) and corroborated cell identities by HCR *in situ* hybridization (Extended Data Fig 6b-

h) to localize fin mesenchyme (*Prrx1$^+$*), dermal fibroblasts (*Prrx1^{low}/Twist3$^+$*), periskeletal cells

(*Chrdl1$^+$*), chondrocytes (*Sox9$^+$*), and tenocytes (*Scx$^+$*) at intermyotomal boundaries. Consistent

with tenocytes as the potential source cells we found tenocytes were depleted in our *Twist3* lineage

labelled populations. We found that genes expressed by tenocytes include *Lfng* and *Meox1*

*(*Extended Data Fig. 6b), which are also expressed during somitic mesoderm development

(Extended Data Fig. 1g), in addition to canonical tenocyte markers such as *Tnmd, Scx*, and *Mkx*

(Fig. 4i). Immunogold labeling of *Col1a2*-labeled cells identified Cherry+ cells between

myomeres that interact with muscle-fiber ends through collagen deposition (Extended Data Fig.

6i-j). Further, whole-body light-sheet imaging of *Col1A2* animals stained for PCNA (proliferative

cells) showed that Cherry$^+$/PCNA$^+$ cells reside in the intermyotomal space in mature tails

(Extended Data Fig. 6k). Together, our results suggest that tenocytes labeled by *Scx, Meox,* or *Lfng*

are a source population for dermo-, myo-, and sclerogenic lineages respectively in the tail. In

addition, based on these results and those of the viral barcoding experiment, we hypothesize that

these tenocytes represent multipotent DMS-progenitors in the mature tail.

To assess the trajectories and molecular features by which DMS-progenitors differentiate

during tail regeneration, we performed scRNA-seq analysis on the *Col1A2*-labeled lineages at one

and two weeks post-amputation (wpa) (Fig. 4g). Consistent with our hypothesis, trajectory analysis

implicated tenocytes as DMS-progenitors (Fig. 4 h, j), and later blastema time points showed

emergence of the myogenic lineage in the *Col1A2* driver dataset (Fig. 4g'). The transcriptional

signatures of all blastema cell clusters were similar to those identified in the mature tail (Fig. 4k,

l). This suggests that no unique cell cluster emerges during regeneration, and that DMS-progenitors

represent a resident progenitor population of the mature tail. Using HCR-ISH of *Lfng$^+$/Meox1$^+$* as

a marker for DMS-progenitors, we examined their localization during regeneration (Extended Data

Fig. 7a-f). At 1 wpa, we found accumulation of DMS-progenitors throughout the blastema

(Extended Data Fig. 7a-c), while at 2 wpa, myomeres separated by DMS-progenitors emerged

(Extended data Fig. 7d-f).

To test whether *Lfng*-expressing cells in the intermyotomal space indeed represent source

cells for regeneration, we generated an *Lfng*-driven Cre-ERT2-expressing transgenic axolotl line.

Conversion of the *Lfng* driver 10 days prior to regeneration showed that Cherry expression was

absent in the fin mesenchyme and was restricted to the inter-myotome population of the medial

axolotl tail (Fig. 4m). Upon tail amputation, the *Lfng* descendants contributed extensively to all

somitic mesodermal lineages of the regenerating tail (Fig. 4n, o, Extended Data Fig. 5e-h),

suggesting faithful labeling of DMS-progenitors. To examine whether *Lfng*-labeled DMS-

progenitors are necessary for tail regeneration, we crossed the *Lfng*-driven CreERT2-expressing

transgenic axolotl line to a *lox*P-dependent NTR2.0 (second-generation of nitroreductase)

transgenic line (*CAGGS:LoxP-nBFP-LoxP-Cherry-T2A-NTR2.0*[19]). Upon nifurpirinol treatment,

we found that ablation of *Lfng*-labeled DMS-progenitors prevented complete tail outgrowth (Fig.

4p-t). We did not quantify ablation efficiency or lineage selectively; therefore, incomplete or off-

target ablation effects cannot be fully excluded. Based on these results, together, we conclude that

Lfng[+] DMS-progenitors contribute to regeneration of somitic lineages and are required for

complete tail outgrowth in our ablation paradigm.

DMS-progenitors represent a non-embryonic identity

Our *Hes7*-mutant and scRNA-seq data suggest that regeneration of somitic mesoderm

lineages in axolotl differs from the embryonic program. This would contrast with regeneration of

limbs in axolotl, where limb connective-tissue cells dedifferentiate to a state that is molecularly

comparable to a limb-bud mesenchyme progenitor[20]. To further investigate this contrast in

processes, we integrated all mature and regeneration timepoints into a single UMAP (Extended

Data Figure 8a-c) and performed scRNAseq on four tail-bud stages (stage 25, 28, 30, 35). Results

showed that, consistent with our earlier Xenium spatial transcriptomic analysis, genes commonly

associated with somitogenesis and resegmentation were expressed during embryonic development

but did not show increased expression during tail regeneration compared to uninjured tails (Fig.

5a-b). However, we did detect expression of the DMS markers *Lfng* and *Meox1* in the combined

mature dataset (Fig. 5b). In addition, we observed expression of two highly conserved markers of

resegmentation (*Uncx4.1* and *Tbx18*) in the embryo via scRNAseq, Xenium spatial

transcriptomics, and HCR-ISH (Fig. 5b, Extended Data Fig. 1h, 8c, d), but neither technique

showed expression of these markers during tail regeneration (Fig. 5b, Extended Data Figure 1i,

2g, n, u 8c, e) Finally, to assess whether tail blastema cells acquire tail-bud-like identities, we

performed quadratic programing on published limb[21] and the above tail development and

regeneration datasets. Results show that unlike limb blastema cells which acquire a limb-bud

identity, tail blastema cells do not acquire a tail-bud identity (Fig. 5c). In summary, DMS-

progenitors have a distinct transcriptional profile and regenerate tail through a process

transcriptionally distinct from that used by embryonic somitic mesodermal cells.

Scalable vertebral segmentation takes place via buckling instability

To understand the distinct mechanisms of vertebral segmentation during regeneration, we

developed a novel vertebrae extirpation assay, which uncouples vertebral segmentation from axis

elongation. By 1 wpi, the injury site collapsed (Fig. 6a, b). Subsequently, by 2 wpi cartilage

condensation was observed and the injury site expanded (Fig. 6a, b). By 3 wpi, a stereotypical

wrinkling pattern had formed on the outer surface of the regenerating cartilage rod (Extended Data

Fig. 9a-g). The wavelength of these wrinkles matches the wavelength of the vertebrae that form at

4 wpi (Fig. 6a, Extended Data Fig. 9a-g) Our combined observations of (i) the collapse of the

injury site, and (ii) subsequent expansion of the cartilage rod and wrinkling before vertebrae

patterning, suggest a buckling instability due to compression, a mechanism that has been proposed

to underlie pattern formation during development of tubular structures in different organisms[22–25].

Buckling instability is a mechanics-based process of periodic pattern formation, which, unlike

morphogen- or cell-based processes, is not intrinsically constrained in its scaling. This property is

particularly relevant given that axolotl can regenerate their tails at all stages of their life. In this

model, under weak compression or a lack of it, a rod maintains a straight and stable shape. Above

a critical level of compression, stresses are relaxed locally, producing periodic surface wrinkling,

with a wavelength that depends only on a few geometrical and mechanical parameters of the rod.

More specifically, we developed a mechanical theory for growth in confined elastic tubes

comprising a shell and a core, suggesting two regimes of wrinkling instability (See materials and

methods section 'Buckling instability models' for details): i) a geometry-dominant model for tubes

with very soft cores, which exhibit very little mechanical resistance to compression; in this model,

the wavelength (λ , which in our theory represents the distance between vertebrae) depends only

on tube geometry, scaling as $\lambda \propto R \left(\frac{h}{(1-v^2)^{\frac{1}{2}} R} \right)^{\frac{1}{2}}$ (λ: wavelength, R:radius, h: shell height, v:

Poisson ratio) and not on any stiffness parameters; and ii) a stiffness-dominant model for tubes

with cores that have a non-negligible stiffness; in this model, the wavelength is dependent on the

relative thickness of the shell and the core, scaling as $\lambda \propto h \left(\frac{E_s}{E_c} \right)^{\frac{1}{3}}$ (λ: wavelength, h: shell height,

E_s: shell Young's modulus, E_c:core Young's modulus), similar to previous theories of buckling

induced wrinkling during the formation of crypts and villi, and epithelial folding[25–28]. To determine

which model best fits vertebral segmentation during axolotl regeneration, we first measured the

mechanical properties of the regenerating tube, i.e., the axolotl cartilage rod, at 4 wpi, using atomic

force microscopy (AFM). We found that the mean Young's modulus of the core is 1.7 kPa, and

that of the shell is 6.9 kPa, which together demonstrate a soft core and stiffer shell (Fig. 6c, e,

Extended Data Fig. 9h, i). We then quantified the shell thickness (h) and tube radius (R), obtaining

mean values of h = 91 μm (Fig. 6f) and R = 450 μm (Extended Data Fig. 9j). Together, these

measurements are consistent with an intermediate regime between the two models described

above, with a theoretical wavelength of ~500 μm, closely matching our experimental observations

in a parameter-free manner (Fig. 6h, Extended Data Fig. 9g). The geometry-dominant model

predicts that the wavelength (λ) should increase as the cartilage rod radius (R) increases. To test

this we took advantage of the variable vertebrae length across the axolotl tail by recording both

the wavelength (λ) and the diameter for each newly formed vertebra at 3 different amputation

planes (4, 10, and 16 myomeres posterior to the cloaca) and used this to plot λ and radius $\sqrt{R}$. We

found that regardless of the amputation plane, our simple geometric prediction of $\lambda \propto \sqrt{R}$ provides

an excellent fit for the data (Fig. 6g, h).

A key prediction of the stiffness-dominant model is that altering the relative contributions

of the shell and core mechanics should change the vertebral wavelength (λ). To test this, we treated

the regenerating cartilage rod in the vertebrae extirpation assay with β-aminopropionitrile (BAPN),

an inhibitor of LOX-mediated collagen crosslinking[29,30]. AFM measurements at 4 wpi

demonstrated that the treatment with 100 μM BAPN significantly lowered the shell's Young's

modulus and reduced the shell thickness (Fig. 6c-f, Extended Data Fig. 9h, i), both predicting a

reduced wavelength, which should result in a larger number of vertebrae within a given length

unit. Indeed, we found that treatment with BAPN resulted in a larger number of regenerating

vertebrae compared to control in the extirpation model (Fig. 6i-k). Furthermore, during tail

regeneration, transient treatment with 10 μM BAPN induced a transient reduction in the $\lambda/\sqrt{R}$

ratio, highlighting the progressive nature of buckling-instability-guided vertebrae patterning

during tail regeneration (Fig. 6l, m, Extended Data Fig. 9k). Interestingly our theory assumes that

external constraints from surrounding tissues prevent other modes of buckling, such as global

deformation of the entire rod, which is expected under free boundary conditions. To test this, we

treated the regenerating tail where one end of the rod is free with 100 μM BAPN. Strikingly, this

induced global spatial oscillations of the entire rod (Fig. 6l-o) that are fully consistent with our

buckling model (Fig. 6j)[31,32], and demonstrates that the cartilage rod is under a compressive load

in both regeneration paradigms.

Together, these results support a quantitatively predictive buckling-instability model for

vertebral segmentation during regeneration, a mechanism that inherently confers scalability to the

system during regeneration (Fig. 6o).

Conclusions

Here we show that regeneration of somitic mesodermal lineages in the axolotl tail proceeds

through mechanisms that are both morphologically and molecularly distinct from those that take

place during embryonic development. In contrast to embryonic somitogenesis, the somitic clock

gene *Hes7* plays no detectable role in vertebral segmentation during regeneration. Instead,

segmentation emerges through a mechanically driven buckling instability. We identify the cellular

source of regenerated somitic mesoderm as *Lfng*[+]/*Meox1*[+] tenocytes, which we term Dermo-Myo-

Sclero (DMS) progenitors, located at the intermyotomal space. This regenerative strategy does not

closely mirror axolotl limb regeneration, where fibroblasts broadly dedifferentiate to an embryonic

limb-bud–like state[20,21]; rather, the tail regenerative strategy is similar to neural regeneration,

where defined adult stem cells undergo lineage-restricted neurogenesis[33].

Previous lineage-tracing studies in larval zebrafish identified a *Col1a2*-expressing connective-

tissue cell population capable of contributing to muscle differentiation[34,35], potentially through

activation of MEOX1[36]. However, because adult zebrafish do not regenerate their primary body

axis, the relevance of this progenitor-like state to vertebral segmentation and axial regeneration in

zebrafish remains unresolved.

Although buckling instability has not been previously proposed as a mechanism for

vertebral segmentation, our findings show that this model is a particularly good fit for vertebral

segmentation during regeneration. Specifically, because the wavelength of buckling is set by tissue

geometry and material properties, vertebral segmentation can automatically scale with the size and

growth state of the regenerating tail. This provides a robust means of generating proportionate,

periodic structures during large-scale tissue regeneration without reliance on a fixed molecular

prepattern. Such mechanically driven scaling mechanisms may represent a general strategy for

regenerating other large, periodic anatomical structures and warrants investigation beyond the

vertebral column.

Our findings show conceptual parallels with vertebral patterning during zebrafish

development[37–40]. In crown teleosts, including zebrafish, classical *Tbx18/Uncx4.1*-based

resegmentation is lost, and vertebral centers are specified through a mechanically mediated process

during which vertical myosepta transmit tension to the notochord, leading to periodic induction of

vertebral elements. While axolotl tail regeneration and zebrafish development employ distinct

molecular players, both rely on mechanical forces that reshape the tissue precursor from which the

vertebral column emerges. Whether this similarity reflects convergent evolution or a shared

ancestral mechanism remains unclear. Tail regeneration is widespread across Euteleostomes,

spanning both Sarcopterygii and Actinopterygii. Euteleostome species capable of tail regeneration

include the banded knifefish, African lungfish, the extinct early tetrapod *Microbrachis pelikani,*

and lizards[41–45]. Notably, vertebral segmentation has been documented in African lungfish and in

Microbrachis pelikani, but has not been observed in lizards[41,42,44,45].

Conceptually, our work reveals that segmented vertebrae can arise through two

fundamentally distinct pathways: an embryonic, clock-based mechanism and a post-embryonic

regenerative mechanism. This distinction has important implications for regenerative medicine, as

it suggests that faithful reconstruction of complex axial structures may not require reactivation of

embryonic programs. Instead, exploiting mechanically guided patterning of adult progenitor

populations may offer an alternative and potentially more accessible route to vertebral column

regeneration in other species, including humans.

Acknowledgments

We would like to acknowledge the valuable services provided by the animal facility, Molecular

Biology Services, and BioOptics facilities at the IMP and IMBA; Histology, and NGS sequencing

core facilities at the Vienna Biocenter; the Comparative Animal Models Core, Light Microscopy

Facility (LMF; RRID:SCR_019166), and Comparative Genomics and Data Science Core at

MDIBL; MDIBL receives support from the COBRE (P30GM154610) and Maine INBRE grant

(P20GM103423) from NIH-NIGMS. Xenium spatial transcriptomics was carried out in the

Genomics and Molecular Biology Shared Resource (RRID:SCR_021293) at Dartmouth Cancer

Center, which is supported by NCI Cancer Center Support Grant 5P30CA023108 and NIH S10

(1S10OD030242) awards. We thank Dr. Stephen Sampson for proofreading this manuscript.

Competing interests

The authors declare that no competing interests exist.

Funding sources

WM is supported by an FWF Lise Meitner fellowship DOI 10.55776/M2444, project grant DOI 10.55776/P34841, and project grant DOI 10.55776/PAT4101524; TD is supported by a scholarship from the Ecole Normale Supérieure de Lyon, France; EMT is supported by an ERC AdG 742046 of the European Commission; PM is supported by grants from the NIH [NIGMS-COBRE (P20GM104318) and ORIP (R21OD031971)] and DFG-527098031.

This research was funded in whole, or in part, by the Austrian Science Fund (FWF) DOI 10.55776/M2444, 10.55776/P34841, 10.55776/PAT4101524 and the European Research Council (ERC) RegenMems 742046. For the purpose of open access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission.

Author contributions

Conceptualization: WM, TG, FF, EH, EMT, PM

Methodology: WM, TG, VSJ, FF, RG, RPS, MP, YST, TYL, TK, JW, JFF, FWK, JHG, EH, PM, MR, OGA, PJT, TD, GD, AA, YTS, TYL, TK, JW, EH, BT, DA

Investigation: WM, TG, VSJ, FF, RG, RPS, SCP, MP, JFF, TD, YST, TYL, TK, FWK, JHG, EH, PM, MR, OGA, TD, GD, AA, YTS, TYL, TK, JW, EH

References:

1. Rodrigo Albors, A. *et al.* Planar cell polarity-mediated induction of neural stem cell expansion during axolotl spinal cord regeneration. *eLife* **4,** e10230 (2015).

2. Cura Costa, E., Otsuki, L., Rodrigo Albors, A., Tanaka, E. M. & Chara, O. Spatiotemporal control of cell cycle acceleration during axolotl spinal cord regeneration. *eLife* **10**, e55665 (2021).

3. Poss, K. D. & Tanaka, E. M. Hallmarks of regeneration. *Cell Stem Cell* **31**, 1244–1261 (2024).

4. Cooke, J. & Zeeman, E. C. A clock and wavefront model for control of the number of repeated structures during animal morphogenesis. *J. Theor. Biol.* **58**, 455–476 (1976).

5. Baker, R. E., Schnell, S. & Maini, P. K. A clock and wavefront mechanism for somite formation. *Developmental Biology* **293**, 116–126 (2006).

6. Teillet, M. A. & Ledouarin, N. M. Consequences of Neural-Tube and Notochord Excision on the Development of the Peripheral Nervous-System in the Chick-Embryo. *Developmental Biology* **98**, 192–211 (1983).

7. Brand-Saberi, B. & Christ, B. 1 Evolution and Development of Distinct Cell Lineages Derived from Somites. in *Current Topics in Developmental Biology* vol. 48 1–42 (Elsevier, 1999).

8. Echeverri, K., Clarke, J. D. W. & Tanaka, E. M. In Vivo Imaging Indicates Muscle Fiber Dedifferentiation Is a Major Contributor to the Regenerating Tail Blastema. *Developmental Biology* **236**, 151–164 (2001).

9. Vincent, C. D., Rost, F., Masselink, W., Brusch, L. & Tanaka, E. M. Cellular dynamics

underlying regeneration of appropriate segment number during axolotl tail

regeneration. *BMC Dev Biol* **15**, 48 (2015).

10. Bussen, M. *et al*. The T-box transcription factor Tbx18 maintains the separation of

anterior and posterior somite compartments. *Genes & Development* **18**, 1209–1221

(2004).

11. Criswell, K. E. & Gillis, J. A. Resegmentation is an ancestral feature of the gnathostome

vertebral skeleton. *bioRxiv* **103**, 69–22 (2019).

12. Piekarski, N. & Olsson, L. Resegmentation in the Mexican axolotl, Ambystoma

mexicanum. *J. Morphol.* **275**, 141–152 (2014).

13. Krol, A. J. *et al*. Evolutionary plasticity of segmentation clock networks. *Development*

138, 2783–2792 (2011).

14. Bessho, Y. *et al*. Dynamic expression and essential functions of Hes7 in somite

segmentation. *Genes Dev.* **15**, 2642–2647 (2001).

15. Fujimuro, T. *et al*. Hes7 3'UTR is required for somite segmentation function. *Scientific*

Reports **4**, 6462 (2014).

16. Biddy, B. A. *et al*. Single-cell mapping of lineage and identity in direct reprogramming.

Nature **564**, 219–224 (2018).

17. Kong, W. *et al*. CellTagging: combinatorial indexing to simultaneously map lineage and

identity at single-cell resolution. *Nat Protoc* 1–25 (2020) doi:10.1038/s41596-019-0247-

2.

18. Trobridge, G. D. Foamy virus vectors for gene transfer. *Expert Opin Biol Ther* **9**, 1427–36

(2009).

19. García-García, D. *et al*. The essential role of connective-tissue cells during axolotl limb

regeneration. Preprint at https://doi.org/10.1101/2025.03.30.645595 (2025).

20. Gerber, T. *et al*. Single-cell analysis uncovers convergence of cell identities during

axolotl limb regeneration. *Science (New York, N.Y.)* **362**, eaaq0681-13 (2018).

21. Lin, T.-Y. *et al*. Fibroblast dedifferentiation as a determinant of successful regeneration.

Developmental Cell **56**, 1541-1551.e6 (2021).

22. Varner, V. D., Gleghorn, J. P., Miller, E., Radisky, D. C. & Nelson, C. M. Mechanically

patterning the embryonic airway epithelium. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 9230–

9235 (2015).

23. Gill, H. K. *et al*. The developmental mechanics of divergent buckling patterns in the

chick gut. *Proc. Natl. Acad. Sci. U.S.A.* **121**, e2310992121 (2024).

24. Shyer, A. E. *et al*. Emergent cellular self-organization and mechanosensation initiate

follicle pattern in the avian skin. *Science* **357**, 811–815 (2017).

25. Hannezo, E., Prost, J. & Joanny, J.-F. Instabilities of Monolayered Epithelia: Shape and

Structure of Villi and Crypts. *Phys. Rev. Lett.* **107**, 078104 (2011).

26. Hannezo, E., Prost, J. & Joanny, J.-F. Theory of epithelial sheet morphology in three

dimensions. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 27–32 (2014).

27. Shyer, A. E. *et al*. Villification: How the Gut Gets Its Villi. *Science* **342**, 212–218 (2013).

28. Santos-Durán, G. N., Cooper, R. L., Jahanbakhsh, E., Timin, G. & Milinkovitch, M. C.

Self-organized patterning of crocodile head scales by compressive folding. *Nature* **637**,

375–383 (2025).

29. Wilmarth, K. R. & Froines, J. R. In vitro and in vivo inhibition of lysyl oxidase

byaminopropionitriles. *Journal of Toxicology and Environmental Health* **37**, 411–423

(1992).

30. Makris, E. A., Responte, D. J., Paschos, N. K., Hu, J. C. & Athanasiou, K. A. Developing

functional musculoskeletal tissues through hypoxia and lysyl oxidase-induced collagen

cross-linking. *Proc. Natl. Acad. Sci. U.S.A.* **111**, (2014).

31. Timoshenko, S., P. & Gere, J., M. *Theory of Elastic Stability*. (McGraw-Hill, 1961).

32. Euler, L. Methodus inveniendi lineas curvas maximi minimive proprietate gaudentes

sive solutio problematis isoperimetrici latissimo sensu accepti. (1744).

33. Lust, K. *et al*. Single-cell analyses of axolotl telencephalon organization, neurogenesis,

and regeneration. *Science* **377**, eabp9262 (2022).

34. Ma, R. C., Jacobs, C. T., Sharma, P., Kocha, K. M. & Huang, P. Stereotypic generation of

axial tenocytes from bipartite sclerotome domains in zebrafish. *PLoS Genet* **14**,

e1007775 (2018).

35. Sharma, P., Ruel, T. D., Kocha, K. M., Liao, S. & Huang, P. Single cell dynamics of

embryonic muscle progenitor cells in zebrafish. *Development* dev.178400 (2019)

doi:10.1242/dev.178400.

36. Nguyen, P. D. *et al.* Muscle Stem Cells Undergo Extensive Clonal Drift during Tissue

Growth via Meox1-Mediated Induction of G2 Cell-Cycle Arrest. *Cell Stem Cell* **21**, 107-

119.e6 (2017).

37. Wopat, S. *et al.* Spine Patterning Is Guided by Segmentation of the Notochord Sheath.

CellReports **22**, 2026–2038 (2018).

38. Wopat, S. *et al.* Notochord segmentation in zebrafish controlled by iterative mechanical

signaling. *Developmental Cell* S1534580724002387 (2024)

doi:10.1016/j.devcel.2024.04.013.

39. Peskin, B. *et al.* Notochordal Signals Establish Phylogenetic Identity of the Teleost

Spine. *Current biology : CB* **30**, 2805-2814.e3 (2020).

40. Lleras Forero, L. *et al.* Segmentation of the zebrafish axial skeleton relies on notochord

sheath cells and not on the segmentation clock. *eLife* **7**, 15–4 (2018).

41. Verissimo, K. M. *et al.* Salamander-like tail regeneration in the West African lungfish.

Proceedings of the Royal Society B: Biological Sciences **287**, 20192939 (2020).

42. Fröbisch, N. B., Bickelmann, C. & Witzmann, F. Early evolution of limb regeneration in

tetrapods: evidence from a 300-million-year-old amphibian. *Proc Biol Sci* **281**,

20141550 (2014).

43. Ellis, M. M. *The Gymnotid Eels of Tropical America*. (Published by the authority of the

Board of Trustees of the Carnegie Institute, Pittsburgh :, 1913).

doi:10.5962/bhl.title.43182.

44. Vonk, A. C. *et al*. Single-cell analysis of lizard blastema fibroblasts reveals phagocyte-

dependent activation of Hedgehog-responsive chondrogenesis. *Nat Commun* **14**, 4489

(2023).

45. Fisher, R. E. *et al*. A Histological Comparison of the Original and Regenerated Tail in the

Green Anole, *Anolis carolinensis*. *The Anatomical Record* **295**, 1609–1619 (2012).

46. Khattak, S. *et al*. Germline transgenic methods for tracking cells and testing gene

function during regeneration in the axolotl. *Stem Cell Reports* **1**, 90–103 (2013).

47. Kawaguchi, A. *et al*. Chromatin states at homeoprotein loci distinguish axolotl limb

segments prior to regeneration. *bioRxiv* 2022.11.14.516253 (2022)

doi:10.1101/2022.11.14.516253.

48. Aulehla, A. *et al*. A β-catenin gradient links the clock and wavefront systems in mouse

embryo segmentation. *Nat Cell Biol* **10**, 186–193 (2008).

49. Fei, J.-F. *et al*. Efficient gene knockin in axolotl and its use to test the role of satellite

cells in limb regeneration. *Proceedings of the National Academy of Sciences of the*

United States of America **114**, 12501–12506 (2017).

50. Fei, J. F. *et al*. Application and optimization of CRISPR-Cas9-mediated genome

engineering in axolotl (Ambystoma mexicanum). *Nat Protoc* **13**, 2908–2943 (2018).

51. Kroll, F. *et al*. A simple and effective F0 knockout method for rapid screening of

behaviour and other complex phenotypes. *eLife* **10**, e59683 (2021).

52. Riquelme-Guzmán, C. & Sandoval-Guzmán, T. Methods for Studying Appendicular

Skeletal Biology in Axolotls. in *Salamanders* (eds Seifert, A. W. & Currie, J. D.) vol. 2562

155–163 (Springer US, New York, NY, 2023).

53. Pende, M. *et al.* A versatile depigmentation, clearing, and labeling method for exploring

nervous system diversity. *Sci Adv* **6**, (2020).

54. Kurth, T. Immunocytochemistry of the amphibian embryo--from overview to

ultrastructure. *Int J Dev Biol* **47**, 373–383 (2003).

55. Kurth, T. *et al.* Electron Microscopy of the Amphibian Model Systems Xenopus laevis

and Ambystoma mexicanum. in *Methods in Cell Biology* vol. 96 395–423 (Elsevier,

2010).

56. Wagner, F. *et al. Human Photoreceptor Cell Transplants Integrate into Human Retina*

Organoids. http://biorxiv.org/lookup/doi/10.1101/2022.08.09.500037 (2022)

doi:10.1101/2022.08.09.500037.

57. Choi, H. M. T. *et al.* Third-generation in situ hybridization chain reaction: multiplexed,

quantitative, sensitive, versatile, robust. *Development* **145**, dev165753 (2018).

58. Bensimon-Brito, A. *et al.* Revisiting in vivo staining with alizarin red S - a valuable

approach to analyse zebrafish skeletal mineralization during development and

regeneration. *BMC Dev Biol* **16**, 2 (2016).

59. Hutter, J. L. & Bechhoefer, J. Calibration of atomic-force microscope tips. *Review of*

Scientific Instruments **64**, 1868–1873 (1993).

60. Kain, L. *et al.* Calibration of colloidal probes with atomic force microscopy for

micromechanical assessment. *Journal of the Mechanical Behavior of Biomedical*

Materials **85**, 225–236 (2018).

61. Andriotis, O. G. *et al*. Nanomechanical assessment of human and murine collagen

fibrils via atomic force microscopy cantilever-based nanoindentation. *Journal of the*

Mechanical Behavior of Biomedical Materials **39**, 9–26 (2014).

62. Hannezo, E., Prost, J. & Joanny, J.-F. Mechanical Instabilities of Biological Tubes. *Phys.*

Rev. Lett. **109**, 018101 (2012).

63. Karam, G. N. & Gibson, L. J. Elastic buckling of cylindrical shells with elastic cores—I.

Analysis. *International Journal of Solids and Structures* **32**, 1259–1283 (1995).

64. Brangwynne, C. P. *et al*. Microtubules can bear enhanced compressive loads in living

cells because of lateral reinforcement. *The Journal of Cell Biology* **173**, 733–741 (2006).

65. Replogle, J. M. *et al*. Combinatorial single-cell CRISPR screens by direct guide RNA

capture and targeted sequencing. *Nat Biotechnol* **38**, 954–961 (2020).

66. Khattak, S. *et al*. Foamy virus for efficient gene transfer in regeneration studies. *BMC*

Dev Biol **13**, 9 (2013).

67. Wallace, H. & Maden, M. The cell cycle during amphibian limb regeneration. *Journal of*

Cell Science **20**, 539–547 (1976).

68. Schreckenberg, G. M. & Jacobson, A. G. Normal stages of development of the axolotl,

Ambystoma mexicanum. *Developmental Biology* **42**, 391–399 (1975).

69. Picelli, S. *et al*. Full-length RNA-seq from single cells using Smart-seq2. *Nat Protoc* **9**,

171–181 (2014).

70. Nowoshilow, S. *et al*. The axolotl genome and the evolution of key tissue formation

regulators. *Nature* 1–21 (2018) doi:10.1038/nature25458.

71. Schloissnig, S. *et al.* The giant axolotl genome uncovers the evolution, scaling, and

transcriptional control of complex gene loci. *Proceedings of the National Academy of*

Sciences **118**, e2017176118 (2021).

72. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21

(2013).

73. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell

transcriptomic data across different conditions, technologies, and species. *Nat*

Biotechnol **36**, 411–420 (2018).

74. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with

Harmony. *Nat Methods* **16**, 1289–1296 (2019).

75. Bergen, V., Lange, M., Peidli, S., Wolf, F. A. & Theis, F. J. Generalizing RNA velocity to

transient cell states through dynamical modeling. *Nat Biotechnol* **38**, 1408–1414

(2020).

76. Wolf, F. A. *et al.* PAGA: graph abstraction reconciles clustering with trajectory inference

through a topology preserving map of single cells. *Genome Biol* **20**, 59 (2019).

77. Angerer, P. *et al.* destiny: diffusion maps for large-scale single-cell data in R.

Bioinformatics **32**, 1241–1243 (2016).

Figure 1: Characterization of axolotl tail regeneration. a-b, Coronal section of a H&E-stained stage-30 axolotl embryo (a) and a 14-dpa tail blastema (b). Arrows indicate epithelialized somites (a), and arrowheads indicate segmented myomeres (b). **c-d,** Vertebral segmentation of the uninjured tail (c) is re-established in stage-matched 5-month regenerated tail (d), dashed line indicates amputation plane. **e,** Quantification of ossified vertebrae numbers posterior to the cloaca in uninjured (c) and stage-matched 5-month regenerated (d) animals (n= 10). **f-k,** Stereoscopic images of a *CAGGS:GFP* transgenic axolotl labeled with Alizarin red S in uninjured (f-h) and regenerated (i-k) tail. **l,** Quantification of muscle vertebrae off-set in the tail of h and k (n=112) (off-set is calculated based on the shortest distance from the middle of a vertebra to the nearest myomere boundary. Statistical analysis e: MZ-test. ns: not significant. l: F-test. ns: not significant, ** p< 0.01, *** p< 0.001. Dashed lines indicate amputation plane (ap). Scale bars: a: 200 μm, b: 500 μm, c-d: 1 cm. f-k: 1000 μm. h', k': 200 μm.

Figure 2: Absence of key somitic clock hallmarks during axolotl tail regeneration. a,
Distribution of 100 genes selected for Xenium spatial transcriptomics based on their function. **b-c,** Analysis of a representative sagittal section from a stage-28 embryo showing histology (b), spatial distribution (b'), and UMAP distribution (c) of identified cell types. **d-e,** Analysis of a representative coronal section from a 2- weeks post-amputation (wpa) tail blastema showing histology (d), spatial distribution (d'), and UMAP distribution (e) of identified cell types. **f-g,** Xenium expression patterns for *Hes7*, *Tbx6*, and *Mesp2* in stage-28 embryo (f), and 2-wpa tail blastema (g). Dashed line indicates amputation plane (ap). **h-k,** Stereoscopic images of Alizarin red S labeling of segmented vertebrae in uninjured noninjected control (h), Hes7 gRNA mix-injected axolotl (i), regenerated non-injected control (j), and regenerated Hes7 gRNA mix-injected axolotl (k). **l,** Scoring of vertebral fusion effects in the tail (h-k) (n=10). **m,** gRNA efficiency of all four Hes7 gRNAs in unamputated and regenerated tails. Statistical analysis: l: Kruskal-Wallis test, followed by Dunn's test. m: one-way ANOVA, followed by Šidák correction. ns: not significant, ** $p < 0.01$. Dashed lines indicate amputation plane (ap). Scale bar: b, d, f-k: 500 μm. f'-g'': 200 μm.

Figure 3: The axolotl tail displays a somite-like lineage potential. **a,** schematic overview of foamy virus barcode labeling vectors. **b,** Experimental overview of infection, amputation and harvesting timepoints. **c,** Representative image of a 7-dpa foamy-virus-infected axolotl tail at 3 days post infection (dpi). **d-e,** UMAP and dotplot representation of different clusters identified in GFP+ cells from foamy-virus-infected tails combining two independent replicates. **f,** Matrix-based depiction of identified clones with a Jaccard similarity index of ≥0.45 (barcodes present in ≥ 2 cells and ≥ 2 UMIs per cell). Each row represents a cell type. Each column represents a clone. Dashed lines indicate amputation plane (ap). Scale bar: c: 500 μm.

Figure 4: DMS-progenitors are necessary for tail regeneration. a-d, Labeling of 4-OHT-

treated *Col1a2:ER-Cre-ER;Caggs:lp-Cherry* and *Twist3:ER-CreER;Caggs:lp-Cherry* axolotl

tails before amputation (a,b), and at 6 wpa (c,d). **e,f**, Quantification of cell-type distribution at

three different positions (mature stump, proximal and distal regenerate, Extended Data Fig. 3g)

of the *Col1a2:ER-Cre-ER;Caggs:lp-Cherry* (e) and *Twist3:ER-CreER;Caggs:lp-Cherry* (f)

regenerated tails. **g,g'**, UMAP integration of cells from 7 and 14 dpa regenerated tails colored by

cell-type identity (left) and by timepoint (right). **h**, Dotplot representation of cell-type marker

genes (columns) across the cell type (rows) identified in g. **i**, Pseudotime estimates by a

diffusion-map analysis are visualized on the UMAP embedding. **j**, Line plots represent the

smoothed gene expression along their pseudotime separated by cell types. **k,l**, Heatmap for the

top 30 markers (rows) of each cell type (columns) in the uninjured scRNA-seq data (see

Extended Data Fig. 6a,b). The genes were used as input for a gene-correlation analysis visualized

as a heatmap with average values within each cell type summarized as a barplot (right). Mature

tail (k) and 14-dpa blastema (l) **m,n,** Labeling of 4-OHT-treated *Lfng:ER-Cre-ER;Caggs:lp-*

Cherry axolotl before amputation (m) and at 6 wpa (n). **o**, Quantification of cell-type distribution

of the regenerated *Lfng:ER-CreER;Caggs:lp-Cherry* tail, including the mature stump, proximal

and distal regenerates. DF: dermal fibroblasts, IF: interstitial fibroblast, PS: peri-skeleton, SK:

skeleton, Noto: notochord, MP: muscle progenitor, MF: muscle fiber. (Extended Data Fig. 3g). **p,**

Schematic depiction of nifurpirinol cell-ablation approach. **q-s**, Tail regenerative response after

nifurpirinol-based cell ablation in *Lfng:ER-Cre-ER;Caggs:lp-Cherry* transgenic axolotl. 4-OHT

treatment (p, p'), and nifurpirinol treatment (q, q') alone do not affect regeneration; only when

both 4-OHT and nifurpirinol treatments are combined is tail regeneration affected (r, r'). **t,**

Length of the regenerating tails shown in p-r. Dashed lines indicate amputation plane (AP). Scale

bars: a-d, p: 500 µm; a'-d'', p': 50 µm. m-n: 1,000 µm. m'-n''. 100 µm. p-s': 4,000 µm.

Figure 5: Tail blastema cells do not acquire a tail-bud identity during tail regeneration. a,b,

Dotplot representation comparing the expression of various somitogenesis (a) and

resegmentation (b) markers between tail-bud, mature-tail, and tail-blastema datasets. **c,**

Comparative quadratic programming shows that limb cells acquire embryonic limb-mesenchyme

identity during limb regeneration (from Lin et al.)[21] but tail cells do not acquire tail-bud identity.

Figure 6: Vertebral segmentation during tail regeneration takes place through buckling

instability. **a**, Stereoscopic images of local injury time series in *Col1a2:TFPnls* (green) axolotl

labeled with Alizarin red S (magenta). Three-vertebrae injury site is marked using a dashed box.

b, Quantification of injury-site length over a 5-week period, relative to the average injury-site

length at 0 wpi. **c-d**, H&E stain of a 4-wpi cross-section after local injury untreated **(c)** and 100

μM BAPN-treated **(d)**. **e**, Young's modulus of the core and shell with and without BAPN

treatment. **f**, Quantification of shell thickness with and without BAPN treatment. **g**, Stereoscopic

image of *Col1a2:TFPnls* (green) axolotl labeled with Alizarin red S (magenta) at 4 wpa,

depicting the recording of wavelength and diameter during tail regeneration. **h**, Quantification of

λ and $\sqrt{R}$ of newly added vertebrae throughout a 7-week regenerative response. Tails were

amputated at +4, +10, and +16 myomeres posterior to the cloaca. **i-j,** Regenerated vertebrae at 5

wpi without **(i)** and with **(j)** 100 μM BAPN treatment. **k**, Quantification of the number of

vertebrae at 5 wpi in the local injury relative to the aspect ratio of the regenerated cartilage rod

($L/\sqrt{R}$) **l-n**, Tail regeneration time series treated with 0 (l), 10 (m), and 100 μM BAPN (n). **o,**

Normalised tortuosity of the regenerated cartilage rod in l-n, Normalised tortuosity is calculated

as $\frac{Path\ length}{Euclidian\ length} - 1$. Statistical analysis: b: one-way ANOVA, followed by Šidák correction, e:

two-way ANOVA, followed by Šidák correction, f: unpaired t test. h, k: simple linear fit, dashed

lines indicate 95% confidence interval, o: two-way ANOVA, with Dunnett's multiple

comparison. Dashed lines indicate amputation plane (ap). Arrows indicate regenerated vertebrae.

Scale bar: a, g: 1,000 μm. c-d: 250 μm. i-j: 500 μm. l-n: 2,000 μm.

Materials and Methods

Animal husbandry, transgenesis, and 4-OH tamoxifen treatment

Axolotls (*Ambystoma mexicanum*) were bred in IMP and MDIBL facilities. All animal handling

and surgical procedures were carried out in accordance with local ethics-committee guidelines.

Animal experiments were performed as approved by the Magistrate of Vienna and the IACUC

committee of MDIBL. "White" refers to a non-transgenic *d/d* strain of axolotl that has leucistic

skin due to the absence of melanocytes. Animals were anaesthetized in 0.03% benzocaine (Sigma)

before amputation and surgery. Axolotl husbandry, transgenesis, and 4-hydroxytamoxifen (4-

OHT) treatment were done as previously described[43].

tgSceI(Mmu.*Col1a2:TFPnls-T2a-ER^T2-Cre-ER^T2*)^Etnka, tgSceI(*CAGGs:LoxP-GFP-LoxP-

Cherry*)^Etnka, tgSceI(*Caggs:LoxP-STOP-LoxP-Cherry*)^Etnka and tgSceI(*Caggs:Lp-BFPnls-LP-

Cherry-T2a-NTR2.0)^Pmx* transgenic lines are described previously[20,46,47]. To generate *Lfng*

transgenic animals, a mouse *Lfng*[48] driver element was cloned at the 5′ end of the GFPnls-T2A-

ER^T2-Cre-ER^T2 (ER-Cre-ER) cassette with flanking ISceI sites. The *Twist3* transgenic animal was

generated by knock-in methodology described previously[49]. *Twist3* is a single-exon gene. First, to

generate a targeting construct, genomic sequence containing 450 bp upstream seq of the 5'UTR of

Twist3 along with its 5'UTR and ORF (without stop codon) (total 1,230 bp) was amplified from

axolotl genomic DNA and was cloned at the 5' end of the MCS-T2a-GFPnls-T2a-ER^T2-Cre-ER^T2

base vector to generate 450bp_upstream_seq-5'UTR-Twist3ORF (without stop codon)-T2a-

GFPnls-T2a-ER^T2-Cre-ER^T2 construct. Next, embryos were co-injected with the targeting

construct, CAS9 protein and gRNA (GCTGCAGCCCATGCAATC-TGG) with a target site 295

bp upstream of the ORF to create dsDNA breaks in both the genome and the targeting construct.

Embryos were screened for nuclear GFP fluorescence at the 3-cm stage and grown up as potential

breeders. Successful germ-line transmission was evaluated by nuclear GFP fluorescence in dermal

fibroblasts across the body, and a line was established. Constructs used for generating transgenic

axolotl driver lines are available at Addgene (200104, 200105, 200106, and 200107).

Xenium spatial transcriptomics

Axolotl embryos were staged to developmental stage 28 after fertilization and fixed overnight in

4% paraformaldehyde (PFA; Sigma-Aldrich, 441244-3KG) at 4°C. Embryos were washed with

phosphate-buffered saline (PBS) and incubated in 2% gelatin (Sigma-Aldrich, G2500-100G) at 37

°C in a water bath for 1 hour and placed under vacuum for 30 minutes at room temperature. The

embryos were washed with ice-cold PBS containing 2% gelatin, and subjected to a sucrose

infiltration series (7.5%, 15%, and 30%). For cryopreservation and dehydration, the embryos were

equilibrated in a 30% sucrose: OCT compound solution (1:1) and embedded in OCT. Samples

were sectioned at a thickness of 20 µm using a cryo-microtome (Leica CM1860 UV). Tails were

collected from animals approximately 3 cm in length that had undergone tail amputation and were

allowed to regenerate for 14 days. On day 14, the animals were anesthetized with 0.03%

benzocaine, and the regenerated tails were harvested using a scalpel (Aspen Surgical Products

371621). The collected tails were arranged and aligned in a cryomold containing OCT compound

(Leica Biosystems 14020108926). Once properly positioned, the OCT was rapidly frozen by

placing the cryomold on a pre-chilled heat block submerged in liquid nitrogen. Cryosections were

prepared at a thickness of 20 µm and subsequently transferred onto Xenium slides for downstream

processing. Xenium panel design, tissue histology, and processing were carried out according to

the manufacturer's instructions for preparing Xenium slides from fresh frozen tissues (10x

Genomics, Protocol: CG000579), with the following modification: slides containing embryo

samples were fixed in methanol for 2hrs at room temperature (rather than the 30min stated in the

protocol). Samples were subjected to probe hybridization, ligation, and amplification steps, and

stained with the Cell Segmentation Add-On Kit (10x Genomics Protocol: CG000749). Xenium

slides were subsequently processed on a Xenium Analyzer instrument running Xenium Software

(v.2.0.1.0) and Onboard Analysis Software (v.2.0.0.10), generating the output data bundle

including principal component analysis (PCA), K-means and graph-based clustering, differential

gene expression analysis, and UMAP (Uniform Manifold Approximation and Projection)-based

dimensionality reduction for downstream analyses. A comprehensive list of all detected transcripts

is available in GEO: GSE313338. The custom panel was designed to target 100 genes associated

with somitogenesis and cell-type classification within the blastema. Each target gene was assigned

6–8 probes, designed based on the provided gene sequences. Final gene selection and probe

numbers were validated by cross-referencing with unfiltered single-cell transcriptomic data to

ensure balanced representation across cell types and to prevent oversaturation of any single

population, which could otherwise obscure distinct fluorescent signals. Slides were stained with

hematoxylin and eosin (H&E) using a Sakura Tissue-Tek Prisma stainer and imaged at 40×

magnification with an Aperio GT450 scanner (Leica). Images were exported from the Xenium

Browser in PNG format, corresponding to subcellular, near single-molecule RNA resolution.

Exported images were imported into Fiji (ImageJ, NIH). Each image was converted to 8-bit

grayscale and inverted to obtain a dark background, after which the images were binarized. Binary

objects were dilated to increase dot visibility.

Xenium cell-feature matrices were processed using a custom Seurat script (version 5.3.0) to

generate a combined cluster profile for the blastema and embryonic timepoints. Data were then

normalized and scaled with the SC Transform package (version 0.4.2), followed by principal

component analysis (PCA) using all genes, with PC cutoffs of 11 and 16. The processed matrices

were subsequently analyzed with Find Neighbors and FindClusters, and dimensionality reduction

was performed with t-SNE and UMAP for visualization.

Xenium data were manually processed within the Xenium Explorer 3 (version 3.2.0). Sub-

epithelial regions of the 14-dpa blastema and embryonic tails were isolated through selections that

were restricted to include the blastema/pre-somitic mesoderm and a segment of uninjured/mature

muscle. Transcripts and cells collected in these regions were parsed using an in-house algorithm

developed using Python3 (version 3.10.14). Within the selection area, cells identified by Xenium

Ranger (version 3.1.1) were used to define sample boundaries. Boundary coordinates were

analyzed to orient local anatomical features, including the distal and proximal edges of the

tail. Xenium data are available at GSE313338.

Generation of *Hes7* mutants

Hes7 mutant animals were generated by co-injecting freshly fertilized *d/d* eggs with RNP

complexes containing a cocktail of four different guide RNAs targeting all three exons of *Hes7*

using standard methods[50,51]. gRNA was prepared using primer templates and *in vitro* RNA

synthesis. gRNA efficiency was assessed by targeted amplification of gRNA target sites

compatible with Interference of CRISPR Edits (ICE) analysis. In brief, gDNA was prepared by

incubating 1- 2 mm of tail tissue with 200 μL of 50mM NaOH at 95°C for 15-20 minutes, and

subsequently neutralizing the solution by mixing it with 50 μL of 1M TRIS-in 50 mM NaOH.

Samples were centrifuged at 15,000 RCF for 6 minutes, and the supernatant was subsequently used

for PCR. 1 μL of gDNA was used to seed 50-μL PCR reactions and cleaned up for Sanger

sequencing using magnetic beads. Sanger sequencing was performed using a primer that is internal

to the amplicon. gRNA efficiency was determined through ICE analysis software (Synthego,

Redwood City). The following primers were used

sgRNA generation:

Common_sgRNA_reverse:

5'AAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTTA

ACTTGCTATTTCTAGCTCTAAAAC 3'

Hes7_gRNA1:   5'

GAAATTAATACGACTCACTATAGGGTCGGTCAGCTTCATGGTGTTTTAGAGCTAGAA

ATAGC 3'

Hes7_gRNA2:

5'GAAATTAATACGACTCACTATAGGGCATGAACCACAGCCTGGGTTTTAGAGCTAG

AAATAGC 3'

Hes7_gRNA3:

5'GAAATTAATACGACTCACTATAGGTCAGTGGCTTCCAGAAGCGTTTTAGAGCTAG

AAATAGC 3'

Hes7_gRNA4:

5'GAAATTAATACGACTCACTATAGGGGGGGTTCCGGAATTTCTGGTTTTAGAGCTAGA

AATAGC 3'

Sequencing:

Hes7_gRNA1_Fwd:5' AACTAGCTCAAACCGGCAGA 3'

Hes7_gRNA1_Rev:5' GAGGCCCGAACAGATATTGA 3'

Hes7_gRNA1_Seq:5' CTCCTGCTGGGGAAGTCAT 3'

Hes7_gRNA2+3_Fwd:5' TGATGCTGTTCCCGTGATAA 3'

Hes7_gRNA2+3_Rev:5' AGGAGGTTGGACCTCTTGGT 3'

Hes7_gRNA2+3_Seq:5' CACGTTGCAGGATCAGGAAT 3'

Hes7_gRNA4_Fwd:5' TTTCTCGCCTCACAAGGTCT 3'

Hes7_gRNA4_Rev:5' AAGTTCCGTTGAGGGTGATG 3'

Hes7_gRNA4_Seq:5' GTGGTGAGGCTGATCCATCT 3'

BAPN treatment

3-Aminopropionitril fumarate (Sigma-Aldrich, A3134-25G) was dissolved in tap water to generate a

stock solution of 100 mM, aliquoted and stored at -80 ˚C. Aliquots were further diluted into tap water

to reach working solutions of 10 and 100 µM. Animals were bathed in BAPN solutions and were

refreshed every other day.

Cell ablation using NTR2.0

Transgenic axolotls were generated by crossing *Lfng:GFPnls-T2A-ER^{T2}-Cre-*

ER^{T2} and *CAGGs:loxP-BFPnls-loxP-Cherry-T2a-NTR2.0* lines to enable tamoxifen-inducible

NTR2.0 expression in *Lfng*-positive cells. Animals (~3.5 cm, 3 months old) were housed

in Holtfreter's solution at 19°C under a 12-hour light/dark cycle. NTR2 expression was induced

by treating animals with 2 µM 4-hydroxytamoxifen (4-OHT, Sigma# H6278-50MG) via 12-hour

immersion on days 0, 2, and 4, followed by maintenance in 20% Holtfreter's until day

10, designated as 0 days post-conversion (dpc). At 0 dpc, tail amputations were performed at the

fifth myotome posterior to the cloaca under 0.03% benzocaine anesthesia. Animals were pre-

assigned into three groups: Group 1 (Nifurpirinol+, 4-OHT–, n=5) as non-converted drug-only

controls; Group 2 (Nifurpirinol–, 4-OHT+, n=6) as converted but no-drug controls; and Group 3

(Nifurpirinol+, 4-OHT+, n=7) as the ablation group targeting *Lfng*-lineage cells via NTR2-

mediated cytotoxicity. 10 µM Nifurpirinol (MedChem Express, HY-135470) was delivered by 24-

hour immersion and prepared from a DMSO stock solution. Imaging was conducted under

consistent conditions at 0, 10, 17, 24, 31, 38, and 45 dpc using a Zeiss AxioZoom V16 microscope

at 20x magnification following anesthesia in 0.03% benzocaine.

Histology

Embryos were fixed in Carnoy's solution, and blastema and mature tissues were fixed in 4% PFA.

H&E staining was done using an Epredia Gemini AS Automated Slide Stainer (Fisher Scientific,

Waltham) with standard protocols. Stained slides were imaged using a Panoramic 250 FLASH II

digital scanner (3DHistech Ltd., New Jersey).

Lineage tracing and analysis

To perform Cre-*Lox*P lineage tracing experiments, Cre driver lines were crossed with either

tgSceI(*CAGGs:LoxP-GFP-LoxP-Cherry*)[Etnka] or tgSceI(*CAGGs:LoxP-STOP-LoxP-Cherry*)[Etnka].

Single- and double-transgenic animals were grown up to 4 cm snout-to-tail size. Animals were

treated three times at alternate days with 2 µM 4-OHT by bathing to obtain indelible Cherry

labeling of cells. Labeled animals were amputated 10 days post-conversion (from the first 4-OHT

treatment), near the 6[th] myotome from the cloaca, and contributions of *Cherry*[+] cells to the

regenerate were accessed by stereoscopic imaging of a regenerated tail every week for the next 6

weeks using an Axio Zoom.V16 (Zeiss, Jena) equipped with an Orca-Flash4.0 camera

(Hamamatsu Photonics, Hamamatsu) and an X-Cite Xylis LED illuminator (Excelitas

Technologies, Waltham). At the end of 6 weeks, tails were harvested near cloaca, and IHC or HCR

analysis was performed.

Antibodies

The primary antibodies used in these studies are rabbit anti-RFP (Rockland# 600-401-379, 1:100),

mouse anti-COL1A2 (DSHB # Sp1.D8, 1:100), mouse anti-PCNA (Calbiochem#NA03, 1:500),

mouse anti-PAX7 (DSHB#Pax7, 1:100), mouse anti-MHC (monoclonal antibody 4A1025, a kind

gift from S. Hughes), goat anti-SOX9 (R&D#AF3075, 1:100), rabbit anti-SOX9

(Chemicon#AB5535, 1:1000), rabbit anti-PRRX1[20] (1:200), and rabbit anti-Laminin (Merck,

#L9393 1:200). Secondary antibodies used in these studies were procured from Thermo Fisher

Scientific (Waltham).

Immunohistochemistry and microscopy

IHC staining was done as previously described[20]. Briefly, 10-μm sections were permeabilized

using phosphate-buffered saline (PBS) containing 0.3% Tween-20 and blocked with PBS with

0.3% Triton-X100 and 2% normal hose serum (Vector Laboratories # S-2000). Slides were

incubated O/N with primary antibodies in blocking buffer in a humidified chamber at room

temperature and subsequently incubated with secondary antibodies and Hoechst at a concentration

of 0.5 μg/mL. Mosaic images of sections were acquired on an Axio Imager.Z2 (Zeiss, Jena)

equipped with an Orca-Flash4.0 camera (Hamamatsu Photonics, Hamamatsu) using a 20x/0.8

plan-apochromat objective. Alcian blue / Alizarin red S staining was performed as previously

described[52]. Stained axolotls were photographed using a Nikon D40 DSLR camera with an AF-S

DX NIKKOR 18-55mm f/3.5-5.6G VR II objective (Nikon, Tokyo) and an iPad Air (4th gen)

(Apple, Cupertino) as a backlight illuminator.

Identification of different cell types based on histology

Collagen 1 alpha 2 (COL1A2) is an extracellular matrix (ECM) protein, and in axolotl it is

expressed in dermal fibroblasts and skeleton cells (28). Staining tail sections with COL1A2

antibodies showed broad labeling of the basal lamina (which demarcates dermal fibroblasts and

basal keratinocytes), skeletal and periskeletal structures, as well as fin mesenchyme. Thus, based

on co-staining of PRRX1 and COL1A2 along with laminin and Pax7 of sister sections, we

identified the following seven distinct connective-tissue subtypes (Extended Data Fig. 4): 1)

"Dermal fibroblasts (DF)" - cells underneath the basal lamina that are COL1A2$^+$ and PRRX1med;

2) "Interstitial fibroblasts (IF)" - cells scattered across the fin mesenchyme that are COL1A2$^+$ and

PRRX1^{high} or PRRX1^{med}; 3) "Periskeletal cells (PS)" - cells at the periphery of skeleton that are

COL1A2$^+$ and PRRX1^{high} or PRRX1^{med}; 4) "Skeletal cells (Sk)" - cells that reside inside

cartilaginous structures and are COL1A2$^+$ and PRRX1^{low}; 5) "Notochordal cells (Noto)" - large

vacuole-containing cells within the embryonic notochord; 6) "Muscle progenitors (MP)" - PAX7$^+$

cells scattered between muscle fibers; and 7) "Muscle fibers (MF)" - MHC$^+$ or laminin$^+$ cells

within the myotome region.

Whole-mount light-sheet imaging of PCNA-labeled cells

In brief, axolotl samples were cleared and labeled using the Deep-Clear protocol followed by

imaging on a custom-built light-sheet system[53]. Samples were fixed in 4 % PFA at 4°C overnight.

Specimens were washed several times with PBS at room temperature to remove fixative. After

PBS washes, specimens were incubated in Solution-1 at 37°C for 1 day under gentle shaking. After

the Solution-1 treatment step of tissue clearing and five short PBS washes at room temperature,

samples were treated with 10 % sheep serum at room temperature for several hours. Samples were

then incubated with primary antibodies in 5 % sheep serum at 4°C (gentle shaking) followed by

additional wash steps followed by incubation with secondary antibodies in 5 % sheep serum at 4°C

(gentle shaking). Solution-1 consists of 8% THEED (Sigma-Aldrich, 87600-100ML), 5% Triton®

X 100 (Roth, 3051.2), 25% urea (Roth, X999.2) and 5% CHAPS (Hopax) in dH$_2$O. Solution-2

was prepared by mixing 50% meglumine diatrizoate (Sigma-Aldrich M5266) in PBS (pH 8.5), and

the refractive index was adjusted to 1.45.

Images were acquired using a custom-built light-sheet system equipped with an Olympus Plan

Achromat 1.0/0.25 objective (N1564200). Sample were illuminated with LightHUB 4 system

(Omicron, Germany) containing the following single-mode lasers: 120mW 405nm LuxX laser,

150mW 488nm LuxX laser, 150mW 561nm Cobalt laser, 100mW 594nm Cobalt laser, and

140mW 647nm LuxX laser. Images were recorded in 16-bit quality as Tiff, with a Kinetix camera

(Teledyne Photometrics, USA) 3200x3200px; pixel size 6.5 μm. Images were processed with

AMIRA (Thermo Scientific, USA) rendering software.

Immunogold labeling and electron microscopy

Samples were fixed in freshly prepared 4% PFA. Longitudinal 50 mm thick vibratome sections

were prepared on a Leica VT 1200 vibratome, and sections with target cells were selected based

on their Cherry fluorescence. Vibratome sections were subjected to pre-embedding immunogold

labeling using Nanogold and silver enhancement[54–56]. In brief, samples were blocked and

permeabilized in 20% normal goat serum (NGS) / PBS / 0.1% Saponin for 2 hours at room

temperature followed by incubation in primary antibody (rabbit anti-RFP, Rockland, # 600-401-

379, 1:100) in 20% NGS/0.05% saponin for 2 days at room temperature. After washes in 20%

NGS/0.05% saponin, the samples were incubated with goat-anti-rabbit nanogold (Fab-fragments,

Nanoprobes, 1:50) in 20% NGS/0.05% saponin overnight at room temperature. After final washes

in 20% NGS/0.05% saponin, the samples were postfixed in 1% glutaraldehyde in PBS, washed in

water, and silver-enhanced using the SE-Kit (Aurion, 1 hour incubation time), followed by washes

in water, post-fixation in 1% osmium tetraoxide/water (1 hour on ice), washes in water, *en bloc*

contrasting with 1% uranyl acetate (1h, on ice), washes in water, and dehydration in a graded series

of ethanol (30%, 50%, 70%, 90%, 96% ethanol/water, 3x 100% ethanol on a molecular sieve). The

samples were infiltrated in mixtures of the Epon substitute EMbed 812 with ethanol (1+2, 1+1,

2+1, 2x pure Epon), flat-embedded on the surface of an empty Epon dummy block, and cured

overnight at 65°C. The region of interest (ROI) was identified on semithin sections (1 mm) stained

with 1% toluidine blue/0.5% borax, and the ROI in the block was trimmed for ultrathin sectioning.

70-nm sections were prepared using a diamond knife (Diatome) and the Leica UC6 ultramicrotome

(Leica Microsystems, Wetzlar). The sections were mounted on Formvar-coated slot grids and

contrasted with 4% uranyl acetate/water for TEM. Finally, sections were imaged with a Jeol

JEM1400Plus transmission electron microscope running at 80kV acceleration voltage and

equipped with a Ruby digital camera (Jeol, Tokyo).

HCR-ISH

Samples were fixed in freshly prepared 4% PFA, prepared for cryo-sectioning, and sectioned at

80-μm thickness. HCR staining was performed according to the previously published protocol[57].

The probes were designed using probe generator software from the Monaghan lab

(https://probegenerator.herokuapp.com/ - the lab is no longer functional) or by the MDIBL

Comparative Genomics and Data Science Core team, and probes were ordered from IDT

(Coralville) as oligo pools. All other HCR reagents were ordered from Molecular Instruments (Los

Angeles).

Whole-mount HCR-FISH imaging of axolotl embryos

Stage-30 embryos were fixed in freshly prepared 4% PFA for 1 hour at RT. Embryos were washed

with PBS for 3 × 5 minutes and depigmented using 3% H_2O_2 and 0.8% KOH in DEPC-H_2O for

20 minutes. Next, embryos were incubated in Solution 1 of the Deep-Clear method[53] for 1 hr at

37°C with shaking at 500 rpm on a tabletop shaker. Embryos were washed with 5x SSC-0.1%

Tween-20 (SSCT) for 3 × 15 minutes at 500 rpm. Probe hybridization and amplification were done

according to the manufacturer's protocol (Molecular Instruments, Los Angeles). Embryos were

mounted in Solution-2 of the Deep-Clear protocol and incubated for 1 hr at room temperature to

match the refractive index before imaging. Images were acquired using a Spinning-disk confocal

unit (CSU-W1, 25 μm pinhole diameter, Yokogawa, Japan) on a Nikon inverted Ti-Eclipse

microscope stand (Nikon Instruments Inc., Japan), equipped with a Nikon CFI Plan Apochromat

Lambda D 10X/0.45 for whole-embryo imaging and a Nikon CFI Plan Apochromat Lambda

20X/0.75 lens for the cropped images. Alexa 647, Cherry and DAPI fluorescence were excited

with 640-nm, 561-nm, and 405-nm laser lines and collected using a 405/488/561/640 dichroic

beamsplitter (Yokogawa) with a ET705/72 (Chroma), ET605/52 (Chroma), and ET436/20

(Chroma) emission filter, respectively. Images were acquired at 2048X2048 pixels in 16 bit with

a Scientific CMOS Zyla 4.2 (Andor Technology, United Kingdom) controlled with NIS AR 5.41

(Nikon Instruments Inc., Japan) software and saved in Nd2 or Tiff file format. A maximum

intensity projection (MIP) of the z-stack was created using Fiji software. Images were acquired

using a Nikon Ti-E Yokogawa CSU-W1 / Nikon C2+ confocal microscope with a 10X Nikon CFI

Plan Apochromat Lambda (NA-0.45) lens for whole-embryo imaging and with a 20X Nikon CFI

Plan Apochromat Lambda (NA-0.75) lens for the cropped images. A MIP of the z-stack was

created using Fiji-ImageJ software.

Vertebrae transplantation

To visualize vertebrae, 5-cm *d/d* and *CAGGs:eGFP* animals were soaked in 0.05% Alizarin red S

(ARS) for 30-60 minutes and rinsed several times in fresh tap water[58]. Subsequently, axolotls were

anaesthetized in 0.03% benzocaine (Sigma) and prepared for surgery. Using a SZX16 stereo-

microscope (Olympus, Tokyo) equipped with a Zyla sCMOS camera (Andor, Belfast) and an X-

Cite Xylis LED illuminator (Excelitas Technologies, Waltham, MA) fluorescent microscope to

visualize vertebrae, and fine forceps, 3-4 vertebrae were carefully extirpated from the *d/d* host.

Subsequently, 3-4 vertebrae were dissected from the donor and grafted into the previously

prepared host. Axolotls were left to recover for 2 weeks, after which tails were amputated through

the GFP$^+$ donor vertebrae and lineage-traced for a total of 6 weeks.

Local injuries

Local injuries were performed on 4-7-cm-long *Col1a2:TFPnls-T2a-ER^{T2}-Cre-ER^{T2}* animals

soaked in 0.05% ARS for 30-60 minutes and rinsed several times in fresh tap water[58].

Subsequently, axolotls were anaesthetized in 0.03% benzocaine (Sigma) and prepared for surgery.

Using a SZX16 stereo-microscope (Olympus, Tokyo) equipped with a Zyla sCMOS camera

(Andor, Belfast) and an X-Cite Xylis LED illuminator (Excelitas Technologies, Waltham, MA)

fluorescent microscope to visualize vertebrae, and fine forceps, 3 vertebrae were carefully

extirpated at 10-15 myomeres posterior to the cloaca. To track the regenerative response, animals

were imaged up to 8 weeks after injury using an Axio Zoom.V16 (Zeiss, Jena) equipped with an

Orca-Flash4.0 camera (Hamamatsu Photonics, Hamamatsu) and an X-Cite Xylis LED illuminator

(Excelitas Technologies, Waltham).

Atomic force microscopy

Regenerated axolotl vertebrae were surgically removed on pre-chilled plates. Regenerated tissue

was flash-frozen in OCT on dry ice. OCT blocks were trimmed and cryo-sectioned at -10°C at a

thickness of 20 μm. Sections were collected on electrocharged slides and stored at -80 ˚C. Atomic

force microscopy (AFM) experiments were conducted on a JPK NanoWizard ULTRA SpeedA

(JPK-Bruker) AFM system equipped with an AXIO Observer.D1 (Zeiss) inverted optical

microscope. MLCT-O10 C (Bruker) tipless cantilevers, of 0.009 N/m measured spring constant

via the thermal noise method[59], were furnished with a borosilicate glass microsphere (SPI Supplies

Uniform Microspheres) 15 μm in diameter as previously described[60].

Immediately after removing a glass slide from the freezer, a 10-30 μl droplet of PBS was placed

on the cryosection to prevent it from drying. Subsequently, a fluid cell was mounted and sealed

with a two-component silicone (picodent twinsil, picodent, Germany), allowing all AFM

micromechanical analysis to be conducted on samples hydrated in 750 μL PBS. Cantilever

sensitivity was calibrated via a contact-based method in PBS, as previously described[60,61].

Deflection (volts) vs. z-displacement (μm) curves were recorded on a clean glass next to the

sample, over a 400 nm x 400 nm scan area with 32 px x 32 px resolution. This was repeated two

to three times until the deviation from the previous sensitivity was less than 1%.

Force curves were collected in both the rim and core (features), visible under brightfield

microscopy. On each of the 15 regions of interest (ROIs), 16 force curves were recorded in force

volume map mode within a 2 µm x 2 µm scan area at 4 px x 4 px resolution. Force curves were

recorded at 1.0 nN applied load (relative setpoint) and with a 2-µm z-length. Measurements were

conducted in 6 cryosections at 4wpi without BAPN treatment (2 animals), and 4 cryosections at

4wpi with BAPN treatment (3 animals). Force curves were analyzed with the Hertz model for

spherical indenters using a custom Matlab program available on Github

(https://github.com/Rufman91/ForceMapAnalysis) as previously described ([60]). A Young's

modulus value was obtained for every force curve. Only indentation moduli values resulting

from fits with $R^2 > 0.96$ were considered for further analysis, and an average Young's modulus

was calculated for each ROI.

1002

Buckling instability models

Buckling instabilities with well-defined length scales, and with smooth oscillations generically

occur under compressive load due to a compromise between two different elastic materials. For

the classical setting of a flat sheet of thickness h and Young modulus E_s, under compression and

tethered to an infinitely large elastic half-plane of Young modulus E_c, simple scaling analysis

considering the sheet-bending modulus $E_s h^3$ and the surrounding modulus E_c means that the

wavelength of the instability must scale as $\lambda \propto h \left(\frac{E_s}{E_c} \right)^{\frac{1}{3}}$, with pre-factors depending on the exact

geometry. In the case of a thin shell of thickness h much smaller than its radius of curvature R,

this limit is expected to hold and predict the buckling instability. More specifically, in our setting,

we first only consider peristaltic deformations, i.e. rotationally symmetric, so that the tube

displacement can be parametrized by $u_r(z)$ where r represents the radial direction and z the

directional along the tube. We consider a resting state for a tube composed of a stiff shell of

thickness h and Young's modulus E_s as well as a softer core of radius R and Young's modulus

E_c. For all materials, we will take the Poisson ratio ν to be close to 0.5, i.e. an incompressibility

assumption classical for biological materials. Interestingly, neglecting the mechanics of the core

reflects a fundamental difference between cylindrical and flat geometries. Computing the elastic

energy of the shell from its strain components[62,63], under the classical Foppl-Von Karman

approximation, reveals that whereas compressed flat sheets buckle according to their

fundamental mode (i.e. system size), the constraints of cylindrical geometry select a finite length

scale for buckling, dependent on the aspect ratio of the shell $\lambda = 2\pi R \left(\dfrac{h}{(1-\nu^2)^{\frac{1}{2}} R}\right)^{\frac{1}{2}}$. In the

regime of a very large radius, we go back to the previous result for flat sheets, dependent on the

ratio of stiffness between core and shell $\lambda = 2\pi h \left(\dfrac{E_s}{E_c}\right)^{\frac{1}{3}}$. The relevance of each regime depends

on the relative mechanics of the shell vs the core, i.e. the dimensionless ratio $\dfrac{E_s h}{E_c R}$. As discussed

in the main text, we have computed these numbers via AFM and quantitative geometrical

measurements, allowing us to quantitatively assess them. Interestingly, we find an intermediate

parameter regime, with a ratio close to 1, as the core is approximately 5 times softer than the

shell (E_c/E_s), but also approximately 4-5 times thicker (R/h). We thus compute the wavelength

under each regime, at 4wpi, based on 8 independent measurements of the ratio R/h, taking in

each case the average measured stiffness ratio (E_s/E_c). Under the geometrical regime, we find

$\lambda = 505 \pm 80\ \mu m$, while under the differential stiffness regime, we find $\lambda = 502 \pm 144\ \mu m$,

and thus, interestingly, we are in an intermediate parameter regime where both models make very

similar predictions regarding the wavelength. Importantly, reducing the stiffness or thickness of

the shell is predicted to decrease the buckling wavelength, again in both regimes of the model,

which is in agreement with the results from BAPN treatment. We also remark that in both

regimes of the model, the wavelength of patterning can scale with geometric parameters, i.e. as

long as R and h are larger in larger animals. As such, the wavelength of segmentation can scale

with animal size based on purely mechanical considerations. This is very different from

predictions of reaction-diffusion models, where additional complex regulatory mechanisms must

be invoked to understand how diffusion coefficients or reaction rates change with animal size.

Interestingly, we find signatures of this geometrical scaling in the data, as discussed in the main

text. In particular, although measuring h systematically across different animal or rod sizes is

complicated, as it requires AFM measurements to be precise with respect to the region of higher

shell stiffness, we find strong correlation between wavelength λ and rod radius R in our tail

regeneration model, as predicted in the model. In the local injury model, the number of vertebrae

N should follow from the model $N = L/\lambda$ where L is the length of the injury. As we predict in

our theory that $\lambda \propto \sqrt{R}$, we predict a scaling relationship between N and $L/\sqrt{R}$, something we

observe experimentally, with an increased prefactor in BAPN treatment related to the decrease of

E_s.

Finally, all models above assume axisymmetric deformations of the rod, i.e. that its overall

position remains centered. This also assumes that we can neglect boundary deformations at the

ends of the rod, i.e. that they are fixed. These assumptions are valid if the external mechanical

constraints acting on the rod are strong. *In vivo*, strong constraints arise from the surrounding

mesenchyme. In particular, in the local injury model, the remaining tissues provide strong

constraints both at the ends of the regenerating region, and around it. However, in the tail

amputation experiments, we hypothesized that external confinement might be much weaker, due

to a free boundary condition at the tip of the tail. If we relax the assumption of axisymmetric

deformations, and consider a third elastic medium surrounding the rod, of stiffness E_m, a second

mode of buckling can generically occur, as observed for instance for microtubules under load[62,64]

where the entire rod buckles out of plane. This is exactly what we observe in the $100\mu M$ BAPN

scenario, providing additional confirmation for the hypothesis of pressure-induced buckling of

the rod also operating under full tail amputation. In this case, and simplifying the rod material

properties to the stiff shell, the wavelength can be written as $\lambda = 2\pi\sqrt{hR/2}\left(\frac{E_s}{E_m}\right)^{1/4}$. This

suggests that for a shell stiffness of around 6kPa, one can explain the global buckling length

scale (1-2mm) with realistic stiffnesses of the surrounding mesenchyme of a few hundred

Pascals.

1068

scRNAseq of mature and blastema tissue

Tissue dissociation, and preparation for scRNAseq was performed as previously described[20]. In

brief, tail tissue was dissociated in a dissociation solution (0.1 units/μL DNaseI, 35 μg/mL Liberase

TM in 0.8X PBS) using fine forceps to continuously tear the tissue apart for approximately 30-40

minutes. Samples were passed over a 70-μm Filcon filter (BD# 340605) and spun down at 2,000

rcf for 3 minutes. Cells were carefully resuspended in 0.8X PBS and flow-sorted with a FACSAria

III (BD Biosciences, Franklin Lakes) using an 85-μm nozzle into 200 μL of RLT buffer (Qiagen,

RNA mini kit). Single-cell sequencing was done according to manufacturer's instructions (10x

Genomics, Pleasanton, CA), with v3 chemistry used for all experiments except for library 65766,

where v2 was used (Extended Data Fig. 8d). We used 5' 10x sequencing to capture viral-barcode-

labeled samples with minor modifications[65]. Ten pmol of a GFP-specific reverse primer and a 10x-

specific overhang was spiked into the 5' 10x droplet capture mix. An aliquot of the cDNA was

amplified using a truseq read1 forward primer and a GFP-specific reverse primer with a nextera

read2 overhang. This generated amplicons to perform targeted sequencing and was integrated with

the data from the 5' 10x single-cell transcriptome analysis.

Custom 5' 10x primers used:

5'_10x_GFP_rev_spikein: 5'
AAGCAGTGGTATCAACGCAGAGTACTGGTGCAGATGAACTTCAGGG
TC 3'

GFP_Truseq_GFP_fwd: 5' CTACACGACGCTCTTCCGATCT 3'

GFP_nxt_GFP_rev: 5'
GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGAACAGCTCCTCGCCCTTG 3'
1091
Viral barcode library production, infection, and clone calling

Viral barcode labeling was performed by modifying existing 3rd generation foamy virus vector

systems based on the CellTag principle with modifications[16,18,66]. A 9-nt random barcode sequence

and a defined 6-nt tag were cloned shortly behind the TSS of a puc2MD9-SFFV:GFP-WPRE[66]

transfer vector plasmid using Gibson assembly. In brief, SFFV:GFP-WPRE plasmids were

amplified in two separate fragments using primers with barcode-containing overhangs, and PCR

products were DpnI-treated and purified using para-magnetic beads, and cloned using Gibson

assembly. Immediately after assembly, samples were heat-shock-transformed into stellar

competent cells (Takara Bio, Kusatsu). SOC outgrowth was performed for 1 hour at 37°C, and a

5-μL aliquot was serially diluted and plated on ampicillin agar plates to estimate cloning

efficiencies reaching a minimum complexity of $2*10^6$ colonies. Foamy virus was produced using

Lenti-X 293T (Takara Bio, Kusatsu) generated as previously described and frozen at -80 °C for

long-term storage[66]. Viral titers were determined by infecting HT-1080 cells with serial dilutions

of previously frozen viral preps in multi-well plates. Four days later, plates were imaged using a

Cell Discoverer 7 (Zeiss, Jena), and GFP[+] cells were scored to estimate viral titers. Concentrated

virus was injected into either mature or blastema tissue using glass capillary needles and a PV830

picopump (WPI, Sarasota, FL) using standard protocols[66]. We flow-sorted GFP[+] cells at 21 dpa in

two independent experiments and performed scRNAseq using the 5' 10X platform. Barcoded GFP

transcripts were recovered by adapting existing gRNA capture and direct sequencing approaches[65].

Gibson PCR primers:

CelltagV1_barcode_Fwd: 5'

AGACTGAGTCGCCCGGGTACCGCGGGCCCGNNNNNNNNNNACCGGTGGATCCACCGG

TCGCCACCATG 3'

CelltagV2_barcode_Fwd: 5'

AGACTGAGTCGCCCGGGTACCGCGGGCCCGNNNNNNNNNNCATCACGGATCCACCGG

TCGCCACCATG 3'

CelltagV3_barcode_Fwd: 5'

AGACTGAGTCGCCCGGGTACCGCGGGCCCGNNNNNNNNNNCGTACAGGATCCACCGG

TCGCCACCATG 3'

Celltag_Uni_Rev: 5' TACCCGGGCGACTCAGTCTGT

AmpR_Fwd: 5' AATAAACCAGCCAGCCGGAA

AmpR_Rev: 5' AAGTTGCAGGACCACTTCTG

1124

Clonal analysis of viral barcode labeling

Non-paraxial mesodermal lineages were excluded from the analysis. Barcodes were extracted and

filtered (present in ≥ 2 cells and ≥ 2 UMIs per cell). Clones were called using a Jaccard similarity

index of ≥ 0.45. The axolotl has a measured cell cycle ranging from 53-103 hours[9,67]. Assuming an

average cell cycle time of 100 hours, tails infected with V1 4 days prior to amputation would have

undergone 5.04 rounds of cell division, tails infected with V2 4 days after amputation would have

undergone 4.08 rounds of division, and tails infected V3 7 days after amputation would have

undergone 3.36 rounds of cell division. This corresponds to maximum clone sizes of 32.89 (V1),

16.91 (V2), and 10.27 (V3). Reducing the Jaccard similarity index threshold results in unusually

large clones being called, which is inconsistent with the known cell-division rate during tail

regeneration.

1136

Called clones are displayed in a matrix to visualize clonal diversity (Fig. 3f). Here each row

represents a specific cell type, and each column represents a clone. To ensure that clones were not

promiscuously called, we computationally combined the two independent replicates, and in this

mixed population called clones using a Jaccard similarity index of ≥0.45. We found that 47/48

clones were exclusively made up of cells originating from either replicate 1 or replicate 2, while

only a single hybrid clone called was made up of 2 cells from replicate 1 and 1 cell from replicate

2 (Extended Data Fig. 4i). This suggests that with a Jaccard similarity index of ≥0.45 we can

reliably call clones in these data-sets and achieve a false discovery rate of 3.6%.

1145

1146 scRNAseq of embryonic tissue

1147 Axolotl embryos were staged to stage 25-35[68] and tail buds were dissected in 0.8X PBS under a

1148 SZX10 stereo-microscope (Olympus, Tokyo). Individual cells were manually collected using P20

1149 pipettes and prepared for Smart-seq2 using standard protocols in a 96-well format[69].

1150

Processing and analysis of scRNAseq data

The quality of the cDNA and resulting sequencing libraries were checked by Bioanalyzer (High

Sensitivity DNA Kit, Agilent, 5067-4626). The libraries were sequenced using an Illumina

NovaSeq PE150. Sequencing reads were mapped against the axolotl genome[70,71] using STARsolo

(STAR 2.7.5a)[72], with parameters mimicking Cell Ranger's (10x Genomics) transcript-counting

strategy. Fluorescent marker genes were manually added to the references. Data analysis was

performed primarily via Seurat v3.1[73]. Ribosomal protein genes and pseudogenes were excluded

from downstream analyses. Generally, cells with a very high or a very? low number of transcripts

counts, as well as those with a high mitochondrial transcript proportion, were excluded (see exact

information in Table 2). When clusters were identified to be still driven only by low RNA counts

in a first screening, cells belonging to these clusters were removed. To enrich connective-tissue

cells, some contaminating cell types such as epidermal cells were identified and removed after an

initial screening using Seurat. Seurat's built-in scale function was used to regress out differences

in nUMI and percent.mt. Scaling and PCA were always performed on all genes. UMAP was

applied to the top principal components (PCs) for visualization. Data integration was performed

for the blastema and the uninjured data sets using Seurat's built-in Harmony[74] function to reduce

the confounding effect of different batches and samples. We performed Harmony using default

parameters on the library level, 100 input PCs for blastema integration, and 50 input PCs for

uninjured integration. The integration of the uninjured and blastema data sets was performed using

Harmony on the library level with slightly adapted parameters (PC=50, theta=2, lambda=9).

1171

Trajectory analysis

To infer directionality of developmental trajectories during regeneration, we used scVelo[75]. To

infer the ratio of spliced and unspliced reads using scVelo, the sequencing data were mapped with

STARsolo and its built-in 'Velocyto' mode[72] to obtain spliced and unspliced count matrices,

respectively. Afterwards, scVelo was used following the standard procedure

(https://scvelo.readthedocs.io/en/stable/VelocityBasics) without dynamical mapping. Fifty input

PCs were used for the moments calculation to obtain the arrow velocity graph. Velocity diffusion

time was calculated and used as a prior for the identification of links and directionality between

the clusters by PAGA.[76] The diffusion pseudotime presented in Figure 4h was calculated using the

standard diffusion map algorithm[77] in order to set the DMS cells manually to 0 and demarcate

them as the starting population.

1183

1184 Statistical analysis

1185 Statistical analysis was performed using a combination of PRISM (version 9.0.0), and R where

1186 appropriate. Student's t-test and ANOVA were performed using PRISM. MZ structural break test

1187 was performed in R.