Calcium-driven In Silico Inactivation of a Human Olfactory Receptor

Lorenza Pirona,[†,¶] Federico Ballabio,[†,¶] Mercedes Alfonso-Prieto,[‡] and Riccardo Capelli[*,†]

†*Department of Biosciences, Università degli Studi di Milano, Via Celoria 26, I-20133 Milano, Italy*

‡*Computational Biomedicine, Institute for Advanced Simulations IAS-5/Institute for Neuroscience and Medicine INM-9, Forschungszentrum Jülich GmbH, Wilhelm-Johnen-Straße, D-54248 Jülich, Germany*

¶*Equal contribution*

E-mail: riccardo.capelli@unimi.it

Abstract

Conformational changes as well as molecular determinants related to the activation and inactivation of olfactory receptors are still poorly understood due to the intrinsic difficulties in the structural determination of this GPCR family. Here, we perform, for the first time, the *in silico* inactivation of the human olfactory receptor 51E2, highlighting the role of calcium in this receptor state transition. Using molecular dynamics simulations, we show that the presence of a divalent ion in the ion binding site stabilizes the receptor in its inactive state. Our findings suggest a novel molecular mechanism for OR inactivation, potentially guiding experimental validation and offering insights into the broader role of divalent ions in GPCR signaling.

Olfactory receptors (ORs) are a family of G protein-coupled receptors (GPCRs) implicated in odor perception.[1] This group of class A GPCRs comprises approximately 400 members, representing half of the GPCRs encoded in the human genome.[2] The function of ORs is not limited to olfaction, as they have been identified in various tissues and have been shown to play a role in a plethora of physiological and pathological processes.[3,4] This important clinical perspective, combined with the fact that non-olfactory GPCRs are the target of 34% of the total number of FDA-approved drugs,[5] makes the study of the OR family one of the most promising fields for biochemistry and pharmacology.

Despite their potential as drug targets,[6] the study of ORs has been hindered by the lack of structural data: the first experimental structure was published in March 2023,[7] consisting of human OR51E2 in its active state, bound to an agonist (propionate) and a mini-G protein. In light of this limitation, the modeling community has approached the OR family with different techniques, from homology modeling[8–11] to *de novo* structure prediction.[12,13] Such computational models have provided useful insights into the mechanism of OR-ligand recognition at the atomistic level, as well as pinpointed crucial residues for OR function, in combination with experimental mutagenesis data.

Recent cryoEM work by Choi *et al.*[14] unveiled, for the first time, both the active and inactive conformations of a chimeric OR, OR52$_{cs}$, which was constructed using a consensus sequence strategy based on 26 members of the OR52 family. The availability of experimental structural data for both conformational states of the same OR receptor opens the opportunity for cross-validating *in silico* models. Furthermore, this enables a computational exploration of the activation or deactivation process, aligning with the approach taken in similar computational studies on other class A GPCRs.[15,16]

In this study, we focus on the deactivation of human OR51E2, leveraging the cryoEM structure of this receptor in its active state.[7] Our primary objective is to simulate the inactivation of the receptor *in silico* over a μs timescale, drawing upon the pioneering work of Dror *et al.* on the β_2-adrenergic receptor.[15] Moreover, OR51E2, along with 90% of the

OR family members, bears two acidic residues in the ion binding site[13] (2.50 and 3.39 using the Ballesteros-Weinstein class A GPCR generic numbering[17]), indicating the potential involvement of charged residues in the ion binding mechanism and conformational dynamics of ORs. This hypothesis is supported by our previous observations of sodium-bound OR51E2 in the inactive state.[13] Accordingly, we aim to elucidate the role of the ion binding site for OR conformational plasticity, building on previous work on sodium binding to other class A GPCRs.[18,19]

Simulations with no ions in the binding site. Initially, the receptor structure was built from its experimentally determined active conformation (PDB code: 8F76[7]), with the agonist and mini-G protein removed. We employed the CHARMM-GUI web server[20,21] for receptor set up and its embedding into a 3:1 POPC:Cholesterol bilayer (see Methods section and Supporting Information for details). Following system preparation, we adopted a multi-step equilibration protocol, applying restraints to preserve the initial fold while relaxing the system (refer to Methods), as done in our previous work.[13] This was followed by five independent unrestrained MD simulations, each lasting 5 μs.

In all five replicas, a partial loss of the receptor fold was consistently observed (see Figure 1a), which is in line with the results obtained when starting from *de novo* structures of OR51E2.[13] The region most impacted by this partial unfolding was the interface between transmembrane helix 6 (TM6) and transmembrane helix 7 (TM7), which widens significantly during the simulation (from 8 Å to 15 Å).

Additionally, the expansion of the TM6-TM7 interface coincides with POPC molecules "diving" into the ion binding pocket, as their positively charged choline groups orient towards the charged residues D69$^{2.50}$ and E110$^{3.39}$ (see Figure 1b). These events clearly indicate a significant charge imbalance near the ion binding pocket, leading to lipid snorkeling through the apolar membrane. Within this structural rearrangement, we observed several phenomena related to the system conformational dynamics: (i) Water molecules freely traversed from the intracellular to the extracellular side of the membrane; (ii) The toggle switch Y251$^{6.48}$

Figure 1: OR51E2 from simulations without ions in the binding pocket. (a) Superposition of the cluster centroid obtained from the five trajectories (blue) started from the cryoEM structure in its active state (white), evidencing the widening of the TM6-TM7 interface. (b) Detail of the ion binding pocket, highlighting the interaction between the charged residues $D69^{2.50}$ and $E110^{3.39}$ and the two POPC headgroups entering the receptor bundle. Additionally, $Y251^{6.48}$ is oriented towards the ion binding pocket. Hydrogen atoms have been omitted for clarity, and the distances refer to the frame shown.

established multiple transient interactions with the charged heads of POPC lipids (both choline and phosphate moieties), $E110^{3.39}$, and $S111^{3.40}$; and (iii) The negatively charged side chains of $D69^{2.50}$ and $E110^{3.39}$, being locally unbalanced, interacted with nearby polar and neutral amino acid side chains, and also attracted the charged groups that entered the

ion binding site (see Table S2 in the Supporting Information). Remarkably, we observed spontaneous binding of a Na^+ ion in two simulations (refer to Table S2 and Figure S1). The ion entered from either the extra- (replica 4) or intracellular (replica 1) side of the membrane, but it is important to note that due to the use of periodic boundary conditions (PBC), the ionic concentration is nearly identical on both sides of the membrane.

As an independent confirmation of our observation of an electrostatic interaction between the ion binding pocket and charged groups, we considered the final frame of 1 μs simulation started from the apo structure of $OR52_{cs}$ (retrieved at https://github.com/sek24/natcomm2023); such simulation was performed under similar conditions to ours (except for the use of KCl instead of NaCl). Also in the case of $OR52_{cs}$, two POPC molecules snorkel into the ion binding pocket. However, in our simulations the two lipid molecules wedge themselves between TM6 and TM7, whilst in the simulations of Choi *et al.* they do between TM5-TM6. In addition, the final $OR52_{cs}$ frame shows a positively charged ion (K^+) located close to $D69^{2.50}$ and $E110^{3.39}$, similar to the spontaneous sodium binding event observed in some of our replica simulations.

Simulations with sodium in the ion binding site. After observing the spontaneous binding of Na^+ in the ion binding site, we carried out a new set of five replica simulations. In these simulations, Na^+ was manually positioned in the ion binding pocket of the initial structure, while retaining the same procedure and parameters as in the previous section. The simulation length was 5 μs per replica. This approach aligns with the observations from our previous work on the *de novo* structures of the same receptor in the inactive state,[13] which maintained their fold only when sodium was present in the ion binding pocket. However, here we began with the cryoEM structure of OR51E2 in the active state[7] and conducted additional and longer replicas.

Overall, the simulations involving Na^+ showed slightly enhanced receptor stability, which is reflected in less POPC snorkeling into the ion binding pocket (Figure S4), reduced water permeability (Table S2), and lower RMSD values (Figure S3) of the C_α atoms of the trans-

Figure 2: OR51E2 from simulations with Na^+ ion in the binding pocket. (a) Superposition of the cluster centroid obtained from the five trajectories (blue) on the cryoEM structure in its active state (white). (b) Detail of the ion binding pocket, highlighting the interactions between Na^+ and $D69^{2.50}$ and $E110^{3.39}$, as well as between POPC and the same two acidic residues. Hydrogen atoms have been omitted for clarity, and the distances refer to the frame shown.

membrane domains with respect to the cryoEM structure. Differently from the simulations with no ions, two trajectories (replicas 1 and 3) did not reveal any POPC snorkeling. In the other three trajectories, we observed the POPC head group moving towards the ion binding site, passing either through the interface between TM5-TM6 and interacting mainly with $E110^{3.39}$ (replica 5 and the last frames of replica 2) or between TM6-TM7, engaging

residues D69$^{2.50}$ and E110$^{3.39}$ (replicas 4 and 5). Interestingly, Na$^+$ unbinding was detected in replica 5 in the last μs, where the ion moved to the intracellular space. Shortly after this event, a second POPC entered the ion binding site. Finally, a new Na$^+$ ion spontaneously translocated from the extracellular space to the ion binding site within 1 μs from the initial dissociation event (Figure S3 and Table S2). This phenomenon supports, once again, the need for a charged group at this location, counteracting the negative charge of the two acidic residues at positions 2.50 and 3.39.

In terms of water permeability, we observed reduced water passing through the protein compared to the simulations with no ions (Table S2). This trend was evident in the first three replicas, particularly in the second trajectory. Considering that the formation of a hydrated pathway spanning the receptor is connected to the reorganization of the H-bonds around the ion binding site,[19] we focused on replica 2 to identify changes in non-bonded interactions at the atomistic level. Namely, we performed a comparative H-bond analysis across all the ten replicas described thus far (five with Na$^+$ and five without) to identify pairs that, in terms of occupancy, exhibited the most significant differences between the second replica and the others. The interaction between D69$^{2.50}$ and Y291$^{7.53}$ was particularly notable. The total occupancy of this H-bond (i.e. considering both oxygen atoms of the side chain of D69$^{2.50}$) in the second replica of the Na$^+$-bound system persisted for 76% of the trajectory. In contrast, replicas 1, 3, 4, and 5, showed occupancies of 19%, 52%, 3%, and 4%, respectively. Extending this analysis to the replicas without the ion in the binding site, this occupancy reached a maximum of 6% (replica 4), while in all the others it was around 1%. Y291$^{7.53}$ has been reported as a conserved residue in class A GPCRs, implicated in the diffusion of Na$^+$ ions,[18,19] the formation of a water pathway inside the receptor, and in the receptor activation mechanism.[22] Another interesting H-bond pattern was observed between S111$^{3.40}$ and Y251$^{6.48}$, which is significantly more frequent in the presence of sodium. This increased persistence is particularly evident in three out of five trajectories with sodium (occupancy of 41%, 78%, 92%, 78%, 4% in replica 1,2,3,4,5 respectively). In contrast, only one replica from

the trajectories lacking ions exhibits a stable H-bond between these two residues (occupancy of 90% in replica 1, 0% in all the others).

To assess the stability of the receptor, we performed an RMSD calculation using the cryoEM structure of OR51E2 in its active state[7] as the reference . Overall, the simulations with Na^+ bound maintained the fold better (i.e. lower RMSD values - see Figure S3) than the simulations without ions (Figure S1), with some minor variability among them. Interestingly, peaks in the RMSD time evolution are linked to lipid snorkeling and increased water permeability in the receptor (Figure S4).

Simulations with calcium in the ion binding site. From the previous simulations we can conclude that the charge imbalance in the ion binding site is still present, even if attenuated, when Na^+ is bound. Hence, we decided to perform another round of five independent simulations, positioning a Ca^{2+} ion in the binding site while maintaining all the simulation parameters used previously (see Methods and Supporting Information). Based on sequence analyses, class A GPCRs containing two acidic residues in the ion binding pocket have been hypothesized to bind Ca^{2+};[23] however, to the best of our knowledge, this possibility has not yet been investigated in the context of ORs.

The simulations with Ca^{2+} bound further support the stabilization of the receptor fold in the presence of an ion in the binding site, as observed in some of the Na^+-bound simulations. In addition to the lower RMSD values (Figure S5), comparison of the RMSF of the three simulation datasets (Figure S7) further confirms increased stability and reduced flexibility of the receptor with calcium bound, particularly in the TM5-TM6-TM7 region. Analysis of the H-bonds, across the five replicas, did not reveal any exclusive interactions that are absent in the trajectories with sodium. Furthermore, in contrast to the simulations with Na^+, $Y291^{7.53}$ is able to form a hydrogen bond with $D69^{2.50}$ in replicas 2 and 4, but only for about 25% of the trajectory. In replicas 1 and 5, $Y291^{7.53}$ establishes stable hydrophobic interactions with $V44^{1.53}$, $L62^{2.43}$ and $I298^{8.50}$. Generally, in the simulations with Ca^{2+}, we observed that the side chain of $Y291^{7.53}$ preferentially orients towards TM1 and TM2. Moreover, the improved

Figure 3: Results of the *in silico* Ca^{2+}-bound OR51E2 simulations compared to the available experimental structures. (a) Superposition of the active (orange) and apo (green) structures of OR52$_{cs}$ solved by Choi *et al.*[14] (b) Comparison of the OR51E2 experimental active structure[7] (white) and the cluster centroid obtained from the five trajectories with Ca^{2+} in the ion binding site (blue). (c) Detail of the ion binding pocket, highlighting the interactions between Ca^{2+} and the two acidic residues, D69$^{2.50}$ and E110$^{3.39}$. Hydrogen atoms have been omitted for clarity, and the distances refer to the frame shown. (d) Histogram of the RMSD of all the simulations performed in this work (without ions, with Na^+, and with Ca^{2+} in the ion binding site) with respect to TM3-TM5-TM6 C_α of the apo form of OR52$_{cs}$.

electrostatic complementarity between the +2 charge of calcium and the -2 charge of the ion binding site appears to preclude lipid snorkeling, as we did not observe any POPC head groups reaching the ion binding site over the total accumulated 25 μs time. In particular, we noticed a tight coordination of the acidic residues D69$^{2.50}$ and E110$^{3.39}$ with the Ca^{2+} ion (Figure 3c). In this context, visual inspection during the simulation reveals a reduced (almost negligible) amount of water passing between the two sides of the membrane through the receptor (see Table S2).

The main global dynamic effect observed during the Ca^{2+}-bound simulations is characterized by the rotational movement of TM5-TM6 around TM3 acting as a pivot (Figure 3b), which happens during the first $0.3 \div 0.7$ μs for all the replicas. Such conformational change appears in agreement with the only experimentally observed active/inactive transition of an OR to date (Figure 3a) and could be considered as the hallmark of inactivation for this family of receptors. Considering this last point, we performed the calculation of RMSD for all the 75 μs of simulations, taking as reference the C_α atoms of TM3, TM5, and TM6 of the $OR52_{cs}$ structure, as labeled with the structure alignment tool in the GPCRdb.[24] We can observe that the RMSD distributions (Figure 3d) are closer to the $OR52_{cs}$ inactive state in the presence of higher-charge ions in the binding site (No ion > Na^+ > Ca^{2+}). Based on this analysis, we can consider these simulations as the first observation of calcium-driven *in silico* inactivation of a human OR.

The conservation of two acidic residues in the ion binding site of $\sim 98\%$ of human ORs,[13] at positions 2.50 and 3.39 (Figure 4a), raises the question of whether not only monovalent, but also divalent ions, can occupy this site and modulate receptor function. The simulations carried out in this work show that indeed calcium is more efficient than sodium at stabilizing the inactive state of a prototypical OR, OR51E2.

The possibility of divalent ions binding to class A GPCRs containing two acidic residues in the ion binding site was already highlighted in the review by Katritch *et al.* as an "outstanding question".[23] Besides the conserved $D^{2.50}$, previous sequence analyses[23,25] have shown that a small subset of non-olfactory class A GPCRs ($\sim 21\%$) bear a second acidic residue in the ion binding site, typically at position 7.49 ($\sim 18\%$) and occasionally at positions 6.44 ($\sim 2\%$) and 3.39 ($\sim 1\%$) (see Figure 4b and Table S3). However, as of January 2024 only the experimental structures of melanocortin receptors[26–32] have shown Ca^{2+} bound to two aspartates at positions 2.50 and 7.49, acting as a cofactor for agonist binding. Instead, experimental structures of proteinase-activated receptors[33,34] and cysteinyl leukotriene receptors,[35] also bearing two aspartates at positions 2.50 and 7.49, have been solved with Na^+

and an antagonist bound.

Figure 4: Sequence logos depicting conservation of residues lining the ion binding site in class A GPCRs. (a) Human olfactory receptors, based on the multiple sequence alignment of 412 sequences in Fierro *et al.*[36] (b) Human non-olfactory class A GPCRs, using the 286 sequences available in the GPCRdb sequence alignment tool[37] as of January 2024. Sequence logos were generated with WebLogo version 3.7.12[38] and amino acids are colored according to their chemical properties.

We thus believe that the importance of calcium ions has been unexplored in the context of GPCRs,[39] in particular for ORs. Despite the great advancements in the experimental determination of GPCR structures, the state of the art in cryo-EM is still unable to discriminate between water and ions.[40] A possible help may come from the use of buffers with high(er) calcium concentrations, focusing our attention on possible ion-protein interactions. Moreover, while most non-olfactory class A GPCRs with a second acidic residue (see Table S3) bear an aspartate at either position 7.49 ($\sim 86\%$) or 6.44 ($\sim 8\%$), the majority of ORs ($\sim 90\%$) have glutamate at position 3.39.[13] This is expected to shift the ion location within the binding site, thus resulting in different effects of ion binding on receptor function. Interestingly, two chemokine receptors (CCR1 and CCR3) and one peptide receptor (QRFP) also display a glutamate at position 3.39, suggesting that calcium might also stabilize the inactive state of these receptors, similarly to what we proposed here for ORs.

In summary, in this work we suggest a molecular mechanism for ORs inactivation, which relies on the presence of calcium in the ion binding site. In this regard, calcium-mediated OR inactivation could constitute a negative feedback mechanism to stop the olfactory signal triggered by odorant binding and subsequent calcium entry through cyclic nucleotide gated channels. We hope that this computationally-driven hypotheses could be validated by means of experimental techniques, highlighting the predictive power of molecular modeling and simulations to suggest biologically relevant structure-function relationships. Such experimental testing could include *in vitro* assays with varying concentrations of calcium in the extracellular medium or structural determination efforts with higher calcium (or divalent metal analogs) concentrations in the buffers used for X-ray, cryo-EM or native mass spectrometry.[41]

Methods

System preparation

The initial configuration of OR51E2 in its active state was obtained from the Protein Data Bank (PDB code: 8F76[7]). This structure was preprocessed using the Protein Preparation Wizard implemented in Schrödinger Maestro 2023-3,[42] which automatically assigns the amino acid protonation states. In particular, both D69$^{2.50}$ and E110$^{3.39}$ were modeled as negatively charged. The complete system was then assembled using the CHARMM-GUI.[20,21] First, a disulfide bond was established between C96$^{3.25}$ and C178$^{45.50}$; then, a cubic box of dimensions $100 \times 100 \times 120\text{Å}^3$ was defined with the receptor embedded in a mixed lipid bilayer composed of POPC and cholesterol (3:1 ratio). The receptor-membrane system was then solvated in water with a NaCl concentration of 0.15 M. In the case of the calcium-bound simulations, we added a single Ca^{2+} ion and neutralized its charge by adding two additional Cl$^-$ ions in the solution. The protein, lipids, and ions were parameterized using the CHARMM36m force field,[43] while water was described using the TIP3P[44] model.

Molecular dynamics simulations

In this work, the equilibration phase of the simulations was conducted following a protocol presented in our previous work;[13] further details are provided in the Supporting Information. For the subsequent 5 μs production phase, we performed unrestrained molecular dynamics (MD) simulations, with a 2 fs time step. To maintain constant temperature and pressure conditions at 310 K and 1 bar, respectively, we employed the velocity rescaling thermostat[45] alongside the semi-isotropic cell rescaling barostat.[46] A total of 15 independent runs were executed, comprising five replicates for each of the three models, i.e. without any ions in the ion binding pocket, with Na^+ bound, and with Ca^{2+} bound. Each of these runs was initiated with distinct initial velocities. The simulations were all carried out using GROMACS,[47] version 2021.2.

Data Availability Statement

Data needed to reproduce the results shown in this paper (structures, topology, GROMACS input files, etc.) and the resulting trajectories are available at Zenodo (https://dx.doi.org/10.5281/zenodo.10589510). All the trajectories computed here for OR51E2 have been also uploaded on the GPCRmd,[48] with the accession codes 1976 (no ions), 1977 (Na^+-bound), and 1978 (Ca^{2+}-bound).

Acknowledgement

We thank Prof. Carlo Camilloni and Dr. Antonella Paladino for fruitful discussion and Prof. Hee-Jung Choi for providing us, before its official deposit, the apo structure of OR52$_{cs}$. We acknowledge EuroHPC Joint Undertaking for awarding access to the Karolina CPU cluster at IT4Innovations, Czech Republic (EHPC-REG-2022R03-049). This research was supported by Università degli Studi di Milano (Piano di supporto di ateneo Linea 2-DBS Capelli) to

Supporting Information Available

The supporting information file contains the extended details of the MD simulations, a table with the Ballesteros-Weinstein numbering of OR51E2, a table with the main events observed in every MD replica, RMSD of all the replicas with respect to experimental active state OR51E2, minimum distance analysis between POPC and the ion binding site, and the RMSF for all the simulations performed.

References

(1) Buck, L.; Axel, R. A novel multigene family may encode odorant receptors: A molecular basis for odor recognition. *Cell* **1991**, *65*, 175–187.

(2) Alexander, S. P. H.; Christopoulos, A.; Davenport, A. P.; Kelly, E.; Mathie, A.; Peters, J. A.; Veale, E. L.; Armstrong, J. F.; Faccenda, E.; Harding, S. D.; Pawson, A. J.; Sharman, J. L.; Southan, C.; Davies, J. A. The concise guide to Pharmacology 2019/20: G protein-coupled receptors. *British Journal of Pharmacology* **2019**, *176*.

(3) Maßberg, D.; Hatt, H. Human olfactory receptors: novel cellular functions outside of the nose. *Physiological Reviews* **2018**, *98*, 1739–1763.

(4) Lee, S.-J.; Depoortere, I.; Hatt, H. Therapeutic potential of ectopic olfactory and taste receptors. *Nature Reviews Drug Discovery* **2018**, *18*, 116–138.

(5) Hauser, A. S.; Attwood, M. M.; Rask-Andersen, M.; Schiöth, H. B.; Gloriam, D. E. Trends in GPCR drug discovery: new agents, targets and indications. *Nature Reviews Drug Discovery* **2017**, *16*, 829–842.

(6) Naressi, R. G.; Schechtman, D.; Malnic, B. Odorant receptors as potential drug targets. *Trends in Pharmacological Sciences* **2023**, *44*, 11–14.

(7) Billesbølle, C. B.; de March, C. A.; van der Velden, W. J. C.; Ma, N.; Tewari, J.; del Torrent, C. L.; Li, L.; Faust, B.; Vaidehi, N.; Matsunami, H.; Manglik, A. Structural basis of odorant recognition by a human odorant receptor. *Nature* **2023**, *615*, 742–749.

(8) de March, C. A.; Yu, Y.; Ni, M. J.; Adipietro, K. A.; Matsunami, H.; Ma, M.; Golebiowski, J. Conserved residues control activation of mammalian G protein-coupled odorant receptors. *Journal of the American Chemical Society* **2015**, *137*, 8611–8616.

(9) Wolf, S.; Jovancevic, N.; Gelis, L.; Pietsch, S.; Hatt, H.; Gerwert, K. Dynamical binding modes determine agonistic and antagonistic ligand effects in the prostate-specific G-protein coupled receptor (PSGR). *Scientific Reports* **2017**, *7*, 16007.

(10) Yu, Y.; Ma, Z.; Pacalon, J.; Xu, L.; Li, W.; Belloir, C.; Topin, J.; Briand, L.; Golebiowski, J.; Cong, X. Extracellular loop 2 of G protein–coupled olfactory receptors is critical for odorant recognition. *Journal of Biological Chemistry* **2022**, *298*, 102331.

(11) Shim, T.; Pacalon, J.; Kim, W.-C.; Cong, X.; Topin, J.; Golebiowski, J.; Moon, C. The Third Extracellular Loop of Mammalian Odorant Receptors Is Involved in Ligand Binding. *International Journal of Molecular Sciences* **2022**, *23*, 12501.

(12) Nicoli, A.; Haag, F.; Marcinek, P.; He, R.; Kreißl, J.; Stein, J.; Marchetto, A.; Dunkel, A.; Hofmann, T.; Krautwurst, D.; Di Pizio, A. Modeling the orthosteric binding site of the G protein-coupled odorant receptor OR5K1. *Journal of Chemical Information and Modeling* **2023**, *63*, 2014–2029.

(13) Alfonso-Prieto, M.; Capelli, R. Machine Learning-Based Modeling of Olfactory Receptors in Their Inactive State: Human OR51E2 as a Case Study. *Journal of Chemical Information and Modeling* **2023**, *63*, 2911–2917.

(14) Choi, C.; Bae, J.; Kim, S.; Lee, S.; Kang, H.; Kim, J.; Bang, I.; Kim, K.; Huh, W.-K.; Seok, C.; Park, H.; Im, W.; Choi, H.-J. Understanding the molecular mechanisms of odorant binding and activation of the human OR52 family. *Nature Communications* **2023**, *14*.

(15) Dror, R. O.; Arlow, D. H.; Maragakis, P.; Mildorf, T. J.; Pan, A. C.; Xu, H.; Borhani, D. W.; Shaw, D. E. Activation mechanism of the β_2-adrenergic receptor. *Proceedings of the National Academy of Sciences* **2011**, *108*, 18684–18689.

(16) Dong, S. S.; Goddard, W. A.; Abrol, R. Conformational and Thermodynamic Landscape of GPCR Activation from Theory and Computation. *Biophysical Journal* **2016**, *110*, 2618–2629.

(17) Ballesteros, J. A.; Weinstein, H. *Methods in neurosciences*; Elsevier, 1995; Vol. 25; pp 366–428.

(18) Miao, Y.; Caliman, A. D.; McCammon, J. A. Allosteric effects of sodium ion binding on activation of the M3 muscarinic G-protein-coupled receptor. *Biophys. J.* **2015**, *108*, 1796–1806.

(19) Vickery, O. N.; Carvalheda, C. A.; Zaidi, S. A.; Pisliakov, A. V.; Katritch, V.; Zachariae, U. Intracellular Transfer of Na+ in an Active-State G-Protein-Coupled Receptor. *Structure* **2018**, *26*, 171–180.e2.

(20) Jo, S.; Kim, T.; Iyer, V. G.; Im, W. CHARMM-GUI: a web-based graphical user interface for CHARMM. *Journal of Computational Chemistry* **2008**, *29*, 1859–1865.

(21) Lee, J. et al. CHARMM-GUI input generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM simulations using the CHARMM36 additive force field. *Journal of Chemical Theory and Computation* **2016**, *12*, 405–413.

(22) Yuan, S.; Filipek, S.; Palczewski, K.; Vogel, H. Activation of G-protein-coupled receptors correlates with the formation of a continuous internal water pathway. *Nature Communications* **2014**, *5*, 4733.

(23) Katritch, V.; Fenalti, G.; Abola, E. E.; Roth, B. L.; Cherezov, V.; Stevens, R. C. Allosteric sodium in class A GPCR signaling. *Trends in Biochemical Sciences* **2014**, *39*, 233–244.

(24) Pándy-Szekeres, G.; Caroli, J.; Mamyrbekov, A.; Kermani, A. A.; Keserű, G. M.; Kooistra, A. J.; Gloriam, D. E. GPCRdb in 2023: state-specific structure models using AlphaFold2 and new ligand resources. *Nucleic Acids Research* **2022**, *51*, D395–D402.

(25) Zarzycka, B.; Zaidi, S. A.; Roth, B. L.; Katritch, V. Harnessing ion-binding sites for GPCR pharmacology. *Pharmacological Reviews* **2019**, *71*, 571–595.

(26) Yu, J. et al. Determination of the melanocortin-4 receptor structure identifies Ca^{2+} as a cofactor for ligand binding. *Science* **2020**, *368*, 428–433.

(27) Heyder, N. A. et al. Structures of active melanocortin-4 receptor–Gs-protein complexes with NDP-α-MSH and setmelanotide. *Cell Research* **2021**, *31*, 1176–1189.

(28) Israeli, H.; Degtjarik, O.; Fierro, F.; Chunilal, V.; Gill, A. K.; Roth, N. J.; Botta, J.; Prabahar, V.; Peleg, Y.; Chan, L. F.; Ben-Zvi, D.; McCormick, P. J.; Niv, M. Y.; Shalev-Benami, M. Structure reveals the activation mechanism of the MC4 receptor to initiate satiation signaling. *Science* **2021**, *372*, 808–814.

(29) Ma, S.; Chen, Y.; Dai, A.; Yin, W.; Guo, J.; Yang, D.; Zhou, F.; Jiang, Y.; Wang, M.-W.; Xu, H. E. Structural mechanism of calcium-mediated hormone recognition and Gβ interaction by the human melanocortin-1 receptor. *Cell Research* **2021**, *31*, 1061–1071.

(30) Zhang, H. et al. Structural insights into ligand recognition and activation of the melanocortin-4 receptor. *Cell Research* **2021**, *31*, 1163–1175.

(31) Feng, W.; Zhou, Q.; Chen, X.; Dai, A.; Cai, X.; Liu, X.; Zhao, F.; Chen, Y.; Ye, C.; Xu, Y.; Cong, Z.; Li, H.; Lin, S.; Yang, D.; Wang, M.-W. Structural insights into ligand recognition and subtype selectivity of the human melanocortin-3 and melanocortin-5 receptors. *Cell Discovery* **2023**, *9*, 81.

(32) Luo, P.; Feng, W.; Ma, S.; Dai, A.; Wu, K.; Chen, X.; Yuan, Q.; Cai, X.; Yang, D.; Wang, M.-W.; Eric Xu, H.; Jiang, Y. Structural basis of signaling regulation of the human melanocortin-2 receptor by MRAP1. *Cell Research* **2023**, *33*, 46–54.

(33) Zhang, C.; Srinivasan, Y.; Arlow, D. H.; Fung, J. J.; Palmer, D.; Zheng, Y.; Green, H. F.; Pandey, A.; Dror, R. O.; Shaw, D. E.; Weis, W. I.; Coughlin, S. R.; Kobilka, B. K. High-resolution crystal structure of human protease-activated receptor 1. *Nature* **2012**, *492*, 387–392.

(34) Cheng, R. K. Y. et al. Structural insight into allosteric modulation of protease-activated receptor 2. *Nature* **2017**, *545*, 112–115.

(35) Luginina, A. et al. Structure-based mechanism of cysteinyl leukotriene receptor inhibition by antiasthmatic drugs. *Science Advances* **2019**, *5*.

(36) Fierro, F.; Suku, E.; Alfonso-Prieto, M.; Giorgetti, A.; Cichon, S.; Carloni, P. Agonist binding to chemosensory receptors: a systematic bioinformatics analysis. *Frontiers in Molecular Biosciences* **2017**, *4*, 63.

(37) Isberg, V.; de Graaf, C.; Bortolato, A.; Cherezov, V.; Katritch, V.; Marshall, F. H.; Mordalski, S.; Pin, J.-P.; Stevens, R. C.; Vriend, G.; Gloriam, D. E. Generic GPCR residue numbers – aligning topology maps while minding the gaps. *Trends in Pharmacological Sciences* **2015**, *36*, 22–31.

(38) Crooks, G. E.; Hon, G.; Chandonia, J.-M.; Brenner, S. E. WebLogo: A Sequence Logo Generator: Figure 1. *Genome Research* **2004**, *14*, 1188–1190.

(39) Zou, R.; Wang, X.; Li, S.; Chan, H. C. S.; Vogel, H.; Yuan, S. The role of metal ions in G protein-coupled receptor signalling and drug discovery. *WIREs Computational Molecular Science* **2021**, *12*.

(40) Pintilie, G.; Chiu, W. Validation, analysis and annotation of cryo-EM structures. *Acta Crystallographica Section D Structural Biology* **2021**, *77*, 1142–1152.

(41) Agasid, M. T.; Sørensen, L.; Urner, L. H.; Yan, J.; Robinson, C. V. The Effects of Sodium Ions on Ligand Binding and Conformational States of G Protein-Coupled Receptors—Insights from Mass Spectrometry. *Journal of the American Chemical Society* **2021**, *143*, 4085–4089.

(42) Schrödinger, LLC Maestro 2023-3, 2023; New York, NY.

(43) Huang, J.; Rauscher, S.; Nawrocki, G.; Ran, T.; Feig, M.; De Groot, B. L.; Grubmüller, H.; MacKerell, A. D. CHARMM36m: an improved force field for folded and intrinsically disordered proteins. *Nature Methods* **2017**, *14*, 71–73.

(44) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. Comparison of simple potential functions for simulating liquid water. *The Journal of Chemical Physics* **1983**, *79*, 926–935.

(45) Bussi, G.; Donadio, D.; Parrinello, M. Canonical sampling through velocity rescaling. *The Journal of Chemical Physics* **2007**, *126*, 014101.

(46) Bernetti, M.; Bussi, G. Pressure control using stochastic cell rescaling. *The Journal of Chemical Physics* **2020**, *153*, 114107.

(47) Abraham, M. J.; Murtola, T.; Schulz, R.; Páll, S.; Smith, J. C.; Hess, B.; Lindahl, E. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* **2015**, *1*, 19–25.

(48) Rodríguez-Espigares, I. et al. GPCRmd uncovers the dynamics of the 3D-GPCRome. *Nature Methods* **2020**, *17*, 777–787.