The DBD-α4 helix of EWS::FLI is required for GGAA microsatellite binding that underlies genome regulation in Ewing sarcoma

Ariunaa Bayanjargal[1,2,3], **Cenny Taslim**[1], **Iftekhar A. Showpnil**[1], **Julia Selich-Anderson**[1], **Jesse C. Crow**[1], **Stephen L. Lessnick**[1,2,3,4,5†], **Emily R. Theisen**[1,2,3,4]*

*For correspondence:
Emily.Theisen@nationwidechildrens.org (ERT)

[1]Center for Childhood Cancer, Abigail Wexner Research Institute at Nationwide Children's Hospital, Columbus, OH, 43205, USA; [2]Medical Scientist Training Program, The Ohio State University, Columbus, OH, 43210, USA; [3]Biomedical Sciences Graduate Program, The Ohio State University, Columbus, OH, 43210, USA; [4]Department of Pediatrics, The Ohio State Univeristy, Columbus, OH, 43210, USA; [5]Division of Pediatric Heme/Onc/BMT, The Ohio State University College of Medicine, Columbus, OH, 43210, USA; [†]Professor Emeritus of Pediatrics at The Ohio State University

Abstract Ewing sarcoma is the second most common bone cancer in children and young adults. In 85% of patients, a translocation between chromosomes 11 and 22 results in a potent fusion oncoprotein, EWS::FLI. EWS::FLI is the only genetic alteration in an otherwise unaltered genome of Ewing sarcoma tumors. The EWS portion of the protein is an intrinsically disordered domain involved in transcriptional regulation by EWS::FLI. The FLI portion of the fusion contains a DNA binding domain shown to bind core GGAA motifs and GGAA repeats. A small alpha-helix in the DNA binding domain of FLI, DBD-α4 helix, is critical for the transcription function of EWS::FLI. In this study, we aimed to understand the mechanism by which the DBD-α4 helix promotes transcription, and therefore oncogenic transformation. We utilized a multi-omics approach to assess chromatin organization, active chromatin marks, genome binding, and gene expression in cells expressing EWS::FLI constructs with and without DBD-α4 helix. Our studies revealed DBD-α4 helix is crucial for cooperative binding of EWS::FLI at GGAA microsatellites. This binding underlies many aspects of genome regulation by EWS::FLI such as formation of TADs, chromatin loops, enhancers and productive transcription hubs.

Introduction

Ewing sarcoma is an aggressive bone-associated tumor currently treated with dose-intense chemotherapy, radiation, and surgery (*Pappo and Dirksen, 2018*). It affects adolescents and young adults with an incidence rate of 3 per million (*Ozaki, 2015*). Of these patients, 25-35% have overt metastatic disease with a recurrence rate of 50-80% (*Gaspar et al., 2015*). Roughly a quarter of patients with localized disease also relapse (*Van Mater and Wagner, 2019*). The five-year survival rate of metastatic and relapsed patients is only 10-30% (*Stahl et al., 2011*). Treatment options for relapsed/metastatic Ewing sarcoma patients have not improved for the last four decades. The lack of efficient and tar-

geted treatment for Ewing sarcoma can be attributed to our poor understanding of how precisely Ewing sarcoma is driven by a fusion oncoprotein called EWS::FLI..

Expression of EWS::FLI results from a translocation between chromosomes 11 and 22 that fuses two genes, EWSR1 and FLI1 (*Delattre et al., 1992*). As a result, the transcription activation domain of EWSR1 and the DNA binding domain of FLI1 fuse together to create a potent transcription factor (*May et al., 1993*). The EWS domain is highly disordered and is important for multimerization and transcriptional hub formation (*Chong et al., 2018*). The FLI portion contains a DNA binding domain that binds at two distinct sites: canonical FLI1 binding sites containing a GGAA core and microsatellites containing GGAA repeats (*Gangwal et al., 2008*). At these GGAA microsatellites, EWS::FLI binding creates de-novo enhancers driving genome-wide reprogramming of enhancers to Ewing-specific enhancers (*Riggi et al., 2014*). Regulation of Ewing-specific enhancers underlies the mechanisms by which EWS::FLI regulates expression of thousands of target genes (*Grunewald et al., 2015*). More specifically, activation of genes involved in proliferation, migration and invasion pathways leads to oncogenic transformation of precursor cells to Ewing sarcoma cells (*Kauer et al., 2009*).

In our recent studies of EWS::FLI, we found a small alpha helix in the DNA binding domain, DBD-α4, to be required for transcription and regulation by the fusion protein (*Boone et al., 2021*). Interestingly, this study did not find any change in chromatin accessibility (ATAC-Seq) and genome localization of EWS::FLI constructs (CUT&RUN) when DBD-α4 helix was deleted leaving the mechanistic basis for the requirement of DBD-α4 in transcription regulation unclear. In parallel studies, we also observed that EWS::FLI expression results in widespread changes to 3D chromatin organization (*Showpnil et al., 2022*). These results together prompted us to consider whether DBD-α4 helix is contributing towards EWS::FLI's abilities to make changes in the epigenome and the local chromatin structure. To assess whether DBD-α4 helix is important in chromatin organization, we used our "knock-down/rescue" system in A673 cells (*Theisen et al., 2019*; *Boone et al., 2021*; *Showpnil et al., 2022*) in conjunction with genomics techniques including RNA-Seq, CUT&Tag, and Micro-C. These sets of experiments allowed us to characterize the involvement of DBD-α4 helix in modulating the chromatin organization, epigenetic reprogramming of enhancers, binding at GGAA microsatellites, and promoting transcription leading to transformation.

Results

DBD-α4 helix of FLI domain is required to restructure 3D chromatin in A673 cells

To assess the mechanism by which DBD-α4 promotes transcription, we depleted endogenous EWS::FLI expression with shRNA (KD) and rescued with previously published DBD and DBD+ EWS::FLI constructs with 3X-FLAG tags (Figure 1A, *Boone et al.* (*2021*)). The main difference between DBD and DBD+ constructs is the presence of DBD-α4 helix in DBD+ (Supplementary Figure 1A). We recapitulated our previous findings that DBD is unable to rescue the same number of Ewing-sarcoma specific genes as DBD+, therefore incapable of driving oncogenic transformation (Supplementary Figure 1B-D, *Boone et al.* (*2021*)).

Next, we sought to understand the role of DBD-α4 helix in 3D chromatin organization. We carried out Micro-C experiments in KD, DBD and DBD+ cells. First, we performed multidimensional scaling (MDS) analysis of the top 1000 interactions of Micro-C matrices at 500kb resolution to assess the global regulation of 3D chromatin organization (Figure 1B). KD replicates clustered together with DBD replicate 1 on both axes and with DBD replicate 2 on y-axis. DBD+ replicates, on the other hand, clustered away from both KD and DBD replicates. These observations suggest that the global chromatin structure of DBD replicates are more similar to KD than DBD+ replicates. We also assessed the contact frequency across the genome in KD, DBD and DBD+ cells at 5kb resolution (Figure 1C). We found that at the mid region of the plot (105-106 bp), DBD+ cells make more contacts in comparison to KD and DBD cells. At this genome-wide level, we observe that DBD cells display similar contact frequency as KD cells suggesting that DBD-α4 helix is required for

Figure 1. DBD-α4 helix of FLI domain is required to restructure chromatin in A673 cell. A. A schematic of DBD and DBD+ constructs used in shRNA knock-down and rescue experiments. B. Multidimensional scaling (MDS) plot of top 1000 interactions (500kb resolution) in each biological replicates. C. Genome-wide interaction frequency (ICE-corrected Micro-C counts) over genomic distance (bp) at 5kb resolution. D. Volcano plot showing differentially interacting regions (DIRs) detected at 500kb resolution for DBD+ replicates versus KD replicates. E. Volcano plot showing DIRs detected at same resolution for DBD replicates versus KD replicates. F. Violin plot of interaction distance (unit length of 500kb) for interactions (p adj < 0.2, log2FC > 0.5) detected in DBD only (mean=54Mbp), in both DBD and DBD+ conditions (mean=39.5Mbp), and in DBD+ only (mean=29.7Mbp). Boxplots depict the minimum, first quartile, median, third quartile, and maximum.* P value < 0.05, *** P value < 0.001

EWS::FLI mediated 3D chromatin restructuring.

To determine if these changes were significant, we conducted differential interaction analysis
using 500kb matrices to compare DBD and DBD+ biological replicates to KD replicates. First, we
detected 14,326 and 65 interactions in DBD+ and DBD cells respectively using cut-off values of ad-
justed p-value < 0.2 and $\log_2 FC > 0.5$. Then, we used cut-off values of adjusted p-value < 0.05 and
$\log_2 FC > 1.5$ to identify differentially interacting regions (DIRs). We found that DBD+ cells have 151
DIRs compared to KD cells (Figure 1D) whereas DBD cells only have 7 DIRs (Figure 1E). When we
used all interactions detected including DIRs for an overlap analysis, we found 31 common interac-
tions detected in both DBD and DBD+ (Figure 1F). We further assessed the distance between these
interactions and identified that DBD unique interactions (mean=54Mbp) were the most distant
followed by common (mean=39.5Mbp) and DBD+ unique interactions (mean=29.7Mbp). Taken to-
gether, these data demonstrate that the DBD-α4 helix of FLI domain is required for EWS::FLI to
restructure 3D chromatin in A673 cells.

101 Altered TAD structure in DBD cells is linked to GGAA microsatellite binding

To investigate the necessity of the DBD-α4 helix for higher-order chromatin organization, we car-
ried out topologically associated domain (TAD) analysis using HiCExplorer suite of tools (*Wolff et al.,*
2020). TADs are functional subunits of chromatin that define regulatory landscape (*Szabo et al.,*
2019). Within these domains, chromatin interactions including those involving regulatory elements
are spatially confined, often by insulators (*Dixon et al., 2012*; *Hou et al., 2012*; *Nora et al., 2012*).
Moreover, genes located in the same TAD are typically co-regulated together (*Dixon et al., 2016*;
Ramírez et al., 2018) and this co-regulation is facilitated by interactions between enhancers and
promoters, which also tend to occur within the confines of a same TAD (*Bonev et al., 2017*). Our ini-
tial step involved identifying TADs in both DBD and DBD+ matrices with hicFindTADs at 10kb, 25kb,
50kb and 100kb resolutions. Subsequently, we computed differential TADs with hicDifferential-
TAD. This analysis entailed a comparison of precomputed TAD regions of DBD and DBD+ matrices
with the same regions of KD matrix. The purpose was to detect TADs that appeared specifically in
response to the expression of DBD and DBD+ constructs in comparison to KD sample (Figure 2A).
At each resolution, the number of TADs detected when DBD+ was expressed was far greater than
when DBD was expressed.

To exclude duplicated TADs when combining TADs from different resolutions, we used the de-
fault threshold value of 5000bp as the minimum boundary difference required to consider two
TADs as distinct. With this approach, we found 6,690 total TADs for DBD and 15,412 total TADs for
DBD+ (Figure 2B). To understand the local biological context of these total TADs in DBD and DBD+
cells, we annotated the whole width of the TADs with CTCF and FLAG CUT&Tag peaks (FDR < 0.05,
FC >8, counts > 80, IDR < 0.01) (Figure 2B). In DBD cells, we detected 11.2% of the TADs bound by
CTCF only, 12.9% bound by DBD FLAG only, 36.3% bound by both CTCF and DBD FLAG, and 39.6%
bound by neither. In contrast, for DBD+, we detected 7.4% and 7.0% of the TADs bound by only
CTCF and only DBD+ FLAG, respectively, 29.0% bound by both and 56.6% bound by neither. These
observations highlight both the increased number of total TADs detected in DBD+ cells and a no-
table increase in the percentage of TADs detected in DBD+ cells with no CTCF or FLAG association.
This suggests a broader role for DBD-α4 helix in genome-wide reorganization of TADs by EWS::FLI,
consistent with the results discussed above relating to Figure 1.

Next, we overlapped total TADs from DBD and DBD+ conditions and identified 2,530 TADs com-
mon to both. There were 4,345 unique TADs in DBD cells and 12,971 unique TADs in DBD+ cells
(Figure 2C). To understand the transcriptional activity driven by the DBD-α4 helix at GGAA repeats,
we focused on the whole width of unique TADs containing DBD and DBD+ FLAG peaks at GGAA
microsatellites (Figure 2D-2H). This analysis narrowed the TADs down to 868 unique ones for DBD
cells and 2,460 unique TADs for DBD+ cells (Figure 2E). We observed that within these TAD sub-
sets containing FLAG peaks at GGAA microsatellites, the intensity of the DBD+ FLAG peaks was
higher compared to DBD FLAG peaks (Figure 2D). Moreover, TADs uniquely bound by DBD+ at

Figure 2. Altered TAD structure in DBD cells is linked to GGAA microsatellite binding. A. Number of TADs detected in DBD and DBD+ compared to KD at resolutions of 10kb, 25kb, 50kb and 100kb. B. Proportion of TADs (compared to KD) bound by FLAG, CTCF, both, or neither. C. Venn diagram of overlap between DBD and DBD+ TADs (compared to KD). D-H. Comparison of DBD and DBD+ unique TADs. D. Binding intensity of unique FLAG peaks (FDR < 0.05, FC > 8, counts > 80, IDR < 0.01) across the width of DBD and DBD+ unique TADs. E. Width of DBD and DBD+ unique TADs in bp. F. Expression level of significantly upregulated genes within unique TADs in DBD and DBD+ bound by FLAG. G. Length of microsatellites bound by unique FLAG peaks in DBD and DBD+ conditions in bp. H. Density of GGAA motif in the microsatellites calculated as (# of motif x 4)/(length of microsatellites) in DBD and DBD+ unique TADs bound by unique FLAG peaks. Boxplots depict the minimum, first quartile, median, third quartile, and maximum. *** P value < 0.001

GGAA microsatellites were not only wider (Figure 2E) but also contained a higher number of dif-
ferentially expressed genes compared to those bound only by DBD (Figure 2F and Supplementary
Figure 2A). Further analysis of GGAA microsatellites bound by DBD and DBD+ peaks within unique
TADs revealed that the DBD+ FLAG peaks were associated with longer microsatellites (Figure 2G)
and denser GGAA motifs (Figure 2H, density calculated by multiplying the number of GGAA motif
by 4 and dividing by the total length of the microsatellites) in contrast those bound by DBD alone.
These findings suggest that DBD+ accesses longer and denser microsatellites within larger TADs
and binds to these sites with higher binding intensity than DBD. Furthermore, these TADs with
DBD+ bound microsatellites show greater changes in genes expression suggesting a functional
role for the DBD-α4 helix in gene regulation within TADs.

TAD boundaries are regions between TADs that act as insulators (*Dixon et al., 2012*) and are
often marked by active chromatin (*Ay et al., 2014*). We conducted a similar analysis as above in a
20kb region around unique TAD boundaries (Supplementary Figure 2B-2G). We observed a compa-
rable percentage of the unique TAD boundaries were bound by only CTCF peaks in DBD and DBD+
cells (36.3% and 32.2%, respectively). The percentage of boundaries with FLAG peaks in DBD and
DBD+ cells were 15.2% and 8.6%, respectively (Supplementary Figure 2B). When we considered
boundaries that overlap with FLAG peaks at microsatellites (Supplementary Figure 2C-2G), we ob-
serve similar findings as those spanning the whole width of TADs (Figure 2D, 2F-2H). The FLAG
peaks at DBD+ specific TAD boundaries were of higher intensity than FLAG peaks at DBD specific
TAD boundaries (Supplementary Figure 2C). Next we assessed gene expression at TAD boundaries
and found DBD+ boundaries contained more significantly regulated genes (Supplementary Figure
2D-2E). The GGAA microsatellites bound by DBD+ FLAG peaks at boundaries were also longer and
denser than DBD bound microsatellites (Supplementary Figure 2F-2G). These data demonstrate

Figure 3. DBD and DBD+ form loops at GGAA microsatellites, but DBD+ rescues more and shorter loops. A. Number of loops detected in DBD and DBD+ compared to KD at resolutions of 1kb at short-range (<50kb), mid-range (50kb-500kb), and long-range (>500kb). B. Venn diagram of overlap between DBD and DBD+ uniquely gained loops (compared to KD). C. Expression level of significant genes overlapped with uniquely gained loop anchors of DBD and DBD+. Means = 0.45 (DBD) and 0.77 (DBD+) D. Peak intensity of unique FLAG peaks (FDR < 0.05, FC > 8, counts > 80, IDR < 0.01) at anchors of uniquely gained loops in DBD and DBD+ cells. Means = 5.01 and 6.30 per DBD and DBD+ E. Length of microsatellites bound by unique FLAG peaks at the anchors of DBD and DBD+ uniquely gained loops in bp. Means = 22.5bp (DBD) and 38bp (DBD+) F. Density of GGAA motif in the microsatellites calculated as (# of motif x 4)/(length of microsatellites) at the anchors of DBD and DBD+ uniquely gained loops bound by unique FLAG peaks. Means = 0.51 (DBD) and 0.70 (DBD+). Boxplots depict the minimum, first quartile, median, third quartile, and maximum.* P value < 0.05, *** P value < 0.001

₁₆₁ that DBD-α4 helix affects overall DNA binding at GGAA microsatellites and gene expression in the
₁₆₂ context of TAD organization.

DBD and DBD+ form loops at GGAA microsatellites, but DBD+ rescues more and shorter loops

₁₆₅ Previously, we observed that EWS::FLI promotes chromatin loop formation at GGAA microsatellites
₁₆₆ leading to enhancer activation and upregulation of gene expression (*Showpnil et al., 2022*). We
₁₆₇ next asked whether DBD-α4 helix participates in functions of EWS::FLI pertaining to the formation
₁₆₈ of loops by calling chromatin loops with Mustache algorithm (*Roayaei Ardakany et al., 2020*) in
₁₆₉ matrices of DBD and DBD+ cells compared to KD cells at 1kb resolution. We detected loops in
₁₇₀ 3 ranges defined as short-range (<50kb), mid-range (50kb-500kb), and long-range (>500kb) loops
₁₇₁ based on the linear genomic distance between loop anchors (Figure 3A). The only range with a
₁₇₂ difference between DBD and DBD+ cells was short-range loops. Specifically, we found that DBD+
₁₇₃ expressed cells gained 1,913 more short-range loops than DBD expressed cells compared to KD.
₁₇₄ Both conditions had a similar number of short-range lost loops (5,100 and 5,463 for DBD and DBD+
₁₇₅ respectively, Figure 3A). For an overlap analysis between DBD and DBD+ conditions, we included all
₁₇₆ lost loops and found 5610 loops that were commonly lost in both conditions (Supplementary Figure
₁₇₇ 3A). When the same overlap analysis was carried out for DBD and DBD+ gained loops of all ranges,
₁₇₈ we found only 219 common loops (Figure 3B). This small overlap of gained loops suggested that
₁₇₉ although they have a similar number of total gained loops, DBD and DBD+ cells contain loops with
₁₈₀ very different anchors. These sets of observations suggest that formation of distinct set of short-

Figure 4. DBD+ rescues de-novo enhancer formation at microsatellites. A. PCA plot of H3K27ac peaks in biological replicates of KD, DBD and DBD+. B. Venn diagram of overlap of H3K27ac peaks (FDR < 0.05, FC > 8, counts > 80, IDR < 0.01) of DBD and DBD+. C. Percentage of H3K27ac peaks at microsatellites in common, DBD unique and DBD+ unique peaks. D. Number of H3K27ac peaks constituting typical and super enhancers called in DBD and DBD+ conditions. E. Constituent size (in bp) of typical and super enhancers in DBD and DBD+ conditions. F. Expression level of significantly upregulated genes at DBD and DBD+ super enhancers. Boxplots depict the minimum, first quartile, median, third quartile, and maximum. Circles depict outliers. *** P value < 0.001.

range loops is a potential mechanism by which DBD-α4 helix restructures chromatin organization
in A673 cells.
Prompted by the above difference in short-range loop formation, we further investigated uniquely
gained loops of all ranges in DBD (9,005) and DBD+ (10,774) conditions (Figure 3B). When individual
anchors of loops were overlapped with GGAA microsatellites, we found 92.4% and 92.2% of DBD
and DBD+ loop anchors were at GGAA microsatellites. We then assessed overall gene expression
at all uniquely gained loop anchors and found DBD+ loop anchors associated with more gene acti-
vation (Figure 3C) and downregulation compared to DBD specific anchors (Supplementary Figure
3B). Next, we focused our analysis on gained loop anchors with FLAG peak at GGAA microsatellites
in DBD and DBD+ cells. We found 102 and 20 FLAG peaks overlapping with DBD and DBD+ loop an-
chors respectively (Figure 3D). At this subset, FLAG peaks had a higher intensity in DBD+ compared
to DBD expressed cells. The GGAA microsatellites overlapping with DBD and DBD+ FLAG peaks at
gained loop anchors were further characterized in length and density (Figure 3E-F). We found that
DBD+ FLAG peaks were bound at GGAA microsatellites that are long and dense in comparison to
DBD bound microsatellites. Taken together, these data indicate that DBD-α4 helix facilitates for-
mation of short-range loops by promoting binding at long and dense GGAA microsatellites, thus
leading to more gene regulation.

DBD+ rescues de-novo enhancer formation at GGAA microsatellites

199 Multiple studies have shown that binding of EWS::FLI at GGAA microsatellites results in genome-
200 wide de-novo enhancer formation (*Riggi et al., 2014*; *Tomazou et al., 2015*; *Sheffield et al., 2017*).

Ewing-specific enhancer establishment at GGAA microsatellites are functionally linked to the oncogenic transformation (*Johnson et al., 2017a*). We therefore sought to understand whether DBD-α4 helix of EWS::FLI has any role in modulating the Ewing-specific enhancer landscape. To assay the enhancer landscape, we collected H3K27ac CUT&Tag data from KD, DBD and DBD+ cells. Principal component analysis of H3K27ac localization shows that the DBD replicates were clustered closer to the KD replicates while being in between the KD and the DBD+ replicates (Figure 4A), suggesting that DBD-α4 helix is required to reshape the enhancer landscape. Next, we did overlap analysis of H3K27ac peaks of DBD and DBD+ cells. There were 7,854 H3K27ac peaks unique to DBD cells, 35,583 peaks common between DBD and DBD+ cells, and 26,317 peaks unique to DBD+ cells (Figure 4B), highlighting the vastly different enhancer landscapes of DBD and DBD+ conditions. Next, we sought to understand the extent of GGAA microsatellite driven H3K27 acetylation by conducting overlap analysis of H3K27ac peaks and GGAA microsatellites that were previously characterized (Figure 4C, *Johnson et al.* (*2017a*)). Interestingly, for each group of H3K27ac peaks we considered (i.e. DBD, common, and DBD+), we found around 40% of the H3K27ac peaks overlapped with GGAA microsatellites. Overall, a significant portion of H3K27ac peaks correlated with GGAA microsatellites in both DBD and DBD+ conditions underscoring the importance of GGAA repeats in formation of novel enhancers by fusions involving the transcriptional activation domain of EWSR1 with the FLI1 DNA binding domain. However, these results suggest that the DBD-α4 helix is required for modulating the enhancer landscape in Ewing cells to specifically promote oncogenic transformation.

Transcription factors have been shown to establish super-enhancers at key genes that drive cellular identity (*Whyte et al., 2013*). We therefore sought to understand whether DBD-α4 helix participates in formation of super-enhancers. We utilized ROSE algorithm to define enhancers and super-enhancers in DBD and DBD+ cells (*Lovén et al., 2013*; *Whyte et al., 2013*). We found 9,480 and 10,370 typical enhancers in DBD and DBD+, respectively (Figure 4D). We also identified 707 and 454 super-enhancers in DBD and DBD+, respectively. Although the super-enhancers associated with DBD+ were fewer in numbers compared to DBD, these super-enhancers were larger in size (Figure 4E) and contained more H3K27ac peaks (Figure 4D). We also assessed the expression level of genes associated with super-enhancers and found DBD+ super-enhancer genes had more gene activation (Figure 4F) and deactivation (Supplementary Figure 4A). These data suggest DBD-α4 helix is required in formation of enhancers and super-enhancers, therefore important in regulation of genes from those highly clustered regions of the genome.

DBD-α4 helix promotes binding at longer and denser GGAA microsatellites

The results of TAD, loop and enhancer analysis hinted towards a mechanism of DBD-α4 helix in binding GGAA microsatellites that differed from our prior CUT&RUN analysis which showed no difference in binding (*Boone et al., 2021*). We decided to further investigate binding patterns of DBD and DBD+ FLAG peaks genome-wide from our CUT&Tag dataset generated for this study. We found 8,118 DBD+ and 10,101 DBD FLAG peaks genome-wide that passed the threshold of FDR < 0.05, FC > 8, counts > 80, IDR < 0.01. When we overlapped the two sets of FLAG peaks, we found 6,102 common, 3,999 DBD unique and 2,016 DBD+ unique FLAG peaks (Figure 5A). When each group of FLAG peaks was overlapped with genomic GGAA microsatellites, we found that 70.68% of DBD+ unique and 73.04% of common peaks bind at GGAA microsatellites (Figure 5B). When we considered DBD unique peaks, we found GGAA microsatellites binding shifted down to 53.04% (Figure 5B, DBD bar). Common FLAG peaks were found to have the highest peak intensity, followed by DBD+ peaks then DBD peaks (Figure 5C). Taken together, these data demonstrate a role of the DBD-α4 helix in binding at a subset of GGAA microsatellites.

We therefore further characterized the GGAA microsatellites bound by these 3 sets of peaks: DBD unique, common and DBD+ unique peaks (Figure 5D-H). We assessed the length of microsatellites and found uniquely DBD bound GGAA microsatellites to be shorter (mean=36.61bp) than uniquely DBD+ bound GGAA microsatellites (43.25bp) (Figure 5D). GGAA microsatellites bound by

Figure 5. DBD-α4 helix of FLI promotes binding at longer and denser GGAA microsatellites. A. Venn diagram of overlap between FLAG peaks (FDR < 0.05, FC > 8, counts > 80, IDR < 0.01) of DBD and DBD+ cells. B. Percentage of FLAG peaks bound at microsatellites in common, DBD unique and DBD+ unique peaks. C. Intensity of peaks in DBD unique (mean=5.75) , DBD common (mean=7.76), DBD+ common (mean=7.75) and DBD+ unique (mean=6.83) FLAG peaks. D. Length (in bp) of GGAA microsatellites bound by DBD unique (mean=36.61), common in both (mean=50.94), and DBD+ unique (mean=43.25) FLAG peaks. E.Total number of GGAA motifs in microsatellites bound by DBD unique (mean=4.66), common in both (mean=8.70), and DBD+ unique (mean=7.78) FLAG peaks. F. Maximum consecutive number of GGAA motifs in microsatellites bound by DBD unique (mean=1.48), common in both (mean=5.21), and DBD+ unique (mean=5.32) FLAG peaks G. Percent of GGAA motif in the microsatellites calculated as (# of motif x 4)/(length of microsatellites) bound by DBD unique (mean=0.54), common in both (mean=0.65), and DBD+ unique (mean=0.69) FLAG peaks. H. Maximum number of insertion (gaps in bp) in microsatellites bounds by DBD unique (mean=10.1), common (mean=9.1), and DBD+ unique (mean=8.3) FLAG peaks. Boxplots depict the minimum, first quartile, median, third quartile, and maximum. * P value < 0.05, ** P value < 0.01, and *** P value < 0.001

²⁵¹ both DBD and DBD+ were the longest (50.94bp) (Figure 5D). When we quantified the total num-
²⁵² ber of GGAA repeats, we found that the mean number of GGAA motifs in uniquely DBD bound
²⁵³ GGAA microsatellites was 4.66 compared to 7.78 in uniquely DBD+ bound GGAA microsatellites
²⁵⁴ and 8.70 in GGAA microsatellites commonly bound in both conditions that were significantly dif-
²⁵⁵ ferent from each other (Figure 5E). Next, we quantified the maximum consecutive GGAA repeats
²⁵⁶ within a microsatellite and found that the DBD unique peaks contained GGAA repeats no longer
²⁵⁷ than 2 repeats (Figure 5F) whereas common and DBD+ unique peaks were localized at approxi-
²⁵⁸ mately 5 consecutive GGAA repeats (mean = 5.21 and 5.32 for common and DBD+, respectively,
²⁵⁹ Figure 5F). Next we plotted GGAA motif density and observed that DBD unique microsatellites are
²⁶⁰ on average (mean=0.54) less dense compared to common (0.65) and DBD+ (0.69) bound GGAA
²⁶¹ microsatellites (Figure 5G). Inverse of GGAA density is the max gap or number of non-GGAA bp in
²⁶² the microsatellites. We then assessed the gap size in each of the groups and found DBD+ binds
²⁶³ microsatellites that contain the least amount of gaps (mean=8.3) followed by common (9.1) then
²⁶⁴ DBD unique peak microsatellites (10.1) (Figure 5H). These analyses demonstrate that the DBD-α4
²⁶⁵ helix is required for effective binding at GGAA microsatellites that are medium in length and denser
²⁶⁶ in its GGAA motifs.

²⁶⁷ **DBD-α4 helix is necessary for productive transcription hub formation at GGAA mi-**
²⁶⁸ **crosatellites**

²⁶⁹ Transcription is a process that is regulated across large genomic distances, time, and space (*Cramer,*
²⁷⁰ *2019*; *Lee and Young, 2013*). One of the emerging concepts that unifies observations of such a reg-
²⁷¹ ulation is the concept of transcriptional hubs. Transcriptional hubs are actively transcribed regions
²⁷² containing clusters of transcription factors and RNA Pol II, and they are highly characterized by en-
²⁷³ hancers (*Lim and Levine, 2021*). EWS::FLI has been shown to form such hubs via the intrinsically dis-
²⁷⁴ ordered region of EWS at GGAA microsatellites (*Chong et al., 2018*). Since DBD-α4 helix is required
²⁷⁵ in effective binding at GGAA microsatellites and the downstream regulation of 3D chromatin, we
²⁷⁶ sought to understand if DBD-α4 helix participates in the function of EWS::FLI in the formation of
²⁷⁷ transcriptional hubs.

²⁷⁸ To characterize EWS::FLI-driven transcription hubs, we turned to a region containing the FCGRT
²⁷⁹ gene since EWS::FLI at this locus is shown to form a hub at GGAA repeats (Figure 6A, *Chong et al.*
²⁸⁰ (*2018*)). The FCGRT gene resides in a 250kb TAD (top panel) in both DBD and DBD+ cells. Con-
²⁸¹ sidering that TADs are highly self-interacting regions and that interactions across different TADs
²⁸² are limited by insulator elements (*Dixon et al., 2012*; *Hou et al., 2012*; *Nora et al., 2012*), we used
²⁸³ TADs as proxy for hubs as we did not expect hubs to extend beyond TAD boundaries. Within the
²⁸⁴ hub, we observed instances where DBD and DBD+ clusters appear in similar locations. In DBD
²⁸⁵ cells, however, we noticed increases in the intensity of all peaks except the FCGRT promoter GGAA
²⁸⁶ microsatellite peak (Figure 6A, magenta tracks). When we zoomed in on this microsatellite, DBD
²⁸⁷ and DBD+ peaks had a similar binding pattern (Figure 6B, magenta and green tracks). Despite this
²⁸⁸ similarity, DBD binding did not facilitate the formation of chromatin loops at the FCGRT promoter
²⁸⁹ as efficiently as DBD+ (inverted red arcs). Furthermore, this lack of loop formation correlated with
²⁹⁰ decreased enhancer marks (Figure 6A, green tracks and bar) and decreased expression of the FC-
²⁹¹ GRT gene (black tracks, padj < 0.05, FC > 2). Even though binding at the FCGRT microsatellite is
²⁹² comparable between DBD and DBD+, DBD is less able to modulate transcription and chromatin at
²⁹³ this microsatellite with high GGAA motif density (0.79, Figure 6B).

²⁹⁴ We also studied the DBD peaks that had higher intensity compared to DBD+ peaks. The majority
²⁹⁵ of these peaks were at microsatellites of varying density (0.35-0.75) and two peaks at the promot-
²⁹⁶ ers of PRR12 and RRAS were not at GGAA repeats (Figure 6A). These peaks were still near GGAA
²⁹⁷ microsatellites with densities of 0.53 and 0.28 for PRR12 and RRAS peaks respectively. Upon analy-
²⁹⁸ sis of the RNA-sequencing data, we found that the expression of genes associated with this subset
²⁹⁹ of peaks were not significantly differentially expressed in both DBD and DBD+ cells compared to
³⁰⁰ KD (Supplementary Table 1). The only gene with significantly changed expression was FCGRT and

Figure 6. DBD-α4 helix promotes formation of transcription hubs by effective binding at microsatellites. A. 250kb region on chr 19 containing FCGRT and other genes. TADs are depicted on 1kb matrices (DBD/KD and DBD+/KD). Uniquely gained loops are shown as red inverted arcs. FLAG CUT&Tag bigwig tracks depicted in magenta. GGAA microsatellites in hg19. CTCF CUT&Tag track is in blue middle row. H3K27ac tracks are in green. Enhancers and super-enhancers are shown as green bars. Gene expression is in black tracks. B. FCGRT promoter region containing GGAA microsatellites. C. NOSIP promoter region containing GGAA repeats. D. Model of EWS::FLI-driven transcription hub

only in DBD+ cells compared to KD. Taken together, these data demonstrate that DBD-α4 helix is required in regulating expression of the FCGRT gene.

Another locus we investigated is CCND1, where hub formation at GGAA repeats is also reported (*Chong et al., 2018*). The CCND1 locus is located at the border of a 700kb TAD in both DBD and DBD+ cells (Supplementary Figure 5). However, in DBD+ cells, formation of two smaller TADs nested within the larger TAD is detected with an increased number of loops (Supplementary Figure 5B). DBD and DBD+ peaks cluster in similar locations across the hub. However, DBD+ binding at MYEOV and CCND1 promoters were the only ones that effectively modulated chromatin at these microsatellites to promote gene expression (black tracks, padj < 0.05, FC >2). The CCND1 promoter microsatellite has a density of 1 meaning that it only consists of GGAA repeats without any gaps, whereas MYEOV microsatellites densities were 0.44 and 0.28. These findings further demonstrate the preference of DBD-α4 helix for binding at dense GGAA repeats and the increased binding intensity seen for DBD without the DBD-α4 helix at shorter and less dense GGAA microsatellites. Taken together, these data provide examples of transcriptional hubs spanning a large genomic distance containing numerous genes and show how transcriptional output at these hubs can be regulated by EWS::FLI binding at GGAA microsatellites.

Discussion

Detailed differential analysis of genomic binding patterns of DBD and DBD+ was not previously performed. In our previous study, we used CUT&RUN to identify DBD and DBD+ peaks and found 90% overlap (*Boone et al., 2021*). In this study, CUT&Tag showed that this overlap is only 60%. This provided us with an opportunity to discern meaningful difference in genomic localization and sequence characteristics of bound loci across three groups: DBD unique, common, and DBD+ unique peaks. Importantly, the methods utilize different enzymes for the digestion step: Micrococcal nuclease (MNase) for CUT&RUN and Tn5 transposase for CUT&Tag. Thus, a few possible factors might explain the differences in peak detection between the two techniques. MNase enzyme has cleavage preferences at sites rich in adenylate, deoxyadenylate, and thymidylate (*Cuatrecasas et al., 1967*). It also shown to have a preference for open chromatin or nucleosome-free regions (*Henikoff et al., 2011*; *Weiner et al., 2010*). Because the specific effects of EWS::FLI binding on GGAA repeats in chromatin is unknown, these loci may be particularly susceptible to excess cleavage by MNase in a way that biases CUT&RUN results. In contrast, the Tn5 transposase is especially efficient at integrating adapters into open chromatin without chromatin degradation and may, therefore, be particularly efficient at capturing binding at GGAA repeat regions. Additionally, CUT&RUN generally has lower signals, higher background, and lower yields compared to CUT&Tag (*Kaya-Okur et al., 2019*). We thus attribute the difference in detection of FLAG peaks in our previous and current studies to the different enzymes and their ability to recognize and access repetitive elements such as GGAA repeats.

With the current results, we favor a model whereby DBD-α4 helix stabilizes collective binding at high density GGAA microsatellites. Our findings underscore the specific characteristics of GGAA repeats bound by EWS::FLI to drive pathogenesis of Ewing sarcoma. At Ewing-specific gene loci such as FCGRT and CCND1 (Figure 6 and Supplementary Figure 5), we found dense microsatellites. This finding was also recapitulated at the genome-wide level as seen in the preference of DBD+ unique peaks for denser microsatellites compared to common peaks (Figure 5G). The DBD+ unique peaks also show a preference of binding at microsatellites that are longer than the DBD unique peaks and shorter than the common peaks (Figure 5D) aligning with our previous findings that an optimal number of GGAA repeats is required for binding by EWS::FLI in Ewing sarcoma transformation (*Johnson et al., 2017b*).

Although our data demonstrates that DBD-α4 helix is required in collective binding at GGAA repeats, we are still unable to decipher the exact mechanism of stabilization at such repeats with our current set of data. There are a few possible mechanisms that exist. DBD-α4 helix could be directly interacting with the DNA or the DNA binding domain of the adjacently bound EWS::FLI

molecule. Another possibility is DBD-α4 helix affects its function through interactions with other epigenetic regulators or transcription machinery proteins. Alternatively, intramolecular interaction with the EWS domain may stabilize DNA binding and could further promote phase condensates. We favor the last mechanism since de-novo ability of EWS::FLI binding GGAA repeats depends on the presence of the EWS domain (*Johnson et al., 2017a*). Further directed studies are needed to address these possible mechanisms.

The multi-omics approach utilized in this study provided us with an opportunity to character-ize transcription hubs driven by EWS::FLI genome-wide. We have shown the clustering of EWS::FLI at GGAA microsatellites underlies the formation of local 3D features such as TADs and chromatin loops. If we use TADs as proxies for hubs, we detected thousands more of these hubs in DBD+ cells compared to DBD cells, highlighting the importance of DBD-α4 helix in binding at dense GGAA re-peats and formation of hubs across the genome. We also observed thousands of loops at unique microsatellites for both DBD and DBD+ cells adding detail to the architecture of hubs, often rep-resented as flower-shaped structures of many loops (*Zhu et al., 2021*). We also probed another aspect of hubs with our H3K27ac CUT&Tag data: the presence of enhancers and super-enhancers. We showed that DBD-α4 helix promotes more active marks genome-wide compared to DBD lead-ing to formation of enhancers and super-enhancer. Finally, we presented FCGRT and CCND1 loci as specific examples of productive transcription hubs driven by EWS::FLI binding at dense GGAA microsatellites. Our study thus uncovered a surprising role for FLI DBD in the process of hub for-mation which is usually attributed to the EWS low complexity domain.

Because EWS::FLI is the sole driver mutation in Ewing sarcoma tumors, it is an attractive ther-apeutic target. However, many approaches to target EWS::FLI have been hampered by common issues that arise with targeting transcription factors, such as its location in the nucleus, its abun-dance in cells, and its lack of an enzymatic pocket to design a small molecule tailored to target it (*Knott et al., 2019*). We propose DBD-α4 helix is a promising therapeutic target as it doesn't directly bind the major groove of the DNA. Additionally, because the DBD-α4 is a helix, it provides a struc-tured region to design small molecules to disrupt its interaction unlike the EWS low complexity domain. Moreover, its importance in maintaining effective binding at GGAA microsatellites offers an opportunity to target EWS::FLI at the most mechanistically important sites in pathogenesis of Ewing sarcoma. Further studies are needed to clarify the mechanism by which the DBD-α4 helix promotes effective binding at GGAA microsatellites and regulates transcription hub formation.

Materials and Methods

Constructs and retroviruses

Mammalian expression constructs used include: Retroviral vectors encoding shRNA for luciferase-RNAi and EWS::FLI-RNAi, as well as cDNA-containing vectors encoding 3xFLAG-EF, 3xFLAG-EF DBD, 3xFLAG-EF DBD+ (*Boone et al., 2021*). The EWS::FLI DBD, and EWS::FLI DBD+ were ordered as gene blocks (Integrated DNA Technologies) and cloned into the pMSCV-hygro plasmid between BamHI and AgeI restriction sites.

Cell culture methods

HEK293-EBNA cells were grown at 37°C, 5% CO2 in Dulbecco's modified Eagle's medium (DMEM, Corning Cellgro 10-013-CV), with 10% heat-inactivated fetal bovine serum (Gibco 16000-044), 1% penicillin/streptomycin/glutamine (P/S/Q, Gibco 10378-016), and 0.3 mg/Ml Geneticin (Gibco 10131-027). A673 was obtained from American Type Culture Collection (ATCC, Manassas, VA). These cells were grown at 37C, 5% CO2 in DMEM with 10% fetal bovine serum, 1% P/S/Q, and 1% sodium pyruvate (Gibco 11360-070). For knockdown of endogenous EWS::FLI in A673, cells were infected with RNAi virus and subsequently infected with the cDNA-containing virus to rescue the cells. After 48 hours, cells were selected with puromycin (100 µg/ul) and hygromycin (150 µg/ul); and allowed to grow for 7-8 days prior to collection for downstream analysis. Cells were tested regularly for

Mycoplasma using the PCR based Universal Mycoplasma Detection Kit (ATCC, 30-1012K). Cell line identities were confirmed by short tandem repeat (STR) profiling (Genetica LabCorp, USA), last performed in February 2022.

Transfection, virus production and transduction

For the generation of retroviruses, HEK293-EBNA cells were co-transfected with retroviral expression plasmids, vesicular stomatitis virus G glycoprotein (VSV-G) and gag/pol packaging plasmids. Briefly, 2.5 x 106 HEK293-EBNA cells were seeded in a 10 cm tissue culture dish the day before transfection, resulting in 60-70% confluency the day of transfection. 10 μg of each plasmid (gag-pol, vsv-g, and transfer plasmid) were combined with 2 ml Opti-MEM I Reduced Serum Medium (Gibco 31985070) and 90 μl MirusBio TransIT-LT1 Transfection Reagent (Mirus MIR2306) and incubated at room temperature for 20 minutes. The transfection mix was then added drop wise to the cells in 3 ml culture medium. Virus-containing supernatant was collected every 4 hours on day 2 (48 hours) and 3 (72 hours) post transfection, pooled, filtered and stored at -80°C. Ewing Sarcoma cells were transduced with viral supernatants using polybrene (8 μg/ml), followed by selection with appropriate antibiotics at 48 hours post infection. In the case of knockdown/rescues, cells were selected with 0.5-2 μg/ml Puromycin and 50-150 μg/ml Hygromycin B (Thermo 10687010) for 7-10 days.

Immunodetection

Whole-cell protein extraction was completed using Pierce™ RIPA buffer (ThermoFisher 88901) supplemented with Protease Inhibitor Cocktail (Sigma P8340-5ML) on ice for 30 minutes. For nuclear extracts cell pellets were resuspended in LB1 buffer (50 mM Tris-HCl pH 7.5, 20 mM NaCl, 1 mM EDTA, 0.5% NP-40, 0.25%, Triton-X 100, 10% Glycerol, 1 mM DTT, Protease Inhibitor Cocktail) and incubated on a nutator for 10 minutes at 4C. After a 5 minute spin at 400xg and 4C, the pellets were washed with LB2 buffer (10 mM Tris-HCl pH 7.5, 20 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, 1 mM DTT, Protease Inhibitor Cocktail) and centrifuged again for 5 minutes at 400xg and 4C. The nuclear pellets were resuspended in a small volume of RIPA buffer, incubated on ice for 30 minutes, followed by a 30 minute centrifugation step at maximum speed and 4C.

Protein concentration was determined using Pierce™ BCA™ Protein Assay Kit (Thermo Scientific 23225). 15-35 μg of protein samples were ran on precast 4-15% gradient gels (Bio Rad) and transferred to nitrocellulose membranes (Invitrogen). 4-15% Mini-PROTEAN® TGX™ precast protein gel (Bio-Rad 4561084) and resolved at 120 V. Proteins were transferred to a nitrocellulose membrane (Thermo IB23002) using the iBlot™ 2 Gel Transfer Device (Thermo IB21001). The membrane was blocked with Odyssey® PBS Blocking Buffer (Li-Cor 927-40003) for 1 hour at room temperature. Immunoblotting was performed overnight at 4C using the following primary antibodies: anti-FLI1 rabbit (Abcam ab15289, 1:1000); monoclonal anti-FLAG M2 mouse (Sigma F1804-200UG, 1:1000); anti--Tubulin [DM1A] mouse (Abcam ab7291, 1:2000), and anti-Lamin B1 [EPR8985(B)] rabbit (Abcam ab133741, 1:1000). The membrane was washed with TBS/0.1% Tween 20 (TBS-T) and incubated with IRDye secondary antibodies (IRDye® 680LT donkey anti-rabbit IgG, IRDye® 800CW goat anti-rabbit IgG (H + L), IRDye® 800CW goat anti-mouse IgG1 specific secondary antibody, Li-Cor 926-68023, 926-32211, 926-32350, 1:2000) for 1 hour at room temperature. After a final wash step with TBS-T the membrane was imaged using the Li-Cor Odyssey CLx Infrared Imaging System. ImageJ software was used to perform densitometry analysis.

Soft agar assays

The anchorage-independent growth capacity of Ewing sarcoma cells was assessed using soft agar assays. Cells were seeded at a density of 7500 cells in 6-cm3 tissue culture dish in duplicate in 0.8% SeaPlaque GTG agarose (Lonza 50111) mixed with Iscove's Modified Dulbecco's medium (Gibco 12200-036) containing 20% FBS, penicillin/streptomycin/glutamine and puromycin/hygromycin. Agars

were imaged at least 14 days after seeding and colony counts were quantified using ImageJ software (V1.51).

RNA-sequencing experiments, data processing, and analysis

RNA-sequencing was performed on 3 biological replicates of KD, DBD and DBD+. Total RNA were extracted using RNeasy Extraction Kit and submitted to the Nationwide Children's Hospital Institute for Genomic Medicine for RNA quality measurement, library preparation, and sequencing. Briefly, cDNA libraries were prepared from total RNA with TruSeq Stranded mRNA Kit (Illumina 20020594) and sequenced on Illumina NovaSeq SP to generate 150-bp paired-end reads. We used in-house RNA-sequencing pipeline to process and analyze the data. Low-quality reads (q < 10) and adapter sequences were trimmed to align to hg19 genome using STAR (*Dobin et al., 2013*). After alignment, the reads were counted and differential analysis performed using DESeq2 (*Love et al., 2014*).

CUT&Tag experiments

CUT&Tag was performed as described in *Kaya-Okur et al.* (*2019*) with slight modifications. 250,000 cells per CUT&Tag condition were bound to BioMag® Plus Concanavalin A-coated magnetic beads (Bangs Laboratories, BP531) and incubated with the primary antibody (anti-FLAG M2 mouse, Sigma F1804-200UG, 1:100) overnight at 4 °C and secondary antibody (rabbit anti-mouse, Abcam ab46540, 1:100) for 1 hour at room temperature.

Adapter-loaded protein A-Tn5 fusion protein was added at a dilution of 1:250 and incubated for 1 hour at room temperature. To activate the Tn5, tagmentation buffer containing MgCl2 was added and samples were incubated for 1 hour at 37°C. Reactions were stopped by addition of EDTA and DNA was solubilized with SDS and Proteinase K for 1 hour at 50°C. Total DNA was purified using Phenol/Chloroform extraction followed by Ethanol precipitation. CUT&Tag libraries were prepared with NEBNext HiFi 2x PCR Master Mix (NEB M0541S) and indexed primers (*Buenrostro et al., 2015*) using a combined Annealing/Extension step at 63°C for 10 seconds and 15 cycles followed by a 1.1X post-amplification AMPure XP (Beckman Coulter, A63880) bead cleanup. The fragment size distributions and concentrations of the final libraries were determined using the High Sensitivity D1000 Screen tape assay and reagents (Agilent, 5067-5584 and 5067-5585) on the Agilent 2200 TapeStation System. Libraries were pooled and sequenced (2 x 150 bp paired end) on the Illumina NovaSeq S1-Xp system (Nationwide Children's Hospital Institute for Genomic Medicine).

Micro-C experiments

Micro-C kits (Catalog 21006) purchased from Dovetail Genomics were used to prepare Micro-C libraries. For each condition, multiple aliquots of $1x10^6$ cells were harvested and frozen at -80°C for at least 30 minutes. Cells were thawed in room temperature and resuspended in first PBS containing 0.3M DSG then in 37% formaldehyde to crosslink DNA. Cells were then digested with various amount of MNase to achieve the digestion profile of 40%-70% mononucleosome peak observed on TapeStation D5000 HS Screen Tape. Conditions that are to be analyzed comparatively were digested to a similar range of mononucleosome peaks (50%-70%). Once desired digestion profiles achieved, the cells were lysed and the chromatin was captured with beads to perform proximity ligation. Libraries were prepared per the protocol of Micro-C kit and each library was indexed with unique primer pairs from IDT (10009816 and 10010147). Micro-C libraries were then shallow sequenced at 7 to 8 million (2 x 150bp) read pairs on Illumina NovaSeq6000 and the QC analysis pipeline provided from Dovetail Genomics were used to assess the quality of each library. Libraries that passed the QC step was then sequenced up to 300M read pairs on NovaSeq6000.

CUT&Tag data processing and analysis

CUT&Tag experiments were carried out for 2 biological replicates of CTCF and H3K27ac and 3 biological replicates for FLAG tagged DBD and DBD+ constructs. An in-house pipeline was used to analyze CUT&Tag data (*Boone et al., 2021*; *Showpnil et al., 2022*). Quality control on raw sequencing

reads were performed with FastQC (v0.11.4) (*Andrews, 2010*). Adapter sequences and/or low qual-
ity reads were trimmed using trim_galore (0.4.4_dev) (*Krueger, 2015*). Reads were aligned to human
(hg19) and spike-in Escherichia coli (Escherichia_coli_K_12_DH10B NCBI 2008-03-17) genomes us-
ing Bowtie2 (v2.3.4.3)11,12 with the following options '–no-unal –no-mixed –no-discordant –dovetail
–phred33 -q -I 10 -X 700'. '–very-sensitive' option was added when aligning to spike-in genome (*Lang-
mead and Salzberg, 2012*). SamTools (v1.9) was used to convert sam to bam with '-bq 10' option (*Li
and Durbin, 2009*). CUT&Tag reads were spike-in normalized using DESeq2's median ratio method
to eliminate bias across different samples, minimize the effect of few outliers and appropriately
account for global occupancy changes (*Anders and Huber, 2010*). Spike-in normalized tracks were
generated and averaged across biological replicates using deepTools (*Ramirez et al., 2016*). Peaks
were called with spike-in normalization and their corresponding IgG as controls accounting for
variation between the biological replicates using MACS2 (v 2.2.7.1) (*Zhang et al., 2008*), DiffBind
(v2.14.0) (*Ross-Innes et al., 2012*; *Stark R., 2011*) and DESeq2 (v1.26.0) (*Love et al., 2014*). All dupli-
cate reads were kept in the analysis. Irreproducible Discovery Rate (IDR) (v 2.0.3) (*Li et al., 2011*)
was used to identify reproducible and consistent peaks across replicates. To ensure high quality
peaks that are most likely to represent biological signals, the final peak lists were generated with
following default thresholds: FDR < 0.05, log2 Fold-Change > 8, mean normalized counts of signal
> 80, and IDR < 0.01.

Micro-C data processing and analysis

Micro-C libraries of 2 biological replicates of KD, DBD and DBD+ were prepared. Sequenced li-
braries were processed per instructions of Dovetail Genomics. Briefly, fastq files were aligned to
hg19 reference genome using BWA-MEM algorithm with options -5SP to map mates independently
(*Li and Durbin, 2010*). Next, parse module from pairtools (**?**) was used to find ligation junctions in
Micro-C libraries with options min-mapq 40 (alignment with mapq < 40 will be marked as multi) and
max-inter-align-gap 30 (if the gap is 30 or smaller, ignore the map, if the gap is >30, mark as "null"
alignment). The parsed pair is then sorted using pairtools sort and PCR duplicates were removed
with pairtools dedup. The pairtools split command was used to split the final .pairsam into .bam
(then sorted with samtools sort) and .pairs files. Using Juicer Tools, .pairs files were converted into
HiC contact matrices (*Durand et al., 2016b*). HiC matrices were then converted to mcool matrices
using hic2cool (*Durand et al., 2016a*).

For MDS plot of individual replicates (Figure 1B), 500kb resolution cool matrices were converted
gi interaction object using hicConvertFormat from HiCExplorer (v.3.7.2) (*Wolff et al., 2020*). Then
using diffHiC package (*Lun and Smyth, 2015*), bins with low average abundance and low absolute
counts were filtered out. Filtered reads were then scaled using library size and bin pairs that were
on the diagonal line were also removed from analysis. Joint normalization of all replicates were
carried out with diffHiC. Specifically, normOffsets function was used to remove trended biases
with loess normalization and then a new set of log2-transformed counts adjusted by the negative
binomial offset were computed. Batch effect was removed using removeBatchEffect from limma
package (*Ritchie et al., 2015*) before plotting the top 1000 interactions. For distance-decay plot (Fi-
gre 1C), individual replicates were combined. First, the combined matrices were normalized using
hicNormalize function from HiCExplorer to scale the libraries to the smallest library (*Wolff et al.,
2020*). Scaled libraries were then plotted for diagnostic plots to determine the thresholds to use
in hicCorrectMatrix function for ICE normalization. ICE-corrected 5kb matrices were then plotted
with hicPlotDistVsCounts. For differentially interacting region analysis (Figure 1D-F), multiHiCcom-
pare package was used (*Stansfield et al., 2019*). Briefly, individual replicates of KD, DBD and DBD+
were loess normalized and pairwise comparison of DBD+ to KD and DBD to KD was done using
QLF (quasi-likelihood) method with batch effect correction. Volcano plots of differentially interact-
ing regions with padj < 0.05 and fold-change > 1.5 plotted for each comparison (DBD+ and DBD vs
KD).

For TAD analysis, individual replicates were combined and then scaled using the smallest library

size with hicNormalize from HiCExplorer (v.3.7.2) (*Wolff et al., 2020*). Then matrices at 10kb, 25kb,
50kb and 100kb resolutions were ICE-corrected with the thresholds determined from diagnostic
plots. First, using hicFindTADs function, we called TADs at the previously mentioned 4 resolutions
for DBD and DBD+ matrices. Then, hicDifferentialTAD used to compute differential TADs by com-
paring the precomputed DBD and DBD+ TAD regions with the same regions of KD matrix. Dif-
ferential TADs from each resolution were combined using hicMergeDomains with default –value
of 5000 to account for duplicated TADs. For CUT&Tag peak annotation, peaks (FDR < 0.05, log2
Fold-Change > 8, mean normalized counts of signal > 80, and IDR < 0.01) were overlapped with
findOverlaps functions from GenomicRanges package (*Lawrence et al., 2013*).

For loop calling, combined replicates at 1kb resolution matrices that were scaled and normal-
ized in same manner for TAD analysis were used with Mustache (v.1.2.0) (*Roayaei Ardakany et al.,*
2020). Mustache uses scale-space theory in computer vision to detect chromatin loops. For differ-
ential loops compared to KD matrix, diff_mustache.py was used to detect loops that were gained
in DBD and DBD+ compared to KD and loops that were lost in DBD and DBD+ compared to KD
matrix.

Statistical analysis

559 When comparing means of two groups, two-sided Student's t test was used. When comparing
560 more than two groups, ANOVA test was used with Tukey honest significant differences test. * P
561 value < 0.05, ** P value < 0.01, and *** P value < 0.001

Supplementary Data

Supplementary Figure 1: A. DBD-α4 helix of FLI1 depicted on dsDNA (PDB) B. Knock-down of endogenous
EWS::FLI detected with FLI1 ab and rescue of wtEF, DBD, and DBD+ detected with FLAG ab. C. A PCA plot of
RNA-seq experiments in A673 cells. D. Representative image of soft agar colony plates and quantification of
three biological replicates.

Supplementary Figure 2. A. Expression level of significant genes ovarlapped with unique TADs in DBD (mean=-0.64) and DBD+ (mean=-1.16) bound by FLAG at GGAA microsatellites. B. Proportion of TAD boundaries bound by FLAG, CTCF or neither. C-G. Comparison of DBD and DBD+ unique TAD boundaries. C. Binding intensity of unique FLAG peaks (FDR < 0.05, FC > 8, counts > 80, IDR < 0.01) at boundaries of DBD and DBD+ unique TADs. D. Expression level of significantly upregulated genes overlapped with boundaries of unique TADs in DBD and DBD+ bound by FLAG at GGAA microsatellites. E. Expression level of significantly downregulated genes overlapped with boundaries of unique TADs in DBD and DBD+ bound by FLAG at GGAA microsatellites. F. Length of microsatellites bound by unique FLAG peaks at the boundaries of DBD and DBD+ conditions in bp. G. Percent of GGAA motif in the microsatellites calculated as (of motif x 4)/(length of microsatellites) at the boundaries of DBD and DBD+ unique TADs bound by unique FLAG peaks. Boxplots depict the minimum, first quartile, median, third quartile, and maximum. *** P value < 0.001

Supplementary Figure 3. A. Venn diagram of overlap between DBD and DBD+ uniquely lost loops (compared to KD). B. Expression level of downregulated genes overlapped with uniquely gained loop anchors of DBD and DBD+. Means = -0.68, -1.25, 0.35, -1.08 * P value < 0.05, *** P value < 0.001. Boxplots depict the minimum, first quartile, median, third quartile, and maximum.

Supplementary Figure 4. A. Expression level of downregulated genes at DBD and DBD+ super enhancers. *** P value < 0.001 Boxplots depict the minimum, first quartile, median, third quartile, and maximum.

Supplementary Figure 5 . A. CCND1 hub in 700kb region on chr 11 in DBD cells. TADs are depicted on 1kb matrices (DBD/KD). Uniquely gained loops are shown as red inverted arcs. FLAG CUT&Tag bigwig tracks depicted in magenta. GGAA microsatellites in hg19. CTCF CUT&Tag track is in blue middle row. H3K27ac tracks are in green. Enhancers and super-enhancers are shown as green bars. Gene expression is in black tracks.

Supplementary Figure 5. B. CCND1 hub in 700kb region on chr 11 in DBD+ cells. TADs are depicted on 1kb matrices (DBD+/KD). Uniquely gained loops are shown as red inverted arcs. FLAG CUT&Tag bigwig tracks depicted in magenta. GGAA microsatellites in hg19. CTCF CUT&Tag track is in blue middle row. H3K27ac tracks are in green. Enhancers and super-enhancers are shown as green bars. Gene expression is in black tracks.

Table 1. Supplementary Table 1. Differential expression of FCGRT hub genes in DBD and DBD+ compared to KD.

Gene Symbol	DBD FC	DBD padj	DBD+ FC	DBD+ padj
ALDH16A1	1.207	0.04635288	1.107	0.329212288
RPL13A	-1.132	0.021442589	-1.209	7.46E-05
RPL13AP5	-1.109	0.490879542	-1.149	0.290788743
RPS11	-1.068	0.337413798	-1.177	0.001887497
FCGRT	1.639	0.0002149	2.44	2.54E-13
RCN3	-1.588	6.85E-06	-1.406	0.000957843
NOSIP	1.065	0.446781941	-1.041	0.636826207
PRRG2	1.117	0.759840545	-1.042	0.916808784
PRR12	1.177	0.006810095	1.088	0.20377503
RRAS	-1.655	1.63E-17	-1.383	5.10E-08
SCAF1	1.065	0.357378849	-1.07	0.283421905
IRF3	1.081	0.4016866	1.196	0.011473037
BCL2L12	1.255	0.006880506	1.359	6.60E-05
PRMT1	1.151	0.015484657	1.145	0.014515775
ADM5	1.122	0.740397828	1.064	0.863762064

Data availability

The sequencing dataset generated and analyzed in this study are available in the Gene Expression Omnibus and accessible at GSE249578. All other data not part of GEO submission are available from the corresponding author on a reasonable request.

Code availability

All codes used in analysis of the sequencing data are compiled in Supplementary files.

Acknowledgements

We thank Ali Snedden, Yuan Zhang, and John Burian at High Performance Computing group at Nationwide Children's Hospital for their support; and Institute for Genomic Medicine at Nationwide Children's Hospital for sequencing support. We are grateful for the comments and discussion from members of Lessnick lab, Theisen lab and Bioinformatics group at Nationwide Children's Hospital. We also thank Dr. Elaine R. Mardis for her thoughtful comments on this manuscript. This research was supported by NIH U54 CA231641 to S.L.L. and NIH T32 CA269052 to A.B.

Conflict of interest statement

S.L.L. declares competing interests as a member of the scientific advisory board for Salarius Pharmaceuticals with financial holdings and receipt of consultation fees. S.L.L. is also a listed inventor on United States Patent No. US 7,939,253 B2, 'Methods and compositions for the diagnosis and treatment of Ewing's sarcoma,' and United States Patent No. US 8,557,532, 'Diagnosis and treatment of drug-resistant Ewing's sarcoma.' E.R.T. has received research funding from Salarius Pharmaceuticals. This support is unrelated to the data presented in this manuscript.

References

Anders S, Huber W. Differential Expression Analysis for Sequence Count Data. Genome Biology. 2010 Oct; 11(10):R106. doi: 10.1186/gb-2010-11-10-r106.

Andrews S, FastQC: A Quality Control Tool for High Throughput Sequence Data [Online]; 2010.

Ay F, Bailey TL, Noble WS. Statistical Confidence Estimation for Hi-C Data Reveals Regulatory Chromatin Con-
tacts. Genome Research. 2014 Jun; 24(6):999–1011. doi: 10.1101/gr.160374.113.

Bonev B, Mendelson Cohen N, Szabo Q, Fritsch L, Papadopoulos GL, Lubling Y, Xu X, Lv X, Hugnot JP, Tanay A,
Cavalli G. Multiscale 3D Genome Rewiring during Mouse Neural Development. Cell. 2017 Oct; 171(3):557–
572.e24. doi: 10.1016/j.cell.2017.09.043.

Boone MA, Taslim C, Crow JC, Selich-Anderson J, Byrum AK, Showpnil IA, Sunkel BD, Wang M, Stanton BZ,
Theisen ER, Lessnick SL. The FLI Portion of EWS/FLI Contributes a Transcriptional Regulatory Function That Is
Distinct and Separable from Its DNA-binding Function in Ewing Sarcoma. Oncogene. 2021 Jul; 40(29):4759–
4769. doi: 10.1038/s41388-021-01876-5.

Buenrostro JD, Wu B, Litzenburger UM, Ruff D, Gonzales ML, Snyder MP, Chang HY, Greenleaf WJ. Single-Cell
Chromatin Accessibility Reveals Principles of Regulatory Variation. Nature. 2015 Jul; 523(7561):486–490. doi:
10.1038/nature14590.

Chong S, Dugast-Darzacq C, Liu Z, Dong P, Dailey GM, Cattoglio C, Heckert A, Banala S, Lavis L, Darzacq X, Tjian
R. Imaging Dynamic and Selective Low-Complexity Domain Interactions That Control Gene Transcription.
Science. 2018 Jul; 361(6400). doi: 10.1126/science.aar2555.

Cramer P. Organization and Regulation of Gene Transcription. Nature. 2019 Sep; 573(7772):45–54. doi:
10.1038/s41586-019-1517-4.

Cuatrecasas P, Fuchs S, Anfinsen CB. Catalytic Properties and Specificity of the Extracellular Nuclease of Staphy-
lococcus Aureus. The Journal of Biological Chemistry. 1967 Apr; 242(7):1541–1547.

Delattre O, Zucman J, Plougastel B, Desmaze C, Melot T, Peter M, Kovar H, Joubert I, Jongt PD, Rouleau G,
Aurias A, Thomas G. Gene Fusion with an ETS DNA-binding Domain Caused by Chromosome Translocation
in Human Tumours. NATURE. 1992; 359:162–165.

Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, Hu M, Liu JS, Ren B. Topological Domains in Mammalian Genomes
Identified by Analysis of Chromatin Interactions. Nature. 2012 Apr; 485(7398):376–80. doi: 10.1038/na-
ture11082.

Dixon JR, Gorkin DU, Ren B. Chromatin Domains: The Unit of Chromosome Organization. Molecular Cell. 2016
Jun; 62(5):668–680. doi: 10.1016/j.molcel.2016.05.018.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. STAR: Ultrafast
Universal RNA-seq Aligner. Bioinformatics. 2013 Jan; 29(1):15–21. doi: 10.1093/bioinformatics/bts635.

Durand NC, Robinson JT, Shamim MS, Machol I, Mesirov JP, Lander ES, Aiden EL. Juicebox Provides a Vi-
sualization System for Hi-C Contact Maps with Unlimited Zoom. Cell Syst. 2016 Jul; 3(1):99–101. doi:
10.1016/j.cels.2015.07.012.

Durand NC, Shamim MS, Machol I, Rao SS, Huntley MH, Lander ES, Aiden EL. Juicer Provides a One-
Click System for Analyzing Loop-Resolution Hi-C Experiments. Cell Syst. 2016 Jul; 3(1):95–8. doi:
10.1016/j.cels.2016.07.002.

Gangwal K, Sankar S, Hollenhorst PC, Kinsey M, Haroldsen SC, Shah AA, Boucher KM, Watkins WS, Jorde LB,
Graves BJ, Lessnick SL. Microsatellites as EWS/FLI Response Elements in Ewing's Sarcoma. Proceedings of
the National Academy of Sciences. 2008 Jul; 105(29):10149–10154. doi: 10.1073/pnas.0801073105.

Gaspar N, Hawkins DS, Dirksen U, Lewis IJ, Ferrari S, Le Deley MC, Kovar H, Grimer R, Whelan J, Claude L, Delattre
O, Paulussen M, Picci P, Sundby Hall K, van den Berg H, Ladenstein R, Michon J, Hjorth L, Judson I, Luksch
R, et al. Ewing Sarcoma: Current Management and Future Approaches Through Collaboration. Journal of
Clinical Oncology. 2015 Sep; 33(27):3036–3046. doi: 10.1200/JCO.2014.59.5256.

Grunewald TG, Bernard V, Gilardi-Hebenstreit P, Raynal V, Surdez D, Aynaud MM, Mirabeau O, Cidre-Aranaz F,
Tirode F, Zaidi S, Perot G, Jonker AH, Lucchesi C, Le Deley MC, Oberlin O, Marec-Berard P, Veron AS, Reynaud
S, Lapouble E, Boeva V, et al. Chimeric EWSR1-FLI1 Regulates the Ewing Sarcoma Susceptibility Gene EGR2
via a GGAA Microsatellite. Nat Genet. 2015 Sep; 47(9):1073–8. doi: 10.1038/ng.3363.

Henikoff JG, Belsky JA, Krassovsky K, MacAlpine DM, Henikoff S. Epigenome Characterization at Single Base-
Pair Resolution. Proceedings of the National Academy of Sciences of the United States of America. 2011 Nov;
108(45):18318–18323. doi: 10.1073/pnas.1110731108.

Hou C, Li L, Qin ZS, Corces VG. Gene Density, Transcription, and Insulators Contribute to the Partition of the Drosophila Genome into Physical Domains. Mol Cell. 2012 Nov; 48(3):471–84. doi: 10.1016/j.molcel.2012.08.031.

Johnson KM, Mahler NR, Saund RS, Theisen ER, Taslim C, Callender NW, Crow JC, Miller KR, Lessnick SL. Role for the EWS Domain of EWS/FLI in Binding GGAA-microsatellites Required for Ewing Sarcoma Anchorage Independent Growth. Proceedings of the National Academy of Sciences. 2017 Sep; 114(37):9870–9875. doi: 10.1073/pnas.1701872114.

Johnson KM, Taslim C, Saund RS, Lessnick SL. Identification of Two Types of GGAA-microsatellites and Their Roles in EWS/FLI Binding and Gene Regulation in Ewing Sarcoma. PLOS ONE. 2017 Nov; 12(11):e0186275. doi: 10.1371/journal.pone.0186275.

Kauer M, Ban J, Kofler R, Walker B, Davis S, Meltzer P, Kovar H. A Molecular Function Map of Ewing's Sarcoma. PLoS ONE. 2009 Apr; 4(4):e5415. doi: 10.1371/journal.pone.0005415.

Kaya-Okur HS, Wu SJ, Codomo CA, Pledger ES, Bryson TD, Henikoff JG, Ahmad K, Henikoff S. CUT&Tag for Efficient Epigenomic Profiling of Small Samples and Single Cells. Nat Commun. 2019 Apr; 10(1):1930. doi: 10.1038/s41467-019-09982-5.

Knott MML, Hölting TLB, Ohmura S, Kirchner T, Cidre-Aranaz F, Grünewald TGP. Targeting the Undruggable: Exploiting Neomorphic Features of Fusion Oncoproteins in Childhood Sarcomas for Innovative Therapies. Cancer and Metastasis Reviews. 2019 Dec; 38(4):625–642. doi: 10.1007/s10555-019-09839-9.

Krueger F. Trim Galore. A wrapper tool around Cutadapt and FastQC to consistently apply quality and adapter trimming to FastQ files. 2015; .

Langmead B, Salzberg SL. Fast Gapped-Read Alignment with Bowtie 2. Nat Methods. 2012 Mar; 9(4):357–9. doi: 10.1038/nmeth.1923.

Lawrence M, Huber W, Pages H, Aboyoun P, Carlson M, Gentleman R, Morgan MT, Carey VJ. Software for Computing and Annotating Genomic Ranges. PLoS Comput Biol. 2013; 9(8):e1003118. doi: 10.1371/journal.pcbi.1003118.

Lee TI, Young RA. Transcriptional Regulation and Its Misregulation in Disease. Cell. 2013 Mar; 152(6):1237–1251. doi: 10.1016/j.cell.2013.02.014.

Li H, Durbin R. Fast and Accurate Short Read Alignment with Burrows-Wheeler Transform. Bioinformatics. 2009 Jul; 25(14):1754–60. doi: 10.1093/bioinformatics/btp324.

Li H, Durbin R. Fast and Accurate Long-Read Alignment with Burrows-Wheeler Transform. Bioinformatics. 2010 Mar; 26(5):589–95. doi: 10.1093/bioinformatics/btp698.

Li Q, Brown JB, Huang H, Bickel PJ. Measuring Reproducibility of High-Throughput Experiments. The Annals of Applied Statistics. 2011; 5(3). doi: 10.1214/11-aoas466.

Lim B, Levine MS. Enhancer-Promoter Communication: Hubs or Loops? Current opinion in genetics & development. 2021 Apr; 67:5–9. doi: 10.1016/j.gde.2020.10.001.

Love MI, Huber W, Anders S. Moderated Estimation of Fold Change and Dispersion for RNA-seq Data with DESeq2. Genome Biol. 2014; 15(12):550. doi: 10.1186/s13059-014-0550-8.

Lovén J, Hoke HA, Lin CY, Lau A, Orlando DA, Vakoc CR, Bradner JE, Lee TI, Young RA. Selective Inhibition of Tumor Oncogenes by Disruption of Super-Enhancers. Cell. 2013 Apr; 153(2):320–334. doi: 10.1016/j.cell.2013.03.036.

Lun AT, Smyth GK. diffHic: A Bioconductor Package to Detect Differential Genomic Interactions in Hi-C Data. BMC Bioinformatics. 2015 Aug; 16:258. doi: 10.1186/s12859-015-0683-0.

May WA, Lesnick SL, Braun BS, Klemsz M, Lewis BC, Lunsford LB, Hromas R, Denny CT. The Ewing's Sarcoma EWS/FLI-1 Fusion Gene Encodes a More Potent Transcriptional Activator and Is a More Powerful Transforming Gene than FLI-1. Molecular And Cellular Biology. 1993; p. 7393–7398.

Nora EP, Lajoie BR, Schulz EG, Giorgetti L, Okamoto I, Servant N, Piolot T, van Berkum NL, Meisig J, Sedat J, Gribnau J, Barillot E, Bluthgen N, Dekker J, Heard E. Spatial Partitioning of the Regulatory Landscape of the X-inactivation Centre. Nature. 2012 Apr; 485(7398):381–5. doi: 10.1038/nature11049.

Ozaki T. Diagnosis and Treatment of Ewing Sarcoma of the Bone: A Review Article. Journal of Orthopaedic Science. 2015; 20(2):250–263. doi: 10.1007/s00776-014-0687-z.

Pappo AS, Dirksen U. Rhabdomyosarcoma, Ewing Sarcoma, and Other Round Cell Sarcomas. J Clin Oncol. 2018 Jan; 36(2):168–179. doi: 10.1200/JCO.2017.74.7402.

Ramirez F, Ryan DP, Gruning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dundar F, Manke T. deepTools2: A next Generation Web Server for Deep-Sequencing Data Analysis. Nucleic Acids Res. 2016 Jul; 44(W1):W160–5. doi: 10.1093/nar/gkw257.

Ramírez F, Bhardwaj V, Arrigoni L, Lam KC, Grüning BA, Villaveces J, Habermann B, Akhtar A, Manke T. High-Resolution TADs Reveal DNA Sequences Underlying Genome Organization in Flies. Nature Communications. 2018 Jan; 9(1):189. doi: 10.1038/s41467-017-02525-w.

Riggi N, Knoechel B, Gillespie SM, Rheinbay E, Boulay G, Suvà ML, Rossetti NE, Boonseng WE, Oksuz O, Cook EB, Formey A, Patel A, Gymrek M, Thapar V, Deshpande V, Ting DT, Hornicek FJ, Nielsen GP, Stamenkovic I, Aryee MJ, et al. EWS-FLI1 Utilizes Divergent Chromatin Remodeling Mechanisms to Directly Activate or Repress Enhancer Elements in Ewing Sarcoma. Cancer Cell. 2014 Nov; 26(5):668–681. doi: 10.1016/j.ccell.2014.10.004.

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. Limma Powers Differential Expression Analyses for RNA-sequencing and Microarray Studies. Nucleic Acids Res. 2015 Apr; 43(7):e47. doi: 10.1093/nar/gkv007.

Roayaei Ardakany A, Gezer HT, Lonardi S, Ay F. Mustache: Multi-Scale Detection of Chromatin Loops from Hi-C and Micro-C Maps Using Scale-Space Representation. Genome Biology. 2020 Sep; 21(1):256. doi: 10.1186/s13059-020-02167-0.

Ross-Innes CS, Stark R, Teschendorff AE, Holmes KA, Ali HR, Dunning MJ, Brown GD, Gojis O, Ellis IO, Green AR, Ali S, Chin SF, Palmieri C, Caldas C, Carroll JS. Differential Oestrogen Receptor Binding Is Associated with Clinical Outcome in Breast Cancer. Nature. 2012 Jan; 481(7381):389–93. doi: 10.1038/nature10730.

Sheffield NC, Pierron G, Klughammer J, Datlinger P, Schönegger A, Schuster M, Hadler J, Surdez D, Guillemot D, Lapouble E, Freneaux P, Champigneulle J, Bouvier R, Walder D, Ambros IM, Hutter C, Sorz E, Amaral AT, de Álava E, Schallmoser K, et al. DNA Methylation Heterogeneity Defines a Disease Spectrum in Ewing Sarcoma. Nature Medicine. 2017 Mar; 23(3):386–395. doi: 10.1038/nm.4273.

Showpnil IA, Selich-Anderson J, Taslim C, Boone MA, Crow JC, Theisen ER, Lessnick SL. EWS/FLI Mediated Reprogramming of 3D Chromatin Promotes an Altered Transcriptional State in Ewing Sarcoma. Nucleic Acids Research. 2022 Sep; 50(17):9814–9837. doi: 10.1093/nar/gkac747.

Stahl M, Ranft A, Paulussen M, Bölling T, Vieth V, Bielack S, Görtitz I, Braun-Munzinger G, Hardes J, Jürgens H, Dirksen U. Risk of Recurrence and Survival after Relapse in Patients with Ewing Sarcoma. Pediatric Blood & Cancer. 2011 Oct; 57(4):549–553. doi: 10.1002/pbc.23040.

Stansfield JC, Cresswell KG, Dozmorov MG. multiHiCcompare: Joint Normalization and Comparative Analysis of Complex Hi-C Experiments. Bioinformatics. 2019 Sep; 35(17):2916–2923. doi: 10.1093/bioinformatics/btz048.

Stark R BG. DiffBind: Differential Binding Analysis of ChIP-Seq Peak Data. . 2011; .

Szabo Q, Bantignies F, Cavalli G. Principles of Genome Folding into Topologically Associating Domains. Science Advances. 2019 Apr; 5(4):eaaw1668. doi: 10.1126/sciadv.aaw1668.

Theisen ER, Miller KR, Showpnil IA, Taslim C, Pishas KI, Lessnick SL. Transcriptomic Analysis Functionally Maps the Intrinsically Disordered Domain of EWS/FLI and Reveals Novel Transcriptional Dependencies for Oncogenesis. Genes & Cancer. 2019 Feb; 10(1-2):21–38. doi: 10.18632/genesandcancer.188.

Tomazou EM, Sheffield NC, Schmidl C, Schuster M, Schönegger A, Datlinger P, Kubicek S, Bock C, Kovar H. Epigenome Mapping Reveals Distinct Modes of Gene Regulation and Widespread Enhancer Reprogramming by the Oncogenic Fusion Protein EWS-FLI1. Cell Reports. 2015 Feb; 10(7):1082–1095. doi: 10.1016/j.celrep.2015.01.042.

Van Mater D, Wagner L. Management of Recurrent Ewing Sarcoma: Challenges and Approaches. OncoTargets and Therapy. 2019 Mar; Volume 12:2279–2288. doi: 10.2147/OTT.S170585.

Weiner LM, Surana R, Wang S. Monoclonal Antibodies: Versatile Platforms for Cancer Immunotherapy. Nature Reviews Immunology. 2010 May; 10(5):317–327. doi: 10.1038/nri2744.

Whyte WA, Orlando DA, Hnisz D, Abraham BJ, Lin CY, Kagey MH, Rahl PB, Lee TI, Young RA. Master Transcription
Factors and Mediator Establish Super-Enhancers at Key Cell Identity Genes. Cell. 2013 Apr; 153(2):307–319.
doi: 10.1016/j.cell.2013.03.035.

Wolff J, Rabbani L, Gilsbach R, Richard G, Manke T, Backofen R, Grüning BA. Galaxy HiCExplorer 3: A Web Server
for Reproducible Hi-C, Capture Hi-C and Single-Cell Hi-C Data Analysis, Quality Control and Visualization.
Nucleic Acids Research. 2020 Jul; 48(W1):W177–W184. doi: 10.1093/nar/gkaa220.

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li W, Liu XS.
Model-Based Analysis of ChIP-Seq (MACS). Genome Biol. 2008; 9(9):R137. doi: 10.1186/gb-2008-9-9-r137.

Zhu I, Song W, Ovcharenko I, Landsman D. A Model of Active Transcription Hubs That Unifies the Roles of Active
Promoters and Enhancers. Nucleic Acids Research. 2021 May; 49(8):4493–4505. doi: 10.1093/nar/gkab235.