Screening of associated bacteria of *Alexandrium catenella* on affecting PSTs

production and influence mechanism

Shanmei Zou*, Xinke Yu, Tiantian Sun, Lina Wei, Xuemin Wu

College of Resources and Environmental Science, Nanjing Agricultural University,

Nanjing 210095, China

* Corresponding authors. E-mail: zousm912@njau.edu.cn

Abstract

Alexandrium causes serious food safety and human death due to paralytic shellfish

toxins (PSTs) production. The associated bacteria can affect PSTs production of

Alexandrium. However, the influencing mechanism is still unclear. Here we firstly

screened functional associated bacteria for affecting PSTs production of

Alexandrium catenella in Yangtze Estuary and further studied their influence on

physiological process and molecular regulation of *A. catenella*. Thirteen bacteria

strains for affecting PSTs production of *A. catenella* were selected. The *A.*

catenella strains co-cultured with different functional associated bacteria all

produced more PSTs than axenic strain with antibiotic treatment. Compared with

axenic *A. catenella*, the non-axenic *A. catenella* produced more algal cells, soluble

sugar, soluble protein and neutral lipid. By RNA-seq, it was found that non-axenic

A. catenella produced more upregulated functional genes than axenic *A. catenella*.

The biosynthesis of cofactors and spliceosome were the dominant different

pathways between axenic and non-axenic *A. catenella* strains. The sxtA

expression was closely related with Arginine and proline metabolism, Arginine

biosynthesis, Fatty acid biosynthesis, TCA cycle and Glutathione metabolism,

which were all downregulated in axenic *A. catenella*. Meantime, the non-axenic *A.*

catenella under nitrogen deprivation produced less PSTs and functional genes

than non-axenic strain under common culture condition, indicating the nitrogen

significance for PSTs production. The detailed signal molecular released by

associated bacteria for regulating PSTs of *A. catenella* needs to be further studied.

Keywords: *Alexandrium catenella*; associated bacteria; PSTs; physiological

process; molecular regulation

1. Introduction

Harmful algal blooms (HABs) occur globally and have produced severe effects on fisheries, aquaculture, tourism and recreation[1, 2]. Dinoflagellates, a large group of phytoplanktonic organisms during marine HABs, play a crucial role in food webs and in balancing ecosystem energy fluxes [3-8]. Many marine dinoflagellates can cause food safety and human fatalities through their production of paralytic shellfish toxins (PSTs) which accumulate in shellfish through the food chain when contaminated seafood is consumed [2, 9-13]. The most widespread dinoflagellates species producing PSTs are in the genus *Alexandrium* [14, 15].

PSTs, mainly produced from dinoflagellate and cyanobacteria, is one of the most powerful neurotoxins such as TTX [16]. Currently, the regulation mechanism for PSTs production in cyanobacteria is clear, which mainly refers to 26 functional genes for PSTs biosynthesis [17]. However, the regulation mechanism for PSTs production of harmful dinoflagellates remains poorly understood due to the exceptionally large genome size and high gene copy number of dinoflagellates [18]. Thus, uncovering the factors that influence the biosynthesis of PSTs toxins in dinoflagellates is not only important for understanding PSTs biosynthesis mechanism but also crucial for seafood safety and human health assurance worldwide [19-23].

The environmental factors, such as light, temperature, salinity, nitrogen or phosphorous, have been shown to affect PSTs content for several *Alexandrium*

species[24]. For example, while nitrogen limitation causes the toxin content to

decrease phosphorous limitation often results in higher toxin cell quotas

[25-32].But the salinity and temperature just have limited effects on toxin

composition [28, 33].

Bacteria directly or indirectly affect the growth and toxin production of

Alexandrium species [34-40]. Originally it was considered that the PSTs in

dinoflagellate was produced by their associated bacteria, e.g., *protogonyaulax* and

Alexandrium [37, 41, 42]. However, the subsequent research has shown that the

"PSTs" produced by bacteria was actually the analogues of PSTs [43, 44]. The

false positive for PSTs detection in bacteria is mainly because of the low

concentration. Subsequently many studies indicated that the PSTs in dinoflagellate

was produced by dinoflagellates alone, but their associated bacteria could affect

their PST production [45-48]. Different bacteria community had different capacity

for affecting PSTs production of dinoflagellate [45]. It has been suggested that the

effects of bacteria on PSTs production of dinoflagellate can be explained by

altered toxin synthesis pathways through algal-bacterium interactions [40]. But

the detailed influencing mechanism is still unclear. It is hypothesized that the

associated bacteria of dinoflagellate affect the PSTs production by releasing some

signals or metabolites that regulate the growth and physiological process of

dinoflagellate [45-47]. Thus, understanding the influencing mechanism of

associated bacteria for PSTs production of dinoflagellate is not only helpful for

underlying the PSTs biosynthesis in dinoflagellate but also useful for controlling

the PSTs production by microbial ways.

Yangtze Estuary is the largest estuary in China, which plays an important role

for industry, agriculture, aquaculture and marine ecology. However, HABs often

happen in Yangtze Estuary due to the environmental pollution and eutrophication,

especially for dinoflagellate. Therefore, the production of PST from

dinoflagellates, especially from *Alexandrium* species, can cause severe shellfish

food poisoning [49]. Due to the difficulty in identifying dinoflagellate by

morphological characters, in this study, the eDNA metabarcoding [50] was

employed to reveal the community composition of dinoflagellate in Yangtze

Estuary. The associated bacteria community diversity of *A. catenella* from

Yangtze Estuary was revealed and the functional bacteria strains for affecting

PSTs production were screened. Then the physiological process and molecular

regulation of *A. catenella* under influence of associated bacteria on PSTs

production were further evaluated.

2. Materials and methods

2.1 Sample collection

A. catenella seawater samples collected from Shanghai (*A. catenella*-C,

121.92348°E, 30.862512°N), Shengsi (*A. catenella*-S, 122.454838°E,

30.700638°N) and Zhoushan (*A. catenella*-Z, 122.413229°E, 29.92582°N) in

Yangtze Estuary of China were firstly pre-filtered through a 200μm pore-size

sieve to remove debris, mesoplankton, and macroplankton. Three subsamples of

equal volume from 0.5 m, 1.0 m and 2.0 m depths were collected and mixed as

one sample from each plot. Then each water sample (~500 ml) was subsequently

filtered through a 0.22μm pore-size polycarbonate membrane (Millipore, Billerica,

MA, USA). The membranes were stored at -80 °C until DNA extraction. The

toxicity of *A. catenella* strains from different locations was detected, where the

stains with the highest toxicity was selected for experiment analysis.

2.2 Dinoflagellate community diversity

The 18S-v4-v5 and 28S primers (Table 1) were used for metabarcoding

eukaryotic plankton and dinoflagellates by Illumina-miseq high throughput

sequencing. DNA was extracted by OMEGA E.Z.N.ATM Mag-Bind Soil DNA

Kit for each sample. PCR amplification of all samples was performed in triplicate.

After sequencing, the fastq files were demultiplexed to separate the samples based

on the indexes. Reads were then analyzed and assigned using QIIME2, where the

PCR primers were removed by cut adapt package and Dada2 was applied for

denoising, dereplicate-sequence, filtering chimera and merging the paired reads

[51]. Taxonomic classification was performed by feature-classifier module with

Naive Bayes [51]. The reference library for ASV classification was referred from

Zou et al [52].

2.3 A. catenella isolation, culture and antibiotic treatment

Vegetative cells preliminarily identified as *A. catenella* based on

morphological characteristics were isolated by pipette under a microscope (40×

zoom), and transferred to a 24-well plates containing culture f/2 medium. These

strains were cultured at 19-26°C, 3000-6000lux light and light to dark ratio of

12D: 12N. Then the cultures were expanded from orifices to 250ml triangulated

vials and maintained in the laboratory using f/2 medium under the same

conditions. The obtained strains were then further detected by 18S sequencing. A

set of publicly available 18S sequences of *Alexandrium* downloaded from

GenBank was added to the new sequences to build phylogenetic trees by the

maximum-likelihood method.

For obtaining axenic strains, cefotaxime sodium and vancomycin

hydrochloride were selected as the antibacterial antibiotics after testing with

different antibiotic mixtures. 30μg/ml cefotaxime sodium and 50μg/ml

vancomycin hydrochloride were added to the sterile f/2 medium for producing

axenic *A. catenella* cultures, which was conducted for triple replicates. Tow

replicates were used to assess the axenic state of the cultures, while the remainder

was used to determine the toxicity. For non-axenic culture with nitrogen

deprivation (non-axenic-ndep) the concentration of NaNO3 was set at 1/4 of the

control group (2.94 mmol/liter).

2.4 Associated bacteria community diversity of A. catenella

The associated bacteria community diversity of *A. catenella* under laboratory

culture condition was analyzed by 16Sv3v4 and 16S metabarcoding respectively

(Table 1). *A. catenella* culture was filtered with a 0.22μm pore-size polycarbonate

membrane to obtain the associated bacteria (Millipore, Billerica, MA, USA).

DNA was extracted by OMEGA E.Z.N.ATM Mag-Bind Soil DNA Kit. PCR

amplification of all samples were triple. The QIIME2 was used for bioinformatics

analysis as above [51].

2.5 Isolation and reseeding of associated bacteria

The 2216E medium was used for isolation of associated bacteria of *A.*

catenella [53]. Each sample was plated in three plates that were incubated at 20°C,

30°C and 37°C respectively, which were then checked for colonies after 24h, 48h

and 72h. Individual bacterial colonies showing different colour, colony size or

morphology were re-isolated on the same medium and further characterized

morphologically and biochemically with microscope and Gram staining method.

Finally, the single clones were further identified by 16Sv3v4 sanger sequencing.

The isolated bacteria were then reseeded to axenic *A. catenella* with

antibiotic treatment, through ways of separate strain and different combined

strains. While *A. catenella* was cultured to logarithmic phase antibiotic treatment

was carried out to get axenic algal solution. The isolated bacterial strains were

cultured to logarithmic phase and then added to the algal solution.

2.6 Toxicity Type and content test

Taking the algal sap from the logarithmic phase, the algal cells were

collected by centrifugation at 6000 rpm·min^{-1} into 50mL polypropylene centrifuge

tubes, which were washed and centrifuged several times using ultrapure water to

ensure that the culture fluid was thoroughly removed. To the washed algal cells,

20mL of acetic acid solution at a concentration of 0.05 mol/L was added. The

algal cells were sonicated under ice bath conditions using an ultrasonic cell

disruptor with the power set at 200W and 20s of sonication paused for 40s, for a

total of 5min. After sonication, the extraction was continued using ultrasonic

assisted extraction for 15 min. the cells were centrifuged at 4000 rpm·min^{-1} for 10

min at 4 °C, and then 1 mL of the supernatant was removed from the centrifuge

tube. The supernatant was transferred to an ultrafiltration centrifuge tube (MWCO

of 10,000u) and centrifuged at 10,000 rpm·min^{-1} for 10min at 4°C. Finally, the

ultrafiltrate was collected into an injection vial and stored at 4°C for subsequent

analysis.

The type and content test of PSTs potentially produced by *A. catenella* and

associated bacteria was detected by liquid chromatography-mass spectrometry

(LC-MS). Samples were analyzed on an Ultimate 3000 ultra-high-pressure liquid

chromatography-Q-Exactive electrostatic field Orbitrap high-resolution mass

spectrometer system. The column was TSK-gel Amide-80 (3µm, 2mm×15cm).

Algal toxin extraction was performed using acetic acid and ultrasound. The

instrumental conditions were determined with reference to the method of Zhang et

al.[54].

2.7 Measurements of growth and biochemical composition

The biomass was detected based on cell number using a hemacytometer

(improved double-Neubauer). Soluble sugar was determined by phenol-sulfuric

acid method. Soluble proteins were determined using the Thomas Brilliant Blue

G-250 staining method. The neutral lipid content was determined by Nile red

fluorescence method.

2.8 RNA isolation, library construction and transcriptome sequencing

The axenic, non-axenic and non-axenic-ndep *A. catenella* strains were

selected for RNA sequencing (RNA-Seq) respectively. Triple replicates of each

strain were cultured under the same conditions and were mixed for RNA

extraction and sequencing. Total RNA was firstly extracted and qualified by

ethanol precipitation protocol and Agilent 2100 bioanalyzer (Thermo Fisher

Scientific, MA, USA) respectively. Then the mRNA was purified, fragmented and

cDNA produced. The cDNA was subsequently index ligated and enriched. The

product was validated on the Agilent Technologies 2100 bioanalyzer for quality

control. Finally, the qualified library was sequenced on the Illumina HiSeq

platform. Clean reads were mapped to the assembled unique genes by Bowtie2

(v2.2.5) [55] and the expression levels of genes was calculated by RSEM (v1.2.8)

and normalized to the FPKM [56].

Functional annotation of genes was achieved by mapping genes to different

databases (NT, NR, GO, COG, KEGG, Swiss-Prot, and InterPro) using the

software BLAST (v2.2.23) [57]. GO annotation was performed by Blast2GO (v

2.5.0) with NR annotations. PossionDis [58] was used to detect different

expression genes (DEGs), and DEGs with a fold change of >2 or <-2 and a

false-discovery rate (FDR) of <0.001 were considered significantly differentially

expressed genes. GO and KEGG enrichment analysis were performed using

Phyper, a function of R. The significant levels of terms and pathways were

corrected by Q value with a rigorous threshold (Q <0.05).

3. Results

3.1 The dinoflagellate community diversity

From the relative abundance of 18S-v4 and 28S ASVs (Figure 1A), it was shown that the Pyrrophyta accounted for a considerable proportion compared to another microalgae phylum, especially higher than Bacillariophyta (Figure 1A). The zooplankton was revealed with a high proportion but the Chlorophyta was uncovered with a low proportion. The dominant dinoflagellate species were mainly included in *Alexandrium*, *Karlodinium*, *Prorocentrum*, Dinophyceae, etc. Thereinto, the *Alexandrium fundyense*, *Alexandrium* sp.*, A. catenella* and *Alexandrium tamarense* dominated high abundance (Figure 1B). The phylogenetic tree showed that the *A. catenella* sequences from Shanghai, Shengsi and Zhoushan (Genbank: OR357631-OR357633) clustered as one clade (Figure 2A). The toxin content of *A. catenella* samples from the three locations was different (Figure 2B), consistent with the difference in bacterioplankton (Figure 2C). The *A. catenella* strain with the highest toxin content from Shengsi was selected for detailed experimental analysis.

3.2 The associated bacteria composition of A. catenella

Here the 16S and 16Sv3v4 assignments were combined together to reveal the associated bacteria composition of *A. catenella*. Six phyla, Bacteroidetes, Proteobacteria, Steptophyta, Firmicutes, Bacteroidota and Pseudomonadota, were detected. The dominant classes were Cyanbacteria, Cytophagia, Deltaproteobacteria, Flavobacteriia, Bacilli, Gammaproteobacteria,

Alphaproteobacteria and Bacteroidia. The main orders included Alteromonadales,

Flavobacteriales, Cytophagales, Rhodobacterales etc. (Figure 3). The dominant

associated bacteria species included *Paenibacillus* sp., *Yangia pacifica*, *Formosa*

sp., *Bacillus humi*, *Alteromonas macleodii*, *Marinobacter* sp., *Cesiribacter* sp.,

Flavobacteriales bacterium etc. Among them, *Marinobacter* accounted for the

highest proportion in 16S and *Paenibacillus* accounted for the highest proportion

in 16Sv3v4 (Figure 4).

3.3 Associated bacteria isolation and capacity of affecting toxin production test

A total of 31 associated bacteria of *A. catenella* were isolated, which were

identified to 13 different bacteria species by 16Sv3v4 sequences (Figure 5A).

These were *A. macleodii, Alteromonas gracilis, Alteromonas marina,*

Thalassospira marina, Alteromonas sp., *Exiguobacterium acetylicum,*

Marinobacter sp.*, Cellvibrio* sp.*, Spongiibacter* sp.*, Saccharospirillum* sp.*,*

Saccharospirillum alexandrill, Marinobacter confluentis, Marinobacterium

rambilcola   (Genbank: OR958743-OR958746,   OR960642-OR960647,

OR976523-OR976525). The 13 bacteria were assigned to four classes, six orders

and eight families (Figure S1).

With LC-MS analysis, it was showed that no PSTs was detected in the mixed

associated bacterial culture since there was no same molecular weight with PSTs

component was found (Figure 6A). On the other hand, a molecular weight of

395.0878 was detected in the algal culture, which was the same as GTX toxin at

the peak time of 6.475min (Figure 6B, C).

The capacity of isolated bacteria for affecting PSTs production of *A.*

catenella was tested separately, together with different combined bacteria strains

(Figure 5B, C). Under normal cultivation conditions, the non-axenic *A. catenella*

strains without antibiotic treatment apparently produced more PSTs than strains

with antibiotic treatment (Figure 5B). Generally, the non-axenic *A. catenella*

strains with separate and combined isolated bacteria all produced more PSTs than

axenic strain, especially for stains with combined bacteria strains (Figure 5B).

Moreover, the capacity of isolated bacteria for affecting PSTs production was

detected under nitrogen deprivation, to select typical bacteria that affects PSTs

production pronouncedly (Figure 5C). It was apparent that the toxin content

decreased for almost all treatments under nitrogen deprivation (Figure 5C).

Among the 13 bacterial strains, it was showed that *A. catenella* produced more

PSTs with five associated bacteria strains, e.g. *E. acetylicum, A.* sp*., A. macleodii,*

S. alexandrill and T. marina (Figure 5B, C).

3.4 The growth and biochemical composition for axenic and non-axenic A.

catenella strains

It was shown that the axenic *A. catenella* strain with antibiotic treatment

grew slower than non-axenic *A. catenella* strain without antibiotic treatment

during the whole growth cycle under normal cultivation conditions (Figure 7A).

Thereinto, the non-axenic *A. catenella* strain with nitrogen deprivation

(non-axenic-ndep) grew the slowest (Figure 7A). The non-axenic *A. catenella*

strain produced more soluble protein, sugar and neutral lipid than axenic *A.*

catenella strain with antibiotic treatment normal cultivation conditions (Figure

7B). On the other hand, the soluble sugar and protein content decreased and the

neutral lipid content increased in the non-axenic *A. catenella* strains with nitrogen

deprivation (non-axenic-ndep), compared with both axenic and non-axenic *A.*

catenella strains under normal cultivation conditions (Figure 7B).

3.5 Genetic regulation for PSTs production with influence of associated bacteria

A total of 45,778,990, 48,518,624 and 43,035,798 reads were generated for

each treatment of axenic, non-axenic and non-axenic-ndep *A. catenella* strains

respectively (Table 2). The clean read ratio of all samples was approximately 97%.

The total number of transcripts for three treatments were 145,580, with a mean

length 923 bp and 1382 N50 length. A total of 126,626 unigenes were obtained for

three treatments, with a mean length of 879 bp and an N50 length of 1327. Thus,

the reads and assembly quality showed a good preformation of transcriptome

sequencing for functional analysis.

Among the annotations of the six databases, the Pfam database showed

higher annotation proportion and more unique annotation (Figure 8A). Among the

transcription factors, the vast majority of unigenes were assigned to C3H,

followed by MYB_superfamily (Figure 8B). The axenic, non-axenic and

non-axenic-ndep strains all produced abundant common and distinct unigenes

(Figure 8C). The up-regulated and down-regulated genes among three *A. catenella*

treatments were different (Figure 8D). The GO functional annotations was

performed for all unigenes where the membrane part, cell part, cellular progress,

metabolic process, binding and catalytic activity accounted for the top

classification categories (Figure S2). The KEGG database assigned the

Carbohydrate metabolism, Amino acid metabolism, Translation and Folding,

sorting and degradation as the top annotation (Figure S3). It was indicated that the

biosynthesis of cofactors was the dominant KEGG pathway annotated between

axenic and non-axenic strains (Figure 9A). The dominant DEGs included

Glutathione metabolism, Plant-pathogen interaction, Arginine and proline

metabolism, Tryptophan metabolism, and spliceosome, etc. (Figure 9B). The PSTs

synthesis genes sxtA was found in *A. catenella* strains of all three treatments,

which were expressed highly in non-axenic strain (Figure 9B). The main

pathways enriched between non-axenic and non-axenic-ndep strains were

Phenylpropanoid biosynthesis, Spliceosome, etc. (Figure S4).

The pathway network related with PSTs production was further constructed

among axenic, non-axenic and non-axenic-ndep strains, including variable

physiological metabolism and PSTs production genes (Figure 10). The pathways

Arginine and proline metabolism, Arginine biosynthesis, Fatty acid biosynthesis,

TCA cycle and Glutathione metabolism were closely related with sxtA expression,

which were all downregulated in axenic strain (Figure 10A). As expected, the

plant hormone signal transduction and Indole-3-acetic acid-amido synthetase were

up-regulated in non-axenic strain (Figure 10A). Similar with non-axenic vs axenic

comparison, generally the non-axenic *A. catenella* strains with nitrogen

deprivation produced more downregulated genes than non-axenic *A. catenella*

strains under normal cultivation conditions in the various pathways of Glycolysis,

TCA cycle, Arginine biosynthesis, Fatty acid biosynthesis, etc. (Figure 10B).

Some potential functional genes connecting various pathways related with PSTs

production were found, like arginine deiminase and ornithine decarboxylase

among Arginine biosynthesis, Arginine and proline metabolism and Glutathione

metabolism.

4. Discussion

The PSTs production by dinoflagellate is associated with bacteria [59, 60].

The PSTs control by microbes is becoming important since the microbes play a

key role in microalgae' growth, metabolism and toxin production [61-63].

Elucidating the associative mechanism between dinoflagellate and associated

bacteria can control the PSTs production of dinoflagellate in environmental ways.

The metabarcoding results here indicated that the dinoflagellate and

Alexandrium were dominant microalgae phylum and species in Yangtze Estuary,

suggesting that the PSTs produced by dinoflagellate in China coast is becoming

serious. Actually, the dominant phytoplankton in Yangtze Estuary has been

transferred from diatom to dinoflagellates in recent years [64]. The

toxin-producing capacity and associated bacteria community diversity of *A.*

catenella from three Yangtze Estuary locations both differed from each other,

indicating that the associated bacteria possibly affect the PSTs production of *A.*

catenella. Green et al. (2010) also found that the associated bacteria diversity of *G.*

catenatum was related with environmental factors and speculated that the

associated bacteria regulated the physiological ecological progress to affect PSTs

production.

The LC-MS detection showed that the PSTs produced by *A. catenella* was

GTX and the associated bacteria did not produce any component of PSTs. This is

consistent with previous reports that the associated bacteria did not produce PSTs

and the "PSTs" detected from bacteria may be the analogues of PSTs or false

positive [43, 44]. Apparently, the PSTs production capacity of axenic *A. catenella*

strains with antibiotic treatment decreased compared with non-axenic strains,

which is consistent with previous reports that the associated bacteria affected the

PSTs production of dinoflagellate [45-48].

The capacity of associated bacteria for affecting PSTs production was further

detected. The *A. catenella* stains co-cultured with separate and combined

associated bacteria both produced higher levels of PSTs than axenic *A. catenella*

stains, especially with combined associated bacteria. Albinsson et al. [45] also

demonstrated that different bacterial communities had different effects on

G.catenatum toxin production. These further confirm that the associated bacteria

affect the PSTs production of dinoflagellate. In this case, it is important to

untangle the regulation mechanism of bacteria for PSTs production of

dinoflagellate.

The axenic *A. catenella* strains grew slowly compared with non-axenic

strains, indicating that the associated bacteria also affect the growth of

dinoflagellate. Likewise, the bacterial associates can modify growth dynamics of

Gymnodinium catenatum [65]. These support the concept that algal–bacterial

interaction is an important structuring mechanism in phytoplankton communities.

Also the contents of soluble sugar, soluble protein and neutral lipid in non-axenic

A. catenella strains were somewhat higher than that in axenic strains. All these

suggest that the associated bacteria affect the PSTs production of dinoflagellate by

regulating growth and physiological metabolism [38-40]. On the other hand, the

PST production capacity of non-axenic *A. catenella* strains under nitrogen

deprivation were both decreased compared to axenic and non-axenic strains under

normal culture conditions, indicating that nitrogen is an important factor for toxin

production of dinoflagellate since it is the main component of PSTs [28-30, 32].

The sxt gene cluster for STX synthesis has been elucidated in cyanobacteria

[17, 66-68]. Among the 26 sxt genes revealed in cyanobacteria, only several sxt

genes were found in dinoflagellate, including sxtA, sxtG and sxtO [69]. The

biosynthesis of PSTs starts with sxtA [66, 69]. However, the functional

mechanism of sxt for PSTs production is still unclear in dinoflagellate. The

polyketide synthase-like (PKS) is another important catalyzing enzyme for carbon

skeleton formation of algae toxins [70]. Here, both sxtA and PKS were both

included in the DEGs. We found that the sxtA expression was related with TCA

cycle, Arginine biosynthesis, proline metabolism, Glutathione metabolism, and

Fatty acid biosynthesis, which were all downregulated in axenic strains. Also,

Arginine is the main elements of PSTs. Thus, here it is clear that the associated

bacteria enhance PSTs production of *A. catenella* by upregulating the expression

of sxtA genes and their associated physiological pathways. Other sxt genes, such

as sxtG and sxtO, were not detected in this study, which may be related with the

intraspecific difference that different strains within one species have different

toxicity and possible regulation mechanism [59, 71]. Consistent with the

physiological analysis, the genetic expression related with PSTs production was

generally downregulated in *A. catenella* strain with nitrogen deprivation,

confirming the nitrogen importance in molecular regulation.

Compared with axenic strain, the Tryptophan metabolism and Plant hormone

signal transduction were also upregulated in non-axenic *A. catenella* strain,

consistent with their growth tendency. This indicates that the associated bacteria

can also promote the growth of *A. catenella* except the toxin production. It has

already been reported that the associated bacteria advanced the growth of diatom

by releasing signal molecules [72, 73]. Finally, some functional genes that were

possibly related with the PSTs production were revealed. So, the future studies

will focus on uncovering the determinant signal molecular released by associated

bacteria and their related regulation pathways for affecting PSTs production of

dinoflagellates.

5. Conclusion

The toxin production capacity, physiological process and molecular

regulation among *A. catenella* strains of axenic, non-axenic and non-axenic with

nitrogen deprivation were compared. Thirteen associated bacteria for affecting

PSTs production of *A. catenella* were selected. Compared with axenic strain with

antibiotic treatment, the non-axenic strain with associated bacteria produced more

PSTs contents, soluble sugar, soluble protein and neutral lipid. The biosynthesis of

cofactors and spliceosome were the dominant pathways between axenic and

non-axenic strains. The pathways Arginine and proline metabolism, Arginine

biosynthesis, Fatty acid biosynthesis, TCA cycle and Glutathione metabolism

were closely related with sxtA expression. The nitrogen plays key role in PSTs

production of *A. catenella* through affecting physiological and molecular

regulation. These findings highlight the importance of associated bacteria in

affecting PSTs production of dinoflagellate through physiological and molecular

regulation.

Credit authorship contribution statement

Shanmei Zou conceived the design and wrote the manuscript. Xinke Yu conducted

the experiments. Shanmei Zou, Xinke Yu, Tiantian Sun analyzed the data. Xinke

Yu and Xuemin Wu collected the samples.

Declaration of competing interest

The authors have no conflict of interest.

Data availability

The NGS sequence data of all samples obtained in this study are deposited in

NCBI. The SRA accession for the NGS sequences is PRJNA992270,

PRJNA998555. The submission information for each sample mainly includes the

reads sequence in fastq form, the sampling location, the sequence platform.

Acknowledgement

The bioinformatics analysis of the project was supported by the Bioinformatics

Center of Nanjing Agricultural University.

References

[1] E. Berdalet, L.E. Fleming, R. Gowen, K. Davidson, P. Hess, L.C. Backer, S.K. Moore, P.

Hoagland, H. Enevoldsen, Marine harmful algal blooms, human health and wellbeing:

challenges and opportunities in the 21st century, J Mar Biol Assoc U K 2015 (2016) 1-31.

[2] G.M. Hallegraef, Harmful algal blooms: a global overview, J manual on harmful marine

microalgae (2004).

[3] N.D. Canini, E.B. Metillo, R.V. Azanza, Monsoon-influenced phytoplankton community

structure in a Philippine mangrove estuary, Tropical Ecology 54 (2013) 331-343.

[4] C. Corinaldesi, M. Barucca, G.M. Luna, A. Dell'Anno, Preservation, origin and genetic

imprint of extracellular DNA in permanently anoxic deep-sea sediments, Molecular

Ecology 20 (2011) 642-654.

[5] Y.M. Lu, S. Wohlrab, M. Groth, G. Glockner, L. Guillou, U. John, Transcriptomic

profiling of Alexandrium fundyense during physical interaction with or exposure to

chemical signals from the *parasite Amoebophrya*, Molecular Ecology 25 (2016)

1294-1307.

[6] N. Rosic, E.Y.S. Ling, C.K.K. Chan, H.C. Lee, P. Kaniewska, D. Edwards, S. Dove, O.

Hoegh-Guldberg, Unfolding the secrets of coral-algal symbiosis, Isme Journal 9 (2015)

844-856.

[7] S. Tas, E. Okus, A review on the Bloom Dynamics of a Harmful Dinoflagellate

Prorocentrum minimum in the Golden Horn Estuary, Turkish Journal of Fisheries and

Aquatic Sciences 11 (2011) 673-681.

[8] C. Totti, S. Accoroni, F. Cerino, E. Cucchiari, T. Romagnoli, Ostreopsis ovata bloom

along the Conero Riviera (northern Adriatic Sea): Relationships with environmental

conditions and substrata, Harmful Algae 9 (2010) 233-239.

[9] R. Guo, H. Wang, Y.S. Suh, J.S. Ki, Transcriptomic profiles reveal the genome-wide

responses of the harmful dinoflagellate *Cochlodinium polykrikoides* when exposed to the

algicide copper sulfate, Bmc Genomics 17 (2016) 1-16.

[10] J.M. Meyer, C. Rodelsperger, K. Eichholz, U. Tillmann, A. Cembella, A. McGaughran,

U. John, Transcriptomic characterisation and genomic glimps into the toxigenic

dinoflagellate *Azadinium spinosum*, with emphasis on polykeitde synthase genes, Bmc

Genomics 16 (2015) 1-14.

[11] A. Persson, B.C. Smith, G.H. Wikfors, M. Quilliam, Grazing on toxic *Alexandrium*

fundyense resting cysts and vegetative cells by the eastern oyster (Crassostrea virginica),

Harmful Algae 5 (2006) 678-684.

[12] M. Tsujino, T. Uchida, Fate of resting cysts of *Alexandrium spp.* ingested by

Perinereis nuntia (Polychaeta) and Theola fragilis (Mollusca), Journal of Experimental

Marine Biology Ecology 303 (2004) 1-10.

[13] Z.H. Wang, X.P. Nie, S.J. Jiang, J.G. Zhao, Y. Cao, Y.J. Zhang, D.Z. Wang, Source

and profile of paralytic shellfish poisoning toxins in shellfish in Daya Bay, South China Sea,

Marine Environmental Research 72 (2011) 53-59.

[14] D.M. Anderson, T.J. Alpermann, A.D. Cembella, Y. Collos, E. Masseret, M. Montresor,

The globally distributed genus *Alexandrium*: Multifaceted roles in marine ecosystems and

impacts on human health, Harmful Algae 14 (2012) 10-35.

[15] A. Cembella, in: Physiological Ecology of Harmful Algal Blooms, Springer-Verlag,

Heidelberg, Anderson, DM, AD Cembella, GM Hallegraeff (Eds.), NATO-Advanced Study

Institute Series, (1998).

[16] M.A. Akbar, N.Y.M. Yusof, N.I. Tahir, A. Ahmad, G. Usup, F.K. Sahrani, H. Bunawan,

Biosynthesis of Saxitoxin in Marine Dinoflagellates: An Omics Perspective, Marine Drugs

18 (2020) 103.

[17] M. Troco, K. Ralf, N. Brett, Characterisation of the paralytic shellfish toxin

biosynthesis gene clusters in *Anabaena circinalis* AWQC131C and *Aphanizomenon sp.*

NH-5, (2009).

[18] S. Zhang, Z.H. Sui, L.P. Chang, K.H. Kang, J.H. Ma, F.R. Kong, W. Zhou, J.G. Wang,

L.L. Guo, H.L. Geng, J. Zhong, Q.X. Ma, Transcriptome de novo assembly sequencing

and analysis of the toxic dinoflagellate *Alexandrium catenella* using the Illumina platform,

Gene 537 (2014) 285-293.

[19] G.M. Hallegraeff, A review of harmful algal blooms and their apparent global increase,

Phycologia 32 (1993) 79-99.

[20] C.A. Scholin, D.M. Anderson, Identification of group-and strain-specific genetic

markers for globally distributed *Alexandrium*(dinophyceae).I.RFLP analysis of SSU rRNA

genes, Journal of Phycology 30 (1994) 744-754.

[21] C.A. Scholin, M. Herzog, M. Sogin, D.M. Anderson, Identification of group-specific

and strain-specific genetic-markers   for globally distributed *Alexandrium* (dinophyceae) .2.

sequence-analysis of a fragment of the LSU ribosomal-RNA gene, Journal of Phycology

30 (1994) 999-1011.

[22] S.E. Shumway, A Review of the Effects of Algal Blooms on Shellfish and Aquaculture,

Journal of the World Aquaculture Society 21 (1990) 65-104.

[23] Z. Su, M. Sheets, H. Ishida, F.H. Li, W.H. Barry, Saxitoxin blocks L-type/(Ca), Journal

of Pharmacology and Experimental Therapeutics 308 (2004) 324-329.

[24] M. Han, H. Lee, D.M. Anderson, B. Kim, Paralytic shellfish toxin production by the

dinoflagellate *Alexandrium pacificum* (Chinhae Bay, Korea) in axenic, nutrient-limited

chemostat cultures and nutrient-enriched batch cultures, Marine Pollution Bulletin 104

(2016) 34-43.

[25] D.M. Anderson, D.M. Kulis, J.J. Sullivan, S. Hall, Toxin composition variations in one

isolate of the dinoflagellate *Alexandrium-fundeyense*, Toxicon 28 (1990) 885-893.

[26] C. Bechemin, D. Grzebyk, F. Hachame, C. Hummert, S.Y. Maestrini, Effect of

different nitrogen/phosphorus nutrient ratios on the toxin content in *Alexandrium minutum*,

Aquatic Microbial Ecology 20 (1999) 157-165.

[27] G.L. Boyer, J.J. Sullivan, R.J. Andersen, P.J. Harrison, F.J.R. Taylor, Effects of

nutrient limitation on toxin production and composition in the marine dinoflagellate

Protogonyaulax-tamarensis, Marine Biology 96 (1987) 123-128.

[28] S.M. Etheridge, C.S. Roesler, Effects of temperature, irradiance, and salinity on

photosynthesis, growth rates, total toxicity, and toxin composition for *Alexandrium*

fundyense isolates from the Gulf of Maine and Bay of Fundy, Deep-Sea Research Part

Ii-Topical Studies in Oceanography 52 (2005) 2491-2500.

[29] E.H. John, K.J. Flynn, Growth dynamics and toxicity of *Alexandrium fundyense*

(Dinophyceae): the effect of changing N : P supply ratios on internal toxin and nutrient

levels, European Journal of Phycology 35 (2000) 11-23.

[30] S. Maestrini, C. Bechemin, D. Grzebyk, C. Hummert, Phosphorus limitation might

promote more toxin content in the marine invader dinoflagellate *Alexandrium minutum*,

Plankton Biology Ecology 47 (2000) 7-11.

[31] P. Marsot, Variations in volume and nutrient (N and P) content of the dinoflagellate

Alexandrium tamarense in N-limited continuous culture, Oceanologica Acta 20 (1997)

639-643.

[32] D.Z. Wang, D.P.H. Hsieh, Growth and toxin production in batch cultures of a marine

dinoflagellate *Alexandrium tamarense* HK9301 isolated from the South China Sea,

Harmful Algae 4 (2005) 401-410.

[33] G. Taroncher-Oldenburg, D.M. Kulis, D.M. Anderson, Coupling of saxitoxin

biosynthesis to the G(1) phase of the cell cycle in the dinoflagellate *Alexandrin fundyense*:

Temperature and nutrient effects, Natural Toxins 7 (1999) 207-219.

[34] W.R. Dantzer, R.E. Levin, Bacterial influence on the production of paralytic shellfish

toxins by dinoflagellated algae, Journal of Applied Microbiology 83 (1997) 464-469.

[35] G.J. Doucette, Interactions between bacteria and harmful algae: a review, Natural

Toxins 3 (1995) 65-74.

[36] G.J. Doucette, C.L. Powell, in: VIII International conference on Harmful algae-

Abstracts and Posters Classification. vp. 1997., (1997).

[37] S. Gallacher, K.J. Flynn, J.M. Franco, E.E. Brueggemann, H.B. Hines, Evidence for

production of paralytic shellfish toxins by bacteria associated with *Alexandrium spp*

(Dinophyta) in culture, Applied and Environmental Microbiology 63 (1997) 239-245.

[38] S. Gallacher, E.A. Smith, Bacteria and paralytic shellfish toxins, Protist 150 (1999)

245-255.

[39] M. Kodama, Paralytic Shellfish Poisoning Toxins: Biochemistry and Origin,

Aqua-BioScience Monographs 3 (2010) 1-38.

[40] E.W. Maas, R.M. Latter, J. Thiele, A.M. Waite, H.J.L. Brooks, Effect of multiple

antibiotic treatments on a paralytic shellfish toxin-producing culture of the dinoflagellate

Alexandrium minutum, Aquatic Microbial Ecology 48 (2007) 255-260.

[41] M. Kodama, T. Ogata, S. Sato, Bacterial production of saxitoxin, Agricultural biological

chemistry 52 (1988) 1075-1077.

[42] E.A. Smith, F.H. Mackintosh, F. Grant, S. Gallacher, Sodium channel blocking (SCB)

activity and transformation of paralytic shellfish toxins (PST) by dinoflagellate-associated

bacteria, Aquatic Microbial Ecology 29 (2002) 1-9.

[43] T.R. Baker, G.J. Doucette, C.L. Powell, G.L. Boyer, F.G. Plumley, GTX(4) imposters:

characterization of fluorescent compounds synthesized by Pseudomonas stutzeri SF/PS

and Pseudomonas/Alteromonas PTB-1, symbionts of saxitoxin-producing *Alexandrium*

spp, Toxicon 41 (2003) 339-347.

[44] C.A. Martins, P. Alvito, M.J. Tavares, P. Pereira, G. Doucette, S. Franca,

Reevaluation of Production of Paralytic Shellfish Toxin by Bacteria Associated with

Dinoflagellates of the Portuguese Coast, Applied Environmental Microbiology 69 (2003)

5693-5698.

[45] M.E. Albinsson, A.P. Negri, S.I. Blackburn, C.J.S. Bolch, Bacterial Community Affects

Toxin Production by *Gymnodinium catenatum*, Plos One 9 (2014).

[46] D.H. Green, M.C. Hart, S.I. Blackburn, C. Bolch, Bacterial diversity of *Gymnodinium*

catenatum and its relationship to dinoflagellate toxicity, Aquatic Microbial Ecology 61

(2010) 73-87.

[47] A.Y.T. Ho, D.P.H. Hsieh, P.Y. Qian, Variations in paralytic shellfish toxin and homolog

production in two strains of *Alexandrium tamarense* after antibiotic treatments, Aquatic

Microbial Ecology 42 (2006) 41-53.

[48] P. Uribe, R.T. Espejo, Effect of Associated Bacteria on the Growth and Toxicity of

Alexandrium catenella, Applied Environmental Microbiology 69 (2003) 659.

[49] Y. Liu, Studies of Phycotoxin Contamination in the Coastal Waters of China and

Preparation of Toxin Reference Materials, Institute of Oceanology, Chinese Academy of

Sciences (2017) 55-158.

[50] A.A. Chariton, Book review: Environmental DNA: For Biodiversity Research and

Monitoring by Pierre Taberlet, Aurélie Bonin, Lucie Zinger and Eric Coissac, Molecular

Ecology 28 (2019) 4549-4551.

[51] E. Bolyen, J.R. Rideout, M.R. Dillon, N.A. Bokulich, C.C. Abnet, G.A. Al-Ghalith, H.

Alexander, E.J. Alm, M. Arumugam, F. Asnicar, Y. Bai, J.E. Bisanz, K. Bittinger, A.

Brejnrod, C.J. Brislawn, C.T. Brown, B.J. Callahan, A.M. Caraballo-Rodriguez, J. Chase,

E.K. Cope, R. Da Silva, C. Diener, P.C. Dorrestein, G.M. Douglas, D.M. Durall, C. Duvallet,

C.F. Edwardson, M. Ernst, M. Estaki, J. Fouquier, J.M. Gauglitz, S.M. Gibbons, D.L.

Gibson, A. Gonzalez, K. Gorlick, J. Guo, B. Hillmann, S. Holmes, H. Holste, C.

Huttenhower, G.A. Huttley, S. Janssen, A.K. Jarmusch, L.J. Jiang, B.D. Kaehler, K. Bin

Kang, C.R. Keefe, P. Keim, S.T. Kelley, D. Knights, I. Koester, T. Kosciolek, J. Kreps,

M.G.I. Langille, J. Lee, R. Ley, Y.X. Liu, E. Loftfield, C. Lozupone, M. Maher, C. Marotz,

B.D. Martin, D. McDonald, L.J. McIver, A.V. Melnik, J.L. Metcalf, S.C. Morgan, J.T. Morton,

A.T. Naimey, J.A. Navas-Molina, L.F. Nothias, S.B. Orchanian, T. Pearson, S.L. Peoples,

D. Petras, M.L. Preuss, E. Pruesse, L.B. Rasmussen, A. Rivers, M.S. Robeson, P.

Rosenthal, N. Segata, M. Shaffer, A. Shiffer, R. Sinha, S.J. Song, J.R. Spear, A.D.

Swafford, L.R. Thompson, P.J. Torres, P. Trinh, A. Tripathi, P.J. Turnbaugh, S. Ul-Hasan,

J.J.J. van der Hooft, F. Vargas, Y. Vazquez-Baeza, E. Vogtmann, M. von Hippel, W.

Walters, W. Walters, Y. Wan, M. Wang, J. Warren, K.C. Weber, C.H.D. Williamson, A.D.

Willis, Z.Z. Xu, J.R. Zaneveld, Y.L. Zhang, Q.Y. Zhu, R. Knight, J.G. Caporaso,

Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2

(vol 37, pg 852, 2019), Nature Biotechnology 37 (2019) 1091-1091.

[52] S. Zou, L. Smith, Comprehensive primer sets and cost efficient multiplex PCR-based

EDNA sequencing for community dynamics of cyanobacteria, eukaryotic phytoplankton

and zooplankton in lake, (2020).

[53] C. Fidalgo, I. Henriques, J. Rocha, M. Taco, A.J.E.S.P.R.I. Alves, Culturable

endophytic bacteria from the salt marsh plant Halimione portulacoides: phylogenetic

diversity, functional characterization, and influence of metal(loid) contamination,

Environmental Science and Pollution Research 23 (2016) 1-15.

[54] X.L. Zhang, B. Feng, Y.Q. Cai, H.J.J.A.M. Yu, Materials, Development and application

of tandem LC-MS method for the determination of paralytic shellfish poisoning toxins for

pollution monitoring, 55 (2011) 51-56.

[55] B. Langmead, S.L. Salzberg, Fast gapped-read alignment with Bowtie 2, Nature

Methods 9 (2012) 357-U354.

[56] C.N. Dewey, L. Bo, RSEM: accurate transcript quantification from RNA-Seq data with

or without a reference genome, BMC Bioinformatics 12 (2011) 323-323.

[57] W. Miller, E. Myers, D. Lipman, Blast (basic local alignment search tool), Encycl.

Genet. Genom. Proteom. Inform 215 (2008) 221.

[58] S. Audic, J.M. Claverie, The significance of digital gene expression profiles, Genome

Research 7 (1997) 986-995.

[59] K.M. Brandenburg, W. Sylke, J. Uwe, K. Anke, J. Jacqueline, K. Bernd, d.W.D.B. Van,

D. Ian, Intraspecific trait variation and trade-offs within and across populations of a toxic

dinoflagellate, Ecology Letters 21 (2018) 1561-1571.

[60] S. Zou, Comparative Transcriptome Analysis of Toxic and Non-Toxic Nassarius

Communities and Identification of Genes Involved in TTX-Adaptation, Toxins 12 (2020)

761.

[61] M.T. Croft, A.D. Lawrence, R.D. Evelyne, M.J. Warren, A.G. Smith, Algae acquire

vitamin B12 through a symbiotic relationship with bacteria, Nature 2015 (2013) 90-93.

[62] Kristy, Deiner, M. Holly, Bik, Elvira, M?chler, Mathew, Seymour, Ana?s,

Lacoursière-Roussel, Environmental DNA metabarcoding: Transforming how we survey

animal and plant communities, Molecular Ecology (2017) 5872-5895.

[63] R. Ramanan, B.-H. Kim, D.-H. Cho, H.-M. Oh, H.-S. Kim, Algae–bacteria interactions:

evolution, ecology and emerging applications, Biotechnology advances 34 (2016) 14-29.

[64] H.M. Li, H.J. Tang, X.Y. Shi, C.S. Zhang, X.L. Wang, Increased nutrient loads from

the Changjiang (Yangtze) River have led to increased Harmful Algal Blooms, Harmful

Algae 39 (2014) 92-101.

[65] C.J. Bolch, T.A. Bejoy, D.H.J.F.i.M. Green, Bacterial associates modify growth

dynamics of the dinoflagellate *Gymnodinium catenatum*, 8 (2017) 670.

[66] R. Kellmann, T.K. Mihali, Y.J. Jeon, R. Pickford, F. Pomati, B.A. Neilan, Biosynthetic

intermediate analysis and functional homology reveal a saxitoxin gene cluster in

cyanobacteria, Applied and Environmental Microbiology 74 (2008) 4044-4053.

[67] T.K. Mihali, W.W. Carmichael, B.A. Neilan, A Putative Gene Cluster from a Lyngbya

wollei Bloom that Encodes Paralytic Shellfish Toxin Biosynthesis, Plos One 6 (2011).

[68] K. Stucken, U. John, A. Cembella, A.A. Murillo, K. Soto-Liebe, J.J. Fuentes-Valdes, M.

Friedel, A.M. Plominsky, M. Vasquez, G. Glockner, The Smallest Known Genomes of

Multicellular and Toxic Cyanobacteria: Comparison, Minimal Gene Sets for Linked Traits

and the Evolutionary Implications, Plos One 5 (2010).

[69] D.Z. Wang, S.F. Zhang, Y. Zhang, L. Lin, Paralytic shellfish toxin biosynthesis in

cyanobacteria and dinoflagellates: A molecular overview, Journal of Proteomics 135 (2016)

132-140.

[70] R.V. Snyder, P.D.L. Gibbs, A. Palacios, L. Abiy, R. Dickey, J.V. Lopez, K.S. Rein,

Polyketide synthase genes from marine dinoflagellates, Marine Biotechnology 5 (2003)

1-12.

[71] R.J. Orr, A. Stüken, T. Rundberget, W. Eikrem, K.S.J.H.A. Jakobsen, Improved

phylogenetic resolution of toxic and non-toxic *Alexandrium* strains using a concatenated

rDNA approach, 10 (2011) 676-688.

[72] S.A. Amin, L.R. Hmelo, H.M. van Tol, B.P. Durham, L.T. Carlson, K.R. Heal, R.L.

Morales, C.T. Berthiaume, M.S. Parker, B. Djunaedi, A.E. Ingalls, M.R. Parsek, M.A.

Moran, E.V. Armbrust, Interaction and signalling between a cosmopolitan phytoplankton

and associated bacteria, Nature 522 (2015) 98-U253.

[73] A.A. Shibl, A. Isaac, M.A. Ochsenkuhn, A. Cárdenas, C. Fei, G. Behringer, M. Arnoux,

N. Drou, M.P. Santos, K.C. Gunsalus, Diatom modulation of select bacteria through use of

two unique secondary metabolites, Proceedings of the National Academy of Sciences 117

(2020) 27445-27455.

Figure legends

Figure 1. The community diversity of dinoflagellate in Yangtze Estuary. (A): ASV mean relative abundance; (B): Dominant dinoflagellate species.

Figure 2. (A): Maximum-likelihood phylogenetic tree of *A. catenella* from 18S. (B): The toxin content of *A. catenella* samples from three locations; (C): The associated bacteria difference of *A. catenella* samples from three locations. The points represent means ± SD of triplicate samples.

Figure 3. Associated bacteria composition of *A. catenella* from 16S and 16Sv3v4 at phylum, class, order taxonomic level.

Figure 4. Associated bacteria composition of *A. catenella* from 16S and 16Sv3v4 at family, genus, species taxonomic level.

Figure 5. (A): Maximum-likelihood phylogenetic tree of isolated associated bacteria of toxic *A. catenella* strain. (B): The influence of isolated bacteria on toxin production of *A. catenella* under normal culture conditions. (C): The influence of isolated bacteria on toxin production of *A. catenella* under nitrogen deprivation conditions. 1-13 indicates the bacteria serial number for reseeding with *A. catenella*.

Figure 6. (A): LC-MS mass spectral peaks for mixed bacterial solution. Vertical coordinate is mean relative abundance, horizontal coordinate is time. (B): LC-MS mass spectral peaks of the toxic *A. catenella* strain; (C): The molecular weight of detected GTX toxins (395.0878) from LC-MS peaks of toxic *A. catenella* strain.

Figure 7. (A): Growth of axenic, non-axenic and non-axenic-ndep *A. catenella*

strains; (B): The content of soluble protein, sugar and neutral lipid from

non-axenic and non-axenic-ndep *A. catenella* strains. The points represent means

± SD of triplicate samples. Different letters above the bars indicate significant

differences (P < 0.05) between growth condition.

Figure 8. (A): Annotation difference of six databases; (B): Transcription factors

annotated from unigenes; (C): Shared and distinct unigenes from axenic,

non-axenic and non-axenic-ndep *A. catenella* strains; (D): Up and down-regulated

genes among axenic, non-axenic and non-axenic-ndep *A. catenella* strains. For

example, non-axenic vs axenic comparison shows that non-axenic strains

generated more downregulated genes (green) than upregulated genes (orange) in

comparison with axenic strain.

Figure 9. (A): Biosynthesis of cofactors annotated between axenic and non-axenic

A. catenella strains from KEGG pathway; (B): The dominant different expression

genes (DEGs) among axenic, non-axenic and non-axenic-ndep *A. catenella*

strains.

Figure 10. The pathway network related with genetic expression of PSTs

production. Red box and circle: upregulated genes, green box and circle:

downregulated genes. For example, "non-axenic vs axenic" means the upregulated

and downregulated genes non-axenic strains produced compared with axenic

strain. The size of box and circle indicates the amount of upregulated and

downregulated genes.

Figure S1. Assignments of isolated associated bacteria at each taxonomic level.

Figure S2. The GO functional annotation performed for all unigenes.

Figure S3. The KEGG pathways assigned.

Figure S4. The main pathways enriched between non-axenic and non-axenic-ndep

A. catenella strains.

Table 1 Primers for PCR amplification

Fragment	Primer Name	Sequence (5'-3')	Reference	Fragment size (bp)
16S-v3v4	16Sv3v4F	CCTACGGGNGGCWGCAG	Klindworth et al., 2012	450-500
	16Sv3v4R	GACTACHVGGGTATCTA ATCC	Klindworth et al., 2013	
16S	16SF174F	CGGTAAGACRGRGGATG CAA	Zou et al., 2020	450
	16SR316R	CRGCTARGACTACTGGG GTA	Zou et al., 2020	
18S-v4-5	v4-387F	GYTGCAGTTAAAAMGCT CGT	Zou et al., 2020	500-520
	v4-730R	TTCAGNCTTGCGACCATA CT	Zou et al., 2020	
28S	1FC1RF27 2F	GAGACCGATAGCRMACA AGT	Zou et al., 2020	400-500
	1FC1RF59 3R	CTYGGTCCGTGTTTCWA GAC	Zou et al., 2020	

Table 2 Sequencing data statistics table

Sample	Raw reads	Clean reads	Error rate (%)	Q20 (%)	Q30 (%)	GC content (%)
non-axenic-ndep	45778990	45067136	0.0234	98.68	95.83	56.25
non-axenic	48518624	48028712	0.024	98.44	95.15	58.51
axenic	43035798	42337016	0.0245	98.25	94.76	56.55

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Figure S1

Figure S2

Figure S3

Figure S4