Microtubules ensure transport of vegetative nuclei and sperm cells by fine-tuning their home positions

Kazuki Motomura[1,2†], Haruna Tsuchi[3†], Marin Komojiri[4], Ayumi Matsumoto[5], Naoya Sugi[3], Daichi Susaki[3], Atsushi Takeda[1], Tetsu Kinoshita[3], Daisuke Maruyama[3]*

1. College of Life Sciences, Ritsumeikan University, Kusatsu, Shiga 525-8577, Japan.
2. JST, PRESTO, Kawaguchi, Saitama 332-0012, Japan.
3. Kihara Institute for Biological Research, Yokohama City University, Maioka-cho 641-12, Totsuka-ku, Yokohama, Kanagawa 244-0813, Japan.
4. Center for Ecological Research, Kyoto University, Otsu, Shiga, 520-2113, Japan.
5. Research Organization of Science and Technology, Ritsumeikan University, Kusatsu, Shiga 525-8577, Japan.

[†]These two authors contributed equally.
*To whom correspondence is addressed. E-mail: dmaru@yokohama-cu.ac.jp

Running head: Microtubules regulate male germ unit positions

Abstract

The pollen tube plays a pivotal role in double fertilization by delivering sperm cells (SCs) to the ovule. In *Arabidopsis thaliana*, a pair of SCs tightly connects with the vegetative nucleus (VN) to form the male germ unit (MGU), which is located in the apical region during pollen-tube growth, keeping the VN ahead of the SCs. MGU transport relies on independent motility of VN and SC pairs. However, the complexity of this dual motive force has hindered our understanding of MGU behavior, including its positioning and nuclear order. We used *Arabidopsis* mutants or transgenic plants that produced semi-motile MGUs with aberrant VNs or SCs to analyze the positioning of VN or SCs after stochastically disconnecting the MGU. In pollen tubes with an immotile SC pair, the VN was ~70 μm away from the tip, whereas in pollen tubes with an immotile VN, the SC pair was ~100 μm away from the tip, implying that the VN and SCs have independent home positions. The position of MGU moved forward owing to the loss of the microtubule-destabilizing kinesin *KINESIN-13A*. Conversely, microtubule depolymerization by oryzalin treatment or introducing mutations in *TUBULIN BETA 4* (*TUB4*) deregulated the position of the MGU and shifted its position backward. In addition, *tub4* plants exhibited reduced fertility. These data indicate a significant role of microtubules in stable MGU positioning to ensure reproductive success.

Introduction

The pollen tube is a tip-growing tissue that delivers the male genome to an ovule during sexual reproduction in flowering plants. In *Arabidopsis thaliana* and other plants that produce tricellular pollen, pollen tube vegetative cells contain two sperm cells (SCs), one of which has a tail-like protrusion tightly associates with the vegetative nucleus (VN) to form a reproductive cell complex known as the male germ unit (MGU). Although MGU vigorously moves back and forth, it gradually moves forward in the net movement accompanying the tip growth of the pollen tube.

In the past decade, genetic evidence of independent motive forces in VN and SC pairs has been shown in *A. thaliana*. Loss-of-function of two WPP domain-interacting tail-anchored protein 1 (WIT1) and WIT2 causes passive migration of VN dragged by the SC pair (Zhou and Meier, 2014). WIT1 and WIT2 are localized on the outer nuclear envelope and redundantly serve as components of the linker of the nucleoskeleton and cytoskeleton (LINC) complex (Tatout et al., 2014; Groves et al., 2020). The LINC complex regulates nuclear migration in plant cells by transmitting the mechanical force generated by motor proteins. Loss of other LINC components resulted in similar VN transport defects, indicating a crucial role of the LINC complex in VN motility (Zhou et al., 2015). Although the key molecules involved in SC motility have

not yet been identified, SC motility is susceptible to alterations in the extracellular

environment of the SC pairs. When the overaccumulation of callose (β-1,3-glucan) was

ectopically induced by the *pHTR10:cals3m* transgene (designated *SC-cal*), which

expresses callose synthase (*cals3m*) from the SC-specific *HTR10* promoter, SC pairs

were observed at the basal region of the pollen tube (Motomura et al., 2021). In these

aberrant pollen tubes, MGU connections tend to be lost, and VNs are often detected in the

apical region, suggesting loss of SC-specific motility in pollen tubes from *SC-cal*

transformants (Motomura et al., 2021).

Active transport of organelles requires cables and motors (Perico and Sparkes,

2018). Pollen tubes contain long actin cables traversing the shank region that are likely to

orient their plus end toward the tip, because the actin polymerization factor formin is

localized at the tip of the pollen tube (Cheung and Wu, 2004; Cheung et al., 2010). In

pollen tubes, various organelles move along actin cables using myosin XI family proteins

(Madison et al., 2015). Myosin XI-I/KAKU1 is the only myosin XI associated with the

nuclear envelope in *Arabidopsis* (Avisar et al., 2009). Myosin XI-I/KAKU1 controls the

spindle shape of the nucleus and LINC-dependent nuclear movement in vegetative

tissues (Tamura et al., 2013). However, unlike *wit1 wit2* double mutant plants, male

fertility was not reduced in *kaku1* plants (Zhou and Meier, 2014). Overall, myosin

function in apical MGU transport is unclear.

The role of microtubules in nuclear transport is unclear, probably because of

relatively moderate abnormalities in MGU transport after treatment with

microtubule-disrupting drugs (Cai et al., 1997). Immunostaining analysis of tubulins and

the GFP-tagged plus-end binding protein AtEB1 showed that pollen tubes contain

longitudinally aligned microtubules in the shank region and shorter cables in the

subapical region (Cheung et al., 2008; Cai and Cresti, 2010). Kinesins with the calponin

homology domain (KCH), which belong to the kinesin 14 subfamily expressed in pollen

tubes, are candidates for motor proteins that drive SCs, in a "tug-of-war" model (Cai et al.,

1997; Schattner et al. 2021; Cai, 2022). *In vitro* experiments have demonstrated that rice

*Os*KCH1 interacts with both microtubules and actin fibers to drive fast (average 69.17 $\pm$

14.77 nm s^{-1}) and slow (average 24.2 $\pm$ 13.42 nm s^{-1}) sliding movements (Schattner et al.,

2021; Walter et al., 2015). Because fast and slow movements appeared simultaneously in

actin fiber fragments railing the same microtubule, the two velocities were assumed to

reflect different motive forces in the parallel and antiparallel sliding of the two fibers. In

the tug-of-war model, KCH1 generates different velocities of back-and-forth movements

by associating on one side with randomly oriented microtubules surrounding the SCs, and

on the other side with uniformly directed long actin cables (Schattner et al., 2021).

Although this model provides a simple explanation based on cytological molecular data,

there is lack of genetic evidence.

Analysis of the apical MGU transport mechanism was hindered by the

behavioral interaction between VN and the SC pair, based on their independent motive

force and physical connection through a tail-like protrusion. Previously, we genetically

dissected the apical transport of the VNs and SCs (Motomura et al., 2021). *SC-cal*

transgenic (immotile SCs) and *wit1 wit2* (immotile VN) plants showed stochastic

disconnection of MGU, producing mobile "orphan" VNs and SC pairs, respectively

(Motomura et al., 2021). The MGU disconnections opened the door to separately monitor

behaviors of VNs and SC pairs, and these plants thus can be key genetic tools to analyze

molecular mechanism of the apical MGU transport. Here, we demonstrate that the orphan

SC pair in *wit1 wit2* plants was at positions distant from pollen-tube tips (~100 μm) than

orphan VNs in *SC-cal* (~70 μm), suggesting that VNs and SCs intrinsically determine

their own "home positions." Different home positions in the VN and SC pairs could

explain the stable MGU order during pollen-tube growth, because the home positions of

each orphan MGU component almost corresponded to those in wild-type MGUs. We

found that microtubules and KINESIN-13A (KIN13A) were involved in MGU

positioning. These data elucidate the basis of a stable MGU positioning system required

for double fertilization.

Results

VN and the SC pair have different home positions in the pollen tube

To examine the behavior of a VN separated from its SC pair, we analyzed *in*

vitro-germinated pollen tubes from transgenic plants carrying the *SC-cal* transgene and

pRPS5A:HISTONE H2B-tdTomato (*RHT*) nuclear marker. In this genetic background,

approximately 80% of tdTomato-labeled MGUs display disconnection of the VN–SC

linkage (Motomura et al., 2021). Apically transported orphan VNs were detected 61.4 ±

29.6 μm away from pollen-tube tip (mean ± SD, Fig. 1, A and B). We analyzed SC pairs

separated from the immotile VN *in vitro*-germinated *wit1 wit2* pollen tubes carrying *RHT*.

Similar to VN in *SC-cal* plants, orphan SC pairs were detected in the apical region (Fig. 1,

C and D, Zhou and Meier, 2014; Motomura et al., 2021). However, orphan SC pairs were

located 95.9 ± 42.6 μm away from the tip of *wit1 wit2* pollen tubes, which was much

basal than those of orphan VN in *SC-cal* plants (Fig. 1, B and D). As previously reported,

we did not observe significant differences in pollen-tube length between wild-type *RHT*,

SC-cal RHT, and *wit1 wit2 RHT* plants, suggesting that differences in the positioning of

VN and SC pair are not associated with the growth activity of pollen tubes (Supplemental

Fig. S1) (Motomura et al., 2021). Thus, the VN and SC pair intrinsically determine their

own "home positions." Interestingly, the home positions of the VN and SC pairs were

considerably close to those of intact MGUs in wild-type *RHT* plants, implying that the

presence of a tail-like connection has a limited effect, at least on the control of the home

position (Fig. 1, A to D).

Microtubule-disrupting drugs disturb positional control of VN and the SC pair

To examine the mechanisms controlling the home positions of the VN and SC

pairs, we analyzed the requirement of microtubules under semi-*in vivo* pollen-tube

growth conditions. Pollinated pistils were incubated on media with or without oryzalin (a

microtubule disrupter) and pollen tubes emerging from the cut style were observed by

microscopy. In *SC-cal RHT*, orphan VN was detected approximately 100 μm from the tip

in the control medium; however, orphan VN was broadly distributed in pollen tubes

treated with 1 μM oryzalin (Fig. 2, A and B). The positional control of orphan SC was

compromised in *wit1 wit2 RHT* (Fig. 2, C and D). These data suggest microtubules play

an important role in the positional control of the VN and SC pairs. To observe how

disturbing the home positions of VN and the SC pair affects MGU behavior, we analyzed

wild-type *RHT* pollen tubes under semi-*in vivo* growth conditions. Consistent with the

observations for the orphan VN and SC pairs, the positions of VN and SN pairs in

wild-type shifted backward and were broadly distributed in the apical region of pollen

tubes on oryzalin-containing medium (Fig. 3, A to C). In approximately 40% of the

pollen tubes, the order of the MGU was inverted after oryzalin treatment, compared with

the control, in which VN was ahead in 90% of the pollen tubes (Fig. 3D). These

observations confirmed the significance of microtubules in stabilizing the MGU structure,

including the VN-ahead order of MGU, by fine-tuning the home positions in the VN and

SC pairs.

To determine the features of microtubule-mediated positional control, we

performed time-lapse imaging of pollen tubes under semi-*in vivo* growth conditions (Fig.

3E). The movement of MGUs in the control pollen tubes was consistent with other

reports; MGUs moved forward toward the apical region as fine saltatory behavior

(Schattner et al., 2021). We named the moments when MGUs began to move forward or

backward as the front critical point (cp) or back cp, respectively, and measured the

distance from the tip to the VN at that moment. In control PTs, both the front cp and back

cp showed little variation, suggesting that MGUs accurately recognized their position and

maintained their home position by switching between forward and backward movements

(Fig. 3, E and G, control). In contrast, under oryzalin treatment, the front cp and back cp

values varied greatly, indicating that the precise control of the MGU position was lost

(Fig. 3, F and G, Oryzalin). However, there was little difference in the MGU velocity

between the control and oryzalin-treated pollen tubes (Fig. 3H). Interestingly, the

duration of backward movement was longer in oryzalin-treated pollen tubes than in

control pollen tubes (Fig. 3I). These results suggest that microtubules contribute to quick

switching between the forward and backward modes of MGUs and consequently control

the home position of the MGU.

***tub4* plants show abnormal MGU positioning and pleiotropic fertilization defects**

To obtain genetic evidence of microtubule-mediated MGU positioning, tubulin

mutants were analyzed. From a characteristic screwed plant body morphology under

microtubule-disrupting conditions, a study identified many semidominant tubulin

mutants using a forward genetic approach (Ishida et al., 2007). The mutant collection

included plants with missense mutations in four *TUBULIN ALPHA* genes (*TUA2*, *TUA3*,

TUA4, and *TUA6*) and four *TUBULIN BETA* genes (*TUB1*, *TUB2*, *TUB3*, and *TUB4*).

Due to its strong expression in mature pollen, we observed *TUB4* mutants (Loraine et al.,

2013). *RHT* was introduced into $tub4^{G96D}$, $tub4^{L250F}$, and $tub4^{S351F}$ homozygous mutants,

and MGU patterns were compared with those of wild-type *RHT* pollen tubes in *in*

vitro-germinated pollen tubes (Fig. 4). In contrast to that observed after oryzalin

treatment, we did not find aberrant MGUs whose SC pairs proceeded with the VN (Figs.

3, A and D; 4A). However, $tub4^{G96D}$, $tub4^{L250F}$, and $tub4^{S351F}$ homozygous plants showed

a broad backward shift in the positions of the VN and SC pairs compared to the wild type

(Fig. 4, B and C).

To examine the fertility of $tub4$ plants, developing siliques from $tub4^{G96D}$,

$tub4^{L250F}$, $tub4^{S351F}$ homozygous, and wild-type Columbia-0 (Col-0) plants were

observed 8 days after manual self-pollination (Supplemental Fig. S2, A and B). Under our

growth conditions, wild-type siliques produced normal seeds in approximately 80% of

the ovules. In contrast, a moderate reduction in fertility was observed in $tub4^{G96D}$ (64.7%)

and $tub4^{L250F}$ (63.0%) plants and severely low fertility (13.6%) in $tub4^{S351F}$ plants

(Supplemental Fig. 2B). To distinguish between paternal and maternal fertility defects,

we performed reciprocal crosses between wild-type and $tub4$ plants (Fig. 5). When $tub4$

plants were pollinated with wild-type pollen, all three mutants produced normal seeds

comparable to those of wild-type plants (Fig. 5, A and C). However, a significant fertility

defect was observed when the wild-type pistils were pollinated with $tub4$ pollen (Fig. 5, A

and B). The reduction in the normal seed ratio was comparable to the results obtained for

autopollination (Fig. 5, A and B; Supplemental Fig. S2B). These results demonstrate male

gametophyte-specific dysfunction in $tub4$ plants.

To evaluate reduced fertility in *tub4* plants, we examined pollen-tube growth

patterns by aniline blue staining of pistils 24 h after pollination (Supplemental Fig. S3, A

to H). In wild-type pistils pollinated with pollen from wild-type plants, the longest pollen

tubes reached the bottom of the pistils (Supplemental Fig. S3A, white arrow), and 90.1%

of the ovules displayed insertion of pollen tubes (n = 201, Supplemental Fig. S3, B and H).

By contrast, *tub4*^{S351F} pollen tubes did not reach the bottom of the pistils (Supplemental

Fig. S3A). Consequently, only 61.2% of ovules received *tub4*^{S351F} pollen tubes (n = 221,

Supplemental Fig. S3, E and H). Remarkably, one-half of the ovules in *tub4*^{S351F} pistils

remained undeveloped even after pollen-tube insertion (comparison of Supplemental Fig.

S3, E and H with Fig. 5B). The ovules that did not develop after pollen-tube insertion

were distinctly smaller than normally fertilized ovules 48 h after pollination.

(Supplemental Fig. S3, I and J, yellow double arrowheads). Failure of double fertilization

allows the ovule to attract a second pollen tube and causes multiple pollen-tube insertions,

known as polytubey (Hater et al., 2020; Sugi and Maruyama, 2023). Consistently, we

observed a slight increase in polytubey in *tub4*^{S351F} pollen tubes (Supplemental Fig. S3, F

and H). In contrast to *tub4*^{S351F}, *tub4*^{G96D} and *tub4*^{L250F} pollen tubes reached the bottom of

the pistils (Supplemental Fig. S3A). The frequency of pollen-tube insertion between

wild-type and *tub4*^{G96D} plants was comparable (84.4%, n = 212, Supplemental Fig. S3, C

and H). By contrast, *tub4*^{*L250F*} pollen tubes exhibited a slight reduction in ovule insertion

(70.5%, n = 268, Supplemental Fig. S3, D and H). In addition, *tub4*^{*L250F*} pollen tubes

displayed polytubey (4.9%) or overgrowth in the ovular micropyle (5.2%) (Supplemental

Fig. S3, F to H). Despite the pleiotropic defects, all three *tub4* mutants displayed a

common phenotype of undeveloped seeds produced following insertion of pollen tubes.

Most likely, a major fertilization defect in *tub4* plants exists during or after pollen-tube

reception.

Loss of KINESIN-13A caused a forward shift in the home position of MGUs

To examine microtubule function in MGU positioning, we analyzed

KINESIN-13 family proteins, which are unique kinesins that lack motor domains and

induce microtubule depolymerization (Oda and Fukuda, 2013). The expression of

KINESIN-13 in pollen tubes was reported in early immunostaining (Wei et al., 2005).

Among the two *KINESIN-13* (*KIN13*) genes encoded in the *Arabidopsis* genome, only

KIN13A is expressed in the pollen and pollen tubes (Qin et al., 2014). Thus, pollen-tube

growth and MGU positions were analyzed in wild-type, *kin13a*, and two

complementation lines (*kin13a*/*eYFP KIN13A*) *in vitro* using medium containing the

DNA fluorescent dye Kakshine532 (Uno et al., 2021). *kin13* pollen tubes displayed

normal MGU order and grew to the same length as wild-type pollen tubes (Fig. 6, A and

B). However, the VN of *kin13* pollen tubes was frequently detected at more tip-close

positions than that in wild-type (Fig. 6, A and C). The forward shift of VN positioning

was rescued in the two complementation lines (Fig. 6, A and C). These data suggest that

KIN13A is a microtubule depolymerizing agent that controls the tip-to-MGU distance.

Observations of tubulin reporters in pollen tubes

248   Inhibitor and mutant analyses suggested the significance of spatiotemporal

microtubule regulation in MGU positioning. We introduced translational sGFP fusion

reporter genes of *TUA1*, *TUA2*, *TUA6*, or *TUB4* into *RHT* and performed time-lapse

imaging of microtubular patterns under semi-*in vivo* growth conditions (Fig. 7). These

reporter lines exhibited a bright GFP signal in the vegetative cell of the pollen tube.

Because the GFP signal in the vegetative cell, the microtubular pattern could not be

clearly determined in the SCs of *pTUA1:sGFP-TUA1*, *pTUA2:sGFP-TUA2*, and

pTUB4:sGFP-TUB4 lines (Fig. 7). Although the *pTUA6:sGFP-TUA6* lines only showed

a strong GFP signal in the SCs, microtubular patterns were barely detectable (Fig. 7). In

contrast, several common microtubular patterns were visible in the vegetative cells of the

growing pollen tube (Fig. 7; Supplemental Videos 1 to 4). First, sGFP-labeled tubulin

formed a moderate gradient from pollen-tube tip to the basal region. Second, actively

moving short microtubules accumulated in front of the MGU, whereas longer cables were

distributed around the basal region of the MGU. Third, longer static cables were found

around the periphery of the plasma membrane. Fourth, rapid microtubule movements

were observed at the center of pollen tubes, which sometimes appeared to associate with

VN or SC pairs (Fig. 7; Supplemental Videos 1 to 4). Unfortunately, quantitative analysis

of the association between microtubules and each MGU component was hindered by

vigorous protoplasmic streaming.

Time-lapse imaging showed clear abnormalities in GFP patterns or pollen-tube

behaviors in the tubulin reporter lines. For example, the *pTUB4:sGFP-TUB4* line showed

severe pollen-tube growth retardation and backward shift of MGU, although visible

microtubules were most frequently observed in the four tubulin reporters (Fig. 7;

Supplemental Fig. S4). Similar but moderate defects were detected in the other reporter

lines. The *pTUA1:sGFP-TUA1* lines often produced granule-shaped structures, in

addition to cable patterns, indicating the formation of nonnative protein aggregates.

These observations suggest formation of artifacts due to the overexpression of

sGFP-tagged tubulin proteins. The number of visible microtubule fragments decreased

during time-lapse imaging. This alteration would be partially due to photobleaching, but

likely shows higher photosensitivity of microtubule dynamics in pollen tubes. Taken

together, the artifacts of tubulin reporter lines, higher photosensitivity, and rapid

protoplasmic streaming demonstrate challenges in analyzing the native dynamics of

microtubules in pollen tubes.

282   Discussion

Apical MGU transport is a key mechanism of pollen tube-dependent sexual

reproduction in flowering plants. In this study, the genetic dissection of MGU motility

indicated that VN and the SC pair tended to be located at different positions in the apical

region of the pollen tube, and MGU positioning was disturbed in *tub4* and *kin13a* plants.

These findings suggest independent home positions of VN and the SC pair produced

through unique saltatory movements modulated by microtubule dynamics.

290   Live imaging of tubulin in pollen tubes

According to antitubulin immunostaining studies in *Nicotiana tabacum*,

Nicotiana alata, Pyrus communis, and *A. thaliana*, pollen tubes show characteristic

microtubular patterns: the tip region contains no microtubules, the subapical region in

front of the MGU displays a distribution of shorter bundles, and the shank region around

and beneath the MGU produces longitudinally aligned cortical microtubules

(Heslop-Harrison et al., 1988; Tiwari and Polito, 1988; Del Casino et al., 1993; Joos et

al., 1994; Åström et al., 1995; Joos et al., 1995; Raudaskoski et al., 2001; Laitiainen et al.,

2002; Yu et al., 2009). In addition to fixed sample observations, microtubule movements

were reported in tobacco pollen tubes transiently expressing GFP-AtEB1a through

microprojectile bombardment (Cheung et al., 2008). During live imaging, pollen tubes

showed a dense signal of GFP-AtEB1a in the subapical region and differential

microtubular dynamics, including static cortical microtubules in the shank and rapid flux

of short microtubule bundles in the subapical region (Cheung et al., 2008). Our time-lapse

imaging of four sGFP-tagged tubulin reporters confirmed these observations (Fig. 7,

Supplemental Videos 1 to 4). However, direct GFP-tagging exhibits a moderate gradient

of tubulins from pollen-tube tip to the basal region and implies a different spatiotemporal

control of the polymerization–depolymerization equilibrium along the longitudinal axis

of the pollen tube.

Microtubule-mediated MGU transport and the tug-of-war model

A study proposed the association of microtubules with MGU using

immunostaining and electron microscopy data (Lancelle et al., 1987). Bright field

analysis of growing pollen tube revealed a dynamic association between MGU and

fibrillar structures (Heslop-Harrison and Heslop-Harrison, 1989b). Our live imaging

confirmed the close presence of microtubules around the MGU (Fig. 7; Supplemental

Videos 1 to 4). Microtubule disruption by oryzalin or colchicine treatment retarded the

apical transport of VN and generative cells in *N. tabacum*, *N. alata, N. sylvestris*, and

Galanthus nivarius (Heslop-Harrison et al., 1988; Joos et al., 1994; Joos et al., 1995;

Åström et al. 1995; Laitiainen et al., 2002). A similar broad MGU distribution phenotype

was observed in oryzalin-treated *Arabidopsis* pollen tubes (Fig. 3), indicative of apical

MGU transport regulated by microtubules. By contrast, cytochalasin D-mediated F-actin

disruption had a more deleterious effect on apical MGU transport (Heslop-Harrison and

Heslop-Harrison, 1989a, 1989b). Thus, microtubules play a subsidiary role in apical

MGU transport. Recently, Schattner et al. proposed a tug-of-war model, in which KCH

kinesins drive bimodal SC behavior, consisting of rapid forward and slow backward

movements (Schattner et al., 2021). The tug-of-war model largely owes to KCH activity,

which generates different velocities from the parallel or antiparallel sliding of F-actin and

microtubules (Walter et al., 2015), suggesting a key role of microtubular dynamics

during MGU transport. A microtubule cage, which is a longitudinally aligned

microtubule bundle in generative cells and SCs, is the primary candidate of a

KCH-associated microtubule scaffold (Schattner et al., 2021). Although microtubule

cages are resistant to oryzalin in several species (Heslop-Harrison et al., 1988; Åström et

al. 1995; Zhang et al., 1995; Laitiainen et al., 2002), *Arabidopsis* pollen tubes appeared

to lack microtubule cages, irrespective of oryzalin treatment, based on observations of

antitubulin immunostaining and microtubule reporters, including *pUBQ14:GFP-TUA6*

and other tubulin markers in this study (Fig. 7) (Yu et al., 2009; Oh et al., 2010). Electron

microscopy analysis revealed that a microtubule cage is an assembly of cortical

microtubules inside generative cells or SCs and is unable to participate in KCH-mediated

cable sliding expected in the tug-of-war model (Lancelle et al., 1987). Finally, our study

demonstrated that in *Arabidopsis*, oryzalin did not affect MGU velocity during forward or

backward movement (Fig. 3H). These data suggest amicrotubule-independent bimodal

MGU movement. Overall, the major generators of bimodal MGU movement are

probably F-actin and myosins, and not a microtubule-dependent system, as argued in

other studies (Cai and Cresti, 2010).

Role of microtubules in MGU positioning

Our time-lapse imaging recorded a longer backward traveling time of MGU

after oryzalin treatment (Fig. 3I). Microtubules support stable apical MGU positioning

presumably by modulating the timing of direction switching during basipetal MGU

movement. This MGU positioning control probably requires the homeostasis of

microtubular patterns dependent on the equilibrium between tubulin polymerization and

depolymerization. Compromised MGU positioning due to oryzalin treatment or *tub4*

mutations should be a representative phenotype of microtubule instability (Figs. 2 to 4).

Interestingly, our sGFP-tagged tubulin reporter lines showed broad MGU positioning

(Fig. 7; Supplemental Fig. S4; Supplemental Video 1). These reporter lines have higher

copies of the tubulin gene, which is expected to increase tubulin polymerization, in

contrast to oryzalin treatment. Yu et al. performed antitubulin immunostaining in

Arabidopsis pollen expressing *TUA1* from *Picea wilsonii* under a strong *LAT52* promoter.

Compared with the wild-type pollen tubes, the transgenic pollen tubes enhanced the

α tubulin signal and increased the accumulation of short microtubules in the

apical-to-subapical region (Yu et al., 2007). Most likely, tubulin overexpression extends

the subapical region occupied with short microtubules in front of the MGU, causing

MGU positioning to shift backward.

Because pollen tubes penetrate with vigorous protoplasmic streaming, they must

have a sophisticated system that maintains a unique equilibrium of microtubule

polymerization–depolymerization in each region. KIN13A can be a component of this

homeostasis. In *Arabidopsis* mesophyll protoplasts, *At*KIN13A is present in the Golgi

apparatus (Lu et al., 2005). This subcellular localization was confirmed in pollen tubes of

N. tabacum, the KIN13A ortholog (*Nt*KIN13As) of which was detected by anti-

*At*KIN13A immunostaining (Wei et al., 2005). Thus, KIN13A was speculated to be

involved in protein secretion at the pollen-tube apex (Wei et al., 2005; Cai and Cresti,

2010). Analysis of metaxylem vessel cells in *Arabidopsis* showed that KIN13A plays a

pivotal role in secondary cell wall pit formation by inducing depolymerization of cortical

microtubules necessary for sliding movement of the cellulose synthase complex (Oda and

Fukuda, 2013). KIN13A is recruited to the cortical microtubules by an active form of Rho

of plants (ROP) GTPase and MICROTUBULE DEPLETION DOMAIN1 (MIDD1)

protein. In pollen tubes, ROP is localized at the apex, where it controls polar protein

secretion required for active tip growth (Hwang and Yang, 2006). In this study, *kin13a*

pollen tubes showed a forward shift of MGU positioning (Fig. 6). It is possible that

KIN13A destroys cortical microtubules at the apical region downstream of ROP signaling,

and this local microtubule instability maintains the size of the subapical region with short

microtubule bundles. Although microtubules do not play a major role in pollen-tube

growth (Joos et al., 1994; Cai and Cresti, 2010), two microtubule-destabilizing proteins

(MICROTUBULE-ASSOCIATED　PROTEIN18　[MAP18]　and　plasma

membrane-associated　cation　binding　protein　1

[PCaP1]/MICROTUBULE-DESTABILIZING PROTEIN25 [MDP25]) have been

shown to control the directional growth of pollen tubes by severing F-actin at the

subapical region (Wang et al., 2007; Nagasaki et al., 2008; Kato et al., 2010; Li et al.,

2011; Zhu et al., 2013; Qin et al., 2014). These microtubule-destabilizing proteins may

control MGU positioning in parallel with the KIN13A pathway.

Independent control of home positions in VN and the SC pair

Upon pollen tube discharge, the pollen tube must include two SCs in a limited

volume of released cellular contents. If MGU positioning is shifted backward, the pollen

tube cannot deliver SCs and fails to accomplish double fertilization. Pollen tubes from

tub4 plants showed reduced fertility even after ovule arrival (Figs. 4 and 5). Pollen tubes

from *SC-cal* plants exhibited reduced fertility due to abnormal SC positioning

(Motomura et al., 2021). Our analysis of *tub4* plants provides additional evidence on the

crucial role of MGU positioning in double fertilization.

Genetic dissection of MGU motility using *SC-cal* and *wit1 wit2* plants revealed

independence of home positions between VN and the SC pair (Fig. 1). Due to their own

home positions, MGUs could maintain the VN-ahead order during transport (Fig. 3D).

Although the VN-ahead MGU order is common in flowering plants (Heslop-Harrison

and Heslop-Harrison, 1989c), its biological significance is unclear. SC pairs or generative

cells may receive less mechanical stress than the opposite order, protected by the

proceeding VN. Concomitantly with a robust VN–SC connection, the VN-ahead MGU

order and different home positions in MGU would ensure stable male genome delivery.

408   Compared with other pollen-tube biology studies, studies of MGU transport

have progressed slowly, probably due to the tight association of F-actin function in

pollen-tube growth and technical difficulty in dissecting bimodal movement. Our data

might change this situation by focusing on microtubule function and proposing the home

position concept. Discovery of independent home positions indicated putative molecular

switches on VN and SCs that perceive relative distance from the tip and determine the

timing to change moving directions. Candidates for switch proteins can be selected from

proteomics data in MGU. To characterize new positioning regulators, monitoring the

native microtubular pattern is necessary. Because the increased copy number of tubulin in

our reporter lines induced an artificial microtubular pattern, more authentic reporter lines

should be generated using in-frame GFP knock-in technology (Miki et al., 2018). These

analyses of the MGU positioning system will contribute to the genetic dissection of basic

bimodal MGU movement in the future.

Methods

Plant materials and growth conditions

Col-0 was used as the background for all plants. *RHT*, *SC-cal RHT*, *wit1 wit2 RHT*, and *Lyn24-mNeongreen RHT* plants have been described (Motomura et al., 2021). *tub4^{G96D}* (CS68892), *tub4^{L250F}* (CS68896), and *tub4^{S351F}* (CS68897) seeds were provided by ABRC (https://abrc.osu.edu). *kin13a* and *kin13a*/*eYFP-KIN13a* plants were gifts from Dr. Yoshihisa Oda (Nagoya University). *ms1* plants were generated using CRISPR/Cas9-mediated genome editing. Sequencing showed insertion of a G at 1,166 bp from the first base of the start codon, A, leading to a frameshift. After confirming Cas9-null status in the T2 generation, T3 *ms1* homozygous mutants were selected based on phenotype, and the pistils of these plants were used for semi-*in vitro* pollen-tube growth.

Seeds were directly sown on soil or were surface sterilized with sterilization solution (2% Plant Preservative Mixture, 0.1% Tween 20), incubated at 4°C for several days, and sown on Murashige and Skoog medium containing 1% sucrose and antibiotics. Approximately 10 d after germination, the plants were transferred to the soil and grown at 22°C under continuous light or standard long-day conditions.

Plasmids and transgenic plants

Plasmids carrying sGFP-tubulin reporter genes were constructed as follows: In each tubulin gene, the promoter sequence and protein coding sequence with 3′-UTR were independently amplified by PCR using primer sets listed in Supplemental Table S1 and Col-0 genome DNA as a template. DNA fragment encoding sGFP with the Gly-Gly-Ser-Gly-Gly-Ser linker was amplified by PCR using mNG_F and GFP_Linker_R as primer sets from the pDM310 plasmid containing

pMYB98:pCOXIV-sGFP (Maruyama et al., 2015). DNA cassettes of the promoter,

protein coding sequence, and sGFP were mixed and subjected to PCR using the forward

primer used for the amplification of the promoter and the reverse primer used for the

amplification of the protein coding sequence. These PCR products were introduced into

pDONR221 by BP reactions to produce four sGFP-tubulin entry vectors (Thermo Fisher,

MA, USA). We performed the LR reaction of the pGWB501 destination vector with

sGFP-tubulin entry vectors carrying *pTUA1:sGFP-TUA1*, *pTUA2:sGFP-TUA2*,

pTUA6:sGFP-TUA6, or *pTUB4:sGFP-TUB4*, and obtained binary vectors pDM772,

pDM693, pDM694, and pDM774, respectively (Nakagawa et al., 2007). The plasmids

were transformed into *RHT* plants using the *Agrobacterium*-mediated floral dipping

method, with *Agrobacterium* GV3101 (Zhang et al., 2006). After kanamycin selection,

T_2 homozygous plants were used for live imaging.

***In vitro* pollen-tube growth**

Mature pollen was collected and observed on pollen germination medium (PGM:

0.01% boric acid, 5 mM $CaCl_2$, 5 mM KCl, 1 mM $MgSO_4$, 10% sucrose; pH adjusted to

7.5 with 1 N KOH) and 1.5% NuSieve GTG agarose supplemented with 10 μM

epibrassinolide) (Muro et al., 2018). Pollen tubes were observed using an upright

microscope (Axio Imager. M1; Zeiss, Oberkochen, Germany). To label DNA in sperm

nuclei and VN, we used PGM containing 1 μM Kakshine532 (Uno et al., 2021). Fiji

(ImageJ image processing package) was used for image processing (Schindelin et al.,

2012).

Semi-*in vitro* pollen-tube growth and oryzalin treatment

Stigmas from *ms1* or emasculated Col-0 pistils were cut with a needle and placed

on the medium, and hand pollinated with pollen from transgenic plants. After 6 h of

incubation, pollen tubes that emerged from the cut pistils were observed using an

Axioscope 5 upright microscope (ZEISS, Oberkochen, Germany) with a CCD camera

(Axiocam 503 mono; ZEISS) or an IX73 inverted microscope (Evident, Tokyo, Japan),

equipped with a 60× objective lens, a spinning disk confocal scanning unit (Yokogawa

Electric, Musashino, Japan), and an sCMOS camera (Zyla 4.2; Andor, Belfast, Northern

Ireland). For oryzalin treatment, PGM was thawed at 65°C and cooled to 37°C before

adding 0.1% 1 mM oryzalin (final concentration 1 μM) or 0.1% ethanol as control and

spread thinly on a dish. Live imaging experiments were performed with an IX83 inverted

microscope (Evident) equipped with a confocal scanner unit (CSU-W1; Yokogawa

Electric, Musashino, Japan), an sCMOS camera (Zyla 4.2 PLUS; Oxford Instruments,

Abingdon, UK), a laser diode illuminator (LDI-7; 89 NORTH, VT, USA), and a silicone

immersion objective lens (UPLSAPO100XS2, Evident, Tokyo, Japan). Stack images

were acquired at 1-mm increments and 20-s time interval and processed with

deconvolution using cellSens Dimension and Maximum Intensity Z-projection in Fiji

(Schindelin et al., 2012).

Seed viability and reciprocal cross

For self-pollination, mutant pistils emasculated a day before were pollinated with

pollen from the same mutant. Pollen from the same mutant was pollinated with each

mutant pistil. For reciprocal cross, Col-0 or mutant pistils emasculated a day before were

pollinated with pollen from Col-0 or mutant anthers. After 8 days, siliques were dissected

and seed viability was determined. Images were captured using MVX10 (Evident, Tokyo,

Japan) under bright field.

Aniline blue staining

Pollen was pollinated on Col-0 pistils emasculated a day before. After 24 or 48 h

of pollination, pistils were fixed with a fixative (ethanol: acetic acid = 9:1) for 3 h.

Fixative was replaced with 1 N sodium hydroxide solution by standing overnight.

Samples were immersed in aniline blue staining solution (0.1% aniline blue, 0.1 M

K_3PO_4) for 10 min. Images were acquired using an IX73 inverted microscope (Evident).

Fluorescence signals were detected with a CFP filter using a mercury vapor lamp with a

10× objective lens to observe pollen-tube growth and excited with a 405 nm laser with a

20× or 40× objective lens to observe pollen-tube attraction to the ovule.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial

or financial relationships that could be understood as a potential conflict of interest.

Author Contributions

KM and DM designed the study, conducted the experiments, analyzed pollen-tube

growth, and drafted the manuscript. HT performed all analyses of tubulin mutants under

the support of NS and DS. MK performed the experiments using oryzalin. AM

performed various quantitative analyses. AT constructed a plasmid for genome editing.

TK provided critical advice and reviewed the manuscript. All authors have contributed

to the manuscript and approved the final version.

Acknowledgements

We thank K. Tamura and I. Mayer for providing the *wit1 wit2* mutant; D. Kurihara for providing *pRPS5A:H2BtdTomato*; Y. Oda for sharing *kin13a* and *kin13a/eYFP-KIN13a* seeds; Y. Sato for providing Kakshine532; A. Saito, Y. Kibayashi, S. Yang, and H. Kakizaki for their technical support. We would like to thank enago for English editing.

Funding

This work was supported by JSPS KAKENHI (Grant nos. JP20H05781, JP20H05422, and JP20H05778 awarded to DM; and Grant nos. JP20K15822, JP22K15147, and 23H04751 awarded to KM); JST PRESTO (Grant no. JPMJPR20D9 awarded to KM); JST FOREST (Grant no. JPMJFR2253 awarded to KM); Yokohama City University (Academic Research Grant to DM; Development Fund to DM; and Strategic Research Promotion Grant no. SK1903 to DM); Japan Science Society (Sasakawa Scientific Research Grant to KM); Takeda Science Foundation (research Grants to KM and DM); Nikki Saneyoshi Foundation (research Grant to KM); and Ritsumeikan Global Innovation Research Organization, Ritsumeikan University (Third-Phase R-GIRO Research Grant to KM and AT).

References

Åström H, Sorri O, Raudaskoski M (1995) Role of microtubules in the movement of the vegetative nucleus and generative cell in tobacco pollen tubes. Sex Plant Reprod 8: 61–69. https://doi.org/10.1007/BF00230890

Avisar D, Abu-Abied M, Belausov E, Sadot E, Hawes C, Sparkes IA (2009) A comparative study of the involvement of 17 Arabidopsis myosin family members on the motility of Golgi and other organelles. Plant Physiol 150: 700–709.

https://doi.org/10.1104/pp.109.136853

Cai G (2022) The legacy of kinesins in the pollen tube 30 years later. Cytoskeleton

(Hoboken) 79: 8–19. https://doi.org/10.1002/cm.21713

Cai G, Cresti M (2010) Microtubule motors and pollen tube growth--still an open

question. Protoplasma 247: 131–143. https://doi.org/10.1007/s00709-010-0214-9

Cai G, Moscatelli A, Cresti M (1997) Cytoskeletal organization and pollen tube growth.

Trends Plant Sci 2: 86–91. https://doi.org/10.1016/S1360-1385(96)10057-1

Cheung AY, Duan QH, Costa SS, De Graaf BH, Di Stilio VS, Feijo J, Wu HM (2008)

The dynamic pollen tube cytoskeleton: live cell studies using actin-binding and

microtubule-binding   reporter   proteins.   Mol   Plant   1:   686–702.

https://doi.org/10.1093/mp/ssn026

Cheung AY, Niroomand S, Zou Y, Wu HM (2010) A transmembrane formin nucleates

subapical actin assembly and controls tip-focused growth in pollen tubes. Proc Natl

Acad Sci U S A 107: 16390–16395. https://doi.org/10.1073/pnas.1008527107

Cheung AY, Wu HM (2004) Overexpression of an *Arabidopsis* formin stimulates

supernumerary actin cable formation from pollen tube cell membrane. Plant Cell 16:

257–269. https://doi.org/10.1105/tpc.016550

Del Casino C, Li YQ, Moscatelli A, Scali M, Tiezzi A, Cresti M (1993) Distribution of

microtubules during the growth of tobacco pollen tubes. Biol Cell 79: 125–132.

https://doi.org/10.1111/j.1768-322X.1993.tb00902.x

Goto C, Tamura K, Nishimaki S, Maruyama D, Hara-Nishimura I (2020) The nuclear

envelope protein KAKU4 determines the migration order of the vegetative nucleus

and   sperm   cells   in   pollen   tubes.   J   Exp   Bot   71:   6273–6281.

https://doi.org/10.1093/jxb/eraa367

Groves NR, Biel A, Moser M, Mendes T, Amstutz K, Meier I (2020) Recent advances in

understanding the biological roles of the plant nuclear envelope. Nucleus 11: 330–

346. https://doi.org/10.1080/19491034.2020.1846836

Hamamura Y, Saito C, Awai C, Kurihara D, Miyawaki A, Nakagawa T, Kanaoka MM,

Sasaki N, Nakano A, Berger F, Higashiyama T (2011) Live-cell imaging reveals the

dynamics of two sperm cells during double fertilization in *Arabidopsis thaliana*. Curr

Biol 21: 497–502. https://doi.org/10.1016/j.cub.2011.02.013

Hater F, Nakel T, Groß-Hardt R (2020) Reproductive multitasking: The female

gametophyte.   Annu   Rev   Plant   Biol   71:   517–546.

https://doi.org/10.1146/annurev-arplant-081519-035943

Heslop-Harrison J, Heslop-Harrison Y (1989a) Cytochalasin effects on structure and

movement   in   the   pollen   tube   of   Iris.   Sex   Plant   Reprod   2   :   27–37.

https://doi.org/10.1007/BF00190116

Heslop-Harrison J, Heslop-Harrison Y (1989b) Conformation and movement of the

vegetative   nucleus   of   the   angiosperm   pollen   tube: Association   with   the   actin

cytoskeleton. J Cell Sci 93: 299–308. https://doi.org/10.1242/jcs.93.2.299

Heslop-Harrison J, Heslop-Harrison Y (1989c) Actomyosin and movement in the

angiosperm pollen tube: An interpretation of some recent results. Sex Plant Reprod 2:

199–207. https://doi.org/10.1007/BF00195579

Heslop-Harrison J, Heslop-Harrison Y, Cresti M, Tiezzi A, Moscatelli A (1988)

Cytoskeletal elements, cell shaping and movement in the angiosperm pollen tube. J

Cell Sci 91: 49–60. https://doi.org/10.1242/jcs.91.1.49

Hwang J-U, Yang Z (2006) Small GTPases and spatiotemporal regulation of pollen tube

growth. In: R Malhó (Ed.), The pollen tube: A cellular and molecular perspective.

Springer, Berlin, Heidelberg 3: 95–116. https://doi.org/10.1007/7089_045

Ishida T, Kaneko Y, Iwano M, Hashimoto T (2007) Helical microtubule arrays in a

collection of twisting tubulin mutants of *Arabidopsis thaliana*. Proc Natl Acad Sci U

S A 104: 8544–8549. https://doi.org/10.1073/pnas.0701224104

Joos, Ute (1994) Van Aken, J, Kristen, U. Microtubules are involved in maintaining the

cellular polarity in pollen tubes of *Nicotiana sylvestris*. Protoplasma 179: 5–15.

https://doi.org/10.1007/s10577-014-9419-7

Joos U, Van Aken J, Kristen U (1995) The anti-microtubule drug carbetamide stops

Nicotiana sylvestris pollen tube growth in the style. Protoplasma 187: 182–191.

https://doi.org/10.1007/BF01280247

Kato M, Nagasaki-Takeuchi N, Ide Y, Maeshima M (2010) An Arabidopsis hydrophilic

Ca2(+) -binding protein with a PEVK-rich domain, PCaP2, is associated with the

plasma membrane and interacts with calmodulin and phosphatidylinositol phosphates.

Plant Cell Physiol 51: 366–379. https://doi.org/10.1093/pcp/pcq003

Laitiainen E, Nieminen K, Vihinen H, Raudaskoski M (2002) Movement of generative

cell and vegetative nucleus in tobacco pollen tubes is dependent on microtubule

cytoskeleton but independent of the synthesis of callose plugs. Sex Plant Reprod 15:

195–204. https://doi.org/10.1007/s00497-002-0155-3

Lancelle SA, Cresti M, Hepler PK (1987) Ultrastructure of the cytoskeleton in

freeze-substituted pollen tubes of *Nicotiana alata*. Protoplasma 140: 141–150.

https://doi.org/10.1007/BF01273723

Li J, Wang X, Qin T, Zhang Y, Liu X, Sun J, Zhou Y, Zhu L, Zhang Z, Yuan M, Mao T

(2011) MDP25, a novel calcium regulatory protein, mediates hypocotyl cell

elongation by destabilizing cortical microtubules in *Arabidopsis*. Plant Cell 23:

4411–4427. https://doi.org/10.1105/tpc.111.092684

Loraine AE, McCormick S, Estrada A, Patel K, Qin P (2013) RNA-seq of Arabidopsis

pollen uncovers novel transcription and alternative splicing. Plant Physiol 162: 1092–

1109. https://doi.org/10.1104/pp.112.211441

Lu L, Lee YRJ, Pan R, Maloof JN, Liu B (2005) An internal motor kinesin is associated

with the Golgi apparatus and plays a role in trichome morphogenesis in *Arabidopsis*.

Mol Biol Cell 16: 811–823. https://doi.org/10.1091/mbc.e04-05-0400

Madison SL, Buchanan ML, Glass JD, McClain TF, Park E, Nebenführ A (2015) Class

XI myosins move specific organelles in pollen tubes and are required for normal

fertility and pollen tube growth in *Arabidopsis*. Plant Physiol 169: 1946–1960.

https://doi.org/10.1104/pp.15.01161

Maruyama D, Völz R, Takeuchi H, Mori T, Igawa T, Kurihara D, Kawashima T, Ueda

M, Ito M, Umeda M, Nishikawa S, Groß-Hardt R, Higashiyama T (2015) Rapid

elimination of the persistent synergid through a cell fusion mechanism. Cell 161:

907–918. https://doi.org/10.1016/j.cell.2015.03.018

Miki D, Zhang W, Zeng W, Feng Z, Zhu JK (2018) CRISPR/Cas9-mediated gene

targeting in Arabidopsis using sequential transformation. Nat Commun 9: 1967.

https://doi.org/10.1038/s41467-018-04416-0

Motomura K, Takeuchi H, Notaguchi M, Tsuchi H, Takeda A, Kinoshita T,

Higashiyama T, Maruyama D (2021) Persistent directional growth capability in

Arabidopsis thaliana pollen tubes after nuclear elimination from the apex. Nat

Commun 12: 2331. https://doi.org/10.1038/s41467-021-22661-8

Muro K, Matsuura-Tokita K, Tsukamoto R, Kanaoka MM, Ebine K, Higashiyama T,

Nakano A, Ueda T (2018) ANTH domain-containing proteins are required for the

pollen tube plasma membrane integrity via recycling ANXUR kinases. Commun Biol 1: 152. https://doi.org/10.1038/s42003-018-0158-8

Nagasaki N, Tomioka R, Maeshima M (2008) A hydrophilic cation-binding protein of *Arabidopsis thaliana*, AtPCaP1, is localized to plasma membrane via N-myristoylation and interacts with calmodulin and the phosphatidylinositol phosphates PtdIns(3,4,5)P(3) and PtdIns(3,5)P(2). FEBS Journal 275: 2267–2282. https://doi.org/10.1111/j.1742-4658.2008.06379.x

Nakagawa T, Suzuki T, Murata S, Nakamura S, Hino T, Maeo K, Tabata R, Kawai T, Tanaka K, Niwa Y, Watanabe Y, Nakamura K, Kimura T, Ishiguro S (2007) Improved Gateway binary vectors: high-performance vectors for creation of fusion constructs in transgenic analysis of plants. Biosci Biotechnol Biochem 71: 2095–2100. https://doi.org/10.1271/bbb.70216

Oda Y, Fukuda H (2013) Rho of plant GTPase signaling regulates the behavior of Arabidopsis kinesin-13A to establish secondary cell wall patterns. Plant Cell 25: 4439–4450. https://doi.org/10.1105/tpc.113.117853

Oh SA, Allen T, Twell D (2010) A ticket for the live show: Microtubules in male gametophyte development. Plant Signal Behav 5: 614–617. https://doi.org/10.4161/psb.11505

Perico C, Sparkes I (2018) Plant organelle dynamics: Cytoskeletal control and membrane contact sites. New Phytol 220: 381–394. https://doi.org/10.1111/nph.15365

Qin T, Liu X, Li J, Sun J, Song L, Mao T (2014) Arabidopsis microtubule-destabilizing protein 25 functions in pollen tube growth by severing actin filaments. Plant Cell 26: 325–339. https://doi.org/10.1105/tpc.113.119768

Qin Y, Leydon AR, Manziello A, Pandey R, Mount D, Denic S, Vasic B, Johnson MA,

Palanivelu R (2009) Penetration of the stigma and style elicits a novel transcriptome

in pollen tubes, pointing to genes critical for growth in a pistil. PLOS Genet 5:

e1000621. https://doi.org/10.1371/journal.pgen.1000621

Raudaskoski M, Åström H, Laitiainen E (2001) Pollen tube cytoskeleton: structure and

function. J Plant Growth Regul 20: 113–130. https://doi.org/10.1007/s003440010015

Schattner S, Schattner J, Munder F, Höppe E, Walter WJ (2020) A tug-of-war model

explains the saltatory sperm cell movement in *Arabidopsis thaliana* pollen tubes by

kinesins with calponin homology domain. Front Plant Sci 11: 601282.

https://doi.org/10.3389/fpls.2020.601282

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,

Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K,

Tomancak P, Cardona A (2012) Fiji: an open-source platform for biological-image

analysis. Nat Methods 9: 676–682. https://doi.org/10.1038/nmeth.2019

Sugi N, Maruyama D (2023) Exploring novel polytubey reproduction pathways

utilizing cumulative genetic tools. Plant Cell Physiol 64: 454–460.

https://doi.org/10.1093/pcp/pcad021

Tamura K, Iwabuchi K, Fukao Y, Kondo M, Okamoto K, Ueda H, Nishimura M,

Hara-Nishimura I (2013) Myosin XI-I links the nuclear membrane to the

cytoskeleton to control nuclear movement and shape in Arabidopsis. Curr Biol 23:

1776–1781. https://doi.org/10.1016/j.cub.2013.07.035

Tatout C, Evans DE, Vanrobays E, Probst AV, Graumann K (2014) The plant LINC

complex at the nuclear envelope. Chromosome Res 22: 241–252.

https://doi.org/10.1007/s10577-014-9419-7

Tiwari SC, Polito VS (1988) Organization of the cytoskeleton in pollen tubes of Pyrus

communis: A study employing conventional and freeze-substitution electron

microscopy, immunofluorescence, and rhodamine-phalloidin. Protoplasma 147: 100–

112. https://doi.org/10.1007/BF01403337

Uno K, Sugimoto N, Sato Y (2021) N-aryl pyrido cyanine derivatives are nuclear and

organelle DNA markers for two-photon and super-resolution imaging. Nat Commun

12: 2650. https://doi.org/10.1038/s41467-021-23019-w

Vogler F, Konrad SS, Sprunck S (2015) Knockin' on pollen's door: Live cell imaging of

early polarization events in germinating Arabidopsis pollen. Front Plant Sci 6: 246.

https://doi.org/10.3389/fpls.2015.00246

Walter WJ, Machens I, Rafieian F, Diez S   (2015) The non-processive rice kinesin-14

OsKCH1 transports actin filaments along microtubules with two distinct velocities.

Nat Plants 1: 15111. https://doi.org/10.1038/nplants.2015.111

Wang X, Zhu L, Liu B, Wang C, Jin L, Zhao Q, Yuan M (2007) Arabidopsis

microtubule-associated protein18 functions in directional cell growth by destabilizing

cortical microtubules. Plant Cell 19: 877–889. https://doi.org/10.1105/tpc.106.048579

Wei L, Liu B, Li Y (2005) Distribution of a kinesin-related protein on Golgi apparatus

of tobacco pollen tubes. Chin Sci Bull 50: 2175–2181.

https://doi.org/10.1007/BF03182668

Yu Y, Li Y, Li L, Lin J, Zheng C, Zhang L (2009) Overexpression of PwTUA1, a

pollen-specific tubulin gene, increases pollen tube elongation by altering the

distribution of alpha-tubulin and promoting vesicle transport. J Exp Bot 60: 2737–

2749. https://doi.org/10.1093/jxb/erp143

Zhang HQ, Bohdanowicz J, Pierson ES, Li YQ, Tiezzi A, Cresti M (1995) Microtubular

organization during asymmetrical division of the generative cell in *Gagea lutea*. J

Plant Res 108: 269–276. https://doi.org/10.1007/BF02344352

Zhang X, Henriques R, Lin SS, Niu QW, Chua NH (2006) Agrobacterium-mediated

transformation of Arabidopsis thaliana using the floral dip method. Nat Protoc 1:

641–646. https://doi.org/10.1038/nprot.2006.97

Zhou X, Groves NR, Meier I (2015) SUN anchors pollen WIP-WIT complexes at the

vegetative nuclear envelope and is necessary for pollen tube targeting and fertility. J

Exp Bot 66: 7299–7307. https://doi.org/10.1093/jxb/erv425

Zhou X, Meier I (2014) Efficient plant male fertility depends on vegetative nuclear

movement mediated by two families of plant outer nuclear membrane proteins. Proc

Natl Acad Sci U S A 111: 11900–11905. https://doi.org/10.1073/pnas.1323104111

Zhu L, Zhang Y, Kang E, Xu Q, Wang M, Rui Y, Liu B, Yuan M, Fu Y (2013) MAP18

regulates the direction of pollen tube growth in Arabidopsis by modulating F-actin

organization. Plant Cell 25: 851–867. https://doi.org/10.1105/tpc.113.110528

Figure 1. The vegetative nucleus and sperm cell pair have different home positions in the pollen tube.
A) Representative images of *in vitro*-germinated pollen tubes from wild-type and *SC-cal* transgenic plants carrying the *pRPS5A:HISTONE H2B-tdTomato* (*RHT*) nuclear marker. **B)** Distance from pollen tube-tip to the vegetative nucleus (Tip-VN) analyzed in (**A**). **C)** Representative images of *in vitro*-germinated pollen tubes from wild-type and *wit1 wit2* plants carrying *RHT*. **D)** Distance from pollen tube-tip to sperm nuclei (Tip-SN) analyzed in (**C**). In (**A**) and (**C**), red fluorescent signals were observed with transmitting light 3 h after pollination. Box-and-whisker plots show median (center line), upper and lower quartiles (box), maximum and minimum (whiskers), and points (solid circles). Asterisks, significance by Welch's *t* test (***$P = 8.4 \times 10^{-6}$). Scale bars, 50 μm.

Figure 2. Microtubule-disrupting drug disturbs home positions of the vegetative nucleus and sperm cell pair.
A) Representative images of an orphan vegetative nucleus under semi-*in vitro*-germinated pollen tubes from the *SC-cal pRPS5A:HISTONE H2B-tdTomato* (*RHT*) plant growing under control or oryzalin-treated conditions. **B)** Distance from pollen tube-tip to vegetative nucleus (Tip-VN) analyzed in (**A**). **C)** Representative images of orphan sperm nuclei from semi-*in vitro*-germinated pollen tubes from the *wit1 wit2 RHT* plant growing under control or oryzalin-treated conditions. **D)** Distance from pollen tube-tip to sperm nuclei (Tip-SN) analyzed in (**C**). In (**A**) and (**C**), red fluorescence signals were observed 6 h after pollination. Box-and-whisker plots show median (center line), upper and lower quartiles (box), maximum and minimum (whiskers), and points (solid circles). Asterisks, significance by Welch's *t* test (***$P = 2.97 \times 10^{-10}$, *$P = 0.014$). Scale bars, 50 μm.

Figure 3. Disruption of male germ unit (MGU) home positions in pollen tubes with disrupted microtubules.

A–D) Pollen tubes from wild-type (WT) *pRPS5A:HISTONE H2B-tdTomato* (*RHT*) nuclear marker were grown under semi-*in vivo* conditions using control medium or microtubule-unstable medium containing 1 µM oryzalin, and nuclear positions were analyzed 6 h after pollination. Representative images of red fluorescence merged with bright field (**A**), distance from pollen-tube tip to the vegetative nucleus (Tip-VN) (**B**), distance from pollen tube-tip to sperm nuclei (Tip-SN) (**C**), and order of MGU (**D**). **E, F)** Kymographs of MGU movement in pollen tubes on control medium (**E**) or oryzalin-containing medium (**F**). Pollen tubes from WT *RHT pACA3:Lyn24-mNeonGreen* plants were grown under semi-*in vivo* conditions, and time-lapse images were captured every 30 s, 6 h after pollination. **G)** Distance from pollen-tube tip to Tip-VN was measured at the front critical point (cp) or back cp analyzed in (**E**) and (**F**). **H, I)** Velocity (**H**) or travel time (**I**) of MGU migration during forward (front) or backward (back) movement analyzed in (**E**) and (**F**). Box-and-whisker plots show median (center line), upper and lower quartiles (box), maximum and minimum (whiskers), and points (solid circles). Asterisks, significance by Welch's *t* test (***$P < 0.001$, *$P = 0.023$). Scale bars, 50 µm.

Figure 4. Abnormal male germ unit (MGU) positions in *tub4* plants.
A) Representative images of *in vitro*-germinated pollen tubes from wild-type (WT) and *tub4* plants carrying the *pRPS5A:HISTONE H2B-tdTomato* (*RHT*) nuclear marker. **B)** Distance from the pollen tube-tip to vegetative nucleus (Tip-VN) analyzed in (**A**). **C)** Distance from the pollen tube-tip to sperm nuclei (Tip-SN) analyzed in (**A**). Red fluorescent signals were observed with transmitting light 6 h after pollination. Box-and-whisker plots show median (center line), upper and lower quartiles (box), maximum and minimum (whiskers), and points (solid circles). Asterisks indicate statistically significant differences detected by Dunnett's test compared with the WT (***$P < 0.001$). Scale bars, 50 μm.

Figure 5. Seed abortion caused by *tub4* pollen tubes.
A) Representative images of seeds 8 d after wild-type (WT) pistils were pollinated with pollen from WT, *tub4^{G96D}*, *tub4^{L250F}*, and *tub4^{S351F}* plants. **B, C)** Percentages of normal seeds in reciprocal crosses between WT and *tub4* plants. Cross-pollinations using *tub4* pollen donors (**B**). Cross-pollinations using *tub4* plants as female parents (**C**). Scale bar, 1 mm. Arrowheads indicate unfertilized ovules or undeveloped seeds. Asterisks indicate statistically significant differences detected by Dunnett's test in comparison with the WT (***$P < 0.001$).

Figure 6. Kinesin-13A modulates male germ unit (MGU) position.
A) Representative images of pollen tubes from wild-type (WT), *kin13a*, *kin13a eYFP-KIN13A* plants. Pollen tubes were *in vitro*-germinated on growth medium containing 1 μM Kakshine532 fluorescent DNA dye and analyzed after 2.5 h incubation. Scale bar, 50 μm. **B, C)** Pollen tube length (**B**) and distance from pollen tube-tip to vegetative nucleus (Tip-VN) (**C**) analyzed in (**A**). Box-and-whisker plots show median (center line), upper and lower quartiles (box), maximum and minimum (whiskers), and points (solid circles). Different alphabets indicate statistically significant differences detected by Tukey's test.

Figure 7. Microtubule dynamics during male germ unit (MGU) transport.
A–D) Microtubules (gray) and nuclei (magenta) in pollen tubes from *pRPS5A:HISTONE H2B-tdTomato* (*RHT*) *pTUB4:sGFP-TUB4* (**A**), *pTUA1:sGFP-TUA1* (**B**), *pTUA2:sGFP-TUA2* (**C**), or *pTUA6:sGFP-TUA6* (**D**) plants. Pollen tubes were grown under semi-*in vivo* conditions at 6 h after pollination. Scale bar, 10 μm.

Parsed Citations

Åström H, Sorri O, Raudaskoski M (1995) Role of microtubules in the movement of the vegetative nucleus and generative cell in tobacco pollen tubes. Sex Plant Reprod 8: 61–69. https://doi.org/10.1007/BF00230890
Google Scholar: Author Only Title Only Author and Title

Avisar D, Abu-Abied M, Belausov E, Sadot E, Hawes C, Sparkes IA (2009) A comparative study of the involvement of 17 Arabidopsis myosin family members on the motility of Golgi and other organelles. Plant Physiol 150: 700–709. https://doi.org/10.1104/pp.109.136853
Google Scholar: Author Only Title Only Author and Title

Cai G (2022) The legacy of kinesins in the pollen tube 30 years later. Cytoskeleton (Hoboken) 79: 8–19. https://doi.org/10.1002/cm.21713
Google Scholar: Author Only Title Only Author and Title

Cai G, Cresti M (2010) Microtubule motors and pollen tube growth--still an open question. Protoplasma 247: 131–143. https://doi.org/10.1007/s00709-010-0214-9
Google Scholar: Author Only Title Only Author and Title

Cai G, Moscatelli A, Cresti M (1997) Cytoskeletal organization and pollen tube growth. Trends Plant Sci 2: 86–91. https://doi.org/10.1016/S1360-1385(96)10057-1
Google Scholar: Author Only Title Only Author and Title

Cheung AY, Duan QH, Costa SS, De Graaf BH, Di Stilio VS, Feijo J, Wu HM (2008) The dynamic pollen tube cytoskeleton: live cell studies using actin-binding and microtubule-binding reporter proteins. Mol Plant 1: 686–702. https://doi.org/10.1093/mp/ssn026
Google Scholar: Author Only Title Only Author and Title

Cheung AY, Niroomand S, Zou Y, Wu HM (2010) A transmembrane formin nucleates subapical actin assembly and controls tip-focused growth in pollen tubes. Proc Natl Acad Sci U S A 107: 16390–16395. https://doi.org/10.1073/pnas.1008527107
Google Scholar: Author Only Title Only Author and Title

Cheung AY, Wu HM (2004) Overexpression of an Arabidopsis formin stimulates supernumerary actin cable formation from pollen tube cell membrane. Plant Cell 16: 257–269. https://doi.org/10.1105/tpc.016550
Google Scholar: Author Only Title Only Author and Title

Del Casino C, Li YQ, Moscatelli A, Scali M, Tiezzi A, Cresti M (1993) Distribution of microtubules during the growth of tobacco pollen tubes. Biol Cell 79: 125–132. https://doi.org/10.1111/j.1768-322X.1993.tb00902.x
Google Scholar: Author Only Title Only Author and Title

Goto C, Tamura K, Nishimaki S, Maruyama D, Hara-Nishimura I (2020) The nuclear envelope protein KAKU4 determines the migration order of the vegetative nucleus and sperm cells in pollen tubes. J Exp Bot 71: 6273–6281. https://doi.org/10.1093/jxb/eraa367
Google Scholar: Author Only Title Only Author and Title

Groves NR, Biel A, Moser M, Mendes T, Amstutz K, Meier I (2020) Recent advances in understanding the biological roles of the plant nuclear envelope. Nucleus 11: 330–346. https://doi.org/10.1080/19491034.2020.1846836
Google Scholar: Author Only Title Only Author and Title

Hamamura Y, Saito C, Awai C, Kurihara D, Miyawaki A, Nakagawa T, Kanaoka MM, Sasaki N, Nakano A, Berger F, Higashiyama T (2011) Live-cell imaging reveals the dynamics of two sperm cells during double fertilization in Arabidopsis thaliana. Curr Biol 21: 497–502. https://doi.org/10.1016/j.cub.2011.02.013
Google Scholar: Author Only Title Only Author and Title

Hater F, Nakel T, Groß-Hardt R (2020) Reproductive multitasking: The female gametophyte. Annu Rev Plant Biol 71: 517–546. https://doi.org/10.1146/annurev-arplant-081519-035943
Google Scholar: Author Only Title Only Author and Title

Heslop-Harrison J, Heslop-Harrison Y (1989a) Cytochalasin effects on structure and movement in the pollen tube of Iris. Sex Plant Reprod 2 : 27–37. https://doi.org/10.1007/BF00190116
Google Scholar: Author Only Title Only Author and Title

Heslop-Harrison J, Heslop-Harrison Y (1989b) Conformation and movement of the vegetative nucleus of the angiosperm pollen tube: Association with the actin cytoskeleton. J Cell Sci 93: 299–308. https://doi.org/10.1242/jcs.93.2.299
Google Scholar: Author Only Title Only Author and Title

Heslop-Harrison J, Heslop-Harrison Y (1989c) Actomyosin and movement in the angiosperm pollen tube: An interpretation of some recent results. Sex Plant Reprod 2: 199–207. https://doi.org/10.1007/BF00195579
Google Scholar: Author Only Title Only Author and Title

Heslop-Harrison J, Heslop-Harrison Y, Cresti M, Tiezzi A, Moscatelli A (1988) Cytoskeletal elements, cell shaping and movement

in the angiosperm pollen tube. J Cell Sci 91: 49–60. https://doi.org/10.1242/jcs.91.1.49
Google Scholar: Author Only Title Only Author and Title

Hwang J-U, Yang Z (2006) Small GTPases and spatiotemporal regulation of pollen tube growth. In: R Malhó (Ed.), The pollen tube: A cellular and molecular perspective. Springer, Berlin, Heidelberg 3: 95–116. https://doi.org/10.1007/7089_045
Google Scholar: Author Only Title Only Author and Title

Ishida T, Kaneko Y, Iwano M, Hashimoto T (2007) Helical microtubule arrays in a collection of twisting tubulin mutants of Arabidopsis thaliana. Proc Natl Acad Sci U S A 104: 8544–8549. https://doi.org/10.1073/pnas.0701224104
Google Scholar: Author Only Title Only Author and Title

Joos, Ute (1994) Van Aken, J, Kristen, U. Microtubules are involved in maintaining the cellular polarity in pollen tubes of Nicotiana sylvestris. Protoplasma 179: 5–15. https://doi.org/10.1007/s10577-014-9419-7
Google Scholar: Author Only Title Only Author and Title

Joos U, Van Aken J, Kristen U (1995) The anti-microtubule drug carbetamide stops Nicotiana sylvestris pollen tube growth in the style. Protoplasma 187: 182–191. https://doi.org/10.1007/BF01280247
Google Scholar: Author Only Title Only Author and Title

Kato M, Nagasaki-Takeuchi N, Ide Y, Maeshima M (2010) An Arabidopsis hydrophilic Ca2(+) -binding protein with a PEVK-rich domain, PCaP2, is associated with the plasma membrane and interacts with calmodulin and phosphatidylinositol phosphates. Plant Cell Physiol 51: 366–379. https://doi.org/10.1093/pcp/pcq003
Google Scholar: Author Only Title Only Author and Title

Laitiainen E, Nieminen K, Vihinen H, Raudaskoski M (2002) Movement of generative cell and vegetative nucleus in tobacco pollen tubes is dependent on microtubule cytoskeleton but independent of the synthesis of callose plugs. Sex Plant Reprod 15: 195–204. https://doi.org/10.1007/s00497-002-0155-3
Google Scholar: Author Only Title Only Author and Title

Lancelle SA, Cresti M, Hepler PK (1987) Ultrastructure of the cytoskeleton in freeze-substituted pollen tubes of Nicotiana alata. Protoplasma 140: 141–150. https://doi.org/10.1007/BF01273723
Google Scholar: Author Only Title Only Author and Title

Li J, Wang X, Qin T, Zhang Y, Liu X, Sun J, Zhou Y, Zhu L, Zhang Z, Yuan M, Mao T (2011) MDP25, a novel calcium regulatory protein, mediates hypocotyl cell elongation by destabilizing cortical microtubules in Arabidopsis. Plant Cell 23: 4411–4427. https://doi.org/10.1105/tpc.111.092684
Google Scholar: Author Only Title Only Author and Title

Loraine AE, McCormick S, Estrada A, Patel K, Qin P (2013) RNA-seq of Arabidopsis pollen uncovers novel transcription and alternative splicing. Plant Physiol 162: 1092–1109. https://doi.org/10.1104/pp.112.211441
Google Scholar: Author Only Title Only Author and Title

Lu L, Lee YRJ, Pan R, Maloof JN, Liu B (2005) An internal motor kinesin is associated with the Golgi apparatus and plays a role in trichome morphogenesis in Arabidopsis. Mol Biol Cell 16: 811–823. https://doi.org/10.1091/mbc.e04-05-0400
Google Scholar: Author Only Title Only Author and Title

Madison SL, Buchanan ML, Glass JD, McClain TF, Park E, Nebenführ A (2015) Class XI myosins move specific organelles in pollen tubes and are required for normal fertility and pollen tube growth in Arabidopsis. Plant Physiol 169: 1946–1960. https://doi.org/10.1104/pp.15.01161
Google Scholar: Author Only Title Only Author and Title

Maruyama D, Völz R, Takeuchi H, Mori T, Igawa T, Kurihara D, Kawashima T, Ueda M, Ito M, Umeda M, Nishikawa S, Groß-Hardt R, Higashiyama T (2015) Rapid elimination of the persistent synergid through a cell fusion mechanism. Cell 161: 907–918. https://doi.org/10.1016/j.cell.2015.03.018
Google Scholar: Author Only Title Only Author and Title

Miki D, Zhang W, Zeng W, Feng Z, Zhu JK (2018) CRISPR/Cas9-mediated gene targeting in Arabidopsis using sequential transformation. Nat Commun 9: 1967. https://doi.org/10.1038/s41467-018-04416-0
Google Scholar: Author Only Title Only Author and Title

Motomura K, Takeuchi H, Notaguchi M, Tsuchi H, Takeda A, Kinoshita T, Higashiyama T, Maruyama D (2021) Persistent directional growth capability in Arabidopsis thaliana pollen tubes after nuclear elimination from the apex. Nat Commun 12: 2331. https://doi.org/10.1038/s41467-021-22661-8
Google Scholar: Author Only Title Only Author and Title

Muro K, Matsuura-Tokita K, Tsukamoto R, Kanaoka MM, Ebine K, Higashiyama T, Nakano A, Ueda T (2018) ANTH domain-containing proteins are required for the pollen tube plasma membrane integrity via recycling ANXUR kinases. Commun Biol 1: 152. https://doi.org/10.1038/s42003-018-0158-8
Google Scholar: Author Only Title Only Author and Title

Nagasaki N, Tomioka R, Maeshima M (2008) A hydrophilic cation-binding protein of Arabidopsis thaliana, AtPCaP1, is localized to plasma membrane via N-myristoylation and interacts with calmodulin and the phosphatidylinositol phosphates PtdIns(3,4,5)P(3) and PtdIns(3,5)P(2). FEBS Journal 275: 2267–2282. https://doi.org/10.1111/j.1742-4658.2008.06379.x
Google Scholar: Author Only Title Only Author and Title

Nakagawa T, Suzuki T, Murata S, Nakamura S, Hino T, Maeo K, Tabata R, Kawai T, Tanaka K, Niwa Y, Watanabe Y, Nakamura K, Kimura T, Ishiguro S (2007) Improved Gateway binary vectors: high-performance vectors for creation of fusion constructs in transgenic analysis of plants. Biosci Biotechnol Biochem 71: 2095–2100. https://doi.org/10.1271/bbb.70216
Google Scholar: Author Only Title Only Author and Title

Oda Y, Fukuda H (2013) Rho of plant GTPase signaling regulates the behavior of Arabidopsis kinesin-13A to establish secondary cell wall patterns. Plant Cell 25: 4439–4450. https://doi.org/10.1105/tpc.113.117853
Google Scholar: Author Only Title Only Author and Title

Oh SA, Allen T, Twell D (2010) A ticket for the live show: Microtubules in male gametophyte development. Plant Signal Behav 5: 614–617. https://doi.org/10.4161/psb.11505
Google Scholar: Author Only Title Only Author and Title

Perico C, Sparkes I (2018) Plant organelle dynamics: Cytoskeletal control and membrane contact sites. New Phytol 220: 381–394. https://doi.org/10.1111/nph.15365
Google Scholar: Author Only Title Only Author and Title

Qin T, Liu X, Li J, Sun J, Song L, Mao T (2014) Arabidopsis microtubule-destabilizing protein 25 functions in pollen tube growth by severing actin filaments. Plant Cell 26: 325–339. https://doi.org/10.1105/tpc.113.119768
Google Scholar: Author Only Title Only Author and Title

Qin Y, Leydon AR, Manziello A, Pandey R, Mount D, Denic S, Vasic B, Johnson MA, Palanivelu R (2009) Penetration of the stigma and style elicits a novel transcriptome in pollen tubes, pointing to genes critical for growth in a pistil. PLOS Genet 5: e1000621. https://doi.org/10.1371/journal.pgen.1000621
Google Scholar: Author Only Title Only Author and Title

Raudaskoski M, Åström H, Laitiainen E (2001) Pollen tube cytoskeleton: structure and function. J Plant Growth Regul 20: 113–130. https://doi.org/10.1007/s003440010015
Google Scholar: Author Only Title Only Author and Title

Schattner S, Schattner J, Munder F, Höppe E, Walter WJ (2020) A tug-of-war model explains the saltatory sperm cell movement in Arabidopsis thaliana pollen tubes by kinesins with calponin homology domain. Front Plant Sci 11: 601282. https://doi.org/10.3389/fpls.2020.601282
Google Scholar: Author Only Title Only Author and Title

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A (2012) Fiji: an open-source platform for biological-image analysis. Nat Methods 9: 676–682. https://doi.org/10.1038/nmeth.2019
Google Scholar: Author Only Title Only Author and Title

Sugi N, Maruyama D (2023) Exploring novel polytubey reproduction pathways utilizing cumulative genetic tools. Plant Cell Physiol 64: 454–460. https://doi.org/10.1093/pcp/pcad021
Google Scholar: Author Only Title Only Author and Title

Tamura K, Iwabuchi K, Fukao Y, Kondo M, Okamoto K, Ueda H, Nishimura M, Hara-Nishimura I (2013) Myosin XI-I links the nuclear membrane to the cytoskeleton to control nuclear movement and shape in Arabidopsis. Curr Biol 23: 1776–1781. https://doi.org/10.1016/j.cub.2013.07.035
Google Scholar: Author Only Title Only Author and Title

Tatout C, Evans DE, Vanrobays E, Probst AV, Graumann K (2014) The plant LINC complex at the nuclear envelope. Chromosome Res 22: 241–252. https://doi.org/10.1007/s10577-014-9419-7
Google Scholar: Author Only Title Only Author and Title

Tiwari SC, Polito VS (1988) Organization of the cytoskeleton in pollen tubes of Pyrus communis: A study employing conventional and freeze-substitution electron microscopy, immunofluorescence, and rhodamine-phalloidin. Protoplasma 147: 100–112. https://doi.org/10.1007/BF01403337
Google Scholar: Author Only Title Only Author and Title

Uno K, Sugimoto N, Sato Y (2021) N-aryl pyrido cyanine derivatives are nuclear and organelle DNA markers for two-photon and super-resolution imaging. Nat Commun 12: 2650. https://doi.org/10.1038/s41467-021-23019-w
Google Scholar: Author Only Title Only Author and Title

Vogler F, Konrad SS, Sprunck S (2015) Knockin' on pollen's door: Live cell imaging of early polarization events in germinating Arabidopsis pollen. Front Plant Sci 6: 246. https://doi.org/10.3389/fpls.2015.00246
Google Scholar: Author Only Title Only Author and Title

Walter WJ, Machens I, Rafieian F, Diez S (2015) The non-processive rice kinesin-14 OsKCH1 transports actin filaments along microtubules with two distinct velocities. Nat Plants 1: 15111. https://doi.org/10.1038/nplants.2015.111

Google Scholar: Author Only Title Only Author and Title

Wang X, Zhu L, Liu B, Wang C, Jin L, Zhao Q, Yuan M (2007) Arabidopsis microtubule-associated protein18 functions in directional cell growth by destabilizing cortical microtubules. Plant Cell 19: 877–889. https://doi.org/10.1105/tpc.106.048579

Google Scholar: Author Only Title Only Author and Title

Wei L, Liu B, Li Y (2005) Distribution of a kinesin-related protein on Golgi apparatus of tobacco pollen tubes. Chin Sci Bull 50: 2175–2181. https://doi.org/10.1007/BF03182668

Google Scholar: Author Only Title Only Author and Title

Yu Y, Li Y, Li L, Lin J, Zheng C, Zhang L (2009) Overexpression of PwTUA1, a pollen-specific tubulin gene, increases pollen tube elongation by altering the distribution of alpha-tubulin and promoting vesicle transport. J Exp Bot 60: 2737–2749. https://doi.org/10.1093/jxb/erp143

Google Scholar: Author Only Title Only Author and Title

Zhang HQ, Bohdanowicz J, Pierson ES, Li YQ, Tiezzi A, Cresti M (1995) Microtubular organization during asymmetrical division of the generative cell in Gagea lutea. J Plant Res 108: 269–276. https://doi.org/10.1007/BF02344352

Google Scholar: Author Only Title Only Author and Title

Zhang X, Henriques R, Lin SS, Niu QW, Chua NH (2006) Agrobacterium-mediated transformation of Arabidopsis thaliana using the floral dip method. Nat Protoc 1: 641–646. https://doi.org/10.1038/nprot.2006.97

Google Scholar: Author Only Title Only Author and Title

Zhou X, Groves NR, Meier I (2015) SUN anchors pollen WIP-WIT complexes at the vegetative nuclear envelope and is necessary for pollen tube targeting and fertility. J Exp Bot 66: 7299–7307. https://doi.org/10.1093/jxb/erv425

Google Scholar: Author Only Title Only Author and Title

Zhou X, Meier I (2014) Efficient plant male fertility depends on vegetative nuclear movement mediated by two families of plant outer nuclear membrane proteins. Proc Natl Acad Sci U S A 111: 11900–11905. https://doi.org/10.1073/pnas.1323104111

Google Scholar: Author Only Title Only Author and Title

Zhu L, Zhang Y, Kang E, Xu Q, Wang M, Rui Y, Liu B, Yuan M, Fu Y (2013) MAP18 regulates the direction of pollen tube growth in Arabidopsis by modulating F-actin organization. Plant Cell 25: 851–867. https://doi.org/10.1105/tpc.113.110528

Google Scholar: Author Only Title Only Author and Title