Transcriptome-wide quantitative profiling of PUS7-dependent pseudouridylation by nanopore direct long read RNA sequencing

Mohit Bansal[1,2,*], Anirban Kundu[2,3], Andrew Gibson[4], Anamika Gupta[1,2], Jane Ding[1,2], Sanjay Varma RudraRaju[5], Sunil Sudarshan[2,3] and Han-Fei Ding[1,2*]

[1]Department of Pathology and [2]O'Neal Comprehensive Cancer Center, [3]Department of Urology, [4]Departments of Medicine, Heersink School of Medicine, University of Alabama at Birmingham, AL 35294, USA, [5]Independent scholar, New York, NY 07932, USA.

*Correspondence: M. Bansal mbansal@uabmc.edu and H.-F. Ding hding@uabmc.edu

Abstract

Pseudouridylation is a prevalent post-transcriptional RNA modification that impacts many aspects of RNA biology and function. The conversion of uridine to pseudouridine (Ψ) is catalyzed by the family of pseudouridine synthases (PUSs). Development of robust methods to determine PUS-dependent regulation of Ψ location and stoichiometry in low abundant mRNA is essential for biological and functional understanding of pseudouridylation. Here, we present a framework, NanoPsiPy, for identifying Ψ sites and quantify their levels in poly-A RNA at single-nucleotide resolution using direct RNA long-read Nanopore sequencing, based on the observation that Ψ can cause characteristic U-to-C basecalling errors in Nanopore direct RNA sequencing data. Our method was able to detect low and high stoichiometric Ψ sites in human mRNA. We validated our method by transcriptome-wide quantitative profiling of PUS7-dependent Ψ sites in poly-A RNA from a *MYCN*-amplified neuroblastoma cell line. We identified 8,625 PUS7-dependent Ψ sites in 1,246 mRNAs that encode proteins involved primarily in ribosome biogenesis, translation, and mitochondrial energy metabolism. Our work provides the first example of using direct RNA long-read Nanopore sequencing for transcriptome-wide quantitative profiling of mRNA pseudouridylation regulated by a PUS. We envision that our method will facilitate functional interrogation of PUSs in biological and pathological processes.

Introduction

Research spanning several decades has demonstrated that RNAs undergo various posttranscriptional modifications, including pseudouridylation[1-3]. Pseudouridine (Ψ) is the first identified RNA modification[4] that constitutes 0.2–0.6% of total uridines in mammalian mRNAs[5]. Ψ can enhance the stability and rigidity of RNAs, influencing cellular processes such as translation, splicing, and telomere maintenance[2,6]. The existence of Ψ in mRNAs remained unknown until 2014-2015[5,7,8], a revelation facilitated by advancements in next-generation sequencing technologies. Human genome encodes thirteen annotated pseudouridylation installation enzymes (pseudouridine synthases, PUSs)[1], highlighting the robust and dynamic nature of Ψ installation in human RNA. The conversion of uridine to Ψ starts with breaking the N1-C1' bond linking the uracil base to the sugar, followed by an 180° base rotation around the N3-C6 axis. The resulting Ψ contains an extra hydrogen bond donor at N1 and a C5-C1' base-sugar linkage[9]. The success of mRNA-based COVID-19 vaccines in which all uridines are replaced by Ψs underscores the crucial role played by Ψ modification in promoting protein expression and increasing the efficacy of mRNA vaccines, showcasing the importance of understanding and harnessing RNA modifications for therapeutic purposes[10]. Ψ does not disrupt canonical base pairing, making it indistinguishable from uridine in hybridization-based methods. Additionally, the molecular weight of Ψ matches that of uridine, complicating direct detection through mass spectrometry[11]. This difficulty is exacerbated when trying to identify Ψ sites within specific mRNA sequences.

Various techniques, such as Pseudo-Seq[7], Ψ -Seq[8], and Ceu-Seq[5] that are based on illumina-short read sequencing, have used the CMC (N-cyclohexyl-N'-β-(4-methylmorpholinium)ethylcarbodiimide)-based approach for transcriptome-wide Ψ mapping.

While CMC-based methods are effective in generating a stop signature during reverse transcription (RT) for Ψ site identification, these methods lack base resolution and stoichiometry information at the modified bases. In contrast to aforementioned techniques, a new generation of methods based on bisulfite-induced deletion signatures at Ψ sites, including RBS-seq[12], BID-seq[13], and PRAISE[14], offer quantitative transcriptome-wide mapping of Ψ sites. However, it is important to note that Ψ-bisulfite adduct formation involves harsh chemical reactions as an intermediate step. Moreover, the read-length limitation of Illumina sequencing also narrows the possibility to examine Ψ distribution in mRNA splice isoforms and the linkage of multiple Ψ sites in single RNA molecules.

We wanted to develop a nanopore direct RNA sequencing method for Ψ site identification and stoichiometry measurement on long native mRNAs without reliance on harsh chemical reactions, providing a more direct and precise means of studying pseudouridylation. Several studies have utilized nanopore-based direct RNA sequencing to directly examine RNA modifications. The detection of Ψ using direct RNA sequencing has been confirmed for ribosomal RNAs (rRNAs) in Saccharomyces cerevisiae[15] and human mRNA[16,11], based on U-to-C basecalling errors. However, it remains to be established whether direct RNA long-read nanopore sequencing is sufficiently robust to determine PUS-dependent regulation of Ψ location and stoichiometry in mRNA. This study was aimed to develop a method to validate the U-to-C basecalling signature as a distinctive feature of Ψ and to quantify transcriptome-wide PUS7-dependent Ψ sites using in vivo and in vitro systems.

Results

Development of NanoPsiPy for identification and quantification of Ψ modification based on U-to-C basecalling errors in nanopore sequencing

We have developed NanoPsiPy, a computational pipeline (https://github.com/vetmohit89/NanoPsiPy), to identify and quantify Ψ levels in RNA based on U-to-C basecalling errors in Nanopore direct RNA sequencing data (Figure 1). Following the alignment of RNA reads to the human reference sequence, NanoPsiPy systematically gathers positional data for each uridine base. Subsequently, it conducts statistical comparisons of U-to-C basecalling errors between two sample groups (control and treatment), identifying and quantifying Ψ modifications at specific locations. As Ψ detection by Nanopore direct RNA sequencing is based on basecalling errors, it is essential to develop a method that accounts only primary alignment of each RNA read (i.e., the alignment with the highest mapping quality) to reduce false positive errors. NanoPsiPy filters out secondary alignments and only uses reads with mapping quality score of 20 or higher, which correspond to the probability of correct alignment of 99% or higher, for downstream analysis and calculating U-to-C basecalling errors. Another potential source of error is the presence of single-nucleotide polymorphisms (SNPs) whereby the base is different from the reference genome or transcriptome sequence. SNPs can cause systematic basecalling errors. NanoPsiPy removes any SNP errors from downstream analysis by comparative analysis of the U-to-C mismatch at each site between control and treatment samples (Figure 1A).

We tested the ability of our nanopore direct RNA sequencing and the NanoPsiPy pipeline to identify and quantify Ψ based on U-to-C basecalling errors using two PUS7-dependent systems. We used lentiviruses expressing a shRNA against human PUS7 (shPUS7-33) to silence

PUS7 expression in the *MYCN*-amplified neuroblastoma cell line BE(2)-C, with shGFP as control (Supplemental Figure 1). It has been shown previously that PUS7 pseudouridylates U5614 in MALAT1 RNA (base position according to NR-002819.4)[17]. In control shGFP BE(2)-C cells, MALAT1 RNA at U5614 showed 75% of RNA reads having a U-to-C basecalling error, which was reduced to 10% in BE(2)-C cells with PUS7 knockdown (Figures 1B-C). To further validate this U-to-C basecalling error as a signature of PUS7-dependent Ψ modification, we performed an in vitro pseudouridylation assay [18] using an in vitro transcribed (IVT) MALAT1 RNA oligo that includes 86 nucleotides upstream and 66 nucleotides downstream of U5614. The RNA oligo was incubated either with immunopurified wild-type human PUS7 or with a catalytically inactive mutant (PUS7 D294A)[19] and analyzed systematically for basecalling features, including mismatches, deletion, insertion and per-base qualities. Approximately 38% of the IVT MALAT1 RNA reads treated with wild-type PUS7 showed a U-to-C basecalling error compared to the 1% error in the RNA reads treated with the mutant PUS7 (Figures 1D-E). We found no significant differences in other basecalling features between the two samples (Figure 1F). These findings confirmed that NanoPsiPy analysis of nanopore direct RNA sequencing data was able to detect and quantify Ψ in RNA by identifying U-to-C basecalling errors.

Nanopore sequencing detects nucleotide bases from their ionic current signature. Our pipeline NanoPsiPy uses Guppy (v6.4.2) for basecalling. By using a pool of different IVT RNA oligos containing only canonical RNA nucleotides, we calculated the basecalling accuracy. We found that our pipeline is highly accurate, with 95.99% for adenine, 97.22% for cytosine, 97.26% for guanine, and 94.51% for uridine (Figure 1G). Additionally, we found that Guppy could reliably identify unmodified and aligned uridine with an average background U-to-C calling error of 2.99% (Figure 1H).

Nanopore direct RNA sequencing method sequences RNA in the 3' to 5' direction, raising the question of whether U-to-C error-based Ψ detection is impacted by the preceding 3' base. Our analysis of IVT RNA data revealed that when the preceding 3' base is U, A, C, or G, there is a background error of 4.8%, 3.07%, 2.07%, or 1.18% in identifying the candidate U as C, respectively (Figure 1I). Further, we observed that the average base quality is low when RNA reads at any location are less than 4 (Figure J). Therefore, to minimize the inclusion of U sites with background errors, we defined a threshold of 10% basecalling error for U-to-C estimation and a minimum of 9 reads at any given U site for Ψ detection and quantification in the whole transcriptome.

Transcriptome-wide identification and quantification of Ψ sites in poly-A RNA from a human neuroblastoma cell line using nanopore direct RNA sequencing and NanoPsiPy

We applied nanopore direct RNA sequencing and NanoPsiPy for de novo detection and quantification of transcriptome-wide Ψ modifications in the *MYCN*-amplified neuroblastoma cell line BE(2)-C. For background control (RNA without modifications), we used the published nanopore direct RNA sequencing data from a HeLa IVT poly-A RNA library that was constructed first by reverse transcription of HeLa poly-A RNAs (with modifications) into cDNAs to remove all modifications and then by in vitro transcription of the cDNAs back into RNAs[11]. We conducted nanopore direct sequencing of poly-A RNA from BE(2)-C shGFP cells and compared the sequencing data with the HeLa IVT poly-A RNA nanopore sequencing data, which showed an overlapping distribution of read lengths (Figure 2A). We then mapped RNA sequencing reads from both datasets to the human genome reference sequence (GRCh38.p14) to identify all the uridine sites common between BE(2)-C and HeLa IVT RNA and assigned the U-

to-C basecalling error as Ψ modification at each site. We defined Ψ sites in BE(2)-C poly-A RNA based on two criteria: 1) the U-to-C basecalling error at each site in BE(2)-C must exceed 10% and 2) the *p* value of U-to-C basecalling error must be < 0.01. Using data from more than 2 millions uridines in BE(2)-C poly-A RNA and more than 3 millions uridines in HeLa IVT RNA, we identified 39,756 Ψ sites in BE(2)-C poly-A RNA with a median Ψ level of 16.67% (Supplement Table 1). To assess the accuracy of our NanoPsiPy pipeline in detecting Ψ sites, we compared the identified BE(2)-C Ψ sites with the previously annotated 1,691 Ψ sites in HeLa by nanopore direct RNA sequencing[11]. Our analysis revealed 517 Ψ sites shared between BE(2)-C and HeLa cells (Supplemental Table 2). Furthermore, we observed a strong correlation (R^2 = 0.618) in the Ψ fraction across these shared Ψ sites (Figure 2B).

Next, we examined the stoichiometric distribution of Ψ levels in BE(2)-C poly-A RNA. Our analysis revealed that approximately 66% of the Ψ sites were modified at the levels of 10-20% and ~11% at the levels of more than 30% (Figure 2C). In addition, our analysis revealed a negative correlation between the levels of Ψ modification and the levels of sequencing coverage (reads) at specific sites: the lower the number of reads, the higher the median Ψ levels (Figure 2D).

We next analyzed the sequence contexts for the BE(2)-C Ψ sites and found that the TRUB1 motifs GUUCA, GUUCC, and GUUCU, and the PUS7 motif UGUAG were the most highly represented (Figure 2E). The median level of Ψ across the top 14 motifs was 18.26% (Figure 2E). For all the Ψ sites, uridine exhibits a strong preference at the -2, -1, and +1 positions (Figure 2F). The most frequent motifs are UUUUG (4.92%), AUUUG (3.78%), UUUUA (3.56%), CUUUG (3.64%), and UUUUU (3.43%) (Figure 2F).

To determine the distribution pattern of the identified Ψ sites in poly-A RNA, we mapped the RNA reads to the transcriptome reference sequence (gencode.v43.transcripts.fa) and observed that within mRNA, Ψ sites were highly enriched in the coding sequence (CDS) and 3′ untranslated region (3'UTR) (Figure 2G), as reported previously using CeU-Seq[5], Bid-seq[13] and Nanopore sequencing methods[11,16]. In addition, we found low levels of Ψ distribution in retained introns (4.46%) and long non-coding RNA (1.14%) (Figure 2G). The limited detection of Ψ sites in the 5′ UTR could be due to a lower level of coverage of the 5′ UTR compared to CDS and 3'UTR as nanopore sequencing moves in the 3' to 5' direction. To address this issue, we generated a metatranscript plot using R2D tools[20] to compare the proportion of significant Ψ sites along mRNA transcripts, which showed comparable Ψ density distribution in the 5'UTR, CDS and 3'UTR (Figure 2H).

Martinej et al.[21] recently reported that Ψs are enriched in proximal introns (within 500 nucleotides of splice sites). In agreement with the report, we found that the Ψ sites in BE(2)-C poly-A RNA were enriched around 200 nucleotides from the exon-exon junctions (Figure 2I).

Transcriptome-wide identification and quantification of PUS7-dependent Ψ sites in BE(2)-C poly-A RNA

We applied our method to map and quantify PUS7-dependent Ψ sites in BE(2)-C transcriptome. We isolated poly-A RNA from control (shGFP) and PUS7 knockdown (shPUS7-33) BE(2)-C cells (Supplemental Figure 1) and prepared libraries for direct RNA sequencing. Two independent shGFP RNA libraries produced a total of ~2.254 million reads, with mean read quality of 9.9 and an average N50 read length of 1,124 bases (defined as the shortest read length needed to cover 50% of the sequenced nucleotides). Two independent shPUS7 RNA libraries

produced a total of 1.44 million reads, with mean read quality of 10.0 and an average N50 read length of 1,351 bases. The median RNA read lengths were 702 and 789 bases for shGFP and shPUS7-33 samples, respectively (Figure 3A and Supplemental Table 3).

To identify PUS7-dependent Ψ sites, we compared U-to-C basecalling errors at uridine sites in shPUS7 poly-A RNA with Ψ sites in shGFP poly-A RNA that were identified as described above (shGFP BE(2)-C vs. HeLa IVT). Employing the two controls (shGFP BE(2)-C and HeLa IVT), we aimed to enhance the precision of our analysis and eliminate potential false positives in calling PUS7-dependent Ψ sites. We identified 8,625 putative PUS7-dependent Ψ sites within protein-coding mRNA, with Ψ stoichiometries ranging from 10.00% to 88.88% and a median Ψ stoichiometry of 16.78%. Additionally, we identified 294 putative PUS7-dependent Ψ sites within retained introns, with Ψ stoichiometries spanning 10.22% to 68.42% and a median Ψ stoichiometry of 18.08%, and 58 putative PUS7 Ψ sites in long non-coding RNA, exhibiting Ψ stoichiometries between 10.28% and 52% and a median Ψ stoichiometry of 18.18% (Supplemental Table 4). Interestingly, single-read analysis revealed a subset of BE(2)-C mRNAs contain multiple PUS7-dependent Ψ sites within each mRNA molecule. We quantified the overall Ψ modification level in each gene by calculating its Ψ-strength (sum of Ψ fractions at all the Ψ sites within a single gene)[13]. PUS7 knockdown resulted in a global reduction in Ψ-strength of PUS7 target mRNAs (Figure 3B).

Analysis of motif frequency and representation for the PUS7-dependent Ψ sites revealed that the previously reported HeLa[11] Ψ motifs UUUAU, AUUUA and UUUAA were the most represented, whereas the Ψ motifs UUUUG, AUUUG and CUUUG were the most frequently detected. The median Ψ stoichiometry of the top 14 motifs was 19.79% while the median stoichiometry of transcriptome-wide PUS7-depended Ψ sites was 16.78% (Figure 3C).

As presented above for transcriptome-wide Ψ sites in shGFP BE(2)-C mRNAs (Figure 2G), PUS7-dependent Ψ sites also showed a similar distribution pattern, with ~48.66% of the Ψ sites in the 3′UTR, 44.98% in the CDS, and 2.43% in the 5′UTR (Figure 3D). Again, the metatranscript plot for PUS7-dependent Ψ sites showed comparable density enrichment in the 5' UTR, CDS, and 3'UTR (Figure 3E). We also found that PUS7-dependent Ψ sites were enriched around 200 nucleotide proximal from the exon-exon junctions (Figure 3F).

We identified 8,625 putative PUS7-dependent Ψ sites in 1,246 unique protein coding genes. To interrogate the functions of these genes, we performed Gene Ontology (GO) analysis, which revealed the PUS7 target genes are involved in ribosome biogenesis, translation, RNA and protein binding, and mitochondrial energy metabolism (Figure 3G and Supplemental Table 5). Further, we identified 57 Ψ sites in 13 unique long non-coding RNAs. GO analysis found highly significant enrichment of GO terms corresponding to RNA processing (Supplemental Table 6).

Discussion

We have developed NanoPsiPy for processing nanopore direct RNA long-read sequencing data to identify Ψ at individual RNA molecule and whole transcriptome levels with location information and stoichiometry quantification. NanoPsiPy abstracts U-to-C basecalling errors to detect and quantify transcriptome-wide Ψ modifications between any two samples using comparison paradigm to minimize false positives. By treating in vitro RNA oligos with wild-type and mutant (D294A) PUS7, our study demonstrated that U-to-C basecalling is the sole characteristic enabling the identification of Ψ in nanopore direct RNA sequencing data. NanoPsiPy weigh the pooled sample analysis to identify transcriptome-wide Ψ, comparing the

average of U-to-C errors between two conditions to minimize false positives due to low coverage that is an issue with direct mRNA sequencing.

Quantitative MS analysis estimated that the Ψ/U ratio is approximately 0.2-0.6% in human mRNA[5], which is similar to the ratio of m[6]A/A at 0.15-0.6% found in the human transcriptome[22-24]. Using NanoPsiPy, we identified 39,756 Ψ sites in poly-A RNA from the human neuroblastoma cell line BE(2)-C, which again is comparable to the 42,116 m[6]A sites detected in poly-A RNA from the human mammary epithelial cell line HMEC using MINES analysis of nanopore direct RNA sequencing data[25]. Recently, Tavakoli et al. reported the identification of 1,691 Ψ sites in HeLa poly-A RNA using PsiNanopore that also relies on U-to-C basecalling errors in direct RNA sequencing [11]. Our analysis identified 517 Ψ sites that are shared between BE(2)-C and HeLa cells, suggesting that a common set of poly-A RNAs are targeted by pseudouridylation in human cancer cell lines of different tissue origins. On the other hand, we identified significantly more Ψ sites in BE(2)-C poly-A RNAs. We suspect that a major reason for the differences in Ψ locations and numbers between HeLa and BE(2)-C cells is the cell-type specific transcriptome profiles because modifications in highly expressed RNAs are most likely to be detected. BE(2)-C cells are of neuronal origin with amplification of the MYCN oncogene, a member of the MYC family of transcription factors that regulates RNA expression of thousands of genes [26,27]. Thus, NanoPsiPy is highly effective in comprehensively identifying Ψ sites and quantifying their levels in the whole transcriptome.

It was reported recently that the median Ψ stoichiometry in HEK293T mRNA is approximately 10%, with 80% of the Ψ sites showing Ψ levels < 20% and 10% showing > 40% [14]. In agreement, our NanoPsiPy analysis revealed ~66% of the Ψ sites in BE(2)-C poly-A RNA having modification levels at 10-20%, 23% at 20-30%, and 11% at > 30%. Interestingly, the

median methylation level of m^{5}C sites in mRNA is in the range of 15-18%. Across various tissues or cell types, the majority (62–70%) of m^{5}C sites exhibited low methylation levels (<20%), and 8-10% showed moderate to high methylation levels (>40%)[28]. Collectively, these findings support the notion that RNA modifications occur at relatively low stoichiometric levels across the entire transcriptome.

Previous studies suggest that Ψ distribution in mRNAs is overrepresented in CDS and 3′UTR and underrepresented in 5'UTR [11,13]. However, our analysis showed that Ψ is proportionally present in all regions, with no evidence of strong bias. In principle, this broad distribution endows Ψ modifications with the ability to influence many aspects of mRNA metabolism and function. In addition, we identified 1,246 PUS7 target mRNAs in the *MYCN*-amplified neuroblastoma cell line BE(2)-C and GO analysis suggest that they are involved primarily in ribosome biogenesis, translation, and mitochondrial energy metabolism. These findings lay the foundation for further investigation of the functional significance of PUS7-dependent pseudouridylation in neuroblastoma and other MYC-driven cancers.

In summary, we have developed a method for quantitative profiling of transcriptome-wide Ψ sites, which allows us to identify and quantify PUS7-dependent Ψ sites at single mRNA molecule resolution with functional implications. We anticipate that NanoPsiPy will facilitate the identification and functional characterization of not only pseudouridylation but also other RNA modifications.

Material and methods

Cell culture and PUS7 knockdown

Neuroblastoma BE(2)-C) cells were cultured in DME/F-12 1:1 (HyClone SH30023) supplemented with 10% Fetal Bovine Serum (Fisher Scientific FB12999102) and 1% Penicillin-Streptomycin (Lonza 17602E). The PUS7 knockdown lentiviral construct shPUS7-33 (TRCN0000061933) was obtained from Sigma-Aldrich. BE(2)-C cells were infected with lentiviruses expressing shGFP or shPUS7-33, and PUS7 knockdown was confirmed by immunoblotting using rabbit anti-PUS7 (1:1000, Thermo Fisher PA5-54983).

Poly-A RNA isolation, library preparation, and nanopore sequencing

Total RNA was obtained through TRIzol extraction, and poly(A) mRNA was isolated using the NEBNext Poly(A) mRNA Magnetic Isolation kit (NEB E7490L). The RNA library for direct RNA sequencing was prepared from ~500 ng of poly-A RNA according to the Oxford Nanopore technology (ONT) direct RNA sequencing protocol version direct-rna-sequencing-sqk-rna002-DRS_9080_v2_revS_14Aug2019-minion. For sequencing, the ONT FLO-MIN106D (R9.4.1) flow cell was primed according to the manufacturer's protocol, and the elute was mixed with an RNA running buffer (RRB) and loaded onto the flow cell. Sequencing was performed using MinKnow (version 19.12.5). Each flow cell was sequenced for 24-72 hours per each run, depending on flowcell exhaustion rate. Two replicates from different passages and different flow cells were used for each biological replicate.

PUS7 immunoprecipitation and purification

All immunoprecipitation steps were performed at 4°C following the protocol of Kundu et al. with some modifications[29]. Briefly, 293FT cells expressing either wild-type (WT) or mutant (D294A) PUS7 (myc-tagged) were lysed in the buffer containing 50 mM HEPES pH 7.5, 130

mM NaCl, 0.3% NP-40, 1 mM EDTA, and 1× protease/phosphatase inhibitor (Thermo Fisher 78440). Approximately 3 mg of protein lysates were mixed with protein A/G-magnetic beads (Thermo Fisher 88802) and 2 μg of either anti-Myc or control IgG antibody and rotated for 16 hours at 4°C. Subsequently, the magnetic beads were pelleted and washed for three times in the buffer containing 20 mM Tris-HCl at pH 8.0, 130 mM NaCl, 0.3% NP-40, and 1 mM EDTA. Then myc-PUS7 (WT or D294A) was eluted from the beads using myc-peptide (Protein Tech 2yp) following manufacture's protocol. Finally, using 30 KDa ultracentrifugal filter, the eluted proteins were concentrated, and buffer exchanged in the 1X pseudouridylation buffer (100 mM Tris-HCL pH-8.0, 100 mM ammonium acerate, 5 mM MgCl2, 0.3 mM EDTA, and 130 mM NaCl). Concentrated proteins were either used immediately or stored in -80°C until use.

In vitro RNA pseudouridylation [18]

The synthetic 192-nucleotide (nt) MALAT1 DNA (Supplemental Table 7) contains 86 nt upstream and 66 nt downstream of the pseudouridine site (U5614, NR_002819.4). A 22-nt sequence containing the T7 promoter and a 25-nt adapter sequence were added to the 5' and 3' ends, respectively. The oligo was obtained from IDT DNA and amplified by PCR using Phusion Polymerase (NEB F630) and purified for in vitro transcription using the MEGAshortscript T7 kit (Invitrogen AM1333). The RNA product was purified using a urea gel, denatured, and refolded. In vitro pseudouridylation was carried out by incubating 15-30 pmol of folded RNA with 1 μg of immunopurified PUS7 or PUS7 D294A. Poly-A tail was added using a Poly (A) tailing kit (Invitrogen AM1350). Library was prepared and loaded into the FLO-MIN106D flow cell for sequencing using MinKnow (version 19.12.5) as described above. Each flow cell was sequenced for 24 h.

Data analysis

All raw fast5 files generated from nanopore sequencing were uploaded to UAB Cheaha cluster for further analysis. Multi-fast5 files were basecalled in real-time by Guppy (6.4.2) using the high accuracy model with minimum quality score of 7. The fastq files were analyzed using the NanoPsiPy where reads were aligned to the reference file (GRCh38.p14 or gencode.v43.transcripts.fa.) using minimap2 (2.17). The mapped reads were piled up to the human reference sequence by samtools (v1.12). The U-to-C basecalling "error" data were extracted and compared between two conditions using NanoPsiPy (http://github.com/vetmohit89/NanoPsiPy.git). NanoPsiPy tool was built around Nanopore_psu tool available at (https://github.com/sihaohuanguc/Nanopore_psU/)[16]. Finally, data was prepared by applying condition when the ratio of C reads from the total of U+C reads of the control group is greater than the ratio of C reads from the total of U+C reads for the treatment group. After filtering the input data, the Chi-square test is applied to a contingency table constructed from the read counts in different categories (control_C, control_U, treatment_C, treatment_U). Yates' continuity correction is applied to improve the accuracy of the results when the sample size is limited and $p < 0.01$ is defined as significant. Gene ontology (GO) analysis was performed using gprofiler2 (https://biit.cs.ut.ee/gprofiler/gost).

Accession Numbers

All FASTQ files generated in this study have been publicly shared and are accessible through the NIH NCBI SRA under the Bio-Project accession PRJNA961708. The data for two HeLa IVT

RNA samples were retrieved from NIH NCBI SRA with the accession numbers SRR23932675 and SRR23929127.

Supplemental Tables description

Supplemental Table 1: Significant Ψ sites identified in BE(2)-C.

Supplemental Table 2: Common Ψ sites identified in BE(2)-C and HeLa poly-A RNA.

Supplemental Table 3: Direct RNA Library description of BE(2)-C shGFP and shPUS7-33.

Supplemental Table 4: Significant PUS7-dependent Ψ sites identified in protein coding, retained RNA and long non-coding RNA of BE(2)-C.

Supplemental Table 5: Gene ontology analysis of protein coding mRNAs containing PUS-dependent Ψ sites

Supplemental Table 6: Gene ontology analysis of long non-coding RNAs containing PUS7-dependent Ψ sites.

Supplemental Table 7: RNA oligo sequences.

Funding

This work is supported by the National Institutes of Health grant R01CA190429 to H.-F.D.

Author contributions

M.B. and H-F.D. conceived the study and designed the experiments with contributions from A.K., A.Gibson, A.Gupta, J.D., and S.S. M.B. performed nanopore RNA sequencing and data analysis with assistance from A.Gibson and S.V.R. A.K. performed PUS7 purification. A.Gupta. and J.D. performed cell culture and immunoblotting. M.B. and H-F.D. wrote the paper with contributions from S.S. and J.D.

H-F.D. supervised and provided funding for the project. All authors read the manuscript and approved its contents.

References:

Li, X., Ma, S. & Yi, C. Pseudouridine: the fifth RNA nucleotide with renewed interests. *Current Opinion in Chemical Biology* **33**, 108-116 (2016).

Borchardt, E. K., Martinez, N. M. & Gilbert, W. V. Regulation and function of RNA pseudouridylation in human cells. *Annual review of genetics* **54**, 309-336 (2020).

Rodell, R., Robalin, N. & Martinez, N. M. Why U matters: detection and functions of pseudouridine modifications in mRNAs. *Trends in Biochemical Sciences* **49**, 12-27 (2024).

Davis, F. F. & Allen, F. W. Ribonucleic acids from yeast which contain a fifth nucleotide. *J Biol Chem* **227**, 907-915 (1957).

Li, X. *et al.* Chemical pulldown reveals dynamic pseudouridylation of the mammalian transcriptome. *Nature chemical biology* **11**, 592-597 (2015).

Garus, A. & Autexier, C. Dyskerin: an essential pseudouridine synthase with multifaceted roles in ribosome biogenesis, splicing, and telomere maintenance. *RNA* **27**, 1441-1458 (2021).

Carlile, T. M. *et al.* Pseudouridine profiling reveals regulated mRNA pseudouridylation in yeast and human cells. *Nature* **515**, 143-146 (2014).

Schwartz, S. *et al.* Transcriptome-wide mapping reveals widespread dynamic-regulated pseudouridylation of ncRNA and mRNA. *Cell* **159**, 148-162 (2014).

De Zoysa, M. D. & Yu, Y.-T. Posttranscriptional RNA pseudouridylation. *The enzymes* **41**, 151-167 (2017).

Morais, P., Adachi, H. & Yu, Y.-T. The critical contribution of pseudouridine to mRNA COVID-19 vaccines. *Frontiers in cell and developmental biology* **9**, 3187 (2021).

Tavakoli, S. *et al.* Semi-quantitative detection of pseudouridine modifications and type I/II hypermodifications in human mRNAs using direct long-read sequencing. *Nature Communications* **14**, 334 (2023).

Khoddami, V. *et al.* Transcriptome-wide profiling of multiple RNA modifications simultaneously at single-base resolution. *Proceedings of the National Academy of Sciences* **116**, 6784-6789 (2019).

Dai, Q. *et al.* Quantitative sequencing using BID-seq uncovers abundant pseudouridines in mammalian mRNA at base resolution. *Nature Biotechnology* **41**, 344-354 (2023).

Zhang, M. *et al.* Quantitative profiling of pseudouridylation landscape in the human transcriptome. *Nature Chemical Biology*, 1-11 (2023).

Begik, O. *et al.* Quantitative profiling of pseudouridylation dynamics in native RNAs with nanopore sequencing. *Nature biotechnology* **39**, 1278-1291 (2021).

Huang, S. *et al.* Interferon inducible pseudouridine modification in human mRNA by quantitative nanopore profiling. *Genome biology* **22**, 1-14 (2021).

Cui, Q. *et al.* Targeting PUS7 suppresses tRNA pseudouridylation and glioblastoma tumorigenesis. *Nature cancer* **2**, 932-949 (2021).

Martinez, N. M., Schaening-Burgos, C. & Gilbert, W. V. in *Methods in enzymology* **658**, 277-310 (Elsevier, 2021).

Guegueniat, J. *et al.* The human pseudouridine synthase PUS7 recognizes RNA with an extended multi-domain binding surface. *Nucleic acids research* **49**, 11810-11822 (2021).

Sethi, A. J., Mateos, P. A., Hayashi, R., Shirokikh, N. & Eyras, E. R2Dtool: Positional interpretation of RNA-centric information in the context of transcriptomic and genomic features. *bioRxiv*, 2022.2009. 2023.509222 (2022).

Martinez, N. M. *et al.* Pseudouridine synthases modify human pre-mRNA co-transcriptionally and affect pre-mRNA processing. *Molecular Cell* **82**, 645-659. e649 (2022).

Dominissini, D. *et al.* Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. *Nature* **485**, 201-206 (2012).

Liu, J. *et al.* A METTL3–METTL14 complex mediates mammalian nuclear RNA N6-adenosine methylation. *Nature chemical biology* **10**, 93-95 (2014).

Ke, S. *et al.* A majority of m6A residues are in the last exons, allowing the potential for 3' UTR regulation. *Genes & development* **29**, 2037-2053 (2015).

Lorenz, D. A., Sathe, S., Einstein, J. M. & Yeo, G. W. Direct RNA sequencing enables m6A detection in endogenous transcript isoforms at base-specific resolution. *RNA* **26**, 19-28 (2020).

Pouliou, M. *et al.* MYCN Amplifications and Metabolic Rewiring in Neuroblastoma. *Cancers* **15**, 4803 (2023).

Dhanasekaran, R. *et al.* The MYC oncogene—the grand orchestrator of cancer growth and immune evasion. *Nature reviews Clinical oncology* **19**, 23-36 (2022).

Huang, T., Chen, W., Liu, J., Gu, N. & Zhang, R. Genome-wide identification of mRNA 5-methylcytosine in mammals. *Nature structural & molecular biology* **26**, 380-388 (2019).

Kundu, A. *et al.* PRDM16 suppresses HIF-targeted gene expression in kidney cancer. *Journal of Experimental Medicine* **217**, e20191005 (2020).

Figure 1. Identification and quantification of Ψ based on NanoPsiPy analysis of nanopore sequencing data and U-to-C basecalling errors.

(A) Flow chart for using the NanoPsiPy pipeline to detect and quantify Ψ modifications based on nanopore direct RNA sequencing. **(B)** Integrated Genome Viewer (IGV) snapshots of aligned nanopore MALAT1 RNA from BE(2)-C shGFP and BE(2)-C shPUS7-33 cells. **(C)** U-to-C mismatch percentage for U5614 based on data from **(B).** U5614 is shown as red and Ψ as blue, correctly aligned bases are shown in gray. Genomic reference sequence is converted to the sense strand and shown as RNA for clarity. **(D)** IGV snapshots of aligned nanopore MALAT1 IVT RNA oligo treated with wildtype or mutant recombinant PUS7. **(E)** U-to-C mismatch percentage for U5614 based on data from **(D)**. **(F)** Comparison of basecalling features between the samples treated with wild-type or mutant recombinant PUS7. **(G-H)** Accuracy of base-calling using a pool of three different IVT RNA oligos. **(I)** Effect of different preceding bases on estimating U-to-C basecalling errors using a pool of three different IVT RNA oligos. **(J)** Average base quality scores with respect to the number of reads for IVT RNA samples. ****$P < 0.0001$.

Figure 2. Transcriptome-wide identification and measurement of Ψ sites in poly-A RNA from BE(2)-C cells using nanopore direct RNA sequencing and NanoPsiPy.

(A) Normalized counts of read lengths for BE(2)-C poly-A RNA (orange) versus HeLa IVT RNA (blue). **(B)** Scatter plot showing distribution of Ψ fraction of 517 Ψ sites shared by BE(2)-C and HeLa poly-A RNA. **(C)** Bar plot showing the proportional distribution of Ψ sites based on Ψ fractions (stoichiometries) for all Ψ sites identified in BE(2)-C poly-A RNA. **(D)** Bar plot showing the distribution of Ψ fractions in BE(2)-C poly-A RNA based on read coverages (reads). **(E)** Motif frequency of most represented and frequently detected Ψ sites with high confidence. **(F)** Motif frequency versus average Ψ fraction for all Ψ sites in BE(2)-C poly-A RNA. **(G)** Distribution of Ψ sites in mRNA regions (5′UTR, CDS and 3'UTR), long non-coding RNA, and retained intron. **(H)** MetaTranscript plot showing the proportion of Ψ along relative positions in the 5'UTR, CDS, and 3'UTR of BE(2)-C mRNA. **(I)** MetaJunction plot showing Ψ sites relative to their distances to the closest downstream (left) and upstream (right) exon-exon junction.

Figure 3. Transcriptome-wide identification and measurement of PUS7-dependent Ψ sites in BE(2)-C cells.

(A) Normalized counts of read lengths for direct BE(2)-C shGFP (orange) versus BE(2)-C shPUS7-33 (blue) poly-A RNA. **(B)** Scatter plot showing that PUS7 knockdown reduced the Ψ strength of Ψ-modified genes, with labeled representative genes (MANF, NDUFS5, CBX3, RPS27A, and HMGB1). **(C)** Motif frequency of the most represented and frequently detected PUS7-dependent Ψ sites with high confidence. **(D)** Distribution of PUS7-dependent Ψ sites in mRNA regions, long non-coding RNA, and retained intron. **(E)** MetaTranscript plot showing the proportion of PUS7-dependent Ψ along relative positions in the 5'UTR, CDS, and 3'UTR of mRNA. **(F)** MetaJunction plot showing PUS7-dependent Ψ sites relative to their distances to the closest downstream (left) and upstream (right) exon-exon junction. **(G)** GO analysis of the 1,246 PUS7 target mRNAs. CC, cellular component; BP, biological process; MF, molecular function.

Supplement Figure 1. Immunoblotting of PUS7 in BE(2)-C cells expressing shGFP or shPUS7-33