Integrative analysis of nanopore direct RNA sequencing data reveals

a role of PUS7-dependent pseudouridylation in regulation of m^{6}A and

m^{5}C modifications

Mohit Bansal[1,2,*], Anirban Kundu[2,3,#], Anamika Gupta[1,2], Jane Ding[1,2], Andrew Gibson[4],

Sanjay Varma RudraRaju[5], Sunil Sudarshan[2,3] and Han-Fei Ding[1,2*]

[1]Department of Pathology and [2]O'Neal Comprehensive Cancer Center, [3]Department of

Urology, [4]Departments of Medicine, Heersink School of Medicine, University of Alabama

at Birmingham, AL 35294, USA, [5]Independent scholar, New York, NY 07850, USA.

[#]Current Address: Department of Urology, University of Arizona in Tucson, AZ, USA.

*Correspondence: M. Bansal mbansal@uabmc.edu and H.-F. Ding hding@uabmc.edu

Abstract

Abstract

Understanding the interactions between different RNA modifications is essential for

unraveling their biological functions. Here, we report NanoPsiPy, a computational

pipeline that employs nanopore direct RNA sequencing to identify pseudouridine (Ψ)

sites and quantify their levels at single-nucleotide resolution. We validated NanoPsiPy

by transcriptome-wide profiling of PUS7-dependent Ψ sites in poly-A RNA and rRNA.

NanoPsiPy leverages Ψ-induced U-to-C basecalling errors in nanopore sequencing

data, allowing detection of both low and high stoichiometric Ψ sites. We identified 8,624

PUS7-dependent Ψ sites in 1,246 mRNAs encoding proteins associated with ribosome

biogenesis, translation, and energy metabolism. Importantly, integrative analysis

revealed that PUS7 knockdown increases global mRNA N^6-methyladenosine (m^6A) and

5-methylcytosine (m^5C) levels, suggesting an antagonistic relationship between Ψ and

these modifications. Our study underscores the potential of nanopore direct RNA

sequencing in revealing the co-regulation of RNA modifications and the capacity of

NanoPsiPy in analyzing pseudouridylation and its impact on other RNA modifications.

Introduction

Research spanning several decades has demonstrated that RNAs undergo various

posttranscriptional modifications, including pseudouridylation[1-3]. Pseudouridine (Ψ) is

the first identified RNA modification[4] that constitutes 0.2–0.6% of total uridines in

mammalian mRNAs[5]. Ψ can enhance the stability and rigidity of RNAs, influencing

cellular processes such as translation, splicing, and telomere maintenance[2,6]. The

existence of Ψ in mRNAs remained unknown until 2014-2015[5,7,8], a revelation facilitated

by advancements in next-generation sequencing technologies. Human genome

encodes thirteen annotated pseudouridylation installation enzymes (pseudouridine

synthases, PUSs)[1], highlighting the robust and dynamic nature of Ψ installation in

human RNA. The conversion of uridine to Ψ starts with breaking the N1-C1' bond

linking the uracil base to the sugar, followed by an 180^o base rotation around the N3-C6

axis. The resulting Ψ contains an extra hydrogen bond donor at N1 and a C5-C1' base-

sugar linkage[9]. The success of mRNA-based COVID-19 vaccines in which all uridines

are replaced by Ψs underscores the crucial role played by Ψ modification in promoting

protein expression and increasing the efficacy of mRNA vaccines, showcasing the

importance of understanding and harnessing RNA modifications for therapeutic

purposes[10]. Ψ does not disrupt canonical base pairing, making it indistinguishable from

uridine in hybridization-based methods. Additionally, the molecular weight of Ψ matches

that of uridine, complicating direct detection through mass spectrometry[11]. This difficulty

is exacerbated when trying to identify Ψ sites within specific mRNA sequences.

Various techniques, such as Pseudo-Seq[7], Ψ -Seq[8], and Ceu-Seq[5] that are

based on illumina-short read sequencing, have used the CMC (N-cyclohexyl-N'-β-(4-

methylmorpholinium)ethylcarbodiimide)-based approach for transcriptome-wide Ψ

mapping. While CMC-based methods are effective in generating a stop signature during

reverse transcription (RT) for Ψ site identification, these methods lack base resolution

and stoichiometry information at the modified bases. In contrast to aforementioned

techniques, a new generation of methods based on bisulfite-induced deletion signatures

at Ψ sites, including RBS-seq[12], BID-seq[13], and PRAISE[14], offer quantitative

transcriptome-wide mapping of Ψ sites. However, it is important to note that Ψ-bisulfite

adduct formation involves harsh chemical reactions as an intermediate step. Moreover,

the read-length limitation of Illumina sequencing also narrows the possibility to examine

Ψ distribution in mRNA splice isoforms and the linkage of multiple Ψ sites in single RNA

molecules.

We wanted to develop a nanopore direct RNA sequencing method for Ψ site

identification and stoichiometry measurement on long native mRNAs without reliance on

harsh chemical reactions, providing a more direct and precise means of studying

pseudouridylation. Several studies have utilized nanopore-based direct RNA

sequencing to directly examine RNA modifications. The detection of Ψ using direct RNA

sequencing has been confirmed for ribosomal RNAs (rRNAs) in Saccharomyces

cerevisiae[15] and human mRNA[16,11], based on U-to-C basecalling errors. However, it

remains to be established whether direct RNA long-read nanopore sequencing is

sufficiently robust to determine PUS-dependent regulation of Ψ location and

stoichiometry in mRNA. This study aimed to develop a method to validate the U-to-C

basecalling signature as a distinctive feature of Ψ and to quantify transcriptome-wide

PUS7-dependent Ψ sites using in vivo and in vitro systems. Furthermore, cross talk

between RNA modifications can significantly impact mRNA expression and translation,

highlighting the intricate regulatory networks within cells. Consequently, we investigated

the interplay between Ψ, N^6-methyladenosine (m6A), and 5-methylcytosine (m5C) to

better understand their interactions.

Results

Development of NanoPsiPy for identification and quantification of Ψ modification

based on U-to-C basecalling errors in nanopore sequencing

We have developed NanoPsiPy, a computational pipeline

(https://github.com/vetmohit89/NanoPsiPy), to identify and quantify Ψ levels in RNA

based on U-to-C basecalling errors in Nanopore direct RNA sequencing data (Figure

1A). Following the alignment of RNA reads to the human reference sequence,

NanoPsiPy systematically gathers positional data for each uridine base. Subsequently,

it conducts statistical comparisons of U-to-C basecalling errors between two sample

groups (control and treatment), identifying and quantifying Ψ modifications at specific

locations. As Ψ detection by Nanopore direct RNA sequencing is based on basecalling

errors, it is essential to develop a method that accounts only primary alignment of each

RNA read (i.e., the alignment with the highest mapping quality) to reduce false positive

errors. NanoPsiPy filters out secondary alignments and only uses reads with mapping

quality score of 20 or higher, which correspond to the probability of correct alignment of

99% or higher, for downstream analysis and calculating U-to-C basecalling errors.

Another potential source of error is the presence of single-nucleotide polymorphisms

(SNPs) whereby the base is different from the reference genome or transcriptome

sequence. SNPs can cause systematic basecalling errors. NanoPsiPy removes any

SNP errors from downstream analysis by comparative analysis of the U-to-C mismatch

at each site between control and treatment samples (Figure 1A).

We tested the ability of our nanopore direct RNA sequencing and the NanoPsiPy

pipeline to identify and quantify Ψ based on U-to-C basecalling errors using two PUS7-

dependent systems. We used lentiviruses expressing a shRNA against human PUS7

(shPUS7-33) to silence PUS7 expression in the *MYCN*-amplified neuroblastoma cell

line BE(2)-C, with shGFP as control (Supplemental Figure 1). It has been shown

previously that PUS7 pseudouridylates U5614 in MALAT1 RNA (base position

according to NR-002819.4)[17]. In control shGFP BE(2)-C cells, MALAT1 RNA at U5614

showed 75% of RNA reads having a U-to-C basecalling error, which was reduced to

10% in BE(2)-C cells with PUS7 knockdown (Figures 1B-C). To further validate this U-

to-C basecalling error as a signature of PUS7-dependent Ψ modification, we performed

an in vitro pseudouridylation assay[18] using an in vitro transcribed (IVT) MALAT1 RNA

oligo that includes 86 nucleotides upstream and 66 nucleotides downstream of U5614.

The RNA oligo was incubated either with immunopurified wild-type human PUS7 or with

a catalytically inactive mutant (PUS7 D294A)[19] and analyzed systematically for

basecalling features, including mismatches, deletion, insertion and per-base qualities.

Approximately 38% of the IVT MALAT1 RNA reads treated with wild-type PUS7 showed

a U-to-C basecalling error compared to the 1% error in the RNA reads treated with the

mutant PUS7 (Figures 1D-E). We found no significant differences in other basecalling

features between the two samples (Figure 1F). These findings confirmed that

NanoPsiPy analysis of nanopore direct RNA sequencing data was able to detect and

quantify Ψ in RNA by identifying U-to-C basecalling errors.

Nanopore sequencing detects nucleotide bases from their ionic current signature.

Our pipeline NanoPsiPy uses Guppy (v6.4.2) for basecalling. By using a pool of

different IVT RNA oligos containing only canonical RNA nucleotides, we calculated the

basecalling accuracy. We found that our pipeline is highly accurate, with 95.99% for

adenine, 97.22% for cytosine, 97.26% for guanine, and 94.51% for uridine (Figure 1G).

Additionally, we found that Guppy could reliably identify unmodified and aligned uridine

with an average background U-to-C calling error of 2.99% (Figure 1H).

Nanopore direct RNA sequencing method sequences RNA in the 3' to 5'

direction, raising the question of whether U-to-C error-based Ψ detection is impacted by

the preceding 3' base. Our analysis of IVT RNA data revealed that when the preceding

3' base is U, A, C, or G, there is a background error of 4.8%, 3.07%, 2.07%, or 1.18% in

identifying the candidate U as C, respectively (Figure 1I). Further, we observed that the

average base quality is low when RNA reads at any location are less than 4 (Figure J).

Therefore, to minimize the inclusion of U sites with background errors, we defined a

threshold of 10% basecalling error for U-to-C estimation and a minimum of 9 reads at

any given U site for Ψ detection and quantification in the whole transcriptome.

Transcriptome-wide identification and quantification of Ψ sites in poly-A RNA

from BE(2)-C cells using nanopore direct RNA sequencing and NanoPsiPy

We applied nanopore direct RNA sequencing and NanoPsiPy for de novo detection and

quantification of transcriptome-wide Ψ modifications in the *MYCN*-amplified

neuroblastoma cell line BE(2)-C. For background control (RNA without modifications),

we used the published nanopore direct RNA sequencing data from a HeLa IVT poly-A

RNA library that was constructed first by reverse transcription of HeLa poly-A RNAs

(with modifications) into cDNAs to remove all modifications and then by in vitro

transcription of the cDNAs back into RNAs[11]. We conducted nanopore direct

sequencing of poly-A RNA from BE(2)-C shGFP cells and compared the sequencing

data with the HeLa IVT poly-A RNA nanopore sequencing data, which showed an

overlapping distribution of read lengths (Figure 2A). We then mapped RNA sequencing

reads from both datasets to the human genome reference sequence (GRCh38.p14) to

identify all the uridine sites common between BE(2)-C and HeLa IVT RNA and assigned

the U-to-C basecalling error as Ψ modification at each site. We defined Ψ sites in

BE(2)-C poly-A RNA based on two criteria: 1) the U-to-C basecalling error at each site

in BE(2)-C must exceed 10% and 2) the p value of U-to-C basecalling error must be <

0.01. Using data from more than 2 million uridines in BE(2)-C poly-A RNA and more

than 3 million uridines in HeLa IVT RNA, we identified 39,756 Ψ sites in BE(2)-C poly-A

RNA with a median Ψ level of 16.67% (Supplemental Table 1). To assess the accuracy

of our NanoPsiPy pipeline in detecting Ψ sites, we compared the identified BE(2)-C Ψ

sites with the previously annotated 1,691 Ψ sites in HeLa by nanopore direct RNA

sequencing[11]. Our analysis revealed 517 Ψ sites shared between BE(2)-C and HeLa

cells (Supplemental Table 2). Furthermore, we observed a strong correlation (R^2 =

0.618) in the Ψ fraction across these shared Ψ sites (Figure 2B).

Next, we examined the stoichiometric distribution of Ψ levels in BE(2)-C poly-A

RNA. Our analysis revealed that approximately 66% of the Ψ sites were modified at the

levels of 10-20% and ~11% at the levels of more than 30% (Figure 2C). In addition, our

analysis revealed a negative correlation between the levels of Ψ modification and the

levels of sequencing coverage (reads) at specific sites: the lower the number of reads,

the higher the median Ψ levels (Figure 2D).

We next analyzed the sequence contexts for the BE(2)-C Ψ sites and found that

the TRUB1 motifs GUUCA, GUUCC, and GUUCU, and the PUS7 motif UGUAG were

the most highly represented (Figure 2E). The median level of Ψ across the top 14 motifs

was 18.26% (Figure 2E). For all the Ψ sites, uridine exhibits a strong preference at the -

2, -1, and +1 positions (Figure 2F). The most frequent k-mers are UUUUG (4.92%),

AUUUG (3.78%), UUUUA (3.56%), CUUUG (3.64%), and UUUUU (3.43%) (Figure 2F).

To determine the distribution pattern of the identified Ψ sites in poly-A RNA, we

mapped the RNA reads to the transcriptome reference sequence

(gencode.v43.transcripts.fa) and observed that within mRNA, Ψ sites were highly

enriched in the coding sequence (CDS) and 3′ untranslated region (3'UTR) (Figure 2G),

as reported previously using CeU-Seq[5], Bid-seq[13] and Nanopore sequencing

methods[11,16]. In addition, we found low levels of Ψ distribution in retained introns

(4.46%) and long non-coding RNA (1.14%) (Figure 2G). The limited detection of Ψ sites

in the 5′ UTR could be due to a lower level of coverage of the 5′ UTR compared to CDS

and 3'UTR as nanopore sequencing moves in the 3' to 5' direction. To address this

issue, we generated a metatranscript plot using R2D tools[20] to compare the proportion

of significant Ψ sites along mRNA transcripts, which showed comparable Ψ density

distribution in the 5'UTR, CDS and 3'UTR (Figure 2H).

Martinej et al.[21] recently reported that Ψs are enriched in proximal introns (within

500 nucleotides of splice sites). In agreement with the report, we found that the Ψ sites

in BE(2)-C poly-A RNA were enriched around 200 nucleotides from the exon-exon

junctions (Figure 2I).

Transcriptome-wide identification and quantification of PUS7-dependent Ψ sites

in poly-A RNA

We applied our method to map and quantify PUS7-dependent Ψ sites in BE(2)-C

transcriptome. We isolated poly-A RNA from control (shGFP) and PUS7 knockdown

(shPUS7-33) BE(2)-C cells (Supplemental Figure 1) and prepared libraries for direct

RNA sequencing. Two independent shGFP RNA libraries produced a total of ~2.254

million reads, with mean read quality of 9.9 and an average N50 read length of 1,124

bases (defined as the shortest read length needed to cover 50% of the sequenced

nucleotides). Two independent shPUS7 RNA libraries produced a total of 1.44 million

reads, with mean read quality of 10.0 and an average N50 read length of 1,351 bases.

The median RNA read lengths were 702 and 789 bases for shGFP and shPUS7-33

samples, respectively (Figure 3A and Supplemental Table 3).

To identify PUS7-dependent Ψ sites, we compared U-to-C basecalling errors at

uridine sites in shPUS7 poly-A RNA with Ψ sites in shGFP poly-A RNA that were

identified as described above (shGFP BE(2)-C vs. HeLa IVT). Employing the two

controls (shGFP BE(2)-C and HeLa IVT), we aimed to enhance the precision of our

analysis and eliminate potential false positives in calling PUS7-dependent Ψ sites. We

identified 8,624 putative PUS7-dependent Ψ sites within protein-coding mRNA, with Ψ

stoichiometries ranging from 10.00% to 88.88% and a median Ψ stoichiometry of

16.78%. Additionally, we identified 294 putative PUS7-dependent Ψ sites within

retained introns, with Ψ stoichiometries spanning 10.22% to 68.42% and a median Ψ

stoichiometry of 18.08%, and 58 putative PUS7 Ψ sites in long non-coding RNA,

exhibiting Ψ stoichiometries between 10.28% and 52% and a median Ψ stoichiometry of

18.18% (Supplemental Table 4). Interestingly, single-read analysis revealed a subset of

BE(2)-C mRNAs contain multiple PUS7-dependent Ψ sites within each mRNA

molecule. We quantified the overall Ψ modification level in each gene by calculating its

Ψ-strength (sum of Ψ fractions at all the Ψ sites within a single gene)[13]. PUS7

knockdown resulted in a global reduction in Ψ-strength of PUS7 target mRNAs (Figure

3B).

Analysis of k-mer frequency and representation for the PUS7-dependent Ψ sites

revealed that the previously reported HeLa[11] Ψ k-mers UUUAU, AUUUA and UUUAA

were the most represented, whereas the Ψ k-mers UUUUG, AUUUG and CUUUG were

the most frequently detected. The median Ψ stoichiometry of the top 14 k-mers was

19.79% while the median stoichiometry of transcriptome-wide PUS7-dependent Ψ sites

was 16.78% (Figure 3C).

As presented above for transcriptome-wide Ψ sites in shGFP BE(2)-C mRNAs

(Figure 2G), PUS7-dependent Ψ sites also showed a similar distribution pattern, with

~48.66% of the Ψ sites in the 3′UTR, 44.98% in the CDS, and 2.43% in the 5′UTR

(Figure 3D). Again, the metatranscript plot for PUS7-dependent Ψ sites showed

comparable density enrichment in the 5' UTR, CDS, and 3'UTR (Figure 3E). We also

found that PUS7-dependent Ψ sites were enriched around 200 nucleotide proximal from

the exon-exon junctions (Figure 3F).

We identified 8,624 putative PUS7-dependent Ψ sites in 1,246 unique protein

coding genes. To interrogate the functions of these genes, we performed Gene

Ontology (GO) analysis, which revealed the PUS7 target genes are involved in

ribosome biogenesis, translation, RNA and protein binding, and mitochondrial energy

metabolism (Figure 3G and Supplemental Table 5). Further, we identified 57 Ψ sites in

13 unique long non-coding RNAs. GO analysis found highly significant enrichment of

GO terms corresponding to RNA processing (Supplemental Table 6).

Comparative analysis of rRNA pseudouridylation in HeLa and BE(2)-C cells

Marchand et al. have identified 104 Ψ sites in HeLa rRNA using HydraPsiSeq[22]. To

further validate our nanopore sequencing method, we performed nanopore sequencing

of rRNA libraries from BE(2)-C cells and obtained 367,424 reads from two BE(2)-C

shGFP replicates and 177,515 reads from two BE(2)-C shPUS7-33 replicates. We

mapped the U-to-C basecalling error of each uridine base in 5.8S, 18S, and 28S rRNAs

and highlighted the previously annotated Ψ sites from HeLa[22] (Figure 4A-C). We

calculated the U-to-C basecalling error for all the 1,510 rRNA uridine bases using direct

RNA Nanopore sequencing. Among them, 1,250 bases showed less than 10% U-to-C

basecalling errors and were therefore excluded from the analysis (Supplemental Table

8). Out of the remaining 260 uridine bases that showed greater than 10% U-to-C

basecalling errors, we identified 97 uridine bases (93.3% of the 104 uridine bases) that

were previously annotated as Ψ sites in HeLa rRNA using HydraPsiSeq[22]

(Supplemental Table 9). The remaining 7 bases (6.7%) with less than 10% U-to-C

basecalling errors in BE(2)-C could be due to cell-type specificity. This result

underscores the high sensitivity of nanopore direct RNA sequencing in detecting

previously annotated Ψ sites. Notably, we did not observe any enrichment of specific k-

mers associated with these 97 Ψ sites in rRNA (Figure 4D).

Additionally, we compared our BE(2)-C rRNA pseudouridylation data with the

HeLa data that were also generated using nanopore direct RNA sequencing[11]. Among

the 103 common Ψ sites identified in the 5.8S and 18S rRNAs of both HeLa and BE(2)-

C cells, 65 sites (63.1%) exhibited more than 10% U-to-C base-calling errors in both cell

lines, indicating a high degree of conservation for these Ψ sites across cell lines. The

median Ψ levels across all Ψ sites in the 5.8S and 18S rRNAs were 25.87% for HeLa

and 18.77% for BE(2)-C, with a correlation coefficient of 0.88 (Figure 4E-G).

(Supplemental Table 10). Also, we found that the UCUGG k-mer is the most

represented k-mer among rRNA Ψ sites identified in both HeLa[11] and BE(2)-C cells

(Figure 4H).

Finally, we identified 15 Ψ sites where the U-to-C basecalling error was

significantly reduced after PUS7 knockdown. Notably, 8 of these 15 sites were

previously identified as Ψ sites in HeLa rRNA (Figure 4I and Supplemental Table 11).

Collectively, our rRNA Ψ analysis demonstrates that nanopore direct RNA

sequencing can reliably identify previously annotated Ψ sites.

PUS7-dependent pseudouridylation represses m^{6}A modification in mRNA

Huang et al. reported recently that transcripts with higher numbers of Ψ sites have

fewer m6A sites, while those with more m^{6}A sites exhibited reduced levels of Ψ

modification, suggesting an antagonistic relationship between Ψ and m6A in mRNA[23]. In

agreement with the finding, our dot blot analysis showed a 1.6-fold increase in m^6A

levels following PUS7 knockdown (Figure 5A). To further investigate the impact of

PUS7-dependent pseudouridylation on m^6A modification, we analyzed the same

nanopore sequencing data for m^6A levels in poly-A RNA from BE(2)-C shGFP and

shPUS7 cells using two independent pipelines, CHEUI[20] and m6anet[24]. We divided the

PUS7-dependent Ψ data into ten fractions, ranging from 0% to 100% in 10%

pseudouridylation stoichiometry increments, which revealed a significant reduction in Ψ

levels across all fractions following PUS7 knockdown (Figure 5B and Supplemental

Table 12). The CHEUI and m6anet tools identified 4,537 and 1,223 m6A sites within

m^6A DRACH motifs, respectively, in BE(2)-C shGFP and shPUS7 mRNA samples. For

the m6anet analysis, we used a probability threshold of 0.9 to select m^6A sites, ensuring

high confidence in the identified sites. Both analyses showed a significant increase in

the numbers of m^6A sites in higher m^6A stoichiometry fractions after PUS7 knockdown

(Figure 5C-D and Supplemental Table 13-14), providing further evidence that PUS7

knockdown increases the global m^6A levels in mRNA.

Next, we examined if the antagonistic relationship between m^6A and Ψ

modifications in mRNAs is specific to certain DRACH motifs. We compared m^6A

stoichiometry between BE(2)-C shGFP and shPUS7 across the seven most common

DRACH m^6A motifs and observed an average increase of 2.60 ± 0.74-fold in m^6A

stoichiometry levels following PUS7 knockdown (Figure 5E).

We further investigated whether this antagonistic relationship between m^6A and

Ψ persists when they co-occur in the same transcripts. We identified a total of 104 m^6A

and Ψ sites on 45 unique mRNA transcripts containing any of the three previously

verified Ψ k-mers (UGUAG, GUUCA, and GUUCC) and m^6A DRACH motifs and then

analyzed m^6A levels using CHEUI. Consistently, we found that as Ψ levels decreased,

m^6A levels increased (Figure 5F-G and Supplemental Table 15).

Collectively, the data presented above suggest that PUS7-dependent

pseudouridylation has a significant role in regulation of m^6A modification in mRNA.

PUS7-dependent pseudouridylation represses m^5C modification in mRNA

Considering above findings on m^6A, we investigated the relationship between m^5C and

Ψ modifications in mRNA. Dot blot analysis revealed a 2.3-fold increase in m^5C levels

following PUS7 knockdown (Figure 6A). Using CHEUI, we identified 37,276 m^5C sites in

BE(2)-C shGFP and shPUS7 poly-A RNA (Supplemental Table 16). Our analysis was

focused on the 3,014 m^5C sites linked to the most frequently occurring k-mers: GACUG

(2.3%), GACUU (2.2%), GACUC (1.8%), GACAC (1.0%), and GACAU (0.9%)

(Supplemental Table 17). We measured the levels of m^5C in 10% gradient fractions of

control and PUS7 knockdown samples, which revealed that PUS7 knockdown

increased the numbers of m^5C sites in poly-A RNAs with higher m^5C stoichiometries

(Figure 6B and Supplemental Table 17). Also, we found that PUS7 knockdown resulted

in a significant increase in m^5C stoichiometry levels in these k-mers (Figure 6C).

We further investigated if this antagonistic relationship between Ψ and m^5C

persists when they co-occur on the same transcripts. To maximize accuracy, we

selected transcripts containing experimentally verified Ψ k-mers (UGUAG, GUUCA, and

GUUCC) and the 5 most frequent m^5C k-mers (GACUG, GACUU, GACUC, GACAC,

and GACAU). We identified a total of 97 m^5C and Ψ sites on 53 unique mRNA

transcripts that meet this criterion. PUS7 knockdown resulted in a 3.7-fold decrease in

Ψ levels in these transcripts (Figure 6D), which was accompanied by a significant

increase in m^5C levels, with the median stoichiometry increasing from 38.5% in shGFP

cells to 55.6% in PUS7 knockdown cells (Figure 6E, Supplemental Table 18).

Taken together, these data suggest that PUS7-dependent pseudouridylation has

a significant role in regulation of m^5C modification in mRNA.

Discussion

We have developed NanoPsiPy for processing nanopore direct RNA long-read

sequencing data to identify Ψ at individual RNA molecule and whole transcriptome

levels with location information and stoichiometry quantification. NanoPsiPy abstracts

U-to-C basecalling errors to detect and quantify transcriptome-wide Ψ modifications

between any two samples using comparison paradigm to minimize false positives. By

treating in vitro RNA oligos with wild-type and mutant (D294A) PUS7, our study

demonstrated that U-to-C basecalling is the sole characteristic enabling the

identification of Ψ in nanopore direct RNA sequencing data. NanoPsiPy weigh the

pooled sample analysis to identify transcriptome-wide Ψ, comparing the average of U-

to-C errors between two conditions to minimize false positives due to low coverage that

is an issue with direct mRNA sequencing.

Quantitative MS analysis estimated that the Ψ/U ratio is approximately 0.2-0.6%

in human mRNA[5], which is similar to the ratio of m^6A/A at 0.15-0.6% found in the human

transcriptome[25-27]. Using NanoPsiPy, we identified 39,756 Ψ sites in poly-A RNA from

the human neuroblastoma cell line BE(2)-C, which again is comparable to the 42,116

m[6]A sites detected in poly-A RNA from the human mammary epithelial cell line HMEC

using MINES analysis of nanopore direct RNA sequencing data[28]. Recently, Tavakoli et

al. reported the identification of 1,691 Ψ sites in HeLa poly-A RNA using PsiNanopore

that also relies on U-to-C basecalling errors in direct RNA sequencing [11]. Our analysis

identified 517 Ψ sites that are shared between BE(2)-C and HeLa cells, suggesting that

a common set of poly-A RNAs are targeted by pseudouridylation in human cancer cell

lines of different tissue origins. On the other hand, we identified significantly more Ψ

sites in BE(2)-C poly-A RNAs. We suspect that a major reason for the differences in Ψ

locations and numbers between HeLa and BE(2)-C cells is the cell-type specific

transcriptome profiles because modifications in highly expressed RNAs are most likely

to be detected. BE(2)-C cells are of neuronal origin with amplification of the MYCN

oncogene, a member of the MYC family of transcription factors that regulates RNA

expression of thousands of genes [29,30]. Thus, NanoPsiPy is highly effective in

comprehensively identifying Ψ sites and quantifying their levels in the whole

transcriptome.

It was reported recently that the median Ψ stoichiometry in HEK293T mRNA is

approximately 10%, with 80% of the Ψ sites showing Ψ levels < 20% and 10% showing

> 40% [14]. In agreement, our NanoPsiPy analysis revealed ~66% of the Ψ sites in BE(2)-

C poly-A RNA having modification levels at 10-20%, 23% at 20-30%, and 11% at >

30%. Interestingly, the median methylation level of m[5]C sites in mRNA is in the range of

15-18%. Across various tissues or cell types, the majority (62–70%) of m[5]C sites

exhibited low methylation levels (<20%), and 8-10% showed moderate to high

methylation levels (>40%)[31]. Collectively, these findings support the notion that RNA

modifications occur at relatively low stoichiometric levels across the entire

transcriptome.

Previous studies suggest that Ψ distribution in mRNAs is overrepresented in

CDS and 3′UTR and underrepresented in 5′UTR [11,13]. However, our analysis indicates

that Ψ is proportionally present in all regions, showing no significant bias. In principle,

this broad distribution endows Ψ modifications with the ability to influence many aspects

of mRNA metabolism and function. In addition, we identified 1,246 PUS7 target mRNAs

in the *MYCN*-amplified neuroblastoma cell line BE(2)-C and GO analysis suggest that

they are involved primarily in ribosome biogenesis, translation, and mitochondrial

energy metabolism. These findings lay the foundation for further investigation of the

functional significance of PUS7-dependent pseudouridylation in neuroblastoma and

other MYC-driven cancers.

A major finding of our study is that PUS7 knockdown markedly reduced Ψ levels

while increased m^6A and m^5C levels in the same transcripts, indicating a crucial role of

PUS7-dependent pseudouridylation in regulation of other RNA modifications. The

underlying molecular mechanism remains to be elucidated. Our findings support a

recent report of an antagonistic relationship between Ψ and m^6A [23] and provide direct

evidence that a decrease in Ψ levels can cause an increase in m^6A and m^5C levels.

In summary, we have developed a method for quantitative profiling of

transcriptome-wide Ψ sites, enabling the identification and quantification of PUS7-

dependent Ψ sites at single mRNA molecule resolution with functional implications. We

anticipate that NanoPsiPy will facilitate the identification and functional characterization

of pseudouridylation and aid in studying cross-talk between RNA modifications.

Material and methods

Cell culture and PUS7 knockdown

Neuroblastoma BE(2)-C) cells were cultured in DME/F-12 1:1 (HyClone SH30023)

supplemented with 10% Fetal Bovine Serum (Fisher Scientific FB12999102) and 1%

Penicillin-Streptomycin (Lonza 17602E). The PUS7 knockdown lentiviral construct

shPUS7-33 (TRCN0000061933) was obtained from Sigma-Aldrich. BE(2)-C cells were

infected with lentiviruses expressing shGFP or shPUS7-33, and PUS7 knockdown was

confirmed by immunoblotting using rabbit anti-PUS7 (1:1000, Thermo Fisher PA5-

54983).

Poly-A RNA isolation, library preparation, and nanopore sequencing

Total RNA was obtained through TRIzol extraction, and poly(A) mRNA was isolated

using the NEBNext Poly(A) mRNA Magnetic Isolation kit (NEB E7490L). The RNA

library for direct RNA sequencing was prepared from ~500 ng of poly-A RNA according

to the Oxford Nanopore technology (ONT) direct RNA sequencing protocol version

direct-rna-sequencing-sqk-rna002-DRS_9080_v2_revS_14Aug2019-minion. For

sequencing, the ONT FLO-MIN106D (R9.4.1) flow cell was primed according to the

manufacturer's protocol, and the elute was mixed with an RNA running buffer (RRB)

and loaded onto the flow cell. Sequencing was performed using MinKnow (version

19.12.5). Each flow cell was sequenced for 24-72 hours per each run, depending on

flowcell exhaustion rate. Two replicates from different passages and different flow cells

were used for each biological replicate. To study rRNA pseudouridine modifications, we

prepared a direct RNA sequencing library from total RNA, similar to the approach used

for poly-A mRNA sequencing. We first added a poly-A tail to the total RNA and then

sequenced it using Flongle flow cells.

PUS7 immunoprecipitation and purification

All immunoprecipitation steps were performed at 4°C following the protocol of Kundu et

al. with some modifications[32]. Briefly, 293FT cells expressing either wild-type (WT) or

mutant (D294A) PUS7 (myc-tagged) were lysed in the buffer containing 50 mM HEPES

pH 7.5, 130 mM NaCl, 0.3% NP-40, 1 mM EDTA, and 1× protease/phosphatase

inhibitor (Thermo Fisher 78440). Approximately 3 mg of protein lysates were mixed with

protein A/G-magnetic beads (Thermo Fisher 88802) and 2 μg of either anti-Myc or

control IgG antibody and rotated for 16 hours at 4°C. Subsequently, the magnetic beads

were pelleted and washed for three times in the buffer containing 20 mM Tris-HCl at pH

8.0, 130 mM NaCl, 0.3% NP-40, and 1 mM EDTA. Then myc-PUS7 (WT or D294A) was

eluted from the beads using myc-peptide (Protein Tech 2yp) following manufacture's

protocol. Finally, using 30 KDa ultracentrifugal filter, the eluted proteins were

concentrated, and buffer exchanged in the 1X pseudouridylation buffer (100 mM Tris-

HCL pH-8.0, 100 mM ammonium acerate, 5 mM MgCl2, 0.3 mM EDTA, and 130 mM

NaCl). Concentrated proteins were either used immediately or stored in -80°C until use.

In vitro RNA pseudouridylation[18]

The synthetic 192-nucleotide (nt) MALAT1 DNA (Supplemental Table 7) contains 86 nt

upstream and 66 nt downstream of the pseudouridine site (U5614, NR_002819.4). A

22-nt sequence containing the T7 promoter and a 25-nt adapter sequence were added

to the 5' and 3' ends, respectively. The oligo was obtained from IDT DNA and amplified

by PCR using Phusion Polymerase (NEB F630) and purified for in vitro transcription

using the MEGAshortscript T7 kit (Invitrogen AM1333). The RNA product was purified

using a urea gel, denatured, and refolded. In vitro pseudouridylation was carried out by

incubating 15-30 pmol of folded RNA with 1 µg of immunopurified PUS7 or PUS7

D294A. Poly-A tail was added using a Poly (A) tailing kit (Invitrogen AM1350). Library

was prepared and loaded into the FLO-MIN106D flow cell for sequencing using

MinKnow (version 19.12.5) as described above. Each flow cell was sequenced for 24 h.

m^6A and m^5C analyses

CHEUI[20] and m6anet[24] were employed to analyze the same nanopore direct RNA

sequencing data from BE(2)-C shGFP and shPUS7 cells to identify and quantify m^6A

levels in transcripts containing DRACH motifs. Likewise, CHEUI was used to analyze

the same nanopore data to identify and quantify m5C levels in transcripts. The FASTQ

files were aligned to the Gencode transcript reference (gencode.v43.transcripts.fa).

Similar to the Ψ analysis, two replicates from different passages and different flow cells

were used for each biological replicate.

Dot blot analysis

Dot blot was performed with either poly-A RNA or total RNA. The poly-A RNA was

isolated from total RNA using a NEB poly-A RNA isolation kit. RNA concentration was

measured using a Qubit (Thermo Fisher Scientific). RNA was dotted on a nitrocellulose

membrane (Bio-Rad). Once the membrane was dried, RNA was cross-linked to the

membrane using 120 mJ/cm2 density of UV light (UVP Hybrilinker HL-2000). The

membrane was washed once with 2× SSC solution (Thermo Fisher Scientific),

incubated with 1× Blot Stain Blue (MilliporeSigma) for 1-2 minutes, and then rinsed

several times with 1× TBST (Tris-buffered saline, pH 7.4, plus 0.1% Tween-20) to

minimize the background. After imaging of the blue RNA dots, the membrane was

blocked with 5% milk for 1 hour, followed by overnight incubation at 4°C with anti-m^{6}A

antibody (Cell Signaling; mAb #56593; 1:1,000 dilution) or anti-m^{5}C antibody

(Proteintech; Cat No. 68301-1; 1:000 dilution). After washing, the membrane was

incubated with HRP-conjugated secondary antibody for 1 hour (1:3,000 dilution) and the

image was captured using a ChemiDoc Imaging System (Bio-Rad).

Data analysis

All raw fast5 files generated from nanopore sequencing were uploaded to UAB Cheaha

cluster for further analysis. Multi-fast5 files were basecalled in real-time by Guppy

(6.4.2) using the high accuracy model with minimum quality score of 7. The fastq files

were analyzed using the NanoPsiPy where reads were aligned to the reference file

(GRCh38.p14 or gencode.v43.transcripts.fa.) using minimap2 (2.17). The mapped

reads were piled up to the human reference sequence by samtools (v1.12). The U-to-C

basecalling "error" data were extracted and compared between two conditions using

NanoPsiPy (http://github.com/vetmohit89/NanoPsiPy.git). NanoPsiPy tool was built

around Nanopore_psu tool available at

(https://github.com/sihaohuanguc/Nanopore_psU/)[16]. Finally, data was prepared by

applying condition when the ratio of C reads from the total of U+C reads of the control

group is greater than the ratio of C reads from the total of U+C reads for the treatment

group. After filtering the input data, the Chi-square test is applied to a contingency table

constructed from the read counts in different categories (control_C, control_U,

treatment_C, treatment_U). Yates' continuity correction is applied to improve the

accuracy of the results when the sample size is limited and $p < 0.01$ is defined as

significant. Gene ontology (GO) analysis was performed using gprofiler2

(https://biit.cs.ut.ee/gprofiler/gost).

Accession Numbers

All Fast5 files generated in this study have been publicly shared and are accessible

through the NIH NCBI SRA under the Bio-Project accession PRJNA961708

(SRR29662300, SRR29662301, SRR29662302 and SRR29662303). The data for two

HeLa IVT RNA samples were retrieved from NIH NCBI SRA with the accession

numbers SRR23932675 and SRR23929127.

Supplemental Tables description

Supplemental Table 1: Significant Ψ sites identified in BE(2)-C poly-A RNA.

Supplemental Table 2: Common Ψ sites identified in BE(2)-C and HeLa poly-A RNA.

Supplemental Table 3: Direct RNA Library description of BE(2)-C shGFP and shPUS7-

33.

Supplemental Table 4: Significant PUS7-dependent Ψ sites in protein coding, retained

RNA and long non-coding RNA of BE(2)-C cells.

Supplemental Table 5: Gene ontology analysis of protein coding mRNAs containing

PUS-dependent Ψ sites

Supplemental Table 6: Gene ontology analysis of long non-coding RNAs containing

PUS7-dependent Ψ sites.

Supplemental Table 7: RNA oligo sequences.

Supplemental Table 8: U-to-C basecalling errors as a proxy for pseudouridine from

BE(2)-C rRNA.

Supplemental Table 9: rRNA Ψ modifications (BE(2)-C vs. HeLa, Marchand et al.,

2020).

Supplemental Table 10: rRNA Ψ modifications (BE(2)-C vs. HeLa, Tavakoli et al.,

2023).

Supplemental Table 11: PUS7-dependent Ψ sites in BE(2)-C rRNA.

Supplemental Table 12: Distribution of Ψ Stoichiometry in poly-A RNA from BE(2)-C

shGFP and shPUS7-33 cells.

Supplemental Table 13: PUS7 knockdown on m^{6}A stoichiometry analyzed by CHEUI.

Supplemental Table 14: PUS7 knockdown on m^{6}A stoichiometry analyzed by m6anet.

Supplemental Table 15: Co-occurrence of PUS7 dependent Ψ and m^{6}A stoichiometry

in the same transcripts from BE(2)-C cells.

Supplemental Table 16: m^{5}C sites in poly-A RNA from BE(2)-C shGFP and shPUS7

cells identified by CHEUI.

Supplemental Table 17: PUS7 knockdown on m^{5}C stoichiometry within the most

frequent m^{5}C k-mers.

Supplemental Table 18: Co-occurrence of PUS7-dependent Ψ and m^{5}C stoichiometry

in the same transcripts from BE(2)-C Cells.

Funding

This work is supported by the National Institutes of Health grant R01CA190429 to H.-

F.D.

Acknowledgment section

AK was partly supported by a Concept Award (Award HT9425-23-1-0339) and an Early

Career Award (HT9425-23-1-0783) from the Department of Defense.

Author contributions

M.B. and H-F.D. conceived the study and designed the experiments with contributions

from A.K., A.Gibson, A.Gupta, J.D., and S.S. M.B. performed nanopore RNA

sequencing and data analysis with assistance from A.Gibson and S.V.R. A.K.

performed PUS7 purification. A.Gupta. and J.D. performed cell culture and

immunoblotting. M.B. and H-F.D. wrote the paper with contributions from S.S. and J.D.

H-F.D. supervised and provided funding for the project. All authors read the manuscript

and approved its contents.

References:

Li, X., Ma, S. & Yi, C. Pseudouridine: the fifth RNA nucleotide with renewed interests. *Current Opinion in Chemical Biology* **33**, 108-116 (2016).

Borchardt, E. K., Martinez, N. M. & Gilbert, W. V. Regulation and function of RNA pseudouridylation in human cells. *Annual review of genetics* **54**, 309-336 (2020).

Rodell, R., Robalin, N. & Martinez, N. M. Why U matters: detection and functions of pseudouridine modifications in mRNAs. *Trends in Biochemical Sciences* **49**, 12-27 (2024).

Davis, F. F. & Allen, F. W. Ribonucleic acids from yeast which contain a fifth nucleotide. *J Biol Chem* **227**, 907-915 (1957).

Li, X. *et al.* Chemical pulldown reveals dynamic pseudouridylation of the mammalian transcriptome. *Nature chemical biology* **11**, 592-597 (2015).

Garus, A. & Autexier, C. Dyskerin: an essential pseudouridine synthase with multifaceted roles in ribosome biogenesis, splicing, and telomere maintenance. *Rna* **27**, 1441-1458 (2021). https://doi.org/10.1261/rna.078953.121

Carlile, T. M. *et al.* Pseudouridine profiling reveals regulated mRNA pseudouridylation in yeast and human cells. *Nature* **515**, 143-146 (2014).

Schwartz, S. *et al.* Transcriptome-wide mapping reveals widespread dynamic-regulated pseudouridylation of ncRNA and mRNA. *Cell* **159**, 148-162 (2014).

De Zoysa, M. D. & Yu, Y.-T. Posttranscriptional RNA pseudouridylation. *The enzymes* **41**, 151-167 (2017).

10   Morais, P., Adachi, H. & Yu, Y.-T. The critical contribution of pseudouridine to

mRNA COVID-19 vaccines. *Frontiers in cell and developmental biology* **9**, 3187

(2021).

11   Tavakoli, S. *et al.* Semi-quantitative detection of pseudouridine modifications and

type I/II hypermodifications in human mRNAs using direct long-read sequencing.

Nature Communications **14**, 334 (2023).

12   Khoddami, V. *et al.* Transcriptome-wide profiling of multiple RNA modifications

simultaneously at single-base resolution. *Proceedings of the National Academy*

of Sciences **116**, 6784-6789 (2019).

13   Dai, Q. *et al.* Quantitative sequencing using BID-seq uncovers abundant

pseudouridines in mammalian mRNA at base resolution. *Nature Biotechnology*

41, 344-354 (2023).

14   Zhang, M. *et al.* Quantitative profiling of pseudouridylation landscape in the

human transcriptome. *Nature Chemical Biology*, 1-11 (2023).

15   Begik, O. *et al.* Quantitative profiling of pseudouridylation dynamics in native

RNAs with nanopore sequencing. *Nature biotechnology* **39**, 1278-1291 (2021).

16   Huang, S. *et al.* Interferon inducible pseudouridine modification in human mRNA

by quantitative nanopore profiling. *Genome biology* **22**, 1-14 (2021).

17   Cui, Q. *et al.* Targeting PUS7 suppresses tRNA pseudouridylation and

glioblastoma tumorigenesis. *Nature cancer* **2**, 932-949 (2021).

18   Martinez, N. M., Schaening-Burgos, C. & Gilbert, W. V. in *Methods in*

enzymology Vol. 658   277-310 (Elsevier, 2021).

19   Guegueniat, J. *et al.* The human pseudouridine synthase PUS7 recognizes RNA

with an extended multi-domain binding surface. *Nucleic Acids Research* **49**,

11810-11822 (2021). https://doi.org/10.1093/nar/gkab934

20   Sethi, A. J., Mateos, P. A., Hayashi, R., Shirokikh, N. & Eyras, E. R2Dtool:

Positional interpretation of RNA-centric information in the context of

transcriptomic and genomic features. *bioRxiv*, 2022.2009. 2023.509222 (2022).

21   Martinez, N. M. *et al.* Pseudouridine synthases modify human pre-mRNA co-

transcriptionally and affect pre-mRNA processing. *Molecular Cell* **82**, 645-659.

e649 (2022).

22   Marchand, V. *et al.* HydraPsiSeq: a method for systematic and quantitative

mapping of pseudouridines in RNA. *Nucleic acids research* **48**, e110-e110

(2020).

23   Huang, S., Wylder, A. C. & Pan, T. Simultaneous nanopore profiling of mRNA

m6A and pseudouridine reveals translation coordination. *Nature Biotechnology*,

1-5 (2024).

24   Hendra, C. *et al.* Detection of m6A from direct RNA sequencing using a multiple

instance learning framework. *Nature methods* **19**, 1590-1598 (2022).

25   Dominissini, D. *et al.* Topology of the human and mouse m6A RNA methylomes

revealed by m6A-seq. *Nature* **485**, 201-206 (2012).

26   Liu, J. *et al.* A METTL3–METTL14 complex mediates mammalian nuclear RNA

N6-adenosine methylation. *Nature chemical biology* **10**, 93-95 (2014).

27   Ke, S. *et al.* A majority of m6A residues are in the last exons, allowing the

potential for 3′ UTR regulation. *Genes & development* **29**, 2037-2053 (2015).

28   Lorenz, D. A., Sathe, S., Einstein, J. M. & Yeo, G. W. Direct RNA sequencing

enables m6A detection in endogenous transcript isoforms at base-specific

resolution. *RNA* **26**, 19-28 (2020).

29   Pouliou, M. *et al.* MYCN Amplifications and Metabolic Rewiring in

Neuroblastoma. *Cancers* **15**, 4803 (2023).

30   Dhanasekaran, R. *et al.* The MYC oncogene—the grand orchestrator of cancer

growth and immune evasion. *Nature reviews Clinical oncology* **19**, 23-36 (2022).

31   Huang, T., Chen, W., Liu, J., Gu, N. & Zhang, R. Genome-wide identification of

mRNA 5-methylcytosine in mammals. *Nature Structural & Molecular Biology* **26**,

380-388 (2019). https://doi.org/10.1038/s41594-019-0218-x

32   Kundu, A. *et al.* PRDM16 suppresses HIF-targeted gene expression in kidney

cancer. *Journal of Experimental Medicine* **217**, e20191005 (2020).

Figure 1. Identification and quantification of Ψ based on NanoPsiPy analysis of nanopore sequencing data and U-to-C basecalling errors.

(A) Flow chart for using the NanoPsiPy pipeline to detect and quantify Ψ based on nanopore direct RNA sequencing. **(B)** Integrated Genome Viewer (IGV) snapshots of aligned nanopore MALAT1 RNA from BE(2)-C shGFP and BE(2)-C shPUS7-33 cells. **(C)** U-to-C mismatch percentage for U5614 based on data from **(B)**. U5614 is shown as red and Ψ as blue, and correctly aligned bases are shown in gray. Genomic reference sequence is converted to the sense strand and shown as RNA for clarity. **(D)** IGV snapshots of aligned nanopore MALAT1 IVT RNA oligo treated with immunopurified wildtype or mutant PUS7. **(E)** U-to-C mismatch percentage for U5614 based on data from **(D)**. **(F)** Comparison of basecalling features between the samples treated with immunopurified wild-type or mutant PUS7. **(G-H)** Accuracy of base-calling using a pool of three different IVT RNA oligos. **(I)** Effect of different preceding bases on estimating U-to-C basecalling errors using a pool of three different IVT RNA oligos. **(J)** Average base quality scores with respect to the number of reads for IVT RNA samples. ****$P < 0.0001$.

Figure 2. Transcriptome-wide identification and quantification of Ψ sites in poly-A

RNA from BE(2)-C cells using nanopore direct RNA sequencing and NanoPsiPy.

(A) Normalized counts of read lengths for BE(2)-C poly-A RNA (orange) versus HeLa

IVT RNA (blue). (**B**) Scatter plot showing distribution of Ψ fractions (stoichiometries) of

517 Ψ sites shared by BE(2)-C and HeLa poly-A RNA. **(C)** Bar plot showing the

proportional distribution of Ψ sites based on Ψ fractions (stoichiometries) for all Ψ sites

identified in BE(2)-C poly-A RNA. **(D)** Bar plot showing the distribution of Ψ fractions in

BE(2)-C poly-A RNA based on read coverages (reads). **(E)** K-mers frequency of most

represented and frequently detected Ψ sites with high confidence. **(F)** K-mers frequency

versus average Ψ fraction for all Ψ sites in BE(2)-C poly-A RNA. **(G)** Distribution of Ψ

sites in mRNA regions (5′UTR, CDS and 3'UTR), long non-coding RNA, and retained

intron. **(H)** MetaTranscript plot showing the proportion of Ψ along relative positions in

the 5'UTR, CDS, and 3'UTR of BE(2)-C mRNA. (**I**) MetaJunction plot showing Ψ sites

relative to their distances to the closest downstream (left) and upstream (right) exon-

exon junction.

Figure 3. Transcriptome-wide identification and quantification of PUS7-dependent

Ψ sites in BE(2)-C cells.

(A) Normalized counts of read lengths for direct BE(2)-C shGFP (orange) versus BE(2)-

C shPUS7-33 (blue) poly-A RNA. (**B**) Scatter plot showing that PUS7 knockdown

reduced the Ψ strength of Ψ-modified genes, with labeled representative genes (MANF,

NDUFS5, CBX3, RPS27A, and HMGB1). **(C)** K-mer frequency of the most represented

and frequently detected PUS7-dependent Ψ sites with high confidence. **(D)** Distribution

of PUS7-dependent Ψ sites in mRNA regions, long non-coding RNA, and retained

intron. **(E)** MetaTranscript plot showing the proportion of PUS7-dependent Ψ along

relative positions in the 5'UTR, CDS, and 3'UTR of mRNA. (**F**) MetaJunction plot

showing PUS7-dependent Ψ sites relative to their distances to the closest downstream

(left) and upstream (right) exon-exon junction. (**G**) GO analysis of the 1,246 PUS7 target

mRNAs. CC, cellular component; BP, biological process; MF, molecular function.

Figure 4. Comparative analysis of rRNA pseudouridylation in HeLa and BE(2)-C cells.

(A-C) Identification and quantification of Ψ sites in 5.8S, 18S and 28S rRNAs from BE(2)-C and HeLa cells. (**D**) Motif analysis of k-mers for Ψ sites shared between BE(2)-C and HeLa rRNAs. **(E)** Distribution of Ψ stoichiometry in HeLa and BE(2)-C rRNAs identified by nanopore direct RNA sequencing. **(F-G)** Scatter plot (**F**) and bar plot (**G**) showing the distribution of Ψ stoichiometry for the 103 Ψ sites common to both BE(2)-C and HeLa rRNAs. **(H)** Most represented Ψ k-mers identified in BE(2)-C and HeLa cells. (**I**) PUS7 dependent Ψ sites in BE(2)-C rRNAs.

Figure 5. PUS7-dependent pseudouridylation represses m6A modification in mRNA

(**A**) m^6A dot blot of 100 ng poly-A RNA from BE(2)-C shGFP and shPUS7 cells (n = 3

biological replicates). Methylene blue blot serves as loading control. **(B)** Bar graph

showing the numbers of Ψ sites across Ψ stoichiometry fractions following PUS7

knockdown. **(C-D)** Bar graphs showing the numbers of m^6A sites across various m^6A

stoichiometry fractions following PUS7 knockdown analyzed by CHEUI (**C**) or m6anet

(**D**) pipelines. **(E)** Violin plot showing m^6A stoichiometry fractions for various m^6A

DRACH k-mers following PUS7 knockdown identified by CHEUI. (**F-G**) Violin plots

showing the effects of PUS7 knockdown on Ψ (**F**) and m^6A (**G**) stoichiometry when they

co-occur on the same transcripts. *$p < 0.05$, ***$p < 0.001$.

Figure 6. PUS7-dependent pseudouridylation represses m^{5}C modification in mRNA

(A) m^{5}C dot blot of 100 ng poly-A RNA from BE(2)-C shGFP and shPUS7 cells (n = 3 biological replicates). Methylene blue blot serves as loading control. (**B**) Bar graph showing the numbers of m^{5}C sites across various m^{5}C stoichiometry fractions following PUS7 knockdown analyzed by CHEUI. (**C**) Violin plot showing PUS7 knockdown on m^{5}C stoichiometry for the most common m^{5}C k-mers analyzed by CHEUI. **(D-E)** Violin plots showing the effects of PUS7 knockdown on Ψ (**D**) and m^{5}C (**E**) stoichiometry when they co-occur on the same transcripts. ***$p < 0.001$.

739   **Supplemental Figure 1.** Immunoblotting of PUS7 in BE(2)-C cells expressing shGFP

740   or shPUS7-33.