High density linkage to physical mapping in a unique Tall x Dwarf

Coconut (*Cocos nucifera* L.) F$_2$ population uncovers a major QTL for

flowering time colocalized with the *FLOWERING LOCUS T (FT)*

Wellington Bruno dos Santos Alves[1,2], Cleso Antônio Patto Pacheco[3] and Dario

Grattapaglia[1,2]

[1]EMBRAPA Recursos Genéticos e Biotecnologia, Brasília, DF, Brazil

[2]Programa de Ciências Genômicas e Biotecnologia, Universidade Católica de Brasília,

Brasília, DF, Brazil

[3]EMBRAPA Tabuleiros Costeiros, Aracaju, SE

ORCID

Dario Grattapaglia: 0000-0002-0050-970X

*Corresponding author: dario.grattapaglia@embrapa.br

Keywords: Coconut, *FLOWERING LOCUS T*, hybrid breeding, SNPs, flowering

Running title: A flowering time QTL colocalized with FLOWERING LOCUS T in coconut

Abstract

The coconut tree crop (*Cocos nucifera* L.) provides vital resources for millions of people worldwide. Coconut germplasm is largely classified into 'Tall' (Typica) and 'Dwarf' (Nana) types. While Tall coconuts are outcrossing, more productive, stress tolerant and late flowering, Dwarf coconut are inbred, flower early with a high rate of bunch emission. Precocity determines earlier production of a plantation and facilitates management and harvest. We used a unique F_2 population generated by open pollination seeds collected in a commercial plantation of F_1 hybrids between Brazilian Green Dwarf from Jiqui (BGDJ) and West African Tall (WAT) to build a linkage map with ~3,000 SNPs, anchored to the existing chromosome-level genome assemblies. The map established the chromosomes numbering correspondence between the two reference genome versions and the relationship between recombination rate, physical distance and gene density. Leveraging the strong segregation for precocity inherited from the Dwarf cultivar in the F_2, a major effect QTL with incomplete dominance was mapped for flowering time. *FLOWERING LOCUS T* (*FT*) gene homologs of coconut previously described as putatively involved in flowering time by alternative splice variant analysis, were colocalized within a ~200kb window of the major effect QTL (LOD=11.86). Our work provides strong phenotype-based evidence for the role of the FT locus as the putative underlying functional variant for the flowering time difference between Dwarf and Tall coconuts. Major effect QTLs were also detected for developmental traits of the palm, plausibly suggesting pleiotropism of the *FT* locus for other precocity traits. Haplotypes of the two SNPs flanking the flowering time QTL inherited from the Dwarf parent BGDJ caused a reduction in the time to flower of around 400 days. These SNPs

46 could be used for high throughput marker assisted selection for early flowering and

47 higher productivity recombinant lines, providing novel genetic material to the coconut

48 industry.

Introduction

The coconut palm (*Cocos nucifera* L.) is a worldwide important tropical tree crop

providing vital resources for millions of farmers and industries with a diverse use of its

oil and fruit products. The only species belonging to the genus *Cocos*, the coconut palm

is a diploid (2n = 32) monoecious, perennial monocotyledon of the family *Arecaceae*

(Palmae), with a relatively large genome size (1C = 2.74 Gb). With its most likely center

of origin in Southeastern Asia, it is currently cultivated in more than 92 tropical countries

[1, 2] covering more than 12 million hectares worldwide [3].

Coconut germplasm is largely classified into 'Tall' (Typica) and 'Dwarf' (Nana)

types. Tall, a.k.a. Giant coconuts, are preferentially outcrossing, genetically diverse and

display high productivity, broad adaptability and tolerance to specific pests and

diseases. Dwarf coconuts are mostly self-pollinating with limited genetic diversity within

each variety but are chosen for their precocity and high rate of bunch emission. Early

flowering of the dwarf varieties is a feature of great agronomic relevance that results

not only in precocity of nut harvest, but also increased production, while facilitating

coconut tree management and harvest operations. While the tall (late) variety takes an

average of 8 years to flower, the early flowering dwarfs typically flower on average 2 to

4 years after planting [4]. The origin of dwarf coconuts has been a matter of debate,

although current consensus based on molecular studies points toward a process of

domestication from the ancestral tall coconut possibly following the appearance of

mutations in flowering time genes resulting in autogamy [4], as well gibberellin

biosynthesis affecting height growth [5, 6].

Despite its global socioeconomic importance, advances in coconut tree

improvement have been modest, hindered by its perennialism, long generational time,

late flowering and large tree size, which demand extensive experimental areas and

continuous long-term investments. Genetic improvement programs have adopted two

main breeding strategies: (1) intravarietal breeding coupled to mass selection and (2)

intervarietal hybridization, the most widely used strategy, which consolidates, into an

intermediate F_1 hybrid, the key features of each variety, such as precocity and short

stature from the dwarfs and the rusticity and production of larger fruits from the tall

varieties [2, 4]. Rare have been studies in F_2 populations derived either from selfing or

cross pollination among F_1 Tall x Dwarf hybrids. The wide segregation for production

traits reported in F_2 generations has discouraged such endeavors. Time to flowering,

pollination behavior, height and the presence of bole - an expansion at the base of the

stem - have been considered to segregate independently. While the former two traits

showed the involvement of several genes, height and bole character segregation

suggested the presence of at least one major gene each, with incomplete dominance [4,

7, 8], although no genetic follow-up testing of such hypothesis was ever reported.

Despite the socioeconomic importance of the coconut tree, advances in mapping

quantitative trait loci (QTLs) of economically important traits have been limited,

restricted to a few studies using Tall x Dwarf F_1 hybrid populations that were typically

readily available [9-13]. In such F_1 experimental populations, segregating variation can

only be captured from the outbred tall coconut, because the dwarf parent, being

essentially an inbred line, will not contribute allelic segregation. Flowering time as well

as other traits related to the dwarf behavior typically display an intermediate phenotype

in the F_1 progeny with no segregation that would allow genetic mapping of major effects

contributed by the dwarf genotype. Studies of F_2 or backcross populations are needed

to advance the molecular understating of key traits in coconut. Furthermore, genetic

mapping in the cited studies has been carried out either with low portability AFLP

markers or a limited set of microsatellites, which do not provide adequate sequence-

level information for deeper investigation of putative genes underlying mapped QTLs.

Only recently has this scenario begun to change with studies employing single

nucleotide polymorphism (SNP) markers for diversity studies [14], linkage mapping to

support genome assembly [15], and whole genome sequencing for genome-wide

studies (GWAS) [6]. Understanding the genetic basis of morphological traits that

determine precocity with a particular interest in early flowering, would represent an

important step towards breeding advancement of coconut.

In the present study, we used a unique segregating F_2 population generated by

open pollination of F_1 hybrids between Brazilian Green Dwarf from Jiqui (BGDJ) and

West African Tall (WAT) to dissect the quantitative variation underlying the genetic

divergence between tall and dwarf coconut types. We built a high-density linkage map

using SNPs derived from genotyping by sequencing with DArTseq, aligned the map to

the current genome assemblies and located QTLs for traits related to precocity, including

a major effect locus for flowering time colocalized with the *FLOWERING LOCUS T* (*FT*).

Materials and methods

Plant material and DNA extraction

The study was carried out with 182 F_2 plants, generated by open pollination among PB-141 F_1 coconut hybrid plants in a ten-year-old commercial plantation managed by Embrapa Tabuleiros Costeiros, Aracaju – SE (10.9513° South, 37.0528° West). PB-141 is an F_1 hybrid derived from the cross between the Brazilian Green Dwarf from Jequi (BGDJ) and West African Tall (WAT). Coconuts were collected from F_1 palms in the central area of the commercial plantation in order to maximize the probability of being effectively derived from intercrossing of F_1 plants. F_2 plants were therefore derived either from selfing or cross pollination between different F_1 plants. Samples of young leaves were collected and used to extract total genomic DNA using an optimized Sorbitol-CTAB-based protocol [16]. DNA concentrations and purity were estimated using a Nanodrop 2000 spectrophotometer (Thermo Scientific).

SNP genotyping

Genomic DNA samples were shipped to the Service of Genetic Analysis for Agriculture (SAGA) laboratory in Mexico for high-throughput genotyping using the DArTSeq method developed by Diversity Arrays Technology Pty Ltd. [17]. DNA complexity reduction was performed using the SbfI/MseI enzyme combination. Samples were processed in digestion/ligation reactions and the scrambled fragments were amplified using two primers with sequence complementary to the ligated adapters and oligonucleotides from the sequencing platform. Subsequently, the clusters were sequenced on an Illumina HiSeq2500 sequencing platform. SNP markers were detected following mapping of sequence reads sampled at adequate depth on the Coconut

136   reference genome [15] according to standard parameters defined in DArTsoft14, an

137   automated genomics data analysis program and DArTdb, a laboratory management

138   system developed and patented by DArT for generating SNP marker data. Codominant

139   SNP markers were scored as "0" for homozygous reference allele, "1" for homozygous

140   alternative allele, and "2" for presence of both the reference and the alternative allele.

Morphological and flowering trait evaluations

Due to restrictions in field availability for planting, only 125 of the 182 F_2 plants

were ultimately planted in a field trial in a site known for its soil homogeneity at

Embrapa Tabuleiros Costeiros research center in June, 2018. Since to replicate plants

were possible for the 125 F_2 plants (n =125 entries), these were deployed in an

augmented randomized complete block design (ARCBD) [18] with c = 3 checks (random

plants of the parents BGDJ, WAT and the F_1 hybrid BGDJxWAT), r= 10 blocks extending

in rows. Because the number of entries was not a multiple of the number of blocks, nine

blocks had 13 entries and the 10th block had 8 entries. Nine traits were ultimately

evaluated in 121 live coconut trees: Number of Leaves (NL), Height of Second Leaf Hem

(cm) (HSLH), Petiole Length (cm) (PL), Rachis Length (cm) (RL), Petiole Width (cm) (PW),

Petiole Thickness (cm) (PT), emergence of the base of the stipe (BOLE), Number of

Bunches with fruits (NBF) and Days to Launch of the First Spathe (DLFS). BOLE was

recorded in a binary fashion with presence of bole when it measured at least 70 cm from

the ground, and absence when at a lower that 70 cm height from the ground. DLFS was

used as a direct quantitative measure of time to flowering. These traits, but BOLE, are

typically used to differentiate dwarf from tall plants in the early stages of life.

Measurements were taken according to the standardized methods described by Santos,

Batugal (19). Trait evaluations were carried out at one and two years after planting, on

June 30, 2019 and July 30, 2020 with evaluation of morphological leaf traits. Flowering

initiation was measured on each individual plant according to the emergence of the first

spathe starting on October 11, 2020 and then by a monthly monitoring to precisely catch

the launch of the first spathe. The number of days for the emergence and opening of

the subsequent spathes until the tenth spathe was recorded until the least evaluation

on August 11, 2022. The final evaluations of the morphological traits were carried out

on May 10, 2022, with the exception of NBF, which was evaluated on August 11, 2022.

Pearson correlations were estimated to assess the correlation between traits using the

Corrplot package available in R [20]. Phenotypic data for all traits but BOLE was analyzed

with the following mixed model: $Y_{ij} = \mu + B_i + E_j + C_{(i)j} + e_{ij}$ where Y is the observed

phenotypic value of the genotype entry (j) in block (i), μ is the overall mean, B is the

fixed block effect, E is the random genotype entry effect, C is the random effect of the

common checks in blocks and e is the random residual effect. Best linear unbiased

predictors (BLUPs) were estimated using the software SELEGEN [21] under model 76.

174   Linkage map construction

The raw SNP calls were filtered for quality for the linkage mapping analyses. The

first filtering step was based on a call rate >0.95 followed by selection of SNP markers

that fit the expected segregation proportions in an F_2, by a Chi-square test under the

null hypothesis of a 1:2:1 segregation ratio with a more relaxed threshold (α>0.001) to

allow including SNPs even with some level of segregation distortion given the

particularity of the mapping population. Linkage analysis and construction of the genetic

map were performed according to the standard F_2 model using JoinMap v3.0 [22]. The

genotypes of the F_2 population were coded, according to the format <hkxhk> for an

unknown linkage phase configuration and three possible genotypic classes. Markers

were assigned to linkage groups (LG) via the clustering module with log-of-probability

(LOD) scores greater than 15, maximum recombination fraction of 0.4, under a Kosambi

mapping function. Linkage maps were drawn and aligned to the assembled genomes

using MAPCHART [23]. For the alignments, recombination distances were expressed in

centiMorgans and physical distances in Mbp.

189 Marey maps and recombination rates in the genome of *Cocos*

190 *nucifera*

To estimate local recombination rates across the 16 coconut chromosomes, the

targeted SNP marker sequences were aligned to the two existing chromosome-level

coconut genome assemblies of Hainan Tall coconut: (1) a draft assembly built using

exclusively short Illumina paired-end sequencing [15] hereafter called the YANG genome

and (2) a reference-grade assembled using long Nanopore single-molecule sequencing

and Hi-C technology [6] hereafter called the WANG genome. The FASTA sequences of

the chromosome assemblies were downloaded from their respective repositories. The

YANG genome from GenBank BIOPROJECT PRJNA374600, and the WANG genome from

the Genome Warehouse in the National Genomics Data Center (Beijing Institute of

Genomics) accession number GWHBEBT00000000, BIOPROJECT PRJCA005463. The

DArTseq sequences with 69 bases were mapped to the sequences of the 16 coconut

pseudo-chromosomes assembled using BLASTn for highly similar sequences (megablast)

with a cutoff E-value <1E-05. DArTSeq sequences that mapped to a unique location in

both genome versions were used in further analyses. The relationships between the

genetic and physical positions of the mapped SNPs were represented by the local slope

of the curve adjusted to the datapoints in MAREY maps [24]. Curves were fitted to the

data using locally weighted scatterplot smoothing (LOESS) polynomial regression with a

smoothing parameter of 0.3 using MareyMap Online [25]. Sporadic abnormal points

from the map that did not fit the expected monotonously increasing relationship

between physical and genetic positions were removed using the available functionality

in the software. The physically mapped markers were used to obtain an estimate of the

effective physical coverage achieved by the genetic map and the recombination rates in

cM/Mb along the chromosomes. Pearson's correlations along windows of 1 Mb were

calculated between the average recombination rate in the 1 Mb interval and the number

of genes as annotated in the coconut chromosomes in the YANG genome [15].

QTL mapping

QTLs detection was performed using the BLUPs for eight traits and the binary 0/1

value for BOLE with R/qtl version 3.5.3, using different methods: interval mapping (IM),

Haley Knott regression (HK) and non-parametric interval mapping for those traits that

violated the assumption of normality [26]. The probability of genotype error (error.prob

= 0.01) was calculated and the threshold logarithm of odds (LOD) score for QTL detection

was determined by genome-wide LOD significance thresholds, setting LOD thresholds at

different alpha levels (α = 0.01; 0.05 and 0.1) using 1900 permutations to allow for the

declaration of QTLs with variable stringency levels in order to control also for Type II

errors.

Co-localization of the FLOWERING LOCUS T (FT) gene with the flowering time QTL

A set of 198 expressed gene sequences in coconut were recently identified as homologues of flowering-time genes in *Arabidopsis*. Five of them, annotated as FT homologs to the *Arabidopsis thaliana* FT gene (AT1G65480) were found to be differentially expressed between seedling and reproductive leaf samples in both tall and dwarf coconut. Alternative splicing analysis showed that the FT gene produces different transcripts in tall compared to dwarf coconut [27]. We took all the 87 *Arabidopsis thaliana* genes present in the annotation of the 198 coconut expressed genes and aligned them to the QTL region on the corresponding chromosomes assembled in the two genome assemblies using BLASTn specifically in the genome stretch from 65 Mbp to 75 Mbp where the major effect QTL was mapped (see below).

Results

SNP markers and linkage mapping

Genotyping by sequencing using DArTseq generated 86,719 raw SNP calls based on the standard parameters used in DArTsoft. After applying a call rate threshold of 95% and a test for adherence to a null hypothesis of 1:2:1 segregation, 3,714 SNPs (4.28% of the raw SNP calls) were retained for further linkage analyses (S1 File). Dominant presence/absence in silico DArTseq variants were not used due to their low information content and considering that an adequate number of co-dominant informative SNPs was already available for linkage mapping purposes. A linkage map was ultimately built with

182 F_2 individuals that correspond to 364 meiotic events sampled in the gametes from

the recombined F_1 parents. All 3,714 SNPs were mapped with LOD ≥15 and maximum

recombination fraction of 0.4, in 16 linkage groups matching the expected 16 coconut

chromosomes. However, for the subsequent assessment of the recombination rates and

QTL mapping, only the SNPs that were physically positioned on the genome assemblies

(see below) were ultimately considered on the linkage map (S2 File). Linkage group

numbering used for our map was based on the physical positioning of SNP markers on

the YANG genome chromosomes which was published first in the literature. The

chromosome numbering used in the WANG genome was different but the

correspondence was established and reported throughout (Table 1). An average of

184.5 SNPs were mapped to each chromosome, varying from a maximum of 329 SNPs

on chromosome 1 to a minimum of 101 SNPs on chromosome 15. The genetic map

spanned a total of 2,124.3 cM in recombination size, with an average linkage map length

of 132.8 cM, and an average intermarker recombination distance between adjacent

marker pairs of 0.742 cM. Chromosome 15 displayed a large recombination gap of 31.2

cM between two denser linkage clusters although still linked by the minimum LOD 15

score threshold adopted (Fig 1). Except for this large gap observed on chromosome 15,

maximum intermarker distance was 6.7 cM on chromosome 7.

Fig 1. Alignment of the linkage map (green) to the two chromosome-level genome

assemblies, the YANG genome assembly (red) [15] and the WANG genome assembly

(blue) [6].

Alignment of the linkage map to the genome assembly

269  The 3,714 DArTseq sequences containing the linkage mapped SNPs were

270 submitted to BLASTn against the chromosome assemblies. On the YANG genome out of

271 the 3,714 linkage mapped DArTSeq SNPs, 2,952 DArTSeq sequences were mapped to

272 unique locations with E-values <7.3e-05. The sequences for the remaining SNPs either

273 aligned to multiple locations or did not align to the assembled coconut chromosomes

274 and were removed from further analyses. On the WANG genome 3,027 DArTSeq

275 sequences were mapped to unique locations with E-values <3.6e-11 (S3 File). On the

276 YANG genome, the linkage map covered 1,035 Mb corresponding to a coverage of

277 approximately 43.14% of the 2,400 Mb *Cocos nucifera* Tall cultivar genome [6]. On the

278 WANG genome, the linkage mapped SNPs covered a total of 2,382 Mb, a 99.27%

279 coverage, more than twice the coverage obtained on the YANG genome with a

280 corresponding lower density of mapped SNPs and long stretches with no segregating

281 SNPs mapped to them. The average estimated physical inter SNP distance were 349kb

282 and 779kb, SNP density of 2.85 and 1.27 SNPs/Mb and maximum physical distances of

283 9.7 and 63.4 Mb respectively on the YANG and WANG genomes (Table 1).

284 **Table 1.** Statistics of the linkage map and its alignment to the two available chromosome-level genome assemblies of coconut (*Cocos nucifera* L.).

	LINKAGE MAP			YANG GENOME						WANG GENOME			
LG #	# of link. map SNPs	Total Size (cM)	Mean interSNP dist (cM)	Chr. #	Phys. dist. covered (Mb	Mean interSNP phys. dist. (Mb)	Number of annotated genes[#]	Average recomb. rate (cM/Mbp)	r^2 (Number of genes and recomb. rate)	Chr. #	Phys. dist. covered (Mb)	Mean interSNP phys. dist. (Mb)	Average recomb. rate (cM/Mbp)
1	329	196.6	0.599 ± 0.470	1	119.197	0.362 ± 0.652	2277	2.27 ± 1.17	0.68	5	177.52	0.570 ± 1.533	2.12 ± 0.94
2	311	199.1	0.642 ± 0.492	2	85.353	0.275 ± 0.411	1997	2.44 ± 0.66	0.72	1	213.40	0.656 ± 2.751	1.97 ± 0.649
3	209	149.2	0.717 ± 0.625	3	82.325	0.396 ± 0.576	1705	1.98 ± 0.65	0.73	4	179.12	0.835 ± 2.527	1.67 ± 1.26
4	222	161.2	0.729 ± 0.536	4	66.748	0.302 ± 0.454	1610	3.05 ± 1.52	0.87	6	170.88	0.762 ± 2.005	2.56 ± 1.46
5	150	147.5	0.989 ± 0.654	5	57.717	0.385 ± 0.624	1104	3.77 ± 2.06	0.89	9	156.70	1.073 ± 5.397	2.98 ± 1.87
6	262	193.4	0.729 ± 0.764	6	81.468	0.311 ± 0.463	2095	2.42 ± 0.74	0.46	3	187.69	0.693 ± 2.894	1.84 ± 0.50
7	215	148.8	0.695 ± 0.686	7	77.453	0.361 ± 0.758	1583	2.13 ± 0.60	0.21	2	188.13	0.776 ± 2.790	1.60 ± 0.76
8	176	95.7	0.547 ± 0.505	8	58.631	0.326 ± 0.668	1214	2.09 ± 1.16	0.77	13	95.96	0.508 ± 1.273	1.56 ± 0.75
9	106	89.9	0.856 ± 0.725	9	59.333	0.554 ± 0.792	1026	2.37 ± 1.33	0.89	11	137.92	1.378 ± 3.871	2.12 ± 1.38
10	144	113.0	0.790 ± 0.493	10	57.105	0.399 ± 0.663	1066	2.90 ± 1.45	0.87	10	144.69	0.969 ± 4.013	2.39 ± 1.53
11	165	72.4	0.785 ± 0.750	11	61.809	0.375 ± 0.617	1379	3.01 ± 1.93	0.61	7	173.86	1.073 ± 3.547	2.64 ± 1.77
12	133	94.9	0.728 ± 0.489	12	42.367	0.302 ± 0.425	868	3.21 ± 1.86	0.86	14	89.87	0.681 ± 2.625	2.88 ± 1.76
13	127	102.6	0.814 ± 0.617	13	60.488	0.479 ± 1.032	847	3.77 ± 2.55	0.87	15	82.46	0.641 ± 1.677	3.63 ± 2.05
14	189	132.8	0.706 ± 0.544	14	56.198	0.296 ± 0.351	1309	3.15 ± 1.35	0.73	8	155.26	0.754 ± 3.799	2.66 ± 1.30
15	101	135.7	1.357 ± 3.139	15	31.114	0.311 ± 0.800	646	5.18 ± 3.33	0.04	12	145.36	1.075 ± 3.394	3.04 ± 1.43
16	113	91.5	0.819 ± 0.664	16	38.139	0.313 ± 0.623	867	3.97 ± 1.48	0.91	16	83.33	0.686 ± 2.239	3.34 ± 1.58
Total	2,952	2,124.3	-		1035.445	-	21,593	2.79 ± 1.65			2,382.141	-	2.32 ± 1.40
Mean ± s.d.	184.5 ± 69.9	132.8 ± 40.8	0.742 ± 0.471		43.14%	0.349 ± 0.652					99.27%	0.779 ± 2.947	

285 # Number of annotated genes from (Yang et al. 2021; BioProject: PRJNA374600)

Marey maps and recombination rates

The linkage maps alignment to the 16 pseudochromosomes was largely colinear with the physical assembly, with only sporadic inconsistencies of the map versus genome positions, 2 in the YANG genome and 20 in the WANG genome (Fig 1; S4 File). As expected, on all linkage groups the alignment to the physically larger WANG genome showed long sections of genome sequence with few or no SNP markers mapped. The Marey maps showed plateaus of restricted recombination, where the recombination distance does not follow the increase in physical distance along the chromosome (Fig 2). These plateaus spanned considerably longer stretches for the Marey maps on the WANG genome. From the Marey maps data, estimates of local recombination rate were obtained for both genomes using LOESS (Locally Weighted Scatterplot Smoothing) (S4 File) and plotted for the YANG genome (Fig 2). Following LOESS treatment of the data, the average local recombination rates were similar: 2.78 cM/Mb for the YANG genome and 2.32 cM/Mb for the WANG genome. The smoother recombination rate curves closely followed expectations. Chromosomal segments of restricted recombination in the Marey maps coincided with local valleys of low recombination rate notably spotted on YANG chromosomes 1, 4, 5, 11 and 14, for example and more accentuated on the corresponding WANG chromosomes.

Fig 2. Marey maps showing the correspondence between the physical position (x-axis, Mb) and the recombination-based position (y-axis, cM) on the 16 Coconut chromosomes of the YANG genome assembly [15] and the WANG genome assembly [6]. Also shown in

the intermediate panels the combined pattern of gene density distribution and the

LOESS curve of recombination rate for the YANG genome.

Average recombination rates per chromosomes varied across chromosomes up

to more than twofold (e.g., 1.98 cM/Mb on chromosome 3 and 5.18 cm/Mb on

chromosome 15 on the YANG genome). However, the correlation between the rates

estimated on the two genome versions was high and significant (r = 0.85; p-value = 3E-

5). The highest rate on chromosome 15 is likely the result of the large linkage gap

observed on this chromosome. The average recombination rate across the entire

genome were similar, at 2.79 and 2.32 cM/Mb for the two genomes (Table 1).

Furthermore, taking advantage of the gene annotation provided for the YANG genome

sequence, we underlaid the frequency distribution of the number of genes in 1 Mb

intervals, equivalent to gene density, to the smoothed recombination rate curve in the

same 1 Mb segment (Fig 2). Visually an almost perfect direct relationship was seen

between gene density and recombination rate. This visual relationship was further

validated by observing high and significant Pearson correlation between average

recombination rate and number of genes in the 1 Mb intervals estimated for the YANG

genome. For all chromosomes but chromosomes 7 and 15 these correlations were high

and significant (p-value <0.001) (Table 1) suggesting, once again, some different

behavior as far as recombination for these two chromosomes.

Trait variation and correlations

The F_2 individuals displayed extensive segregation for all traits measured, as

expected in a recombinant F_2 population of divergent cultivars. Raw phenotypic values

and BLUPs are provided in S5 File. The continuous traits showed an approximately

normal distribution, although with some skewness, most notably for NBF and DLFS (Fig

3). BOLE expression fit a 3:1 segregation ratio with 98 plants showing no bole and 23

with bole (Chi-square 2.31; p-value = 0.128). Days to Launch of the First Spathe (DLFS)

varied from a minimum of 812 days for the precocious flowering plants to a maximum

of 1608 days for the late blooming ones with an average of 1287 ± 214.5 days. One plant

therefore displayed 2.2 standard deviations below the mean and 13 flowered with less

than 1000 days, corresponding to more than one standard deviation below the mean.

Eight of these very early flowering plants displayed the highest numbers (≥13) fruit per

bunch (NBF). Significant negative correlations were seen between DLFS and NL, HSLH,

BOLE and NBF, and a significant positive correlation with PL (Fig 4). Positive significant

correlations were seen between five of the six leaf morphology traits (NL, HSHL, RL, PW

and PT), except PL that showed no correlation with all other five traits. Positive and

significant correlations were also observed between NBF and NL, HSLH, RL and BOLE.

Coconut trees with a recorded bole flowered on average after 1098 days and produced

an average of 9.5 fruit per bunch while plants without bole flowered on average after

1347 days, and produced 5.96 fruit per bunch.

Fig 3. Histogram of the coconut phenotypic trait distributions in the F_2 mapping

population.

Fig 4. Matrix of genotypic correlations among the measured traits in the F_2 mapping

population

Table 2. Summary of QTL mapping results in the coconut F2 population.

	Days to Launch of First Spathe (DLFS)	Bole emergence (BOLE)	Number of bunches with fruit (NBF)	Petiole Length (PL)
Chromosome	3	3	1	3
LOD score threshold (percentile)	5.14 (99%)	4.47 (95%)	5.14 (99%)	5.14 (99%)
# SNPs in QTL region	52	1	1	22
QTL region position (cM)	118.928 - 149.166	132.62	57.16	123.503 – 136.125
Right SNP	100287206_T>A-55	-	-	100287206_T>A-55
Left SNP	100273632_T>G-15	-	-	100273632_T>G-15
Peak LOD	11.86	4.64	5.68	7.79
Peak LOD SNP	100226499_C>T-34	100228320_G>A-40	100237259_A>G-48	100226499_C>T-34
Target SNP allele (Origin; effect)	T (Dwarf; reduction)	A (Dwarf; increase)	A (Dwarf; increase)	T (Dwarf; reduction)
Genome position (bp)	73,147,443	75,613,473	19,206,916	73,147,443
Variance explained (%)	64.0	3.12	0.43	22.3

Mapping and haplotype analysis of a major flowering time QTL

QTL detection was carried out with three methods, IM, CIM and HK to assess

consistency of results and minimize type I errors. Significant QTL peaks were declared

based on the LOD limit analysis for the different alpha levels to allow the detection of

suggestive QTLs in order to minimize Type II errors. The significance thresholds indicated

LOD limits of 4.15, 4.49 and 5.22 respectively for the different alpha levels ($\alpha = 0.01$;

0.05 and 0.1) (Table 2). Results with CIM showed a considerably larger number of 16

putatively significant QTLs in 11 chromosomes when compared to the results with IM

and HK. These CIM results are not discussed further but are provided (S6 File). Genome-

wide QTL profile results across the 16 chromosomes obtained with the three QTL

detection methods, for the four traits for which significant QTLs were detected are

provided for a full visualization of results. The LOD limit thresholds are specified by the

differentially dotted lines (Fig 5).

Fig 5. Genome-wide QTL mapping profiles for the mapped traits (Y axis) obtained with

the three different QTL mapping methods implemented (X axis)

Overall, consistent QTLs across the methods were detected on chromosome 3

for DFLS, BOLE and PL and on chromosome 1 for NBF. A genomic segment at the tip of

chromosome 3 spanning 52 significantly linked SNPs to DFLS stands out as a major effect

QTL for flowering time, explaining an estimated 64% of the variation in DLFS. The

alignment of the linkage map to the chromosome 3 of the YANG genome and

chromosome 4 of the WANG genome allowed a detailed haplotype analysis of this

segment. In the YANG genome starts at SNP 100285601_C>G-14 with LOD 4.99 located

at 117.99 cM corresponding to 68.84 Mbp, and ends at SNP 100270789_A>G-59 with

LOD 6.53 located at 148.17 cM corresponding to position 82.32 Mbp (S3 File). In the

WANG genome it spans 55 SNPs including the 52 mapped to the YANG genome form

position 162.7 Mb to the end of the chromosome at 178.5 Mb. Within this segment, the

highest QTL peak (LOD 11.86) is located at SNP 100226499_C>T-34, positioned at 129.11

cM and 73.15 Mbp, flanked by SNPs 100287206_T>A-55 and 100273632_T>G-15. At SNP

100226499_C>T the T allele inherited from the Dwarf parent AVeBrJ causes a major

reduction in the time to flower, suggesting incomplete dominance of the locus.

Heterozygous genotype G/T confers a reduction from 1460 to 1390 days to the launch

of the first spathe, while the homozygous T/T genotype confers a further reduction of

almost 300 days, as the coconut trees bearing this genotype flowered on average after

1110 days (Fig 6). In addition to acting on early flowering, the same genomic segment is

involved in the control of BOLE and PL. The LOD peak for both QTLs for BOLE and PL was

located at the same SNP 100228320_G>A-40, at 132.62 cM and 75.61 Mbp. The co-

localization of QTLs for three traits suggests a pleiotropic action of this major effect QTL

segment on chromosome 3. Some additional suggestive QTLs were detected for PL on

chromosomes 1, 2, 4, 6 that were significant at $\alpha = 0.1$ based on the IM analysis but that

did not reach significance with the two other QTL detection methods. Finally, a single

QTL was also found for NBF on chromosome 1 whose Dwarf allele confers an increase

in the number of fruits per bunch.

Fig 6. Detailed view of the major effect QTL region for flowering time, petiole length and

bole presence in the stem on chromosome 3 of the YANG genome. The *A. thaliana*

FLOWERING LOCUS T was colocalized with the leading SNPs for the QTL.

Co-localization of the flowering time QTL with the FLOWERING

LOCUS T (FT)

Out of the 87 *Arabidopsis thaliana* genes involved in flowering-time pathways

that were found to be homologous to 198 coconut genes [27], only the complete coding

sequence of the *Arabidopsis thaliana* putative flowering signals mediating protein FT

(At1g65480) produced strongly significant alignment to the flowering time QTL mapped

on linkage group 3. The FT gene (At1g65480) complete cds mRNA (accession

AY065378.1) with 840 bp produced two hits in the YANG genome: (i) at 70,554,038 to

70,554,249 bp with bit score 108, 71% identity, e-value = 3e-23 and 0/212 gaps (0%) and

(ii) at 73,265,231 to 73,265,450 bp with bit score 88.7, 70% identity, e-value = 3e-17 and

3/222 (1%) gaps (Fig 6; S7 File). These hits are located respectively at 137 kb from SNP

100241292_A>G-33 LOD 9.55 and 117 kb from SNP 100226499_C>T-34 this last one had

the highest LOD score (LOD 11.86) in the QTL region. When the FT gene (At1g65480)

was blasted against the corresponding chromosome 4 in the WANG genome it also

produced two significant hits at 164.688 Mb (e-value = 7E-21) and 168.065 Mb (e-value

= 7E-15) that correspond to the genomic positions at the tip of the chromosome (Fig 1;

S8 File), further corroborating the finding. This co-localization result between the

flowering time QTL and the FLOWERING LOCUS T (FT) gene when taken together with

the splice variant analysis reported earlier [27], strongly suggests that the underlying

candidate gene to the flowering time QTL mapped in this study could be an FT homolog

in *Cocos nucifera.*

425 Discussion

We have generated a dense linkage map for *Cocos nucifera* based on a novel

approach for the crop, by using an F_2 population derived from open pollination

intercrossing of F_1 hybrids between Brazilian Green Dwarf from Jequi (BGDJ) and West

African Tall (WAT). Almost 3,000 genotyping by sequencing DArTSeq sequence derived

SNPs were positioned on the existing chromosome-level coconut genome assemblies.

By aligning the linkage map to the genome sequences, we provided an independent

assessment of the contiguity of the current coconut reference genomes, and a detailed

analysis of the relationship between recombination rate and physical distance for the

corresponding coconut chromosomes. Furthermore, based on the segregation observed

for traits related to precocious behavior inherited from the Dwarf cultivar, we mapped

a major effect QTL for flowering time on chromosome 3. Flowering Locus T (FT) gene

homologs of coconut, previously described as involved in flowering time by alternative

splice variant analysis, were co-localized with the molecular phenotype mapped QTL,

providing evidence for its role as the putative underlying gene.

440 Exploiting the unique genetics of the Tall x Dwarf coconut hybrid

441 for linkage mapping in an outbred F_2.

The linkage map reported in our study is the first one in coconut built from an F_2

segregating population from a Tall x Dwarf hybrid. Previous maps either used hybrid F_1

or backcross (BC1) populations [9, 10, 13, 15] or a cross between Tall cultivars[11].

Differently from a conventional F_2 populations generated by selfing an individual F_1

plant, the particular genetics of coconut allowed using an open pollinated progeny of an

F_1 hybrid field. The parental F_1 plants composed a commercial production plantation

with seeds resulting from controlled pollination between a pollen mix of several Tall

coconut parents on female flowers of several Dwarf plants. In the particular case of

coconut, we took advantage of the fact that the Dwarf variety is essentially a fixed inbred

line with very little if any residual heterozygosity, while the Tall parents are allogamous,

heterozygous and genetically diverse [6, 14, 28]. The F_1 plants SNP genotypes would

potentially differ only in what was inherited from the Tall parents, but SNP markers fixed

in the Tall variety and polymorphic in relation to the Dwarf would segregate in a 1:2:1

ratio in the F_2, irrespective of what Tall parent generated each different F_1 plant

intercrossed to produce the F_2 population. This premise proved valid, and 3,714 SNPs

called with a call rate >95%, fit the expected segregation and were linkage mapped.

Approximately 20% of the sequences of linkage mapped DArTSeq mapped to multiple

locations in the two genomes and were left out of subsequent analysis, as no objective

criteria allowed choosing one location over another. Therefore, for QTL mapping

analyses we used only around 3,000 linkage mapped SNPs that could be anchored to

unique positions on the 16 assembled chromosomes of the two genome versions.

The linkage map assembled in the 16 expected linkage groups spanned a total of

2,124.3 cM, slightly smaller than the linkage map reported earlier by Yang et al. [15] with

2,365 cM built from a backcross (BC_1) population with 216 individuals and 8,402 SNPs

retained with >80% call rate. Linkage group recombination sizes of our map were also

similar to those reported by Yang et al. in the BC_1 map (Spearman rank correlation 0.72),

including the only large recombination gap in chromosome 15, starting at around the 40

cM position (Fig 1). This recombination gap was hypothesized to be due to reduced

heterozygosity resulting from a selection sweep in favor of the allele transmitted from

the Dwarf parent observed in the YANG genome assembly [15].

Although our map was built from a smaller number of individuals than the BC_1

map used to anchor the YANG genome [15], those were derived from two recombined

gametes from the F_1 parents, therefore sampling twice as many meiotic events as in a

BC1 population. The number of recombination events per meiosis captured in the

mapping population is one of the main drivers of mapping accuracy, with an F_2 being the

most efficient configuration [29, 30]. Additionally, for improved linkage map quality, we

decided to use only SNPs with call rate >95% aware of the potential reproducibility

challenges of marker data derived from genotyping by sequencing methods in

heterozygous genomes [31]. From the raw data, 10,576 DArTSeq markers were retained

at >95% call rate, and the main detractor of the final SNPs number was the test for the

1:2:1 segregation. This substantiates the variability in the F_1 parents contributed by the

Tall grandparental pollen pool. This observation indicates that a truly self-pollinated

progeny from a single F_1 plant would possibly allow mapping a considerably larger

number of DArTSeq SNP markers.

486   The F_2 linkage map provides an independent assessment of the

487   properties of the coconut genome assemblies

The DArTseq method using the rare cutter SbfI combined to the frequent MseI

enzyme proved effective for complexity reduction and enrichment of the sequenced

pool for low copy genomic regions across most of the large coconut genome, generating

a large number of segregating markers [17]. Our mapping and data quality control

approach proved effective to order SNPs markers along the linkage groups into robust

maps, highly colinear with the genome sequences (Table 1). The physical distance

covered by our F_2 linkage map was 1035.4 Mb on the YANG genome, slightly higher than

the 1,020 Mb of the Yang et al. BC_1 map, while on the WANG genome, the mapped SNPs

covered 2,382 Mb, reaching a 99.27% coverage. As expected, very few segregating

DArTSeq sequences and the corresponding SNPs were mapped to long physical

stretches covering several hundred Mb of DNA of the reference-grade WANG genome

assembly produced by nanopore long read sequencing (Fig 1). These regions were not

sampled in the YANG genome, assembled exclusively with short read sequencing. The

extensive genome gaps with no mapped SNPs and low gene density (Fig 2), likely

correspond to heterochromatin, typically found at centromeres and telomeres, known

to be relatively gene poor, mostly consisting of repetitive DNA sequences [32], where

the DArTSeq method by principle tends not to sample restriction fragments. Performing

linkage mapping in those heterochromatic regions will prove challenging, requiring

approaches that can target the occasional low-copy regions scattered in the repetitive

DNA. Using a selfed F_1 progeny might prove valuable in this respect to sample much

larger number of segregating DArTSeq SNPs, possibly targeting low-copy interspersed

segments in these heterochromatic regions of the coconut genome.

The alignment of the linkage maps to the genome assembly indicated consistent

ordering between the Joinmap estimated positions of the SNPs, and their relative

physical order along the assembled chromosomes, with only sporadic inconsistencies

seen most notably at the tip of some chromosomes (Fig 1). These localized

inconsistencies could be due either to erroneous linkage map ordering or misassembled

genome scaffolds. Historically, the majority of the inconsistencies between the physical

and genetic map order pointed to errors in the physical map order [33], making linkage

maps effective tools to correct genome assemblies [34]. However, sequencing

technologies and assembly algorithms have improved tremendously in recent years. The

alignment of the linkage map to two independent genome assemblies, one of them built

with long reads, indicates that those few SNPs that show inconsistent positions in

relation to both genome assemblies, for example at the tips of chromosomes 8, 9 and

12, most likely were inaccurately mapped.

The largely consistent genetic to physical position of the mapped SNPs led us to

carry out further analysis of this relationship for each coconut chromosome by

estimating local recombination rates in cM/Mb, given by the slope of the curve built

with graphical Marey maps. Additionally, the gene density histogram distribution

annotated in the YANG genome was merged to the Marey maps for a combined view

(Fig 2). In all chromosomes, the Marey maps showed genomic stretches of constrained

recombination, where the recombination distance does not follow the linear increase in

physical distance and gene density drops. This observation was evidently more distinct

in the WANG genome Marey maps that included the heterochromatic regions of the

genome. These regions are typically considered to be the putative location of

centromeres. Considering the WANG chromosome numbering for now, these were

located more toward the extremes of the chromosome (chromosomes 3, 7, 8, 9, 10, 12,

13 and 16) that would correspond to acrocentric or even telocentric chromosomes. In

the other chromosomes these regions are found more toward the middle (Fig 2).

Centromeres are essential for faithful segregation of chromosomes both in mitosis and

meiosis and are typically rich in repetitive sequences in many eukaryotes [35].

Equivalent observations to ours on the putative centromere positions was reported in

the YANG genome [15], establishing their correlation to the presence of repetitive rich

DNA regions that were challenging to assemble. A few localized negative recombination

rate estimates were also observed when the LOESS curve under passed the red dotted

line, for example on chromosomes 7, 8, 9 and 12 (Fig 2). These localized negative

estimates coincide with the inconsistent genetic to physical map alignments as

described above and are usually located at the tip of chromosomes. All our results taken

together, on the collinearity between our linkage map and both physical genomes

provide an independent experimental validation of the genome assemblies currently

available for coconut and establish the correspondence of chromosome numbering used

in the two reports (Fig 2).

550　Colocation of a major effect QTL with the FLOWERING LOCUS T

551　(FT) provides evidence for its role in early flowering

Little is known about the underlying mechanisms of early flowering in dwarf

coconut. Knowledge on the complex physiology and genetics of flowering initiation has

come mainly from annual plants including the model species *Arabidopsis* as well as crop

plants, which typically flower and set seeds within a single year [36, 37]. Less is known

about flowering time in perennial plants which typically have distinguishable transition

from a juvenile to an adult phase, the developmental stage that determines the

emergence of flower meristems. It is now well established, however, that the formation

of flower structures is largely conserved across flowering plants, annual and perennial,

with relatively minor variations in the role of key developmental genes [38]. Amongst

the several genes identified as involved in regulation of flowering time, the expression

of FLOWERING LOCUS T (FT) and its homologues has been shown to accelerate the onset

of flowering in a number of plant species [39]. During the course of our study, we came

across a transcriptomics report in coconut showing that the *FLOWERING LOCUS T* (FT)

gene generates different transcripts in tall compared to dwarf coconut, with the shorter

splice variant of FT present exclusively in dwarf varieties but absent in most tall coconuts

[27]. The authors suggested that this FT splice variant could be the putative causal

mechanism for flowering time differentiation between the dwarf and tall coconut types,

although they recommended further investigation and validation to confirm this effect.

By collocating a phenotype-derived major QTL, detected agnostically to any prior gene-

centered hypothesis, our study provides independent validation of the FT gene as the

underlying major functional variants controlling flowering time between the two

coconut varieties.

574   The *FT* locus likely controls other precocity traits in Dwarf
575   coconut

A number of traits were targeted in previous QTL mapping studies in coconut,

such as yield, fruit components, early germination and cuticular wax [9-13]. Flowering

time, however could not be approached because it did not segregate in intermediate F_1

hybrid families and less so in progeny of crosses between Tall cultivars. Two studies to

date have reported on the patterns of genetic variation in F_2 populations of coconut

hybrids, although without attempting QTL mapping. Ample segregation was reported

for a number of vegetative and morphological flowering traits, including height growth

and the presence of a bole, for which the presence of at least one major gene each, with

incomplete dominance was suggested [4, 7]. Curiously, despite its importance as a trait

directly linked to production, time to flower was not assessed. Our study is therefore

the first one to tackle flowering time in coconut by a QTL mapping approach facilitated

by the strong segregation seen in the F_2. In our experiment we could not, however,

measure height growth with confidence and include this trait in the mapping work.

Recently a GWAS was carried out in a collection of Tall and Dwarf cultivars identifying a

major effect QTL for height growth [6]. The leading SNP for the GWAS hit was mapped

to WANG chromosome 12, co-located in the promoter region of the gibberellin (GA)

biosynthetic enzymes GA 20-oxidases (GA20ox) further backed by expression analysis,

validating previous reports [5]. These results strongly support the role of GA20ox in

controlling height growth difference between Tall and Dwarf coconuts. Taken together

with the detection of a major flowering time QTL on chromosome 4 of WANG genome

(chromosome 3 of YANG) (Figs 5 and 6) and its colocation with the FT gene, these results

also suggest that height growth and flowering time in coconut involve different major

effect loci in line with the early proposal of two independent major loci [4] , although it

cannot be ruled out that the FT gene is also involved in height growth (see below).

Petiole length (PL) and the presence of a bole (BOLE) were also assessed in our

study. Both traits are typically used as morphological markers to identify dwarf plants

[2]. In fact, PL and BOLE showed significant direct (0.466) and inverse (-0.502) correlation

with flowering time (DLFS) (Fig 4). Major effect QTLs were also detected for these traits

in the same exact region on chromosome 3 suggesting that the *FT* gene might also be

affecting these traits (Fig 5). The *FT/TFL1* gene family has been extensively studied in

flowering plants. It constitutes a major target of evolution in nature, as a single essential

gene that has played a central role in plant diversification and adaptation [40]. FT

homologs have been identified in many species, demonstrating general conservation of

functions across gymnosperms and angiosperms. It encodes a major florigen, a key

flowering hormone in controlling flowering time. Studies in a number of species have

revealed diverse roles of the FT/TFL1 gene family in plant developmental processes

other than flowering regulation, with a distinct effect on stem and leaf development,

mainly in the inhibition of stem and leaf growth [41, 42]. It is therefore plausible that *FT*

is directly affecting PL and the formation of a bole structure in coconut as well.

Conclusions: implications for coconut advanced breeding lines

From the conventional coconut breeding perspective, offspring of F_1 hybrids

would in principle not be desirable for commercial production due to the wide variation

in yield traits among F_2 coconut trees. Nevertheless, the two previous studies of F_2

populations have reported individuals displaying high husked nut weights, indicating

little or no inbreeding depression, likely resulting from the inbred nature of the ancestral

Dwarf grandparent. These results indicated good potential of extracting recombined

lines from F_2 populations with desirable characters [7, 8]. Still, no developments have

been made to implement this breeding strategy in coconut possibly due to the long time

necessary to develop and eventually propagate such recombined lines.

The results of our study open a fundamentally new perspective in terms of

marker assisted selection (MAS) in coconut breeding. Knowledge of the position and

the genotypes at the SNP markers linked to the flowering time QTL would easily allow

selecting hybrid F_2 plants homozygous for the early flowering haplotype. Homozygosity

for the favorable alleles at SNPs 100226499_C>T-34, 100287206_T>A-55 and

100273632_T>G-15, inherited from the Dwarf parent BGDJ caused a reduction in the

time to flower around 400 days in our experiment. Our data also showed a significant

negative correlation between the necessary days to flowering (DLFS) and NBF (Fig 4).

Early flowering plants displayed some of the highest numbers of fruit per bunch and a

separate QTL was mapped in chromosome 1 which could be selected by a set of linked

SNPs flanking the lead SNP 100237259_A>G-48 (Table 2). This small set of SNPs could be

used for high throughput inexpensive MAS screening of F_2 plants at ultra-early stages of

development or even at the embryo stage. Juvenile tissue of these selected individuals

could be induced to somatic embryogenesis [43] or, more successfully, to shoot culture

[44] for clonal propagation. This would capture the full genetic value of these plants

allowing regenerating thousands of plants indefinitely, providing the coconut industry

with a solution for their current need of homogeneous quality planting material.

Acknowledgments

643 We would like thank the entire staff of field technicians EMBRAPA Tabuleiros Costeiros

644 who were involved in establishing and managing the several field trials, and collecting

645 and organizing the data used in this study. We thank Dr. Orzenil B. Silva-Junior and Dr.

646 Roberto Togawa for assistance with the BLAST analysis against the YANG and Wang et

647 al. genomes respectively.

References

1.   Gunn BF, Baudouin L, Olsen KM. Independent Origins of Cultivated Coconut (Cocos
nucifera L.) in the Old World Tropics. Plos One. 2011;6(6). doi: 10.1371/journal.pone.0021143.
PubMed PMID: WOS:000292033700039.
2.   Batugal P, Bourdeix R, Baudouin L. Coconut Breeding. In: Jain SM, Priyadarshan PM,
editors. Breeding Plantation Tree Crops: Tropical Species. New York, NY: Springer New York;
2009. p. 327-75.
3.   FAOSTAT statistical database [Internet]. Food and Agricultural Organization, Rome (Date
accessed: March 10, 2022). 2021 [cited March 10, 2022]. Available from:
https://search.library.wisc.edu/catalog/999890171702121.
4.   Perera L, Baudouin L, Mackay I. SSR markers indicate a common origin of self-pollinating
dwarf coconut in South-East Asia under domestication. Scientia Horticulturae. 2016;211:255-62.
doi: 10.1016/j.scienta.2016.08.028. PubMed PMID: WOS:000386409800031.

5.   Boonkaew T, Mongkolsiriwatana C, Vongvanrungruang A, Srikulnath K, Peyachoknagul
S. Characterization of GA20ox genes in tall and dwarf types coconut (Cocos nucifera L.). Genes
& Genomics. 2018;40(7):735-45. doi: 10.1007/s13258-018-0682-4.

6.   Wang S, Xiao Y, Zhou Z-W, Yuan J, Guo H, Yang Z, et al. High-quality reference genome
sequences of two coconut cultivars provide insights into evolution of monocot chromosomes
and differentiation of fiber content and plant height. Genome Biology. 2021;22(1):304. doi:
10.1186/s13059-021-02522-9.

7.   Namboothiri CGN, Niral V, Parthasarathy VA, Muralidharan K. Segregation, phenotypic
and genotypic variance and heritability in the F2 population of coconut (*Cocos nucifera*). Indian
Journal of Agricultural Sciences. 2011;81(10):954-6.

8.   Fernando WMU, Perera P. Evaluation of genotypes arising from F2 generation
segregations in dwarf × tall crosses of cocos nucifera. Cord. 1997;13(2):26-35.

9.   Herrán A, Estioko L, Becker D, Rodriguez MJB, Rohde W, Ritter E. Linkage mapping and
QTL analysis in coconut (Cocos nucifera L.). Theoretical and Applied Genetics. 2000;101(1):292-
300. doi: 10.1007/s001220051482.

10.   Lebrun P, Baudouin L, Bourdeix R, Konan JL, Barker JHA, Aldam C, et al. Construction of
a linkage map of the Rennell Island Tall coconut type (Cocos nucifera L.) and QTL analysis for
yield characters. Genome. 2001;44(6):962-70. doi: 10.1139/gen-44-6-962. PubMed PMID:
WOS:000172264900006.

11.   Riedel M, Riederer M, Becker D, Herran A, Kullaya A, Arana-Lopez G, et al. Cuticular wax
composition in Cocos nucifera L.: physicochemical analysis of wax components and mapping of
their QTLs onto the coconut molecular linkage map. Tree Genetics & Genomes. 2009;5(1):53-
69. doi: 10.1007/s11295-008-0168-7. PubMed PMID: WOS:000261265100005.

12.   Baudouin L, Lebrun P, Konan JL, Ritter E, Berger A, Billotte N. QTL analysis of fruit
components in the progeny of a Rennell Island Tall coconut (Cocos nucifera L.) individual.
Theoretical and Applied Genetics. 2006;112(2):258-68. doi: 10.1007/s00122-005-0123-z.
PubMed PMID: WOS:000234026000007.

13.   Ritter E, Rodriguez MJB, Herran A, Estioko L, Becker D, Rohde W. Analysis of quantitative
trait locis (QTL) based on linkage maps in Coconut (Cocos nucifera L.). In: Arencibia AD, editor.
Plant Genetic Engineering: Towards the Third Millennium. Developments in Plant Genetics and
Breeding. 52000. p. 42-8.

14.   Muñoz-Pérez JM, Cañas GP, López L, Arias T. Genome-wide diversity analysis to infer
population structure and linkage disequilibrium among Colombian coconut germplasm.
Scientific Reports. 2022;12(1):2958. doi: 10.1038/s41598-022-07013-w.

15.   Yang Y, Bocs S, Fan H, Armero A, Baudouin L, Xu P, et al. Coconut genome assembly
enables evolutionary analysis of palms and highlights signaling pathways involved in salt
tolerance. Communications Biology. 2021;4(1):105. doi: 10.1038/s42003-020-01593-x.

16.   Inglis PW, Pappas MdCR, Resende LV, Grattapaglia D. Fast and inexpensive protocols for
consistent extraction of high quality DNA and RNA from challenging plant and fungal samples
for high-throughput SNP genotyping and sequencing applications. PLosOne.
2018;13(10):e0206085. doi: 10.1371/journal.pone.0206085.

17.   Sansaloni C, Petroli C, Jaccoud D, Carling J, Detering F, Grattapaglia D, et al. Diversity
Arrays Technology (DArT) and next-generation sequencing combined: genome-wide, high
throughput, highly informative genotyping for molecular breeding of *Eucalyptus*. BMC
Proceedings. 2011;5(Suppl 7):P54. PubMed PMID: doi:10.1186/1753-6561-5-S7-P54.

18.   Crossa J, Federer † W. I.4 Screening Experimental Designs for Quantitative Trait Loci,
Association Mapping, Genotype-by Environment Interaction, and Other Investigations. Frontiers
in Physiology. 2012;3.

19.   Santos GA, Batugal P, Othman A, Baudouin L, Labouisse J-P. Manual on standardized
research techniques in coconut breeding. 1996.

20.   Wei T, Simko V, Levy M, Xie Y, Jin Y, Zemla J, et al. R package 'corrplot': Visualization of
a Correlation Matrix (Version 0.92). 2021.

21.   Resende MDV. Software Selegen-REML/BLUP: a useful tool for plant breeding. Crop
Breed Appl Biotechnol. 2016;16(4):330-9. doi: 10.1590/1984-70332016v16n4a49. PubMed
PMID: WOS:000391702900009.
22.   Van Ooijen JW, Voorrips RE. JoinMap 3.0, Software for the calculation of genetic linkage
maps. P R International. 2001; Wageningen, The Netherlands.
23.   Voorrips RE. MapChart: Software for the graphical presentation of linkage maps and
QTLs. Journal of Heredity. 2002;93(1):77-8. doi: DOI 10.1093/jhered/93.1.77. PubMed PMID:
ISI:000175627300018.
24.   Chakravarti A. A Graphical Representation of Genetic and Physical Maps - the Marey
Map. Genomics. 1991;11(1):219-22. doi: Doi 10.1016/0888-7543(91)90123-V. PubMed PMID:
ISI:A1991GA48800029.
25.   Siberchicot A, Bessy A, Guéguen L, Marais GA. MareyMap Online: A User-Friendly Web
Application and Database Service for Estimating Recombination Rates Using Physical and
Genetic Maps. Genome Biology and Evolution. 2017;9(10):2506-9. doi: 10.1093/gbe/evx178.
26.   Broman KW, Wu H, Sen Ś, Churchill GA. R/qtl: QTL mapping in experimental crosses.
Bioinformatics. 2003;19(7):889-90. doi: 10.1093/bioinformatics/btg112 %J Bioinformatics.
27.   Xia W, Liu R, Zhang J, Mason AS, Li Z, Gong S, et al. Alternative splicing of flowering time
gene FT is associated with halving of time to flowering in coconut. Scientific Reports.
2020;10(1):11640. doi: 10.1038/s41598-020-68431-2.
28.   Santos P, Venancio TM, dos Santos PHD, Ramos HCC, Aredes FAS, Azevedo AON, et al.
Genotyping-by-sequencing technology reveals directions for coconut (Cocos nucifera L.)
breeding strategies for water production. Euphytica. 2020;216(3). doi: 10.1007/s10681-020-
02582-1. PubMed PMID: WOS:000519369500001.
29.   Ferreira A, da Silva MF, Silva L, Cruz CD. Estimating the effects of population size and
type on the accuracy of genetic maps. Genetics and Molecular Biology. 2006;29(1):187-92. doi:
10.1590/s1415-47572006000100033. PubMed PMID: WOS:000236280300033.
30.   Liu HB. Statistical Genomics, Linkage, Mapping and QTL Analysis. Boca Raton, Florida:
CRC; 1998. 611 p.
31.   Myles S, Chia JM, Hurwitz B, Simon C, Zhong GY, Buckler E, et al. Rapid Genomic
Characterization of the Genus Vitis. Plos One. 2010;5(1):e8219. doi:
10.1371/Journal.Pone.0008219. PubMed PMID: ISI:000273554600001.
32.   Allshire RC, Madhani HD. Ten principles of heterochromatin formation and function.
Nature Reviews Molecular Cell Biology. 2018;19(4):229-44. doi: 10.1038/nrm.2017.119.
33.   DeWan AT, Parrado AR, Matise TC, Leal SM. The Map Problem: A Comparison of Genetic
and Sequence-Based Physical Maps. The American Journal of Human Genetics. 2002;70(1):101-
7. doi: https://doi.org/10.1086/324774.
34.   Fierst JL. Using linkage maps to correct and scaffold de novo genome assemblies:
methods, challenges, and computational tools. Frontiers in Genetics. 2015;6. doi:
10.3389/fgene.2015.00220.
35.   Achrem M, Szucko I, Kalinka A. The epigenetic regulation of centromeres and telomeres
in plants and animals. Comparative Cytogenetics. 2020;14(2):265-311. doi:
10.3897/CompCytogen.v14i2.51895. PubMed PMID: WOS:000546682500001.
36.   Komeda Y. GENETIC REGULATION OF TIME TO FLOWER IN ARABIDOPSIS THALIANA.
Annual Review of Plant Biology. 2004;55(1):521-35. doi:
10.1146/annurev.arplant.55.031903.141644.
37.   Roux F, Touzet P, Cuguen J, Le Corre V. How to be early flowering: an evolutionary
perspective. Trends in Plant Science. 2006;11(8):375-81. doi:
https://doi.org/10.1016/j.tplants.2006.06.006.
38.   Khan MRG, Ai X-Y, Zhang J-Z. Genetic regulation of flowering time in annual and
perennial plants. WIREs RNA. 2014;5(3):347-59. doi: https://doi.org/10.1002/wrna.1215.

39.   Putterill J, Varkonyi-Gasic E. FT and florigen long-distance flowering control in plants. Current Opinion in Plant Biology. 2016;33:77-82. doi: https://doi.org/10.1016/j.pbi.2016.06.008.

40.   Pin PA, Nilsson O. The multifaceted roles of FLOWERING LOCUS T in plant development. Plant, Cell & Environment. 2012;35(10):1742-55. doi: https://doi.org/10.1111/j.1365-3040.2012.02558.x.

41.   Wickland DP, Hanzawa Y. The FLOWERING LOCUS T/TERMINAL FLOWER 1 Gene Family: Functional Evolution and Molecular Mechanisms. Molecular Plant. 2015;8(7):983-97. doi: 10.1016/j.molp.2015.01.007. PubMed PMID: WOS:000357615600003.

42.   Freytes SN, Canelo M, Cerdan PD. Regulation of Flowering Time: When and Where? Current Opinion in Plant Biology. 2021;63. doi: 10.1016/j.pbi.2021.102049. PubMed PMID: WOS:000711158100002.

43.   Pérez-Núñez MT, Chan JL, Sáenz L, González T, Verdeil JL, Oropeza C. Improved somatic embryogenesis from Cocos nucifera (L.) plumule explants. In Vitro Cellular & Developmental Biology - Plant. 2006;42(1):37-43. doi: 10.1079/IVP2005722.

44.   Wilms H, De Bièvre D, Longin K, Swennen R, Rhee J, Panis B. Development of the first axillary in vitro shoot multiplication protocol for coconut palms. Scientific Reports. 2021;11(1):18367. doi: 10.1038/s41598-021-97718-1.

Supporting information

S1 File. DArTSeq SNP genotypic data for the full 3714 linkage mapped SNPs in the F_2 population

S2 File. Genotypic data for the 2952 DArTSeq SNP markers linkage mapped and physically mapped to unique positions on the coconut genome assemblies

S3 File. BLAST results of the DArTSeq SNP corresponding sequences, target SNP type and position, against the YANG and WANG genomes.

S4 File. Local recombination rates estimated following the Marey maps construction and LOESS smoothing between the linkage map and the two assembled genomes

S5 File. Raw phenotypic data and estimated BLUPs following the mixed model analysis for the evaluated traits in the F_2 population

S6 File. QTL mapping results obtained with the composite interval mapping method

S7 File. BLAST alignment result of the *Arabidopsis thaliana* putative flowering signals mediating protein FT (At1g65480) mRNA, complete cds against the YANG chromosome 3

S8 File. BLAST alignment result of the *Arabidopsis thaliana* putative flowering signals mediating protein FT (At1g65480) mRNA, complete cds against the WANG chromosome 4

Figure 1

Figure 2

Figure 3

	HSLH	PL	RL	PW	PT	BOLE	NBF	DLFS
NL	0.772	-0.091	0.519	0.368	0.224	0.368	0.421	-0.484
HLH		0.119	0.706	0.435	0.362	0.430	0.369	-0.349
PL			0.043	0.053	0.146	-0.207	-0.348	0.479
RL				0.514	0.505	0.292	0.255	-0.132
PW					0.403	0.141	0.066	-0.067
PT						0.081	-0.044	0.110
BOLE							0.347	-0.417
NBF								-0.519

Figure 4

Figure 5

Figure 6