Sexually Dimorphic Influence of Circadian Pacemaker Neurons on Behavioral Rhythms

Aishwarya Ramakrishnan Iyer[1,2], Eva Scholz-Carlson[1,2], Evardra Bell[2], Grace Biondi[2], Shlesha Richhariya[3], and Maria P. Fernandez[1,2,*]

1- Department of Biology, Indiana University Bloomington. Bloomington, IN
2- Department of Neuroscience and Behavior, Barnard College. New York City, NY
3- HHMI, Brandeis University. Waltham, MA

*Corresponding author: fernanm@iu.edu

Abstract

The circadian system regulates the timing of multiple molecular, physiological, metabolic, and behavioral phenomena. In *Drosophila,* as in other species, most of the research on how the timekeeping system in the brain controls the timing of behavioral outputs has been conducted in males, or sex has not been included as a biological variable. A critical set of circadian pacemaker neurons in *Drosophila* release the neuropeptide pigment-dispersing factor (PDF), which functions as a key output factor in the network with complex effects on other clock neurons. Lack of *Pdf* or its receptor, *PdfR,* results in most flies displaying arrhythmicity in activity–rest cycles under constant conditions. However, our results show that female circadian rhythms are less affected by mutations in both *Pdf* and *PdfR*. Crispr-Cas9-mediated mutagenesis of *Pdf,* specifically in ventral lateral neurons (LN_vs), also has a greater effect on male rhythms. We tested the influence of M-cells on the circadian network and showed that speeding up the molecular clock specifically in M-cells led to sexually dimorphic phenotypes, with a more pronounced effect on male rhythmic behavior. Our results suggest that the female circadian system is more resilient to manipulations of M-cells and the PDF pathway, suggesting that circadian timekeeping is more distributed across the clock neuron network in females.

Keywords: circadian rhythms, sex differences, PDF, *Drosophila,* sleep/wake cycles, oscillator hierarchies

Introduction

Differences in neuronal circadian timekeeping between sexes remain relatively unexplored, despite the expanding body of research highlighting the influence of sex on the mechanisms underlying neuronal control of behavior [1]. In mammals, steroid hormones display daily, clock-driven changes in abundance, and while these sex hormones are not required to maintain rhythms, they differentially influence the amplitude of activity behavior between the sexes [1, 2]. Furthermore, structural and functional sex differences have been observed in brain areas that receive direct input from the brain's circadian timekeeping center [2, 3]. Research in humans has also revealed significant sexual dimorphism: men tend to have lower-amplitude endogenous rhythms than women [4], are less resilient to nocturnal sleep disruptions, and spend less time asleep [5].

In mammals, the main circadian pacemaker resides in the suprachiasmatic nuclei (SCN), which in mice consist of a network of ~20,000 neurons (reviewed in [6]). The *Drosophila* circadian clock network has ~150 neurons and is the functional equivalent of the mammalian SCN [7, 8]. Each circadian clock neuron has an intracellular molecular timekeeping mechanism based on a transcriptional□translational feedback loop: the genes *Clock* (*Clk*) and *cycle* (*cyc*) promote rhythmic transcription of several key genes, including *Period* (*Per*) *and Timeless* (*Tim*), which build up daily and inhibit their own transcription [9]. Multiple kinases that act on components of these clock proteins and can affect the pace of the molecular clock have been identified. One such kinase is *Doubletime* (DBT), which binds to and phosphorylates PER, regulating its nuclear accumulation [10, 11].

The fly clock network consists of *lateral neurons (LNs)*, which include ventrolateral (LN_v), dorsolateral (LN_d), and lateral posterior neurons (LPNs), as well as three groups of *dorsal neurons* (*DN1, DN2, and DN3*), some of which can be further subdivided [8, 12-15]. The ventral and dorsal LNs are sufficient to produce the normal endogenous bimodal rhythm of sleep and activity [16, 17]. The four small LN_vs (s-LN_vs) are usually referred to as morning cells (M-cells) since they control the morning peak of activity under light□dark cycles (LDs). These cells are also essential for maintaining rhythmicity under free-running conditions [18, 19]. The evening peak is controlled by the LN_ds and a *Pdf*-negative LN_v, the 5th LN_v (E-cells) [16, 20, 21]. Some

DNs also contribute to the timing and amount of sleep via the modulation of M and E cells [22-
25].

The release of the circadian neuropeptide Pigment Dispersing Factor (PDF) by s-LN$_v$s is
essential for endogenous circadian timekeeping. A *Pdf* null mutation, *Pdf^{01}*, results in a
substantial fraction of arrhythmic flies [18], desynchronization of molecular oscillations [26, 27],
and phase changes in the electrical activity of some clock clusters, most notably the LN$_d$s [28].
Loss of *PdfR* also leads to loss of behavioral rhythms [29-31]. Interestingly, PDF and PDFR also
regulate behaviors that are sex-specific or sexually dimorphic. Rival-induced long mating
durations require PDF expression in s-LN$_v$s, PDFR expression in a subset of LN$_d$s, and NPF
expression in LN$_d$s [32]. PDF controls rhythms in the sexually dimorphic pheromone profiles
produced by oenocytes [33] and is involved in long-term mating suppression in males [34]. Both
PDF and PDFR contribute to geotactic behaviors [29], and the phenotypes of *Pdf^{01}* mutants are
sexually dimorphic, with males showing a more extreme negative geotaxis phenotype [35].

Sexual dimorphism in *Drosophila* sleep/wake cycles has been studied mostly under LD
cycles. Males exhibit lower levels of activity and more sleep during the light phase [36-38]. This
increase in midday sleep is due to the activity of a subset of sleep-promoting DN1s, which are
more active in males [36] and receive input from the male-specific P1 neurons that control male
courtship [37]. Unlike studies on circadian rhythms, *Drosophila* sleep research often involves
only females. Males also have an earlier and more pronounced morning peak and a larger phase
angle between the morning peak and the evening peak [39]. Under conditions of constant
darkness and temperature (DD), males of several wild-type strains have a small but significant
reduction in the free-running period (FRP) relative to females of the same strain [39]. Moreover,
males are more likely to retain a bimodal activity pattern in DD [39]. A recent transcriptomic
analysis of *fruitless* (*fru*)-expressing neurons revealed clusters that are enriched for circadian
clock genes [40]. Interestingly, only the male dataset revealed coexpression of *fru*- and circadian-
related genes. A previous study reported that DN1s express the male-specific Fru^M protein [41]
and that the number of cells in the DN1$_p$s cluster is sexually dimorphic [42]. In addition, the E3
subset of LN$_d$s has been shown to be dimorphic in its expression of the neuropeptide NPF [43,
44].

Given the sexually dimorphic roles of neuropeptides, including PDF, in other behaviors [45], we asked whether females were similarly affected by manipulations of the *Pdf/PdfR* pathway. We found that female circadian rhythms are less affected by null mutations in both *Pdf* and *PdfR* and that similar effects are observed via CRISPR-Cas9-mediated *Pdf* mutagenesis, specifically in the LN$_v$s. Moreover, speeding up the molecular clock in the LN$_v$s via expression of *DBT^s* leads to an advance of the morning peak in males but not in females, and the pace of the free-running period of activity is significantly shortened only in males. Taken together, our results show that the female circadian system is more resilient to manipulations in the PDF pathway and suggest that *Pdf+* neurons play a more dominant role in the male than in the female circadian network.

Results

Mutations in PDF and PDFR lead to sexually dimorphic phenotypes.

A null mutation in *Pdf* results in pronounced behavioral phenotypes in *Drosophila* males [18]. We assayed the locomotor activity rhythms of *Pdf^{01}* females under free-running conditions (DD) and found that a large proportion of the experimental females were still rhythmic (Fig. 1 A-C). The rhythmic power of experimental flies was significantly reduced in both sexes (Fig. 1D), but the effect was less pronounced in females (Fig. 1E), suggesting that *Pdf^{01}* females have more consolidated rhythms than *Pdf^{01}* males. Mutant females that were rhythmic had a slightly, but significantly, shorter free-running period than the controls (Table 1). This phenotype was not always observed in experimental males (Table 1), consistent with a recent study [46]. Sleep cycles under DD also appeared to be more consolidated in females (Fig. 1F). *Pdf* mutants have increased sleep, and this effect is mediated by PDF acting on the LN$_v$s themselves [47]. We found that both sexes show an increase in total sleep in LD, but the effect was more pronounced in females (Fig. 1G-I). While the increase in sleep in males was most prominent at midday, females exhibited increased sleep throughout most of the light phase (Fig. 1G).

To rule out the presence of remnant PDF expression in *Pdf^{01}* females, we stained the brains of control and experimental males and females with an anti-PDF antibody. We did not observe any

traces of PDF in experimental flies of either sex, even with increased laser intensity (Fig. S1A). PDF accumulates rhythmically in the dorsal termini of the s-LN$_v$ projections in a time-of-day-dependent manner both in LD and DD [18, 19]. To determine if there were differences in the amplitude of PDF cycling between the sexes in a wild-type background (Canton-S), we dissected control males and females on the third day under DD at 6 timepoints over a 24-hour cycle. Using a COSINOR-based curve fitting method [48], we found that both males and females have clear 24-hour rhythms in PDF cycling in their dorsal projections, with no significant sex differences in amplitude (Fig. S1B-C, Table 2).

Next, we asked whether the effects of a *Pdf* receptor mutation (*PdfR*) on activity and sleep were also sexually dimorphic. The expression of PDFR, a GPCR, can be detected in most clock neurons, with the exception of 3 LNds, half DN1ps, and some DN3s [49], which coincides with *Cryptochrome* expression in clock neurons [49]. The *han^{5304}* mutant is a *PdfR* hypomorph and exhibits *Pdf^{01}*-like behavioral phenotypes under both LD 12:12 and DD [29-31]. Under DD, both *Han* males and females showed a significant reduction in rhythmicity compared with the controls, but there was a greater proportion of rhythmic females (~65%) than males (~16%) (Fig. 1J). The free-running period of the experimental flies was significantly shorter for both sexes (Table 3), as reported previously for males. Rhythmic power was significantly lower than that of the controls for both *han^{5304}* males and females (Fig. 1K), but the effect was more pronounced in males, suggesting that females have more consolidated rhythms (Fig. 1L). Similar to the effect of the *Pdf* mutation, *han^{5304}* flies showed significantly higher levels of LD sleep than controls, and this effect was also more pronounced in females (Fig. 1M☐O). Taken together, these results suggest that female circadian rhythms are less affected by the loss of both *Pdf* and *PdfR*. However, the LD sleep phenotypes were more pronounced in females.

CRISPR-Cas9-mediated *Pdf* mutagenesis has more pronounced effects on male behavior.

In the *Pdf* null mutant, background effects could contribute to the sexual dimorphism observed in behavioral rhythms. We therefore employed a tissue-specific CRISPR-Cas9-mediated knockout of *Pdf* in both males and females, as described in a recent study that focused on males [50]. To assess the efficiency of the manipulation, we stained for PDF in flies that

constitutively expressed *Pdf* gRNA and Cas9 in *Pdf*+ neurons. This experiment was conducted at 28°C, as this temperature was more effective at mimicking the behavioral phenotypes of the Pdf^{01} mutant males and allows direct comparison with adult-specific manipulation, which requires switching from 18°C to 28°C.

PDF was reduced in the s-LN$_v$s in both sexes (Fig. 2 A-C), although in most experimental brains, we noted faint staining in the dorsal projections of at least one s-LN$_v$ in at least one brain hemisphere (Fig. 2A). We quantified PDF intensity in the cell bodies of the sLNvs and found that the signal intensity was reduced in both sexes in a similar manner (Fig. 2B-D). PDF expression within the large LN$_v$s was less affected and could be detected in 2–3 l-LNv cell bodies in most brains (Fig. 2A). In addition to behavioral phenotypes, Pdf^{01} mutation leads to pronounced misrouting of s-LN$_v$ projections in male flies [51] . We employed a transgene expressing a red fluorescent protein under the *Pdf* regulatory sequence [52] and observed faint projections occasionally defasciculating from the main bundle in one or both hemispheres in *Pdf >Pdf*-g; *Cas9* flies of both sexes (Fig. 2A, middle panels).

We analyzed activity☐rest rhythms in *Pdf >Pdf*-g; *Cas9* flies and found that while experimental males were largely arrhythmic, a greater proportion of experimental females remained rhythmic (Fig. 3A-B). The free-running period of the experimental males was not significantly different from that of the controls, but a wide range of periods were observed. Compared with parental controls, *Pdf >Pdf*-g; *Cas9* females had significantly shorter free-running periods (Fig. 3C). The rhythmic power was significantly lower in experimental flies of both sexes (Fig. 3D), but the effect was less pronounced in females (Fig. 3E). Sleep under LD was not affected (Fig. S2 A-B), whereas sleep under DD was similarly increased in males and females (Fig. S2 C-E).

We next restricted the CRISPR mutagenesis of *Pdf* to the small LN$_v$s via a specific driver from the split-Gal4 collection generated by the Rubin Laboratory (*SS00681*-Gal4). *Pdf* knockdown in the s-LNv resulted in most males being arrhythmic (~30% rhythmicity), whereas the experimental females were ~45% rhythmic (Fig. S3A). The percentage of rhythmic flies was significantly lower than that of both controls for experimental males but not for females (Fig.

S3A). The free-running period of the experimental males was shorter than that of their respective control flies, whereas the experimental females did not differ from their controls (Fig. S3B). The rhythmic power of the males was lower than that of the controls, but was not different between the experimental females and their controls (Fig. S3C). This suggests that PDF from the s-LNv is important for the behavioral and sex-specific differences observed in Pdf^{01} mutants.

We next conducted adult-specific CRISPR-Cas9 knockdown of Pdf (Fig S4). Although a previous study in males did not establish a connection between developmental misrouting caused by Pdf downregulation and the disruption of behavioral rhythms in adulthood [51], sex-specific differences observed upon Pdf loss may result from developmental effects. The AGES system was previously employed to rule out the developmental effects of the loss of Pdf [50], but a recent study showed that AUXIN exposure induces short- and long-term changes in physiology and behavior [53]. Therefore, we employed a temperature-sensitive Gal80 variant with ubiquitous expression to conditionally inhibit Gal4-mediated expression of the Pdf guide and Cas9 [54]. This allows for the temporal control of UAS transgenes since Gal80^{ts} is active at lower temperatures but inactive at higher temperatures. We raised the flies at 18°C and transferred them to 28°C immediately after eclosion so that Pdf knockout occurred only in adult flies. Consistent with what was reported for males by Gorostiza et al. [51] via RNA-mediated interference (RNAi) of Pdf, we did not observe misrouting in experimental flies (Fig. S4A). While PDF levels were significantly reduced in both males and females, the manipulation was substantially more effective in males (Fig S4B-C). Therefore, although the circadian behavioral phenotypes were again more pronounced in males than in females (Fig S4D-F), this is likely due to higher remaining PDF expression in experimental females.

Speeding up the M-cell clock leads to a coherent period shortening only in males.

Next, we sought to determine if the influence of the Pdf-releasing cells themselves was sexually dimorphic. While PDF is released from both large and small LN$_v$s, only s-LN$_v$s (Morning cells) play key roles in regulating free-running rhythm properties [16]. In males, manipulations that change the pace of the clock specifically in the LN$_v$s result in changes in the phase of the morning peak of activity and in the free-running period [20, 55]. We expressed the

Doubletime 'short' (*DBT*^s) allele [56] under the *Pdf-Gal4* driver (Fig. 4A) and analyzed the effects on behavior in both sexes. We found that $Pdf > DBT^s$ males, but not females, have an advanced phase of the morning peak of activity (M-peak, Fig. 4B-C). This suggests that under LD cycles, the M-oscillator is more effective at setting the phase of male than female behavior.

Under free-running conditions, both $Pdf > DBT^s$ male and female flies showed a significantly lower percentage of rhythmic flies than their controls (Fig. 4D), but there were fewer rhythmic females (~40%) than males (~65%) (Fig. 4D-E). The free-running period of most $Pdf > DBT^s$ males was ~18.5 hours (Fig. 4F), whereas the $Pdf > DBT^s$ females showed a large proportion of individuals with a period of ~24 hours, reflecting the pace of the molecular clock in the rest of the clock network (Fig. 4F). Some $Pdf > DBT^s$ males (~23%) and a smaller proportion of females (~6%) also presented complex rhythms (Table 4). The average period value of the second period component (which has a lower power value) was ~24.07 ± 0.4 for the experimental males and 19.7 ± 0.4 for the experimental females. Neither male nor female control flies exhibited complex rhythms (Table 4). Among rhythmic flies, both $Pdf > DBT^s$ males and females had lower rhythmic power than the controls (Fig. 4G), with no difference between the sexes (Fig. 4H). As a control, we expressed *DBT*^s via *Clk856-Gal4* which is expressed in most clock neurons (Fig. S5A). Speeding up the molecular clock in most clock neurons significantly advanced both the morning and evening activity peaks of both males and females (Fig. S5B-D). There were no significant differences between experimental males and females in the percentage of rhythmic flies, rhythmic power, or shortening of free-running periods following DBT expression in all clock neurons (Fig. S5E-G).

These results support the notion that M-cells are more dominant in the male than in the female circadian network. Previous studies have shown that blocking synaptic neurotransmission by expressing the tetanus toxin light chain (TeTxLC) in small and large LN$_v$s affects male activity rhythms, likely in a *Pdf*-independent manner [57-59]. We analyzed male and female *Pdf >* TeTxLC flies and found that neither sex significantly changed the ability to maintain rhythmicity under free-running conditions (Fig S6A-B). Both sexes significantly lengthened the free-running period, but the effect was more pronounced in males (Fig. S6C-D). Rhythmic power was not affected in experimental flies of either sex (Fig. S6E).

We next asked whether changing the pace of the clock in LNds (E-cells) via DBT^S expression

also had sexually dimorphic effects on behavior. These cells can be subdivided into at least 3

different clusters on the basis of their anatomy [8], physiology [55], transcriptomic profiles [13]

and connectivity patterns [15]. The PDFR-expressing E1 and E2 clusters have been shown to

regulate evening activity under LD [60, 61] and to be able to maintain free-running activity

rhythms in the absence of a functional clock in M cells [62, 63], whereas the behavioral role of

the E3 cluster remains unknown. We used the MB122-B split-gal4 driver to target the E1 and E2

subsets (Fig. 5A) and found that while expressing DBT^S in this group of evening cells

significantly advanced the phase of the E-peak in experimental flies of both sexes (Fig. 5B-C),

the effect was more pronounced in females (Fig. 5D). Speeding up of clocks in the E1 + E2 LN_ds

did not significantly alter the free-running period or rhythmic power of experimental flies of

either sex (Fig. 5E-F). M-cells have been shown to be the dominant oscillators in DD and to

regulate rhythm properties such as persistence and the free-running period of endogenous

locomotor rhythms to a large extent [16, 20, 21, 55], although manipulations of other clock cells

can affect rhythm properties to some extent [55, 64]. Thus, speeding up the clock in the PDFR$^+$

E1 and E2 clusters leads to similar behavioral phenotypes in males and females under free-

running conditions. Taken together, these results indicate that the relative influence of the M and

E subsets of clock neurons are sexually dimorphic.

272
273   **Discussion**
274

The critical importance of considering sex as a biological variable has gained increasing

recognition in biomedical research [65, 66]. Bias toward male subjects is particularly prevalent

in neuroscience, with single-sex studies using male animals outnumbering those using female

animals at a ratio of 5.5:1 [67]. This disparity extends to chronobiology, resulting in a limited

understanding of how sex affects circadian organization in the nervous system. However, work

from several laboratories has revealed sexual dimorphism within the SCN and in its input and

output pathways [1, 68]. Sex differences in SCN morphology have been described in both animal

models and humans, and sex differences in SCN electrical activity and steroid hormone receptors

have also been reported (reviewed in [2]). Notably, sex differences in the number of SCN

neurons that express the neuropeptide vasoactive intestinal polypeptide (VIP) and in *Vip* mRNA *e*xpression have been reported (reviewed in [2]). The roles of the mammalian VIP and the *Drosophila* PDF in circadian physiology are highly similar, although neither these peptides nor their receptors are sequence orthologs [69].

Several studies have shown the importance of PDF in generating coherent rhythms of ~24-hour periodicity. Here, we report that females lacking *Pdf* or its receptor *PdfR* are more likely to maintain consolidated activity-rest behavior than males. This could be because of sex differences in PDF signaling mechanisms, PDFR expression, or the influence of other clock neurons within the network. We did not detect sex differences in the overall levels or temporal patterns of PDF accumulation, suggesting that these differences might not contribute to the sexually dimorphic effects of the loss of *Pdf* expression. In males, other neuropeptides are known to act in concert with PDF to maintain consolidated rhythms in the network, although none of them have as profound an effect as PDF in regulating activity-rest rhythms in DD [46, 70]. Single mutants of DH31 and CCHamide1 do not affect activity rhythms by themselves, but the double mutants of these neuropeptides along with Pdf^{01} ($DH31^{01}/Pdf^{01}$ and $Pdf^{01}/CCHa^{SK8}$) are almost completely arrhythmic, suggesting that these neuropeptides act hierarchically in the network, with PDF being at the top of that hierarchy [46, 70]. The importance of PDF relative to other peptides released by clock neurons may also be sexually dimorphic.

To overcome the possible pleiotropic effects of Pdf^{01} and *PdfR* mutations, we employed a constitutive knockout of *Pdf* in ventral lateral neurons via the CRISPR-Cas9 method. Although CRISPR manipulation was only partially effective at eliminating PDF expression, it nevertheless produced phenotypes reminiscent of those produced by the Pdf^{01} mutation. We observed faint staining in the dorsal projections of at least one s-LN$_v$ in at least one hemisphere in most brains, and PDF staining in a single s-LN$_v$ projection reaching the dorsal brain has been shown to be sufficient for behavioral rhythms [71]. Experimental flies in which *Pdf* was knocked out starting at the onset of promoter expression early in development showed extensive misrouting of their dorsal termini, similar to what has been reported for Pdf^{01} males [51]. Instances of s-LN$_v$ misrouting have also been observed in other core clock mutants, such as per^{01} and tim^{01} [72] and cyc^{01} [73]. We observed misrouting in experimental flies of both sexes (data not shown), and no

correlation between misrouting and behavioral phenotypes was found by others for Pdf^{01} males [51]. Importantly, $Pdf > Pdfg;Cas9$ manipulation recapitulates the sexually dimorphic circadian phenotypes of Pdf^{01} mutants: a larger fraction of females are rhythmic, and females exhibit greater rhythm power.

In males, changing the speed of the M cell clock leads to phase changes in the morning peak under LD [60]. To determine whether M-cell manipulations also have sexually dimorphic effects on behavior, we sped up the molecular clock by expressing the *Doubletime short* allele ($DBT^{s)}$). Surprisingly, our results revealed that speeding up the clock of M-cells advances the phase of the morning peak in males, whereas the female morning peak phase is not affected. These results support previous studies conducted in males on the role of M-cells in regulating the morning peak of activity [16, 21, 60] and suggest that M-cells are unable to regulate the phase of morning activity in the same way in females. In DD, males had largely coherent short-term rhythms, and the majority (65%) were rhythmic. In contrast, only 40% of the females were rhythmic, and their period showed a bimodal distribution, with most flies exhibiting an ~24-hour period. These findings further support the notion that M-cells are less dominant than the circadian clock network in females. A possible explanation for this is that other clock neurons are able to "resist" their influence, and the conflict between the fast-paced M cell clock and the ~24 clock other clock neurons is what leads to greater arrhythmicity in females.

The expression of a tetanus toxin light chain construct in flies blocks neurotransmission by binding and cleaving the synaptic protein Synaptobrevin [74]. Expressing tetanus toxin in ventral lateral neurons does not affect behavioral rhythmicity and lengthens the free-running period in males, as reported in previous studies [58, 59, 75]. The behavioral phenotypes resulting from the blockade of synaptic transmission differ from those resulting from the loss of PDF [18] or the ablation of *Pdf*-expressing neurons [21], possibly because tetanus toxin affects classical transmission and not the dense core vesicle-mediated release of neuropeptides such as PDF [76]. Abrogating the dorsal termini of the small ventral lateral neurons, where most of the output synapses are found [15, 76], does not result in behavioral phenotypes similar to those of Pdf^{01} under either LD or DD [22]. Our results show that blocking synaptobrevin-dependent synaptic transmission in M-cells does not affect rhythmicity but rather lengthens the free-running period.

The period lengthening is more pronounced in males, supporting the notion that M-cells have a
greater influence on the circadian network in males.

Cryptochrome and *PdfR*-expressing clusters of evening cells—the sNPF-expressing E1 cluster
and the ITP-expressing E2 cluster [77]—have roles in setting the phase of the E-peak under LD
and sustaining behavioral rhythms in the absence of a functional molecular clock in M-cells [62,
63]. To test whether these cells have a differential influence on the network in males and females,
we expressed DBT^S under a driver that is expressed specifically in the E1 and E2 subsets of
LN_ds. Our results showed that speeding up the clocks in the E1+E2 clusters resulted in a phase
advance in the evening peak of activity in both sexes, but the effect was more pronounced in
females. Neither MB122 > DBT^S males nor females presented phenotypes under free-running
conditions. A possible reason for the behavioral differences observed between the sexes could be
redundancy in females, such that the network is not as dependent on PDF or M-cells for
timekeeping. This finding suggests that the female network could have a more distributed mode
of timekeeping throughout the circadian clock network.

Across species, sex differences in the circadian timing system are largely related to the
regulation of reproduction-related behaviors. In mammals, the SCN determines the timing of the
release of reproductive hormones and influences the timing of mating (reviewed in [68]) and
aggression [78]. In *Drosophila*, the circadian clock controls the timing of sex-specific and
sexually dimorphic behaviors, such as male courtship [79] and female sexual receptivity [80] and
egg laying [81]. This regulation of rhythmic behaviors requires connectivity between clock
neurons and downstream sex-specific circuits. For example, the $DN1_p$ cluster, which has been
shown to be more active in males [36], is functionally connected to the male-specific *fru*-
expressing P1 neurons that regulate male courtship [37]. In females, Allostatin C-producing
$DN1_p$s have been shown to connect to downstream targets to control rhythms in oogenesis [82],
and the Janelia female hemibrain connectome revealed that the LN_ds form connections with the
doublesex-expressing PC1 cluster [15]. Our data suggest that the relative hierarchy of circadian
oscillators is sexually dimorphic, with a less dominant M oscillator in females. This pattern of
circadian timekeeping may serve an adaptive purpose, ensuring the precise timing of essential

female-specific behaviors crucial for reproductive fitness, such as sexual receptivity and egg

laying.

379
Acknowledgments

We are grateful to Charlotte Helfrich-Forster, Abhilash Lakshman, Cahir O'Kane, Jeff Price,

and Paul Taghert for helpful discussions and Orie Shafer and members of the Fernandez Lab for

helpful comments on the manuscript. We also thank Justin Blau, Aljoscha Nern, Gerry Rubin,

Michael Rosbash, and Paul Taghert for sharing fly lines. The mouse anti-PDF antibody was

obtained from the Developmental Studies Hybridoma Bank, created by the NICHD of the NIH

and maintained at The University of Iowa, Department of Biology, Iowa City, IA 52242. Stocks

obtained from the Bloomington *Drosophila* Stock Center (NIH P40OD018537) were used in this

study. This work was supported by an NIH grant (R01NS118012) and an NSF Grant (IOS-

2239994) to M.P.F. and a Beckman Foundation Scholarship to E.S-C.

Materials and methods

Fly lines and rearing.

All the genotypes were reared on corn syrup soy media (Archon Scientific; Durham, NC) under LD 12:12 cycles at 25°C unless specified otherwise (see figure legends for details). The fly lines used in this study were *Canton-S*, w^{1118}, Pdf^{01}, $PdfR^{01}$, *Pdf-RFP*, *Pdf-Gal4; tub-Gal80^{ts}*, UAS-*Cas9;* UAS-*Pdfg*, *Pdf-Gal4*, UAS-*DBT^{s}, UAS TeTxLC, Clk856Gal4, s-LNvGal4,* and *MB122B-Gal4.* See the fly lines and reagents table below for more details.

Activity☐rest behavior recording and analysis.

Individual male and virgin female flies (3–5 days old) were housed in glass locomotor tubes containing 2% agar☐4% sucrose food on one end and yarn on the other end. Locomotor activity was recorded using Drosophila activity monitors (DAM, Trikinetics, Waltham, United States of America). The experiments were conducted in Tritech or Percival incubators under controlled light and temperature conditions. Flies were entrained to 12:12 LD cycles for at least 5 days and then transferred to constant darkness (DD) for at least 7 days at a constant temperature of 25°C unless otherwise specified (see figure legends for details). The raw data obtained from the DAM system were scanned and binned into activity counts of 15 min intervals via the DAM File scan. The data were analyzed via the CLOCKLAB software (Actimetrics, Wilmette, IL).

The values of period and rhythmic power were calculated for a period of 7 days via a chi-square periodogram with a cutoff of *p =0.01*. The rhythmic power for each designated rhythmic fly was determined by subtracting the chi-square significance value from the power of the periodogram. Flies that did not exhibit a periodicity peak above the significance threshold were categorized as "arrhythmic," and their period and rhythmic power were not included in the analysis. The values of the morning and evening peaks were calculated via PHASE software [83]. The total LD sleep values for all the genotypes were calculated for a period of 3 days (LD days 2--4) via the PHASE software. Representative actograms were generated via ClockLab, and activity plots were generated via PHASE. The period, rhythmic power, total sleep, and phase values of all the flies for a particular experimental genotype were compared against the background or parental controls via either the Mann☐Whitney test or the Kruskal☐Wallis

ANOVA followed by the Dunn's multiple comparisons test. The details of the statistical
comparisons and the number of flies used in a given experiment are indicated in their respective
figure legends. The number of rhythmic flies of the experimental genotype was compared against
their respective background or parental controls via Fisher's exact test. All the statistical analyses
were performed via GraphPad Prism 9.0.

Immunohistochemistry

The brains of adult male or female flies were dissected in ice-cold Schneider's insect media
(S2) and fixed immediately after dissection in 2% paraformaldehyde (PFA) in S2 media for 30
minutes at room temperature. The fixed brains were washed (three washes of 10 minutes each)
with 0.3% PBS-Triton X 100 (PBS-TX) and then treated with blocking solution (5% normal goat
serum made in 0.3% PBS-TX) for 1 hr at room temperature. The brains were then incubated with
primary antibodies at 4°C for 24 hrs. The primary antibodies used were anti-PDF (mouse,
1:3000, C7, DSHB) and anti-RFP (rabbit, 1:2000, Rockland Immunochemicals). After
incubation, the brains were subjected to six washes with 0.3% PBS-TX and incubated with Alexa
Fluor-conjugated secondary antibodies overnight at 4°C. The following secondary antibodies
were used: goat anti-mouse 488 (1:3000, Invitrogen) and goat anti-rabbit 568 (1:3000,
Invitrogen). After incubation, the brain samples were washed 6 times with 0.3% PBS-TX,
cleaned and mounted on a clean glass slide using Vectashield mounting media.

Image acquisition and analysis

The brains were imaged via a confocal microscope (Olympus FV3000) with an Olympus
UPLanXApo 20x or 40x objective. Image analysis was performed via Fiji software [84]. In the
samples, small and large ventral lateral neurons were classified on the basis of their anatomical
locations and expression of the PDF. PDF intensities in these cells were measured by selecting
the slice of the Z-stack that showed the maximum intensity, drawing a region of interest (ROI)
around the cells and measuring their intensities. Three to four separate background values were
also measured around each cell, and the final intensity was taken as the difference between the
cell intensity and the average background.

For quantification of the PDF in the dorsal projections, a rectangular box was drawn as the
ROI starting from the point where the PDF projection turns into the dorsal brain and the intensity
is measured. Three to four background values were also measured around the projection. The
intensity values obtained from both hemispheres for each cell type for each brain were averaged
and used for statistical analysis. PDF intensity from the s-LN$_v$ was compared between the
experimental and control genotypes via a Mann☐Whitney test. To estimate different aspects of
rhythmicity in PDF oscillations in the dorsal termini of s-LN$_v$ in males and females, we used a
COSINOR-based curve-fitting method (Cornelissen, 2014). COSINOR analysis was
implemented via the CATCosinor function from the CATkit package written for R [85].

Fly lines and reagents.

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms/Strains		
Canton-S	Bloomington *Drosophila* Stock Center	BDSC 64349
w^{1118};+;+	Bloomington *Drosophila* Stock Center	BDSC 3605
yw;Pdf-RFP,Pdf-Gal4;Tub-gal80^{ts}	Justin. Blau, NYU	
w;;Pdf^{01}	Paul Taghert, Wash U Med. School	
w PdfR^{01};;	Paul Taghert, Wash U Med. School	
w; Pdf-gal4; +	Paul Taghert, Wash U Med. School	
;UAS Cas9; pdfg	Michael Rosbash, Brandeis University	BDSC 99650
w;+;DBT^s	Jeffrey Price, University of Missouri	

w; MB122B-Gal4; +	Gerry Rubin, HHMI Janelia Research Campus	
w; Clk856-Gal4; +	Orie Shafer, CUNY ASRC	
s-LNv Gal4	Gerry Rubin, Janelia Research Campus	SS00681-Gal4
w; UAS TeTxLC; +	Cahir O'Kane, University of Cambridge	
Antibodies		
Rabbi anti-RFP (1:2000)	Rockland	#600-401-379-RTU
Mouse anti-PDF (1:3000)	Developmental Studies Hybridoma Bank	
Anti-rabbit Alexa-568 (1:3000)	Thermo Fisher	A11036
Anti-mouse Alexa-488 (1:3000)	Thermo Fisher	A11029
Software		
Fiji	http://fiji.sc	RRID: SCR_002285
MATLAB R2022b	MathWorks, Natick	RRID: SCR_001622
GraphPad Prism 9.0	GraphPad Software	RRID: SCR_002798
DAM FileScan	Trikinetics	
ClockLab	Actimetrics	RRID:SCR_014309
Chemicals, Peptides, and Recombinant Proteins		
Vectashield Mounting Medium	Vector Laboratories	#H-1000-10
Premix PBS Buffer (10x)	Sigma☐Aldrich	Cat# 11666789001
2% Paraformaldehyde (PFA)	Sigma☐Aldrich	47608-250ML-F

Triton- X-100	Bio Basic	CAS#9002-93-1
Schneider's *Drosophila* Medium (S2)	Thermo Fisher	21720024
Other		
DAM2 *Drosophila* Activity Monitors	Trikinetics	
DAM Drosophila Environmental Monitors	Trikinetics	

Figure legends.

Figure 1: Mutations in *Pdf* and *PdfR* lead to sexually dimorphic phenotypes. (A)
Representative double-plotted actograms of w^{1118} (left) and Pdf^{01} (right) male flies subjected to 6
days of LD followed by seven days of DD. **(B)** Representative actograms of w^{1118} (left) and Pdf^{01}
(right) female flies subjected to six days of LD followed by seven days of DD. **(C)** Percentages
of rhythmic flies are plotted for control (w^{1118}) and Pdf^{01} males (left, n = 86 (w^{1118}), n = 81
(Pdf^{01})) and females (right, n = 75 (w^{1118}), n = 101 (Pdf^{01})). The error bars represent the SEM
values plotted across three replicate experiments. **(D)** Rhythmic power of control (w^{1118}) and
Pdf^{01} male and females was calculated via a Chi-square Periodogram. **(E)** The differences in
rhythmic power between experimental males and females and their respective controls are
plotted. **(F)** Average sleep plots of flies over seven days in DD are plotted for male and female
control (w^{1118}) and experimental (Pdf^{01}) flies. **(G)** Average sleep plots under LD 12:12 for the
control (w^{1118}) and experimental (Pdf^{01}) groups are plotted for males (left) and females (right).
The plots are averaged over flies and days for a period of three days under LD 12:12. **(H)** Total
sleep values under LD conditions are plotted for male (left) and female (right) control (w^{1118}) and
experimental (Pdf^{01}) flies. **(I)** The differences in total LD sleep values between experimental
males and females and their respective controls are plotted. **(J)** Percentages of rhythmic flies are
plotted for the control (w^{1118}) and han^{5304} males (left, n = 86 (w^{1118}), n = 115 (han^{5304})) and
females (right, n = 74 (w^{1118}), n = 94 (han^{5304})) **(K).** Rhythmic power of the control (w^{1118}) and

han^5304 males and females calculated via the chi-square periodogram are plotted. **(L)** The
differences in rhythmic power between experimental males and females and their respective
controls are plotted. **(M)** Average sleep plots under LD 12:12 for control (w^{1118}) and
experimental (*han^5304*) flies are plotted for males (left) and females (right). The plots are
averaged over flies and days for a period of three days under LD 12:12. **(N)** Total sleep values
under LD conditions are plotted for male (left) and female (right) control (w^{1118}) and
experimental (*han^5304*) flies. **(O)** The differences in total LD sleep values between experimental
males and females and their respective controls are plotted. Statistical comparisons were
performed between the control and experimental flies of both sexes using a Mann☐Whitney U
test. The box plots extend from the 25^{th} to 75^{th} percentile, with whiskers extending from the
smallest to the largest value, and each point represents data from a single fly. Combined data
from at least three replicate experiments are plotted. * p < 0.05, ** p < 0.01, *** p < 0.001.

Figure 2: Tissue-specific CRISPR-mediated *Pdf* manipulation leads to a reduction in PDF
levels and the misrouting of s-LN_v dorsal termini in both sexes. (A) Representative confocal
images of control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^ts*) (top) and experimental (*Pdf-RFP, Pdf-Gal4;*
tub-Gal80^ts > Cas9; Pdf-g) (middle, males; bottom, females) flies stained with RFP and PDF
antibodies. Experimental flies show a significant reduction in PDF levels in the sLN_vs (white
arrows, PDF channel) and misrouting of the s-LN_v dorsal termini (white arrows, RFP channel).
(B) Representative confocal images of the small and large LN_vs of control (*Pdf-RFP, Pdf-Gal4;*
tub-Gal80^ts) (top) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^ts > Cas9; Pdf-g*) (bottom)
brains stained with RFP and PDF antibodies. **(C)** Quantification of PDF levels from s-LN_v cell
bodies in control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^ts*) and experimental (*Pdf-RFP, Pdf-Gal4; tub-*
Gal80^ts > Cas9; Pdf-g) flies are plotted for males (left) and females (right). *n* > 12 brains for all
genotypes. **(C)** Differences in the PDF intensity values of experimental males and females from
their respective parental controls. Statistical comparisons were performed between the control
and experimental flies of both sexes using the Mann☐Whitney U test. The box plots extend from
the 25^{th} to 75^{th} percentile, with whiskers extending from the smallest to the largest value, and
each point represents data from a single fly. Combined data from at least two replicate
experiments are plotted. * p < 0.05, ** p < 0.01, *** p < 0.001. Scale bars = 50 μm.

Figure 3: CRISPR-Cas9-mediated *Pdf* mutagenesis has more pronounced effects on male behavior. (A) Representative actograms of control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*) and *(UAS Cas9; Pdf-g)* and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) males (left) and females (right) are plotted for six days of LD followed by nine days of DD. **(B)** Percentages of rhythmic flies are plotted for control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*) and (*UAS Cas9; Pdf-g*) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) males (left, $n = 85$ (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*), $n = 89$ (*UAS Cas9; Pdf-g*), $n = 106$ (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) and females (right, $n = 79$ (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*), $n = 78$ (*UAS Cas9; Pdf-g*), $n = 85$ (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*). The error bars represent SEM values plotted across three replicate experiments. **(C)** Free-running periods of control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g*) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) males (left) and females (right) calculated via the chi-square periodogram are plotted. **(D)** Rhythmic power of control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g*) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) males (left) and females (right) calculated via the chi-square periodogram are plotted. **(E)** The differences in rhythmic power of experimental males and females from their respective controls are plotted. Flies were kept at 28°C throughout development, and as adults, experiments were conducted at 28°C. Statistical comparisons were performed between the control and experimental flies of both sexes using the Kruskal☐Wallis test followed by Dunn's multiple comparisons test for panels 3B-3D and Mann-Whitney U test for Fig. 3E. The box plots extend from the 25^{th} to 75^{th} percentile, with whiskers extending from the smallest to the largest value, and each point represents data from a single fly. Combined data from at least three replicate experiments are plotted. ** $p < 0.01$, *** $p < 0.001$.

Figure 4: Speeding up the clock in M-cells leads to sexually dimorphic phenotypes. (A) Depiction of the adult *Drosophila* brain hemisphere indicating the clock cell subsets (colored) having a faster running molecular clock. **(B)** Average activity plots of control (*Pdf-Gal4*) and (UAS-*DBT^s*) and experimental (*Pdf > DBT^s*) flies are plotted for males (left) and females (right). The plots are averaged over flies and days for a period of three days under LD 12:12. **(C)** The

phase of the morning peak of activity under LD 12:12 is plotted for control (*Pdf-Gal4*), (UAS-

DBT^s) and experimental (*Pdf >DBT^s*) males (left, *n* = 116 (*Pdf-Gal4*), *n* = 120 (UAS-*DBT^s*), *n* =

88 (*Pdf > DBT^s*)) and females (right, *n* = 85 (*Pdf-Gal4*), *n* = 109 (UAS-*DBT^s*), *n* = 73 (*Pdf >*

DBT^s)). **(D)** Percentages of rhythmic flies are plotted for controls (*Pdf-Gal4), (UAS-DBT^s),* and

experimental (*Pdf > DBT^s)* males (left, *n* = 90 (*Pdf-Gal4*), *n* = 108 (*UAS-DBT^s*), *n* = 93 (*Pdf >*

DBT^s) and females (right, *n* = 66 (*Pdf-Gal4*), *n* = 83 (*UAS-DBT^s*), *n* = 70 (*Pdf > DBT^s).* The error

bars represent the SEM values plotted across three replicate experiments. **(E)** Representative

actograms of control (UAS-*DBT^s*) and experimental (*Pdf > DBT^s*) males (left) and females

(right) plotted for six days of LD followed by ten days of DD. **(F)** Free-running periods of

control (*Pdf-Gal4), (UAS-DBT^s)* and experimental (*Pdf > DBT^s)* males (left) and females (right)

calculated via the chi-square periodogram are plotted. **(G)** Rhythmic power of control (*Pdf-*

Gal4), (UAS-DBT^s) and experimental (*Pdf > DBT^s)* males (left) and females (right) calculated via

the chi-square periodogram are plotted. **(H)** The differences in rhythmic power between

experimental males and females and their respective controls are plotted. Statistical comparisons

were performed between the control and experimental flies of both sexes using the

Kruskal☐Wallis test followed by Dunn's multiple comparisons test for panels 4C, 4F, and 4G

and Mann-Whitney U test for Fig. 4H. The box plots extend from the 25^{th} to 75^{th} percentile, with

whiskers extending from the smallest to the largest value, and each point represents data from a

single fly. Combined data from at least three replicate experiments are plotted. * $p < 0.05$, ** $p <$

0.01, *** $p < 0.001$.

Figure 5: Speeding up the clock in an E-cell subset leads to a more advanced evening peak

in females. (A) Depiction of the adult *Drosophila* brain hemisphere indicating the clock cell

subsets (colored) having a faster running molecular clock. **(B)** Average activity plots of control

(*MB122B-Gal4*) and (*UAS-DBT^s*) and experimental (*MB122B > DBT^s*) flies are plotted for males

(left) and females (right). The plots are averaged over flies and days for a period of three days

under LD 12:12. **(C)** Phase of the evening peak of activity under LD 12:12 for controls

(*MB122B-Gal4*) and (*UAS-DBT^s*) and experimental (*MB122B > DBT^s*) males (left, *n* = 92

(*MB122B-Gal4*), *n* = 93 (*UAS-DBT^s*), *n* = 93 (*MB122B > DBT^s*)) and females (right, *n* = 85

(*MB122B-Gal4*), *n* = 87 (*UAS-DBT^s*), *n* = 89 (*MB122B > DBT^s*)). **(D)** The differences in the

phase of the E peak between experimental males and females and their respective controls are

plotted. **(E)** Free-running period of control (*MB122B-Gal4* and *UAS-DBT^s)* and experimental

(*MB122B > DBT^s)* males (left) and females (right) calculated via the chi-square periodogram are

plotted. **(F)** Rhythmic power of control (*MB122B-Gal4 and UAS-DBT^s)* and experimental

(*MB122B > DBT^s)* males (left) and females (right) calculated via the chi-square periodogram are

plotted. Statistical comparisons were performed between the control and experimental flies of

both sexes using the Kruskal□Wallis test followed by Dunn's multiple comparisons test for

panels 5C, 5E, and 5F and Mann-Whitney U test for Fig. 5D. The box plots extend from the 25th

to 75th percentile, with whiskers extending from the smallest to the largest value, and each point

represents data from a single fly. Combined data from at least three replicate experiments are

plotted. ** p < 0.01, *** p < 0.001.

Supplementary figures

Figure S1: PDF rhythmic accumulation is similar in males and females (related to Figure

1.)

(A) Representative confocal images of control (*w^{1118}*) (top) and experimental (*Pdf^{01}*) (bottom)

male and female flies stained with the PDF antibody. **(B)** Scatterplots of PDF staining intensities

of the s-LN$_v$ dorsal projections of both male and female flies plotted at different time points over

a 24-hr cycle on DD day 3. Each dot represents the mean PDF intensity value averaged over both

hemispheres of one brain. The cyan and pink lines are the best fit cosine curves from the

parameters that were extracted via COSINOR analysis. See Table 2 for more details. **(C)**

Amplitude values of PDF oscillation obtained from COSINOR curve fits are plotted for male and

female flies. The error bars are 95% CI values calculated from the standard error obtained from

COSINOR analysis. The overlapping error bars indicate that the amplitude values of males and

females are not significantly different. n > 7 brain samples/time point.

Figure S2: CRISPR-Cas9-mediated *Pdf* mutagenesis has similar effects on male and female

sleep. (Related to Figure 3.) (A) Average sleep plots under LD 12:12 of control (*Pdf-RFP, Pdf-*

Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} >*

Cas9; Pdf-g) males (left) and females (right) are plotted. The plots are averaged over flies and

days for a period of 3 days under LD 12:12. **(B)** Total sleep values under LD conditions are plotted for control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g)* and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g)* males (left) and females (right). **(C)** Average sleep plots of flies over seven days in DD are plotted for male and female control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g)* and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) flies. **(D)** Total sleep values under DD conditions are plotted for control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g)* and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g)* males (left) and females (right). **(E)** The differences in the total DD sleep values of experimental males and females from their respective controls are plotted. Statistical comparisons were performed between the control and experimental flies of both sexes using the Kruskal☐Wallis test followed by Dunn's multiple comparisons test for panels S2B and S2D and Mann-Whitney U test for Panel S2E. The box plots extend from the 25th to 75th percentile, with whiskers extending from the smallest to the largest value, and each point represents data from a single fly. Combined data from at least three replicate experiments are plotted. ** $p < 0.01$, *** $p < 0.001$.

Figure S3: CRISPR-Cas9-mediated *Pdf* mutagenesis in small ventral lateral neurons has more pronounced effects on male circadian behavior. (A) Percentages of rhythmic flies are plotted for control (*s-LNv-Gal4) and (UAS Cas9; Pdf--g)* and experimental (*s-LNv > Cas9; Pdf--g)* males (left, $n = 16$ for all genotypes) and females (right, $n = 16$ for all genotypes). **(B)** Free-running period of control (*s-LNv-Gal4 and UAS Cas9; Pdf-g)* and experimental (*s-LNv > Cas9; Pdf-g)* males (left) and females (right) calculated via the chi-square periodogram are plotted. **(C)** Rhythmic power of control (*s-LNv-Gal4 and UAS Cas9; Pdf-g)* and experimental (*s-LNv > Cas9; Pdf-g)* males (left) and females (right) calculated via the chi-square periodogram are plotted. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Statistical comparisons were performed between the control and experimental flies of both sexes using the Kruskal☐Wallis test followed by Dunn's multiple comparisons test. The box plots extend from the 25th to 75th percentile, with whiskers extending from the smallest to the largest value, and each point represents data from a single fly. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure S4: Adult-specific CRISPR☐Cas9-mediated *Pdf* mutagenesis is more effective in males. (Related to Figure 3.) (A) Representative confocal images of control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*) (top) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) male (middle) and female (bottom) flies stained with RFP and PDF antibodies. The experimental flies presented a significant reduction in PDF levels but no misrouting of the s-LN$_v$ dorsal termini. **(B)** Quantification of PDF intensity from s-LN$_v$ cell bodies in control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) flies are plotted for males (left) and females (right). The data points represent the intensity from a single brain sample. *n* > 10 brains for all genotypes. **(C)** The differences in PDF intensity values of experimental males and females from their respective controls are plotted. **(D)** Percentages of rhythmic flies are plotted for control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*) and (*UAS Cas9; Pdf-g*) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) males (left, *n* = 103 (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*), *n* = 109 (*UAS Cas9; Pdf-g*), *n* = 115 (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) and females (right, *n* = 96 (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*), *n* = 90 (*UAS Cas9; Pdf-g*), *n* = 119 (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*). **(E)** Free-running periods of control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g*) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) males (left) and females (right) calculated via the chi-square periodogram are plotted. **(F)** Rhythmic power of control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g*) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) males (left) and females (right) calculated via the chi-square periodogram are plotted. Flies were raised at 18°C until eclosion and were then transferred to 28°C as adults. The flies were kept at 28°C for a period of four to five days before their activity-rest behavior was recorded at 28°C. Statistical comparisons were performed between the control and experimental flies of both sexes using the Kruskal☐Wallis test followed by Dunn's multiple comparisons test for panels S4E-F and Mann-Whitney U test for panels S4B-C. The box plots extend from the 25th to 75th percentile, with whiskers extending from the smallest to the largest value, and each point represents data from a single fly. Combined data from at least three replicate experiments are plotted. Scale bars represent 50 μm. * p < 0.05, ** p < 0.01, *** p < 0.001.

Figure S5: Speeding up the clock in all clock cells leads to a phase advance and shortening of the free-running period of activity rhythms (related to Figures 4 and 5). (A) Diagram of

an adult *Drosophila* brain hemisphere indicating the clock cell subsets (blue) expressing $DBT^{s.}$ **(B)** Average activity plots of control (*Clk856-Gal4*) and (UAS-*DBT^s*) and experimental (*Clk856 > DBT^s)* flies are plotted for males (left) and females (right). The plots are averaged over flies and days for a period of three days under LD 12:12. **(C)** Phases of the morning peak of activity under LD 12:12 for controls (*Clk856-Gal4*) and (*UAS-DBT^s*) and experimental (*Clk856 > DBT^s*) males (left, $n = 62$ (*Clk856-Gal4*), $n = 57$ (*UAS--DBT^s*), $n = 60$ (*Clk856 > DBT^s*)) and females (right, $n = 59$ (*Clk856-Gal4*), $n = 59$ (*UAS--DBT^s*), $n = 50$ (*Clk856 > DBT^s*)) are plotted. **(D)** Phase of the evening peak of activity under LD 12:12 for controls (*Clk856--Gal4*) and (*UAS-DBT^s*) and experimental (*Clk856 > DBT^s*) males (left) and females (right) are plotted. **(E)** Percentages of rhythmic flies are plotted for control (*Clk856--Gal4*) and (*UAS-DBT^s*) and experimental (*Clk856 > DBT^s*) males (left) and females (right). The error bars represent the SEM values plotted across two replicate experiments. **(F)** Free-running periods of rhythmic flies calculated via the chi-square period are plotted for controls (*Clk856--Gal4*) and (*UAS-DBT^s*) and experimental (*Clk856 > DBT^s*) males (left) and females (right). **(G)** Rhythmic power of flies calculated via the chi-square periodogram is plotted for controls (*Clk856--Gal4*) and (*UAS-DBT^s*) and experimental (*Clk856 > DBT^s*) males (left) and females (right). Statistical comparisons were performed between the control and experimental flies of both sexes via Kruskal Wallis test followed by Dunn's multiple comparisons test. The box plots extend from the 25th to 75th percentile, with whiskers extending from the smallest to the largest value, and each point represents data from a single fly. Combined data from at least two replicate experiments are plotted. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure S6: Blocking neurotransmission in *Pdf*-expressing cells leads to more pronounced lengthening of the free-running period in males. (Related to Figure 6). (A) Representative actograms of control *(Pdf-Gal4)*, and (*UAS TeTxLC*) and experimental (*Pdf > TeTxLC*) males (left) and females (right) are plotted for five days of LD followed by eight days of DD. **(B)** Percentage of rhythmic flies are plotted for control (*Pdf-Gal4*), and (*UAS TeTxLC*), and experimental (*Pdf > TeTxLC*) males (left, n = 56 (*Pdf-Gal4*), n = 51 (UAS *TeTxLC*), n = 52 (*Pdf > TeTxLC*) and females (right, n = 53 (*Pdf-Gal4*), n = 52 (*UAS TeTxLC*), n = 52 (*Pdf > TeTxLC*). Error bars are SEM values plotted across two replicate experiments. **(C)** Free-running period of control (*Pdf-Gal4* and UAS *TeTxLC*) and experimental (*Pdf > TeTxLC*) males (left) and females

(right) calculated using the Chi-square Periodogram are plotted. **(D)** The difference in free-
running period of experimental males and females from their respective controls are plotted. **(E)**
Rhythmic power of control (*Pdf-Gal4* and *UAS TeTxLC*) and experimental (*Pdf > TeTxLC*)
males (left) and females (right) calculated using the Chi-square periodogram are plotted.
Statistical comparisons were performed between the control and experimental flies for both sexes
using the Kruskal-Wallis test followed by the Dunn's multiple comparisons test for all panels
except S6D where comparisons were made using the Mann Whitney U test. The box plots extend
from the 25^{th} to 75^{th} percentile, with whiskers extending from the smallest to the largest value,
and each point representing data from a single fly. Combined data from at least two replicate
experiments are plotted.

Table 1: Table representing the % rhythmicity, phase of the E peak, free-running period and
rhythmic power of w^{1118} and Pdf^{01} males and females.

Pdf^{01}				
Genotype	% Rhythmicity ± SEM	E-peak phase ± SEM	Free-running period ± SEM	Rhythmic power ± SEM
w^{1118} (male)	97.3 ± 2.66	11.92 ± 0.07	23.8 ± 0.04	97.8 ± 3.88
Pdf^{01} (male)	30.3 ± 5.05	11.72 ± 0.07	23.72 ± 0.13	22.1 ± 3.24
w^{1118} (female)	92.8 ± 3.27	11.71 ± 0.07	24.1 ± 0.04	109.9 ± 5.63
Pdf^{01} (female)	56.9 ± 3.63	11.85 ± 0.08	23.9 ± 0.09	49.63 ± 4

720
721
722
723 **Table 2:** Cosinor analysis parameters for *Canton-S* males and females.
724
725

Cosinor parameters for PDF cycling in s-LNv dorsal termini					
Genotype	p value	PR	Mesor ± s.e.	Amplitude ± s.e.	Phase ± s.e.
Canton-S (male)	2.02e–06	42.1	17.15 ± 2.67	22.38 ± 3.79	-4.71 ± 9.64
Canton-S (female)	7.33e–07	37.554	14.59 ± 2.9	24.47 ± 4.07	-19.13 ± 9.65

726
727
728

729 **Table 3:** Table representing the % rhythmicity, phase of the E-peak, free-running period and
730 rhythmic power of w^{1118} and $PdfR^{01}$ males and females.

731
732

$PdfR^{01}$				
Genotype	% Rhythmicity ± SEM	E-peak phase ± SEM	Free-running period ± SEM	Rhythmic power ± SEM
w^{1118} (male)	97.3 ± 2.66	11.68 ± 0.07	23.8 ± 0.03	103.1 ± 4.15

PdfR^{01} (male)	16.4 ± 1.64	10.33 ± 0.04	21.8 ± 0.12	17.46 ± 2.28
w^{1118} (female)	92.7 ± 3.25	11.45 ± 0.07	24.1 ± 0.05	89.13 ± 4.75
PdfR^{01} (female)	65.4 ± 5.8	10.36 ± 0.07	22.4 ± 0.07	47.3 ± 3.5

Table 4: Table representing the % rhythmicity, % complex rhythms, free-running period and rhythmic power of *control and Pdf >* males and females.

Pdf > DBT^s			
Genotype	% Rhythmicity ± SEM	% Complex rhythms ± SEM	Free-running period ± SEM	Rhythmic power ± SEM
Pdf-Gal4 (male)	95.86 ± 2.73	0	24.3 ± 0.03	111.8 ± 5.58
UAS DBT^s (male)	96.79 ± 2.21	0	23.9 ± 0.01	134.8 ± 4.44
Pdf > DBT^s (male)	69.87 ± 6.89	23.73 ± 5.51	18.9 ± 0.14	49.48 ± 3.54
Pdf-Gal4 (female)	93.56 ± 4.02	0	24.4 ± 0.03	122 ± 6.76
UAS DBT^s (female)	89.45 ± 2.74	0	24.2 ± 0.03	89.7 ± 5.19
Pdf > DBT^s (female)	38.41 ± 2.74	5.54 ± 3.45	22.5 ± 0.58	24.8 ± 3.55

References

1. Anderson ST, FitzGerald GA. Sexual dimorphism in body clocks. Science. 2020;369(6508):1164-5. doi: doi:10.1126/science.abd4964.
2. Bailey M, Silver R. Sex differences in circadian timing systems: implications for disease. Front Neuroendocrinol. 2014;35(1):111-39. Epub 2013/11/30. doi: 10.1016/j.yfrne.2013.11.003. PubMed PMID: 24287074; PubMed Central PMCID: PMCPMC4041593.
3. Vida B, Hrabovszky E, Kalamatianos T, Coen CW, Liposits Z, Kalló I. Oestrogen Receptor α and β Immunoreactive Cells in the Suprachiasmatic Nucleus of Mice: Distribution, Sex Differences and Regulation by Gonadal Hormones. Journal of Neuroendocrinology. 2008;20(11):1270-7. doi: https://doi.org/10.1111/j.1365-2826.2008.01787.x.
4. Lyall LM, Wyse CA, Graham N, Ferguson A, Lyall DM, Cullen B, et al. Association of disrupted circadian rhythmicity with mood disorders, subjective wellbeing, and cognitive function: a cross-sectional study of 91 105 participants from the UK Biobank. Lancet Psychiatry. 2018;5(6):507-14. Epub 2018/05/20. doi: 10.1016/S2215-0366(18)30139-1. PubMed PMID: 29776774.
5. Bixler EO, Papaliaga MN, Vgontzas AN, Lin HM, Pejovic S, Karataraki M, et al. Women sleep objectively better than men and the sleep of young women is more resilient to external stressors: effects of age and menopause. J Sleep Res. 2009;18(2):221-8. Epub 2009/03/24. doi: 10.1111/j.1365-2869.2008.00713.x. PubMed PMID: 19302341; PubMed Central PMCID: PMCPMC3594776.
6. Hastings MH, Maywood ES, Brancaccio M. The Mammalian Circadian Timing System and the Suprachiasmatic Nucleus as Its Pacemaker. Biology (Basel). 2019;8(1). Epub 20190311. doi: 10.3390/biology8010013. PubMed PMID: 30862123; PubMed Central PMCID: PMCPMC6466121.
7. Helfrich-Forster C. The neuroarchitecture of the circadian clock in the brain of Drosophila melanogaster. Microsc Res Tech. 2003;62(2):94-102. Epub 2003/09/11. doi: 10.1002/jemt.10357. PubMed PMID: 12966496.
8. Schubert FK, Hagedorn N, Yoshii T, Helfrich-Forster C, Rieger D. Neuroanatomical details of the lateral neurons of Drosophila melanogaster support their functional role in the circadian system. J Comp Neurol. 2018;526(7):1209-31. Epub 2018/02/10. doi: 10.1002/cne.24406. PubMed PMID: 29424420; PubMed Central PMCID: PMCPMC5873451.
9. Yu W, Hardin PE. Circadian oscillators of Drosophila and mammals. J Cell Sci. 2006;119(Pt 23):4793-5. Epub 2006/11/30. doi: 10.1242/jcs.03174. PubMed PMID: 17130292.
10. Kloss B, Price JL, Saez L, Blau J, Rothenfluh A, Wesley CS, et al. The Drosophila clock gene double-time encodes a protein closely related to human casein kinase Iepsilon. Cell. 1998;94(1):97-107. Epub 1998/07/23. doi: 10.1016/s0092-8674(00)81225-8. PubMed PMID: 9674431.
11. Price JL, Blau J, Rothenfluh A, Abodeely M, Kloss B, Young MW. double-time is a novel Drosophila clock gene that regulates PERIOD protein accumulation. Cell. 1998;94(1):83-95. Epub 1998/07/23. doi: 10.1016/s0092-8674(00)81224-6. PubMed PMID: 9674430.
12. Helfrich-Forster C. Organization of endogenous clocks in insects. Biochem Soc Trans. 2005;33(Pt 5):957-61. Epub 2005/10/26. doi: 10.1042/BST20050957. PubMed PMID: 16246020.

13. Ma D, Przybylski D, Abruzzi KC, Schlichting M, Li Q, Long X, et al. A transcriptomic
taxonomy of Drosophila circadian neurons around the clock. eLife. 2021;10:e63056. doi:
10.7554/eLife.63056.
14. Shafer OT, Helfrich-Forster C, Renn SC, Taghert PH. Reevaluation of Drosophila
melanogaster's neuronal circadian pacemakers reveals new neuronal classes. J Comp Neurol.
2006;498(2):180-93. Epub 2006/07/21. doi: 10.1002/cne.21021. PubMed PMID: 16856134;
PubMed Central PMCID: PMCPMC2596765.
15. Shafer OT, Gutierrez GJ, Li K, Mildenhall A, Spira D, Marty J, et al. Connectomic analysis of
the Drosophila lateral neuron clock cells reveals the synaptic basis of functional pacemaker
classes. Elife. 2022;11. Epub 2022/06/30. doi: 10.7554/eLife.79139. PubMed PMID: 35766361;
PubMed Central PMCID: PMCPMC9365390.
16. Grima B, Chelot E, Xia R, Rouyer F. Morning and evening peaks of activity rely on different
clock neurons of the Drosophila brain. Nature. 2004;431(7010):869-73. Epub 2004/10/16. doi:
10.1038/nature02935. PubMed PMID: 15483616.
17. Rieger D, Shafer OT, Tomioka K, Helfrich-Forster C. Functional analysis of circadian
pacemaker neurons in Drosophila melanogaster. J Neurosci. 2006;26(9):2531-43. Epub
2006/03/03. doi: 10.1523/JNEUROSCI.1234-05.2006. PubMed PMID: 16510731.
18. Renn SC, Park JH, Rosbash M, Hall JC, Taghert PH. A pdf neuropeptide gene mutation and
ablation of PDF neurons each cause severe abnormalities of behavioral circadian rhythms in
Drosophila. Cell. 1999;99(7):791-802. Epub 2000/01/05. PubMed PMID: 10619432.
19. Park JH, Helfrich-Forster C, Lee G, Liu L, Rosbash M, Hall JC. Differential regulation of
circadian pacemaker output by separate clock genes in Drosophila. Proc Natl Acad Sci U S A.
2000;97(7):3608-13. Epub 2000/03/22. doi: 10.1073/pnas.070036197. PubMed PMID:
10725392; PubMed Central PMCID: PMCPMC16287.
20. Stoleru D, Peng Y, Nawathean P, Rosbash M. A resetting signal between Drosophila
pacemakers synchronizes morning and evening activity. Nature. 2005;438(7065):238-42. Epub
2005/11/11. doi: 10.1038/nature04192. PubMed PMID: 16281038.
21. Stoleru D, Peng Y, Agosto J, Rosbash M. Coupled oscillators control morning and evening
locomotor behaviour of Drosophila. Nature. 2004;431(7010):862-8. Epub 2004/10/16. doi:
10.1038/nature02926. PubMed PMID: 15483615.
22. Fernandez MP, Pettibone HL, Bogart JT, Roell CJ, Davey CE, Pranevicius A, et al. Sites of
Circadian Clock Neuron Plasticity Mediate Sensory Integration and Entrainment. Curr Biol.
2020;30(12):2225-37 e5. Epub 2020/05/11. doi: 10.1016/j.cub.2020.04.025. PubMed PMID:
32386535; PubMed Central PMCID: PMCPMC7314648.
23. Guo F, Chen X, Rosbash M. Temporal calcium profiling of specific circadian neurons in
freely moving flies. Proc Natl Acad Sci U S A. 2017;114(41):E8780-E7. Epub 2017/10/05. doi:
10.1073/pnas.1706608114. PubMed PMID: 28973886; PubMed Central PMCID:
PMCPMC5642695.
24. Lamaze A, Stanewsky R. DN1p or the "Fluffy" Cerberus of Clock Outputs. Front Physiol.
2019;10:1540. Epub 2020/01/24. doi: 10.3389/fphys.2019.01540. PubMed PMID: 31969832;
PubMed Central PMCID: PMCPMC6960142.
25. Collins B, Kane Elizabeth A, Reeves David C, Akabas Myles H, Blau J. Balance of Activity
between LNvs and Glutamatergic Dorsal Clock Neurons Promotes Robust Circadian Rhythms in
Drosophila. Neuron. 2012;74(4):706-18. doi: https://doi.org/10.1016/j.neuron.2012.02.034.

26. Yoshii T, Wulbeck C, Sehadova H, Veleri S, Bichler D, Stanewsky R, et al. The neuropeptide
pigment-dispersing factor adjusts period and phase of Drosophila's clock. J Neurosci.
2009;29(8):2597-610. Epub 2009/02/27. doi: 10.1523/JNEUROSCI.5439-08.2009. PubMed
PMID: 19244536.
27. Peng Y, Stoleru D, Levine JD, Hall JC, Rosbash M. Drosophila free-running rhythms require
intercellular communication. PLoS Biol. 2003;1(1):E13. Epub 2003/09/17. doi:
10.1371/journal.pbio.0000013. PubMed PMID: 12975658; PubMed Central PMCID:
PMCPMC193604.
28. Liang X, Holy TE, Taghert PH. A Series of Suppressive Signals within the Drosophila
Circadian Neural Circuit Generates Sequential Daily Outputs. Neuron. 2017;94(6):1173-89 e4.
Epub 2017/05/30. doi: 10.1016/j.neuron.2017.05.007. PubMed PMID: 28552314; PubMed
Central PMCID: PMCPMC5502710.
29. Mertens I, Vandingenen A, Johnson EC, Shafer OT, Li W, Trigg JS, et al. PDF receptor
signaling in Drosophila contributes to both circadian and geotactic behaviors. Neuron.
2005;48(2):213-9. Epub 2005/10/26. doi: 10.1016/j.neuron.2005.09.009. PubMed PMID:
16242402.
30. Hyun S, Lee Y, Hong ST, Bang S, Paik D, Kang J, et al. Drosophila GPCR Han is a receptor for
the circadian clock neuropeptide PDF. Neuron. 2005;48(2):267-78. Epub 2005/10/26. doi:
10.1016/j.neuron.2005.08.025. PubMed PMID: 16242407.
31. Lear BC, Merrill CE, Lin JM, Schroeder A, Zhang L, Allada R. A G protein-coupled receptor,
groom-of-PDF, is required for PDF neuron action in circadian behavior. Neuron. 2005;48(2):221-
7. Epub 2005/10/26. doi: 10.1016/j.neuron.2005.09.008. PubMed PMID: 16242403.
32. Kim WJ, Jan LY, Jan YN. A PDF/NPF neuropeptide signaling circuitry of male Drosophila
melanogaster controls rival-induced prolonged mating. Neuron. 2013;80(5):1190-205. doi:
10.1016/j.neuron.2013.09.034. PubMed PMID: 24314729; PubMed Central PMCID:
PMCPMC3867312.
33. Krupp JJ, Billeter JC, Wong A, Choi C, Nitabach MN, Levine JD. Pigment-dispersing factor
modulates pheromone production in clock cells that influence mating in drosophila. Neuron.
2013;79(1):54-68. Epub 2013/07/16. doi: 10.1016/j.neuron.2013.05.019. PubMed PMID:
23849197; PubMed Central PMCID: PMCPMC3955580.
34. Inami S, Sato S, Kondo S, Tanimoto H, Kitamoto T, Sakai T. Environmental Light Is Required
for Maintenance of Long-Term Memory in Drosophila. J Neurosci. 2020;40(7):1427-39. Epub
2020/01/15. doi: 10.1523/JNEUROSCI.1282-19.2019. PubMed PMID: 31932417; PubMed
Central PMCID: PMCPMC7044726.
35. Toma DP, White KP, Hirsch J, Greenspan RJ. Identification of genes involved in Drosophila
melanogaster geotaxis, a complex behavioral trait. Nat Genet. 2002;31(4):349-53. Epub
2002/06/04. doi: 10.1038/ng893. PubMed PMID: 12042820.
36. Guo F, Yu J, Jung HJ, Abruzzi KC, Luo W, Griffith LC, et al. Circadian neuron feedback
controls the Drosophila sleep--activity profile. Nature. 2016;536(7616):292-7. Epub 2016/08/02.
doi: 10.1038/nature19097. PubMed PMID: 27479324; PubMed Central PMCID:
PMCPMC5247284.
37. Chen D, Sitaraman D, Chen N, Jin X, Han C, Chen J, et al. Genetic and neuronal
mechanisms governing the sex-specific interaction between sleep and sexual behaviors in

Drosophila. Nat Commun. 2017;8(1):154. Epub 2017/07/30. doi: 10.1038/s41467-017-00087-5.
PubMed PMID: 28754889; PubMed Central PMCID: PMCPMC5533705.
38.   Kunst M, Hughes ME, Raccuglia D, Felix M, Li M, Barnett G, et al. Calcitonin gene-related
peptide neurons mediate sleep-specific circadian output in Drosophila. Curr Biol.
2014;24(22):2652-64. Epub 2014/12/03. doi: 10.1016/j.cub.2014.09.077. PubMed PMID:
25455031; PubMed Central PMCID: PMCPMC4255360.
39.   Helfrich-Forster C. Differential control of morning and evening components in the activity
rhythm of Drosophila melanogaster--sex-specific differences suggest a different quality of
activity.   J   Biol   Rhythms.   2000;15(2):135-54.   Epub   2000/04/13.   doi:
10.1177/074873040001500208. PubMed PMID: 10762032.
40.   Palmateer CM, Artikis C, Brovero SG, Friedman B, Gresham A, Arbeitman MN. Single-cell
transcriptome profiles of Drosophila fruitless-expressing neurons from both sexes. eLife.
2023;12:e78511. doi: 10.7554/eLife.78511.
41.   Fujii S, Amrein H. Ventral lateral and DN1 clock neurons mediate distinct properties of
male sex drive rhythm in Drosophila. Proc Natl Acad Sci U S A. 2010;107(23):10590-5. Epub
2010/05/26. doi: 10.1073/pnas.0912457107. PubMed PMID: 20498055; PubMed Central
PMCID: PMCPMC2890825.
42.   Hanafusa S, Kawaguchi T, Umezaki Y, Tomioka K, Yoshii T. Sexual interactions influence the
molecular oscillations in DN1 pacemaker neurons in Drosophila melanogaster. PLoS One.
2013;8(12):e84495. Epub 2013/12/25. doi: 10.1371/journal.pone.0084495. PubMed PMID:
24367668; PubMed Central PMCID: PMCPMC3867508.
43.   Lee G, Bahn JH, Park JH. Sex- and clock-controlled expression of the neuropeptide F gene
in Drosophila. Proc Natl Acad Sci U S A. 2006;103(33):12580-5. Epub 20060807. doi:
10.1073/pnas.0601171103.   PubMed   PMID: 16894172;   PubMed   Central   PMCID:
PMCPMC1567921.
44.   Hermann C, Yoshii T, Dusik V, Helfrich-Forster C. Neuropeptide F immunoreactive clock
neurons modify evening locomotor activity and free-running period in Drosophila melanogaster.
J Comp Neurol. 2012;520(5):970-87. Epub 2011/08/10. doi: 10.1002/cne.22742. PubMed PMID:
21826659.
45.   Nässel DR, Zandawala M. Recent advances in neuropeptide signaling in Drosophila, from
genes to physiology and behavior. Prog Neurobiol. 2019;179:101607. Epub 2019/03/25. doi:
10.1016/j.pneurobio.2019.02.003. PubMed PMID: 30905728.
46.   Kuwano R, Katsura M, Iwata M, Yokosako T, Yoshii T. Pigment-dispersing factor and
CCHamide1 in the Drosophila circadian clock network. Chronobiol Int. 2023;40(3):284-99. Epub
2023/02/15. doi: 10.1080/07420528.2023.2166416. PubMed PMID: 36786215.
47.   Parisky KM, Agosto J, Pulver SR, Shang Y, Kuklin E, Hodge JJ, et al. PDF cells are a GABA-
responsive wake-promoting component of the Drosophila sleep circuit. Neuron.
2008;60(4):672-82. Epub 2008/11/29. doi: 10.1016/j.neuron.2008.10.042. PubMed PMID:
19038223; PubMed Central PMCID: PMCPMC2734413.
48.   Cornelissen G. Cosinor-based rhythmometry. Theoretical Biology and Medical Modelling.
2014;11(1):16. doi: 10.1186/1742-4682-11-16.
49.   Im SH, Li W, Taghert PH. PDFR and CRY signaling converge in a subset of clock neurons to
modulate the amplitude and phase of circadian behavior in Drosophila. PLoS One.

2011;6(4):e18974. Epub 2011/05/12. doi: 10.1371/journal.pone.0018974. PubMed PMID:
21559487; PubMed Central PMCID: PMCPMC3084726.

50. Richhariya S, Shin D, Le JQ, Rosbash M. Dissecting neuron-specific functions of circadian
genes using modified cell-specific CRISPR approaches. Proc Natl Acad Sci U S A.
2023;120(29):e2303779120. Epub 2023/07/10. doi: 10.1073/pnas.2303779120. PubMed PMID:
37428902; PubMed Central PMCID: PMCPMC10629539.

51. Gorostiza EA, Ceriani MF. Retrograde Bone Morphogenetic Protein Signaling Shapes a Key
Circadian Pacemaker Circuit. The Journal of Neuroscience. 2013;33(2):687-96. doi:
10.1523/jneurosci.3448-12.2013.

52. Ruben M, Drapeau MD, Mizrak D, Blau J. A mechanism for circadian control of pacemaker
neuron excitability. J Biol Rhythms. 2012;27(5):353-64. doi: 10.1177/0748730412455918.
PubMed PMID: 23010658; PubMed Central PMCID: PMCPMC4019749.

53. Fleck SA, Biswas P, DeWitt ED, Knuteson RL, Eisman RC, Nemkov T, et al. Auxin Exposure
Disrupts Feeding Behavior and Fatty Acid Metabolism in Adult Drosophila. bioRxiv. 2023. Epub
2023/08/30. doi: 10.1101/2023.08.15.553385. PubMed PMID: 37645868; PubMed Central
PMCID: PMCPMC10462055.

54. McGuire SE, Mao Z, Davis RL. Spatiotemporal gene expression targeting with the TARGET
and gene-switch systems in Drosophila. Sci STKE. 2004;2004(220):pl6. Epub 2004/02/19. doi:
10.1126/stke.2202004pl6. PubMed PMID: 14970377.

55. Yao Z, Shafer OT. The Drosophila circadian clock is a variably coupled network of multiple
peptidergic units. Science. 2014;343(6178):1516-20. Epub 2014/03/29. doi:
10.1126/science.1251285. PubMed PMID: 24675961; PubMed Central PMCID:
PMCPMC4259399.

56. Muskus MJ, Preuss F, Fan J-Y, Bjes ES, Price JL. Drosophila DBT Lacking Protein Kinase
Activity Produces Long-Period and Arrhythmic Circadian Behavioral and Molecular Rhythms.
Molecular and Cellular Biology. 2007;27(23):8049-64. doi: 10.1128/mcb.00680-07.

57. Kaneko M, Park JH, Cheng Y, Hardin PE, Hall JC. Disruption of synaptic transmission or
clock-gene-product oscillations in circadian pacemaker cells of Drosophila cause abnormal
behavioral rhythms. J Neurobiol. 2000;43(3):207-33. Epub 2000/06/08. doi: 10.1002/(sici)1097-
4695(20000605)43:3<207::aid-neu1>3.0.co;2-0. PubMed PMID: 10842235.

58. Blanchardon E, Grima B, Klarsfeld A, Chelot E, Hardin PE, Preat T, et al. Defining the role of
Drosophila lateral neurons in the control of circadian rhythms in motor activity and eclosion by
targeted genetic ablation and PERIOD protein overexpression. Eur J Neurosci. 2001;13(5):871-
88. Epub 2001/03/27. doi: 10.1046/j.0953-816x.2000.01450.x. PubMed PMID: 11264660.

59. Umezaki Y, Yasuyama K, Nakagoshi H, Tomioka K. Blocking synaptic transmission with
tetanus toxin light chain reveals modes of neurotransmission in the PDF-positive circadian clock
neurons of Drosophila melanogaster. J Insect Physiol. 2011;57(9):1290-9. Epub 2011/06/29. doi:
10.1016/j.jinsphys.2011.06.004. PubMed PMID: 21708159.

60. Yao Z, Bennett AJ, Clem JL, Shafer OT. The Drosophila Clock Neuron Network Features
Diverse Coupling Modes and Requires Network-wide Coherence for Robust Circadian Rhythms.
Cell Rep. 2016;17(11):2873-81. Epub 2016/12/16. doi: 10.1016/j.celrep.2016.11.053. PubMed
PMID: 27974202; PubMed Central PMCID: PMCPMC5161247.

61. Brown MP, Verma S, Palmer I, Zuniga AG, Rosensweig C, Keles MF, et al. A subclass of
evening cells promotes the switch from arousal to sleep at dusk. bioRxiv. 2023. Epub

2023/09/11. doi: 10.1101/2023.08.28.555147. PubMed PMID: 37693540; PubMed Central PMCID: PMCPMC10491161.

62. Delventhal R, O'Connor RM, Pantalia MM, Ulgherait M, Kim HX, Basturk MK, et al. Dissection of central clock function in Drosophila through cell-specific CRISPR-mediated clock gene disruption. Elife. 2019;8. Epub 2019/10/16. doi: 10.7554/eLife.48308. PubMed PMID: 31613218; PubMed Central PMCID: PMCPMC6794090.

63. Schlichting M, Diaz MM, Xin J, Rosbash M. Neuron-specific knockouts indicate the importance of network communication to Drosophila rhythmicity. Elife. 2019;8. Epub 2019/10/16. doi: 10.7554/eLife.48301. PubMed PMID: 31613223; PubMed Central PMCID: PMCPMC6794074.

64. Bulthuis N, Spontak KR, Kleeman B, Cavanaugh DJ. Neuronal Activity in Non-LNv Clock Cells Is Required to Produce Free-Running Rest:Activity Rhythms in Drosophila. J Biol Rhythms. 2019;34(3):249-71. Epub 2019/04/18. doi: 10.1177/0748730419841468. PubMed PMID: 30994046; PubMed Central PMCID: PMCPMC7153773.

65. Shansky RM, Murphy AZ. Considering sex as a biological variable will require a global shift in science culture. Nature Neuroscience. 2021;24(4):457-64. doi: 10.1038/s41593-021-00806-8.

66. Kokras N, Dalla C. Sex differences in animal models of psychiatric disorders. British Journal of Pharmacology. 2014;171(20):4595-619. doi: https://doi.org/10.1111/bph.12710.

67. Beery AK, Zucker I. Sex bias in neuroscience and biomedical research. Neurosci Biobehav Rev. 2011;35(3):565-72. Epub 2010/07/14. doi: 10.1016/j.neubiorev.2010.07.002. PubMed PMID: 20620164; PubMed Central PMCID: PMCPMC3008499.

68. Joye DAM, Evans JA. Sex differences in daily timekeeping and circadian clock circuits. Semin Cell Dev Biol. 2022;126:45-55. Epub 20210514. doi: 10.1016/j.semcdb.2021.04.026. PubMed PMID: 33994299; PubMed Central PMCID: PMCPMC8589873.

69. Taghert PH, Nitabach MN. Peptide neuromodulation in invertebrate model systems. Neuron. 2012;76(1):82-97. Epub 2012/10/09. doi: 10.1016/j.neuron.2012.08.035. PubMed PMID: 23040808; PubMed Central PMCID: PMCPMC3466441.

70. Goda T, Umezaki Y, Alwattari F, Seo HW, Hamada FN. Neuropeptides PDF and DH31 hierarchically regulate free-running rhythmicity in Drosophila circadian locomotor activity. Scientific Reports. 2019;9(1):838. doi: 10.1038/s41598-018-37107-3.

71. Helfrich-Forster C. Robust circadian rhythmicity of Drosophila melanogaster requires the presence of lateral neurons: a brain-behavioral study of disconnected mutants. J Comp Physiol A. 1998;182(4):435-53. Epub 1998/04/08. PubMed PMID: 9530835.

72. Fernandez MP, Berni J, Ceriani MF. Circadian remodeling of neuronal circuits involved in rhythmic behavior. PLoS Biol. 2008;6(3):e69. Epub 2008/03/28. doi: 10.1371/journal.pbio.0060069. PubMed PMID: 18366255; PubMed Central PMCID: PMCPMC2270325.

73. Biondi G, McCormick G, Fernandez MP. The Drosophila Circadian Clock Gene Cycle Controls Development of Clock Neurons. bioRxiv. 2023:2023.10.29.564626. doi: 10.1101/2023.10.29.564626.

74. Sweeney ST, Broadie K, Keane J, Niemann H, O'Kane CJ. Targeted expression of tetanus toxin light chain in Drosophila specifically eliminates synaptic transmission and causes behavioral defects. Neuron. 1995;14(2):341-51. Epub 1995/02/01. doi: 10.1016/0896-6273(95)90290-2. PubMed PMID: 7857643.

75. Kaneko M, Helfrich-Forster C, Hall JC. Spatial and temporal expression of the period and
timeless genes in the developing nervous system of Drosophila: newly identified pacemaker
candidates and novel features of clock gene product cycling. J Neurosci. 1997;17(17):6745-60.
Epub 1997/09/01. PubMed PMID: 9254686.
76. Hofbauer B, Zandawala M, Reinhard N, Rieger D, Werner C, Evers JF, et al. The
neuropeptide pigment-dispersing factor signals independently of Bruchpilot-labelled active
zones in daily remodelled terminals of Drosophila clock neurons. European Journal of
Neuroscience. 2024;59(10):2665-85. doi: https://doi.org/10.1111/ejn.16294.
77. Johard HAD, Yoishii T, Dircksen H, Cusumano P, Rouyer F, Helfrich-Förster C, et al.
Peptidergic clock neurons in Drosophila: Ion transport peptide and short neuropeptide F in
subsets of dorsal and ventral lateral neurons. Journal of Comparative Neurology.
2009;516(1):59-73. doi: 10.1002/cne.22099.
78. Todd WD, Fenselau H, Wang JL, Zhang R, Machado NL, Venner A, et al. A hypothalamic
circuit for the circadian control of aggression. Nat Neurosci. 2018;21(5):717-24. Epub 20180409.
doi: 10.1038/s41593-018-0126-0. PubMed PMID: 29632359; PubMed Central PMCID:
PMCPMC5920747.
79. Fujii S, Krishnan P, Hardin P, Amrein H. Nocturnal male sex drive in Drosophila. Curr Biol.
2007;17(3):244-51. Epub 2007/02/06. doi: 10.1016/j.cub.2006.11.049. PubMed PMID:
17276917; PubMed Central PMCID: PMCPMC2239012.
80. Sakai T, Ishida N. Circadian rhythms of female mating activity governed by clock genes in
Drosophila. Proc Natl Acad Sci U S A. 2001;98(16):9221-5. Epub 2001/07/27. doi:
10.1073/pnas.151443298. PubMed PMID: 11470898; PubMed Central PMCID: PMCPMC55401.
81. T M, Hari Dass S, Sharma VK. Egg-laying rhythm in Drosophila melanogaster. J Genet.
2008;87(5):495-504. Epub 2009/01/17. doi: 10.1007/s12041-008-0072-9. PubMed PMID:
19147938.
82. Zhang C, Daubnerova I, Jang YH, Kondo S, Žitňan D, Kim YJ. The neuropeptide allatostatin C
from clock-associated DN1p neurons generates the circadian rhythm for oogenesis. Proc Natl
Acad Sci U S A. 2021;118(4). doi: 10.1073/pnas.2016878118. PubMed PMID: 33479181;
PubMed Central PMCID: PMCPMC7848730.
83. Persons JL, Abhilash L, Lopatkin AJ, Roelofs A, Bell EV, Fernandez MP, et al. PHASE: An
Open-Source Program for the Analysis of DrosophilaPhase, Activity, and Sleep Under
Entrainment. J Biol Rhythms. 2022;37(4):455-67. Epub 2022/06/22. doi:
10.1177/07487304221093114. PubMed PMID: 35727044; PubMed Central PMCID:
PMCPMC10362883.
84. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
source platform for biological-image analysis. Nature Methods. 2012;9:676. doi:
10.1038/nmeth.2019
https://www.nature.com/articles/nmeth.2019#supplementary-information.
85. Lee Gierke C, Cornelissen G. Chronomics analysis toolkit (CATkit). Biol Rhythm Res.
2016;47(2):163-81. doi: 10.1080/09291016.2015.1094965.
1045

Pdf >Cas9 + Pdf-g

MB122 > DBT^S

A
MB122> DBT^S

PDFR+ E-cells = ~18.5
Other clocks = ~24 hr

B
♂ MB122-Gal4
UAS-DBT^S
MB122>DBT^S

♀ MB122-Gal4
UAS-DBT^S
MB122 >DBT^S

Activity counts /30 min
Zeitgeber Time (ZT)

C
♂ MB122-Gal4
UAS-DBT^S
MB122>DBT^S

♀ MB122Gal4
UAS-DBT^S
MB122 >DBT^S

E-peak (hrs.)

D
Δ (E-phase)
***
♂ ♀

E
♂
Free-running period
MB122-Gal4
UAS-DBT^S

♀
Free-running period
MB122-Gal4
UAS-DBT^S

F
♂
Rhythmic power
MB122-Gal4
UAS-DBT^S

♀
Rhythmic power
MB122-Gal4
UAS-DBT^S