The Circadian Neuropeptide PDF has Sexually Dimorphic Effects on Activity Rhythms

Aishwarya R. Iyer[1], Eva Scholz-Carlson[1], Evardra Bell[1], Grace Biondi[1], Shlesha Richhariya[2], and Maria P. Fernandez[1,3]

1- Department of Neuroscience and Behavior, Barnard College. New York, NY 10027
2- Department of Biology, Brandeis University. Waltham, MA 02454
3- Corresponding author: mfernand@barnard.edu

Abstract

The circadian system regulates the timing of multiple molecular, physiological, metabolic, and behavioral phenomena. In *Drosophila* as in other species, most of the research on how the timekeeping system in the brain controls timing of behavioral outputs has been conducted in males, or sex was not included as a biological variable. The main circadian pacemaker neurons in *Drosophila* release the neuropeptide Pigment Dispersing Factor (PDF), which functions as a key synchronizing factor in the network with complex effects on other clock neurons. Lack of *Pdf* or its receptor, *PdfR,* results in most flies displaying arrhythmicity in activity-rest cycles under constant conditions. However, our results show that female circadian rhythms are less affected by mutations in both *Pdf* and *PdfR*. Crispr-Cas9 mutagenesis of *Pdf* specifically in the ventral lateral neurons (LN$_v$s) also has a greater effect on male rhythms. We tested the influence of the M-cells over the circadian network and show that speeding up the molecular clock specifically in the M-cells leads to sexually dimorphic phenotypes, with a more pronounced effect on male rhythmic behavior. Our results suggest that the female circadian system is more resilient to manipulations of the PDF pathway and that circadian timekeeping is more distributed across the clock neuron network in females.

Keywords: circadian rhythms, sex differences, PDF, neuropeptides, *Drosophila,* sleep/wake cycles

Introduction

The differences in neuronal circadian timekeeping between sexes remain relatively unexplored, despite the expanding body of research highlighting the influence of sex on the mechanisms underlying neuronal control of behavior [1]. In mammals, steroid hormones display daily, clock-driven changes in abundance, and while these sex hormones are not required to maintain rhythms, they influence the amplitude and response to photic stimuli [1, 2]. Furthermore, structural and functional sex differences have been observed in brain areas that receive direct input from the brain's circadian timekeeping center [2, 3]. Research in humans has also revealed significant sexual dimorphism: men tend to have lower-amplitude endogenous rhythms than women [4], and they are less resilient to nocturnal sleep disruptions and spend less time asleep [5].

In mammals, the main circadian pacemaker resides in the suprachiasmatic nuclei (SCN), which in mice consist of a network of ~20,000 neurons (reviewed by [6]). The *Drosophila* circadian clock neuron network involves ~150 neurons and is the functional equivalent of the mammalian SCN [7, 8]. Each circadian clock neuron contains an intracellular molecular timekeeping mechanism based on a transcriptional-translational feedback loop: the genes *Clock* (*Clk*) and *cycle* (*cyc*) promote rhythmic transcription of several key genes, including *Period* (*Per*) *and Timeless* (*Tim*), which inhibit their own transcription [9]. Multiple kinases that act on components of these clock proteins and can affect the pace of the molecular clock have been identified. One such kinase is *Doubletime* (DBT), which binds to and phosphorylates PER, hindering its nuclear accumulation [10, 11].

The fly's clock network consists of lateral neurons (LNs), which include ventroateral (LN$_v$), dorsolateral (LN$_d$), and lateral posterior neurons (LPNs), as well as three groups of dorsal neurons (DN1, DN2, and DN3), some of which can be further subdivided [8, 12-16]. The ventral and dorsal LNs are sufficient to produce the normal endogenous bimodal rhythm of sleep and activity [17, 18]. The four small LN$_v$s (s-LN$_v$s) are usually referred to as morning cells (M-cells) since they control the morning peak of activity under light-dark cycles (LD). These cells are also essential to maintain rhythmicity under free-running conditions [19, 20]. The evening peak is controlled by the LN$_d$s and a *Pdf*-negative LN$_v$, the 5th LN$_v$ (E-cells) [17, 21, 22]. Some DNs also contribute to timing and amount of sleep via modulation of the M and E cells [23-27].

Release of the circadian neuropeptide Pigment Dispersing Factor (PDF) by the s-LN$_v$s is essential for endogenous circadian timekeeping. A *Pdf* null mutation, *Pdf^{01}*, results in a substantial fraction of arrhythmic flies [19], desynchronization of molecular oscillations [28, 29], and phase changes in electrical activity of some clock clusters, most notably the LNds [30]. Loss of *PdfR* also leads to loss of behavioral rhythms [31-33]. Interestingly, PDF and PDFR also regulate behaviors that are sex-specific or sexually dimorphic. Rival induced long mating duration requires PDF expression in the s-LN$_v$s, PDFR expression in a subset of LN$_d$s, and NPF expression in the LN$_d$s [34]. PDF controls rhythms in the sexually dimorphic pheromone profiles produced by the oenocytes [35], and is involved in long term mating suppression in males [36]. Both PDF and PDFR contribute to geotactic behaviors [37] and the phenotypes of *Pdf^{01}* mutants are sexually dimorphic, with males showing a more extreme negative geotaxis phenotype [38].

Sexual dimorphism in *Drosophila* sleep/wake cycles has been studied mostly under LD cycles. Males show lower activity and more sleep during the light phase [39-42]. This higher midday activity is due to the activity of a subset of sleep-promoting DN1s, which is more active in males [39] and receives input from the male-specific P1 neurons that control male courtship [40]. Unlike studies on circadian rhythms, *Drosophila* sleep research often involves only females. Males also have an earlier and more pronounced morning peak and a larger phase angle between the morning and the evening peak [43]. Under conditions of constant darkness and temperature (DD), males of several wild type strains have a small but significant reduction in free-running period relative to females of the same strain [43]. Moreover, males are more likely to retain a bimodal activity pattern in DD [43]. A recent transcriptomic analysis of *fruitless* (*fru*) expressing neurons revealed clusters that are enriched in circadian clock genes [44]. Interestingly, only the male dataset had some *fru+* clusters with multiple genes having known circadian functions. A previous study reported that the DN1s express the male-specific Fru^M protein [45], and that the number of cells in the DN1$_p$s cluster is sexually dimorphic in cell number [46]. In addition, the E3 subset of LN$_d$s has been shown to be dimorphic in expression of the neuropeptide NPF [47, 48].

Given the sexually dimorphic roles of neuropeptides, including PDF, in other behaviors [49], we asked if females were similarly affected by manipulations of the *Pdf/PdfR* pathway. We found that

female circadian rhythms are less affected by null mutations in both *Pdf* and *PdfR*, and that similar

effects are observed via CRISPR-Cas9 *Pdf* mutagenesis specifically in the LN_vs. Moreover,

speeding up the molecular clock in the LN_vs via expression of *DBT^s* leads to an advance of the

morning peak in males but not in females, and the pace of the free-running period of activity is

significantly shortened only in males. Taken together, our results show that the female circadian

system is more resilient to manipulations in the PDF pathway and suggest that the *Pdf+* neurons

play a more dominant role over the male than over the female circadian network.

Results

Mutations in PDF and PDFR lead to sexually dimorphic phenotypes.

A null mutation in *Pdf* results in molecular, physiological, and behavioral phenotypes in males

[50]. We assayed the activity rhythms of *Pdf^01*females under free-running conditions (DD) and

found that a remarkably large proportion of the experimental females were still rhythmic (Fig. 1

A-C). We calculated the difference in rhythmic power of the experimental flies relative to their

respective controls and found that the effect is less pronounced in females than in males (Fig. 1 D-

E), suggesting that experimental females have more consolidated rhythms. Mutant females that

were still rhythmic had a slightly but significantly shorter free-running period than did the controls

(Table 1). This phenotype, previously described in *Pdf^01* males, was not observed in experimental

males, consistent with the findings of a recent study [51].

Pdf mutants have increased sleep, and this effect is mediated by PDF function on the LN_vs

themselves [52]. We found that both sexes show an increase in total sleep in LD, but the effect is

more pronounced in females (Fig. 1F-I). While the sleep increase in males was most prominent at

midday, females exhibited increased sleep throughout most of the light phase (Fig. 1G). To rule

out the presence of remnant PDF expression in *Pdf^01* females we stained brains of control and

experimental males and females with an anti-PDF antibody. We did not observe any traces of PDF

in experimental flies of either sex (Fig. S1A).

PDF accumulates rhythmically in the s-LN$_v$ dorsal termini in a time-of-day dependent manner both in LD and DD [19, 20], and can be released from the projections and the soma [53]. To examine if there are differences in the phase and amplitude of PDF cycling between the sexes in a wild type background, we dissected *Canton-S* males and females on the third day under DD at 6 timepoints over a 24-hour cycle. Using a COSINOR-based curve fitting method [54], we found that both males and females have a significant 24-hour rhythms in PDF cycling in their dorsal projections, with no significant sex differences in phase or amplitude (Fig. S1B-C, Table 2).

Next, we asked if the effects of a *Pdf* receptor mutation (*PdfR*) on activity and sleep were also sexually dimorphic. Expression of PDFR, a GPCR, can be detected in most of the clock neurons except for 3 LNds, half of DN1ps, and some DN3s [55], coinciding with *Cryptochrome* expression in clock neurons [55]. However, *PdfR* mRNA appears to be expressed in all clock neurons [56]. The *han^5304* mutant is a *PdfR* hypomorph and exhibits behavioral phenotypes under both LD 12:12 and DD [31-33]. Under DD, both *Han* males and females had a significant reduction in rhythmicity compared to the controls, but there was a greater proportion of rhythmic females (~65%) than males (~16%) (Fig. 1J). The free-running period of experimental flies was significantly shorter for both sexes (Table 3), as reported previously for males. Rhythmic power is significantly lower than controls for both *han^5304* males and females (Fig. 1K), but the effect was more pronounced in males, suggesting that females have more consolidated rhythms (Fig. 1L). Sleep plots for the first week under DD show that both males and females sleep more than controls (Fig. 1E). Similar to the effect of the *Pdf* mutation, *han^5304* flies showed significantly higher levels of LD sleep compared to controls, and the effect was again more pronounced in females (Fig. 1 M-O). Taken together, the results suggest that female circadian rhythms are less affected by the loss of both *Pdf* and *PdfR*, while at least under LD, the sleep phenotypes are more pronounced in females.

CRISPR-Cas9 mediated *Pdf* mutagenesis has more pronounced effects on male behavior.

In the *Pdf* null mutant background, other genetic factors could contribute to the sexual dimorphism observed in behavioral rhythms. We therefore employed a tissue-specific CRISPR-Cas9 mediated knockout of *Pdf* in both males and females, as described in a recent study that focused on males [57]. To assess the efficiency of the manipulation we stained for PDF in flies that

constitutively expressed *Pdf* gRNA and Cas9 in *Pdf+* neurons. This experiment was conducted at

28°C, as this temperature was more effective at mimicking the behavioral phenotypes of the *Pdf*[01]

mutant than was 25°C (described below), and to allow direct comparison with an adult-specific

manipulations.

PDF was significantly reduced in the s-LN$_v$s in both sexes (Fig. 2 A-B). However, in most brains

we noted faint staining in the dorsal projections of at least one s-LN$_v$ in at least one hemisphere.

The PDF signal within the large LN$_v$s was higher and could be detected in two or three l-LNv cell

bodies in both hemispheres in the majority of brains. In addition to behavioral phenotypes, the

Pdf[01] mutation leads to pronounced misrouting of the s-LN$_v$ projections in male flies [58]. We next

employed a transgene expressing a red fluorescent protein under a *Pdf* regulatory sequence [59]

and observed some s-LN$_v$ projections occasionally defasciculating from the main axonal bundle

in one or both hemispheres in *Pdf >Pdf*-g; *Cas9* flies of both sexes (Fig. 2A). The extent of the

reduction in PDF intensity of experimental flies relative to the controls was similar between the

sexes (Fig. 2C).

We then analyzed activity-rest rhythms in *Pdf >Pdf*-g; *Cas9* flies and found that experimental

males were mostly arrhythmic, whereas a greater proportion of experimental females remained

rhythmic (Fig. 3A-B). The free-running period of experimental males was not significantly

different from one of the controls, possibly because of the wide range of periods exhibited. Females

had significantly shorter free-running period compared to parental controls (Fig. 3C). The

rhythmic power was significantly lower in experimental flies of both sexes (Fig. 3D) but the effect

was less pronounced in females (Fig. 3E), indicating that females are able to better consolidate

activity rhythms. At 25°C, both males and females exhibited reduction in percent rhythmicity but

the majority of the flies were rhythmic (Fig S3A). Both males and females showed a shortening of

their free-running period (Fig. S3B). Similar to what we observed at 28°C, the rhythmic power of

the experimental flies of both the sexes was significantly reduced (Fig. S3C) but females were less

affected (Fig. S3D).

We next conducted an adult specific Crispr-Cas9 knockdown of *Pdf* (Fig S4). While a previous

study conducted in males did not find a correlation between the misrouting induced by the loss of

Pdf during development and the loss of behavioral rhythms in adults [60], it is possible that the

sex differences in Pdf^{01} and *Pdf >Pdf*-g; *Cas9* flies were due to developmental effects. The AGES

system was previously employed to rule of a developmental contribution to the *Pdf* mutagenesis

phenotypes [57], but a recent study showed that AUXIN exposure short- and long-term changes

in physiology and behavior [61]. Therefore, we employed a temperature-sensitive Gal80 variant

with ubiquitous expression to conditionally inhibit Gal4-mediated expression of the *Pdf* guide and

Cas9 [62]. This allows for the temporal control of UAS transgenes since Gal80^{ts} is active at lower

temperatures but inactive at higher temperatures. We raised flies at 18˚C and transferred them to

28˚C immediately after eclosion, so that *Pdf* mutagenesis only occurs in adult flies. Consistent

with what was reported for males by Gorostiza *et al*. [60] using RNA mediated interference (RNAi)

of *Pdf*, we did not observe misrouting in flies of either sex (Fig. S4A). While PDF levels were

significantly reduced in both males and females, the manipulation was more effective in males

(Fig S4A-B). Therefore, although the circadian behavioral phenotypes were again more

pronounced in males than in females (Fig S4C-E), this is likely due to higher remaining PDF

expression in experimental females.

Speeding up the M-cell clock leads to a coherent period shortening only in males

Next, we sought to determine if the influence of the *Pdf* releasing cells themselves was sexually

dimorphic. While PDF is released from both the large and small LN$_v$s, only the s-LN$_v$s (M-cells)

play key roles in regulating free-running rhythm properties [17]. In males, manipulations that

change the pace of the clock specifically in the LN$_v$s result in changes in the phase of the morning

peak of activity and in free running period [21, 63]. We expressed the *Doubletime* 'short' (*DBT*^{s})

allele [64] under the *Pdf-Gal4* driver (Fig. 4A) and analyzed the effects on behavior in both sexes.

Interestingly, we found that *Pdf > DBT*^{s} males but not females have an advanced phase of the

morning peak of activity (M-peak, Fig. 4B-C). This suggests that under LD cycles, the M-oscillator

is more effective at setting the phase of male than female behavior.

Under free-running conditions, both *Pdf > DBT*^{s} males and females flies showed a significantly

lower percentage of rhythmic flies compared to their controls (Fig. 4D-E), but there were fewer

rhythmic females (~40%) than males (~65%) (Fig. 4D). A possible reason for this could be that

the rest of the clock network, where the pace of the molecular clock remained unaltered, is able to
resist the influence of the M-cells to a greater extent in females. The free-running period (FRP) of
Pdf > DBT[s] males was ~18.5 hours, with very few exceptions (Fig. 4F). However, *Pdf > DBT*[s]
females showed a bimodal distribution, with some individuals showing a FRP consistent with the
pace of the M-cells (~18-18.5 hrs.), whereas a large proportion of females had FRP of ~24 hours,
consistent with the pace of the molecular clock in the rest of the clock network, which was not
affected by the genetic manipulation (Fig. 4F). Among rhythmic flies, both *Pdf > DBT*[s] males and
females had lower rhythmic power than the controls (Fig. 4G), with no difference between the
sexes (Fig. 4H). These results support the notion that M-cells are more dominant over the male
than the female circadian network. Previous studies have shown that M-cells' influence over the
rest of the clock neurons is mediated via PDF [65], so it is possible that the differential role of M
cells in males and females is precisely due to the sexually dimorphic effects of PDF.
230
231 **Speeding up the E-cell clock leads to a more advanced evening peak in females**
232
We next asked if changing the pace of the clock in the evening cells (E-cells) via DBT[S]
expression also had sexually dimorphic effects on behavior. These cells can be subdivided into at
least 3 different clusters, based on their anatomy [8], physiology [63], transcriptomics profiles [14]
and connectivity patterns [16]. The PDFR-expressing E1 and E2 clusters have been shown to
regulate evening activity under LD [65, 66] and to be able to maintain free-running activity
rhythms in the absence of a functional clock in the M cells [67, 68], while the behavioral role of
the E3 cluster remains unknown. We used the MB122-B split-gal4 driver to target the E1 and E2
subsets (Fig. 5A). We found that expressing DBT[S] in the E1+E2 subsets of evening cells
significantly advanced the phase of E-peak in experimental flies of both sexes (Fig. 5B-C).
Remarkably, this effect was more pronounced in females (Fig. 5D).
243
Speeding up of clocks in the E1 + E2 LN$_d$s did not significantly alter the free-running period or
rhythmic power of experimental flies of either sex (Fig. 5E-F). M-cells have been shown to be the
most dominant oscillators in DD and to regulate rhythm properties like persistence and free-
running period to a large extent [17, 21, 22, 63], although manipulations of other clock cells can
affect rhythm properties to some extent. Thus, speeding up the clock in the PDFR[+] E1 and E2

clusters leads to similar behavioral phenotypes in males and females under free-running

conditions.

Discussion

The critical role of sex as a biological variable has gained increasing recognition in biomedical research [69, 70]. Male bias is particularly prevalent in neuroscience, with single-sex studies using male animals outnumber those using females by a ratio of 5.5 to 1 [71]. This disparity extends to chronobiology, leading to a lack of knowledge regarding how temporal organization in the nervous system is influenced by sex. However, work from several laboratories has revealed sexual dimorphism in the SCN, as well as in its input and output pathways [1, 72]. Sex differences in the SCN morphology have been described in both animal models and humans, as well as sex differences in SCN electrical activity and steroid hormone receptors. In addition, sex differences in the number of SCN neurons that express the key neuropeptide vasoactive intestinal polypeptide (VIP) and in *Vip* mRNA *e*xpression have also been reported (reviewed in [2]).The roles of the mammalian VIP and the *Drosophila* PDF to circadian physiology are highly similar, although neither these peptides nor their receptors are sequence orthologs [73].

Several studies over the years have shown the importance of PDF in generating coherent rhythms of ~2hour periodicity. Here, we report that females lacking *Pdf* or its receptor *PdfR* are more likely to be able to maintain consolidated activity-rest behavior than males. This could be because of sex differences in PDF signaling mechanisms, PDFR expression, or the roles of other clock neurons within the network. We did not detect sex differences in the overall levels or temporal pattern of PDF accumulation, suggesting that these differences might not contribute to the sexually dimorphic effects of the loss of *Pdf* expression. Another possibility is that in the female circadian system, other neuropeptides or neurotransmitters can maintain network synchronization even in the absence of PDF. In males, other neuropeptides have been shown to act in concert with PDF to maintain consolidated rhythms in the network, although none of them have as profound an effect as PDF in regulating activity-rest rhythms in DD [51, 74]. Single mutants of DH31 and CCHamide1 do not affect activity rhythms by themselves but the double mutants of these neuropeptides along with Pdf^{01} ($DH31^{01}Pdf^{01}$ and $Pdf^{01}/CCHa^{SK8}$) are almost completely arrhythmic, indicating that these neuropeptides act hierarchically in the network, with PDF being on top of that hierarchy [51, 74]. The importance of PDF relative to other peptides released by clock neurons may also be sexually dimorphic.

To overcome possible pleiotropic effects of Pdf^{01} and $PdfR$ mutations, we employed a

constitutive knockout of Pdf in the ventral lateral neurons using the CRISPR-Cas9 method.

Although the Crispr manipulation was only partially effective at eliminating PDF expression, it

was able to replicate the phenotypes of the Pdf^{01} mutation. We observed faint staining in the dorsal

projections of at least one s-LN$_v$ in at least one hemisphere in most brains, and PDF staining in a

single s-LN$_v$ projection reaching the dorsal brain has been shown to be sufficient for behavioral

rhythms [75]. Experimental flies in which Pdf was mutagenized starting at the onset of the

promoter expression early in development showed extensive misrouting of their dorsal termini,

similar to what has been reported for Pdf^{01} males [60]. Instances of s-LN$_v$ misrouting have also

been observed in other core clock mutants such as per^{01} and tim^{01} [76], and cyc^{01} [77]. We did not

find evidence of sex differences in the severity of the misrouting phenotype caused by the loss of

$Pdf,$ but the percentage of brains with misrouting was higher in experimental females (data not

shown), and no correlation between misrouting and behavioral phenotypes was found by others

for pdf^{01} males [60]. Importantly, the $Pdf > Pdfg;Cas9$ manipulation recapitulates the sexually

dimorphic circadian phenotypes of Pdf^{01} mutants: a larger fraction of females are rhythmic and

females exhibit a higher rhythm power.

In males, changing the speed of the M cell clock leads to phase changes in the morning peak

under LD [65]. To determine if M-cell manipulations also have sexually dimorphic effects on

behavior, we sped up the molecular clock by expressing the $Doubletime\ short$ allele, DBT^s.

Surprisingly, our results showed that speeding up the clock of M-cells advances the phase of the

morning peak in males, while the female morning peak phase is not affected. These results support

previous studies conducted in males on the role of M-cells in regulating the morning peak of

activity [17, 22, 65], and suggest that the M cells are unable to regulate the phase of the morning

activity in the same way in females. In DD, males had largely coherent short period rhythms and

the majority (65%) were rhythmic. In contrast, only 40% of the females were rhythmic, and their

FRP showed a bimodal distribution, with most flies exhibiting a ~24-hour period. This further

supports the notion that the M-cells are less dominant over the circadian clock network in females.

One possible explanation is that other clock neurons are able to "resist" their influence, and the

conflict between the fast-paced M cell clock and the ~24 clock other clock neurons is what leads
to higher arrhythmicity in the females.

The Cryptochrome and *PdfR*-expressing clusters of evening cells – the sNPF expressing E1
cluster and the ITP-expressing E2 cluster [78] – have roles in setting the phase of the E-peak under
LD and sustaining behavioral rhythms in the absence of a functional molecular clock in the M-
cells [67, 68]. To test if these cells have a differential influence over the network in males and
females, we expressed DBT^S under a driver that is expressed specifically in the E1 and E2 subsets
of LN_ds. Our results showed that speeding up the clocks in the E1+E2 clusters resulted in a phase
advance in evening peak of activity in both the sexes, but the effect was more pronounced in the
females. Neither MB122 > DBT^S males nor females had phenotypes under free running conditions.
A possible reason for the behavioral differences seen between the sexes could be redundancy in
females, such that the network is not as dependent on PDF and therefore on M-cells for
timekeeping. This would mean that the female network could have more distributed mode of
timekeeping throughout the circadian clock network.

Across species, sex differences in the circadian timing system are largely related to the
regulation of reproduction relevant behaviors. In mammals, the SCN determines the timing of the
release of reproductive hormones and influences timing of mating (reviewed in [72]) and
aggression [79] . In *Drosophila*, the circadian clock controls the timing of sex-specific and sexually
dimorphic behaviors, such as male courtship [80] and aggression [81] and female sexual
receptivity [82] and egg-laying [83]. This regulation of rhythmic behaviors requires connectivity
between clock neurons and downstream sex specific circuits. For example, the $DN1_p$ cluster, which
has been shown to be more active in males [39], is functionally connected to the male-specific *fru*-
expressing P1 neurons that regulate male courtship [40]. In females, Allostatin C- producing
$DN1_p$s have been shown to connect to downstream targets to control rhythms in oogenesis [84],
and the Janelia female hemibrain connectome revealed that the LN_ds form connections *doublesex*-
expressing PC1 cluster [16]. Our data suggest that the relative hierarchy of circadian oscillators is
sexually dimorphic, with a less dominant M oscillator in females. This pattern of circadian
timekeeping may serve an adaptive purpose, ensuring precise timing of essential female-specific
behaviors crucial for reproductive fitness, such as sexual receptivity and egg laying.

Acknowledgments

We are grateful to Abhilash Lakshman, Jeff Price, Orie Shafer, and Paul Taghert for helpful discussions members of the Fernandez Lab for helpful comments on the manuscript. We also thank Justin Blau, Aljoscha Nern, Michael Rosbash, Gerry Rubin, and Paul Taghert for sharing fly lines. The mouse anti-PDF antibody was obtained from the Developmental Studies Hybridoma Bank, created by the NICHD of the NIH and maintained at The University of Iowa, Department of Biology, Iowa City, IA 52242. Stocks obtained from the Bloomington *Drosophila* Stock Center (NIH P40OD018537) were used in this study. This work was supported by NIH grant (R01NS118012), an NSF Grant (IOS-2239994) to M.P.F. and Barnard Internal Funds (SRI) to E.S-C., G.B. and E.B.

Materials and Methods

Fly lines and rearing

All genotypes were reared on standard cornmeal medium under LD (12 hr Light: 12 hr Dark) cycles and 25°C, unless specified otherwise (see figure legends for details). The fly lines used in this study were *Canton-S, w^{1118}, Pdf^{01}, PdfR^{01}, w; Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}, w*;UAS-*Cas9;* UAS-*Pdfg*, w; *Pdf-Gal4, w;;*UAS-*DBT^{s},* and *w;MB122B-Gal4;*.

Activity-rest behavior recording and analysis

Individual male and virgin female flies (3–5 days old) were housed in glass locomotor tubes with sucrose agar food on one end and yarn on the other end. Locomotor activity was recorded using the Drosophila Activity Monitors (DAM, Trikinetics, Waltham, United States of America). Experiments were conducted in Tritech or Percival incubators under controlled light and temperature conditions. Flies were entrained to 12:12 LD cycles for at least 5 days, and then transferred to constant darkness (DD) for at least 7 days, at a constant temperature of 25°C, unless otherwise specified (see figure legends for details). Raw data obtained from the DAM system were scanned and binned into activity counts of 15 min interval using DAM File scan. Data was analyzed using the CLOCKLAB software (Actimetrics, Wilmette, IL).

Values of period and rhythmic power were calculated for a period of 8 days using the Chi-square periodogram with a cut-off of *p =0.01*. The Rhythmic Power for each designated rhythmic fly was determined by subtracting the Chi-squared significance value from the Power of the Periodogram. Flies that did not exhibit a periodicity peak above the significance threshold were categorized as "arrhythmic," and their period and rhythmic power were not included in the analysis. Values of the Morning and Evening peaks was calculated using the PHASE software [85]. Total LD sleep values for all genotypes were calculated for a period of 3 days (LD days 2-4). Representative actograms were generated using ClockLab and activity plots were generated using PHASE. The period, rhythmic power, total sleep, phase values of all the flies for a particular experimental genotype were compared against the background or parental controls either using the Mann-Whitney test or the Kruskal-Wallis ANOVA followed by the Dunn's multiple comparisons test. The details on the statistical comparisons and the no. of flies used in a given experiment are indicated in their

respective figure legends. The number of rhythmic flies of the experimental genotype were

compared against their respective background or parental controls using the Fisher's exact test. All

statistical analysis was done using GraphPad Prism 9.0.

Immunohistochemistry

Brains from adult male or female flies (6-8 days old) were dissected in ice-cold Schneider's Insect

media (S2) and fixed immediately after dissection in 2%Paraformaldehyde (PFA) in S2 media for

30 minutes at room temperature. The fixed brains were washed (3 washes of 10 mins each) with

0.3% PBS-Triton X 100 (PBS-TX) and then treated with the blocking solution (5% Normal Goat

Serum made in 0.3% PBS-TX) for 1-hr at room temperature. The brains were then incubated with

primary antibodies at 4°C for 24-48 hrs. The primary antibodies used were anti-PDF (mouse,

1:3000, C7, DSHB), anti-RFP (rabbit, 1:2000, Rockland). After incubation, the brains were given

6 washes with 0.3% PBS-TX and incubated with Alexa-fluor conjugated secondary antibodies

overnight or for 24-48 hrs. at 4°C. The following secondary antibodies were used, goat anti-mouse

488 (1:3000, Invitrogen), goat anti-rabbit 568 (1:3000, Invitrogen). After incubation, the brain

samples were washed 6 times with 0.3% PBS-TX, cleaned and mounted on a clean, glass slide

using Vectashield mounting media.

Image acquisition and analysis

The slides were imaged using a confocal microscope (Olympus FV3000) using the 20X objective.

Image analysis was performed using the Fiji software [86]. In the samples, small and large ventral

lateral neurons were classified based on their anatomical locations and expression of PDF. PDF

intensity in these cells were measured by selecting the slice of the Z-stack that showed the

maximum intensity, drawing a Region of Interest (ROI) around the cells and measuring their

intensities. 3–4 separate background values were also measured around each cell and the final

intensity was taken as the difference between the cell intensity and the background. For

quantification of PDF in the dorsal projections, a rectangular box was drawn as ROI starting from

the point where the PDF projection turns into the dorsal brain and intensity was measured. 3–4

background values were also measured around the projection. The intensity values obtained from

both the hemispheres for each cell type for each brain was averaged and used for statistical

analysis. PDF intensity from the s-LN$_v$ were compared between the experimental and control

genotypes using the Mann-Whitney test. To estimate different aspects of rhythmicity in PDF

oscillations in the dorsal termini of s-LN$_v$ in males and females, we used a COSINOR based curve-

fitting method [54] COSINOR analysis was implemented using the CATCosinor function from the

CATkit package written for R [87].

Figure legends

Figure 1: Mutations in *Pdf* and *PdfR* lead to sexually dimorphic phenotypes. (A)

Representative actograms of *w^{1118}* (left) and *Pdf^{01}* (right) male flies for 6 days of LD followed by

7 days of DD. **(B)** Representative actograms of *w^{1118}* (left) and *Pdf^{01}* (right) female flies for 6 days

of LD followed by 7 days of DD. **(C)** Percentage of rhythmic flies are plotted for control (*w^{1118}*)

and *Pdf^{01}* males (left, *n* = 86 (*w^{1118}*), *n* = 81 (*Pdf^{01}*)) and females (right, *n* = 75 (*w^{1118}*), *n* = 101

(*Pdf^{01}*)). Error bars are SEM values plotted across 3 replicate experiments. **(D)** Rhythmic power

of control (*w^{1118}*) and *Pdf^{01}* male and females calculated using the Chi-square periodogram are

plotted. **(E)** The difference in rhythmic power of experimental males and females from their

respective controls are plotted. **(F)** Average sleep plots of flies over 7 days in DD are plotted for

male and female control (*w^{1118}*) and experimental (*Pdf^{01}*) flies. **(G)** Average sleep plots under LD

12:12 of control (*w^{1118}*) and experimental (*Pdf^{01}*) are plotted for males (left) and females (right).

The plots are averaged over flies and days for a period of 3 days under LD 12:12. **(H)** Total sleep

values under LD are plotted for male (left) and female (right) flies of control (*w^{1118}*) and

experimental (*Pdf^{01}*) flies. **(I)** The difference in total LD sleep values of experimental males and

females from their respective controls are plotted. **(J)** Percentage of rhythmic flies are plotted for

control (*w^{1118}*) and *han^{5304}* males (left, *n* = 86 (*w^{1118}*), *n* = 115 (*han^{5304}*)) and females (right, *n* = 74

(*w^{1118}*), *n* = 94 (*han^{5304}*)). **(K)** Rhythmic power of control (*w^{1118}*) and *han^{5304}* male and females

calculated using the Chi-square periodogram are plotted. **(L)** The difference in rhythmic power of

experimental males and females from their respective controls are plotted. **(M)** Average sleep plots

under LD 12:12 of control (*w^{1118}*) and experimental (*han^{5304}*) flies are plotted for males (left) and

females (right). The plots are averaged over flies and days for a period of 3 days under LD 12:12.

(N) Total sleep values under LD are plotted for male (left) and female (right) flies of control (*w^{1118}*)

and experimental (*han^{5304}*) flies. **(O)** The difference in total LD sleep values of experimental males

and females from their respective controls are plotted. Free-running period and rhythmic power

values are calculated using Chi square periodogram in CLOCKLAB for a period of 7 days. Total

sleep values are calculated for 3 days in LD using PHASE. Statistical comparisons were performed

between the control and experimental flies for both sexes using the Mann-Whitney U test. The box

plots extend from the 25^{th} to 75^{th} percentile, with whiskers extending from the smallest to the

largest value, and each point representing data from a single fly. Combined data from at least 3

replicate experiments are plotted.

Figure 2: Tissue-specific CRISPR mediated *Pdf* manipulation leads to a reduction in PDF

levels and misrouting of s-LN$_v$ dorsal termini. (A) Representative confocal images of control

(*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*) (top) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} >*

Cas9; Pdf-g) (middle) male flies stained with RFP and PDF antibodies. Representative confocal

images of experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) (bottom) female flies

stained with RFP and PDF antibodies. The experimental flies show significant reduction in PDF

levels and misrouting of the s-LN$_v$ dorsal termini (arrows) as seen in the RFP channel for both the

sexes. **(B)** Quantification of PDF intensity from s-LN$_v$ cell bodies in control (*Pdf-RFP, Pdf-Gal4;*

tub-Gal80^{ts}) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) flies are plotted

for males (left) and females (right). Data points represent intensity from a single brain sample. *n* >

12 brains for all genotypes. **(C)** The difference in PDF intensity values of experimental males and

females from their respective controls are plotted.

Figure 3: CRISPR-Cas9 mediated *Pdf* mutagenesis has more pronounced effects on male

behavior. (A) Representative actograms of control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*), and *(UAS*

Cas9; Pdf-g) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) males (left) and

females (right) are plotted for 6 days of LD followed by 9 days of DD. **(B)** Percentage of rhythmic

flies are plotted for control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}), and (UAS Cas9; Pdf-g),* and

experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g)* males (left, *n* = 85 (*Pdf-RFP, Pdf-*

Gal4; tub-Gal80^{ts}), *n* = 89 (*UAS Cas9; Pdf-g*), *n* = 106 (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9;*

Pdf-g) and females (right, *n* = 79 (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*), *n* = 78 (*UAS Cas9; Pdf-g*), *n*

= 85 (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g).* Error bars are SEM values plotted across 3

replicate experiments. **(C)** Free-running period of control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and*

UAS Cas9; Pdf-g) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g)* males (left)

and females (right) calculated using the Chi-square periodogram are plotted. **(D)** Rhythmic power

of control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g)* and experimental (*Pdf-RFP,*

Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g) males (left) and females (right) calculated using the Chi-

square periodogram are plotted **(E)** The difference in rhythmic power of experimental males and

females from their respective controls are plotted. Flies were kept at 28°C throughout development

and as adults. The activity-rest behavior was recorded at 28°C. Free-running period and rhythmic

power values are calculated using Chi square periodogram in CLOCKLAB for a period of 7 days.

Statistical comparisons were performed between the control and experimental flies for both sexes

using the Kruskal-Wallis ANOVA followed by the Dunn's multiple comparisons test. The box plots

extend from the 25th to 75th percentile, with whiskers extending from the smallest to the largest

value, and each point representing data from a single fly. Combined data from at least 3 replicate

experiments are plotted.

Figure 4: Speeding up the clock in M-cells leads to sexually dimorphic phenotypes. (A)

Depiction of the adult *Drosophila* brain hemisphere indicating the clock cell subsets (colored)

having a faster running molecular clock. **(B)** Average activity plots of control (*Pdf-Gal4*) and

(UAS-*DBT^s*) and experimental (*Pdf > DBT^s*) flies are plotted for males (left) and females (right).

The plots are averaged over flies and days for a period of 3 days under LD 12:12. **(C)** The phase

of the morning peak of activity under LD 12:12 is plotted for control (*Pdf-Gal4*), (UAS-*DBT^s*) and

experimental (*Pdf >DBT^s*) males (left, *n* = 116 (*Pdf-Gal4*), *n* = 120 (UAS-*DBT^s*), *n* = 88 (*Pdf >*

DBT^s)) and females (right, *n* = 85 (*Pdf-Gal4*), *n* = 109 (UAS-*DBT^s*), *n* = 73 (*Pdf > DBT^s*)). **(D)**

Percentage of rhythmic flies are plotted for controls (*Pdf-Gal4), (UAS-DBT^s),* and experimental

(*Pdf > DBT^s)* males (left, *n* = 90 (*Pdf-Gal4*), *n* = 108 (*UAS-DBT^s*), *n* = 93 (*Pdf > DBT^s*) and females

(right, *n* = 66 (*Pdf-Gal4*), *n* = 83 (*UAS-DBT^s*), *n* = 70 (*Pdf > DBT^s). Error bars are SEM values

plotted across 3 replicate experiments. **(E)** Representative actograms of control (UAS-*DBT^s*) and

experimental (*Pdf > DBT^s*) males (left) and females (right) plotted for 6 days of LD followed by

10 days of DD. **(F)** Free-running period of control (*Pdf-Gal4), (UAS-DBT^s)* and experimental (*Pdf*

> DBT^s) males (left) and females (right) calculated using the Chi-square periodogram are plotted.

(G) Rhythmic power of control (*Pdf-Gal4), (UAS-DBT^s)* and experimental (*Pdf > DBT^s)* males

(left) and females (right) calculated using the Chi-square periodogram are plotted. **(H)** The

difference in rhythmic power of experimental males and females from their respective controls are plotted. Free-running period and rhythmic power values are calculated using Chi-square periodogram in Clocklab for a period of 7 days. Activity plots and phase of the M-peak were calculated using PHASE. Statistical comparisons were performed between the control and experimental flies for both sexes using the Kruskal-Wallis ANOVA followed by the Dunn's multiple comparisons test. The box plots extend from the 25^{th} to 75^{th} percentile, with whiskers extending from the smallest to the largest value, and each point representing data from a single fly. Combined data from at least 3 replicate experiments are plotted.

Figure 5: Speeding up the clock in a E-cell subset leads to a more advanced evening peak in females. (A) Depiction of the adult *Drosophila* brain hemisphere indicating the clock cell subsets (colored) having a faster running molecular clock. **(B)** Average activity plots of control (*MB122B-Gal4*) and (*UAS-DBT^s*) and experimental (*MB122B > DBT^s*) flies are plotted for males (left) and females (right). The plots are averaged over flies and days for a period of 3 days under LD 12:12. **(C)** Phase of the evening peak of activity under LD 12:12 for controls (*MB122B-Gal4*) and (*UAS-DBT^s*) and experimental (*MB122B > DBT^s*) males (left, $n = 92$ (*MB122B-Gal4*), $n = 93$ (*UAS-DBT^s*), $n = 93$ (*MB122B > DBT^s*)) and females (right, $n = 85$ (*MB122B-Gal4*), $n = 87$ (*UAS-DBT^s*), $n = 89$ (*MB122B > DBT^s*)). **(D)** The difference in the phase of E-peak of experimental males and females from their respective controls are plotted. **(E)** Free-running period of control (*MB122B-Gal4* and *UAS-DBT^s)* and experimental (*MB122B > DBT^s)* males (left) and females (right) calculated using the Chi-square periodogram are plotted. **(F)** Rhythmic power of control (*MB122B-Gal4 and UAS-DBT^s)* and experimental (*MB122B > DBT^s)* males (left) and females (right) calculated using the Chi-square periodogram are plotted. Free-running period and rhythmic power values are calculated using Chi-square periodogram in CLOCKLAB for a period of 7 days. Activity plots and phase of the M-peak were calculated using PHASE. Statistical comparisons were performed between the control and experimental flies for both sexes using the Kruskal-Wallis ANOVA followed by the Dunn's multiple comparisons test. The box plots extend from the 25^{th} to 75^{th} percentile, with whiskers extending from the smallest to the largest value, and each point representing data from a single fly. Combined data from at least 3 replicate experiments are plotted.

Supplementary figures

Figure S1: PDF rhythmic accumulation is similar in male and female brains. (Related to Figure 1.) (A) Representative confocal images of control (w^{1118}) (top) and experimental (Pdf^{01}) male (middle) and female (bottom) flies stained with PDF antibodies. **(B)** Scatterplots of PDF staining intensities in s-LN$_v$ dorsal projections of both male and female flies plotted at different time-points over a 24-hr cycle on DD day3. Each dot represents the mean PDF intensity value averaged over both the hemispheres of one brain. The cyan and pink lines are the best fit Cosine curves from the parameters that were extracted using COSINOR analysis. See Table 2 for more details. **(C)** Amplitude values of PDF oscillation obtained from COSINOR curve fits are plotted for male and female flies. Error bars are 95% CI values calculated from the standard error obtained from COSINOR analysis. Overlapping error bars indicate that amplitude values of males and females are not significantly different from controls. COSINOR analysis was implemented using the CATCosinor function from the CATkit package written for R [87]. n > 7 brain samples.

Figure S2: CRISPR-Cas9 mediated *Pdf* mutagenesis has similar effects on male and female sleep. (Related to Figure 3.) (A) Average sleep plots under LD 12:12 of control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g)* and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g)* males (left) and females (right) are plotted. The plots are averaged over flies and days for a period of 3 days under LD 12:12 at 28°C. **(B)** Total sleep values under LD are plotted for control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g)* and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g)* males (left) and females (right). **(C)** Average sleep plots of flies over 7 days in DD are plotted for male and female control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g*) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) flies. **(D)** Total sleep values under DD are plotted for control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g)* and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g)* males (left) and females (right). **(E)** The difference in the total DD sleep values of experimental males and females from their respective controls are plotted.

Figure S3: CRISPR-Cas9 mediated *Pdf* mutagenesis has more pronounced effects on male circadian behavior at 25C. (Related to Figure 3.) (A) Percentage of rhythmic flies are plotted for control (*Pdf-Gal4), and (UAS Cas9; Pdf-g),* and experimental (*Pdf > Cas9; Pdf-g*) males (left,

$n = 36$ (*Pdf-Gal4*), $n = 49$ (*UAS Cas9; Pdf-g*), $n = 52$ (*Pdf > Cas9; Pdf-g*) and females (right, $n = $

33 (*Pdf-Gal4*), $n = 46$ (*UAS Cas9; Pdf-g*), $n = 35$ (*Pdf > Cas9; Pdf-g*). Error bars are SEM values

plotted across 2 replicate experiments. **(B)** Free-running period of control (*Pdf-Gal4 and UAS*

Cas9; Pdf-g) and experimental (*Pdf > Cas9; Pdf-g*) males (left) and females (right) calculated

using the Chi-square periodogram are plotted. **(C)** Rhythmic power of control (*Pdf-Gal4 and UAS*

Cas9; Pdf-g) and experimental (*Pdf > Cas9; Pdf-g*) males (left) and females (right) calculated

using the Chi-square periodogram are plotted. **(D)** The difference in the rhythmic power values of

experimental males and females from their respective controls are plotted.

Figure S4: Adult specific CRISPR-Cas9 mediated *Pdf* mutagenesis has more pronounced

effects on male circadian behavior. (Related to Figure 3.) (A) Representative confocal images

of control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*) (top) and experimental (*Pdf-RFP, Pdf-Gal4; tub-*

Gal80^{ts} > Cas9; Pdf-g) male (middle) and female (bottom) flies stained with RFP and PDF

antibodies. The experimental flies show significant reduction in PDF levels but no misrouting of

the s-LN$_v$ dorsal termini. **(B)** Quantification of PDF intensity from s-LN$_v$ cell bodies in control

(*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-*

g) flies are plotted for males (left) and females (right). Data points represent intensity from a single

brain sample. $n > 10$ brains for all genotypes. **(C)** The difference in PDF intensity values of

experimental males and females from their respective controls are plotted. **(D)** Percentage of

rhythmic flies are plotted for control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}), and (UAS Cas9; Pdf-g*),

and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) males (left, $n = 103$ (*Pdf-RFP,*

Pdf-Gal4; tub-Gal80^{ts}), $n = 109$ (*UAS Cas9; Pdf-g*), $n = 115$ (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} >*

Cas9; Pdf-g) and females (right, $n = 96$ (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts}*), $n = 90$ (*UAS Cas9; Pdf-*

g), $n = 119$ (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*). Error bars are SEM values plotted

across 3 replicate experiments. **(E)** Free-running period of control (*Pdf-RFP, Pdf-Gal4; tub-*

Gal80^{ts} and UAS Cas9; Pdf-g) and experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*)

males (left) and females (right) calculated using the Chi-square periodogram are plotted. **(F)**

Rhythmic power of control (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} and UAS Cas9; Pdf-g*) and

experimental (*Pdf-RFP, Pdf-Gal4; tub-Gal80^{ts} > Cas9; Pdf-g*) males (left) and females (right)

calculated using the Chi-square periodogram are plotted. Flies were raised at 18°C until eclosion

and were then transferred to 28°C as adults. Adult flies were kept at 28°C for a period of 7 days

before recording activity-rest behavior at 28°C. Free-running period and rhythmic power values

are calculated using Chi square periodogram in CLOCKLAB for a period of 7 days. Statistical

comparisons were performed between the control and experimental flies for both sexes using the

Kruskal-Wallis ANOVA followed by the Dunn's multiple comparisons test. The box plots extend

from the 25th to 75th percentile, with whiskers extending from the smallest to the largest value, and

each point representing data from a single fly. Combined data from at least 3 replicate experiments

are plotted.

References
619

1.   Anderson ST, FitzGerald GA. Sexual dimorphism in body clocks. Science.
2020;369(6508):1164-5. Epub 2020/09/05. doi: 10.1126/science.abd4964. PubMed PMID:
32883849.
2.   Bailey M, Silver R. Sex differences in circadian timing systems: implications for disease.
Front Neuroendocrinol. 2014;35(1):111-39. Epub 2013/11/30. doi: 10.1016/j.yfrne.2013.11.003.
PubMed PMID: 24287074; PubMed Central PMCID: PMCPMC4041593.
3.   Vida B, Hrabovszky E, Kalamatianos T, Coen CW, Liposits Z, Kalló I. Oestrogen Receptor α
and β Immunoreactive Cells in the Suprachiasmatic Nucleus of Mice: Distribution, Sex Differences
and Regulation by Gonadal Hormones. Journal of Neuroendocrinology. 2008;20(11):1270-7. doi:
https://doi.org/10.1111/j.1365-2826.2008.01787.x.
4.   Lyall LM, Wyse CA, Graham N, Ferguson A, Lyall DM, Cullen B, et al. Association of
disrupted circadian rhythmicity with mood disorders, subjective wellbeing, and cognitive
function: a cross-sectional study of 91 105 participants from the UK Biobank. Lancet Psychiatry.
2018;5(6):507-14. Epub 2018/05/20. doi: 10.1016/S2215-0366(18)30139-1. PubMed PMID:
29776774.
5.   Bixler EO, Papaliaga MN, Vgontzas AN, Lin HM, Pejovic S, Karataraki M, et al. Women sleep
objectively better than men and the sleep of young women is more resilient to external stressors:
effects of age and menopause. J Sleep Res. 2009;18(2):221-8. Epub 2009/03/24. doi:
10.1111/j.1365-2869.2008.00713.x. PubMed PMID: 19302341; PubMed Central PMCID:
PMCPMC3594776.
6.   Hastings MH, Maywood ES, Brancaccio M. The Mammalian Circadian Timing System and
the Suprachiasmatic Nucleus as Its Pacemaker. Biology (Basel). 2019;8(1). Epub 20190311. doi:
10.3390/biology8010013. PubMed PMID: 30862123; PubMed Central PMCID: PMCPMC6466121.
7.   Helfrich-Forster C. The neuroarchitecture of the circadian clock in the brain of Drosophila
melanogaster. Microsc Res Tech. 2003;62(2):94-102. Epub 2003/09/11. doi: 10.1002/jemt.10357.
PubMed PMID: 12966496.
8.   Schubert FK, Hagedorn N, Yoshii T, Helfrich-Forster C, Rieger D. Neuroanatomical details
of the lateral neurons of Drosophila melanogaster support their functional role in the circadian
system. J Comp Neurol. 2018;526(7):1209-31. Epub 2018/02/10. doi: 10.1002/cne.24406.
PubMed PMID: 29424420; PubMed Central PMCID: PMCPMC5873451.
9.   Yu W, Hardin PE. Circadian oscillators of Drosophila and mammals. J Cell Sci. 2006;119(Pt
23):4793-5. Epub 2006/11/30. doi: 10.1242/jcs.03174. PubMed PMID: 17130292.
10.   Kloss B, Price JL, Saez L, Blau J, Rothenfluh A, Wesley CS, et al. The Drosophila clock gene
double-time encodes a protein closely related to human casein kinase Iepsilon. Cell.
1998;94(1):97-107. Epub 1998/07/23. doi: 10.1016/s0092-8674(00)81225-8. PubMed PMID:
9674431.
11.   Price JL, Blau J, Rothenfluh A, Abodeely M, Kloss B, Young MW. double-time is a novel
Drosophila clock gene that regulates PERIOD protein accumulation. Cell. 1998;94(1):83-95. Epub
1998/07/23. doi: 10.1016/s0092-8674(00)81224-6. PubMed PMID: 9674430.
12.   Helfrich-Forster C. Organization of endogenous clocks in insects. Biochem Soc Trans.
2005;33(Pt 5):957-61. Epub 2005/10/26. doi: 10.1042/BST20050957. PubMed PMID: 16246020.

13.   Dubowy C, Sehgal A. Circadian Rhythms and Sleep in Drosophila melanogaster. Genetics.
2017;205(4):1373-97. Epub 2017/04/01. doi: 10.1534/genetics.115.185157. PubMed PMID:
28360128; PubMed Central PMCID: PMCPMC5378101.
14.   Ma D, Przybylski D, Abruzzi KC, Schlichting M, Li Q, Long X, et al. A transcriptomic
taxonomy of Drosophila circadian neurons around the clock. eLife. 2021;10:e63056. doi:
10.7554/eLife.63056.
15.   Shafer OT, Helfrich-Forster C, Renn SC, Taghert PH. Reevaluation of Drosophila
melanogaster's neuronal circadian pacemakers reveals new neuronal classes. J Comp Neurol.
2006;498(2):180-93. Epub 2006/07/21. doi: 10.1002/cne.21021. PubMed PMID: 16856134;
PubMed Central PMCID: PMCPMC2596765.
16.   Shafer OT, Gutierrez GJ, Li K, Mildenhall A, Spira D, Marty J, et al. Connectomic analysis of
the Drosophila lateral neuron clock cells reveals the synaptic basis of functional pacemaker
classes. Elife. 2022;11. Epub 2022/06/30. doi: 10.7554/eLife.79139. PubMed PMID: 35766361;
PubMed Central PMCID: PMCPMC9365390.
17.   Grima B, Chelot E, Xia R, Rouyer F. Morning and evening peaks of activity rely on different
clock neurons of the Drosophila brain. Nature. 2004;431(7010):869-73. Epub 2004/10/16. doi:
10.1038/nature02935. PubMed PMID: 15483616.
18.   Rieger D, Shafer OT, Tomioka K, Helfrich-Forster C. Functional analysis of circadian
pacemaker neurons in Drosophila melanogaster. J Neurosci. 2006;26(9):2531-43. Epub
2006/03/03. doi: 10.1523/JNEUROSCI.1234-05.2006. PubMed PMID: 16510731.
19.   Renn SC, Park JH, Rosbash M, Hall JC, Taghert PH. A pdf neuropeptide gene mutation and
ablation of PDF neurons each cause severe abnormalities of behavioral circadian rhythms in
Drosophila. Cell. 1999;99(7):791-802. Epub 2000/01/05. PubMed PMID: 10619432.
20.   Park JH, Helfrich-Forster C, Lee G, Liu L, Rosbash M, Hall JC. Differential regulation of
circadian pacemaker output by separate clock genes in Drosophila. Proc Natl Acad Sci U S A.
2000;97(7):3608-13. Epub 2000/03/22. doi: 10.1073/pnas.070036197. PubMed PMID:
10725392; PubMed Central PMCID: PMCPMC16287.
21.   Stoleru D, Peng Y, Nawathean P, Rosbash M. A resetting signal between Drosophila
pacemakers synchronizes morning and evening activity. Nature. 2005;438(7065):238-42. Epub
2005/11/11. doi: 10.1038/nature04192. PubMed PMID: 16281038.
22.   Stoleru D, Peng Y, Agosto J, Rosbash M. Coupled oscillators control morning and evening
locomotor behaviour of Drosophila. Nature. 2004;431(7010):862-8. Epub 2004/10/16. doi:
10.1038/nature02926. PubMed PMID: 15483615.
23.   Fernandez MP, Pettibone HL, Bogart JT, Roell CJ, Davey CE, Pranevicius A, et al. Sites of
Circadian Clock Neuron Plasticity Mediate Sensory Integration and Entrainment. Curr Biol.
2020;30(12):2225-37 e5. Epub 2020/05/11. doi: 10.1016/j.cub.2020.04.025. PubMed PMID:
32386535; PubMed Central PMCID: PMCPMC7314648.
24.   Guo F, Chen X, Rosbash M. Temporal calcium profiling of specific circadian neurons in
freely moving flies. Proc Natl Acad Sci U S A. 2017;114(41):E8780-E7. Epub 2017/10/05. doi:
10.1073/pnas.1706608114. PubMed PMID: 28973886; PubMed Central PMCID:
PMCPMC5642695.
25.   Lamaze A, Stanewsky R. DN1p or the "Fluffy" Cerberus of Clock Outputs. Front Physiol.
2019;10:1540. Epub 2020/01/24. doi: 10.3389/fphys.2019.01540. PubMed PMID: 31969832;
PubMed Central PMCID: PMCPMC6960142.

26.   Collins B, Kane Elizabeth A, Reeves David C, Akabas Myles H, Blau J. Balance of Activity
between LNvs and Glutamatergic Dorsal Clock Neurons Promotes Robust Circadian Rhythms in
Drosophila. Neuron. 2012;74(4):706-18. doi: https://doi.org/10.1016/j.neuron.2012.02.034.
27.   Women, Minorities, and Persons with Disabilities in Science
and Engineering: 2021. Special Report NSF 21-321. Alexandria, VA: National Science Foundation:
National Center for Science and Engineering Statistics, 2021.
28.   Yoshii T, Wulbeck C, Sehadova H, Veleri S, Bichler D, Stanewsky R, et al. The neuropeptide
pigment-dispersing factor adjusts period and phase of Drosophila's clock. J Neurosci.
2009;29(8):2597-610. Epub 2009/02/27. doi: 10.1523/JNEUROSCI.5439-08.2009. PubMed PMID:
19244536.
29.   Peng Y, Stoleru D, Levine JD, Hall JC, Rosbash M. Drosophila free-running rhythms require
intercellular   communication.   PLoS   Biol.   2003;1(1):E13.   Epub   2003/09/17.   doi:
10.1371/journal.pbio.0000013.   PubMed   PMID: 12975658;   PubMed   Central   PMCID:
PMCPMC193604.
30.   Liang X, Holy TE, Taghert PH. A Series of Suppressive Signals within the Drosophila
Circadian Neural Circuit Generates Sequential Daily Outputs. Neuron. 2017;94(6):1173-89 e4.
Epub 2017/05/30. doi: 10.1016/j.neuron.2017.05.007. PubMed PMID: 28552314; PubMed
Central PMCID: PMCPMC5502710.
31.   Mertens I, Vandingenen A, Johnson EC, Shafer OT, Li W, Trigg JS, et al. PDF Receptor
Signaling in Drosophila Contributes to Both Circadian and Geotactic Behaviors. Neuron.
2005;48(2):213-9. doi: https://doi.org/10.1016/j.neuron.2005.09.009.
32.   Hyun S, Lee Y, Hong ST, Bang S, Paik D, Kang J, et al. Drosophila GPCR Han is a receptor for
the circadian clock neuropeptide PDF. Neuron. 2005;48(2):267-78. Epub 2005/10/26. doi:
10.1016/j.neuron.2005.08.025. PubMed PMID: 16242407.
33.   Lear BC, Merrill CE, Lin JM, Schroeder A, Zhang L, Allada R. A G protein-coupled receptor,
groom-of-PDF, is required for PDF neuron action in circadian behavior. Neuron. 2005;48(2):221-
7. Epub 2005/10/26. doi: 10.1016/j.neuron.2005.09.008. PubMed PMID: 16242403.
34.   Kim WJ, Jan LY, Jan YN. A PDF/NPF neuropeptide signaling circuitry of male Drosophila
melanogaster controls rival-induced prolonged mating. Neuron. 2013;80(5):1190-205. doi:
10.1016/j.neuron.2013.09.034.   PubMed   PMID: 24314729;   PubMed   Central   PMCID:
PMCPMC3867312.
35.   Krupp JJ, Billeter JC, Wong A, Choi C, Nitabach MN, Levine JD. Pigment-dispersing factor
modulates pheromone production in clock cells that influence mating in drosophila. Neuron.
2013;79(1):54-68. Epub 2013/07/16. doi: 10.1016/j.neuron.2013.05.019. PubMed PMID:
23849197; PubMed Central PMCID: PMCPMC3955580.
36.   Inami S, Sato S, Kondo S, Tanimoto H, Kitamoto T, Sakai T. Environmental Light Is Required
for Maintenance of Long-Term Memory in Drosophila. J Neurosci. 2020;40(7):1427-39. Epub
2020/01/15. doi: 10.1523/JNEUROSCI.1282-19.2019. PubMed PMID: 31932417; PubMed Central
PMCID: PMCPMC7044726.
37.   Mertens I, Vandingenen A, Johnson EC, Shafer OT, Li W, Trigg JS, et al. PDF receptor
signaling in Drosophila contributes to both circadian and geotactic behaviors. Neuron.
2005;48(2):213-9. Epub 2005/10/26. doi: 10.1016/j.neuron.2005.09.009. PubMed PMID:
16242402.

38.   Toma DP, White KP, Hirsch J, Greenspan RJ. Identification of genes involved in Drosophila
melanogaster geotaxis, a complex behavioral trait. Nat Genet. 2002;31(4):349-53. Epub
2002/06/04. doi: 10.1038/ng893. PubMed PMID: 12042820.
39.   Guo F, Yu J, Jung HJ, Abruzzi KC, Luo W, Griffith LC, et al. Circadian neuron feedback controls
the Drosophila sleep–activity profile. Nature. 2016;536:292. doi: 10.1038/nature19097
https://www.nature.com/articles/nature19097#supplementary-information.
40.   Chen D, Sitaraman D, Chen N, Jin X, Han C, Chen J, et al. Genetic and neuronal mechanisms
governing the sex-specific interaction between sleep and sexual behaviors in Drosophila. Nat
Commun. 2017;8(1):154. Epub 2017/07/30. doi: 10.1038/s41467-017-00087-5. PubMed PMID:
28754889; PubMed Central PMCID: PMCPMC5533705.
41.   Kunst M, Hughes ME, Raccuglia D, Felix M, Li M, Barnett G, et al. Calcitonin gene-related
peptide neurons mediate sleep-specific circadian output in Drosophila. Curr Biol.
2014;24(22):2652-64. Epub 2014/12/03. doi: 10.1016/j.cub.2014.09.077. PubMed PMID:
25455031; PubMed Central PMCID: PMCPMC4255360.
42.   Asahina K. Sex differences in Drosophila behavior: Qualitative and Quantitative
Dimorphism. Curr Opin Physiol. 2018;6:35-45. Epub 2018/11/06. doi:
10.1016/j.cophys.2018.04.004. PubMed PMID: 30386833; PubMed Central PMCID:
PMCPMC6205217.
43.   Helfrich-Forster C. Differential control of morning and evening components in the activity
rhythm of Drosophila melanogaster--sex-specific differences suggest a different quality of activity.
J Biol Rhythms. 2000;15(2):135-54. Epub 2000/04/13. doi: 10.1177/074873040001500208.
PubMed PMID: 10762032.
44.   Palmateer CM, Artikis C, Brovero SG, Friedman B, Gresham A, Arbeitman MN. Single-cell
transcriptome profiles of Drosophila fruitless-expressing neurons from both sexes. eLife.
2023;12:e78511. doi: 10.7554/eLife.78511.
45.   Fujii S, Amrein H. Ventral lateral and DN1 clock neurons mediate distinct properties of
male sex drive rhythm in Drosophila. Proc Natl Acad Sci U S A. 2010;107(23):10590-5. Epub
2010/05/26. doi: 10.1073/pnas.0912457107. PubMed PMID: 20498055; PubMed Central PMCID:
PMCPMC2890825.
46.   Hanafusa S, Kawaguchi T, Umezaki Y, Tomioka K, Yoshii T. Sexual interactions influence the
molecular oscillations in DN1 pacemaker neurons in Drosophila melanogaster. PLoS One.
2013;8(12):e84495. Epub 2013/12/25. doi: 10.1371/journal.pone.0084495. PubMed PMID:
24367668; PubMed Central PMCID: PMCPMC3867508.
47.   Lee G, Bahn JH, Park JH. Sex- and clock-controlled expression of the neuropeptide F gene
in Drosophila. Proc Natl Acad Sci U S A. 2006;103(33):12580-5. Epub 20060807. doi:
10.1073/pnas.0601171103. PubMed PMID: 16894172; PubMed Central PMCID:
PMCPMC1567921.
48.   Hermann C, Yoshii T, Dusik V, Helfrich-Forster C. Neuropeptide F immunoreactive clock
neurons modify evening locomotor activity and free-running period in Drosophila melanogaster.
J Comp Neurol. 2012;520(5):970-87. Epub 2011/08/10. doi: 10.1002/cne.22742. PubMed PMID:
21826659.
49.   Nässel DR, Zandawala M. Recent advances in neuropeptide signaling in Drosophila, from
genes to physiology and behavior. Prog Neurobiol. 2019;179:101607. Epub 2019/03/25. doi:
10.1016/j.pneurobio.2019.02.003. PubMed PMID: 30905728.

50.   Shafer OT, Yao Z. Pigment-Dispersing Factor Signaling and Circadian Rhythms in Insect
Locomotor Activity. Curr Opin Insect Sci. 2014;1:73-80. Epub 2014/11/12. doi:
10.1016/j.cois.2014.05.002. PubMed PMID: 25386391; PubMed Central PMCID:
PMCPMC4224320.

51.   Kuwano R, Katsura M, Iwata M, Yokosako T, Yoshii T. Pigment-dispersing factor and
CCHamide1 in the Drosophila circadian clock network. Chronobiol Int. 2023;40(3):284-99. Epub
2023/02/15. doi: 10.1080/07420528.2023.2166416. PubMed PMID: 36786215.

52.   Parisky KM, Agosto J, Pulver SR, Shang Y, Kuklin E, Hodge JJ, et al. PDF cells are a GABA-
responsive wake-promoting component of the Drosophila sleep circuit. Neuron. 2008;60(4):672-
82. Epub 2008/11/29. doi: 10.1016/j.neuron.2008.10.042. PubMed PMID: 19038223; PubMed
Central PMCID: PMCPMC2734413.

53.   Klose MK, Bruchez MP, Deitcher DL, Levitan ES. Temporally and spatially partitioned
neuropeptide release from individual clock neurons. Proc Natl Acad Sci U S A. 2021;118(17). Epub
2021/04/21. doi: 10.1073/pnas.2101818118. PubMed PMID: 33875606; PubMed Central PMCID:
PMCPMC8092580.

54.   Cornelissen G. Cosinor-based rhythmometry. Theoretical Biology and Medical Modelling.
2014;11(1):16. doi: 10.1186/1742-4682-11-16.

55.   Im SH, Li W, Taghert PH. PDFR and CRY signaling converge in a subset of clock neurons to
modulate the amplitude and phase of circadian behavior in Drosophila. PLoS One.
2011;6(4):e18974. Epub 2011/05/12. doi: 10.1371/journal.pone.0018974. PubMed PMID:
21559487; PubMed Central PMCID: PMCPMC3084726.

56.   Reinhard N, Fukuda A, Manoli G, Derksen E, Saito A, Möller G, et al. Synaptic and
peptidergic connectomes of the Drosophila circadian clock. bioRxiv.
2023:2023.09.11.557222. doi: 10.1101/2023.09.11.557222.

57.   Richhariya S, Shin D, Le JQ, Rosbash M. Dissecting neuron-specific functions of circadian
genes using modified cell-specific CRISPR approaches. Proc Natl Acad Sci U S A.
2023;120(29):e2303779120. Epub 2023/07/10. doi: 10.1073/pnas.2303779120. PubMed PMID:
37428902; PubMed Central PMCID: PMCPMC10629539.

58.   Gorostiza EA, Depetris-Chauvin A, Frenkel L, Pírez N, Ceriani María F. Circadian Pacemaker
Neurons Change Synaptic Contacts across the Day. Current Biology. 2014;24(18):2161-7. doi:
https://doi.org/10.1016/j.cub.2014.07.063.

59.   Ruben M, Drapeau MD, Mizrak D, Blau J. A mechanism for circadian control of pacemaker
neuron excitability. J Biol Rhythms. 2012;27(5):353-64. doi: 10.1177/0748730412455918.
PubMed PMID: 23010658; PubMed Central PMCID: PMCPMC4019749.

60.   Gorostiza EA, Ceriani MF. Retrograde Bone Morphogenetic Protein Signaling Shapes a Key
Circadian Pacemaker Circuit. The Journal of Neuroscience. 2013;33(2):687-96. doi:
10.1523/jneurosci.3448-12.2013.

61.   Fleck SA, Biswas P, DeWitt ED, Knuteson RL, Eisman RC, Nemkov T, et al. Auxin Exposure
Disrupts Feeding Behavior and Fatty Acid Metabolism in Adult Drosophila. bioRxiv. 2023. Epub
2023/08/30. doi: 10.1101/2023.08.15.553385. PubMed PMID: 37645868; PubMed Central
PMCID: PMCPMC10462055.

62.   McGuire SE, Mao Z, Davis RL. Spatiotemporal gene expression targeting with the TARGET
and gene-switch systems in Drosophila. Sci STKE. 2004;2004(220):pl6. Epub 2004/02/19. doi:
10.1126/stke.2202004pl6. PubMed PMID: 14970377.

63.   Yao Z, Shafer OT. The Drosophila circadian clock is a variably coupled network of multiple
peptidergic   units.   Science.   2014;343(6178):1516-20.   Epub   2014/03/29.   doi:
10.1126/science.1251285.   PubMed   PMID: 24675961;   PubMed   Central   PMCID:
PMCPMC4259399.
64.   Muskus MJ, Preuss F, Fan J-Y, Bjes ES, Price JL. Drosophila DBT Lacking Protein Kinase
Activity Produces Long-Period and Arrhythmic Circadian Behavioral and Molecular Rhythms.
Molecular and Cellular Biology. 2007;27(23):8049-64. doi: 10.1128/mcb.00680-07.
65.   Yao Z, Bennett AJ, Clem JL, Shafer OT. The Drosophila Clock Neuron Network Features
Diverse Coupling Modes and Requires Network-wide Coherence for Robust Circadian Rhythms.
Cell Rep. 2016;17(11):2873-81. Epub 2016/12/16. doi: 10.1016/j.celrep.2016.11.053. PubMed
PMID: 27974202; PubMed Central PMCID: PMCPMC5161247.
66.   Brown MP, Verma S, Palmer I, Zuniga AG, Rosensweig C, Keles MF, et al. A subclass of
evening cells promotes the switch from arousal to sleep at dusk. bioRxiv. 2023. Epub 2023/09/11.
doi: 10.1101/2023.08.28.555147.   PubMed   PMID: 37693540;   PubMed   Central   PMCID:
PMCPMC10491161.
67.   Delventhal R, O'Connor RM, Pantalia MM, Ulgherait M, Kim HX, Basturk MK, et al.
Dissection of central clock function in Drosophila through cell-specific CRISPR-mediated clock
gene disruption. Elife. 2019;8. Epub 2019/10/16. doi: 10.7554/eLife.48308. PubMed PMID:
31613218; PubMed Central PMCID: PMCPMC6794090.
68.   Schlichting M, Diaz MM, Xin J, Rosbash M. Neuron-specific knockouts indicate the
importance of   network communication   to Drosophila rhythmicity. Elife.   2019;8. Epub
2019/10/16. doi: 10.7554/eLife.48301. PubMed PMID: 31613223; PubMed Central PMCID:
PMCPMC6794074.
69.   Shansky RM, Murphy AZ. Considering sex as a biological variable will require a global shift
in science culture. Nature Neuroscience. 2021;24(4):457-64. doi: 10.1038/s41593-021-00806-8.
70.   Kokras N, Dalla C. Sex differences in animal models of psychiatric disorders. British Journal
of Pharmacology. 2014;171(20):4595-619. doi: https://doi.org/10.1111/bph.12710.
71.   Beery AK, Zucker I. Sex bias in neuroscience and biomedical research. Neurosci Biobehav
Rev.   2011;35(3):565-72.   Epub   2010/07/14.   doi: 10.1016/j.neubiorev.2010.07.002.   PubMed
PMID: 20620164; PubMed Central PMCID: PMCPMC3008499.
72.   Joye DAM, Evans JA. Sex differences in daily timekeeping and circadian clock circuits.
Semin Cell Dev Biol. 2022;126:45-55. Epub 20210514. doi: 10.1016/j.semcdb.2021.04.026.
PubMed PMID: 33994299; PubMed Central PMCID: PMCPMC8589873.
73.   Taghert PH, Nitabach MN. Peptide neuromodulation in invertebrate model systems.
Neuron.   2012;76(1):82-97.   Epub   2012/10/09.   doi: 10.1016/j.neuron.2012.08.035.   PubMed
PMID: 23040808; PubMed Central PMCID: PMCPMC3466441.
74.   Goda T, Umezaki Y, Alwattari F, Seo HW, Hamada FN. Neuropeptides PDF and DH31
hierarchically   regulate   free-running   rhythmicity   in   Drosophila   circadian   locomotor   activity.
Scientific Reports. 2019;9(1):838. doi: 10.1038/s41598-018-37107-3.
75.   Helfrich-Forster C. Robust circadian rhythmicity of Drosophila melanogaster requires the
presence of lateral neurons: a brain-behavioral study of disconnected mutants. J Comp Physiol A.
1998;182(4):435-53. Epub 1998/04/08. PubMed PMID: 9530835.
76.   Fernandez MP, Berni J, Ceriani MF. Circadian remodeling of neuronal circuits involved in
rhythmic   behavior.   PLoS   Biol.   2008;6(3):e69.   Epub   2008/03/28.   doi:

10.1371/journal.pbio.0060069. PubMed PMID: 18366255; PubMed Central PMCID:
PMCPMC2270325.

77. Biondi G, McCormick G, Fernandez MP. The Drosophila Circadian Clock Gene
Cycle Controls Development of Clock Neurons. bioRxiv. 2023:2023.10.29.564626.
doi: 10.1101/2023.10.29.564626.

78. Johard HAD, Yoishii T, Dircksen H, Cusumano P, Rouyer F, Helfrich-Förster C, et al.
Peptidergic clock neurons in Drosophila: Ion transport peptide and short neuropeptide F in
subsets of dorsal and ventral lateral neurons. The Journal of Comparative Neurology.
2009;516(1):59-73. doi: 10.1002/cne.22099.

79. Todd WD, Fenselau H, Wang JL, Zhang R, Machado NL, Venner A, et al. A hypothalamic
circuit for the circadian control of aggression. Nat Neurosci. 2018;21(5):717-24. Epub 20180409.
doi: 10.1038/s41593-018-0126-0. PubMed PMID: 29632359; PubMed Central PMCID:
PMCPMC5920747.

80. Fujii S, Krishnan P, Hardin P, Amrein H. Nocturnal male sex drive in Drosophila. Curr Biol.
2007;17(3):244-51. Epub 2007/02/06. doi: 10.1016/j.cub.2006.11.049. PubMed PMID:
17276917; PubMed Central PMCID: PMCPMC2239012.

81. Chowdhury B, Wang M, Gnerer JP, Dierick HA. The Divider Assay is a high-throughput
pipeline for aggression analysis in Drosophila. Commun Biol. 2021;4(1):85. Epub 2021/01/21. doi:
10.1038/s42003-020-01617-6. PubMed PMID: 33469118; PubMed Central PMCID:
PMCPMC7815768.

82. Sakai T, Ishida N. Circadian rhythms of female mating activity governed by clock genes in
Drosophila. Proc Natl Acad Sci U S A. 2001;98(16):9221-5. Epub 2001/07/27. doi:
10.1073/pnas.151443298. PubMed PMID: 11470898; PubMed Central PMCID: PMCPMC55401.

83. T M, Hari Dass S, Sharma VK. Egg-laying rhythm in Drosophila melanogaster. J Genet.
2008;87(5):495-504. Epub 2009/01/17. doi: 10.1007/s12041-008-0072-9. PubMed PMID:
19147938.

84. Zhang C, Daubnerova I, Jang YH, Kondo S, Žitňan D, Kim YJ. The neuropeptide allatostatin
C from clock-associated DN1p neurons generates the circadian rhythm for oogenesis. Proc Natl
Acad Sci U S A. 2021;118(4). doi: 10.1073/pnas.2016878118. PubMed PMID: 33479181; PubMed
Central PMCID: PMCPMC7848730.

85. Persons JL, Abhilash L, Lopatkin AJ, Roelofs A, Bell EV, Fernandez MP, et al. PHASE: An
Open-Source Program for the Analysis of DrosophilaPhase, Activity, and Sleep Under
Entrainment. J Biol Rhythms. 2022;37(4):455-67. Epub 2022/06/22. doi:
10.1177/07487304221093114. PubMed PMID: 35727044; PubMed Central PMCID:
PMCPMC10362883.

86. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
source platform for biological-image analysis. Nature Methods. 2012;9:676. doi:
10.1038/nmeth.2019
https://www.nature.com/articles/nmeth.2019#supplementary-information.

87. Lee Gierke C, Cornelissen G. Chronomics analysis toolkit (CATkit). Biol Rhythm Res.
2016;47(2):163-81. doi: 10.1080/09291016.2015.1094965.

A

Pdf > Cas9 + Pdf-g

Pdf-RFP, Pdf-Gal4; tub-Gal80ts (Control)

PDF | RFP

♂ *Pdf > Cas9 + Pdf-g* (experimental)

PDF | RFP

♀

PDF | RFP

B

♂

PDF intensity (a.u.)

***

| Pdf-Gal4 | + | + |
| UAS-Cas9+ UAS-Pdf-g | | + |

♀

PDF intensity (a.u.)

***

| Pdf-Gal4 | + | + |
| UAS-Cas9+ UAS-Pdf-g | | + |

C

Δ (PDF intensity)

♂   ♀

Pdf >Cas9 + Pdf-g

A

B

Pdf > Cas9 + Pdf-g (25°C)

Pdf > Cas9 + Pdf-g (adult specific)