Neurotrophic factor Neuritin modulates T cell electrical and metabolic state for the balance of tolerance and immunity

Hong Yu[1,2*#], Hiroshi Nishio[1, 2, 3, #], Joseph Barbi[1,2,4, #], Marisa Mitchell-Flack[1,2], Paolo D. A. Vignali[1,2,5], Ying Zheng[1, 2], Andriana Lebid[1,2], Kwang-Yu Chang[1,6], Juan Fu[1,2], Makenzie Higgins[1], Ching-Tai Huang[2, 7], Xuehong Zhang[8], Zhiguang Li[8], Lee Blosser[1], Ada Tam[1], Charles G. Drake[2,9], and Drew M. Pardoll[1,2]

[1] Bloomberg-Kimmel Institute for Cancer Immunotherapy, Immunology and Hematopoiesis Division, Department of Oncology, Johns Hopkins University School of Medicine, Baltimore, Maryland.
[2] The Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University School of Medicine, Baltimore, Maryland.
[3] Current address: Department of Obstetrics and Gynecology, Keio University School of Medicine, Tokyo 160-8582, Japan
[4] Current address: Department of Immunology, Roswell Park Comprehensive Cancer Center, Buffalo, NY14263, USA
[5] Current address: University of Pittsburgh, Carnegie Mellon.
[6] Current address: National Institute of Cancer Research, National Health Research Institutes, Tainan, Taiwan
[7] Current address: Infectious Diseases, Department of Medicine, Chang Gung Memorial Hospital, Taiwan
[8] Institute of Cancer Stem Cell, Cancer Center, Dalian Medical University, Dalian 116044, China.
[9] Current address: Division of Hematology and Oncology, Herbert Irving Comprehensive Cancer Center, Columbia University Medical Center, New York, New York 10032.
# These authors contributed equally to this work.
*Correspondence to hyu13@jhmi.edu.

Abstract:

The adaptive T cell response is accompanied by continuous rewiring of the T cell's electric and

metabolic state. Ion channels and nutrient transporters integrate bioelectric and biochemical

signals from the environment, setting cellular electric and metabolic states. Divergent electric and

metabolic states contribute to T cell immunity or tolerance. Here, we report that neuritin (Nrn1)

contributes to tolerance development by modulating regulatory and effector T cell function. Nrn1

expression in regulatory T cells promotes its expansion and suppression function, while expression

in the T effector cell dampens its inflammatory response. Nrn1 deficiency causes dysregulation

of ion channel and nutrient transporter expression in Treg and effector T cells, resulting in

divergent metabolic outcomes and impacting autoimmune disease progression and recovery.

These findings identify a novel immune function of the neurotrophic factor Nrn1 in regulating the

T cell metabolic state in a cell context-dependent manner and modulating the outcome of an

immune response.

Introduction

Peripheral T cell tolerance is important in restricting autoimmunity and minimizing collateral damage during active immune reactions and is achieved via diverse mechanisms, including T cell anergy, regulatory T (Treg) cell mediated suppression, and effector T (Te) cell exhaustion or deletion (ElTanbouly and Noelle, 2021). Upon activation, Treg and conventional T cells integrate environmental cues and adapt their metabolism to the energetic and biosynthetic demands, leading to tolerance or immunity. Tolerized versus responsive T cells are characterized by differential metabolic states. For example, T cell anergy is associated with reduced glycolysis, whereas activated T effector cells exhibit increased glycolysis (Buck et al., 2017; Geltink et al., 2018; Peng and Li, 2023; Zheng et al., 2009). Cellular metabolic states depend on electrolyte and nutrient uptake from the microenvironment (Chapman and Chi, 2022; Olenchock et al., 2017). Ion channels and nutrient transporters, which can integrate environmental nutrient changes, affect the cellular metabolic choices and impact the T cell functional outcome (Babst, 2020; Bohmwald et al., 2021; Ramirez et al., 2018). Each cell's functional state would correspond with a set of ion channels and nutrient transporters supporting their underlying metabolic requirements. The mechanisms coordinating the ion channel and nutrient transporter expression changes to support the adaptive T cell functional state in the immune response microenvironment remain unclear.

Nrn1, also known as candidate plasticity gene 15 (CPG15), was initially discovered as a neurotrophic factor linked to the neuronal cell membrane through a glycosylphosphatidylinositol (GPI) anchor (Nedivi et al., 1998; Zhou and Zhou, 2014). It is highly conserved across species, with 98% overall homology between the murine and human protein. Nrn1 plays multiple roles in neural development, synaptic plasticity, synaptic maturation, neuronal migration, and survival (Cantallops et al., 2000; Javaherian and Cline, 2005; Nedivi et al., 1998; Putz et al., 2005; Zito et al., 2014). In the immune system, Nrn1 expression has been found in Foxp3[+] Treg and follicular regulatory T cells (Tfr) (Gonzalez-Figueroa et al., 2021; Vahl et al., 2014), T cells from transplant tolerant recipients (Lim et al., 2013), anergized CD8 cells or CD8 cells from tumor-infiltrating lymphocytes in mouse tumor models (Schietinger et al., 2012; Schietinger et al., 2016; Singer et al., 2016), and in human Treg infiltrating breast cancer tumor tissue (Plitas et al., 2016). Soluble Nrn1 can be released from Tfr cells and act directly on B cells to suppress autoantibody development against tissue-specific antigens (Gonzalez-Figueroa et al., 2021). Despite the observation of Nrn1 expression in Treg cells and T cells from tolerant environments (Gonzalez-

Figueroa et al., 2021; Lim et al., 2013; Plitas et al., 2016; Schietinger et al., 2012; Schietinger et
al., 2016; Singer et al., 2016), the roles of Nrn1 in T cell tolerance development and Treg cell
function have not been explored, and functional mechanism of Nrn1 remains elusive. This study
demonstrates that the neurotrophic factor Nrn1 can moderate T cell tolerance and immunity
through both Treg and Te cells, impacting Treg cell expansion and suppression while controlling
inflammatory response in Te cells.

Results

Nrn1 expression and function in T cell anergy.

To explore the molecular mechanisms underlying peripheral tolerance development, we
utilized a system we previously developed to identify tolerance-associated genes (Huang et al.,
2004). We compared the gene expression patterns associated with either a T effector/memory
response or tolerance induction triggered by the same antigen but under divergent *in vivo*
conditions (Huang et al., 2004). Influenza hemagglutinin (HA) antigen-specific TCR transgenic
CD4 T cells were adoptively transferred into WT recipients with subsequent HA-Vaccinia virus
(VacHA) infection to generate T effector/memory cells while tolerogenic HA-specific CD4s were
generated by transfer into hosts with transgenic expression of HA as self-antigen (C3-HA mice,
Figure 1A.)(Huang et al., 2004). One of the most differentially expressed genes upregulated in the
anergy-inducing condition was Nrn1. Nrn1 expression was significantly higher among cells
recovered from C3-HA hosts vs. cells from VacHA infected mice at all time points tested by qRT-
PCR (Figure 1A). To further confirm the association of Nrn1 expression with T cell anergy, we
assessed Nrn1 expression in naturally occurring anergic polyclonal CD4$^+$ T cells (Ta), which can
be identified by surface co-expression of Folate Receptor 4 (FR4) and the ecto-5'-nucleotidase
CD73 (Ta, CD4$^+$CD44$^+$FR4^{hi}CD73^{hi} cells)(Kalekar et al., 2016). Nrn1 expression was
significantly higher in Ta than in naïve CD4 (Tn, CD4$^+$CD62L$^+$CD44$^-$FR4$^-$CD73$^-$) and antigen-
experienced cells (Te, CD4$^+$CD44$^+$FR4$^-$CD73$^-$) under steady-state conditions measured by both
qRT-PCR and western blot (Figure 1B). Given that Treg cells, like anergic cells, have roles in
maintaining immune tolerance, we queried whether Nrn1 is also expressed in Treg cells. Nrn1
expression can be detected in nTreg and induced Treg (iTreg) cells generated *in vitro* (Figure 1C).

To evaluate Nrn1 expression under pathological tolerant conditions (Cuenca et al., 2003),
we evaluated Nrn1 expression in T cells within the tumor microenvironment. Nrn1 expression in

murine Treg cells and non-Treg CD4$^+$ cells from tumor infiltrates were compared to the Treg cells

and non-Treg CD4$^+$ T cells isolated from peripheral blood. Nrn1 mRNA level was significantly

increased among tumor-associated Treg cells and non-Treg CD4 cells compared to cells from

peripheral blood (Figure 1-figure supplement 1A). Consistent with our findings in the mouse tumor

setting, the Treg and non-Treg T cells from human breast cancer infiltrates reveal significantly

higher Nrn1 expression compared to the peripheral blood Treg and non-Treg cells (Figure 1-figure

supplement 1B) (Plitas et al., 2016).

CD4$^+$ T cells may pass through an effector stage after activation before reaching an anergic

state (Adler et al., 1998; Chen et al., 2004; Huang et al., 2003; Opejin et al., 2020). To evaluate

the potential role of Nrn1 expression in T cell tolerance development, we further examined Nrn1

expression kinetics after T cell activation. Nrn1 expression was significantly induced after CD4$^+$

T cell activation (Figure 1D). Using an Nrn1-specific, monoclonal antibody, Nrn1 can be detected

on activated CD4$^+$ and CD8$^+$ cells (Figure 1D, Figure 1-figure supplement 1C). The significant

enhancement of Nrn1 expression after T cell activation suggests that Nrn1 may contribute to the

process of T cell tolerance development and/or maintenance. Although Treg cells express Nrn1,

we were not able to consistently detect substantial cell surface Nrn1 expression (Figure 1D, Figure

1-figure supplement 1D), likely due to Nrn1 being produced in a soluble form or cleaved from the

cell membrane (Gonzalez-Figueroa et al., 2021).

To understand the functional implication of Nrn1 expression in immune tolerance, we

analyzed Nrn1-deficient (Nrn1$^{-/-}$) mice (Fujino et al., 2011). In the first evaluation of the Nrn1$^{-/-}$

colony, Nrn1$^{-/-}$ mice had consistently reduced body weight compared to heterozygous Nrn1$^{+/-}$ and

WT (Nrn1$^{+/+}$) mice (Figure 1-figure supplement 2A). The lymphoid tissues of Nrn1$^{-/-}$ mice were

comparable to their Nrn1$^{+/-}$ and WT counterparts except for a slight reduction in cell number that

was observed in the spleens of Nrn1$^{-/-}$ mice, likely due to their smaller size (Figure 1-figure

supplement 2B). Analysis of thymocytes revealed no defect in T cell development (Figure 1-figure

supplement 2C), and a flow cytometric survey of the major immune cell populations in the

peripheral lymphoid tissue of these mice revealed similar proportions of CD4, CD8 T cells, B cells,

monocytes and dendritic cells (DCs) (Figure 1-figure supplement 2D). Similarly, no differences

were found between the proportions of anergic and Treg cells in Nrn1$^{-/-}$, Nrn1$^{+/-}$ and WT mice

(Figure 1-figure supplement 2E, F), suggesting that Nrn1 deficiency does not significantly affect

anergic and Treg cell balance under steady state. Additionally, histopathology assessment of lung,

heart, liver, kidney, intestine and spleen harvested from 13 months old Nrn1$^{-/-}$ and Nrn1$^{+/-}$ littermates did not reveal any evidence of autoimmunity (data not shown). The comparable level of anergic and Treg cell population among Nrn1$^{-/-}$, Nrn1$^{+/-}$ and WT mice and lack of autoimmunity in Nrn1$^{-/-}$ aged mice suggest that Nrn1 deficiency is not associated with baseline immune abnormalities or overt dysfunction. Due to the similarity between Nrn1$^{+/-}$ and WT mice, we have used either Nrn1$^{+/-}$ or WT mice as our control depending on mice availability and referred to both as "ctrl" in the subsequent discussion.

To evaluate the relevance of Nrn1 in CD4^{+} T cell tolerance development, we employed the classic peptide-induced T cell anergy model (Vanasek et al., 2006). Specifically, we crossed OVA antigen-specific TCR transgenic OTII mice onto the Nrn1$^{-/-}$ background. Nrn1$^{-/-}$_OTII^{+} or control_OTII^{+} (ctrl_OTII^{+}) cells marked with Thy1.1^{+} congenic marker (Thy1.1^{+}Thy1.2^{-}), were co-transferred with polyclonal WT Tregs (marked as Thy1.1^{-}thy1.2^{+}), into TCRα knockout mice (TCRα$^{-/-}$), followed by injection of soluble OVA peptide to induce clonal anergy (Figure 1E) (Chappert and Schwartz, 2010; Martinez et al., 2012; Mercadante and Lorenz, 2016; Shin et al., 2014). On day 13 after cell transfer, the proportion and number of OTII cells increased in the Nrn1$^{-/-}$_OTII compared to the ctrl_OTII hosts (Figure 1F). Moreover, Nrn1$^{-/-}$_OTII cells produced increased IL2 than ctrl_OTII upon restimulation (Figure 1G). Anergic CD4 Tconv cells can transdifferentiate into Foxp3^{+} pTreg cells *in vivo* (DL, 2017; Kalekar et al., 2016; Kuczma et al., 2021). Consistent with reduced anergy induction, the proportion of Foxp3^{+} pTreg among Nrn1$^{-/-}$_OTII was significantly reduced (Figure 1H). In parallel with the phenotypic analysis, we compared gene expression between Nrn1$^{-/-}$_OTII and ctrl_OTII cells by RNA Sequencing (RNASeq). Gene set enrichment analysis (GSEA) revealed that the gene set on T cell anergy was enriched in ctrl relative to Nrn1$^{-/-}$_OTII cells (Figure 1I)(Safford et al., 2005). Also, consistent with the decreased transdifferentiation to Foxp3^{+} cells, the Treg signature gene set was prominently reduced in Nrn1$^{-/-}$_OTII cells relative to the ctrl (Figure 1J). Anergic T cells are characterized by inhibition of proliferation and compromised effector cytokines such as IL2 production (Choi and Schwartz, 2007). The increased cell expansion and cytokine production in Nrn1$^{-/-}$_OTII cells and the reduced expression of anergic and Treg signature genes all support the notion that Nrn1 is involved in T cell anergy development.

Anergic T cells are developed after encountering antigen, passing through a brief effector stage, and reaching an anergic state (Chappert and Schwartz, 2010; Huang et al., 2003; Silva

Morales and Mueller, 2018; Zha et al., 2006). Enhanced T cell activation, defective Treg cell

conversion or expansion, and heightened T effector cell response may all contribute to defects in

T cell anergy induction and/or maintenance (Chappert and Schwartz, 2010; Huang et al., 2003;

Kalekar et al., 2016; Silva Morales and Mueller, 2018; Zha et al., 2006). We first examined early

T cell activation to understand the underlying cause of defective anergy development in Nrn1$^{-/-}$

cells. Nrn1$^{-/-}$ CD4$^+$ cells showed reduced T cell activation, as evidenced by reduced CellTrace

violet dye (CTV) dilution, activation marker expression, and Ca^{++} entry after TCR stimulation

(Figure 1-figure supplement 3A, B, C). The reduced early T cell activation observed in Nrn1$^{-/-}$

CD4 cells suggests that the compromised anergy development in Nrn1$^{-/-}$_OTII cells was not caused

by enhanced early T cell activation. The defective pTreg generation and/or enhanced effector T

cell response may contribute to compromised anergy development.

Compromised Treg expansion and suppression in the absence of Nrn1.

The significant reduction of Foxp3$^+$ pTreg among Nrn1$^{-/-}$_OTII cells could be caused by

the diminished conversion of Foxp3$^-$ Tconv cells to pTreg and/or diminished Treg cell expansion

and persistence. To understand the cause of pTreg reduction in Nrn1$^{-/-}$_OTII cells (Figure 1H), we

turned to the induced Treg (iTreg) differentiation system to evaluate the capability of Foxp3$^+$ Treg

development and expansion in Nrn1$^{-/-}$ cells. Similar proportions of Foxp3$^+$ cells were observed in

Nrn1$^{-/-}$ and ctrl cells under the iTreg culture condition (Figure 2A), suggesting that Nrn1 deficiency

does not significantly impact Foxp3$^+$ cell differentiation. To examine the capacity of iTreg

expansion, Nrn1$^{-/-}$ and ctrl iTreg cells were restimulated with anti-CD3, and we found reduced live

cells over time in Nrn1$^{-/-}$ iTreg compared to the ctrl (Figure 2B). The reduced live cell number in

Nrn1$^{-/-}$ was accompanied by reduced Ki67 expression (Figure 2C). Although Nrn1$^{-/-}$ iTregs

retained a higher proportion of Foxp3$^+$ cells three days after restimulation, however, when taking

into account the total number of live cells, the actual number of live Foxp3$^+$ cells was reduced in

Nrn1$^{-/-}$ (Figure 2D). Treg cells are not stable and are prone to losing Foxp3 expression after

extended proliferation (Feng et al., 2014; Floess et al., 2007; Li et al., 2014; Zheng et al., 2010).

The increased proportion of Foxp3$^+$ cells was consistent with reduced proliferation observed in

Nrn1$^{-/-}$ cells. Thus, Nrn1 deficiency can lead to reduced iTreg cell proliferation and persistence *in*

vitro.

The defects observed in iTreg cell expansion *in vitro* prompt further examination of Nrn1$^-$

$^{-/-}$ nTreg expansion and suppression function *in vivo*. To this end, we tested the suppression capacity of congenically marked (CD45.1$^-$CD45.2$^+$) Nrn1$^{-/-}$ or ctrl nTreg toward CD45.1$^+$CD45.2$^-$ responder cells in Rag2$^{-/-}$ mice (Figure 2E). The CD45.1$^+$CD45.2$^-$ responder cells devoid of Treg cells were splenocytes derived from Foxp3DTRGFP (FDG) mice pretreated with diphtheria toxin (DT) (Kim et al., 2007; Workman et al., 2011). DT treatment caused the deletion of Treg cells in FDG mice (Kim et al., 2007). Although the CD45.1$^-$CD45.2$^+$ Nrn1$^{-/-}$ and ctrl cell proportions were not significantly different among hosts splenocytes at day 7 post transfer (Figure 2F), Nrn1$^{-/-}$ cells retained a higher Foxp3$^+$ cell proportion and reduced Ki67 expression comparing to the ctrl (Figure 2G, H). These findings were similar to our observation of iTreg cells *in vitro* (Figure 2C, D). Nrn1$^{-/-}$ Tregs also showed reduced suppression toward CD45.1$^+$ responder cells, evidenced by increased CD45.1$^+$ proportion and cell number in host splenocytes (Figure 2I).

To evaluate the functional implication of Nrn1$^{-/-}$ Treg suppression in disease settings, we challenged the Rag2$^{-/-}$ hosts with the poorly immunogenic B16F10 tumor (Figure 2E). Tumors grew much slower in Nrn1$^{-/-}$ Treg recipients than those reconstituted with ctrl Tregs (Figure 2J). Moreover, the number of CD45.1$^+$ cells in tumor-draining lymph nodes and spleens increased significantly in Nrn1$^{-/-}$ Treg hosts compared to the ctrl group (Figure 2K). Consistently, the CD45.1$^+$ responder cell proportion among tumor lymphocyte infiltrates (TILs) was also increased (Figure 2L), accompanied by an increased proportion of IFNγ$^+$ cells among CD8 TILs from Nrn1$^{-/-}$ Treg hosts (Figure 2M). The increased expansion of CD45.1$^+$ responder cells and reduced tumor growth further confirmed the reduced suppressive capacity of Nrn1$^{-/-}$ Treg cells.

Nrn1 impacts Treg cell electrical and metabolic state

To understand the molecular mechanisms associated with Nrn1$^{-/-}$ Treg cells, we compared gene expression between Nrn1$^{-/-}$ and ctrl iTregs under resting (IL2 only) and activation (aCD3 and IL2) conditions by RNASeq. GSEA on gene ontology database and clustering of enriched gene sets by Cytoscape identified three clusters enriched in resting Nrn1$^{-/-}$ iTreg (Figure 3A, Figure 3-figure supplement Table 1)(Shannon et al., 2003; Subramanian et al., 2005). The "neurotransmitter involved in membrane potential (MP)" and "sodium transport" clusters involved gene sets on the ion transport and cell MP regulation (Figure 3A, Figure 3-figure supplement Table 1). MP is the difference in electric charge between the interior and the exterior of the cell membrane (Abdul Kadir et al., 2018; Blackiston et al., 2009; Ma et al., 2017). Ion channels and transporters for Na$^+$

and other ions such as K^+, Cl^- *et al.* maintain the ion balance and contribute to cell MP (Blackiston

et al., 2009). MP change can impact cell plasma membrane lipid dynamics and affect receptor

kinase activity (Zhou et al., 2015). The enrichment of "receptor protein kinase" gene set clusters

may reflect changes caused by MP (Figure 3A, Figure 3-figure supplement Table 1). Gene set

cluster analysis on activated iTreg cells also revealed the enrichment of the "ion channel and

receptor" cluster in Nrn1$^{-/-}$ cells (Figure 3B, Figure 3-figure supplement table 2), supporting the

potential role of Nrn1 in modulating ion balances and MP.

The "Neurotransmitter receptor activity involved in regulation of postsynaptic membrane

potential" gene set was significantly enriched under resting and activation conditions (Figure 3C

and D; Figure 3-figure supplement Table 3). The α-amino-3-hydroxy-5-methyl-4-

isoxazolepropionic acid receptor (AMPAR) subunits Gria2 and Gria3 are the major components

of this gene set and showed increased expression in Nrn1$^{-/-}$ cells (Figure 3D). AMPAR is an

ionotropic glutamate receptor that mediates fast excitatory synaptic transmission in neurons. Nrn1

has been reported as an accessory protein for AMPAR (Pandya et al., 2018; Schwenk et al., 2012;

Subramanian et al., 2019), although the functional implication of Nrn1 as an AMPAR accessory

protein remains unclear. The enrichment of MP related gene set prompted the examination of

electric status, including MP level and ion channel expressions. We examined the relative MP level

by FLIPR MP dye, a lipophilic dye able to cross the plasma membrane, which has been routinely

used to measure cell MP changes (Dvorak et al., 2021; Joesch et al., 2008; Nik et al., 2017;

Whiteaker et al., 2001). When the cells are depolarized, the dye enters the cells, causing an increase

in fluorescence signal. Conversely, cellular hyperpolarization results in dye exit and decreased

fluorescence. Compared to ctrl iTreg cells, Nrn1$^{-/-}$ exhibits significant hyperpolarization under

both resting and activation conditions (Figure 3E). Consistent with the MP change, the "MF_metal

ion transmembrane transporter activity" gene set, which contains 436 ion channel related genes,

was significantly enriched and showed a different expression pattern in Nrn1$^{-/-}$ iTregs (Figure 3F;

Figure 3-figure supplement 1A and B). The changes in cellular MP and differential expression of

ion channel and transporter genes in Nrn1$^{-/-}$ implicate the role of Nrn1 in the balance of electric

state in the iTreg cell.

MP changes have been associated with changes in amino acid (AA) transporter expression

and nutrient acquisition, which in turn influences cellular metabolic and functional state (Yu et al.,

2022). To understand whether MP changes in Nrn1$^{-/-}$ are associated with changes in nutrient

acquisition and thus the metabolic state, we surveyed AA transport-related gene expression using
the "Amino acid transmembrane transporter activity" gene set and found differential AA
transporter gene expression between Nrn1$^{-/-}$ and ctrl iTregs (Figure 3G). Upon AA entry through
transporters, the electric charge carried by these molecules may transiently affect cell membrane
potential. Differential AA transporter expression patterns may have different impacts on cellular
MP upon AA entry. Thus, we loaded Nrn1$^{-/-}$ and ctrl iTreg with FLIPR MP dye in the HBSS
medium and tested cellular MP change upon exposure to MEM AAs. The AA-induced cellular
MP change was reduced in Nrn1$^{-/-}$ compared to the ctrl, reflective of differential AA transporter
expression patterns (Figure 3H). Electrolytes and AAs entry are critical regulators of mTORC1
activation and T cell metabolism (Liu and Sabatini, 2020; Saravia et al., 2020; Sinclair et al., 2013;
Wang et al., 2020). We examined mTORC1 activation at the protein level by evaluating mTOR
and S6 phosphorylation via flow cytometry. We found reduced phosphorylation of mTOR and S6
in activated Nrn1$^{-/-}$ iTreg cells (Figure 3-figure supplement 1C). We further performed a nutrient-
sensing assay to evaluate the role of ion and nutrient entry in mTORC1 activation. Nrn1$^{-/-}$ and ctrl
iTreg cells were starved for one hour in a nutrient-free buffer, followed by adding RPMI medium
with complete ions and nutrients, and cultured for two more hours. While adding the medium with
nutrients clearly increased the mTOR and S6 phosphorylation, the degree of change was
significantly less in Nrn1$^{-/-}$ than in the ctrl (Figure 3I). Consistently, GSEA on Hallmark gene sets
reveal reduced gene set enrichment relating to the mTORC1 signaling, corroborating the reduced
pmTOR and pS6 detection in Nrn1$^{-/-}$ cells. Moreover, Nrn1$^{-/-}$ cells also showed reduced expression
of glycolysis, fatty acid metabolism, and oxidative phosphorylation related gene sets under both
resting and activating conditions (Figure 3J, Figure 3-figure supplement 1D), indicating changes
in metabolic status. Since previous work has identified mTORC1 to be an important regulator of
aerobic glycolysis and given that our GSEA data suggested changes in glycolysis (Figure 3J)
(Salmond, 2018), we performed the seahorse assay and confirmed reduced glycolysis among Nrn1^{-}
$^{/-}$ cells (Figure 3K). Examination of mitochondrial bioenergetic function revealed a similar oxygen
consumption rate (OCR) between Nrn1$^{-/-}$ and ctrl cells (Figure 3K). Thus, Nrn1 expression can
affect the iTreg electric state, influence ion channel and nutrient transporter expression, impact
nutrient sensing, modulate metabolic state, and contribute to Treg expansion and suppression
function.
We have observed significant changes in the electrical and metabolic state among Nrn1$^{-/-}$

iTreg compared to the ctrl. Because Nrn1 can be expressed on the cell surface, one question arises

whether the changes observed in Nrn1$^{-/-}$ cells were caused by the functional deficiency of Nrn1 or

arose secondary to potential changes in cell membrane structure originating at the Nrn1$^{-/-}$ naïve T

cell stage. To answer this question, we first examined potential changes in electrical and metabolic

status among Nrn1$^{-/-}$ naïve CD4 T cells. The Nrn1$^{-/-}$ naïve CD4 T cells showed similar resting MP

and AA-induced MP changes compared to the ctrl cells (Figure 3-supplement figure 2 A, B). We

also observed comparable glycolysis and mitochondrial bioenergetic function between Nrn1$^{-/-}$

naïve CD4 T cells and the ctrl (Figure 3-supplement figure 2 C). These results suggest the electrical

and metabolic state in Nrn1$^{-/-}$ T cells are comparable to the ctrl cells at the naïve cell stage. To

further rule out the possibility that the observed changes in Nrn1$^{-/-}$ iTreg are secondary to

developmental structural changes, not Nrn1 functional deficiency, we differentiated WT T cells in

the presence of antagonistic Nrn1 antibody and compared to the WT ctrl and Nrn1$^{-/-}$ iTreg cells.

WT iTreg cells differentiated in the presence of Nrn1 antibody exhibit reduced resting MP, similar

to Nrn1$^{-/-}$ cells (Figure 3-supplement figure 2 D). Moreover, upon restimulation, WT iTreg cells

differentiated under Nrn1 antibody blockade showed a similar phenotype as Nrn1$^{-/-}$ cells, with

reduced live cell number, reduced Ki67 expression, and increased Foxp3^{+} cell proportion among

live cells (Figure 3-supplement figure 2 E, F). These results suggest that Nrn1 functional

deficiency likely contributes to the electrical and metabolic state change observed in Nrn1$^{-/-}$ iTreg

cells.

Nrn1 impact effector T cell inflammatory response.

CD4^{+} T cells can pass through an effector stage on their way to an anergic state (Huang et

al., 2003). Since Nrn1 expression is significantly induced after T cell activation (Figure 1D), Nrn1

might influence CD4^{+} effector (Te) cell differentiation, affecting anergy development. Nrn1 may

exert different electric changes due to distinct ion channel expression contexts in Te cells than in

Tregs. We first evaluated Nrn1$^{-/-}$ Te cell differentiation *in vitro*. Nrn1 deficient CD4 Te cells

showed increased Ki67 expression, associated with increased cytokine TNFα, IL2, and IFNγ

expression upon restimulation (Figure 4A). To evaluate Nrn1$^{-/-}$ Te cell response *in vivo*, we crossed

Nrn1$^{-/-}$ with FDG mice and generated Nrn1$^{-/-}$_FDG and ctrl_FDG mice, which enabled the

elimination of endogenous Treg cells (Figure 4B). Deleting endogenous Foxp3^{+} Treg cells using

DT will cause the activation of self-reactive T cells, leading to an autoimmune response (Kim et

al., 2007; Nystrom et al., 2014). Upon administration of DT, we observed accelerated weight loss in Nrn1$^{-/-}$_FDG mice, reflecting enhanced autoimmune inflammation (Figure 4C). Examination of T cell response revealed a significant increase in Ki67 expression and inflammatory cytokine TNFα, IL2, and IFNγ expression among Nrn1$^{-/-}$ CD4 cells on day 6 post DT treatment (Figure 4D), consistent with the findings *in vitro*. The proportion of Foxp3^{+} cells was very low on day 6 post DT treatment and comparable between Nrn1$^{-/-}$ and the ctrl (Figure 4E), suggesting that the differential Te cell response was not due to the impact from Treg cells. Thus, Nrn1 deficiency enhances Te cell response *in vitro* and *in vivo*.

To identify molecular changes responsible for Nrn1$^{-/-}$ Te phenotype, we compared gene expression between Nrn1$^{-/-}$ and ctrl Te cells by RNASeq. GSEA and Cytoscape analysis identified a cluster of gene sets on "membrane repolarization", suggesting that Nrn1 may also be involved in the regulation of MP under Te context (Figure 4F, Figure 4-figure supplement Table 4) (Shannon et al., 2003; Subramanian et al., 2005). While the "membrane_repolarization" gene set was enriched in Nrn1$^{-/-}$ (Figure 4G), the "Neurotransmitter receptor activity involved in regulation of postsynaptic membrane potential" gene set was no longer enriched, but the AMPAR subunit Gria3 expression was still elevated in Nrn1$^{-/-}$ Te cells (Figure 4-figure supplement 1A). Although MP in Te cells was comparable between Nrn1$^{-/-}$ and ctrl (Figure 4H), the "MF_metal ion transmembrane transporter activity" gene set was significantly enriched in Nrn1$^{-/-}$ with different gene expression patterns (Figure 4I, Figure 4-figure supplement 1B), indicative of different electric state. The significant enrichment of ion channel related genes in Nrn1$^{-/-}$ Te cells was in line with the finding in Nrn1$^{-/-}$ iTreg cells, supporting the notion that Nrn1 expression may be involved in ion balance and MP modulation.

Examination of nutrient transporters revealed that the "Amino acid transmembrane transporter activity" gene set was significantly enriched in Nrn1$^{-/-}$ cells than the ctrl (Figure 4J). We further examined AA entry-induced cellular MP change in Nrn1$^{-/-}$ and ctrl Te cells. AA entry caused enhanced MP change among Nrn1$^{-/-}$ Te than the ctrl, in contrast with the finding under iTreg cell context (Figure 4K). Along with the enrichment of ion channel and nutrient transporter genes (Figure 4I and J), we found enhanced mTOR and S6 phosphorylation in Nrn1$^{-/-}$ Te cells (Figure 4-supplemental figure 1C). We also compared nutrient sensing capability between Nrn1$^{-/-}$ and ctrl Te cells, as outlined in Figure 3I. Nrn1$^{-/-}$ Te showed increased mTOR and S6 phosphorylation after sensing ions and nutrients in RPMI medium (Figure 4L), confirming the

differential impact of ions and nutrients on Nrn1$^{-/-}$ and ctrl Te cells. GSEA on Hallmark collection
showed enrichment of mTORC1 signaling gene set (Figure 4M), corroborating with increased
pmTOR and pS6 detection in Nrn1$^{-/-}$ Te cells. Along with increased mTORC1 signaling, Nrn1$^{-/-}$
Te cells also showed enrichment of gene sets on glycolysis and proliferation (Figure 4M).
Evaluation of metabolic changes by seahorse confirmed increased glycolysis in Nrn1$^{-/-}$ cells, while
the OCR remained comparable between Nrn1$^{-/-}$ and ctrl (Figure 4N). These *in vitro* studies on Te
cells indicate that Nrn1 deficiency resulted in the dysregulation of the electrolyte and nutrient
transport program, impacting Te cell nutrient sensing, metabolic state, and the outcome of
inflammatory response.

Nrn1 deficiency exacerbates autoimmune disease

The coordinated reaction of Treg and Te cells contributes to the outcome of the immune
response. We employed the experimental autoimmune encephalomyelitis (EAE), the murine
model of multiple sclerosis (MS) to evaluate the overall impact of Nrn1 on autoimmune disease
development. Upon EAE induction, the incidence and time to EAE onset in Nrn1$^{-/-}$ mice were
comparable to the ctrl mice, but the severity, disease persistence, and body weight loss were
increased in Nrn1$^{-/-}$ mice (Figure 5A). Exacerbated EAE was associated with significantly
increased CD45^{+} cell infiltrates, increased CD4^{+} cell number, increased proportion of MOG-
specific CD4 cells, and reduced proportion of Foxp3^{+} CD4 cells in the Nrn1$^{-/-}$ spinal cord (Figure
5B-E). Moreover, we also observed increased proportions of IFNγ^{+} and IL17^{+} CD4 cells in Nrn1$^{-/-}$
mice remaining in the draining lymph node compared to the ctrl mice (Figure 5F). Thus, the
results from EAE corroborated with earlier data and confirmed the important role of Nrn1 in
establishing immune tolerance and modulating autoimmunity.

Discussion

T cell expansion and functional development depend on adaptive electric and metabolic
changes, maintaining electrolyte balances, and appropriate nutrient uptake. The negative charge of
the plasma membrane, ion channel expression pattern, and function are key characteristics
associated with the cellular electric state in different systems, impacting cell proliferation and
function (Blackiston et al., 2009; Emmons-Bell and Hariharan, 2021; Kiefer et al., 1980; Monroe
and Cambier, 1983; Sundelacruz et al., 2009). The electrolytes and nutrients, including amino

acids, metabolites, and small peptides transported through ion channels and nutrient transporters,

are also regulators and signaling agents impacting the choice of cellular metabolic pathways and

functional outcomes (Hamill et al., 2020). In this study, we report that the neurotropic factor Nrn1

expression influences CD4 T cell MP, ion channels, and nutrient transporter expression patterns,

contributing to differential metabolic states in Treg and Te cells. Nrn1 deficiency compromises

Treg cell expansion and suppression while enhancing Te cell inflammatory response, exacerbating

autoimmune disease.

Bioelectric controls have been defined as a type of epigenetics that can control information

residing outside of genomic sequence (Levin, 2021). The sum of ion channels and pump activity

generates the ionic gradient across the cell membrane, establishing the MP level and bioelectric

state. Cells with the same MP can have different ion compositions, and the same ion channel may

have a differential impact on MP when in combination with different ion channels (Abdul Kadir

et al., 2018). Consistent with this notion, Nrn1 deficiency has differential impacts on the cellular

electric state under the Treg and Te cells with different ion channel combinations. Altered MP was

detected in Nrn1 deficient Treg cells (Figure 3E), while comparable MP was observed between

Nrn1$^{-/-}$ and ctrl Te cells (Figure 4H). The MP level determined by ion channels and pump activity

can influence the nutrient transport pattern, establishing a metabolic and functional state matching

the MP level (Blackiston et al., 2009; Emmons-Bell and Hariharan, 2021; Kiefer et al., 1980;

Monroe and Cambier, 1983; Sundelacruz et al., 2009; Yu et al., 2022). Yu et al. reported that

macrophage MP modulates plasma membrane phospholipid dynamics and facilitates cell surface

retention of nutrient transporters, thus supporting nutrient uptake and impacting the inflammatory

response (Yu et al., 2022). Nutrient transport is key to T cell fate decisions and has been considered

signal 4 to T cell fate choices (Chapman and Chi, 2022; Long et al., 2021). The changes in ion

channel related gene expression and MP level in Nrn1$^{-/-}$ cells were accompanied by differential

expression of AA transporter genes and nutrient sensing activity that impacted mTORC1 pathway

activation and cellular glycolytic state (Figures 3 and 4). These results corroborate previous

observations on the connection of MP in nutrient acquisition and metabolic change and support

the role of Nrn1 in coordinating T cell electric and metabolic adaptation (Yu et al., 2022).

Although Nrn1, as a small GPI-anchored protein, does not have channel activity by itself,

it has been identified as one of the components in the AMPAR complex (Pandya et al., 2018;

Schwenk et al., 2012; Subramanian et al., 2019). Na$^+$-influx through the AMPA type ionotropic

glutamate receptor can quickly depolarize the postsynaptic compartment and potentiate synaptic

transmission in neurons. We have observed increased expression of AMPAR subunits in Nrn1$^{-/-}$

iTreg and Te cells (Figure 3D, Figure 4-figure supplement 1), implicating potential change in

AMPAR activity in Nrn1$^{-/-}$ under Treg and Te cell context. Glutamate secreted by proliferating

cells may influence T cell function through AMPAR. High glutamate levels are detected at the

autoimmune disease site and tumor interstitial fluid (Bonnet et al., 2020; McNearney et al., 2004;

Sullivan et al., 2019). Moreover, AMPAR has been implicated in exacerbating autoimmune

disease (Bonnet et al., 2015; Sarchielli et al., 2007). The increased expression of AMPAR subunits

in Nrn1$^{-/-}$ cells supports the potential connection of Nrn1 and AMPAR and warrants future

investigation on the possibility that Nrn1 functions through AMPAR, impacting T cell electric

change. Besides AMPAR, Nrn1 has been reported to function through the insulin receptor and

fibroblast growth factor pathway (Shimada et al., 2016; Yao et al., 2012). Subramanian et al have

suggested that rather than a traditional ligand with its cognate receptor, Nrn1 may function as an

adaptor to receptors to perform diverse cell-type-specific functions (Subramanian et al., 2019).

Our results do not rule out these possibilities.

Overall, we found that Nrn1 expression in Treg and Te cells can impact cellular electric

state, nutrient sensing, and metabolism in a cell context dependent manner. The predominant

enrichment of ion channel related gene sets in both Treg and Te cell context underscores the

importance of Nrn1 in modulating ion balance and MP. The changes in ion channels and nutrient

transporter expression in Treg and Te cells and associated functional consequences highlight the

importance of Nrn1 in coordinating cell metabolic changes through channels and transporters

during the adaptive response and contribute to the balance of tolerance and immunity.

Materials and Methods

Mouse models.

The Nrn1$^{-/-}$ mice (Fujino et al., 2011), Foxp3DTRGFP (FDG)(Kim et al., 2007), and TCRα$^{-/-}$ mice

were obtained from the Jackson Laboratory. OTII mice on Thy1.1^{+} background was kindly

provided by Dr. Jonathan Powell. Rag2$^{-/-}$ mice were maintained in our mouse facility. 6.5 TCR

transgenic mice specific for HA antigen and C3HA mice (both on the B10.D2 background) have

been described previously (Huang et al., 2004). Nrn1$^{-/-}$ mice were crossed with OTII mice to

generate Nrn1$^{-/-}$_OTII^{+} mice, ctrl_OTII^{+} mice. Nrn1$^{-/-}$ mice were also crossed with FDG mice to

generate Nrn1$^{-/-}$_FDG and ctrl_FDG mice. All mice colonies were maintained in accordance with

the guidelines of Johns Hopkins University and the institutional animal care and use committee

Antibodies and Reagents.

We have used the following antibodies: Anti-CD3 (17A2), anti-CD4 (RM4-5), anti-CD8a (53-6.7),

anti-CD25 (PC61), anti-CD45.1 (A20), anti-CD45.2 (104), anti-CD62L (MEL-14), CD73

(TY/111.8), anti-CD90.1 (OX-7), anti-CD90.2 (30-H12), anti-TCR V□□5.1, 5.2 (MR9-4), anti-

PD1 (29F.1A12), anti-IFNγ (XMG1.2), anti-IL17α (TC11-18H10.1), anti-TNFα (MP6-XT22),

anti-Tbet (4B10), anti-Ki67 (16A8) were purchased from Biolegend. Anti-CD44 (IM7), CD45

(30-F11), anti-CD69(H1.2F3) were purchased from BD Bioscience. Anti-FOXP3 (FJK-16s) was

purchased from eBioscience. The flow cytometry data were collected using BD Celesta (BD

Biosciences) or Attune Flow Cytometers (ThermoFisher). Data were analyzed using FlowJo (Tree

Star) software.

Mouse monoclonal anti-Nrn1 antibody (Ab) against Nrn1 was custom-made (A&G

Pharmaceutical). The specificity of anti-Nrn1 Ab was confirmed by ELISA, cell surface staining

of Nrn1 transfected 293T cells, and western blot of Nrn1 recombinant protein and brain protein

lysate from WT mice or Nrn1$^{-/-}$ mice (data not shown). OVA$_{323-339}$ peptide and MOG$_{35-55}$ was

purchased from GeneScript. Incomplete Freund's adjuvant (IFA) and Mycobacterium tuberculosis

H37Ra (killed and desiccated) were purchased from Difco. Pertussis toxin was purchased from

List Biological Laboratories and diphtheria toxin was obtained from Millipore-Sigma.

Cell purification and culture.

Naïve CD4 cells were isolated from the spleen and peripheral lymph node by a magnetic bead-

based purification according to the manufacturer's instruction (Miltenyi Biotech). Purified CD4

cells were stimulated with plate-bound anti-CD3 (5ug/ml, Bio-X-Cell) and anti-CD28 (2ug/ml,

Bio-X-Cell) for 3 days, in RPMI1640 medium supplemented with 10%FBS, HEPES,

penicillin/streptomycin, MEM Non-Essential Amino Acids, and β-mercaptoethanol. For iTreg cell

differentiation, cells were stimulated in the presence of human IL2 (100u/ml, PeproTech), human

TGFβ (10ng/ml, PeproTech), anti-IL4, and anti-IFNγ antibody (5ug/ml, Clone 11B11 and clone

XMG1.2, Bio-X-Cell) in 10% RPMI medium. CD4$^+$ Te cells were differentiated without

additional cytokine or antibody for three days, followed by additional culture for 2 days in IL2

100u/ml in 10%RPMI medium. nTreg cells were isolated by sorting from the FDG CD4$^+$ fraction

based on Foxp3$^+$GFP and CD25 expression (CD4$^+$CD25$^+$GFP$^+$). Alternatively, nTreg cells were

enriched from CD4 cells by positive selection using the CD4$^+$CD25$^+$ Regulatory T Cell Isolation

Kit from Miltenyi.

Self-antigen induced tolerance model.

1x10^6 HA-specific Thy1.1$^+$ 6.5 CD4 cells from donor mice on a B10.D2 background were

transferred into C3-HA recipient mice, where HA is expressed as self-antigen in the lung; or into

WT B10.D2 mice followed by infection with Vac-HA virus (1x10^6 pfu). HA-reactive T cells were

recovered from the lung-draining lymph node of C3-HA host mice or WT B10.D2 Vac-HA

infected mice at indicated time points by cell sorting. RNA from sorted cells was used for qRT-

PCR assay examining Nrn1 expression.

Peptide-induced T cell anergy model.

5x10^5 Polyclonal Treg cells from CD45.1$^+$ C57BL/6 mice were mixed with 5x10^6 thy1.1$^+$ OTII

cells from Nrn1$^{-/-}$_OTII or ctrl_OTII mice and transferred by *i.v.* injection into TCRα$^{-/-}$ mice.

100ug of OVA$_{323-339}$ dissolved in PBS was administered *i.v.* on days 1, 4, and 7 after cell transfer.

Host mice were harvested on day 13 after cell transfer, and cells from the lymph node and spleen

were further analyzed.

In vivo **Treg suppression assay.**

nTreg cells from CD45.2$^+$CD45.1$^-$ Nrn1$^{-/-}$ or ctrl mice (5x10^5/mouse) in conjunction with CD45.1$^+$

splenocytes (2x10^6/mouse) from FDG mice were cotransferred i.p. into Rag2$^{-/-}$ mice. The CD45.1$^+$

splenocytes were obtained from FDG mice pretreated with DT for 2 days to deplete Treg cells.

Treg suppression toward CD45.1$^+$ responder cells was assessed on day 7 post cell transfer.

Alternatively, 7 days after cell transfer, Rag2$^{-/-}$ hosts were challenged with an *i.d.* inoculation of

B16F10 cells (1x10^5). Tumor growth was monitored daily. Treg-mediated suppression toward

anti-tumor response was assessed by harvesting mice day 18-21 post-tumor inoculation.

Induction of autoimmunity by transient Treg depletion.

To induce autoimmunity in Nrn1$^{-/-}$_FDG and ctrl_FDG mice, 1ug/mouse of DT was administered

i.p. for two consecutive days, and the weight loss of treated mice was observed over time.

EAE induction

EAE was induced in mice by subcutaneous injection of 200 μg MOG_{35-55} peptide with 500 μg M.

tuberculosis strain H37Ra (Difco) emulsified in incomplete Freund Adjuvant oil in 200ul volume

into the flanks at two different sites. In addition, the mice received 400 ng pertussis toxin (PTX;

List Biological Laboratories) *i.p*. at the time of immunization and 48 h later. Clinical signs of EAE

were assessed daily according to the standard 5-point scale (Miller et al., 2007): normal mouse; 0,

no overt signs of disease; 1, limp tail; 2, limp tail plus hindlimb weakness; 3, total hindlimb

paralysis; 4, hindlimb paralysis plus 75% of body paralysis (forelimb paralysis/weakness); 5,

moribund.

ELISA

MaxiSorp ELISA plates (ThermoScientific Nunc) were coated with 100 μl of 1μg/ml anti-mIL-2

(BD Pharmingen #554424) at 4°C overnight. Coated plates were blocked with 200μl of blocking

solution (10%FBS in PBS) for 1hr at room temperature (RT) followed by incubation of culture

supernatant and mIL-2 at different concentrations as standard. After 1hr, plates were washed and

incubated with anti-mIL-2-biotin (BD Pharmingen #554426) at RT for 1hr. After 1hr, plates were

incubated with 100μl of horseradish peroxidase-labeled avidin (Vector Laboratory, #A-2004)

1μg/ml for 30min. After washing, samples were developed using the KPL TMB Peroxidase

substrate system (Seracare #5120-0047) and read at 405 or 450 nm after the addition of the stop

solution.

Quantitative RT-PCR

RNA was isolated using the RNeasy Micro Kit (Qiagen 70004) following the manufacturer's

instructions. RNA was converted to cDNA using the High-Capacity cDNA Reverse Transcription

Kit (ThermoFisher Scientific #4368814) according to the manufacturer's instructions. The primers

of murine genes were purchased from Integrated DNA Technology (IDT). qPCR was performed

using the PowerUp SYBR Green Master Mix (ThermoFisher Scientific #A25780) and the Applied

Biosystems StepOnePlus 96-well real-time PCR system. Gene expression levels were calculated

based on the Delta-Delta Ct relative quantification method. Primers used for Nrn1 PCR were as

follows: GCGGTGCAAATAGCTTACCTG   (forward);   CGGTCTTGATGTTCGTCTTGTC
(reverse).

Ca^{++} flux and Membrane potential measurement
To measure Ca^{++} flux, CD4 cells were loaded with Fluo4 dye at 2μM in the complete cell culture
medium at 37°C for 30min. Cells were washed and resuspended in HBSS Ca^{++} free medium and
plated into 384 well glass bottom assay plate (minimum of 4 wells per sample). Ca^{++} flux was
measured using the FDSS6000 system (Hamamatsu Photonics). To measure store-operated
calcium entry (SOCE), after the recording of the baseline T cells Ca^{++} fluorescent for 1min,
thapsigargin (TG) was added to induce store Ca^{++} depletion, followed by the addition of Ca^{++} 2μM
in the extracellular medium to observe Ca^{++} cellular entry.
Membrane potential was measured using FLIPR Membrane Potential Assay kit (Molecular devices)
according to the manufacturer's instructions. Specifically, T cells were loaded with FLIPR dye by
adding an equal volume of FLIPR dye to the cells and incubated at 37°C for 30 minutes. Relative
membrane potential was measured by detecting FLIPR dye incorporation using flow cytometry.
To measure changes of MP after AAs transport, T cells were plated and loaded with FLIPR dye at
37°C for 30 minutes in 384 well glass bottom assay plate (minimum of 6 wells per sample). After
recording the baseline T cell MP for 1min, MEM AAs (Gibco MEM Amino Acids #11130-051)
were injected into each well, and the change of MP was recorded for 5 min.

Extracellular flux analysis (Seahorse assays).
Real-time measurements of extracellular acidification rate (ECAR) and oxygen consumption rate
(OCR) were performed using an XFe-96 Bioanalyser (Agilent). T cells (2×10^5 cells per well;
minimum of four wells per sample) were spun into previously poly-d-lysine-coated 96-well plates
(Seahorse) in complete RPMI-1640 medium. ECAR was measured in RPMI medium in basal
condition and in response to 25mM glucose, 1μM oligomycin, and 50mM of 2-DG (all from Sigma
Aldrich). OCR was measured in RPMI medium supplemented with 25mM glucose, 2mM L-
glutamine, and 1mM sodium pyruvate, under basal condition and in response to 1μM oligomycin,
1.5μM of carbonylcyanide-4-(trifluoromethoxy)-phenylhydrazone (FCCP) and 1μM of rotenone
and antimycin (all from Sigma Aldrich).

RNAseq and data analysis

RNASeq samples: 1. Anergic T cell analysis. Ctrl and Nrn1$^{-/-}$ OTII cells were sorted from the host mice (n=3 per group). 2. iTreg cell analysis. In vitro differentiated Nrn1$^{-/-}$ and ctrl iTreg cells were replated in resting condition (IL2 100u/ml) or stimulation condition (IL2 100u/ml and aCD3 5ug/ml). Cells were harvested 20 hr after replating for RNASeq analysis. 3. Effector T cells. Nrn1$^{-/-}$ and ctrl CD4 Tn cells were activated for 3 days (aCD3 5ug/ml, aCD28 2ug/ml), followed by replating in IL2 medium (100u/ml). Te cells were harvested two days after replating and subjected to RNASeq analysis.

RNA-sequencing analysis was performed by Admera Health (South Plainfield, NJ). Read quality was assessed with FastQC and aligned to the Mus Musculus genome (Ensembl GRCm38) using STAR aligner (version 2.6.0)(Dobin et al., 2013). Aligned reads were counted using HTSeq (version 0.9.0)(Anders et al., 2015), and the counts were loaded into R (The R Foundation). DESeq2 package (version 1.24.0)(Love et al., 2014) was used to normalize the raw counts. GSEA was performed using public gene sets (HALLMARK, and GO)(Subramanian et al., 2005). Cytoscape was used to display enriched gene sets cluster (Shannon et al., 2003).

Statistical analysis. All numerical data were processed using Graph Pad Prism 10. Data are expressed as the mean +/- the SEM, or as stated. Statistical comparisons were made using an unpaired student t-test or ANOVA with multiple comparison tests where 0.05 was considered significant, and a normal distribution was assumed. The p values are represented as follows: * p<0.05; ** p<0.01; *** p<0.001, **** p<0.0001.

Acknowledgements:

This research is supported by grants from the Bloomberg-Kimmel Institute of JHU, the Melanoma Research Alliance, the National Institutes of Health (RO1AI099300 and RO1AI089830), and the Department of Defense (PC130767). JB's research was supported by a Crohn's and Colitis Foundation of America Research Fellowship, the Melanoma Research Foundation, and NCI grant P30CA016056.

We thank Dennis Gong for data processing and critical reading of the manuscript. We thank Dr. Elly Nedivi for providing polyclonal Nrn1 antibody and Dr. Fan Pan for reagent support. We thank Drs. Franck Housseau, Chien-Fu Hung for the constructive discussion of the project and the manuscript. We thank Drs. Hao Shi and Hongbo Chi for critical reading of the manuscript and

helpful suggestions. We thank Dr. Rachel Helm for manuscript editing. We thank Drs. Richard L.

Huganir, Bian Liu and Hana Goldschmidt for constructive discussion on Nrn1 and AMPAR

connection.

Author contributions:

H. Y. was involved in all aspects of this study, including planning and performing experiments,

analyzing and interpreting data, and writing the manuscript. H.N. and J.B. were involved in

performing experiments and data interpretation. P.V., Y.Z, and A.L. analyzed Nrn1 expression in

Treg cells and performed Treg suppression and functional assay. M.M. was involved in Nrn1

expression, Treg suppression assay, and manuscript writing. C. H. and C.D. conducted the anergy

and Te cell differential gene expression study; Y.Z., J.F., and K.C. conducted an autoimmune

inflammation study, and M.H. helped with mouse colony genotyping. X.Z. and Z.L. contributed

to bioinformatic analysis. D.M.P. oversaw the project and was involved in data interpretation,

manuscript preparation, and funding support.

Competing interests:

C.D. is a co-inventor on patents licensed from JHU to BMS and Janssen and is currently an

employee of Janssen Research. D.M.P. is a consultant for Compugen, Shattuck Labs, WindMIL,

Tempest, Immunai, Bristol-Myers Squibb, Amgen, Janssen, Astellas, Rockspring Capital,

Immunomic, and Dracen; owns founders equity in ManaT Bio Inc., WindMIL, Trex, Jounce, Enara,

Tizona, Tieza, and RAPT; and receives research funding from Compugen, Bristol-Myers Squibb,

and Enara. All other authors do not have conflicting financial interests.

Data availability:

The raw RNA sequencing data has been deposited under the GEO accession no. GSE121908 and

GSE224083.

References:

Abdul Kadir, L., M. Stacey, and R. Barrett-Jolley. 2018. Emerging Roles of the Membrane
Potential: Action Beyond the Action Potential. *Front Physiol* 9:1661.
Adler, A.J., D.W. Marsh, G.S. Yochum, J.L. Guzzo, A. Nigam, W.G. Nelson, and D.M. Pardoll.
1998. CD4+ T cell tolerance to parenchymal self-antigens requires presentation by bone
marrow-derived antigen-presenting cells. *The Journal of experimental medicine*
187:1555-1564.
Anders, S., P.T. Pyl, and W. Huber. 2015. HTSeq--a Python framework to work with high-
throughput sequencing data. *Bioinformatics* 31:166-169.
Babst, M. 2020. Regulation of nutrient transporters by metabolic and environmental stresses.
Curr Opin Cell Biol 65:35-41.
Blackiston, D.J., K.A. McLaughlin, and M. Levin. 2009. Bioelectric controls of cell proliferation:
ion channels, membrane voltage and the cell cycle. *Cell Cycle* 8:3527-3536.
Bohmwald, K., N.M.S. Gálvez, C.A. Andrade, V.P. Mora, J.T. Muñoz, P.A. González, C.A. Riedel,
and A.M. Kalergis. 2021. Modulation of Adaptive Immunity and Viral Infections by Ion
Channels. *Front Physiol* 12:736681.
Bonnet, C.S., S.J. Gilbert, E.J. Blain, A.S. Williams, and D.J. Mason. 2020. AMPA/kainate
glutamate receptor antagonists prevent posttraumatic osteoarthritis. *JCI insight* 5:
Bonnet, C.S., A.S. Williams, S.J. Gilbert, A.K. Harvey, B.A. Evans, and D.J. Mason. 2015.
AMPA/kainate glutamate receptors contribute to inflammation, degeneration and pain
related behaviour in inflammatory stages of arthritis. *Annals of the rheumatic diseases*
74:242-251.
Buck, M.D., R.T. Sowell, S.M. Kaech, and E.L. Pearce. 2017. Metabolic Instruction of Immunity.
Cell 169:570-586.
Cantallops, I., K. Haas, and H.T. Cline. 2000. Postsynaptic CPG15 promotes synaptic maturation
and presynaptic axon arbor elaboration in vivo. *Nature neuroscience* 3:1004-1011.
Chapman, N.M., and H. Chi. 2022. Metabolic adaptation of lymphocytes in immunity and
disease. *Immunity* 55:14-30.
Chappert, P., and R.H. Schwartz. 2010. Induction of T cell anergy: integration of environmental
cues and infectious tolerance. *Current opinion in immunology* 22:552-559.
Chen, T.C., S.P. Cobbold, P.J. Fairchild, and H. Waldmann. 2004. Generation of anergic and
regulatory T cells following prolonged exposure to a harmless antigen. *Journal of*
immunology (Baltimore, Md. : 1950) 172:5900-5907.
Choi, S., and R.H. Schwartz. 2007. Molecular mechanisms for adaptive tolerance and other T cell
anergy models. *Seminars in immunology* 19:140-152.
Cuenca, A., F. Cheng, H. Wang, J. Brayer, P. Horna, L. Gu, H. Bien, I.M. Borrello, H.I. Levitsky, and
E.M. Sotomayor. 2003. Extra-lymphatic solid tumor growth is not immunologically
ignored and results in early induction of antigen-specific T-cell anergy: dominant role of
cross-tolerance to tumor antigens. *Cancer research* 63:9007-9015.
DL, K.L.a.M. 2017. Relationship between CD4 Regulatory T Cells and Anergy In Vivo. *Journal of*
immunology (Baltimore, Md. : 1950) 198:2527.
Dobin, A., C.A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, and
T.R. Gingeras. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29:15-21.

Dvorak, V., T. Wiedmer, A. Ingles-Prieto, P. Altermatt, H. Batoulis, F. Bärenz, E. Bender, D.
704   Digles, F. Dürrenberger, L.H. Heitman, I.J. AP, D.B. Kell, S. Kickinger, D. Körzö, P. Leippe,
705   T. Licher, V. Manolova, R. Rizzetto, F. Sassone, L. Scarabottolo, A. Schlessinger, V.
706   Schneider, H.J. Sijben, A.L. Steck, H. Sundström, S. Tremolada, M. Wilhelm, M. Wright
707   Muelas, D. Zindel, C.M. Steppan, and G. Superti-Furga. 2021. An Overview of Cell-Based
708   Assay Platforms for the Solute Carrier Family of Transporters. *Frontiers in pharmacology*
709   12:722889.
710   ElTanbouly, M.A., and R.J. Noelle. 2021. Rethinking peripheral T cell tolerance: checkpoints
711   across a T cell's journey. *Nat Rev Immunol* 21:257-267.
712   Emmons-Bell, M., and I.K. Hariharan. 2021. Membrane potential regulates Hedgehog signalling
713   in the Drosophila wing imaginal disc. *EMBO reports* 22:e51861.
714   Feng, Y., A. Arvey, T. Chinen, J. van der Veeken, G. Gasteiger, and A.Y. Rudensky. 2014. Control
715   of the inheritance of regulatory T cell identity by a cis element in the Foxp3 locus. *Cell*
716   158:749-763.
717   Floess, S., J. Freyer, C. Siewert, U. Baron, S. Olek, J. Polansky, K. Schlawe, H.D. Chang, T. Bopp, E.
718   Schmitt, S. Klein-Hessling, E. Serfling, A. Hamann, and J. Huehn. 2007. Epigenetic control
719   of the foxp3 locus in regulatory T cells. *PLoS biology* 5:e38.
720   Fujino, T., J.H. Leslie, R. Eavri, J.L. Chen, W.C. Lin, G.H. Flanders, E. Borok, T.L. Horvath, and E.
721   Nedivi. 2011. CPG15 regulates synapse stability in the developing and adult brain. *Genes*
722   *& development* 25:2674-2685.
723   Geltink, R.I.K., R.L. Kyle, and E.L. Pearce. 2018. Unraveling the Complex Interplay Between T Cell
724   Metabolism and Function. *Annual review of immunology* 36:461-488.
725   Gonzalez-Figueroa, P., J.A. Roco, I. Papa, L. Núñez Villacís, M. Stanley, M.A. Linterman, A. Dent,
726   P.F. Canete, and C.G. Vinuesa. 2021. Follicular regulatory T cells produce neuritin to
727   regulate B cells. *Cell*
728   Hamill, M.J., R. Afeyan, M.V. Chakravarthy, and T. Tramontin. 2020. Endogenous Metabolic
729   Modulators: Emerging Therapeutic Potential of Amino Acids. *iScience* 23:101628.
730   Huang, C.T., D.L. Huso, Z. Lu, T. Wang, G. Zhou, E.P. Kennedy, C.G. Drake, D.J. Morgan, L.A.
731   Sherman, A.D. Higgins, D.M. Pardoll, and A.J. Adler. 2003. CD4+ T cells pass through an
732   effector phase during the process of in vivo tolerance induction. *Journal of immunology*
733   *(Baltimore, Md. : 1950)* 170:3945-3953.
734   Huang, C.T., C.J. Workman, D. Flies, X. Pan, A.L. Marson, G. Zhou, E.L. Hipkiss, S. Ravi, J.
735   Kowalski, H.I. Levitsky, J.D. Powell, D.M. Pardoll, C.G. Drake, and D.A. Vignali. 2004. Role
736   of LAG-3 in regulatory T cells. *Immunity* 21:503-513.
737   Javaherian, A., and H.T. Cline. 2005. Coordinated motor neuron axon growth and
738   neuromuscular synaptogenesis are promoted by CPG15 in vivo. *Neuron* 45:505-512.
739   Joesch, C., E. Guevarra, S.P. Parel, A. Bergner, P. Zbinden, D. Konrad, and H. Albrecht. 2008. Use
740   of FLIPR membrane potential dyes for validation of high-throughput screening with the
741   FLIPR and microARCS technologies: identification of ion channel modulators acting on
742   the GABA(A) receptor. *J Biomol Screen* 13:218-228.
743   Kalekar, L.A., S.E. Schmiel, S.L. Nandiwada, W.Y. Lam, L.O. Barsness, N. Zhang, G.L. Stritesky, D.
744   Malhotra, K.E. Pauken, J.L. Linehan, M.G. O'Sullivan, B.T. Fife, K.A. Hogquist, M.K.
745   Jenkins, and D.L. Mueller. 2016. CD4(+) T cell anergy prevents autoimmunity and
746   generates regulatory T cell precursors. *Nature immunology* 17:304-314.

Kiefer, H., A.J. Blume, and H.R. Kaback. 1980. Membrane potential changes during mitogenic
stimulation of mouse spleen lymphocytes. *Proceedings of the National Academy of*
Sciences of the United States of America 77:2200-2204.
Kim, J.M., J.P. Rasmussen, and A.Y. Rudensky. 2007. Regulatory T cells prevent catastrophic
autoimmunity throughout the lifespan of mice. *Nature immunology* 8:191-197.
Kuczma, M.P., E.A. Szurek, A. Cebula, V.L. Ngo, M. Pietrzak, P. Kraj, T.L. Denning, and L.
Ignatowicz. 2021. Self and microbiota-derived epitopes induce CD4(+) T cell anergy and
conversion into CD4(+)Foxp3(+) regulatory cells. *Mucosal Immunol* 14:443-454.
Levin, M. 2021. Bioelectric signaling: Reprogrammable circuits underlying embryogenesis,
regeneration, and cancer. *Cell* 184:1971-1989.
Li, X., Y. Liang, M. LeBlanc, C. Benner, and Y. Zheng. 2014. Function of a Foxp3 cis-element in
protecting regulatory T cell identity. *Cell* 158:734-748.
Lim, D.G., Y.H. Park, S.E. Kim, S.H. Jeong, and S.C. Kim. 2013. Diagnostic value of tolerance-
related gene expression measured in the recipient alloantigen-reactive T cell fraction.
Clinical immunology (Orlando, Fla.) 148:219-226.
Liu, G.Y., and D.M. Sabatini. 2020. mTOR at the nexus of nutrition, growth, ageing and disease.
Nat Rev Mol Cell Biol 21:183-203.
Long, L., J. Wei, S.A. Lim, J.L. Raynor, H. Shi, J.P. Connelly, H. Wang, C. Guy, B. Xie, N.M.
Chapman, G. Fu, Y. Wang, H. Huang, W. Su, J. Saravia, I. Risch, Y.D. Wang, Y. Li, M. Niu, Y.
Dhungana, A. Kc, P. Zhou, P. Vogel, J. Yu, S.M. Pruett-Miller, J. Peng, and H. Chi. 2021.
CRISPR screens unveil signal hubs for nutrient licensing of T cell immunity. *Nature*
600:308-313.
Love, M.I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome biology* 15:550.
Ma, Y., K. Poole, J. Goyette, and K. Gaus. 2017. Introducing Membrane Charge and Membrane
Potential to T Cell Signaling. *Frontiers in immunology* 8:1513.
Martinez, R.J., N. Zhang, S.R. Thomas, S.L. Nandiwada, M.K. Jenkins, B.A. Binstadt, and D.L.
Mueller. 2012. Arthritogenic self-reactive CD4+ T cells acquire an FR4hiCD73hi anergic
state in the presence of Foxp3+ regulatory T cells. *Journal of immunology (Baltimore,*
Md. : 1950) 188:170-181.
McNearney, T., B.A. Baethge, S. Cao, R. Alam, J.R. Lisse, and K.N. Westlund. 2004. Excitatory
amino acids, TNF-alpha, and chemokine levels in synovial fluids of patients with active
arthropathies. *Clinical and experimental immunology* 137:621-627.
Mercadante, E.R., and U.M. Lorenz. 2016. Breaking Free of Control: How Conventional T Cells
Overcome Regulatory T Cell Suppression. *Frontiers in immunology* 7:193.
Miller, S.D., W.J. Karpus, and T.S. Davidson. 2007. Experimental Autoimmune Encephalomyelitis
in the Mouse. *Current protocols in immunology* CHAPTER:Unit-15 11.
Monroe, J.G., and J.C. Cambier. 1983. B cell activation. I. Anti-immunoglobulin-induced receptor
cross-linking results in a decrease in the plasma membrane potential of murine B
lymphocytes. *The Journal of experimental medicine* 157:2073-2086.
Nedivi, E., G.Y. Wu, and H.T. Cline. 1998. Promotion of dendritic growth by CPG15, an activity-
induced signaling molecule. *Science (New York, N.Y.)* 281:1863-1866.

Nik, A.M., B. Pressly, V. Singh, S. Antrobus, S. Hulsizer, M.A. Rogawski, H. Wulff, and I.N. Pessah.
2017. Rapid Throughput Analysis of GABA(A) Receptor Subtype Modulators and Blockers
Using DiSBAC(1)(3) Membrane Potential Red Dye. *Mol Pharmacol* 92:88-99.
Nystrom, S.N., D. Bourges, S. Garry, E.M. Ross, I.R. van Driel, and P.A. Gleeson. 2014. Transient
Treg-cell depletion in adult mice results in persistent self-reactive CD4(+) T-cell
responses. *European journal of immunology* 44:3621-3631.
Olenchock, B.A., J.C. Rathmell, and M.G. Vander Heiden. 2017. Biochemical Underpinnings of
Immune Cell Metabolic Phenotypes. *Immunity* 46:703-713.
Opejin, A., A. Surnov, Z. Misulovin, M. Pherson, C. Gross, C.A. Iberg, I. Fallahee, J. Bourque, D.
Dorsett, and D. Hawiger. 2020. A Two-Step Process of Effector Programming Governs
CD4(+) T Cell Fate Determination Induced by Antigenic Activation in the Steady State.
Cell reports 33:108424.
Pandya, N.J., C. Seeger, N. Babai, M.A. Gonzalez-Lozano, V. Mack, J.C. Lodder, Y. Gouwenberg,
H.D. Mansvelder, U.H. Danielson, K.W. Li, M. Heine, S. Spijker, R. Frischknecht, and A.B.
Smit. 2018. Noelin1 Affects Lateral Mobility of Synaptic AMPA Receptors. *Cell reports*
24:1218-1230.
Peng, M., and M.O. Li. 2023. Metabolism along the life journey of T cells. *Life Metab* 2:
Plitas, G., C. Konopacki, K. Wu, P.D. Bos, M. Morrow, E.V. Putintseva, D.M. Chudakov, and A.Y.
Rudensky. 2016. Regulatory T Cells Exhibit Distinct Features in Human Breast Cancer.
Immunity 45:1122-1134.
Putz, U., C. Harwell, and E. Nedivi. 2005. Soluble CPG15 expressed during early development
rescues cortical progenitors from apoptosis. *Nature neuroscience* 8:322-331.
Ramirez, G.A., L.A. Coletto, C. Sciorati, E.P. Bozzolo, P. Manunta, P. Rovere-Querini, and A.A.
Manfredi. 2018. Ion Channels and Transporters in Inflammation: Special Focus on TRP
Channels and TRPC6. *Cells* 7:
Safford, M., S. Collins, M.A. Lutz, A. Allen, C.T. Huang, J. Kowalski, A. Blackford, M.R. Horton, C.
Drake, R.H. Schwartz, and J.D. Powell. 2005. Egr-2 and Egr-3 are negative regulators of T
cell activation. *Nature immunology* 6:472-480.
Salmond, R.J. 2018. mTOR Regulation of Glycolytic Metabolism in T Cells. *Frontiers in cell and*
developmental biology 6:122.
Saravia, J., J.L. Raynor, N.M. Chapman, S.A. Lim, and H. Chi. 2020. Signaling networks in
immunometabolism. *Cell research* 30:328-342.
Sarchielli, P., M. Di Filippo, A. Candeliere, D. Chiasserini, A. Mattioni, S. Tenaglia, M. Bonucci,
and P. Calabresi. 2007. Expression of ionotropic glutamate receptor GLUR3 and effects
of glutamate on MBP- and MOG-specific lymphocyte activation and chemotactic
migration in multiple sclerosis patients. *Journal of neuroimmunology* 188:146-158.
Schietinger, A., J.J. Delrow, R.S. Basom, J.N. Blattman, and P.D. Greenberg. 2012. Rescued
tolerant CD8 T cells are preprogrammed to reestablish the tolerant state. *Science (New*
York, N.Y.) 335:723-727.
Schietinger, A., M. Philip, V.E. Krisnawan, E.Y. Chiu, J.J. Delrow, R.S. Basom, P. Lauer, D.G.
Brockstedt, S.E. Knoblaugh, G.J. Hammerling, T.D. Schell, N. Garbi, and P.D. Greenberg.
2016. Tumor-Specific T Cell Dysfunction Is a Dynamic Antigen-Driven Differentiation
Program Initiated Early during Tumorigenesis. *Immunity* 45:389-401.

Schwenk, J., N. Harmel, A. Brechet, G. Zolles, H. Berkefeld, C.S. Muller, W. Bildl, D. Baehrens, B.
Huber, A. Kulik, N. Klocker, U. Schulte, and B. Fakler. 2012. High-resolution proteomics
unravel architecture and molecular diversity of native AMPA receptor complexes.
Neuron 74:621-633.
Shannon, P., A. Markiel, O. Ozier, N.S. Baliga, J.T. Wang, D. Ramage, N. Amin, B. Schwikowski,
and T. Ideker. 2003. Cytoscape: a software environment for integrated models of
biomolecular interaction networks. *Genome Res* 13:2498-2504.
Shimada, T., T. Yoshida, and K. Yamagata. 2016. Neuritin Mediates Activity-Dependent Axonal
Branch Formation in Part via FGF Signaling. *The Journal of neuroscience : the official*
journal of the Society for Neuroscience 36:4534-4548.
Shin, D.S., A. Jordan, S. Basu, R.M. Thomas, S. Bandyopadhyay, E.F. de Zoeten, A.D. Wells, and F.
Macian. 2014. Regulatory T cells suppress CD4+ T cells through NFAT-dependent
transcriptional mechanisms. *EMBO reports* 15:991-999.
Silva Morales, M., and D. Mueller. 2018. Anergy into T regulatory cells: an integration of
metabolic cues and epigenetic changes at the Foxp3 conserved non-coding sequence 2.
F1000Research 7:
Sinclair, L.V., J. Rolf, E. Emslie, Y.B. Shi, P.M. Taylor, and D.A. Cantrell. 2013. Control of amino-
acid transport by antigen receptors coordinates the metabolic reprogramming essential
for T cell differentiation. *Nature immunology* 14:500-508.
Singer, M., C. Wang, L. Cong, N.D. Marjanovic, M.S. Kowalczyk, H. Zhang, J. Nyman, K. Sakuishi,
S. Kurtulus, D. Gennert, J. Xia, J.Y. Kwon, J. Nevin, R.H. Herbst, I. Yanai, O. Rozenblatt-
Rosen, V.K. Kuchroo, A. Regev, and A.C. Anderson. 2016. A Distinct Gene Module for
Dysfunction Uncoupled from Activation in Tumor-Infiltrating T Cells. *Cell* 166:1500-
1511.e1509.
Subramanian, A., P. Tamayo, V.K. Mootha, S. Mukherjee, B.L. Ebert, M.A. Gillette, A. Paulovich,
S.L. Pomeroy, T.R. Golub, E.S. Lander, and J.P. Mesirov. 2005. Gene set enrichment
analysis: a knowledge-based approach for interpreting genome-wide expression
profiles. *Proceedings of the National Academy of Sciences of the United States of*
America 102:15545-15550.
Subramanian, J., K. Michel, M. Benoit, and E. Nedivi. 2019. CPG15/Neuritin Mimics Experience
in Selecting Excitatory Synapses for Stabilization by Facilitating PSD95 Recruitment. *Cell*
reports 28:1584-1595.e1585.
Sullivan, M.R., L.V. Danai, C.A. Lewis, S.H. Chan, D.Y. Gui, T. Kunchok, E.A. Dennstedt, M.G.
Vander Heiden, and A. Muir. 2019. Quantification of microenvironmental metabolites in
murine cancers reveals determinants of tumor nutrient availability. *eLife* 8:
Sundelacruz, S., M. Levin, and D.L. Kaplan. 2009. Role of membrane potential in the regulation
of cell proliferation and differentiation. *Stem Cell Rev Rep* 5:231-246.
Vahl, J.C., C. Drees, K. Heger, S. Heink, J.C. Fischer, J. Nedjic, N. Ohkura, H. Morikawa, H. Poeck,
S. Schallenberg, D. Riess, M.Y. Hein, T. Buch, B. Polic, A. Schonle, R. Zeiser, A. Schmitt-
Graff, K. Kretschmer, L. Klein, T. Korn, S. Sakaguchi, and M. Schmidt-Supprian. 2014.
Continuous T cell receptor signals maintain a functional regulatory T cell pool. *Immunity*
41:722-736.

Vanasek, T.L., S.L. Nandiwada, M.K. Jenkins, and D.L. Mueller. 2006. CD25+Foxp3+ regulatory T
cells facilitate CD4+ T cell clonal anergy induction during the recovery from
lymphopenia. *Journal of immunology (Baltimore, Md. : 1950)* 176:5880-5889.
Wang, Y., A. Tao, M. Vaeth, and S. Feske. 2020. Calcium regulation of T cell metabolism. *Current*
opinion in physiology 17:207-223.
Whiteaker, K.L., S.M. Gopalakrishnan, D. Groebe, C.C. Shieh, U. Warrior, D.J. Burns, M.J.
Coghlan, V.E. Scott, and M. Gopalakrishnan. 2001. Validation of FLIPR membrane
potential dye for high throughput screening of potassium channel modulators. *J Biomol*
Screen 6:305-312.
Workman, C.J., L.W. Collison, M. Bettini, M.R. Pillai, J.E. Rehg, and D.A. Vignali. 2011. In vivo
Treg suppression assays. *Methods in molecular biology (Clifton, N.J.)* 707:119-156.
Yao, J.J., X.F. Gao, C.W. Chow, X.Q. Zhan, C.L. Hu, and Y.A. Mei. 2012. Neuritin activates insulin
receptor pathway to up-regulate Kv4.2-mediated transient outward K+ current in rat
cerebellar granule neurons. *The Journal of biological chemistry* 287:41534-41545.
Yu, W., Z. Wang, X. Yu, Y. Zhao, Z. Xie, K. Zhang, Z. Chi, S. Chen, T. Xu, D. Jiang, X. Guo, M. Li, J.
Zhang, H. Fang, D. Yang, Y. Guo, X. Yang, X. Zhang, Y. Wu, W. Yang, and D. Wang. 2022.
Kir2.1-mediated membrane potential promotes nutrient acquisition and inflammation
through regulation of nutrient transporters. *Nature communications* 13:3544.
Zha, Y., R. Marks, A.W. Ho, A.C. Peterson, S. Janardhan, I. Brown, K. Praveen, S. Stang, J.C.
Stone, and T.F. Gajewski. 2006. T cell anergy is reversed by active Ras and is regulated
by diacylglycerol kinase-alpha. *Nature immunology* 7:1166-1173.
Zheng, Y., G.M. Delgoffe, C.F. Meyer, W. Chan, and J.D. Powell. 2009. Anergic T cells are
metabolically anergic. *Journal of immunology (Baltimore, Md. : 1950)* 183:6095-6101.
Zheng, Y., S. Josefowicz, A. Chaudhry, X.P. Peng, K. Forbush, and A.Y. Rudensky. 2010. Role of
conserved non-coding DNA elements in the Foxp3 gene in regulatory T-cell fate. *Nature*
463:808-812.
Zhou, S., and J. Zhou. 2014. Neuritin, a neurotrophic factor in nervous system physiology.
Current medicinal chemistry 21:1212-1219.
Zhou, Y., C.O. Wong, K.J. Cho, D. van der Hoeven, H. Liang, D.P. Thakur, J. Luo, M. Babic, K.E.
Zinsmaier, M.X. Zhu, H. Hu, K. Venkatachalam, and J.F. Hancock. 2015. SIGNAL
TRANSDUCTION. Membrane potential modulates plasma membrane phospholipid
dynamics and K-Ras signaling. *Science (New York, N.Y.)* 349:873-876.
Zito, A., D. Cartelli, G. Cappelletti, A. Cariboni, W. Andrews, J. Parnavelas, A. Poletti, and M.
Galbiati. 2014. Neuritin 1 promotes neuronal migration. *Brain Structure and Function*
219:105-118.
909

Figure 1. Nrn1 expression and function in anergic T cells.

(**A**) Experimental scheme identifying Nrn1 in anergic T cells and qRT-PCR confirmation of Nrn1

expression in HA-specific CD4 cells recovered from HA-expressing host vs WT host activated

with Vac_HA virus. (**B**) qRT-PCR and western blot detecting Nrn1 expression in naïve

CD4$^+$CD62L^{hi}CD44^{lo} Tn cell, CD4 effector CD4$^+$Foxp3$^-$CD44^{hi}CD73$^-$FR$^-$ Te cells and CD4

anergic CD4$^+$Foxp3$^-$CD44^{hi}CD73$^+$FR$^+$ Ta cells. (**C**) Nrn1 expression was measured by qRT-PCR

and western blot among naïve CD4$^+$ T cells, CD4$^+$Foxp3$^+$ nTreg, and *in vitro* generated iTregs.

(**D**) Nrn1 expression was detected by qRT-PCR and flow cytometry among WT naïve CD4$^+$ cells

and activated CD4$^+$ cells on days 1, 2, and 3 after activation. Nrn1$^{-/-}$ CD4 cells were also stained

for Nrn1 three days after activation. qPCR Data are presented as average $\pm$ SEM. *p<0.05,

**p<0.01, ***p<0.001, ****p<0.0001. Triplicates were used. Ordinary one-way ANOVA was

performed for multi-comparison.

(**E-J**). Anergy induction *in vivo*. (**E**) Experimental outline evaluating anergy development *in vivo*:

2x10^6 Thy1.1$^+$ Nrn1$^{-/-}$ or ctrl CD4 OTII T cells were co-transferred with 5x10^5 Thy1.2$^+$Thy1.1$^-$

WT Treg cells into TCR$\alpha^{-/-}$mice. Cells were recovered on day 13 post-transfer. (**F**) Proportions

and numbers of OTII cells recovered from recipient spleen; (**G**) IL2 secretion from OTII cells

upon *ex vivo* stimulation with OVA peptide. (**H**) Foxp3$^+$ cell proportion among Thy1.1$^+$ Nrn1$^{-/-}$ or

ctrl CD4 cells. (**I & J**) Nrn1$^{-/-}$ vs ctrl OTII cells recovered from the peptide-induced anergy model

were subjected to bulk RNASeq analysis. GSEA comparing the expression of signature genes for

anergy (**I**) and Treg (**J**) among ctrl and Nrn1$^{-/-}$ OTII cells.

Data are presented as mean $\pm$SEM and representative of 3 independent experiments (N$\geq$4 mice

per group). *p<0.05, **p<0.01, ***p<0.001. Unpaired Student's t-tests were performed.

Figure 2. Reduced proliferation and suppression function in Nrn1$^{-/-}$ Treg cells.

(**A**). Proportion of Foxp3$^+$ cells three days after *in vitro* iTreg differentiation. (**B-D**) iTreg cell expansion after restimulation. (**B**) The number of live cells from day 1 to day 3 after iTreg cell restimulation with anti-CD3. (**C**) Ki67 expression among CD4$^+$Foxp3$^+$ cells day 3 after restimulation. (**D**) Foxp3$^+$ cell proportion and number among live CD4$^+$ cells day 3 after restimulation. Triplicates in each experiment, data represent one of four independent experiments. (**E-M**). Nrn1$^{-/-}$ or ctrl nTreg cells expansion and suppression *in vivo*. (**E**) The experimental scheme. CD45.2$^+$ nTreg T cells from Nrn1$^{-/-}$ or ctrl were transferred with CD45.1$^+$ FDG splenocytes devoid

of Tregs into Rag2$^{-/-}$ host. Treg cell expansion and suppression toward FDG CD45.1$^+$ responder

cells were evaluated on day 7 post cell transfer. Alternatively, B16F10 tumor cells were inoculated

on day 7 after cell transfer and monitored for tumor growth. (**F-J**) CD45.2$^+$ cell proportion (**F**),

Foxp3 retention (**G**), and Ki67 expression among Foxp3$^+$ cells (**H**) at day 7 post cell transfer. (**I**)

CD45.1$^+$ cell proportion and number in the spleen of Nrn1$^{-/-}$ or ctrl Treg hosts day 7 post cell

transfer. (**J-L**)**.** Treg cell suppression toward anti-tumor response. (**J**) Tumor growth curve and

tumor size at harvest from Nrn1$^{-/-}$ or ctrl nTreg hosts. (**K**) CD45.1$^+$ cell count in tumor draining

lymph node (LN) and spleen. (**L**) the proportion of CD45.1$^+$ cells among CD45$^+$ tumor lymphocyte

infiltrates (TILs). (**M**) IFNγ% among CD8$^+$ T cells in TILs. n$\geq$5 mice per group. (**F-I**) represents

three independent experiments, (**J-M**) represents two independent experiments. Data are presented

as mean $\pm$SEM *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. Unpaired Student's t-tests were

performed.

Figure.3. Nrn1 expression impacts Treg cell electrical and metabolic state.

(**A-C**). Gene sets clusters enriched in Nrn1[-/-] and ctrl iTreg cells. Gene sets cluster analysis via

Cytoscape was performed on Gene ontology Molecular Function (GO_MF) gene sets. The results

cutoff: p-value ≤0.05 and FDR q-value ≤0.1. (**A**) Gene sets cluster in Nrn1[-/-] iTreg cells cultured

under resting condition (IL2 only) (Figure 3-figure supplement Table 1). (**B**) Gene sets clusters in

Nrn1$^{-/-}$ and ctrl iTreg cells reactivated with anti-CD3 (Figure 3-figure supplement Table 2). (**C**)

Comparison of enriched gene sets in Nrn1$^{-/-}$ under resting vs. activating condition (Figure 3-figure

supplement Table 3). (**D-F**) Changes relating to cell electric state. (**D**) Enrichment of

"GOMF_Neurotransmitter receptor activity involved in the regulation of postsynaptic membrane

potential" gene set and enriched gene expression heatmap. (**E**) Membrane potential was measured

in Nrn1$^{-/-}$ and ctrl iTreg cells cultured in IL2 or activated with anti-CD3 in the presence of IL2.

Data represent three independent experiments. (**F**) Enrichment of "GOMF_Metal ion

transmembrane transporter activity" gene set and enriched gene expression heatmap (Figure 3-

figure supplement 1A). (**G-K**). Metabolic changes associated with Nrn1$^{-/-}$ iTreg. (**G**) Heatmap of

differentially expressed amino acid (AA) transport-related genes (from "MF_Amino acid

transmembrane transporter activity" gene list) in Nrn1$^{-/-}$ and ctrl iTreg cells. (**H**) AAs induced MP

changes in Nrn1$^{-/-}$ and ctrl iTreg cells. Data represent three independent experiments. (**I**)

Measurement of pmTOR and pS6 in iTreg cells that were deprived of nutrients for 1h and refed

with RPMI for two hours. (**J**) Hallmark gene sets significantly enriched in Nrn1$^{-/-}$ and ctrl iTreg.

NOM p-val<0.05, FDR q-val<0.25. (**K**) Seahorse analysis of extracellular acidification rate

(ECAR) and oxygen consumption rate (OCR) in Nrn1$^{-/-}$ and ctrl iTreg cells. n=6~10 technical

replicates per group. Data represent three independent experiments. **p<0.01, ***p<0.001,

****p<0.0001. Unpaired student t-test for two-group comparison. Unpaired t-test (H, K), two-way

ANOVA (E, I). ns, not significant.

Figure 4. Nrn1 deficiency affects Te cell response.

(**A**) Comparison of cell proliferation and cytokine expression in Nrn1$^{-/-}$ and ctrl Te cells. Data represent one of three independent experiments. (**B-E**) An enhanced autoimmune response in Nrn1$^{-/-}$ mice *in vivo*. (**B**) Experimental scheme. Nrn1$^{-/-}$ mice were crossed with FDG mice and

Nrn1$^{-/-}$_FDG or ctrl_FDG mice were obtained. The autoimmune response was induced by injecting

DT i.p. to delete endogenous Treg cells. Mice's weight change was monitored after disease

induction. (**C**) Relative body weight change after autoimmune response induction. (**D**) Mice were

harvested six days after DT injection and assessed for ki67, cytokine TNFα, IL2, and IFNγ

expression in CD4$^+$ cells. (**E**) Foxp3 expression among CD4$^+$ cells day six post DT treatment. n$\geq$5

mice per group. Data represent four independent experiments. (**F-I**) Changes relating to ion

balances in Te cells. (**F**) Gene sets clusters from GSEA of GO_MF and GO_Biological process

(GO_BP) results in Nrn1$^{-/-}$ and ctrl Te cells (Figure 4-figure supplement Table 4). (**G**) Enrichment

of "GOBP_ membrane repolarization" gene set and enriched gene expression heatmap. (**H**)

Membrane potential measurement in Te cells. Data represent two independent experiments. (**I**)

Enrichment of "GOMF_Metal ion transmembrane transporter activity" gene set and heatmap of

differential gene expression pattern (Figure 4-figure supplement 1B). (**J-N**) Metabolic changes

associated with Nrn1$^{-/-}$ Te cell. (**J**) Enrichment of "GOMF_amino acid transmembrane transporter

activity" gene set and differential gene expression heatmap. (**K**) AAs induced MP changes in Te

cells. Data represent two independent experiments. (**L**) Measurement of pmTOR and pS6 in Te

cells after nutrient sensing. Data represent three independent experiments. (**M**) Enriched Hallmark

gene sets (p<0.05, FDR q<0.25). (**N**) Seahorse analysis of extracellular acidification rate (ECAR)

and oxygen consumption rate (OCR) in Nrn1$^{-/-}$ and ctrl Te cells. n$\geq$6 technical replicates per group.

Data represent three independent experiments. Error bars indicate $\pm$SEM. *p<0.05, **p<0.01,

***p<0.001, ****p<0.0001, unpaired Student's t-test was performed for two-group comparison.

1014

1015

1016

1017
Figure 5. Nrn1 deficiency exacerbates autoimmune EAE disease.

(**A**) Aggravated body weight loss and protracted EAE disease in Nrn1$^{-/-}$ mice. (**B**) CD45$^+$ cell

number in the spinal cord infiltrates. (**C**) CD4$^+$ cell number in the spinal cord infiltrates. (**D**)

Mog$_{38-49}$/IA^b tetramer staining of spinal cord infiltrating CD4 cells. (**E**) Foxp3$^+$ proportion among

CD4$^+$ cells in spinal cord infiltrates. (**F**) IFNγ$^+$ and IL17$^+$ cell proportion among CD4$^+$ cells in

draining lymph nodes. n$\geq$5 mice per group. Data represent three independent experiments. The P

value was calculated by 2way ANOVA for (**A**). The p-value was calculated by the unpaired student

t-test for (**B-F**). *P<0.05, **P<0.01.

1026
1027

1028 **Figure supplements**

1029

1030 **Neurotrophic factor Neuritin modulates T cell electrical and metabolic state for the balance**

1031 **of tolerance and immunity**

1032 **Hong Yu** *et al.*

1033

1034

1035

Figure 1-figure supplement 1. Nrn1 expression in T cells from tumor environment and
during early T cell activation. (A-B). Nrn1 expression in tumor infiltrates. (A) Comparison of
Nrn1 expression by qRT-PCR among Tregs and non-Treg CD44^{hi}CD4$^+$ cells recovered either from
B16 melanoma infiltrates or from peripheral blood of Foxp3DTRgfp mice bearing subcutaneous
B16 melanomas. **(B)** Comparison of Nrn1 expression in breast tumor-infiltrating Treg (T-Treg)
and Te (T-CD4$^+$Foxp3$^-$) cells vs. peripheral blood Treg (P-Treg) and Te (PBMC CD4$^+$Foxp3$^-$)
cells. Data derived from the "Regulatory T Cells Exhibit Distinct Features in Human Breast Cancer"
report (Plitas et al., 2016). **(C)** Nrn1 cell surface detection on day 2 activated CD4 and CD8 cells
by flow cytometry. **(D).** Detection of Nrn1 expression in nTreg cells by qRT-PCR and cell surface
Nrn1 staining.

1046

1047

1048

1049

Figure 1-figure supplement 2. Nrn1^{-/-} mice body weight and immune cell profile analysis compared to Nrn1^{+/-} and WT mice. (**A**) Average body weight of 10-12 wk old age and sex-matched Nrn1^{-/-}, Nrn1^{+/-} and WT mice. (**B**) Thymus and peripheral lymphoid tissue total cell count. (**C-D**) Immune cell frequencies in the thymus and spleen. (**E**) Proportion of CD4$^+$Foxp3$^-$ CD44$^+$FR4^{hi} CD73^{hi} anergic T cells among splenocytes CD4 cell population. (**F**) FOXP3$^+$cell frequency among CD4 cells in the spleen. The immune profile assessment used n>3 mice/group of Nrn1^{-/-} and control mice. *P* values were calculated by one-way analysis of variance (ANOVA). *$P<0.05$. **$P<0.01$.

1062

Figure 1-figure supplement 3. Compromised T cell activation in Nrn1$^{-/-}$ cells. **(A)** Cell tracker dye violet (CTV) dilution in Nrn1$^{-/-}$ or ctrl CD4 cells after stimulation with plate-bound aCD3 (5ug/ml) and soluble aCD28 (2ug/ml); **(B)** Cell surface activation markers CD25, CD44, CD69, and PD1 expression day 2 after naïve CD4$^+$ cell activation. Unpaired student's t-test, *p<0.05, **p<0.01. Data represent three independent experiments. **(C)** Store-operated Ca^{++} entry (SOCE) was examined on day 2 activated CD4$^+$ cells labeled with Fluo-4 dye. Representative graph and mean $\pm$ SEM of SOCE induced by CD4 cell stimulation with 1uM thapsigargin (TG) in Ca^{++} free HBSS (0mM Ca^{++}) followed by addition of 2mM Ca^{++}. Representative graphs of Ca^{++} influx from three independent experiments (****p<0.0001).

1072

1073

1074 **Figure 3-figure supplement Table 1. Gene sets enriched in Nrn1$^{-/-}$ iTreg cells cultured under**

1075 **the resting condition.**

Nrn1$^{-/-}$	**Sodium transport**
	SODIUM_CHANNEL_ACTIVITY
	VOLTAGE_GATED_SODIUM_CHANNEL_ACTIVITY
	MONOATOMIC_ANION_MONOATOMIC_CATION_SYMPORTER_ACTIVITY
	SODIUM_ION_TRANSMEMBRANE_TRANSPORTER_ACTIVITY
	MONOATOMIC_ANION_SODIUM_SYMPORTER_ACTIVITY
	SOLUTE_SODIUM_SYMPORTER_ACTIVITY
	ORGANIC_ACID_SODIUM_SYMPORTER_ACTIVITY
	SOLUTE_MONOATOMIC_CATION_SYMPORTER_ACTIVITY
	Neurotransmitter and Membrane potential
	POSTSYNAPTIC_NEUROTRANSMITTER_RECEPTOR_ACTIVITY
	NEUROTRANSMITTER_RECEPTOR_ACTIVITY_INVOLVED_IN_REGULATION _OF_POSTSYNAPTIC_MEMBRANE_POTENTIAL
	Receptor kinase activity
	TRANSMEMBRANE_RECEPTOR_PROTEIN_KINASE_ACTIVITY
	TRANSMEMBRANE_RECEPTOR_PROTEIN_TYROSINE_KINASE_ACTIVITY

1076   Gene sets involved in Figure. 3A clusters enriched in Nrn1$^{-/-}$ iTreg cells cultured under the resting condition.

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094 **Figure 3-figure supplement Table 2: Gene sets enriched in Nrn1[-/-] iTreg cells cultured under**

1095 **the reactivating condition.**

		Ion channel and receptor
Nrn1[-/-]		LIGAND_GATED_MONOATOMIC_CATION_CHANNEL_ACTIVITY
		LIGAND_GATED_MONOATOMIC_ION_CHANNEL_ACTIVITY
		NEUROTRANSMITTER_RECEPTOR_ACTIVITY
		GABA_RECEPTOR_ACTIVITY
		TRANSMITTER_GATED_CHANNEL_ACTIVITY
		INTRACELLULAR_LIGAND_GATED_MONOATOMIC_ION_CHANNEL_ACTIVITY
		G_PROTEIN_COUPLED_AMINE_RECEPTOR_ACTIVITY
		LIGAND_GATED_MONOATOMIC_CATION_CHANNEL_ACTIVITY
		POSTSYNAPTIC_NEUROTRANSMITTER_RECEPTOR_ACTIVITY
		NEUROTRANSMITTER_RECEPTOR_ACTIVITY_INVOLVED_IN_REGULATION_OF_POSTSYNAPTIC_MEMBRANE_POTENTIAL
		LIGAND_GATED_CALCIUM_CHANNEL_ACTIVITY
		Receptor kinase activity
		TRANSMEMBRANE_RECEPTOR_PROTEIN_KINASE_ACTIVITY
		TRANSMEMBRANE_RECEPTOR_PROTEIN_TYROSINE_KINASE_ACTIVITY
ctrl		**Chaperone protein folding**
		ATP_DEPENDENT_PROTEIN_FOLDING_CHAPERONE
		PROTEIN_FOLDING_CHAPERONE
		UNFOLDED_PROTEIN_BINDING
		Translation regulator activity
		TRANSLATION_REGULATOR_ACTIVITY_NUCLEIC_ACID_BINDING
		TRANSLATION_REGULATOR_ACTIVITY

1096 Gene sets involved in Figure 3B clusters enriched in Nrn1[-/-] and ctrl iTreg cells reactivated with

1097 anti-CD3.

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

Figure 3-figure supplement Table 3: Comparison of gene sets enriched in Nrn1[-/-] iTreg cells cultured under the resting and TCR restimulation condition.

Nrn1[-/-]	Resting and TCR restimulation
	TRANSMEMBRANE_RECEPTOR_PROTEIN_KINASE_ACTIVITY
	TRANSMEMBRANE_RECEPTOR_PROTEIN_TYROSINE_KINASE_ACTIVITY
	POSTSYNAPTIC_NEUROTRANSMITTER_RECEPTOR_ACTIVITY
	NEUROTRANSMITTER_RECEPTOR_ACTIVITY_INVOLVED_IN_REGULATION_OF_POSTSYNAPTIC_MEMBRANE_POTENTIAL
	Resting
	SODIUM_CHANNEL_ACTIVITY
	VOLTAGE_GATED_SODIUM_CHANNEL_ACTIVITY
	MONOATOMIC_ANION_MONOATOMIC_CATION_SYMPORTER_ACTIVITY
	SODIUM_ION_TRANSMEMBRANE_TRANSPORTER_ACTIVITY
	MONOATOMIC_ANION_SODIUM_SYMPORTER_ACTIVITY
	SOLUTE_SODIUM_SYMPORTER_ACTIVITY
	ORGANIC_ACID_SODIUM_SYMPORTER_ACTIVITY
	SOLUTE_MONOATOMIC_CATION_SYMPORTER_ACTIVITY
	TCR restimulation
	LIGAND_GATED_MONOATOMIC_CATION_CHANNEL_ACTIVITY
	LIGAND_GATED_MONOATOMIC_ION_CHANNEL_ACTIVITY
	NEUROTRANSMITTER_RECEPTOR_ACTIVITY
	GABA_RECEPTOR_ACTIVITY
	TRANSMITTER_GATED_CHANNEL_ACTIVITY
	INTRACELLULAR_LIGAND_GATED_MONOATOMIC_ION_CHANNEL_ACTIVITY
	G_PROTEIN_COUPLED_AMINE_RECEPTOR_ACTIVITY
	LIGAND_GATED_CALCIUM_CHANNEL_ACTIVITY

Listing of gene sets involved in Figure. 3C Venn diagram, comparisons of gene set cluster components in Nrn1[-/-] under resting vs. TCR restimulation condition.

1121

Figure 3-figure supplement 1. Heatmap of differentially expressed genes and Hallmark gene set enrichment. (A) Heatmap of differentially expressed genes in "GOMF_Metal ion transmembrane transporter activity" gene set from Nrn1[-/-] and ctrl iTreg cells cultured under the resting condition. (B) Heatmap of differentially expressed genes in "GOMF_Metal ion transmembrane transporter activity" gene set from reactivated Nrn1[-/-] and ctrl iTreg cells. (C) Detection of pmTOR and pS6 in Nrn1[-/-] and ctrl iTreg cells. Data represents three independent experiments. **p<0.01. Unpaired Student's t-tests were performed. (D). Enrichment of Hallmark gene set in activated Nrn1[-/-] and ctrl iTreg cells (P<0.05, FDR q<0.25).

1130

Figure 3-figure supplement 2. Characterization of Nrn1$^{-/-}$ naïve CD4 T cells and effect of

Nrn1 blockade on WT iTreg cell differentiation and expansion. (**A**) Resting MP in Nrn1$^{-/-}$ and

ctrl naïve CD4$^+$ T cells. (**B**) AAs induced MP change in Nrn1$^{-/-}$ and ctrl naïve CD4$^+$ T cells. (**C**)

Seahorse analysis of extracellular acidification rate (ECAR) and oxygen consumption rate (OCR)

in Nrn1$^{-/-}$ and ctrl iTreg cells. n=6~10 technical replicates per group. Data represent three

independent experiments. (**D-F**) WT iTreg cells differentiated in the presence of Nrn1 antibody

blockade. **(D).** MP in Nrn1$^{-/-}$ and WT iTreg cells differentiated in the presence or absence of anti-

Nrn1 antibody. **(E).** Live cell number and proportion of Ki67 expressing cells after anti-CD3

restimulation among Nrn1$^{-/-}$ and WT iTreg cells differentiated in the presence or absence of anti-

Nrn1 antibody. **(F).** Foxp3^{+} cell proportion three days after anti-CD3 restimulation in Nrn1$^{-/-}$ and

WT iTreg cells differentiated and restimulated in the presence or absence of anti-Nrn1 antibody.

Data represent three independent experiments. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Ordinary one-way ANOVA was performed for multi-comparison.

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169 **Figure 4-figure supplement Table 1: Gene sets enriched in Nrn1$^{-/-}$ and ctrl Te cells.**

Nrn1$^{-/-}$	**Monoamine transport**
	IMPORT_INTO_CELL
	MONOAMINE_TRANSPORT
	CATECHOLAMINE_TRANSPORT
	NEUROTRANSMITTER_REUPTAKE
	Membrane repolarization
	CELL_CELL_SIGNALING_INVOLVED_IN_CARDIAC_CONDUCTION
	CARDIAC_MUSCLE_CELL_ACTION_POTENTIAL_INVOLVED_IN_CONTRACTION
	MEMBRANE_REPOLARIZATION
	CARDIAC_MUSCLE_CELL_MEMBRANE_REPOLARIZATION
	CARDIAC_MUSCLE_CELL_ACTION_POTENTIAL
	CARDIAC_CONDUCTION
	MEMBRANE_REPOLARIZATION_DURING_CARDIAC_MUSCLE_CELL_ACTION_POTENTIAL
	Cell junction
	CELL_CELL_JUNCTION_ASSEMBLY
	ADHERENS_JUNCTION_ORGANIZATION
Ctrl	**Transporter & Oxidoreduction**
	ACTIVE_TRANSMEMBRANE_TRANSPORTER_ACTIVITY
	PROTON_TRANSMEMBRANE_TRANSPORTER_ACTIVITY
	OXIDOREDUCTASE_ACTIVITY_ACTING_ON_NAD_P_H
	PRIMARY_ACTIVE_TRANSMEMBRANE_TRANSPORTER_ACTIVITY
	OXIDOREDUCTION_DRIVEN_ACTIVE_TRANSMEMBRANE_TRANSPORTER_ACTIVITY
	ELECTRON_TRANSFER_ACTIVITY
	Sensory perception
	SENSORY_PERCEPTION_OF_CHEMICAL_STIMULUS
	SENSORY_PERCEPTION_OF_SMELL

1170 Gene sets involved in Fig. 4F gene sets clusters enriched in Nrn1$^{-/-}$ and ctrl Te cells.

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

Figure 4-figure supplement 1. Heatmap of enriched genes in Te cells. (A) Differentially

expressed genes in "GOMF_Neurotransmitter receptor activity involved in the regulation of

postsynaptic membrane potential" gene set in Nrn1$^{-/-}$ and ctrl Te cells. **(B)** Heatmap of

differentially expressed genes in "MF_metal ion transmembrane transporter activity" in Nrn1$^{-/-}$

and ctrl Te cells. **(C).** Detection of pmTOR and pS6 in Nrn1$^{-/-}$ and ctrl iTreg cells. Data represents

three independent experiments.   **p<0.01, ****p<0.0001. Unpaired Student's t-tests were

performed.

1190

1191

1192

1193 **Reference for supplement figure.**

1194 Plitas, G., C. Konopacki, K. Wu, P.D. Bos, M. Morrow, E.V. Putintseva, D.M. Chudakov, and

1195 A.Y. Rudensky. 2016. Regulatory T Cells Exhibit Distinct Features in Human Breast Cancer.

1196 *Immunity* 45:1122-1134.

1197

1198