Comparative transcriptomic analyses reveal differences in the responses of diploid and triploid eastern oysters to environmental stress during a summer mortality event

Authors:

Rujuta V. Vaidya[1], Sarah Bodenstien[2], Dildorakhon Rasulova[1], Jerome F. La Peyre[3], and Morgan W. Kelly[1]

Contact Information:

1. Department of Biological Sciences, Louisiana State University, Baton Rouge, LA, USA

2. School of Animal Sciences, Louisiana State University Agricultural Center, Baton Rouge, LA 70803, United States of America

Abstract:

Triploid oysters are commonly used as the basis for production in the aquaculture of eastern oysters along the U.S East and Gulf of Mexico coasts. While they are valued for their rapid growth, incidents of triploid mortality during summer months have been well documented in eastern oysters, especially at low salinity sites. Various factors, including parental contribution, gametogenesis failures, as well as cellular and physiological issues stemming from their polyploid state have been proposed as potential causes of triploid mortality, but the exact molecular and physiological mechanisms remain unresolved. We used comparative transcriptomics to examine global transcriptomic responses at the onset of summer mortality (July 2020) in diploid and triploid oysters outplanted to a high and low salinity site, to detect unique gene expression patterns associated with outplant site, maternal broodstock, and ploidy. Diploid dams from three different salinity regimes were used to generate diploid and triploid

oyster lines that were spawned at two hatcheries and were outplanted to high (annual mean salinity = 19.4 ± 6.7) and low (annual mean salinity = 9.3 ± 5.0) salinity sites. Our DEG analyses comparing diploid and triploid oysters detected very few DEGs across the three dams, indicating that diploid parentage wasn't a key contributing factor to triploid mortality. We detected a higher number of DEGs across outplant sites and cohorts, indicating stronger influence of these two factors on triploid mortality. At the low salinity site where triploid oysters experienced high mortality, we observed downregulation of GO terms related to calcium signaling, ciliary activity, and cell cycle check points. These GO terms implicate mis regulation of the low salinity stress response and problems during cell division as key contributors to elevated summer mortality in triploid oysters.

Keywords: *Crassostrea virginica*, ploidy, salinity, gene expression, mortality, Gulf of Mexico

1. Introduction:

Polyploidy is the heritable condition in which a typically diploid cell or organism inherits an additional set (or sets) of chromosomes (de Sousa et al., 2016). Instances of polyploidy occur when chromosomes fail to separate normally during cell division, resulting in gametes that have unequal distribution of chromosomes. Having extra sets of chromosomes affects many structural and physiological aspects of the cell – polyploid individuals are generally bigger in size, have faster growth rates, and offer better yields in commercially valuable species (Piferrer et al., 2009). The prospect of bigger marketable product has generated great interest in the generation of polyploids in agriculture and aquaculture, including in commercially valuable bivalves like the eastern oyster. Generation of triploid and tetraploid eastern oysters was first reported in 1981 (Stanley et al., 1981) and in later years, several standardized protocols for hatchery production of

triploid and tetraploid oysters have been published (Wadsworth et al., 2019; H. Yang, 2022, Hudson, K., 2019) and adapted by oyster farms as their primary basis for production. Compared to the natural diploid oysters, triploid oysters generally grow faster, and have reduced gametogenesis as a consequence of their triploidy (Allen Jr. and Downing, 1986, 1990; Wadsworth et al., 2019a; Matt and Allen, 2021). Unlike diploid oysters, who invest considerable energy reserves in reproduction, triploids oysters instead channel that energy into their growth, resulting in better meat quality during summer (Barber and Mann, 1991; Degremont et al., 2012;Walton et al., 2013). All of these 'triploid advantages' have made them a desirable alternative to natural diploid oysters, and triploids have been adapted by many hatcheries as their primary basis for production (J. Brianik & Allam, 2023)

However, in recent years there have been reports of triploid eastern oysters experiencing higher mortality than diploid oysters during spring and summer months. This phenomenon, collectively termed as 'triploid mortality' has been observed along the US mid-Atlantic and Gulf of Mexico (GoM) coast.(George et al., 2023; Houssin et al., 2019; Matt et al., 2020; Wadsworth et al., 2019). While in some cases the greater mortality of triploid oysters have been clearly associated with low salinity, in other cases, the causes remain unknown. Moreover, the exact underlying cellular mechanisms causing these triploid mortality events remain unresolved. A contributor to triploid mortality in low salinity conditions could be their increased cell sizes, caused by their excess genomic content. These larger cell sizes increase cell volume and volume to surface area ratios, which in turn negatively affects volume regulation under fluctuating salinity (Comai, 2005; Guo & Allen, 1994). It is also possible that altered gene copy numbers in triploids disrupt gene regulation and inhibit adaptive cellular stress responses. This disruption of stress response

pathways has been observed in a variety of other taxa with elevated ploidy, ranging from corn to salmon, and seems a plausible mechanism for increased stress susceptibility in triploid oysters (Auger et al., 2005; Ching et al., 2010) Jamieson et al., 2009). The poor summer performance of triploids could also be influenced by parental stocks used to create them. Eastern oysters show variation in salinity tolerance across the northern Gulf of Mexico (nGOM) and using parents susceptible to salinity stress could lead to production of triploids that perform poorly at low salinity sites (Marshall et al., 2021a; Marshall et al., 2021b; Swam et al., 2022). Studies in the Eastern oysters and Sydney rock oysters have shown that triploids bred from parents selected for desirable traits perform better than triploids bred from parents without those traits (Callam et al., 2016; R. E. Hand et al., 2004).

Lastly, another possible cause of triploid mortality could be cellular damage due to defective reproductive effort in response to environmental spawning cues. The timing of triploid mortality coincides with gametogenesis and the spawning season of diploids and flawed reproductive effort has been proposed as a probable mechanism for triploid mortality in the Pacific oyster, *Crassotrea gigas* (Houssin et al., 2019; Q. Yang et al., 2022). Triploids eastern oysters are not completely sterile, but show reduced gonadal development compared to diploid oysters (Matt & Allen, 2021). It is possible that during summer months, triploid oysters initiate gametogenesis in response to spawning cues, and are unable to complete it owing to either cell division errors because of their odd number of chromosomes, or due to incomplete gonadal development. Reproduction is a costly process for bivalves (J. Brianik & Allam, 2023), and additional challenge of investing in a metabolically costly endeavor in addition to stressful environmental condition could potentially lead to their increased mortality.

Our study focuses on understanding the summer triploid mortality in GoM eastern oysters using comparative transcriptomics. The transcriptome (set of RNA transcripts produced by a cell, tissue, or organism) itself is a phenotype, and can change in response to variation in the environment or developmental stages. Comparative transcriptomic approaches quantify variation in transcriptomes across individuals, environmental conditions, and developmental stages to gain insights into mechanisms generating phenotypic variation and pathways employed to respond to environmental changes (DeBiasse & Kelly, 2016). We sought to compare the global transcriptomic response of diploid and triploid oysters outplanted to a high (annual mean salinity $= 19.4 \pm 6.7$) and low ((annual mean salinity $= 9.3 \pm 5.0$) salinity site, following a significant increase in mortality to detect possible unique gene expression patterns associated with both ploidies. To test for parental contributions to triploid performance, we used diploid dams adapted to three different salinity regimes to generate triploid oyster lines for our experiments. To keep the comparisons uniform, the same diploid brood stocks were used to generate the diploid oyster lines. Both oyster lines were spawned at two hatcheries and were outplanted to intermediate and low salinity sites. Environmental data (salinity and temperature) and morphological measurements including growth, *Perkinsus marinus* (dermo) infection prevalence, and mortality were performed and are explained in Bodenstein et al., 2023. We compared transcriptomic profiles of diploid and triploid oysters to identify differentially expressed genes in response to site, maternal broodstock, and ploidy.

2. Methods:

a) Oyster crossing and outplanting:

The detailed cross strategy and outplanting designs are explained in Bodenstein et al., 2023 and is summarized in Figure 1. Briefly, to establish diploid parental broodstocks adapted to different salinity regimes, oysters were collected from each of three Louisiana public oyster grounds that had different salinity regimes: Calcasieu Lake (CL, annual mean salinity = 16.2 ± 2.8 annual mean salinity of 16.2 ± 2.8 [± standard deviation (SD), n= 10, 2009–2018]), Sister Lake (SL, annual mean salinity = 11.2 ± 5.5), Vermilion Bay (VB, annual mean salinity = 7.4 ± 1.6). The tetraploids used in this study were LSU 4DGNL17 (maintained at LSURF) and AU 4MC18 (maintained at the Grand Bay Oyster Park, Alabama). Oysters were conditioned and spawned in June 2019 LSUL and at the Auburn University Shellfish Laboratory (AUSL) in Dauphin Island, AL in July 2019 as described in Bodenstein et al., 2023. Depending on where they were spawned, F1 progeny was sorted into LSU (Louisiana State University) or AU (Auburn) cohorts. A total of six crosses were produced at each hatchery. The diploid F1 crosses were produced by crossing male and female oysters from each wild broodstock population (CL, SL, VB) and the triploid F1 crosses were produced by crossing females from wild broodstock populations with males from two tetraploid lines (one from each hatchery, Fig. 1). Diploid and triploid crosses were generated from the same females at the Auburn hatchery and were therefore half-siblings. However, at the LSU hatchery, diploid and triploid crosses were not half-siblings and, in certain cases, were produced on separate days. Oysters were reared and transported to their final outplant sites as described in Bodenstein et al., 2023. Oysters from both cohorts were outplanted to a low (near the Louisiana Universities Marine Consortium (LUMCON), annual mean salinity = 9.3 ± 5.0) and a moderate salinity site (LSURF, annual mean salinity = 19.4 ± 6.7) in Louisiana. Oysters from Auburn cohort were outplanted in November 2019 at the two sites and oysters from LSU cohort were outplanted at LSURF in December 2019 and at LUMCON in

January 2020. At each site, four replicate baskets containing 80 oysters each were outplanted for each of the six crosses of each ploidy in the two cohorts, a total of 48 baskets per site. Growth and mortality measurements were performed monthly till November 2020 and are described in Bodenstein et al., 2023.

b) RNA sequencing:

On 8[th] July, 2020, eight individuals per cross were sampled at both outplant sites. Approximately 0.5-cm^2 piece of gill tissue was dissected in the field from each individual and preserved with Invitrogen RNAlater. Total RNA was extracted using an E.Z.N.A. Total RNA Kit I (Omega BIO-TEK; VWR catalog no. 101319) following the manufacturer's instructions. The yield and quantity were assessed using a NanoDrop 2000 spectrophotometer. Total RNA extracted from the 192 individuals were sent to the University of Texas at Austin's Genomic Sequencing and Analysis Facility where RNA quality control was confirmed using a 2100 Agilent Bioanalyzer on a Eukaryote Total RNA Nano chip and libraries were produced using the Tag-sequencing approach (Meyer et al., 2011). The resulting 192 libraries were sequenced equally across two lanes of an Illumina HiSeq 2500 platform, with 100-bp single-end reads. Adapter sequences were trimmed using trimmomatic (version 0.39) (Bolger et al., 2014) and base pairs with quality scores below 35 were removed. The trimmed reads were then mapped to the *C. virginica* reference genome (Gómez-Chiarri et al., 2015) with known haplotigs removed (https://github.com/jpuri tz/ Oyste rGenomeProject/tree/maste r/Haplo tig_Masked_Genome) using star rna-seq aligner (single pass option). Reads were mapped to gene features with the options (-- quantMode GeneCounts –outFilterScoreMinOverLread 0.50 --

outFilterMatchNminOverLread 0.50) to adjust for poly-A tail contamination (Sirovy et al., 2021) to finally generate a count matrix using ReadsPerGene.out.tab output (supplementary file 1).

c) Differential Gene Expression (DEG), Weighted Gene Correlation Network Analysis (WGCNA), and aneuploidy analysis:

To maximize the true positive rate and specificity, we used both parametric and non-parametric tests to obtain a single consensus list of DEGs (Costa-Silva et al., 2017) for our pairwise comparisons. DEGs were calculated using DESeq2 (v 1.24.0) (Love et al., 2014), EdgeR (Robinson et al., 2010), and NOISeq (Tarazona et al., 2011) and the R package VennDiagram was used to find consensus lists of DEGs detected by all three methods for all comparisons. DEGs were identified using a false discovery rate (FDR)-adjusted p-value cutoff of 0.01. DEGs were calculated for pairwise comparisons comparing ploidy, site, cohort, and dams. Gene ontology enrichment was also performed using GO_MWU (Wright et al., 2015), using the Fisher's Exact Test ($p<0.05$) the scripts available at:

https://github.com/z0on/GO_MWU/blob/maste r/GO_MWU.R. The R package WGCNA (Langfelder & Horvath, 2008) was used to identify clusters of genes with highly correlated expression patterns. We normalized the data using DESeq2 variance stabilizing transformation (VST) function and filtered out genes with fewer than five counts per million in 80% of all samples. The WGCNA was run using 9172 genes that passed this criterion, using a soft threshold value of 12, a minimum module size of 30, and a signed adjacency matrix. The network was then correlated to site, cohort, ploidy, and dams variables converted to binary format. Module eigengenes, defined as first principle component of a given module (Langfelder & Horvath, 2008) were visualized using ggplot2 and GO enrichment analysis was conducted using

GO_MWU. To measure the magnitude of the difference between transcriptomic responses of our diploid and triploid oysters, we weighted the samples by the percent of variation explained by each axis and then calculated the shift (Euclidean distances) in multivariate space between diploid and triploid samples matched for their cohort, site, and dams. We also assessed for instances of aneuploidy across all 10 chromosomes. To do that, we first categorized our samples into diploid and triploid pairs by matching their site, cohort, and dam, i.e. a diploid oyster from Auburn cohort and VB dam broodstock and was outplanted to Grand Isle was paired with a triploid oyster from Auburn cohort and VB dam broodstock and was outplanted to Grand Isle. For every pair, we then plotted normalized log transformed Counts Per Millions (CPM) values such that each point on the plot represented a transcript and X and Y coordinates of every point were CPM values of that transcript in the diploid and triploid oyster respectively. For each pair compared, ten such plots (one for each chromosome) were made, and the slope of the regression line was calculated. As the CPM values were normalized to the library sizes, change in the slope for any chromosome for a particular sample pair was considered as an indicator of aneuploidy occurring at that chromosome.

3. Results:

3.2 Principal Component Analysis and aneuploidy analysis:

Detailed mortality and growth measurements are described in Bodenstein et al.,2023. Overall, triploids showed higher mortality at low salinity site (LUMCON) (Fig. 2). At both high and low salinity sites, the cumulative mortality of triploids was higher than diploids, and triploids from LSU cohort experienced higher cumulative mortality than AU cohort. Maternal stock (dams) did not influence cumulative mortality of triploids at either site.

The first two principal component axes accounted for 61% and 6% variance in samples respectively (Fig. 3a). Out of the four variables, outplant site had the greatest effect on sample clustering, followed by cohort and ploidy. No clustering was observed based on dams. The difference in the magnitude of transcriptomic response, calculated as the Euclidean distances between points in the PCA representing diploid and triploid oysters matched for their cohort, site, and dams, was largest in the LSU cohort at the LUMCON site (fig. 3b). Overall, the difference was higher in diploid vs triploid oyster comparisons at LUMCON site, regardless of their cohorts. To check if the differences in mortality were a result of aneuploidy in triploid oysters, we plotted normalized log transformed CPM values of diploid and triploid oysters on X and Y axis respectively. Slope values for all ten chromosomes for each sample pair are summarized in supplementary table 1 . We do not see change in slopes of any chromosomes for any of our sample pairs.

3.3 DEG and WGCNA:

Differentially expressed genes (FDR<0.05) were calculated using pairwise comparisons in EDGER, NOISeq, and DESeq2. DEG lists from all three approaches were compared and only genes that were detected by all three approaches were used as DEGs for downstream analysis. The results of our DEG analyses are summarized in Table 1. In all the comparisons performed, the highest number of DEGs (9022) were obtained for the pairwise comparison between triploid oysters from Auburn and LSU cohort at low salinity site. All pairwise comparisons for dams (within and among ploidy) yielded some of the lowest number of DEGs. For our first set of comparisons, we compared expression profiles of diploid and triploid oysters, focusing on one experimental variable at a time. For example, to compare transcriptomic response of diploid and

triploid oysters at Grand Isle outplant site, we compared expression profiles of diploid and triploid oysters outplanted to Grand Isle site, regardless of their cohorts and dams. Similar comparison was made for the LUMCON site as well. To parse out cohort specific differences, we also compared diploid and triploid oysters from two cohorts similarly, this time regardless of their site and dams. Lastly, similar comparisons were performed for each of the three dams. Overall, we observe higher number of DEGs for LUMCON site (1233) and LSU cohort (867), and dams yielded some of the lowest number of DEGs across all comparisons. Functional enrichment for these DEGs showed 8 significantly upregulated GO terms for LUMCON site, related to calcium ion binding and microtubule based process. On the other hand, no GO terms were detected for Grand Isle site, LSU and AU cohort and the three dams.

For our next set of comparisons, we compared expression profiles of diploid and triploid oysters belonging to a specific site and cohort to check for potential site or cohort specific expression patterns that may have been obscured in our first set of comparisons. For LUMCON site, we detected higher DEGs for oysters from LSU cohort (1812) than their Auburn counterparts (709). GO analysis for LSU + LUMCON DEGs revealed upregulation of GO terms related to microtubules, cell cycle, and calcium binding by triploid oysters. Oysters from LSU cohorts also showed higher DEGs at Grand Isle site (286) than their Auburn counterparts (125). GO analysis performed on LSU + GI DEGs revealed downregulation of GO terms related to reactive oxygen species and superoxide metabolic process by triploid oysters. GO analysis for Auburn + LUMCON and Auburn + GI DEGs yielded no GO terms. For our last set of comparisons, we compared ploidy-specific expression patterns of diploid and triploid oysters separately for each site and cohort. Overall, triploid oysters showed a higher number of DEGs across sites and

cohorts than diploid oysters. Across two outplant sites, triploid oysters downregulated GO terms related to cell cycle regulation and amino acid metabolism and upregulated GO terms related to ion transport. Across two cohorts, triploid oysters upregulated GO terms related to apoptosis and downregulated GO terms related to DNA replication.

We also performed Weighted Gene Correlated Network Analysis (WGCNA) to identify groups of genes displaying similar expression patterns in response to site, ploidy, cohort, and dams. Fig. 4a shows 11 modules labeled as color names and the degree of their correlation with traits of interest. We found five (grey60, brown, greenyellow, red, and blue) modules associated with site and two (midnightblue and blue) with cohort. The module blue showed significant association with both site and cohort. Fig. 4b shows eigengene expression plots for brown, grey60, red, and blue module with lines indicating samples grouped by their ploidies and cohorts, revealing ploidy and cohort specific expression patterns in these modules. The blue module was negatively correlated with site and cohort and contained 2209 genes that were downregulated by oysters from LUMCON site and LSU cohort. GO analysis for this module revealed an overrepresentation of GO terms related to calcium ion binding and actin binding. The brown module showed strong negative correlation for site, i.e. genes from this module were downregulated by oysters from LUMCON site when compared with oysters from Grand Isle. GO analysis revealed enrichment of GO terms related to plasma membrane, lipid binding, cell adhesion, calcium binding, and extracellular matrix. On the other hand, Grey60 module contained 119 genes and was positively correlated to site. Genes from this module were upregulated by oysters from LUMCON when compared with Grand Isle. Go analysis for this module revealed overrepresentation of GO terms related to TCA cycle and cysteine type

endopeptidase inhibitor activity among others. Red module (2060 genes) also showed positive correlation to site and genes from this module were upregulated by oysters from LUMCON site when compared with oysters from Grand Isle. GO terms overrepresented in this module were related to initiation and regulation of translation, calcium ion binding, cytoskeleton protein binding, and cellular response to stimulus.

4. Discussion:

The primary aim of our study was to understand molecular and physiological causes of elevated summer mortality in triploid oysters. Contrary to our hypothesis, our DEG analyses comparing diploid and triploid oysters detected very few DEGs across the three dams, indicating that maternal stocks used to breed oysters weren't the key factor contributing to differences in their summer mortality. This lack of influence of maternal stock on triploid oysters was also observed in the growth and morphological measurements performed on these oysters by Bodenstein et al 2023. Our analyses also did not detect instances of aneuploidy, which has previously been suggested as a cause of problems in polyploids. On the other hand, comparisons of diploid and triploid oysters across outplant sites and cohorts revealed a significant number of DEGs and GO terms related to key biological processes, indicating the stronger influence of these two factors on summer mortality. Our inter - and intra - ploidy analyses reveal that diploid and triploid oysters share at least some of their cellular response to low salinity and summer conditions, as seen through enrichment of common GO terms and transcripts. The results from DEG comparisons also match with the correlation patterns observed in WGCNA - the modules significantly associated with site and cohort were enriched for biological processes detected in DEG comparisons.

Intra-ploidy variation in transcriptomic response to salinity/outplant sites:

Our within ploidy comparisons, performed separately for diploid and triploid oysters revealed that cell division was significantly affected in both diploid and triploid oysters at the low salinity site, which experienced greater mortality. We observe that at low salinity site, diploid as well as triploid oysters from the higher mortality LSU cohort upregulate GO terms enriched for DNA replication and repair, cell cycle regulation, and response to stimulus. Presence of these common GO terms indicates that regardless of their ploidies, both diploid and triploid oysters respond to low salinity cues by altering or arresting their cell division. Upregulation of genes involved in cell cycle checkpoints in the gill tissue has been previously reported in mussels (Lockwood & Somero, 2011) and killfish (Kültz & Avila, 2001; Whitehead et al., 2011), indicating that cell cycle regulation is a conserved response to low salinity across species. Changes in the salinity have multi-faceted impact on a cell – they can change the osmotic balance inside the cell, activate the cellular signaling pathways that detect changes in salinity, and affect structure and functions of proteins and nucleic acids in the cell. Oysters are osmoconformers, however ability to osmoconform to abrupt and large changes in salinity, however, may take days and is a function of the initial salinity (i.e., salinity oysters are held prior to a change is salinity) and the rate, magnitude and direction of the salinity change (Shumway, 1996; S. C. Hand & Stickle, 1977; M. La Peyre et al., 2003; M. K. La Peyre et al., 2009; McFarland et al., 2013).

Therefore, response to salinity stress has been proposed to be a complex process that targets multiple aims including the initiation of a damage control response to repair the protein and DNA damage caused by salinity change, and also initiating a response to remove molecules or

cells that have been damaged beyond repair (Hochachka and Somero, 2002). The enrichment of DNA repair GO terms we see for LSU cohort therefore suggests that they were employing similar stress response mechanisms to combat low salinity stress. Arresting growth during salinity stress is a way of ensuring that damaged DNA isn't passed down in the cell division cycles – it gives the cell enough time to repair the DNA damage and then progress to cell division. The upregulation of cell cycle checkpoints, which are essentially 'quality check' points to ensure the structural integrity on the DNA before progressing to the next step, are also an indicator of LSU triploids coping with low salinity stress.

This common response across ploidies presents an interesting case as in our next set of comparisons, we observe that when compared to each other, triploids downregulate these GO terms when compared with the diploids. The shared transcriptomic response to salinity and its decreased magnitude in triploids, could mean that triploids mount a weaker adaptive response to salinity, resulting in greater cellular damage.

Along with these shared patterns, we also observed some ploidy-specific gene expression patterns for triploid oysters. At LUMCON, triploid oysters from the LSU cohort upregulated GO terms and transcripts involved in immune response compared to their Auburn counterparts. Even though both cohorts experienced dermo infection, LSU cohort had a higher number of infected individuals (higher prevalence) which could explain this LSU-specific upregulation.

Inter-ploidy variation in transcriptomic response to salinity/outplant sites:

For our next set of comparisons, we computed differentially expressed genes between diploid and triploid oysters for each site, cohort, and dams. To account for the potential effects of each cohort and site, we also compared DEGs between diploid and triploid oysters from the same cohort for each site. Despite finding many DEGs, we did not find any GO terms for oysters from LSU cohort at Grand Isle and for oysters from Auburn cohort at LUMCON. Absence of GO terms in these comparisons could be interpreted as absence of any known GO terms enriched for DEGs found in these comparisons. The remaining two categories did yield DEGs as well as GO terms. At Grand Isle, triploid oysters from AUBURN cohort upregulated GO terms related to reactive oxygen species metabolic process and superoxide metabolic process.

At LUMCON, triploid oysters from the LSU cohort downregulated genes involved in three key processes - calcium signaling, ciliary activity, and cell cycle checkpoints. This cohort specific response was also seen by Bodenstein et al 2023 in their mortality measurements performed on these oysters – mortality levels for triploids from LSU cohort were highest at low salinity site. The GO terms and transcripts enriched for these processes hints at some degree of crosstalk among these three processes, suggesting a coordinated stress response in response to salinity, led by calcium signaling. Intracellular calcium is noted for its role as a messenger molecule that facilitates intra and intercellular communication. Specifically, increase in intracellular calcium in response to abiotic stress is a hallmark of plant stress response, and a similar increase in calcium and its receptors (partners) in response to ocean acidification has been noted in the Pacific oyster as well (Xin et al., 2022). Unlike their diploid counterparts, triploid oysters were unable to upregulate genes involved in calcium signaling, indicating triploid-specific problems in detecting and responding to low salinity stress. In our study, triploid oysters from LSU cohort downregulated several key members of calcium-mediated stress response pathway such as

Calmodulin-a, Histidine-rich calcium binding protein (HRC), and cadherin EGF LAG seven-pass G-type receptor 1(CELSR1). Calmodulin-a, is a well-studied calcium-binding protein that influences key physiological processes like cell proliferation, differentiation, and apoptosis by interacting with its many downstream targets (Xin et al., 2022). Histidine-rich calcium binding protein (HRC), which is important for calcium homeostasis and cadherin EGF LAG seven-pass G-type receptor 1(CELSR1), a type of G protein-coupled transmembrane receptors instrumental for communication between extracellular and intracellular environment, were also downregulated by triploid oysters (Arvanitis et al., 2011; Wang et al., 2014).

Furthermore, genes encoding calcium dependent proteins responsible for structural elements of intra and extracellular space like fibrillin1 & 2, annexin A4, cadherin-23 were also downregulated by triploid oysters. Changes in cell shape and volume do occur in response to salinity, and downregulation of these cytoskeleton elements by triploid oysters may have hindered their effort to maintain their cell shape and osmotic balance under low salinity conditions (Zhang et al., 2017), Hochachka and Somero, 2002). Taken together, downregulation of these calcium related genes shows that low salinity had a domino-like effect on triploid oysters, affecting several key biological processes in them.

The decrease in ciliary activity was marked by downregulation of numerous genes coding for cytoplasmic and axonemal dyneins. As one of the three major cytoskeletal motor proteins, cytoplasmic and axonemal dyneins have been shown to be associated with sensory and motility functions of the cilia, respectively. Upregulation of transcripts involved in ciliary activity at low salinity has been observed in diploid *C. virginica*, and has been hypothesized to help in maintaining homeostasis and avoiding valve closure (Jones et al., 2019). The lack of upregulation of these genes in triploid oysters, combined with their increased mortality at the low

salinity site hints at sub-optimal ciliary function in their gill tissue. Additionally, morphological measurements performed on these oysters by Bodenstein et al reveal that triploids have larger gill area than diploids, after correcting for their size differences. Bigger cell size and cell volume of triploid cells has been shown to cause disruption of ion channels and decrease their sensitivity to environment, and negatively affect volume regulation under fluctuating salinity (Guo & Allen, 1994; Haim-Abadi et al., 2023). It is possible that the difference in gill area is due to triploids having larger cells than diploids and is in turn at least partially responsible for disruption of their cellular homeostasis under low salinity stress. Changes in ciliary activity as a response to salinity stress have been observed in marine bivalves (Johnson et al., 2021; Maynard et al., 2018), although we are unclear as to why we observe this for only one cohort of triploid oysters in our study. It is possible that triploid oysters from LSU cohort were more severely affected by salinity due to differences in their rearing conditions as compared to their Auburn counterparts.

Along with ciliary activity, triploid oysters from LSU cohort also downregulated genes involved in cell cycle and its regulation. Genes such as KIF2A, KIF3C, NEK4, NEK6, CDK1, TNKS1, HAUS augmin-like complex subunit 3, and MAPKKK which play key roles in different stages of cell cycle including but not limited to maintenance of chromosome structure, DNA replication and repair, formation and maintenance of spindle fiber, normal progression of cell division, and cell cycle checkpoints (Chang & Karin, 2001; Fry et al., 2012; Hsiao & Smith, 2008; Kültz & Burg, 1998; Lawo et al., 2009; Sheng et al., 2018) were downregulated by triploids at low salinity site. Genes with regulatory functions during cell division like MCM7, which ensures that the replication of chromosomal DNA occurs only once before cell division and CHTF18, which ensures fidelity of transmitting chromosomes from one generation to next were also downregulated in triploid oysters (Berkowitz et al., 2012; Onesti & MacNeill, 2013). Lastly,

genes involved in DNA repair (SMCs and RTEL1) were also downregulated by triploid oysters at low salinity site.

Downregulation of several key players involved in cell cycle reveals/highlights/indicates that gill tissue cells of triploid oysters were encountering errors at multiple stages of cell division. As mentioned in the section about intra-ploidy comparisons, upregulation of genes involved in cell cycle was a shared transcriptomic response observed in both ploidies, but its magnitude appears to be lower in triploid oysters when compared to diploids. This difference in magnitudes could be due to polyploid nature of triploid oysters. Cell division in polyploid cells, especially those with odd number of chromosomes such as triploids is prone to errors such as chromosome loss and aneuploid cells (Comai, 2005). Additionally, progression through cell cycle does depend on environmental stimuli like thermal and osmotic stress (Hochachka and Somero, 2002). Taken together, it appears that the triploid oysters at low salinity sites were particularly challenged with the dual threat of unfavorable environmental conditions and unfavorable cellular milieu for optimal cell division, and therefore had higher mortality rates than diploids.

The cohort effect in our interploidy comparisons could be attributed to differences in the two cohorts. At the low salinity site, field mortality measurements from Bodenstein et al., 2023 show higher cumulative mortalities for triploids from LSU cohort, but do not see a cohort-specific effect on diploid mortality. The cohort effect was also observed in growth rates – at low salinity sites, both diploids and triploids from LSU cohort had slower growth rates. Since the point of distinction between two cohorts was hatchery conditions, we think that early life experience could be influencing growth and survival in later life. Stressors present in the early life environment of oysters have been shown to have an impact on their later life survival (Donelan et al., 2021; Hettinger et al., 2012; Spencer et al., 2020), and the cohort specific differences in

growth rates and survival we see could be resulting due to different rearing conditions experienced by oyster larvae at the two hatcheries.

The elevated mortality of triploids oysters observed in our study appears to be a result of their cellular defects that were exacerbated by unfavorable environmental conditions. Triploid oysters have been shown to lose their growth advantage over diploids in stressful environments (Callam et al., 2016; Davis, 1994; Wadsworth et al., 2019b) and are also more vulnerable to multi-stressor scenarios (George et al., 2023; J. Brianik & Allam, 2023; Li et al., 2022). Additionally, as our sampling time was during peak summer (July 2020), both diploid and triploid oysters in our study were exposed to environmental spawning cues. It is possible that triploid oysters in our study responded to spawning cues by investing in gametogenesis, and the addition of this metabolically costly process while experiencing abiotic stress threw their energy budget into a disarray. As we used gill tissue for our transcriptomics analyses, we were unable to detect gene expression changes related to reproduction in our experiment. Future transcriptomic studies using gonadal tissue samples will be helpful to detect transcriptomic signatures of reproductive effort in triploid oysters. Furthermore, triploid oysters in our study did show gonadal development but were not tested for formation of viable gametes. This distinction is important, as previous studies have shown that even though triploid *C. virginica* develops gonads, it fails to produce viable gametes and shows errors in synchrony of gametogenesis (Matt & Allen, 2021). The cost of this unsuccessful reproductive effort and its impact on survival of triploid oysters remains unexplored. Future studies aimed at exploring tradeoffs of this imperfect gametogenesis in triploids will be helpful in clarifying the contribution of reproduction in triploid mortality. Lastly, previous studies have shown occurrences of aneuploidy and subsequent chromosome loss in triploid oysters, both of which can cause cell death by disrupting normal patterns of gene

expression (De Sousa et al., 2023). Even though we did not detect any incidents of aneuploidy, our test was an indirect measure and was performed on gill tissues of the diploid and triploid oysters. Future studies involving direct measures like karyotyping gonadal cells of triploid oysters across their developmental stages will be helpful in understanding whether these genomic irregularities exist and contribute to their mortality.

Conflict of interest: The authors declare that they have no conflict of interest.

Acknowledgements: We thank Dr. Brian Callam, F. Scott Rikard, and staff members of the Michael Voisin Oyster Research Lab and Hatchery and Auburn University Shellfish Lab for producing the oysters used in this study.

Funding information: This work was partly supported by Sea Grant Marine Aquaculture Grant Program (NA18OAR4170350) to J. La Peyre with additional support provided by National Institute of Food and Agriculture, United States Department of Agriculture (Hatch project, LAB94509).

References:

Arvanitis, D. A., Vafiadaki, E., Sanoudou, D., & Kranias, E. G. (2011). Histidine-rich calcium binding protein: The new regulator of sarcoplasmic reticulum calcium cycling. *Journal of Molecular and Cellular Cardiology*, *50*(1), 43–49. https://doi.org/10.1016/j.yjmcc.2010.08.021

Auger, D. L., Gray, A. D., Ream, T. S., Kato, A., Coe, E. H., & Birchler, J. A. (2005). Nonadditive Gene Expression in Diploid and Triploid Hybrids of Maize. *Genetics*, *169*(1), 389–397. https://doi.org/10.1534/genetics.104.032987

Berkowitz, K. M., Sowash, A. R., Koenig, L. R., Urcuyo, D., Khan, F., Yang, F., Wang, P. J.,
Jongens, T. A., & Kaestner, K. H. (2012). Disruption of Chtf18 Causes Defective Meiotic
Recombination in Male Mice. *PLoS Genetics*, *8*(11), e1002996.
https://doi.org/10.1371/journal.pgen.1002996

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina
sequence data. *Bioinformatics*, *30*(15), 2114–2120.
https://doi.org/10.1093/bioinformatics/btu170

Callam, B. R., Allen, S. K., & Frank-Lawale, A. (2016). Genetic and environmental influence on
triploid Crassostrea virginica grown in Chesapeake Bay: Growth. *Aquaculture*, *452*, 97–
106. https://doi.org/10.1016/j.aquaculture.2015.10.027

Chang, L., & Karin, M. (2001). Mammalian MAP kinase signalling cascades. *Nature*,
410(6824), 37–40. https://doi.org/10.1038/35065000

Ching, B., Jamieson, S., Heath, J. W., Heath, D. D., & Hubberstey, A. (2010). Transcriptional
differences between triploid and diploid Chinook salmon (Oncorhynchus tshawytscha)
during live Vibrio anguillarum challenge. *Heredity*, *104*(2), 224–234.
https://doi.org/10.1038/hdy.2009.108

Comai, L. (2005). The advantages and disadvantages of being polyploid. *Nature Reviews
Genetics*, *6*(11), 836–846. https://doi.org/10.1038/nrg1711

Costa-Silva, J., Domingues, D., & Lopes, F. M. (2017). RNA-Seq differential expression
analysis: An extended review and a software tool. *PLOS ONE*, *12*(12), e0190152.
https://doi.org/10.1371/journal.pone.0190152

de Sousa, J. T., Allen, S. K., Baker, H., & Matt, J. L. (2016). Aneuploid progeny of the
American oyster, *Crassostrea virginica* , produced by tetraploid × diploid crosses:

Another example of chromosome instability in polyploid oysters. *Genome*, *59*(5), 327–

338. https://doi.org/10.1139/gen-2015-0222

DeBiasse, M. B., & Kelly, M. W. (2016). Plastic and Evolved Responses to Global Change:

What Can We Learn from Comparative Transcriptomics?: Table 1. *Journal of Heredity*,

107(1), 71–81. https://doi.org/10.1093/jhered/esv073

Donelan, S. C., Breitburg, D., & Ogburn, M. B. (2021). Context-dependent carryover effects of

hypoxia and warming in a coastal ecosystem engineer. *Ecological Applications*, *31*(4),

e02315. https://doi.org/10.1002/eap.2315

Fry, A. M., O'Regan, L., Sabir, S. R., & Bayliss, R. (2012). Cell cycle regulation by the NEK

family of protein kinases. *Journal of Cell Science*, jcs.111195.

https://doi.org/10.1242/jcs.111195

George, M. N., Cattau, O., Middleton, M. A., Lawson, D., Vadopalas, B., Gavery, M., &

Roberts, S. B. (2023). Triploid Pacific oysters exhibit stress response dysregulation and

elevated mortality following heatwaves. *Global Change Biology*, gcb.16880.

https://doi.org/10.1111/gcb.16880

Gómez-Chiarri, M., Warren, W. C., Guo, X., & Proestou, D. (2015). Developing tools for the

study of molluscan immunity: The sequencing of the genome of the eastern oyster,

Crassostrea virginica. *Fish & Shellfish Immunology*, *46*(1), 2–4.

https://doi.org/10.1016/j.fsi.2015.05.004

Guo, X., & Allen, S. K. (1994). Reproductive Potential and Genetics of Triploid Pacific Oysters,

Crassostrea gigas (Thunberg). *The Biological Bulletin*, *187*(3), 309–318.

https://doi.org/10.2307/1542288

Haim-Abadi, G., Golan-Lev, T., Koren, A., & Benvenisty, N. (2023). Generation, genomic characterization, and differentiation of triploid human embryonic stem cells. *Stem Cell Reports*, *18*(5), 1049–1060. https://doi.org/10.1016/j.stemcr.2023.04.001

Hand, R. E., Nell, J. A., & Thompson, P. A. (2004). Studies on triploid oysters in Australia. *Aquaculture*, *233*(1–4), 93–107. https://doi.org/10.1016/j.aquaculture.2003.09.017

Hand, S. C., & Stickle, W. B. (1977). Effects of tidal fluctuations of salinity on pericardial fluid composition of the American oyster Crassostrea virginica. *Marine Biology*, *42*(3), 259–271. https://doi.org/10.1007/BF00397750

Hettinger, A., Sanford, E., Hill, T. M., Russell, A. D., Sato, K. N. S., Hoey, J., Forsch, M., Page, H. N., & Gaylord, B. (2012). Persistent carry-over effects of planktonic exposure to ocean acidification in the Olympia oyster. *Ecology*, *93*(12), 2758–2768. https://doi.org/10.1890/12-0567.1

Houssin, M., Trancart, S., Denechere, L., Oden, E., Adeline, B., Lepoitevin, M., & Pitel, P.-H. (2019). Abnormal mortality of triploid adult Pacific oysters: Is there a correlation with high gametogenesis in Normandy, France? *Aquaculture*, *505*, 63–71. https://doi.org/10.1016/j.aquaculture.2019.02.043

Hsiao, S. J., & Smith, S. (2008). Tankyrase function at telomeres, spindle poles, and beyond. *Biochimie*, *90*(1), 83–92. https://doi.org/10.1016/j.biochi.2007.07.012

J. Brianik, C., & Allam, B. (2023). The need for more information on the resistance to biological and environmental stressors in triploid oysters. *Aquaculture*, *577*, 739913. https://doi.org/10.1016/j.aquaculture.2023.739913

Johnson, K. M., Jones, H. R., Casas, S. M., La Peyre, J. F., & Kelly, M. W. (2021). Transcriptomic signatures of temperature adaptation in the eastern oyster *Crassostrea*

virginica. *Journal of Evolutionary Biology*, *34*(8), 1212–1224.

https://doi.org/10.1111/jeb.13789

Jones, H. R., Johnson, K. M., & Kelly, M. W. (2019). Synergistic Effects of Temperature and

Salinity on the Gene Expression and Physiology of Crassostrea virginica. *Integrative and*

Comparative Biology, *59*(2), 306–319. https://doi.org/10.1093/icb/icz035

Kültz, D., & Avila, K. (2001). Mitogen-activated protein kinases are in vivo transducers of

osmosensory signals in fish gill cells. *Comparative Biochemistry and Physiology Part B:*

Biochemistry and Molecular Biology, *129*(4), 821–829. https://doi.org/10.1016/S1096-

4959(01)00395-5

Kültz, D., & Burg, M. (1998). Evolution of Osmotic Stress Signaling *Via* Map Kinase Cascades.

Journal of Experimental Biology, *201*(22), 3015–3021.

https://doi.org/10.1242/jeb.201.22.3015

La Peyre, M. K., Gossman, B., & La Peyre, J. F. (2009). Defining Optimal Freshwater Flow for

Oyster Production: Effects of Freshet Rate and Magnitude of Change and Duration on

Eastern Oysters and Perkinsus marinus Infection. *Estuaries and Coasts*, *32*(3), 522–534.

https://doi.org/10.1007/s12237-009-9149-9

La Peyre, M., Nickens, A., Volety, A., Tolley, G., & La Peyre, J. (2003). Environmental

significance of freshets in reducing Perkinsus marinus infection in eastern oysters

Crassostrea virginica: Potential management applications. *Marine Ecology Progress*

Series, *248*, 165–176. https://doi.org/10.3354/meps248165

Langfelder, P., & Horvath, S. (2008). WGCNA: An R package for weighted correlation network

analysis. *BMC Bioinformatics*, *9*(1), 559. https://doi.org/10.1186/1471-2105-9-559

Lawo, S., Bashkurov, M., Mullin, M., Ferreria, M. G., Kittler, R., Habermann, B., Tagliaferro, A., Poser, I., Hutchins, J. R. A., Hegemann, B., Pinchev, D., Buchholz, F., Peters, J.-M., Hyman, A. A., Gingras, A.-C., & Pelletier, L. (2009). HAUS, the 8-Subunit Human Augmin Complex, Regulates Centrosome and Spindle Integrity. *Current Biology*, *19*(10), 816–826. https://doi.org/10.1016/j.cub.2009.04.033

Li, Y., Jiang, K., & Li, Q. (2022). Comparative transcriptomic analyses reveal differences in the responses of diploid and triploid Pacific oysters (Crassostrea gigas) to thermal stress. *Aquaculture*, *555*, 738219. https://doi.org/10.1016/j.aquaculture.2022.738219

Lockwood, B. L., & Somero, G. N. (2011). Transcriptomic responses to salinity stress in invasive and native blue mussels (genus Mytilus): TRANSCRIPTOMES OF INVASIVE VS. NATIVE MUSSELS. *Molecular Ecology*, *20*(3), 517–529. https://doi.org/10.1111/j.1365-294X.2010.04973.x

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550. https://doi.org/10.1186/s13059-014-0550-8

Matt, J. L., & Allen, S. K. (2021). A classification system for gonad development in triploid Crassostrea virginica. *Aquaculture*, *532*, 735994. https://doi.org/10.1016/j.aquaculture.2020.735994

Matt, J. L., Guévélou, E., Small, J. M., & Allen, S. K. (2020). A field test investigating the influence of brood stock origin and ploidy on the susceptibility of Crassostrea virginica to "triploid mortality" in the Chesapeake Bay. *Aquaculture*, *526*, 735375. https://doi.org/10.1016/j.aquaculture.2020.735375

Maynard, A., Bible, J. M., Pespeni, M. H., Sanford, E., & Evans, T. G. (2018). Transcriptomic

    responses to extreme low salinity among locally adapted populations of Olympia oyster (

    Ostrea lurida). *Molecular Ecology*, *27*(21), 4225–4240.

    https://doi.org/10.1111/mec.14863

McFarland, K., Donaghy, L., & Volety, A. (2013). Effect of acute salinity changes on

    hemolymph osmolality and clearance rate of the non-native mussel, Perna viridis, and the

    native oyster, Crassostrea virginica, in Southwest Florida. *Aquatic Invasions*, *8*(3), 299–

    310. https://doi.org/10.3391/ai.2013.8.3.06

Meyer, E., Aglyamova, G. V., & Matz, M. V. (2011). Profiling gene expression responses of

    coral larvae (Acropora millepora) to elevated temperature and settlement inducers using a

    novel RNA-Seq procedure: RNA-Seq EXPRESSION PROFILING CORAL LARVAE.

    Molecular Ecology, no-no. https://doi.org/10.1111/j.1365-294X.2011.05205.x

Onesti, S., & MacNeill, S. A. (2013). Structure and evolutionary origins of the CMG complex.

    Chromosoma, *122*(1–2), 47–53. https://doi.org/10.1007/s00412-013-0397-x

Piferrer, F., Beaumont, A., Falguière, J.-C., Flajšhans, M., Haffray, P., & Colombo, L. (2009).

    Polyploid fish and shellfish: Production, biology and applications to aquaculture for

    performance improvement and genetic containment. *Aquaculture*, *293*(3–4), 125–156.

    https://doi.org/10.1016/j.aquaculture.2009.04.036

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: A Bioconductor package for

    differential expression analysis of digital gene expression data. *Bioinformatics*, *26*(1),

    139–140. https://doi.org/10.1093/bioinformatics/btp616

Sheng, L., Hao, S.-L., Yang, W.-X., & Sun, Y. (2018). The multiple functions of kinesin-4

   family motor protein KIF4 and its clinical potential. *Gene*, *678*, 90–99.

   https://doi.org/10.1016/j.gene.2018.08.005

Sirovy, K. A., Johnson, K. M., Casas, S. M., La Peyre, J. F., & Kelly, M. W. (2021). Lack of

   genotype-by-environment interaction suggests limited potential for evolutionary changes

   in plasticity in the eastern oyster, *Crassostrea virginica*. *Molecular Ecology*, *30*(22),

   5721–5734. https://doi.org/10.1111/mec.16156

Spencer, L. H., Venkataraman, Y. R., Crim, R., Ryan, S., Horwith, M. J., & Roberts, S. B.

   (2020). Carryover effects of temperature and PCO_2 across multiple Olympia oyster

   populations. *Ecological Applications*, *30*(3), e02060. https://doi.org/10.1002/eap.2060

Stanley, J. G., Allen, S. K., & Hidu, H. (1981). Polyploidy induced in the American oyster,

   Crassostrea virginica, with cytochalasin B. *Aquaculture*, *23*(1–4), 1–10.

   https://doi.org/10.1016/0044-8486(81)90002-8

Tarazona, S., García-Alcalde, F., Dopazo, J., Ferrer, A., & Conesa, A. (2011). Differential

   expression in RNA-seq: A matter of depth. *Genome Research*, *21*(12), 2213–2223.

   https://doi.org/10.1101/gr.124321.111

Wadsworth, P., Wilson, A. E., & Walton, W. C. (2019). A meta-analysis of growth rate in

   diploid and triploid oysters. *Aquaculture*, *499*, 9–16.

   https://doi.org/10.1016/j.aquaculture.2018.09.018

Wang, X.-J., Zhang, D.-L., Xu, Z.-G., Ma, M.-L., Wang, W.-B., Li, L.-L., Han, X.-L., Huo, Y.,

   Yu, X., & Sun, J.-P. (2014). Understanding cadherin EGF LAG seven-pass G-type

   receptors. *Journal of Neurochemistry*, *131*(6), 699–711.

   https://doi.org/10.1111/jnc.12955

Whitehead, A., Galvez, F., Zhang, S., Williams, L. M., & Oleksiak, M. F. (2011). Functional
Genomics of Physiological Plasticity and Local Adaptation in Killifish. *Journal of
Heredity*, *102*(5), 499–511. https://doi.org/10.1093/jhered/esq077

Wright, R. M., Aglyamova, G. V., Meyer, E., & Matz, M. V. (2015). Gene expression associated
with white syndromes in a reef building coral, Acropora hyacinthus. *BMC Genomics*,
16(1), 371. https://doi.org/10.1186/s12864-015-1540-2

Xin, X., Liu, C., Liu, Z., Zhang, Y., Gao, Y., Zhu, T., Wang, L., & Song, L. (2022). Calmodulin
regulates the calcium homeostasis in mantle of Crassostrea gigas under ocean
acidification. *Frontiers in Marine Science*, *9*, 1050022.
https://doi.org/10.3389/fmars.2022.1050022

Yang, H. (2022). Performance and Fecundity of Triploid Eastern Oysters Crassostrea virginica
(Gmelin, 1791) and Challenges for Tetraploid Production. *Journal of Shellfish Research*,
40(3). https://doi.org/10.2983/035.040.0305

Yang, Q., Yu, H., & Li, Q. (2022). Disruption of cell division prevents gametogenesis in triploid
Pacific oysters (Crassostrea gigas). *Aquaculture*, *560*, 738477.
https://doi.org/10.1016/j.aquaculture.2022.738477

Zhang, X., Wen, H., Wang, H., Ren, Y., Zhao, J., & Li, Y. (2017). RNA-Seq analysis of salinity
stress–responsive transcriptome in the liver of spotted sea bass (Lateolabrax maculatus).
PLOS ONE, *12*(3), e0173238. https://doi.org/10.1371/journal.pone.0173238

Table 1: Number of DEGs detected for each pairwise comparison performed using EdgeR,
DESeq2, and NOISeq. R package VennDiagram was used to find common DEGs detected by all
three methods for each comparison.

Comparison	EdgeR	DESeq2	NOISeq	Common DEGs	Up-regulated DEGs	Down-regulated DEGs
2n LSU vs 3n LSU at GI	817	1860	702	289	113	176
2n LSU vs 3n LSU at LU	4683	5522	2864	1812	1000	812
2n AU vs 3n AU at GI	689	1925	198	125	26	99
2n AU vs 3n AU at LU	1886	2652	1628	709	273	436
2n AU vs 2n LSU at LUMCON	10690	10180	6246	3779	503	3346
2n AU vs 2n LSU at GI	1368	2232	1274	785	612	173
3n AU vs 3n LSU at LUMCON	10476	11810	13621	9022	5437	3585
3n AU vs 3n LSU at GI	1594	1371	2657	1065	677	388
2n vs 3n at GI	787	1371	776	281	117	164
2n vs 3n at LU	2589	3783	2867	1233	594	639
2n vs 3n at AU	1352	2187	2678	826	368	458
2n vs 3n at LSU	2282	3348	2243	867	310	557
Dam CL : 2n vs 3n	1595	2252	297	161	154	7
Dam VB : 2n vs 3n	1419	2137	93	58	40	18
Dam SL : 2n vs 3n	1576	2073	22	16	2	14

Figure 1: Cross strategy used to generate diploid and triploid lines for this study. Detailed cross strategy and number of oysters used for each cross are described in Bodenstein et al., 2023.

Figure 2: Cumulative mortality of diploid and triploid oysters outplanted to (A) Grand Isle and (B) LUMCON sites, replotted from Bodenstein et al., 2023. The X-axis indicates timepoints in months for which mortality data was collected, and the Y-axis represents cumulative mortality. Oysters are grouped together based on their ploidy and cohorts. Purple line: triploid oysters from LSU cohort, green dotted line: triploid oysters from Auburn cohort, blue dotted line: diploid oysters from LSU cohort, orange line: diploid oysters from Auburn cohort.

Figure 3: A) PCA showing clustering based on site and ploidy; B) transcriptomic shift measured as distance in multivariate space between diploid and triploid samples matched for their site, cohort, and dams.

Figure 4: a) Heatmap of WGCNA module-trait relationship results. The X-axis shows the experimental parameters of interest used for correlations. Red indicates positive correlations and blue indicates negative correlations between a module and experimental parameter. b) Eigengene expression plots for four modules showing significant correlation to site and cohort in WGCNA, with samples grouped together by their cohorts and ploidies; and GO terms with highest p-values in those corresponding modules. Orange: diploids from Auburn cohort, green: diploids from LSU cohort, blue: triploids from Auburn cohort, purple: triploids from LSU cohort.

Figure 1: Cross strategy used to generate diploid and triploid lines for this study. Detailed cross strategy and number of oysters used for each cross are described in Bodenstein et al., 2023.

Figure 2: Cumulative mortality of diploid and triploid oysters outplanted to (A) Grand Isle and (B) LUMCON sites, replotted from Bodenstein et al., 2023. The X-axis indicates timepoints in months for which mortality data was collected, and the Y-axis represents cumulative mortality. Oysters are grouped together based on their ploidy and cohorts. Purple line: triploid oysters from LSU cohort, green dotted line: triploid oysters from Auburn cohort, blue dotted line: diploid oysters from LSU cohort, orange line: diploid oysters from Auburn cohort.

Figure 3: A) PCA showing clustering based on site and ploidy; B) transcriptomic shift measured as distance in multivariate space between diploid and triploid samples matched for their site, cohort, and dams.

Figure 4: a) Heatmap of WGCNA module-trait relationship results. The X-axis shows the experimental parameters of interest used for correlations. Red indicates positive correlations and blue indicates negative correlations between a module and experimental parameter. b) Eigengene expression plots for four modules showing significant correlation to site and cohort in WGCNA, with samples grouped together by their cohorts and ploidies; and GO terms with highest p-values in those corresponding modules. Orange: diploids from Auburn cohort, green: diploids from LSU cohort, blue: triploids from Auburn cohort, purple: triploids from LSU cohort.

Supporting information:

Supplementary file 1: Count matrix used in downstream differential gene expression analysis and WGCNA.

Supplementary file 2: Enriched GO terms detected in differential gene expression analysis and WGCNA.

Supplementary table 1: Regression line slopes calculated for ten chromosomes for each diploid and triploid oyster pair, matched for their site, cohort, and dam.