The domesticated transposon protein L1TD1 associates with its ancestor L1 ORF1p to promote LINE-1 retrotransposition

Gülnihal Kavaklıoğlu[1], Alexandra Podhornik[1], Terezia Vcelkova[1], Jelena Marjanovic[1], Mirjam A. Beck[1], Trinh Phan-Canh[2], Theresia Mair[3], Claudia Miccolo[4], Aleksej Drino[1], Gerda Egger[3], Susanna Chiocca[4], Miha Modic[5,6], Christian Seiser[1*]

Division of Cell and Developmental Biology, Center for Anatomy and Cell Biology, Medical University of Vienna, Vienna, Austria

Medical University of Vienna, Max Perutz Labs Vienna, Campus Vienna Biocenter, Vienna, Austria

Department of Pathology, Medical University of Vienna, Vienna, Austria

Department of Experimental Oncology, IEO, European Institute of Oncology IRCCS, Milan, Italy

National Institute of Chemistry, Ljubljana, Slovenia

Dementia Research Institute at King's College London and The Francis Crick institute, London, UK

[*]Corresponding author:

Christian Seiser
Center for Anatomy and Cell Biology, Division for Cell and Developmental Biology, Medical University of Vienna, Schwarzspanierstrasse 17, 1090 Vienna, Austria
e-mail: christian.seiser@meduniwien.ac.at

ABSTRACT

Repression of retrotransposition is crucial for the successful fitness of a mammalian organism. The domesticated transposon protein L1TD1, derived from LINE-1 ORF1p, is an RNA-binding protein that is expressed only in some cancers and early embryogenesis. In human embryonic stem cells it is found to be essential for maintaining pluripotency. In cancer, L1TD1 expression is highly correlative with malignancy progression and as such considered a potential prognostic factor for tumors. However, its molecular role in cancer remains largely unknown. Our findings reveal that DNA hypomethylation induces the expression of L1TD1 in HAP1 human tumor cells. L1TD1 depletion significantly modulates both the proteome and transcriptome and thereby reduces cell viability. Notably, L1TD1 associates with LINE-1 transcripts and interacts with LINE-1 ORF1p protein, thereby facilitating LINE-1 retrotransposition. Our data suggest that L1TD1 collaborates with its ancestral LINE-1 ORF1p as an RNA chaperone, ensuring the efficient retrotransposition of LINE-1 retrotransposons, rather than directly impacting the abundance of L1TD1 targets. In this way, L1TD1 might have an important role not only during early development but also in tumorigenesis.

INTRODUCTION

Molecular domestication of transposable elements (TEs) or TE-derived sequences give rise to novel genes and regulatory sequences in the genome contributing to both genetic and epigenetic variation in an organism (Jangam *et al*, 2017; Miller *et al*, 1992; Smit, 1999). TEs are key drivers of genome instability and thus are constantly exposed to silencing mechanisms by the host. Yet, co-option of TEs thorough evolution has created so-called domesticated genes beneficial to the gene network in a wide range of organisms (Modzelewski et al., 2022; Jangam *et al.*, 2017). One example of such domesticated gene is LINE-1 type transposase containing domain 1 (*L1TD1*), which is the only domesticated protein-coding gene that originated from LINE-1 (McLaughlin *et al*, 2014). L1TD1 shares sequence resemblance with LINE-1 ORF1 protein (L1 ORF1p) that is required for retrotransposition (Martin, 2006) and L1TD1 was recently shown to facilitate the dissolution of stress granules (Jin *et al*, 2024). L1 ORF1p acts as nucleic acid chaperone with binding capacity to both RNA and DNA (Martin, 2006) and contributes to forming L1 ribonucleoproteins (L1-RNPs), key factors for retrotransposition of LINE-1 elements, but its co-factors are yet to be elucidated (Wallace *et al*, 2008).

L1TD1, also named Embryonic Stem Cell Associated Transcript 11 (*ECAT11*), is expressed predominantly only in early embryogenesis and germ cells and gets rapidly downregulated upon the exit from pluripotency (Iwabuchi *et al*, 2011; Mitsui *et al*, 2003). Although it is dispensable for mouse development (Iwabuchi *et al.*, 2011), L1TD1 is essential for the maintenance of human pluripotency (Narva *et al*, 2012; Emani *et al*, 2015). In the context of human embryonic stem cells (hESCs), L1TD1 regulates translation of a distinct set of mRNAs (Jin *et al.*, 2024), including core pluripotency factors (Emani *et al.*, 2015). It has been proposed, that during mammalian evolution the *L1TD1* gene was under positive selection due to the function in the maintenance of pluripotency in some species (McLaughlin *et al.*, 2014). Apart from early development, L1TD1 is expressed in germ cell tumors and colorectal cancers. In this context, L1TD1 was identified as a possible prognostic marker for medulloblastoma (Santos *et al*, 2015) and colorectal tumors (Chakroborty *et al*, 2019), where its expression is highly correlative with malignancy progression (Urh *et al*, 2021).

DNA methyltransferase (DNMT) inhibitors are successfully used as anti-cancer drugs and two recent reports suggest that the lethal response of human tumor cells to these compounds is based to a large extent on the activation of an anti-viral immune response to endogenous retroviral transcripts (Chiappinelli *et al*, 2015; Roulois *et al*, 2015). DNMT inhibitor treatment of non-small cell lung carcinoma (NSCLC) patient-derived cells resulted in up-regulation of L1TD1 and reduced cell viability indicating an epigenetic control of L1TD1 (Altenberger *et al*, 2017). Accordingly,

xenograft tumors with NSCLCs overexpressing L1TD1 showed decreased tumor growth suggesting a negative impact of L1TD1 expression on tumor viability. In contrast, L1TD1 was shown to be required for cell viability in medulloblastoma (Santos *et al.,* 2015). Hence, the exact mechanistic function of this domesticated transposon protein in human tumor cells is still unrevealed.

Here, we aimed at elucidating the molecular function of L1TD1 and its potential role for the control of viability in human tumor cells. To this end, we activated L1TD1 expression by inducing DNA hypomethylation *via* deletion of the DNA maintenance methyltransferase DNMT1 in the nearly haploid human cancer cell line HAP1 (Carette *et al*, 2011). First, we found that L1TD1 expression and LINE-1 activation correlate with local DNA hypomethylation. Next, we identified L1TD1-associated RNAs by RIP-seq, which revealed transcripts and proteins with differential expression in the absence of L1TD1 by transcriptome and proteome analyses. We showed that L1TD1 protein binds to L1-RNPs. Importantly, L1TD1 facilitated LINE-1 retrotransposition in HAP1 cells. Our results demonstrated that upon DNA hypomethylation L1TD1 affects gene expression and cell viability and cooperates with its ancestor protein L1 ORF1p in the control of LINE-1 retrotransposition.

MATERIALS AND METHODS

Cell lines and cell culture

HAP1 is a near-haploid human cell line derived from the KBM-7 chronic myelogenous leukemia (CML) cell line (Andersson *et al*, 1987; Carette *et al*., 2011). HAP1 cells were cultured in Iscove's Modified Dulbecco Medium (IMDM, Sigma Aldrich, I3390) supplement with 10% Fetal Bovine Serum (Sigma Aldrich, F7524) and 1% penicillin/streptomycin (Sigma Aldrich, P4333).

The human malignant papillary serous carcinoma cell line OV-90 (Gralewska *et al*, 2021) was cultured in MCDB 105 (Sigma Aldrich, 117) and Medium 199 (Sigma Aldrich, M4530) in a 1:1 ratio supplemented with 15% Fetal Bovine Serum (Sigma Aldrich, F7524) and 1% penicillin/streptomycin (Sigma Aldrich, P4333). The cells were kept in a humidified incubator at 37°C and 5% CO_2.

Gene editing

DNMT1 knockout (KO) HAP1 cells harboring a 20 bp deletion in the exon 4 of the *DNMT1* gene were generated using CRISPR/Cas9 gene editing. DNMT1/L1TD1 double knockout (DKO) HAP1 cells were generated by CRISPR/Cas9 gene editing resulting in a 13 bp deletion in exon 4 of the L1TD1 gene. CRISPR/Cas9 gene editing was performed by Horizon Genomics (Horizon Discovery).

Viability and Apoptosis Assay

Both assays were performed as described previously (Hess *et al*, 2022). For the viability assay $2x10^3$ cells of each cell line were seeded in triplicates in solid white 96-well plates and allowed to attach to the plates overnight. Cell viability was measured using CellTiter-Glo Luminiscent Cell Viability Assay (Promega, G7571) and Glomax Discover plate reader (Promega). The measurement was performed every 24 hours starting from the day after seeding (24h, 48h, 72h). For the apoptosis assay $5x10^6$ cells were collected 48hours after seeding, fixed for 20 min 2% paraformaldehyde (Merck Life Science) and for 30 min in 75% ethanol. Cells were permeabilized with 0.1% TritonX100 (Merck Life Science) for 10 min and blocked in 10% donkey serum (Merck Life Science) for 30 min and then incubated with 1:50 diluted anti-cleaved caspase 3 antibody in phosphate-buffered saline (PBS) (Cell Signaling, 9661) for 1 hour. Cells were washed and incubated with 1:400 Alexa 488 anti-rabbit secondary antibody (Jackson Immuno Research, 751-545-152). Samples were acquired using FACSCelesta flow cytometer (BD Bioscience) and the data were analyzed using FlowJo software 10.6.1 (BD Bioscience)

Immunoprecipitation

HAP1 cells pellets harvested from 15-cm culture plates were lysed in Hunt Buffer (20mM Tris/HCl ph 8.0, 100mM NaCl, 1mM EDTA, 0.5% NP-40) supplemented with Complete Protease inhibitor cocktail (Roche, 11697498001), Complete Phosphatase inhibitor cocktail (Roche, 11836145001), 10mM sodium fluoride, 10mM β-glycerophosphate, 0.1mM sodium molybdate and 0.1mM PMSF. The lysis was achieved by repeating freeze-thaw three times followed by centrifugation 12.000g for 15 min at 4 °C. Cell lysate was collected and the protein concentration was measured using Bradford assay. 40ul of Dynabeads^{TM} Protein G beads (ThermoFisher Scientific, 10004D) was blocked with 10% BSA in a rotor at 4°C for 1 hour. 1mg of cell lysate was incubated with blocked Protein G beads and monoclonal L1 ORF1p antibody (EMD Millipore, MABC1152) at 12 rpm at 4°C overnight. The next day, the beads were washed 3 times with Hunt buffer.

Western Blot Analysis

Immunoprecipitated complex was eluted from the beads with 1x SDS loading dye followed by heat-incubation at 95°C for 5 minutes for detection of proteins. For input samples, 20μg whole cell lysate was similarly denatured as IP samples. Proteins were separated in SDS-polyacrylamide gel and transferred onto nitrocellulose membrane (Amersham Protran, GE10600001, Sigma Aldrich) by wet transfer method. The membrane was blocked with the blocking solution (1x PBS, 1% polyvinylpyrrolidone, 3% non-fat dried milk, 0.1% tween-20, 0.01% sodium azide, pH 7.4) and incubated with the corresponding antibodies listed in **Supplementary Table 6**. To detect the protein of interest, ECL Western blotting detection reagents were used with a FUSION FX chemiluminescence imaging system.

Mass spectrometry analysis

Proteomes of DNMT1 KO and DNMT1/L1TD1 DKO cells were analyzed by quantitative TMT (Tandem Mass Tag) multiplex mass spectrometry analysis. Proteins were precipitated from cell extracts with acetone. After reduction and alkylation of Cys under denaturing conditions, proteins were digested with LysC/trypsin overnight. Peptides were cleaned-up using Oasis MCX (Waters) and labeled with TMT6plex (Thermo Fisher) according to manufacturer's protocol. Equal amounts of each channel were mixed, desalted and fractionated by neutral pH reversed phase chromatography. Equal interval fractions were pooled and analysed by LC-MS3 on an Ultimate 3000 RSLCnano LC coupled to an Orbitrap Eclipse™ Tribrid™ mass spectrometer via a FAIMS Pro ion mobility interface (all Thermo Fisher). Data were analyzed with MaxQuant 1.6.7.0.

(Tyanova *et al*, 2016). Differentially enriched proteins were determined with LIMMA (Ritchie *et al*, 2015). Workflow and analysis are described in detail in the **Supplementary information**.

RNA Immunoprecipitation

The native RIP protocol was followed as previously described in (Yang *et al*, 2017). In brief, cells were resuspended in 1ml Polysomal Lysis Buffer (100mM KCl, 5mM $MgCl_2$, 10mM HEPES (pH 7.0), 0.5% NP-40, 1mM DTT) supplemented with Protector RNase inhibitor (Sigma Aldrich, 3335399001) and protease inhibitor cocktail (Roche, 11697498001). Cell lysis was facilitated with 27G needle and syringe at least 7 times. The lysates were centrifuged at 12.000g for 15 min at 4℃. 1% input was taken from the cell lysate for RIP-seq experiment. The supernatant was cleared via pre-incubation with Dynabeads™ Protein G beads (ThermoFisher Scientific, 10004D) rotating for 1h at 4℃. For immunoprecipitation 1mg of pre-cleaned cell lysate was incubated with 4µg mouse monoclonal L1TD1 antibody (R&D systems, MAB8317) at 4℃ at 20rpm on the rotor overnight. 40ul Dynabeads™ Protein G beads per IP was blocked with 10 % bovine serum albumin (BSA) for 1h at 4℃. Blocked beads were added to the lysate and antibody mixture rotating for 1h at 4℃. Next, the beads were pelleted and washed 3 times with polysomal lysis buffer. 20% of the RIP was used for Western blot analysis to confirm immunoprecipitation of L1TD1. 1% of the RIP was saved as input. All experiments were performed in biological triplicates.

RNA isolation

To extract total RNA, immunoprecipitated RNA and 1% input RNA Monarch RNA Cleanup Kit (New England Biolabs, T2047L) was used according to the manufacturers' instructions. In the case of RIP samples, the beads were first treated with DNase I (New England Biolabs, M0303S) at 37℃ for 15 minutes and 1ml TRIzol (Thermo Fischer Scientific, 15596018) was directly added on. Upon addition of 150 µl chloroform the mixture was vortexed vigorously for 15 sec and centrifuged for at 12.000g for 15 minutes at 4℃. Aqueous phase was transferred to a new Eppendorf tube and mixed with 1 volume of EtOH (>95%). The mixture was loaded on the RNA columns and spun for 1 minute. The columns were washed 2 times with the washing buffer (supplemented with the kit) and eluted in 20ul nuclease free water. RNA concentration was measured using the Qubit RNA High Sensitivity kit (Thermo Fischer Scientific, Q32852)

cDNA library preparation

The cDNA libraries for the RIP-seq experiment were prepared as described in Smart-seq3 protocol by Hageman-Jensen *et al.*, 2020

cDNA synthesis and qRT-PCR

1ug of total RNA was reverse transcribed using iScript cDNA synthesis Kit (Bio-Rad, 1708891). Reverse transcription reaction was performed as follows: 25℃ for 5 minutes, 46℃ for 30 minutes, 95℃ for 5 minutes. Resulting cDNA was diluted to 1:10 with nuclease free water. 5μl of the cDNA was used for SYBR Green qPCR Master mix (Bio-Rad 1725275) together with the primers used in this study which were listed in **Supplementary Table 5**.

Methylight Assay

Genomic DNA was extracted from DNMT1 KO, DNMT1/L1TD1 DKO and WT HAP1 cells using the Wizard Genomic DNA isolation kit (Promega) following the manufacturer's protocol. Next, 1 μg of genomic DNA per sample was subjected to bisulfite conversion using the EZ DNA Methylation Kit (Zymo research Cat. no. D5001) following the manufacturer's instructions. Bisulfite-treated DNA was eluted in ddH$_2$O to a final concentration of 10 ng/μL and stored at -20°C until further use. The MethyLight method was performed as previously described by Campan *et al.* (Campan *et al*, 2009) using primers for *L1TD1* and *LINE-1*; *ALU* was used as a reference. The primer sequences are provided in **Supplementary Table 5**. The MethyLight reactions were done in technical triplicates. Each reaction contained 7.5 μL of 2X TaqMan Universal PCR Master Mix (Thermo Fisher Cat. No. 4324018), 300 nM of each primer and 100 nM of probe. For the *L1TD1* reactions 50ng and for the *LINE-1* reactions 20 ng of bisulfite-converted DNA were used in a total volume of 15 μL. The reactions were performed on a BioRad CFX96 thermocycler with an initial incubation at 95°C for 10 minutes, followed by 50 cycles of 95°C for 15 sec and 60°C for 1 minute. Each run included the individual samples, a serial dilution of fully methylated DNA, a non-methylated DNA control (Human Methylated & Non-Methylated (WGA) DNA Set D5013 Zymo research), and a non-template control. The PMR (percentage of methylation reference) of each individual sample was calculated using the following formula: 100 x [(GENE-X mean value) sample/ (ALU mean value) sample] / [(GENE-X mean value)100% methylated / (ALU mean value)100% methylated].

RNA-seq and RIP-seq Analyses

RNA sequencing data were analyzed using the RNA-Seq pipeline of IMP/IMBA Bioinfo core facility (ii-bioinfo@imp.ac.at). The pipeline is based on the nf-core/rnaseq pipeline [doi: 10.5281/zenodo.1400710] and is built with nextflow (Ewels *et al*, 2020).The RNA-seq analysis was performed as described in the following : Adapters were clipped with trimgalore (Martin,

2011). Abundant sequence fractions (rRNA) were removed using (bowtie2) (Langmead & Salzberg, 2012). Cleaned raw reads were mapped against the reference genome (hg38,Homo_sapiens.GRCh38.107.gtf) with STAR (reverse_stranded, STAR) (Dobin *et al*, 2013). Mapped reads were assigned to corresponding genes using featureCounts (Liao *et al*, 2014). Abundances were estimated through Kallisto (Bray *et al*, 2016) and Salmon (Patro *et al*, 2017). Analysis of differentially expressed genes was performed using DESeq2 (Love *et al*, 2014). TEtranscript software (Jin *et al*, 2015) was used to identify differentially expressed TE-derived sequences. In the case of RIP-seq, differential enrichment of transcripts relative to input as well as to the negative control (HAP1 DNMT1/L1TD1 DKO) was calculated using DEseq2. Only transcripts enriched ($\log_2$FC>2, adjusted p-value < 0.05) in both assays (DNMT1 KO *versus* input and DNMT1 KO *versus* DNMT1/L1TD1 DKO) were considered as L1TD1 associated RNAs.

L1 Retrotransposition Assay

Transfection of HAP1 cells with pJJ101, pJJ105, EGFP, pLenti6.2, pCEP4 was performed using polyethyleneimine (PEI)/NaCl solution. Briefly, 0.2×10^6 HAP1 cells/well were seeded into 6-well plates. 24 hours later and at 70% confluence, medium was replaced by fresh IMDM without penicillin/streptavidin followed by transfection. For each transfection, 100µl of 150mM NaCl, 7µl of PEI were mixed with DNA/NaCl solution (100µl of 150mM NaCl, 4µg of plasmid DNA), incubated at room temperature for 30 minutes, followed by adding the mixture in a drop-wise fashion onto cells. 18 hours post-transfection, the media was replaced with complete IMDM including 1% penicillin/streptavidin. Transfection efficiency was monitored 48 hours after transfection using fluorescent microscopy.

The retrotransposition assay was performed as described in (Kopera *et al,* 2016) with the following modification. To measure retrotransposition efficiency, 2×10^5 cells per well were seeded in 6-well culture plates and cultured at 37°C overnight. On day 0, 4ug of pJJ101/L1.3, pJJ105/L1.3, pLenti6.2, pCEP4 vectors described in Kopera *et al*., 2016 were transfected separately using PEI/NaCl in DNMT1 KO and DNMT1/L1TD1 DKO cells. To correct for potential differences in transfection efficiency and susceptibility to blasticidin in DNMT1 KO and DNMT1/L1TD1 DKO cells, we additionally used blasticidin deaminase containing vector (pLenti6.2). The following day (d1) the media was replaced with IMDM medium containing 1% penicillin/streptavidin. On d3, 2×10^5 cells per well seeded into 6-well culture dishes for each genotype and condition. Blasticidin treatment 10ug/ml was started on d4 and the cells were cultured 37°C for 9 days without medium change. Blasticidin resistant colonies were fixed on d13 with 4% Methanol-free formaldehyde (Thermo Fischer Scientific, 28908) at room temperature for 20 minutes and stained with 0.1% bromophenol blue (w/v) at room temperature for 1 hour. The pictures of the wells containing the

colonies were taken in FX chemiluminescence imaging system. The pictures were further processed inFiji Software following Analyze Particles function (Schindelin *et al*, 2012). The colony counts were obtained from three technical replicates per transfection. Mean colony counts were calculated and adjusted retrotransposition mean was calculated by adjusting for blasticidin resistant colonies in in blasticidin vector (pLenti6.2) transfected cells.

Indirect Immunofluorescence Staining

The protocol with minor modification from Sharma *et al.* (Sharma *et al*, 2016) was performed for immunostaining. Briefly, 75x10^3 HAP1 cells per well were plated on cover slips in 12 well plates. Next day, the cells were washed with 1x PBS and fixed with 4% methanol-free formaldehyde at room temperature for 10 minutes. The cells were washed with 1x PBS/Glycine pH 7.4 and permeabilized with 0.1% Triton-X in1x PBS/Glycine for 3 minutes. The samples were blocked with 1x PBS/Glycine/1% BSA at room temperature for 1 hour. After blocking, the cells were stained primary antibodies in blocking solution at 4 ℃ overnight. On the following day, the cells were washed three times with 1x PBS/Glycine/1% BSA for 5 minutes and stained with secondary antibody in blocking solution for 2 hours in dark. After a wash with PBS/Glycine, they were incubated with DAPI (Sigma Aldrich, D9542, 1:10000) in PBS/Glycine at room temperature for 10 minutes and washed three times with 1x PBS/Glycine pH 7.4. The slides were mounted with ProLong Gold Mountant (Thermo Fischer Scientific, P36930) and dried. The staining was monitored by Olympus Confocal Microscope.

RESULTS

L1TD1 expression is activated through DNA hypomethylation in HAP1 cells

To address the function of L1TD1 in human tumor cells, we generated a defined cellular model by using the nearly haploid tumor cell line HAP1 cells. By CRISPR/Cas-9 mediated gene editing, we deleted *DNMT1* in these cells resulting in HAP1 DNMT1 knock-out (KO) cells. As shown in **Figure 1A**, DNA methylation-specific PCR (MethyLight) analysis revealed significantly reduced DNA methylation at the *L1TD1* promoter in DNMT1 KO cells. DNA hypomethylation at the *L1TD1* gene correlated with a strong induction of L1TD1 mRNA and protein expression in DNMT1 KO cells (**Figure 1B-D)**. To analyze the role of L1TD1 in a DNMT1 null background, we ablated L1TD1 in KO cells by gene editing resulting in HAP1 DNMT1/L1TD1 double knock-out (DKO) cells (**Figure 1C**). Knowing that L1TD1 is highly expressed in human germ cell tumors, we used the ovarian cancer cell line OV-90 as a positive control (**Figure 1D-E**). In accordance with previously published data with human embryonic stem cells (hESCs), indirect immunofluorescence analysis revealed that L1TD1 was preferentially localized in the cytosol of HAP1 DNMT1 KO cells and OV-90 cells (**Figure 1E**). L1TD1 was reported to localize in P-bodies in hESCs (Narva *et al*, 2012). We detected a similar granular localization of L1TD1 in HAP1 DNMT1 KO cells (**Figure 1E**). DNMT1 depletion did not affect cell proliferation but additional loss of L1TD1 in DNMT1/L1TD1 DKO cells led to a significant reduction in the number of viable cells, and increased apoptosis (cleaved caspase 3) and DNA damage (γ-H2AX) (**Figure 1F-H**). These results suggest that L1TD1 expression is regulated by DNA methylation and its depletion affects cell viability in DNMT1-deficient HAP1 cells.

L1TD1 binds to LINE-1 transcripts and a subset of mRNAs

L1TD1 was previously shown to act as RNA-binding protein by binding to its own transcript (Narva *et al*., 2012) and selected group of mRNAs under L1TD1 translational control (Jin *et al*., 2024). To determine the L1TD1 binding repertoire in the context of L1TD1- and DNMT-dependent cell viability and proliferation control, we performed RNA immunoprecipitation (RIP) followed by sequencing (RIP-seq) (**Figure 2A**). L1TD1-containing RNP complexes (L1TD1-RNPs) were immunoprecipitated in DNMT1 KO cells by an L1TD1- specific antibody and DNMT1/L1TD1 DKO cells were used as a negative control. We identified 597 transcripts enriched ($\log_2$FC>2, adjusted p-value < 0.05) in L1TD1-RNPs when compared to the input (**Figure 2B** and **Suppl. Table S1**). Upon stringent analysis excluding non-specific binders and artifacts (**Suppl. Figure S1A** and **Suppl. Table S1**), 228 transcripts were specifically detected in L1TD1-RNPs. Importantly, *L1TD1*

mRNA was detected as one of the top transcripts in L1TD1-RNPs (**Figure 2B**).

To validate the RIP-seq results, individual transcripts selected from a list of top hits were amplified by gene-specific qRT-PCR analysis after repeating the RIP. Transcripts of *L1TD1*, *ARMC1* and *SYNJ2BP* were specifically enriched in L1TD1-RNPs obtained from DNMT1 KO cells when compared to DNMT1/L1TD1 DKO cells. Of note, L1TD1 was also associated with the transcript of *YY2*, a retrotransposon-derived paralogue of yin yang 1 (YY1) (Tahmasebi *et al*, 2016), whereas the negative control *GAPDH* did not show such enrichment (**Figure 2C**). Gene Ontology analyses of 228 common hits indicated enriched groups in cell division and cell cycle and, in accordance with a recent study (Jin *et al.*, 2024), enriched transcripts encoding zinc finger and in particular KRAB domain proteins which have been implicated in the restriction of endogenous retroviruses (Rosspopoff & Trono, 2023) (**Suppl. Figure S1B**).

Analysis of RIP-Seq using TEtranscript designed for the analysis of transposon-derived sequences (Jin *et al.*, 2015), in the differentially expressed sets indicated an association of L1TD1 with *LINE-1* transcripts (**Figure 2D** and **Suppl. Table S2**), which was also validated by qRT-PCR analysis of L1TD1 RIP samples (**Figure 2E**). In addition, we also identified *ERV* and *AluY* elements as L1TD1-RNP-associated transcripts **(Suppl. Table S2)**. These combined results suggest that L1TD1 interacts with a set of transcripts including RNAs with functions in cell division. Furthermore, L1TD1 has kept the ability to associate with *LINE-1* transcripts, and potentially regulating them similar to its ancestor protein L1 ORF1p (L1 ORF1p) (Wei *et al*, 2001).

Loss of L1TD1 modulates the proteome and transcriptome of HAP1 cells

To explore the cellular function of L1TD1, we analyzed the proteomes and transcriptomes of DNMT1 KO and DNMT1/L1TD1 DKO cells. Comparison of the DNMT1 KO and DNMT1/L1TD1 DKO proteomes revealed 98 upregulated and 131 downregulated proteins in DNMT1/L1TD1 DKO ($log_2FC \geq 1$, adj. p-value ≤ 0.05) (**Figure 3A** and **Suppl. Table S3**). Gene Ontology Enrichment Analysis revealed that differentially expressed proteins were mainly associated with regulation of cell junction and cysteine-type endopeptidase activity involved in apoptotic process (**Suppl. Fig. S2A**). To investigate whether the corresponding protein abundance of mRNAs contained in L1TD1-RNPs is affected by L1TD1 expression, we compared proteomic data with the RIP-seq data. Comparison of differentially expressed proteins with transcripts associated with L1TD1 identified only L1TD1 (**Suppl. Fig. S2B**). The lack of overlap between RIP-seq and proteomics data (except L1TD1) suggested that modulation of the abundance of proteins encoded by L1TD1-associated transcripts is not the main function of L1TD1.

Interestingly, we discovered that the L1 transposon protein ORF1p was up-regulated in the

absence of L1TD1 (**Suppl. Table S3**). Western blot analysis confirmed the increased expression of L1 ORF1p in HAP1 DNMT1/L1TD1 DKO cells when compared to DNMT1 KO cells (**Figure 3B**), while LINE-1 transcript levels showed no significant change (**Suppl. Figure S3A**). This indicates that L1TD1 attenuates the expression of L1 ORF1p suggesting a potential regulatory crosstalk between L1TD1 with its ancestor protein.

To investigate a potential effect of L1TD1 on the expression levels of associated RNAs, comparative RNA-seq analysis was performed with DNMT1 KO and DNMT1/L1TD1 DKO cells. Upon loss of L1TD1, we identified 323 up-regulated and 321 down-regulated transcripts, respectively (log_2-fold change ≥ 1; adjusted p-value < 0.05) (**Figure 3B** and **Suppl. Table S4**). About 25% of the differentially expressed proteins (35% of upregulated and 21% of downregulated proteins) identified in the proteome analysis were also significantly deregulated at the RNA level (**Suppl. Table S3**). However, none of the deregulated transcripts, except *L1TD1*, was detected in L1TD1-RNPs (**Figure 3B** and **Suppl. Tables S1** and **S4**). This finding was corroborated by qRT-PCR analysis of L1TD1-RNP-associated transcripts (**Suppl. Figure S3B**) suggesting that the cellular function of L1TD1 might be independent of a deregulation of its associated mRNAs.

Based on our observations implying that L1TD1 does not exert its cellular function as translational regulator while being associated with transposon RNAs **(Figure 2D)**, we used TEtranscript (Jin *et al.*, 2015) to identify differentially expressed TE-derived sequences transcriptome-wide. Loss of L1TD1 resulted in upregulation of specific SINEs such as AluY elements (**Suppl. Figure S3C** and **Suppl. Table S4).** Taken together, these data suggest that L1TD1 affects the abundance of transposon transcripts in DNMT1-deficient HAP1 cells.

L1TD1 interacts with the LINE-1 ORF1p protein

An interaction of L1TD1 with *LINE-1* RNA (**Figure 2D**) and increased L1 ORF1p levels in L1TD1-depleted cells (**Figure 3B**) imply a direct association of L1TD1 with L1 ORF1p. We thus hypothesized that L1TD1 associates with L1-RNPs *via* L1 ORF1p. Therefore, we immunoprecipitated L1 ORF1p from HAP1 DNMT1KO and OV-90 cells and tested the IPs for the presence of endogenous L1TD1. Indeed, L1TD1 was co-precipitated with L1 ORF1p from DNMT1 KO cells (**Figure 3D**). Next, we showed that the association of L1TD1 with L1 ORF1p was RNA-independent **(Figures S4A** and **S4B**).

In agreement with their association in protein complexes, indirect immunofluorescence experiments revealed a partial co-localization of L1TD1 and L1 ORF1p in both cell lines (**Figure 3E** and **Suppl. Figure S4C)**. In summary, our results suggest L1TD1 associates with its ancestor ORF1p and this association does not depend on RNA intermediates.

L1TD1 has a positive impact on LINE-1 retrotransposition

Since L1TD1 associates with *LINE-1* RNA and modulates L1 ORF1p expression, we next asked whether the domesticated transposon protein can impact LINE-1 retrotransposition in HAP1 cells. To this end, we carried out plasmid-based retrotransposition assays (Kopera *et al.*, 2016) in DNMT1 KO and DNMT1/L1TD1 DKO cells (**Figure 4A-B)**. Following transfection and blasticidin selection, the rate of retrotransposition of the reporter construct was assessed through counting blasticidin-resistant colonies, transfected either with a retrotransposition-competent reporter construct or a retrotransposition-deficient control. An average of 506 colonies were observed in DNMT1 KO cells in three independent experiments whereas DNMT1/L1TD1 DKO cells yielded over 5-fold fewer colonies when transfected with the retrotransposition-competent reporter construct (86 colonies on average) (**Figures 4C-D**). No blasticidin resistant clones were obtained for the negative controls, but comparable retrotransposition rates were observed for retrotransposition-competent reporter constructs in three independent retrotransposition assays (**Suppl. Figures S5A-B**). Combined, these data suggest that L1TD1 enhances LINE-1 retrotransposition in DNMT1-deficient HAP1 cells and this effect is mediated through the association with ORF1p and LINE-1 RNA.

Discussion

In this study we show that the domesticated transposon protein L1TD1 promotes LINE-1 retrotransposition. Numerous cellular factors have been shown in the past to affect retrotransposition (Ariumi, 2016; Goodier, 2016; Pizarro & Cristofari, 2016; Protasova *et al*, 2021). However, many of these proteins are part of host defense mechanisms and restrict retrotransposition, and it was speculated that L1TD1 also limits L1 retrotransposition (McLaughlin *et al.*, 2014). Unexpectedly, we identified L1TD1 as one of the cellular factors that has a positive impact on L1 retrotransposition, most likely by interacting with L1 RNPs and acting as RNA chaperone. L1TD1 deletion resulted in reduced L1 retrotransposition despite upregulation of ORF1p indicating that the upregulation of the transposon protein cannot fully compensate the loss of L1TD1. Interestingly, a similar effect has been previously described for the nonsense-mediated decay factor UPF1 (Taylor *et al*, 2013). UPF1 knockdown increased the amount of L1 mRNA and proteins but simultaneously reduced the effectiveness of the retrotransposition. A positive effect of L1TD1 on retrotransposition was recently observed upon L1TD1 overexpression in HeLa cells (Jin *et al.*, 2024). However, the authors considered this as artefact of ectopic L1TD1 expression. L1TD1 was lost or pseudogenized multiple times during mammalian evolution and has evolved under positive selection during primate and mouse evolution suggesting an important function during development (McLaughlin *et al.*, 2014). L1 elements can retrotranspose in the germline, in embryonic stem cells and in the early embryo (Ariumi, 2016; Goodier, 2016; Pizarro & Cristofari, 2016; Protasova *et al.*, 2021). Mouse embryos with repressed LINE-1 expression show impaired embryonic development, suggesting that LINE-1 is part of the developmental program of early embryos (Jachowicz *et al*, 2017). The simultaneous L1TD1 expression during early embryogenesis might ensure activation of L1 retrotransposition at this developmental stage (Iwabuchi *et al.*, 2011).

To gain insight into the regulatory function of L1TD1 in tumor cells, we performed RIP-seq, transcriptome and proteome analyses in HAP1 cells. The RIP-seq approach identified, in addition to the known associated *L1TD1* mRNA (Narva *et al.*, 2012), a defined set of transcripts comprising mRNAs, lncRNAs and transposable elements including *LINE-1* transcripts. These findings indicate that L1TD1 has not only inherited its function as RNA-binding protein but also its affinity for transposon transcripts. Recently, Jin *et al.* published a study on the impact of L1TD1 on the translation in hESCs and identified L1TD1-bound RNAs by CLIP-seq (Jin *et al.*, 2024). Interestingly, the majority of L1TD1-associated transcripts in HAP1 cells (69%) identified in our study were also reported as L1TD1 targets in hESCs suggesting a conserved binding affinity of

this domesticated transposon protein across different cell types.

The same study (Jin *et al.*, 2024) showed that L1TD1 localizes to high-density RNP condensates and enhances the translation of a subset of mRNAs enriched in the condensates. Our mass spectrometry data showed that L1TD1 ablation leads to a significantly changed proteome. However, none of the proteins encoded by the mRNAs associated with L1TD1 showed significantly changed abundance in the absence of L1TD1 suggesting that this is not a direct effect of L1TD1 association with the respective mRNAs. The different outcomes might be due to different cell types and methods used in the two studies. Similarly, and in accordance with the hESC study (Jin *et al.*, 2024), we did not find an overlap between RNAs associated with L1TD1 and their deregulation at the RNA level. This suggests that in HAP1 cells L1TD1 does not have a global effect on RNA stability and translation of its associated RNAs. This is consistent with a recent study on L1 bodies, where RNA processing association of L1 body components did not correlate with RNA stability or translational control of the RNA targets of L1 bodies (De Luca *et al*, 2023).

Of note, L1TD1 associated with *SINE* transcripts such as AluY and L1TD1 ablation positively affected AluY transposon expression. This relatively young subfamily of SINE elements is still retrotransposition-competent and their mobilization depends on the activity of LINE proteins (Roy-Engel, 2012; Styles & Brookfield, 2009). These observations suggest that L1TD1 has not only inherited its function as an RNA-binding protein but also the affinity to *LINE-1* and *SINE* transcripts. L1TD1 association with *AluY* transcripts could potentially be connected to modulation of LINE-1 and SINE-1 transposon activities.

LINE-1 is the only protein-coding transposon that is active in humans and LINE-1 overexpression and retrotransposition are hallmarks of cancers (Kazazian & Moran, 2017; Mendez-Dorantes & Burns, 2023). Similarly to the epigenetic control of the ancestral LINE-1 elements (Sanchez-Luque *et al*, 2019; Zhou *et al*, 2020), L1TD1 expression in HAP1 cells is restricted by DNA methylation and DNA hypomethylation induced by DNMT1 ablation results in robust L1TD1 RNA and protein expression. This observation is in line with the findings in the NSCLC cancer model, where pharmacological DNMT inhibition resulted in L1TD1 induction (Altenberger *et al.*, 2017). In contrast to the negative effect of L1TD1 on cell viability and tumor growth in the NSCLC xenograft model, we report that L1TD1 deletion in DNMT1-deficient cells led to reduced cell viability and increased apoptosis in HAP1 cancer cells. This is in accordance with the observation that L1TD1 has a positive impact on cell viability in medullablastoma (Santos *et al.*, 2015) and suggests a (cancer) cell type-specific effect of L1TD1 that might be related to the DNA methylation state of the tumors.

Taken together we present here a novel cellular function for a domesticated transposon protein

by showing that L1TD1 associates with L1-RNPs and promotes LINE-1 retrotransposition. This function might have a beneficial effect during early development but could also impact on tumorigenesis.

Acknowledgments

We would like to thank Wolfgang Sommergruber for the screen of human tumor cells for L1TD1 expression and Jernej Ule for help with the analysis of RNPs. We also want to thank Luisa Seufert, Stephanie Schneider, Christina Maria Schuh and Marlene Müller for help with the characterization of transgenic HAP1 cells, Brigitte Gundacker, Urska Janjos and Milena Mijovic for professional technical support and Dorothea Anrather for help with data analysis. We are also grateful to Wolfgang Miller, Heinz Fischer and Matthias Schaefer for numerous helpful discussions.
GL and TM were supported by the Austrian Science Foundation (FWF, doc.funds grant DOC59), CS was supported by the Austrian Science Foundation (P34998, DOC32) and the Austrian Research Promotion Agency (FFG): Bridge Early Phase project 5722451. GK were PhD students of the doc.funds program (DOC32) supported by the Austrian Science Foundation (FWF). TPC was supported by a Student Fellowship from the Ernst Mach Grant - ASEA-UNINET through the Austrian Academic Exchange (OeAD).

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol *et al*, 2022) partner repository with the dataset identifier PXD047402. RNA-seq and RIP-seq data were submitted to GEO under accession numbers GSE254459 and GSE254460.

REFERENCES

Altenberger C, Heller G, Ziegler B, Tomasich E, Marhold M, Topakian T, Mullauer L, Heffeter P, Lang G, End-Pfutzenreuter A *et al* (2017) SPAG6 and L1TD1 are transcriptionally regulated by DNA methylation in non-small cell lung cancers. *Mol Cancer* 16: 1

Andersson BS, Beran M, Pathak S, Goodacre A, Barlogie B, McCredie KB (1987) Ph-positive chronic myeloid leukemia with near-haploid conversion in vivo and establishment of a continuously growing cell line with similar cytogenetic pattern. *Cancer Genet Cytogenet* 24: 335-343

Ariumi Y (2016) Guardian of the Human Genome: Host Defense Mechanisms against LINE-1 Retrotransposition. *Front Chem* 4: 28

Bray NL, Pimentel H, Melsted P, Pachter L (2016) Near-optimal probabilistic RNA-seq quantification. *Nature biotechnology* 34: 525-527

Campan M, Weisenberger DJ, Trinh B, Laird PW (2009) MethyLight. *Methods Mol Biol* 507: 325-337

Carette JE, Raaben M, Wong AC, Herbert AS, Obernosterer G, Mulherkar N, Kuehne AI, Kranzusch PJ, Griffin AM, Ruthel G *et al* (2011) Ebola virus entry requires the cholesterol transporter Niemann-Pick C1. *Nature* 477: 340-343

Chakroborty D, Emani MR, Klen R, Bockelman C, Hagstrom J, Haglund C, Ristimaki A, Lahesmaa R, Elo LL (2019) L1TD1 - a prognostic marker for colon cancer. *BMC Cancer* 19: 727

Chiappinelli KB, Strissel PL, Desrichard A, Li H, Henke C, Akman B, Hein A, Rote NS, Cope LM, Snyder A *et al* (2015) Inhibiting DNA Methylation Causes an Interferon Response in Cancer via dsRNA Including Endogenous Retroviruses. *Cell* 162: 974-986

De Luca C, Gupta A, Bortvin A (2023) Retrotransposon LINE-1 bodies in the cytoplasm of piRNA-deficient mouse spermatocytes: Ribonucleoproteins overcoming the integrated stress response. *PLoS genetics* 19: e1010797

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29: 15-21

Emani MR, Narva E, Stubb A, Chakroborty D, Viitala M, Rokka A, Rahkonen N, Moulder R, Denessiouk K, Trokovic R *et al* (2015) The L1TD1 protein interactome reveals the importance of post-transcriptional regulation in human pluripotency. *Stem Cell Reports* 4: 519-528

Ewels PA, Peltzer A, Fillinger S, Patel H, Alneberg J, Wilm A, Garcia MU, Di Tommaso P, Nahnsen S (2020) The nf-core framework for community-curated bioinformatics pipelines. *Nature biotechnology* 38: 276-278

Goodier JL (2016) Restricting retrotransposons: a review. *Mob DNA* 7: 16

Gralewska P, Gajek A, Marczak A, Rogalska A (2021) Metformin Affects Olaparib Sensitivity through Induction of Apoptosis in Epithelial Ovarian Cancer Cell Lines. *International journal of molecular sciences* 22

Hess L, Moos V, Lauber AA, Reiter W, Schuster M, Hartl N, Lackner D, Boenke T, Koren A, Guzzardo PM *et al* (2022) A toolbox for class I HDACs reveals isoform specific roles in gene regulation and protein acetylation. *PLoS genetics* 18: e1010376

Iwabuchi KA, Yamakawa T, Sato Y, Ichisaka T, Takahashi K, Okita K, Yamanaka S (2011) ECAT11/L1td1 is enriched in ESCs and rapidly activated during iPSC generation, but it is dispensable for the maintenance and induction of pluripotency. *PloS one* 6: e20461

Jachowicz JW, Bing X, Pontabry J, Boskovic A, Rando OJ, Torres-Padilla ME (2017) LINE-1 activation after fertilization regulates global chromatin accessibility in the early mouse embryo. *Nature genetics* 49: 1502-1510

Jangam D, Feschotte C, Betran E (2017) Transposable Element Domestication As an Adaptation to Evolutionary Conflicts. *Trends Genet* 33: 817-831

Jin SW, Seong Y, Yoon D, Kwon YS, Song H (2024) Dissolution of ribonucleoprotein condensates by the embryonic stem cell protein L1TD1. *Nucleic acids research*

Jin Y, Tam OH, Paniagua E, Hammell M (2015) TEtranscripts: a package for including transposable elements in differential expression analysis of RNA-seq datasets. *Bioinformatics* 31: 3593-3599

Kazazian HH, Jr., Moran JV (2017) Mobile DNA in Health and Disease. *N Engl J Med* 377: 361-370

Kopera HC, Larson PA, Moldovan JB, Richardson SR, Liu Y, Moran JV (2016) LINE-1 Cultured Cell Retrotransposition Assay. *Methods Mol Biol* 1400: 139-156

Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nature methods* 9: 357-359

Liao Y, Smyth GK, Shi W (2014) featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 30: 923-930

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome biology* 15: 550

Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *2011* 17: 3

Martin SL (2006) The ORF1 protein encoded by LINE-1: structure and function during L1 retrotransposition. *Journal of biomedicine & biotechnology* 2006: 45621

McLaughlin RN, Jr., Young JM, Yang L, Neme R, Wichman HA, Malik HS (2014) Positive selection and multiple losses of the LINE-1-derived L1TD1 gene in mammals suggest a dual role in genome defense and pluripotency. *PLoS genetics* 10: e1004531

Mendez-Dorantes C, Burns KH (2023) LINE-1 retrotransposition and its deregulation in cancers: implications for therapeutic opportunities. *Genes & development* 37: 948-967

Miller WJ, Hagemann S, Reiter E, Pinsker W (1992) P-element homologous sequences are tandemly repeated in the genome of Drosophila guanche. *Proceedings of the National Academy of Sciences of the United States of America* 89: 4018-4022

Mitsui K, Tokuzawa Y, Itoh H, Segawa K, Murakami M, Takahashi K, Maruyama M, Maeda M, Yamanaka S (2003) The homeoprotein Nanog is required for maintenance of pluripotency in mouse epiblast and ES cells. *Cell* 113: 631-642

Modzelewski AJ, Gan Chong J, Wang T, He L. Mammalian genome innovation through transposon domestication. *Nat Cell Biol* 2022;24: 1332–1340

Narva E, Rahkonen N, Emani MR, Lund R, Pursiheimo JP, Nasti J, Autio R, Rasool O, Denessiouk K, Lahdesmaki H *et al* (2012) RNA-binding protein L1TD1 interacts with LIN28 via RNA and is required for human embryonic stem cell self-renewal and cancer cell proliferation. *Stem cells* 30: 452-460

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C (2017) Salmon provides fast and bias-aware quantification of transcript expression. *Nature methods* 14: 417-419

Perez-Riverol Y, Bai J, Bandla C, Garcia-Seisdedos D, Hewapathirana S, Kamatchinathan S, Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M *et al* (2022) The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences. *Nucleic acids research* 50: D543-D552

Pizarro JG, Cristofari G (2016) Post-Transcriptional Control of LINE-1 Retrotransposition by Cellular Host Factors in Somatic Cells. *Front Cell Dev Biol* 4: 14

Protasova MS, Andreeva TV, Rogaev EI (2021) Factors Regulating the Activity of LINE1 Retrotransposons. *Genes (Basel)* 12

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK (2015) limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic acids research* 43: e47

Rosspopoff O, Trono D (2023) Take a walk on the KRAB side: (Trends in Genetics, 39:11 p:844-857, 2023). *Trends Genet*

Roulois D, Loo Yau H, Singhania R, Wang Y, Danesh A, Shen SY, Han H, Liang G, Jones PA, Pugh TJ *et al* (2015) DNA-Demethylating Agents Target Colorectal Cancer Cells by Inducing Viral Mimicry by Endogenous Transcripts. *Cell* 162: 961-973

Roy-Engel AM (2012) LINEs, SINEs and other retroelements: do birds of a feather flock together? *Front Biosci (Landmark Ed)* 17: 1345-1361

Sanchez-Luque FJ, Kempen MHC, Gerdes P, Vargas-Landin DB, Richardson SR, Troskie RL, Jesuadian JS, Cheetham SW, Carreira PE, Salvador-Palomeque C *et al* (2019) LINE-1 Evasion of Epigenetic Repression in Humans. *Molecular cell* 75: 590-604 e512

Santos MC, Silva PB, Rodini CO, Furukawa G, Marco Antonio DS, Zanotto-Filho A, Moreira JC, Okamoto OK (2015) Embryonic Stem Cell-Related Protein L1TD1 Is Required for Cell Viability, Neurosphere Formation, and Chemoresistance in Medulloblastoma. *Stem Cells Dev* 24: 2700-2708

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B *et al* (2012) Fiji: an open-source platform for biological-image analysis. *Nature methods* 9: 676-682

Sharma R, Rodic N, Burns KH, Taylor MS (2016) Immunodetection of Human LINE-1 Expression in Cultured Cells and Human Tissues. *Methods Mol Biol* 1400: 261-280

Smit AF (1999) Interspersed repeats and other mementos of transposable elements in mammalian genomes. *Current opinion in genetics & development* 9: 657-663

Styles P, Brookfield JF (2009) Source gene composition and gene conversion of the AluYh and AluYi lineages of retrotransposons. *BMC Evol Biol* 9: 102

Tahmasebi S, Jafarnejad SM, Tam IS, Gonatopoulos-Pournatzis T, Matta-Camacho E, Tsukumo Y, Yanagiya A, Li W, Atlasi Y, Caron M *et al* (2016) Control of embryonic stem cell self-renewal and differentiation via coordinated alternative splicing and translation of YY2. *Proceedings of the National Academy of Sciences of the United States of America* 113: 12360-12367

Taylor MS, LaCava J, Mita P, Molloy KR, Huang CR, Li D, Adney EM, Jiang H, Burns KH, Chait BT *et al* (2013) Affinity proteomics reveals human host factors implicated in discrete stages of LINE-1 retrotransposition. *Cell* 155: 1034-1048

Tyanova S, Temu T, Cox J (2016) The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. *Nat Protoc* 11: 2301-2319

Urh K, Zlajpah M, Zidar N, Bostjancic E (2021) Identification and Validation of New Cancer Stem Cell-Related Genes and Their Regulatory microRNAs in Colorectal Cancerogenesis. *Biomedicines* 9

Wallace N, Wagstaff BJ, Deininger PL, Roy-Engel AM (2008) LINE-1 ORF1 protein enhances Alu SINE retrotransposition. *Gene* 419: 1-6

Wei W, Gilbert N, Ooi SL, Lawler JF, Ostertag EM, Kazazian HH, Boeke JD, Moran JV (2001) Human L1 retrotransposition: cis preference versus trans complementation. *Molecular and cellular biology* 21: 1429-1439

Yang CC, Chen YT, Chang YF, Liu H, Kuo YP, Shih CT, Liao WC, Chen HW, Tsai WS, Tan BC (2017) ADAR1-mediated 3' UTR editing and expression control of antiapoptosis genes fine-tunes cellular apoptosis response. *Cell Death Dis* 8: e2833

Zhou W, Liang G, Molloy PL, Jones PA (2020) DNA methylation enables transposable element-driven genome expansion. *Proceedings of the National Academy of Sciences of the United States of America* 117: 19359-19366

FIGURE LEGENDS

Figure 1. DNA hypomethylation results in activation of L1TD1 expression and loss of L1TD1 affects cell viability in HAP cells. **(A)** Quantification of DNA methylation levels at the *L1TD1* promoter in HAP1 wildtype (WT), DNMT1 KO and DNMT1/L1TD1 DKO cells using the MethyLight assay. DNA methylation is shown as percentage of methylation ratio (PMR). **(B)** qRT-PCR analysis of *L1TD1* mRNA expression in HAP1 wildtype (WT), DNMT1 KO and DNMT1/L1TD1 DKO cells. *GAPDH* was used as a normalization control and relative *L1TD1* mRNA levels in DNMT1 KO cells were set to 1. All data in the figure are shown as a mean of ± SD of 3 biological replicates. **(C)** Western blot analysis of L1TD1 levels in HAP1 WT, DNMT1 KO and DNMT1/L1TD1 DKO cells. LAMIN B was used as a loading control. **(D)** Western blot analysis of L1TD1 protein expression in HAP1 WT and DNMT1 KO cells and OV-90 cells. β-actin was used as a loading control. **(E)** Indirect immunofluorescent staining of L1TD1 in HAP1 DNMT1 KO and DNMT1/L1TD1 DKO cells and OV-90 cells. Nuclear DNA was stained with DAPI (blue). **(F)** Cell viability analysis of HAP1 WT, DNMT1 KO and DNMT1/L1TD1 DKO cells using the CellTiter-Glo assay measured over 72h. **(G)** Bar graph representing the percentage of apoptotic cells of cultured cell lines quantified by flow cytometry analysis of cleaved caspase 3. **(H)** Western blot analysis of γH2AX levels in HAP1 WT, DNMT1 KO and DNMT1/L1TD1 DKO cells in nuclear extracts. Antibodies specific for histone H3 C-terminus and LAMIN B were used as loading controls. **(A-B, F-G)** Statistical significance was determined using one-way ANOVA with Tukey's multiple comparison correction. * p≤0.05, ** p ≤0.01, ** p ≤0.01, *** p ≤0.001, ns = not significant.

Figure 2. RIP-seq identifies a set of RNAs and transposon transcripts associated with L1TD1. **(A)** Schematic representation of the RNA immunoprecipitation sequencing (RIP-seq) method (Figure created with BioRender.com). L1TD1-RNA complexes were isolated from HAP1 KO cell extracts with an L1TD1-specific antibody, RNA was isolated from complexes and input and cDNA libraries were prepared using the Smart-seq3 protocol. Sequencing data was analyzed by DEseq2 and TEtranscript software, separately. **(B)** Volcano plot showing L1TD1-associated transcripts as a result of DESeq2 analysis (cut-off log2FC > 2 and adj. p-value < 0.05). Selected hits are highlighted in green. **(C)** RIP-qPCR analysis confirms the interaction of L1TD1 to the transcripts *L1TD1*, *ARMC1*, *YY2*. The bar graphs represent the fold enrichments of the transcripts in the IP samples normalized to input samples in the indicated cells. *GAPDH* was used as a negative control for the top hit transcripts. **(D)** Volcano plot showing L1TD1-associated transposon transcripts as result of the TEtranscript analysis with a log2FC > 1 and adj. p-value < 0.05. **(E)**

RIP-qPCR analysis confirms the association of L1TD1 with *LINE-1* transcripts. Statistical significance was determined using paired two-tailed t-test. All data in the figure are shown as a mean of ± SD of 3 biological replicates. * p ≤0.05, ** p ≤0.01, *** p ≤0.001, **** p ≤0.0001, ns = not significant.

Figure 3. L1TD1 cross-talk with its ancestor L1 ORF1p. (A) Proteome analysis: Volcano plot displaying proteomics of HAP1 DNMT1 KO and DNMT1/L1TD1 DKO cells determined by mass spectrometry. Differentially abundant proteins were plotted as DNMT1/L1TD1 DKO over DNMT1 KO (log2FC ≥ 1, adj. p-value < 0.05 (red) and log2FC ≤ -1, adj. p-value < 0.05 (blue)). **(B)** Transcriptome analysis: Volcano plot illustrating DESeq2 analysis of RNA-seq performed in HAP1 DNMT1 KO and DNMT1/L1TD1 DKO cells. Differentially expressed genes are plotted as DNMT1/L1TD1 DKO over DNMT1 KO (log2FC ≥ 1, adj. p-value < 0.05 (red) and log2FC ≤ -1, adj. p-value < 0.05 (blue)). **(C)** Western blot analysis illustrating protein levels of L1TD1 and L1 ORF1p in HAP1 WT, DNMT1 KO and DNMT1/L1TD1 DKO cells. β-actin was used as loading control. **(D)** Physical interaction of L1 ORF1p and L1TD1. L1 ORF1p was immunoprecipitated with an L1 ORF1p-specific antibody from whole cell extracts prepared from DNMT1 KO and DNMT1/L1TD1 DKO cells and precipitated L1 RNP complexes were analyzed on a Western blot. IgG was used as a negative IP control and β-actin was used as loading control. **(E)** Confocal microscopy images of indirect immunofluorescence co-stainings using mouse ORF1p (green) and rabbit L1TD1 (red) antibodies in DNMT1 KO cells. Nuclear DNA was stained with DAPI.

Figure 4. L1TD1 promotes L1 retrotransposition. (A) Schematic representation of plasmids used for retrotransposition (Figure modified from (Kopera *et al.*, 2016)) (Figure created with BioRender.com). The pJJ101/L1.3 construct contains the full length human L1.3 element with a blasticidin deaminase gene (mblast) inserted in antisense within the 3'UTR. The *mblast* gene is disrupted with an intron and *mblast* expression occurs only when L1 transcript is expressed, reverse transcribed and inserted into the genome. The pJJ105/L1.3 plasmid contains a mutation in the reverse transcriptase (RT) resulting in defective retrotransposition. The backbone plasmid pCEP4 was used as additional negative control. The blasticidin deaminase gene containing plasmid pLenti6.2 was used as transfection/selection control. **(B)** Workflow of retrotransposition assay. DNMT1 KO and DNMT1/L1TD1 DKO cells were separately transfected with pJJ101 and control plasmids. Equal number of cells were seeded for each condition. Blasticidin selection (10μg/ml) was started at day 4 and resistant colonies were counted on day 13. **(C)** Bar graph showing the average number of retrotransposition events per 10^6 cells seeded in three

independent experiments. Blasticidin resistant colonies in pLenti6.2 transfected cells were used for normalization. Statistical significance was determined using unpaired t-test. All data in the figure are shown as a mean of ± SD of three independent experiments, **** p ≤0.0001. **(D)** Representative pictures of bromophenol blue stainings of resistant colonies for each genotype and each transfection.

SUPPLEMENTARY FIGURE LEGENDS

Supplementary Figure 1. L1TD1 interacts with a specific set of mRNAs and transposon transcripts. (A) Volcano plot shows the L1TD1-associated transcripts by L1TD1 enriched in DNMT1 KO IP over DNMT1/L1TD1 DKO IP with a log2FC ≥ 2 and adj. p-val < 0.05. The transcripts of *L1TD1*, *SYNJ2BP*, *ARMC1* are highlighted in green. **(B)** and **(C)** DAVID Gene Ontology enrichment of 228 common transcripts identified as L1TD1 targets in both DESeq2 comparisons (DNMT1 KO *versus* input and DNMT1 KO *versus* DNMT1/L1TD1 DKO) in terms of biological processes **(B)** and sequence features **(C)**.

Supplementary Figure 2. GSEA analysis of deregulated proteins in HAP1 DNMT1/L1TD1 DKO versus DNMT1 KO cells. (A) Gene ontology enrichment analysis of differentially expressed proteins upon loss of L1TD1. **(B)** Venn plot of the overlap of mRNAs identified in the RIP-seq analysis and differentially expressed proteins (up and down) identified in the mass spectrometry analysis. The single common mRNA is L1TD1.

Supplementary Figure 3. Ablation of L1TD1 leads to changes in the transcriptome of HAP1 cells. (A and **B)** qRT-PCR analyses of selected L1TD1-associated transcripts. Statistical significance was determined using unpaired two-tailed t-test. All data in the figure are shown as mean of ± SD of 3 biological replicates. ** p ≤0.01, ns not significant. **(C)** Volcano plot illustrating TEtranscript analysis of RNA-seq performed in HAP1 DNMT1 KO and DNMT1/L1TD1 DKO cells.

Supplementary Figure 4. RNA-independent interaction of L1TD1 and ORF1p in HAP1 and OV-90 cells. (A) Western blot analysis of ORF1p IPs with and without RNase A/T$_1$ digestion in DNMT1 KO HAP1 cells using antibodies specific for ORF1p and L1TD1. DNMT1/L1TD1 DKO HAP1 cells were used as a negative IP control and β-actin was used as Western blot loading control. **(B)** Western blot analysis of ORF1p IPs with and without RNase A/T$_1$ digestion in OV-90

cells using antibodies specific for ORF1p and L1TD1 in OV-90 cells. Mouse IgG was used as a negative control for the IP and β-actin was used as Western blot loading control. **(C)** Indirect immunofluorescence in OV-90 cells marks ORF1p (green) and L1TD1 (red) in cytoplasmic granules.

Supplementary Figure 5. L1TD1 enhances LINE-1 retrotransposition. (A) The bar graphs separately show the number of retrotransposition events per 10^6 cells seeded for three independent biological replicates. Blasticidin resistant colonies in pLenti6.2 transfected cells were used for normalization. Statistical significance was determined using unpaired t-test. All data in the figure are shown as a mean of ± SD of 3 technical replicates. ** p ≤0.01**** p ≤0.0001. **(B)** Representative pictures of bromophenol blue stainings of resistant colonies for transfection with the backbone plasmid pCEP4 and the blasticidin deaminase gene containing plasmid pLenti6.2 used as transfection/selection control.

Figure 1

Figure 2

Figure 3

Figure 4

A

B

C **D**

Supplementary Figure S1

A

B

C

Supplementary Figure S2

A

B

Supplementary Figure S3

Supplementary Figure S4

A

B

C

Supplementary Figure S5

A

B