The lnc-FANCI-2 intrinsically restricts RAS signaling in HPV16-infected cervical cancer

Haibin Liu[1a,b], Lulu Yu[1a], Vladimir Majerciak[1], Thomas Meyer[2], Ming Yi[3], Peter F. Johnson[4],
Maggie Cam[2], Douglas R. Lowy[5], and Zhi-Ming Zheng[1]*

[1]Tumor Virus RNA Biology Section, HIV Dynamics and Replication Program, CCR/NCI/NIH

[2]Collaborative Bioinformatics Resource, CCR/NCI/NIH

[3]NCI RAS Initiative, Cancer Research Technology Program, Frederick National Laboratory for
Cancer Research, Frederick, MD, USA.

[4]Mouse Cancer Genetics Program, CCR/NCI/NIH

[5]Laboratory of Cellular Oncology, CCR/NCI/NIH

[a]Co-first authors

Running title: lnc-FANCI-2 regulates RAS signaling

[b]Present address: Center for Emerging Infectious Diseases, Wuhan Institute of Virology, and

Center for Biosafety Mega-Science, Chinese Academy of Sciences, Wuhan, Hubei, 430207,

China

*Corresponding: zhengt@exchange.nih.gov

ABSTRACT

We recently discovered the expression of a long noncoding RNA, lnc-FANCI-2, coinciding with cervical lesion progression from CIN1, CIN2-3 to cervical cancer. Viral E7 of high-risk HPVs and host transcription factor YY1 are two major factors promoting lnc-FANCI-2 expression. To explore possible roles of lnc-FANCI-2 in HPV-induced cervical carcinogenesis, we ablated the expression of *lnc-FANCI-2* in the HPV16-positive cervical cancer cell line, CaSki. Knock-out (KO) single cell clones expressed HPV16 oncogenes normally but displayed altered cell morphology and proliferation when compared with the parental cells. Proteomic profiling of cytosolic and secreted proteins from the parental and KO cells showed that lnc-FANCI-2 regulates expression of a subset of cell soluble receptors responsible for cell signaling, epithelial mesenchymal transition, and IFN responses. RNA-seq analyses revealed that, relative to the parental cells, lnc-FANCI-2 KO cells exhibited significantly increased RAS signaling but decreased IFN pathways. In the KO cells, phosphorylated Akt and Erk1/2, two important RAS pathway effectors, were increased more than 3-fold, accompanied by increase of IGFBP3, MCAM, VIM, and CCND2 (cyclin D2) and decrease of RAC3 expression. High levels of lnc-FANCI-2 and lower levels of MCAM in cervical cancer patients are associated with improved survival. We found that lnc-FANCI-2 interacts specifically with 32 host proteins, including H13, HNRH1, K1H1, MAP4K4, and RNPS1, and knockdown of MAP4K4 in CaSki cells led to an increase of phosphorylation of Erk1/2 and somewhat Akt. In summary, a key function of lnc-FANCI-2 is to intrinsically regulate RAS signaling to impact cervical lesion progression and affect cervical cancer prognosis.

Significance

Expression of lnc-FANCI-2 is related to cervical lesion progression. Knock-out (KO) or knock-down (KD) of lnc-FANCI-2 expression in HPV16-positive cervical cancer cells CaSki significantly increases RAS signaling, phosphorylation of Akt and Erk1/2, and the epithelial mesenchymal transition factors. lnc-FANCI-2 KO also regulates the expression of a subset of cell soluble receptor proteins IGFBP3 and MCAM. A high level of lnc-FANCI-2 and lower level of MCAM in cervical cancer patients are associated with improved survival. lnc-FANCI-2 in CaSki cells interacts specifically with 32 host proteins, including MAP4K4. KD of MAP4K4 expression in CaSki cells led to an increase in phosphorylation of Erk1/2 and Akt. Thus, one lnc-FANCI-2 function is to intrinsically regulate RAS signaling to impact cervical lesion progression.

INTRODUCTION

Cervical cancer is a leading cause of mortality in women worldwide. It is the second most

diagnosed cancer among women, with an estimated 661,000 new cases worldwide each year, and

the third most frequent cause of cancer-related death, accounting for 348,000 deaths annually [1].

Persistent infections in the cervical epithelium with the cancer-associated alpha human

papillomaviruses (high-risk HPVs), in particular HPV16 and HPV18 with the sustained

expression of viral oncoproteins, E6 and E7, contribute to squamous epithelial cell tumorigenesis

[2-4]. Approximate 31.3% of high-grade neoplastic cervical lesions progress to invasive cervical

cancer in 30 years [5], while most lesions resolve spontaneously. The progression to cervical

cancer typically takes 15 to 20 years after initial infection (https://www.who.int/news-room/fact-

sheets/detail/cervical-cancer). These observations suggest that active host defense systems play

important roles in preventing malignant transformation of neoplastic cervical lesions, and

additional factors besides E6 and E7 are necessary for cervical cancer development. Indeed, one

tumor suppressor network, the Fanconi anemia (FA) pathway [6], can be activated by HPV-

induced DNA damage. FA proteins including FANCA, FANCD2 and FANCI are elevated upon

HPV infection [7-9] and the activated FA pathway restricts HPV replication and guards host

genome stability [10]. FANCD2 mutation stimulates HPV16 and HPV31 genome amplification

and also promotes cervical and vaginal cancer development in HPV16 E7 transgenic mouse [11,

12].

HPV-positive cervical cancer tissues in general exhibit wild-type p53, wild-type K-RAS,

and no overexpression of RAS genes [13-16]. Viral oncoprotein E7 plays a major role in the

immortalization of primary epithelial cells mainly by inactivation of pRb family of proteins [17],

whereas E6 mediates degradation of tumor suppressor p53 and activation of human telomerase

reverse transcriptase (hTERT) transcription [18-20]. However, high-risk E6 and E7 are necessary

but not sufficient to transform fully the immortalized cells into malignant cells [21-25] and

additional oncogenic stress is needed. For example, the HRAS[G12V], a constitutively activated

RAS GTPase mutation, triggers malignant transformation of E6/E7-expressing keratinocytes

[26-29]. Normal RAS regulates cell proliferation, cell differentiation and cell adhesion through

cellular signal transduction from growth-factor receptors such as insulin-like growth factors

(IGFs) and insulin-like growth factor binding proteins (IGFBP1-6) [30-32]. Thus, overactive

RAS signaling might facilitate transformation of E6/E7-expressing keratinocytes. In transgenic

mice, HPV16 E6 and E7 can promote the development of spontaneous epithelial skin tumors, but

not spontaneous tumors of the reproductive tract [33-36]. Prolonged estrogen treatment is

required [37]. Estrogen receptor (ER) activation stimulates the mitogen-activated protein kinase

(MAPK/Erk) and phosphoinositide 3-kinase (Pl3K/Akt) pathways, both of which are mediated

by RAS [38]. However, the molecular mechanism underlying estrogen-induced cervical cancer

in E6 and E7 transgene mice is not fully understood.

Long non-coding RNAs (lncRNAs) are RNAs over 200 nucleotides (nt) in length lacking

a coding capacity for translation into functional proteins. Although several lncRNAs have been

proposed to function in diverse biological processes, evidence is still lacking to support the

functionality of the majority of lncRNAs [39, 40].

We recently reported an increased level of lnc-FANCI-2 expression in high-risk HPV-

positive cervical intraepithelial neoplasia (CIN) and invasive cervical cancer (ICC) tissues [8].

At least 14 RNA isoforms of lnc-FANCI-2 are expressed by usage of two alternative promoters,

alternative RNA splicing, and alternative selection of two polyadenylation sites from the

genomic locus adjacent to the FANCI on Chr5. Ironically, the lnc-FANCI-2 was wrongly

annotated by NCBI as a *MIR9-3HG* in the recently updated RefSeq database

(https://www.ncbi.nlm.nih.gov/refseq/) and two alternative lnc-FANCI-2 promoters identified

for expression of lnc-FANCI-2 are ~10 kb downstream of the non-expressible miR9-3 gene [8].

Both lnc-FANCI-2 and FANCI are up-regulated simultaneously in neoplastic cervical lesions

and cervical cancer by high-risk HPV infections. E6 but, in particular E7 are responsible for the

enhanced expression of lnc-FANCI-2, the transcription of which is also regulated by YY1[8].

HPV infection increases YY1 levels but decreases the expression of p53-dependent miR-29a

which targets the YY1 3′ UTR. Viral E7 interacts with YY1 and facilitates YY1 transactivation

of lnc-FANCI-2 promoter [8]. In situ hybridization has revealed that lnc-FANCI-2 is

preferentially cytoplasmic, but its function in HPV infected cells has not been characterized. In

this report, we demonstrate that lnc-FANCI-2 in HPV16-infected cells controls RAS signaling

by interaction with MAP4K4 and other RNA-binding proteins. Ablation of lnc-FANCI-2 in the

cells promotes RAS signaling and phosphorylation of Akt and Erk. High levels of lnc-FANCI-2

and low level of MCAM expression in cervical cancer patients correlate with improved survival,

indicating that lnc-FANCI-2 plays a critical role in regulating RAS signaling to affect cervical

cancer progression and patient outcomes.

RESULTS

Differential expression of lnc-FANCI-2 in cervical cancer tissues and its derived cell lines

As lnc-FANCI-2 expression is up-regulated along with cervical lesion progression by high-risk

HPV infections [8], we further verified the increased expression of lnc-FANCI-2 in cervical

cancer tissues by RNAscope single molecule RNA *in situ* hybridization (RNA-ISH) using a lnc-

FANCI-2 antisense probe spanning over the major isoform of lnc-FANCI-2 nt 359-1713 region

(GenBank MT669800.1). We found both cytoplasmic and nuclear locations of the increased lnc-

FANCI-2 ISH signals within the tumor nest in HPV16-infected cervical squamous cell

carcinoma tissues, but not much so in its adjacent tissue areas (Fig. 1A/B). The increased

expression of lnc-FANCI-2 became obvious in human foreskin keratinocytes with HPV16 or

HPV18 infection when compared to the HFK cells without HPV infection (Fig. 1C).

However, we subsequently demonstrated the high expression of lnc-FANCI-2 in two

HPV16-infected cervical cancer cell lines (SiHa and CaSki) and two HPV16-infected low-grade

cervical lesion-derived cervical keratinocyte subclone lines (W12 20861 and W12 20863 cells),

but not in two HPV18-infected cervical cancer cell lines (HeLa and C4II cells), nor other HPV-

negative cell lines of HCT116 (colorectal cancer cells), BCBL-1 (body cavity B lymphoma

cells), HEK293 (Ad5 E1/E2-immortalized human kidney cells), and HaCaT (spontaneously

immortalized human epidermal cells) (Fig. S1A). Interestingly, the high expression of lnc-

FANCI-2 was observed in a HPV-negative cervical cancer cell line C33A cells with mutations of

both p53 and RB genes [41] (Fig. S1A). Northern blot confirmed the increased expression of lnc-

FANCI-2 in C33A cells but no expression in HeLa cells (Fig. S1B).

Subcellular distribution of lnc-FANCI-2 was characterized by cell fractionation and

Western blot assays using nuclear SRSF3 and cytoplasmic GAPDH protein as an indication of

fractionation efficiency (Fig. S1C). By RT-qPCR analysis of the fractionated total RNA, we

demonstrated that lnc-FANCI-2 RNA is mainly cytoplasmic in HPV16-positive CaSki cells, but

nuclear in HPV16-positive SiHa and HPV-negative C33A cells (Fig. S1C). The differential

subcellular distributions of lnc-FANCI-2 RNA from CaSki to SiHa and C33A cells (Fig. S1D)

were further confirmed by RNAscope RNA-ISH using the lnc-FANCI-2 antisense probe

described above.

lnc-FANCI-2 regulates proliferation of HPV-transformed cervical cancer cells

lnc-FANCI-2 RNA is transcribed mainly from a proximal promoter TSS2, but also from

an alternative minor, distal promoter TSS1. Two highly conserved YY1 binding motifs upstream

of the TSS2 are essential for the TSS2 transcriptional activity [8] (Fig. 2A). To elucidate the

function of lnc-FANCI-2 in high-risk HPV-infected cervical cancer cells, we knocked out (KO)

lnc-FANCI-2 expression in HPV16-positive CaSki cells using CRISPR/Cas9 deletion of either a

3-kb promoter region encompassing both TSS1 and TSS2 or an 86-bp region containing two

YY1-binding motifs in the TSS2 promoter (Fig. 2A). Two tested gRNAs with high KO

efficiency were selected and cloned into a modified CRISPR/Cas9 expression vector to express

both 5′-specific gRNA and 3'-specific gRNA simultaneously for efficient genome editing [42].

CaSki cells stably transfected with the dual gRNA expression vector were generated. Through a

serial dilution, several single cell clones were isolated and verified for homozygous deletion by

PCR screening. PCR genotyping indicated successful homozygous deletion of the promoter or

YY1-binding motifs, respectively (Fig. 2B).

Loss of lnc-FANCI-2 expression in the single-cell clones was examined by RT-qPCR

(Fig. 2C). When compared to all control clones transfected and selected from an empty vector

containing no gRNA, all single cell clones with deletion of YY1-binding motifs (ΔYY1) showed

70-80% reduction of lnc-FANCI-2 expression, whereas the single cell clones with deletion of

both TSS1 and TSS2 promoters (ΔPr) showed >90% reduction (Fig. 2C). Northern blot analysis

confirmed the decrease in levels of the two major lnc-FANCI-2 isoforms (Fig. 2D), with an

abundant 4-kb lnc-FANCI-2 RNA derived from a distal polyadenylation site pA2 and a less

abundant 2-kb of lnc-FANCI-2 RNA derived from a proximal polyadenylation site pA1[8] (Fig.

2A and 2D). Reduced lnc-FANCI-2 RNA expression was also confirmed by RNAscope RNA-

ISH (Fig. 2E and 2F). The parental (WT) CaSki cells expressed ~10 copies of lnc-FANCI-2

RNA per cell [8], whereas the copy number of lnc-FANCI-2 RNA in the ΔYY1-D5 cells

dropped to ~5 copies and to ~2 copies per cell in the ΔPr-A9 cells (each averaged from 200

cells).

Although the WT CaSki cells preferentially grow as distinct cluster cell islands, the lnc-

FANCI-2 KO cells displayed a dispersed cell growth pattern and some cells exhibited an

irregular or spindle-like cell morphology (Fig. S2). The KO cells also grew slower than the

parental CaSki cells (Fig. 1G), formed fewer colonies with reduced size (Fig. 1H), and showed

decreased migration capacities (Fig. 1I), with ΔPr-A9 cells having a more severe phenotype than

ΔYY1-D5 cells. Subsequently, the ΔPr-A9 and ΔPr-B3 cells were further examined for their

HPV16 E6 and E7 expression and the ΔPr-A9 for cell senescence. We found both ΔPr-A9 and

ΔPr-B3 cells, when compared with parental WT CaSki cells, exhibited some small, variable

changes in the expression of E6, E7, p53 (E6 downstream target) and E2F1 (E7 downstream

target) proteins (Fig. S3A), most likely from sampling in the assays, but the ΔPr-A9 cells

displayed significant increase in cell senescence (Fig. S3B). Altogether, the data indicate that

lnc-FANCI-2 RNA is important for cell growth, colony formation, and cell migration.

lnc-FANCI-2 regulates the expression and secretion of cell soluble receptors

Considering that the altered expression of cell membrane proteins and secreted factors in

the lnc-FANCI-2 KO cells might contribute to the observed different cell morphology and

growth properties, we performed a Proteome Profiler Human sReceptor Array analysis to

examine possible changes in expression of 105 well-characterized soluble protein receptors

being involved in cell signaling using total cell lysates and cell culture supernatants from ΔPr-A9

and WT CaSki cells.

Using 30% cut-off (FC -/+0.3), we identified from the ΔPr-A9 cell lysate nine proteins

with increased expression compared to the WT CaSki cells, including PODXL2, ECM1,

NECTIN2, MCAM, ADAM9, ADAM10, CDH5, ITGA5 and NOTCH1, and six proteins with

decreased expression including ITGB6, CDH13, LGALS3BP, TIMP2, ADAM8 and SCARF2

(Fig. 3A and 3B). We also found from the ΔPr-A9 cell culture supernatant five proteins with

increased expression, including ADAM9, NECTIN2, ADAM10, CDH5 and ECM1, and the

decreased expression of four proteins, including CRELD2, SDC1, SDC4 and TIMP2 (Fig. 3A

and 3B). By immunoblot assays, we verified selectively the increase in PODXL2, MCAM, and

ECM1 and the decrease in ADAM8 and TIMP2 from the ΔPr-A9 cell lysates and the increase in

ECM1 and the decrease in TIMP2 in the ΔPr-A9 cell culture supernatant (Fig. 3C). As ADAM8

is proteolytically processed into two protein isoforms for cell adhesion [43] and migration[44],

we confirmed by immunoblot the decreased expression of all three sizes of ADAM8 protein in

the ΔPr-A9 cell lysate over the WT CaSki cells (Fig. 3C).

lnc-FANCI-2 regulates the expression of genes involved in RAS signaling

We next conducted genome-wide RNA-seq analyses of WT and lnc-FANCI-2 KO cells

(four samples/group) to determine the transcriptomic consequences of lnc-FANCI-2 deficiency.

We obtained ~120 million mappable RNA reads to the human reference genome hg38 from each

sample. Analysis of RNA-seq reads-coverage map by Integrative Genomics Viewer (IGV)

showed ~90% reduction of lnc-FANCI-2 expression in ΔPr-A9 cells and ~72% decrease in

ΔYY1-D5 cells compared to the WT CaSki cells (Fig. 4A), consistent with the results shown in

Fig. 2C-F.

Hierarchical clustering analysis showed more global transcriptional similarity between

ΔYY1-D5 and ΔPr-A9 cells (Fig. 4B). By applying a threshold of fold change (FC) ≥ 1.8 or FC

≤ -1.8 with FDR ≤ 0.01, we found 1230 genes in ΔPr-A9 and 797 genes in ΔYY1-D5 cells with

significantly differential expression relative to the WT CaSki cells expressing ~15890 genes.

Nine (*Italic*) of 15 (60%) soluble receptor proteins in cell lysates (PODXL2, ECM1, *NECTIN2,*

MCAM, ADAM9, CDH5, ADAM10, *ITGA5, NOTCH1*, SCARF2, ADAM8, *TIMP2, LGALS3BP*,

CDH13, and *ITGB6*) exhibited consistent changes in expression by both RNA-seq and protein

array assays. The most significantly affected genes are shown in Volcano plots (Fig. 4C and

Table S1). Among these, 211 were upregulated and 189 were downregulated in both ΔPr-A9 and

Δ YY1-D5 cells (Fig. 4D and Table S2). By applying more stringent criteria with FPKM ≥7 in at

least one of four RNA-seq samples as a cutoff, we profiled the genes with large expression

differences in both ΔPr-A9 and ΔYY1-D5 cells. The results displayed in the heatmap of Fig. 4E

included 52 upregulated, and 47 downregulated genes excluding lnc-FANCI-2. Subsequently, we

selectively verified by RT-PCR the decreased expression of ITGB6, NLRP2, PLAC8, PSG4,

PSMB9, and SERPINB1 and the increased expression of CFH, CNTN5, EMP3, IGFBP3,

PTGS2, and SERPINB2 in ΔPr-A9 cells (Fig. 4F), as well as the increased expression of

IGFBP3 protein, a RAS signaling driver [30-32], in ΔPr-A9 cells by immunoblot (Fig. 4G).

By performing the Gene Set Enrichment Analysis (GSEA) on the Hallmark gene sets,

which provide more refined and concise inputs for GSEA [45], we found the most significantly

upregulated pathways in both ΔPr-A9 and ΔYY1-D5 cells, when compared with the WT CaSki

cells, were KRAS signaling and epithelial mesenchymal transition (EMT). The most

significantly downregulated pathways in both ΔPr-A9 and ΔYY1-D5 cells over the WT CaSki

cells were interferon gamma (IFN-γ) and interferon alpha (IFN-α) responses (Fig. 5A). GSEA

plots for ΔPr-A9 shows 29 of 111 genes, including IGFBP3 [30-32], in RAS signaling and 39 of

130 genes in EMT (Fig. 5B and 5C, Table S3) were significantly upregulated, whereas 37 of 139

genes in IFN-γ response and 24 of 71 genes in IFN-α response were significantly downregulated

(Fig. 5D and 5E, Table S3). Similar enriched gene sets with increased RAS signaling and EMT

and decreased IFN-γ and IFN-α responses were observed in ΔYY1-D5 cells (Fig. S4, Table S3).

Since ΔPr-A9 cells exhibit higher KO efficiency of lnc-FANCI-2 and a higher number of

differentially expressed genes (DEGs) and show more severe phenotypic changes in cell

proliferation, migration, and colony formation than ΔYY1-D5 cells, the ΔPr-A9 cells were

primarily used for further focused studies on RAS signaling in this report.

lnc-FANCI-2 regulates RAS GTPase activities to phosphorylate RAS signaling effectors

Akt and Erk

RAS activation triggers two major downstream signal transduction pathways,

Raf/Mek/Erk and PI3K/Akt (Fig. S5), to transduce signals from extracellular stimuli to the cell

nucleus where specific effector genes are activated for their corresponding functions [30-32, 46,

47]. To further confirm the GSEA data and verify that two major signaling pathways of RAS are

activated, we first examined and compared the RAS GTPase activities in ΔPr-A9 cells and the

WT CaSki cells. We found a significant increase in RAS GTPase activity in ΔPr-A9 cells (Fig.

6A, top bar graphs), indicating that the endogenous lnc-FANCI-2 RNA in CaSki cells suppresses

not only the expression of IGFBP3 (Fig. 4F and 4G), the most abundant IGFBP, but also RAS

activation (Fig. 6A, top bar graphs). Since ΔPr-A9 cells had undergone long term selection and

adaption during single cell screening, the increased RAS GTPase activity might result from

selection pressure for cell survival. However, by transient siRNA knockdown (KD) of lnc-

FANCI-2 expression in the WT CaSki cells, we obtained a similar, albeit weaker, increase of

RAS GTPase activity (Fig. 6A, lower bar graphs). This result suggests that the increased RAS

GTPase activity in ΔPr-A9 cells was unrelated to the persistent selection pressure, but rather the

deficiency of lnc-FANCI-2.

Next, we examined activation of the PI3K/Akt and Raf/Mek/Erk pathways [48-50]. We

observed a 3-fold increase in phosphorylated Akt (p-Akt) and 2.5-fold increase of p-Erk1/2

(p44/p42 MAPK), after being normalized to total Akt and Erk levels, in ΔPr-A9 cells over the

WT CaSki cells (Fig. 6B). These results were confirmed in ΔYY1-D5 cells (Fig. S6A). The

increased p-Akt and p-Erk (mostly p-Erk2/p42) was accompanied by elevated expression of

MCAM, VIM, and CCND2, and decreased expression of RAC3 (Fig. 6B and Fig. S6A). These

randomly chosen potential RAS effector proteins facilitate soluble receptor functions, cell

proliferation, and EMT. Moreover, transient siRNA knockdown of lnc-FANCI-2 in the WT

CaSki cells also led to the increased levels of p-Akt and p-Erk (Fig. 6C) and increased

expression of MCAM and VIM at 48 h and 96 h post-transfection (Fig. 6C). These observations

indicate that a transient loss of lnc-FANCI-2 in CaSki cells is sufficient to trigger RAS signaling.

The same siRNA transfection in HeLa cells, a lnc-FANCI-2 negative cell line (Fig. S1A and

S1B), exhibited no effect on p-Akt or p-Erk1/2 levels (Fig. S6B), but in SiHa cells containing

mainly nuclear lnc-FANCI-2 (Fig. S1C and S1D) enhanced the expression of p-Akt and p-Erk1/2

(Fig. S6B), verifying the specific effect of lnc-FANCI-2 depletion on RAS signaling pathways

independently of predominant cellular distribution of lnc-FANCI-2. The lnc-FANCI-2 KO-

mediated increase in p-Akt and p-Erk was sensitive to PI3K inhibitor LY294002 [51] (Fig. 6D,

lanes 4-6) and MEK1/2 inhibitor U0126 [52] (Fig. 6D, lanes 7-9), respectively. The inhibitors

also decreased the expression of VIM and MCAM and cell proliferation, in particular by

MEK1/2 inhibitor U0126 (Fig. 6D and 6E). The effects of LY294002 on cell proliferation were

similar from ΔPr-A9 to the WT CaSki cells (Fig. 6E).

lnc-FANCI-2 inhibits the expression of IGFBP3 and MCAM (CD146 or MUC18)

Given that IGFBP-3 is the most abundant IGFBP among all six IGFBP members in potentiation of IGF action and PI3K/AKT activities [30] and a reduced expression of IGFBP-3 mRNA level is associated with progression to cervical cancer [53], we further investigated the effect of lnc-FANCI-2 on the expression of IGFBP3 as lnc-FANCI-2 effector gene. As shown in Fig. 4F-G and Fig. S7A, the increased RNA and protein expression of IGFBP3 appears in the lnc-FANCI-2 KO ΔPr-A9 cells over the parental WT CaSki cells, indicating a suppressive effect of lnc-FANCI-2 on IGFBP3 expression. This suppressive function of lnc-FANCI-2 was further confirmed in lnc-FANCI-2 rescue experiments in the ΔPr-A9 cells (Fig. 7A and 7B). By transient expression of one major isoform of lnc-FANCI-2 RNA (a-PA2, GenBank: MT669800.1) [8] in the ΔPr-A9 cells, lnc-FANCI-2 (red) and IGFBP3 RNA (green) at 24 h post transfection were detected by RNAscope RNA-ISH using each specific antisense RNA probe (Fig. 7A). We demonstrated that the cells with rescued expression of cytoplasmic lnc-FANCI-2 RNA displayed much reduced expression of cytoplasmic IGFBP3 RNA (Fig. 7B).

As shown in Fig. 4E and Fig. S7B, the increased RNA expression of MCAM (CD146 or MUC18) [54] appears to be transcriptional or posttranscriptional in both ΔPr-A9 and ΔYY1-D5 cells. We confirmed by RT-PCR the lnc-FANCI-2 KO-mediated increase of MCAM RNA expression in ΔPr-A9 cells (Fig. 7C). A rescue experiment was further performed by transient expression of one major isoform of lnc-FANCI-2 RNA (a-PA2, GenBank: MT669800.1) [8] in ΔPr-A9 cells to determine both nuclear and cytoplasmic MCAM expression [55, 56] in individual lnc-FANCI-2 expressing cells. ΔPr-A9 cells ectopically expressing diffused cytoplasmic lnc-FANCI-2 RNA, as detected by RNAscope RNA-ISH, exhibited a marked reduction in nuclear MCAM protein (Fig. 7D-7E). Quantitative analyses of the ΔPr-A9 cells with

or without transient lnc-FANCI-2 RNA expression showed significant reduction of nuclear

MCAM protein in the 35 cells with rescued lnc-FANCI-2 expression when compared to the 54

cells expressing no lnc-FANCI-2 (Fig. 7F). Due to RNAscope procedures, the membrane-bound

and cytoplasmic MCAM protein was mostly removed by protease III treatment when RNA-ISH

was performed and thus only the nuclear signal of MCAM protein remained was detectable by IF

staining. Using cell fractionation, we detected a cleaved MCAM of ~46 kDa mainly in the

nuclear fraction by Western blot (Fig. 7G). The increased nuclear MCAM expression in ΔPr-A9

cells was proportional to the cytoplasmic and total MCAM levels when compared with the WT

CaSki cells (Fig. 7G). The nuclear MCAM signal appeared as a proteolytic cleavage product

presumably by the increased metalloprotease ADAM9 and ADAM10 proteins and decreased

metalloprotease inhibitor TIMP2 (Fig. 3), as observed in other studies [56].

The significance of an inverse correlation of lnc-FANCI-2 and MCAM in CaSki cells

was further investigated by analyzing their expression in 304 cervical cancer samples from the

TCGA dataset. There was a significant negative correlation in lnc-FANCI-2 (being wrongly

assigned as LINC00925 by NCBI) [8] and MCAM RNA levels (Fig. 7H). The opposite effects of

lnc-FANCI-2 and MCAM on cervical cancer survival (Fig. 7I and Fig. S8) show that the cervical

cancer patients with a higher level of lnc-FANCI-2 [8] but a lower level of MCAM in the

cervical tissue (Fig. S8 and Fig. 7I) exhibited a better survival prognosis.

Regulatory roles of lnc-FANCI-2-mediated increase of MCAM and IGFBP3 on RAS

signaling

To further dissect the lnc-FANCI-2-associated expression of MCAM and IGFBP3 on

RAS signaling, we examined MCAM and IGFBP3 on phosphorylation of Akt and Erk1/Erk2 in

CaSki cells in the presence or absence of lnc-FANCI-2. As shown in Figure 7J, we found,

although KD or KO of lnc-FANCI-2 promotes phosphorylation of Akt and Erk (Fig. 7J, lane 2),

that KD of MCAM expression in the parental WT CaSki cells in the absence of lnc-FANCI-2

(Fig. 7J, lane 3) or in ΔPr-A9 cells (Fig. 7J, lane 5) led to reduction of Akt and Erk

phosphorylation. These data suggest that MCAM is a lnc-FANCI-2 effector but also could be a

trigger of signal transduction as reported [54, 57].

IGFBP3 protein has been viewed as a RAS signaling regulator [30-32], but recent studies

show that IGFBP3 has a variety of intracellular ligands involved in many unexpected functions

[58, 59]. Thus, KD or KO of lnc-FANCI-2-mediated IGFBP3 expression on RAS signaling in

the parental WT CaSki cells or ΔPr-A9 cells was examined by Western blot after siRNA KD of

IGFBP3 expression. As shown in Figure 7K, KD of IGFBP3 expression was found to increase

phosphorylation of Erk1/2 by ~70% (lane 4) to ~60% (lane 6), but not much so for Akt

phosphorylation (lanes 4 and 6). Instead, KD of IGFBP3 expression could prevent lnc-FANCI-2

KD-enhanced Akt phosphorylation in the parental WT CaSki cells (Fig. 7K, compare lane 2 and

lane 3). These data suggest a separate role of IGFBP3 on phosphorylation of Erk1/2 from Akt in

the presence or absence of lnc-FANCI-2.

MAP4K4 association with lnc-FANCI-2 RNA in CaSki cells regulates RAS signaling and

phosphorylation of Akt and Erk

We speculated lnc-FANCI-2 restriction on RAS signaling through interactions with

cellular proteins. Subsequently, comprehensive identification of lnc-FANCI-2-binding proteins

in the parental WT CaSki cells was performed by an IRPCRP protocol (modified from published

ChIRP [60]) in combination with mass spectrometry (Fig. 8A). In this protocol, the lnc-FANCI-

2-binding proteins in the parental WT CaSki cells were covalently crosslinked to lnc-FANCI-2

RNA via UV irradiation and pulled down by pooled 30 antisense oligos crossing over the entire

lnc-FANCI-2. The RNA extracted from the pulldowns were verified to be lnc-FANCI-2 specific

by RT-PCR in the absence (-) or presence (+) of reverse transcriptase (RT) (Fig. 8B). The lnc-

FANCI-2-binding proteins in the pulldowns were then subjected to LC-MS/MS analyses. We

identified 32 specific lnc-FANCI-2-binding proteins, including H13, HNRH1, K1H1, MAP4K4,

and RNPS1 (Fig. 8C, Table S4).

MAP4K4, a serine/threonine protein kinase, stimulates cancer cell proliferation, invasion

and migration, and is recently characterized as a novel MAPK/ERK pathway regulator [61-63]

and negatively regulates RAS signaling by binding to Ras p21 protein activator 1 or RASA1

[64, 65]. Although the exact MAK4K4 binding site on lnc-FANCI-2 was not explored, many

enzymes turn out an RNA-binding enzyme [66, 67]. Thus, whether MAP4K4 association with

lnc-FANCI-2 could regulate PI3K/Akt or MAPK/Erk signal transduction was investigated in the

parental WT CaSki cells. We demonstrated that siRNA KD of MAP4K4 expression in the

parental WT CaSki cells, as seen for KD of lnc-FANCI-2, led to ~40% or more increase of

phosphorylation of both Akt and Erk1/2 and increased expression of MCAM and IGFBP3 (Fig.

8D, compare lane 2 to lane 3). Interestingly, siRNA KD of MAP4K4 expression in the lnc-

FANCI-2 KO ΔPr-A9 cells had no effect on expression of IGFBP3, only a minimal effect on

MCAM, but strongly on pErk1/2 (Fig. S7C). The data suggest that, through RNA-protein

interactions, the increased lnc-FANCI-2 RNA in cells and their association with MAP4K4 and

other cellular proteins could be one arm to control RAS signaling and gene expression of its

effectors (Fig. 8E).

DISCUSSION

HPV oncoproteins E6 and E7 are necessary but not sufficient for development of cervical cancer. Indeed, early studies indicated that other oncogenic stress, such as activated RAS mutant, is needed to trigger malignant transformation of E6 and E7 expressing cells [21, 26, 68, 69] and tumorigenesis [27, 28, 70-74]. Moreover, E7 has been also reported to repress phosphorylation of Akt and Akt-mediated signaling [75]. In this study, we show that lnc-FANCI-2, whose expression is highly dependent on E7 and YY1 [8], intrinsically restricts RAS GTPase activities and phosphorylation of Akt and Erk presumably by interacting with cellular factors in HPV16-positive CaSki cells. Interestingly, early reports indicated that p-Akt attenuation by E7 could be abolished by introduction of H73E mutation in the E7 CR3 domain [75]. This E7 CR3 domain is also essential to interact with transcription factor YY1 to activate lnc-FANCI-2 transcription [8]. Thus, it would be presumable that the reported p-Akt attenuation by E7 might be mediated by the increased expression of lnc-FANCI-2. More importantly, these findings also suggest that additional oncogenic stress, such as activated RAS mutant, is required to overcome the lnc-FANCI-2 restriction on RAS signaling for the early observed malignant transformation [21, 26, 68, 69] and tumorigenesis [27, 28, 70-74] of E6 and E7 expressing cells. In CaSki cells, loss of lnc-FANCI-2 in this report was found to promote RAS signaling but reduction of IFN responses. However, high RAS-AKT-ERK signaling induces cell senescence and inhibits tumor cell survival as shown in this report and in a lung cancer study [76].

Some lncRNAs function as RAS regulators [77, 78] by their ability to sequester RAS-targeting miRNAs, including MALAT1/ miR-217, RMRP/miR-206, and KRAS1P/miR-143/let-7 [78]. Conceivably, lncRNAs may also regulate RAS signaling through other mechanisms beyond miRNAs. In this report, loss of lnc-FANCI-2 in HPV16-positive CaSki and SiHa cells

promotes RAS signaling and expression of RAS signaling effectors (Fig. 8E). By profiling the

proteins associated with lnc-FANCI-2 in CaSki cells, we identified a serine/threonine protein

kinase MAP4K4 as one of major lnc-FANCI-2-binding proteins that may partially mediate the

lnc-FANCI-2 restriction on RAS signaling. MAP4K4 is a negative RAS signaling regulator in

the context of the early embryo [62, 64] and lymphatic vascular development by interacting with

RASA1 [65]. Indeed, silencing of MAP4K4 in parental WT CaSki cells, like KD or KO of lnc-

FANCI-2, led to partially increase of both p-Akt and p-Erk1/2 (Fig. 8D) as reported [64, 65].

Interestingly, KD of MAP4K4 in ΔPr-A9 cells led to further increased expression of p-Erk1/2

but not p-Akt, suggesting a negative correlation of MAP4K4 activity in interaction with lnc-

FANCI-2 on Erk1/2 phosphorylation. Thus, our data suggest that MAP4K4 protein may

function coordinately with lnc-FANCI-2 RNA in the CaSki cells to restrict one arm but not the

other of RAS signaling.

All HR-HPVs are capable of inducing lnc-FANCI-2 expression by experimental infection

of keratinocytes [8]. HPV[+] cervical cancer tissues express high level of lnc-FANCI-2. As

expected, all HPV16[+] cervical cancer and pre-cancer cell lines examined, such as CaSki, SiHa,

W12 subclone cells 20861(HPV16[+], integrated) and 20863 (HPV16[+], episomal), produce lnc-

FANCI-2. Unexpectedly, we found no expression of lnc-FANCI-2 in isogenic HPV18[+] HeLa

and C4II cells, nor in colorectal cancer cell line HCT116, adenovirus 5 DNA-transformed human

embryonic kidney cell line HEK293, spontaneously transformed aneuploid immortal

keratinocyte cell line HaCaT, and KSHV-infected B cell lymphoma cell line BCBL-1 cells

which are all HPV-negative cell lines. It remains to know what negatively regulates lnc-FANCI-

2 expression in those cell lines and more HPV18[+] cell lines should be examined for lnc-FANCI-

2 expression. However, we did find that a HPV-negative cervical cancer cell line, C33A cells

expressing mutant p53 and mutant pRB [41], produce a high level of lnc-FANCI-2 (Fig. S1A

and S1B). Using dual luciferase report assays, we did find lnc-FANCI-2 promoter activity in

response to YY1 binding in CaSki and C33A cells, but not in HeLa and HCT116 cells,

suggesting the presence of a repressive factor (s) in the latter two cell lines (data not shown). KD

of E2F1 had no effect on lnc-FANCI-2 promoter activity in CaSki cells [8]. In the lnc-FANCI-2

producing cells, we noticed that cellular location of lnc-FANCI-2 also varies from cell types,

with lnc-FANCI-2 predominantly in the nucleus in SiHa and C33A cells, but in the cytoplasm in

CaSki cells, although both cytoplasmic and nuclear lnc-FANCI-2 restrict RAS signaling. This

suggests that a variety of nuclear and cytoplasmic functions of lnc-FANCI-2 remains to be

explored.

IGFBP3 is a major IGF carrier and enhances both IGF- and EGF-RAS signaling [31, 79]

to activate phosphatidylinositol 3-kinase (PI3K)/AKT and the mitogen-activated protein kinase

(MAPK)/ERK pathways [80]. Surprisingly, we find a suppressive role of IGFBP3 on Erk

phosphorylation, but not much on Akt, in the presence or absence of lnc-FANCI-2 in HPV16-

positive CaSki cells. Although ~99% of circulating IGF-1 is bound to IGFBPs, predominantly to

IGFBP-3, the most abundant IGFBP in human serum, case-control studies showed significantly

lower IGF-1 and IGFBP-3 serum levels in the patients with invasive cervical cancer over the

control group [81, 82]. A reduced expression of IGFBP-3 mRNA level appears to be associated

with progression to cervical cancer [53]. Consistently with lnc-FANCI-2's suppressive effect on

IGFBP3 expression in our study, an increased level of lnc-FANCI-2 expression is associated

step-wisely with high-risk HPV-positive cervical intraepithelial neoplasia and invasive cervical

cancer tissues [8].

It is obvious from our RNA-seq and human soluble receptor array analysis that KO of

lnc-FANCI-2 in CaSki cells affects the expression of a large set of genes, including the

expression of increased NECTIN2, ADAM9, ADAM10, ITGA5, MCAM, PODXL2, ECM1, but

decreased ADAM8, TIM2, and ITGB6, which are responsible for RAS signaling, EMT pathway,

and IFN responses. For example, the dysregulation of RAS signaling and ADAM protein activity

is implicated in various cancers. ADAM proteins can modulate RAS signaling by cleaving and

releasing ligands that activate or inactivate RAS-related pathways [83-86]. Some ADAM

proteins are Involved in the migration and invasion of cancer cells, and its loss can promote the

degradation of KRAS [87]. ADAMs are essential for extracellular signaling and regulation of

cell adhesion and their catalytic activity is strongly correlated to Src-homology 3 (SH3) domain

binding of SH3 proteins [85, 86, 88]. In addition to the notified activation of PI3K/Akt and

Raf/Mek/Erk pathways, one of the major RAS signaling effectors in lnc-FANCI-2 KO or KD

cells was found to be MCAM.

MCAM binds various cellular surface receptors or co-receptors to trigger signal

transduction, cell proliferation and motility, tumor angiogenesis, and tumor cell metastasis [54,

57]. It is an integral, highly glycosylated membrane protein with three common protein isoforms

[54, 89, 90]. Two membrane-anchored forms with a long (p113) or short (p76) cytoplasmic tail

produced by alternative RNA splicing and a proteolytically cleaved soluble form without

transmembrane and cytoplasmic regions [54, 90, 91]. In addition to its roles in cell signal

transduction [54, 57, 89, 90], the short isoform produced by an intramembrane cleavage may be

directed towards the nucleus to regulate gene transcription [56]. We found that CaSki cells

express mainly a cytoplasmic form of full-length MCAM (~113 kDa) and a small nuclear form

(~46 kDa), but not the short ~76 kDa isoform. Moreover, lnc-FANCI-2 KO cells exhibit high

MCAM levels, which was most likely a result of the increased RAS signaling. Consistently, lnc-

FANCI-2 and MCAM levels are negatively correlated in cervical cancer patients, and low levels

of MCAM and high levels of lnc-FANCI-2 indicate better prognosis. This suggests that

increased MCAM might promote cervical tumorigenesis in vivo. Given that the cells with lnc-

FANCI-2 KO exhibit increased levels of IGFBP3 and MCAM and the ΔPr-A9 cells with rescued

lnc-FANCI-2 RNA expression displayed remarkable reduction of IGFBP3 and MCAM

expression, our current model (Fig. 8E) highlights how the HPV16-enhanced expression of lnc-

FANCI-2 in cervical cancer cells might function as a negative regulator by binding to various

RNA-binding proteins to block RAS signaling and reduce the expression of RAS signaling

effectors such as IGFBP3 and MCAM, etc.

475   In conclusion, this study provides the first evidence that one of lnc-FANCI-2 functions is

to maintain epithelial functional integrity by repressing RAS signaling in preventing malignant

transformation of neoplastic cervical lesions. In addition, more extensive studies are undergoing

to elucidate the role of lnc-FANCI-2 in regulation of IFN responses, which are not covered in

this report. Since transactivation of lnc-FANCI-2 expression is largely dependent on host

transcription factor YY1 of which expression is regulated by HPV oncoproteins and host miR-

29a, dissecting their transcription network on lnc-FANCI-2 and its effectors may further provide

novel insights involved in host homeostasis against HPV infection-induced carcinogenesis. Our

observations also highlight additional, unexpected players beyond viral E6 and E7 that drive

cells toward malignant transformation.

MATERIALS AND METHODS

Cell cultures and treatment

CaSki, a HPV16-positive (HPV16[+]) human cervical cancer cell line with wild-type p53, pRb,

HRAS, KRAS, and NRAS (https://depmap.org/portal/cell_line/ACH-001336?tab=mutation) was

obtained from the American Type Culture Collection (ATCC, Manassas, VA), maintained in

Dulbecco's modified Eagle medium (DMEM) (Thermo Fisher Scientific) supplemented with

10% fetal bovine serum (FBS) and 1% penicillin-streptomycin, and grown in a cell culture

incubator at 37°C with 5% CO_2.

SiHa (HPV16[+]), HeLa (HPV18[+]), C4II (HPV18[+]), C33A (no HPV), and other HPV-

negative cell lines, HEK293T (Ad5 E1/E2-immortalized human kidney cells), and HaCaT

(spontaneously immortalized human epidermal cells), were all obtained from ATCC and grown

in Dulbecco's modified Eagle's medium (DMEM) with 10% FBS at 37°C and 5% CO_2. HCT116

cell line, a colorectal cancer cell line, was a gift from Dr. Bert Vogelstein of Johns Hopkins

University and was grown in McCoy's 5A medium with 10% FBS at 37°C and in 5% CO_2. W12

subclones 20861 (HPV16[+], integrated) and 20863 (HPV16[+], episomal) were a gift from Dr. Paul

Lambert of University of Wisconsin, Madison, and grown on mitomycin C-pretreated NIH3T3

feeder cells in F12 medium at 37°C and in 5% CO_2. BCBL-1 (body cavity B lymphoma cells

with KSHV infection) was obtained from the AIDS Research and Reference Reagent Program,

Division of AIDS, NIAID, NIH and cultured in RPMI 1640 containing 10% FBS at 37°C and in

5% CO_2.

CaSki cells were treated with 20 μM phosphoinositide 3-kinase (PI3K) inhibitor,

LY294002 (Cell Signaling Technology, Danvers, MA, #9901) or 10 μM mitogen-activated

protein kinase kinase (MEK) inhibitor, U0126 (Cell Signaling Technology, Danvers, MA,

#9903) and examined for their proliferation and viability by CCK-8 assay and RAS effector

expression by Western blot.

Raft tissues and cervical tissue sections

Raft tissues with or without HPV16 or HPV18 infection and cervical cancer tissue sections from Zhejiang University Women's hospital were described in our previous reports [8, 92].

siRNAs, plasmid constructions and transfections

Synthetic double-stranded siRNA targeting the lnc-FANCI-2 transcript is listed in a supplementary Table S5. Human MAP4K4 siRNA (#M-003971), human IGFBP3 siRNA (#M-004777) and non-targeting control siRNA(#D-001210-01) were purchased from Dharmacon, Inc (Lafayette, CO). Each siRNA transfection into CaSki or other type of cells was carried out by LipoJet in vitro transfection reagent (SignaGen, Frederick, MD, USA). Plasmid pHBL03 has a short isoform of lnc-FANCI-2 cDNA (lnc-FANCI-2a-PA1, GenBank Acc. No. MT669801). Overlapping PCR products generated from 5' and 3' RACE PCR products with a primer pair of oHBL 17 and oHBL 18 were digested and inserted into pcDNA3 at Hind III and Xba I sites. The insertion was verified by sequencing. Plasmid pHBL05 has a long isoform of lnc-FANCI-2 (lnc-FANCI-2a-PA2, GenBank Acc. No. MT669800). PCR products generated from genomic DNA with oHBL 19 and oHBL 23 were digested and swapped into pHBL03 to replace lnc-FANCI-2a-PA1 at Bsmb I and Xba I sites.

CRISPR/Cas9 system was used to knockout lnc-FANCI-2 gene from CaSki cells. Two lnc-FANCI-2 specific guide RNAs (gRNAs) were designed by an on-line CRISPR design tool developed by the Zhang lab (http://crispr.mit.edu). A modified cloning strategy was used to clone the two gRNA into pSpCas9(BB)-2A-Puro vector (Addgene, Watertown, MA, #62988) and described in our previous publication [42]. This strategy allows all three components (two gRNAs and Cas9) express from the same plasmid to ensure the transfected cells expressing two

532   gRNAs simultaneously knocking out the targeted gene [42]. The sequences of primers to make

533   guide RNAs were listed in Supplementary Table S5.

534   Knockout of lnc-FANCI-2 gene in CaSki cells, PCR screening and genotyping

Two gRNAs each under control by a U6 promoter in a pSpCas9(BB)-2A-Puro vector were used

to create deletion in CaSki cells. Plasmid pHBL25 which contains gRNA 1 and 6 was used to

delete the entire promoter region of lnc-FANCI-2 and plasmid pHBL27 which contains gRNA 3

and 4 was used to delete only two YY1-binding motifs spanning over an 87-bp region in the lnc-

FANCI-2 promoter. CaSki cells transfected with the gRNA-expressing vectors were then under 2

weeks of puromycin selection (0.1 µg/ml). The puromycin-resistant CaSki cells were diluted to a

final concentration of 0.5 cells per 100 µl, plated 100 µl of diluted cells into each well of a 96-

well plate, and grew under continuous purymycin selection. The colonies were inspected to

ensure the clones were grown from one cell, and refed or re-plated as needed in 3 weeks of

expansion.

A direct PCR was performed for number of single cell clones as described [42]. Briefly,

several hundreds of cells expanded from a single cell clone were resuspended in 50 µl of

phosphate-buffered saline (PBS) and then frozen-thawed on dry ice for three times. Cell lysates

were treated with 4 µl of Qiagen protease (QIAGEN, Hilden, Germany, #1017782,) at 56°C for

10 min, and the protease was inactivated at 95°C for 5 mins. The resulting product was directly

used for PCR screening. P1 + P5 were used for verification of promoter deletion and P3 + P4 for

YY1 motifs deletion. To ensure the selected single cell clones displaying homozygous knockout

of lnc-FANCI-2, total cell DNA was isolated using a QIAamp DNA Blood Minikit (QIAGEN,

Hilden, Germany, #51106) and then the homozygous deletion was screened and the cells with

homozygous KO were selected and confirmed by PCR. The primers used for genotyping are

listed in supplementary Table S5. At least two single cell clones either with lnc-FANCI-2

promoter deletion or with YY1 motif deletion were selected. Three single cell clones from the

cells transfected with an empty vector, pSpCas9(BB)-2A-Puro, were also selected and served as

controls for the genotyping screening.

Cell proliferation and viability assay

Cell proliferation and viability were determined by a CCK-8 assay (Dojindo Molecular

Technologies, Rockville, MD). For proliferation assay, 500 µl of cell suspension (50,000

cells/well) were dispensed into individual wells in a 24-well plate. Cells were then treated with

inhibitors (20 µM LY294002, or 10 µM U0126 respectively) after 24 h culture. At each time

point of 24 h, 48 h and 72 h, 50 µl of CCK-8 solution was added in three wells of each group and

incubated for 1 h in a cell culture incubator at 37°C. Cell proliferation was determined by

absorbance at 450 nm, and normal medium was used to subtract background. Percentage of cell

viability was calculated by the following formula: cell viability % = OD450 of inhibitors treated

sample / OD450 of untreated sample × 100%.

Colony formation, cell migration assay and senescence analysis

CaSki Cells were seeded at 1×10^4, 5×10^3 or 1×10^3 cells/well in 6-well plates. After 24 h of

cell growth, the culture medium in each well was replaced by DMEM containing 1%

methylcellulose and 10% FBS. The plates were fixed with 3.7% formaldehyde and stained with

1% crystal violet in 2 weeks. Transwell cell migration assay was performed as previously

described [93]. Briefly, CaSki cells were re-suspended in serum free DMEM containing 0.1%

BSA. 1×10^5 cells in 100 ul were then plated on top of the filter membrane in a transwell insert

(SARSTEDT, Nümbrecht, GERMANY, # 83.3932.800) and incubate for 10 min at 37 °C to

577 allow the cells to settle down, then 600 µl of complete media containing 10% FBS were added

578 into the bottom of the lower chamber in a 24-well plate. After 24 h in culture, the cells on the top

579 of the insert membrane were removed, and cells on the bottom were fixed, stained by 0.2%

580 crystal violet and then counted.

581  Cell senescence was determined by Cellular Senescence Assay (MilliporeSigma,

582 Burlington, MA, # QIA117-1KIT). Briefly, CaSki Cells were plated at 2×10^5 cells/well in 6-

583 well plates and cultured for 2 days to 50% confluence. Cells were washed once with 2 ml PBS

584 and fixed with 0.5 ml of Fixative Solution at room temperature for 10 to 15 min. Then, cells

585 were washed twice with 2 ml PBS, and 1 ml Staining Solution was added to each well. Cells

586 were incubated at 37°C overnight without CO_2 and protected from the light. Blue stained cells

587 were then counted under light microscopy.

588 **Nuclear and cytoplasmic fractionation**

589 CaSki and SiHa cells were fractionated by using Nuclei EZ Prep Kit (Sigma-Aldrich, #NUC-

590 101) following the manufacturer's protocol. The details of the method could be found in our

591 previous publication [94]. Western blot analysis for nuclear protein SRSF3 and cytoplasmic

592 protein GAPDH was used to determine the fractionation efficiency. The fractionated cytoplasmic

593 and nuclear RNAs were used for detection of lnc-FANCI-2 RNA with human GAPDH RNA

594 serving as a control for RNA fractionation efficiency of the cytoplasmic RNAs.

595 **RT-qPCR**

596 Detection of lnc-FANCI-2 by RT-qPCR was performed as described [8]. Pre-designed primers

597 for lnc-FANCI-2 are listed in Table S5. Briefly, 2 µg total RNA was converted to cDNA using

598 Superscript First-stand Synthesis kit (Thermo Fisher Scientific, Waltham, MA, #11904018).

qPCR was performed using TaqMan gene expression Master Mix (Applied Biosystems,

Waltham, MA, #4369016) on a StepOne Plus Real-Time PCR system (Applied Biosystems).

TaqMan Gene Expression Assay (Applied Biosystems, #4331182) for GADPH

(Hs02758991_g1) was served as an internal control. Data were plotted as fold change over the

control group using the $2^{-\Delta\Delta Ct}$ method by which the data was normalized first to the values for

GAPDH and then to the median value for control samples. Data are presented as a bar graph with

mean ± SD for each group.

Northern blot

To validate lnc-FANCI-2 KO in CaSki cells, Northern blot was performed using polyA$^+$ mRNA

isolated from 100 μg of total CaSki RNA with PolyATtract mRNA Isolation Systems (Promega,

#Z5310) or 10 μg of total CaSki total RNA as described previously [8]. Briefly, RNA samples

were denatured in NorthernMax Formaldehyde loading dye (Thermo Fisher Scientific,

#AM8552) at 75°C for 15 min, then separated in an 1% formaldehyde-containing agarose gel

and transferred onto a GeneScreen Plus hybridization transfer membrane (Perkin Elmer,

Waltham, MA, #NEF987001PK). RNAs on the membrane were crosslinked by exposing to UV

light, and then prehybridized with PerfectHyb Plus hybridization buffer (Sigma-Aldrich, St.

Louis, MO, #H7003) for 2 h at 42°C. Specific oligos against lnc-FANCI-2 or GAPDH listed in

Table S5 were labeled with [γ-^{32}P]-ATP using T4 PNK (Thermo Fisher Scientific, Waltham,

MA, #18004–010) and added into hybridization buffer for overnight incubation at 42°C. The

membrane was then washed with 2× SSPE/0.5% SDS solution for 5 mins, followed by twice

washes with 0.5× SSPE/0.5% SDS each for 15 min at 42°C and then exposed to a

PhosphorImager screen.

RNA-seq and data analysis

Total RNA was from WT CaSki cells, ΔYY1-D5 and ΔPr-A9 cells were extracted using TRIzol

reagent (Thermo Fisher Scientific, Waltham, MA, #15596018). Total ribo-minus RNA-seq

libraries, four samples in each group, were prepared with TruSeq Stranded Total RNA Library

Kit and then subjected to pair-end sequencing using Illumina-HiSeq3000/4000 platform.

The obtained reads were processed using the CCBR Pipeliner utility

(https://github.com/CCBR/Pipeliner). Briefly, reads were trimmed from adapters and low-quality

bases using Cutadapt (version 1.18) (https://bioweb.pasteur.fr/packages/pack@cutadapt@1.18)

before alignment to the custom reference genome described below. The transcripts were aligned

using STAR v2.5.2b in two-pass mode [95]. Expression levels were quantified using RSEM

(RNA-Seq by Expectation-Maximization) (version 1.3.0) [96] with a custom gene annotation

described below.

The custom reference genome allowing quantification of both HPV16 and host

expression used in this alignment consisted of the human reference genome (hg38/Dec.

2013/GRCh38) with a HPV16 FASTA (https://pave.niaid.nih.gov/) sequence added as an

additional pseudochromosome. The custom gene annotation used for gene expression

quantification consisted of a concatenation of the hg38 GENCODE annotation version 30 [97]

and the HPV16 genome, with one other notable alteration. The hg38 v30 gene annotation is

incorrect at the location of the lnc-FANCI-2 locus in the hg38 genome at chr15:89378104-

89398487, which affects quantification of this gene. To correct this annotation and provide

accurate quantification, therefore, we first performed BLAST against hg38 using the sequences

of the 14 known isoform transcripts of lnc-FANCI-2 (MT669800 for lnc-FANCI-2a-PA2,

MT669801 for lnc-FANCI-2a-PA1, MT669802 for lnc-FANCI-2b-PA2, MT669803 for lnc-

FANCI-2b-PA1, MT669804 for lnc-FANCI-2c-PA2, MT669805 for lnc-FANCI-2c-PA1,

MT669806 for lnc-FANCI-2d-PA2, MT669807 for lnc-FANCI-2d-PA1, MT669808 for lnc-

FANCI-2e-PA2, MT669809 for lnc-FANCI-2e-PA1, MT669810 for lnc-FANCI-2f-PA2,

MT669811 for lnc-FANCI-2f-PA1, MT669812 for lnc-FANCI-2g-PA2, and MT669813 for lnc-

FANCI-2g-PA1). The results were used to determine the precise exon start and stop sites for

each of the isoforms. These were then used to manually adjust the default hg38 v30 gene

annotations at the locus indicated above to account for the correct structure of the lnc-FANCI-2

locus and its 14 isoforms.

Downstream analysis and visualization were performed within the NIH Integrated

Analysis Portal (NIDAP) using R programs developed on the Foundry platform (Palantir

Technologies). Briefly, raw counts data produced by RSEM were imported into the NIDAP

platform, genes were filtered for low counts (<1 cpm) and the voom algorithm [98] from the

Limma R package (version 3.40.6) [99] was used for quantile normalization and calculation of

differentially expressed genes. Pre-ranked GSEA was performed using the Molecular Signatures

Database version 6.2 [100, 101] and the fgsea package [102]. Raw data and the analyzed RNA-

seq data supporting the findings in this study have been deposited in the NCBI GEO database

(the accession#: GSE190904).

Human Soluble Receptor Array

To detect the changes of soluble receptors expressed and released from parental WT CaSki cells

and lnc-FANCI-2 KO cells, Proteome Profiler Human sReceptor (Soluble Receptor) Array

(R&D Systems, Minneapolis, MN, # ARY012) was performed according to a manufacturer

protocol. Briefly, 7×10^6 WT CaSki cells and ΔPr-A9 cells were cultured in 15 ml of DMEM

supplemented with 10% FBS in a T75 flask for 24 h. Cell culture supernatants were then

collected and centrifuged at 4,000 × g for 15 min to remove cell debris. Cells were rinsed by 10

ml of PBS and solubilized in Lysis Buffer 17 at 4°C for 30 min. The cell lysis was then

centrifuged at 14,000 × g for 5 min to remove cell debris. Protein concentration of cell lysis was

measured by Micro BCA Protein Assay (PIERCE, Waltham, MA, #23235). Next, two sets of N

and C membranes were blocking by Buffer 8/1 in 4-Well Multi-dish for 1 h on a rocking

platform. 500 µl of culture supernatant or 100 µg cell lysate generated from the WT CaSki cells

or ΔPr-A9 cells were diluted in 3 ml of Buffer 8/1 and then applied to each membrane after three

washes with 1× Wash Buffer. The membranes were incubated overnight at 4 °C on a rocking

platform and washed three times to remove unbound materials followed by incubation with their

specific cocktail of biotinylated detection antibodies for 2 h at room temperature on a rocking

platform. 2 ml of diluted Streptavidin-HRP was then applied to each membrane for 30 min at

room temperature on a rocking platform after three times of wash. 1 ml of the prepared Chemi

Reagent Mix was added onto the membranes after three times of wash. The signal was then

captured by ChemiDoc Touch Imaging System (Bio-Rad). Quantification of the relative protein

expression in ΔPr-A9 cells compared to the WT cells was determined in Image Lab software

(Bio-Rad). The average signal of duplicate spots subtracting background value from negative

control spots represented the levels of each protein. The relative change in protein levels was

then determined by comparing ΔPr-A9 cells to the WT cells.

Antibodies and Immunoblot

Immunoblot analysis was performed for individual proteins by using the following antibodies:

anti-MCAM (#17564-1-AP), anti-E2F1 (#12171-1-AP), anti-MAP4K4 (#55247-1-AP), and anti-

p53 (#10442-1-AP) antibodies were from ProteinTech (Rosemont, IL). Anti-tubulin (#T5201)

antibodies were from Sigma-Aldrich (St. Louis, MO). Anti-hnRNP C1/C2 (#Ab10294) and anti-

RAC3 (#ab129062) antibodies were from Abcam (Cambridge, United Kingdom). Anti- CCND2

(Cyclin D2, #3741), anti-Akt (#9272), anti-phospho-Akt (#9271), anti-Erk1/2 (p44/42 MAPK,

#4695), anti-phospho-Erk1/2 (#4370), and anti-IGFBP3 (D1U9C, #25864) were from Cell

Signaling Technology (Danvers, MA). Anti-VIM (#MA5-11883) was from Thermo Fisher

Scientific (Waltham, MA). Anti- PODXL2 (#AF1524), anti-ECM1 (#MAB39371), anti-TIMP2

(#AF971) and anti-ADAM8 (#AF1031) were from R&D Systems (Minneapolis, MN). Anti-

SRSF3 (#NBP2-76892) was from Novus Biologicals (Centennial, CO). Rabbit polyclonal anti-

HPV16 E7 (#GTX133411) was from GeneTex (Irvine, CA).

RAS GTPase Chemi ELISA

RAS GTPase activity in cell lysate of the WT CaSki or ΔPr-A9 cells was determined by RAS

GTPase Chemi ELISA Kit (Active Motif, Carlsbad, CA, # 52097) according to the manufacturer

protocol. Briefly, 2×10^7 cells were solubilized in 500 µl of Complete Lysis/Binding buffer at

4°C for 15 min. Protein concentration of cell extract was then measured by Micro BCA Protein

Assay (PIERCE, #23235) after cell debris was removed by centrifuged at 14,000 × g for 10 min.

2 µg of GST-Raf-RBD in 50 µl of Complete Lysis/Binding buffer was coated in each well of a

96-well plate by incubating for 1 h at 4°C with 100 rpm agitation. After three times of wash, 50

µg of extract in 50 µl Complete Lysis/Binding buffer was then added in each well. HeLa (EGF

treated) extract was served as a positive control. The plate was incubated for 1 h at room

temperature with 100 rpm agitation. The active RAS protein (GST-Raf-RBD binding) in each

well was then detected by incubating with primary RAS antibody and three times of wash, and

then, HRP-conjugated secondary antibody and three times of wash. Chemiluminescence in each

well was read in luminometer by adding 50 µl room-temperature Chemiluminescent Working

Solution.

RNA *in situ* Hybridization (RNA-ISH) and Immunofluorescence (IF) Staining

Endogenous lnc-FANCI-2 in cervival cancer tissues and their derived cell lines was examined by

RNAScope Multiplex Fluorescent V2 Assay (Advanced Cell Diagnostics, Minneapolis, MN,

#323100) as described previously [8]. A custom-designed probe targeting to nt 359-1713 of

GenBank Acc. No. MT669800.1 transcript for lnc-FANCI-2a-PA2 was utilized. Dual staining of

lnc-FANCI-2 and IGFBP RNA was performed according to RNAscope Multplex Fluorescent v2

Manual part 2 (#323100), and the channel one probe for IGFBP3 (#310351) and channel three

probe for lnc-FANCI-2 (#509061-C3) were applied.

For RNA-ISH combined IF staining of MCAM protein, 3×10^5 of CaSki cells were

grown on a glass coverslip in a 6-well plate for 24 h before transfection. Plasmid pHBL05

containing a long isoform of lnc-FANCI-2a-PA2 (GenBank Acc. No. MT669800.1) were

transfected into cells with LipoD 293 DNA in vitro transfection reagent. Cultured Adherent Cell

Sample Preparation for the RNAscope Multplex Fluorescent v2 was performed according to a

manufacturer protocol (Advanced Cell Diagnostics, Minneapolis, MN, MK-50-010). Briefly, the

cells were washed with PBS and fixed by 10% neutral buffered formalin at room temperature for

30 min. The cells were washed with PBS three times and dehydrated by 50% ethanol for 5 min,

70% ethanol for 5 min, 100% ethanol for 5 min, and 100% ethanol for 10 min. The cells were

then rehydrated by 70% ethanol for 2 min, 50% ethanol for 2 min, and PBS for 10 min. The cells

were then applied to hydrogen peroxide at RT for 10 min and washed with distilled water for

three times. The cells were then applied to Protease III (1:15 dilution with PBS) at RT for 10

min. The cells were ready for RNAscope Multplex Fluorescent v2 Manual part 2 (#323100). IF

was performed after RNA-ISH but before counterstaining with DAPI. The cells were washed

with PBS, and then were blocked with 2% bovine serum albumin (BSA) in PBS for 1 h at 37°C

or overnight at 4°C. The cells were incubated with the MCAM primary antibody (ProteinTech,

Rosemont, IL, #17564-1-AP, diluted 1:200 in 2% BSA blocking buffer) for 2 h at 37°C. An

Alexa Fluor-conjugated secondary antibody (1: 500, ThermoFisher Scientific) was diluted in a

blocking solution and incubated for 1 h at 37˚C. The slides were washed with PBS for three

times. DAPI was used for nuclei counterstaining before mounting in a Prolong Gold Antifade

mounting medium (Thermo Fisher Scientific, #P36934).

Confocal images were collected using a Zeiss LSM710 laser-scanning microscope equipped with

a 63 × Plan-Apochromat (N.A. 1.4) objective lens. Three dimensional distributions of lnc-

FANCI-2 were generated by Z stacks using ZEN 2.3 software (Zeiss).

Isolation of lnc-FANCI-2 RNA-protein complex by RNA Purification (IRPCRP) and LC-

MS/MS analysis

To identify the lnc-FANCI-2 RNA-associated proteins, we pulled down lnc-FANCI-2 RNA from

WT CaSki cells using IRPCRP, a modified ChIRP protocol from the published method [103],

followed by mass spectrometry [60]. Briefly, a total of 30 lnc-FANCI-2 anti-sense oligo probes

(oLLY496-oLLY525), each with biotinTEG at 3' end, across entire lnc-FANCI-2 RNA

(GenBank Acc. No MT669800.1) were designed using the online probe designer

(singlemoleculefish.com). Two probe pools were used in IRPCRP assays: the probe pool 1 was

the mixed oligos in even numbers (oLLY496

/498/500/502/504/506/508/510/512/514/516/518/520/522/524) and the probe pool 2 mixed with

the oligos in odd numbers

(oLLY497/499/501/503/505/507/509/511/513/515/517/519/521/523/525) (Table S5). CaSki

cells were seeded and harvested at 24 h and the cell lysates from 200 million cells were used for

each IRPCRP reaction. Instead of formaldehyde crosslink in the standard protocol [103], UV

cross-linking (254 nm, energy at 480 mJ/cm^2) was applied in our study to minimize non-specific

binding. After UV cross-linking, the cells were lysed on ice in 1× radioimmunoprecipitation

assay (RIPA) buffer (Boston Bioproducts, Ashland, MA, #BP-115) (50 mM Tris base/Tris-HCl

[pH 7.4], 150 mM NaCl, 0.5% sodium deoxycholate, 0.1% sodium dodecyl sulfate [SDS], 1%

NP-40) supplemented with protease inhibitors (complete mini-EDTA-free protease inhibitor

cocktail, Millipore Sigma, Burlington, MA, #469315900) and RNase inhibitor (Thermo Fisher

Scientific, #AM2694) for 30 min, followed by a brief sonication. The obtained cell lysates were

spin for 10 min of centrifugation at 20,000 × g at 4°C and the collected cell lysate supernatants

were pre-absorbed with C-1 magnetic beads (Thermo Fisher Scientific, #65002) for 1 h at room

temperature before proceeding to oligo probe hybridization.

Two different hybridization protocols were used to pull down lnc-FANCI-2 RNA. One

had the pre-absorbed cell lysate incubated with two separate probe pools in hybridization buffer

(750 mM NaCl, 0.1% SDS, 50 mM Tris-Cl pH 7.0, 1 mM EDTA, 15% formamide) at room

temperature overnight and the pre-washed C-1 magnetic beads was then added and incubated for

additional 4 h; the other had two separate oligo probe pools immobilized onto pre-washed C-1

magnetic beads first at room temperature for 1 h and then mixed with the pre-absorbed cell

lysates in hybridization buffer overnight at room temperature, along with the negative control-

beads only IRPCRP reaction without oligo probes serving as a control in the modified protocol.

After hybridization, the beads were washed with 1× wash buffer (2× NaCl and sodium citrate,

0.1% SDS) for 5 times and resuspend in 1 ml lysis buffer. 100 µl beads from each IRPCRP were

set aside for RNA isolation of lnc-FANCI-2 pulldown efficiency and the leftover beads were sent

for mass spectrometry analysis. The input and IRPCRP RNA were isolated using a MIRNeasy

mini kit (Qiagen, #217004) following the standard protocol after proteinase K (Millipore Sigma,

#71049) digestion.

For mass spectrometry analysis, the IRPCRP beads were processed for trypsin/LysC

digestion before submitted to Thermo Scientific Orbitrap Exploris 240 Mass Spectrometer and a

Thermo Dionex UltiMate 3000 RSLCnano System for Proteinomics. Peptides from trypsin

digestion were loaded onto a peptide trap cartridge at a flow rate of 5 µl/min. The trapped

peptides were eluted onto a reversed-phase Easy-Spray Column PepMap RSLC, C18, 2 µM,

100A, 75 µm × 250 mm (Thermo Scientific) using a linear gradient of acetonitrile (3-36%) in

0.1% formic acid. The elution duration was 110 min at a flow rate of 0.3 µl/min. Eluted peptides

from the Easy-Spray column were ionized and sprayed into the mass spectrometer, using a Nano

Easy-Spray Ion Source (Thermo Fisher Scientific) under the following settings: spray voltage,

1.6 kV, Capillary temperature, 275 °C. Other settings were empirically determined. Raw data

files were searched against human protein sequences database using the Proteome Discoverer 2.4

software (Thermo Fisher Scientific, San Jose, CA) based on the SEQUEST algorithm. All the

protein peptides identified in the IRPCRP pulldowns were summarized in Table S4.

ACKNOWLEDGEMENTS

We thank Louise T. Chow of the University of Alabama at Birmingham for her critical reading

of the manuscript. We thank Craig Meyers of Penn State University Hershey Medical Center for

providing HPV16- and HPV18- infected raft cultures, Xing Xie and Yang Li of Zhejiang

University Women's Hospital of China for cervical tissue sections, and Johannes G. Schweizer

from Arbor Vita corporation for anti-HPV16 E6-specific antibodies. This study was supported

by the Intramural Research Program of the NIH, National Cancer Institute, Center for Cancer

Research.

DATA AVAILABILITY

NCBI GEO database Acc. No. GSE190904 for CaSki cell RNA-seq.

AUTHORS CONTRIBUTIONS: H. L., L. Y., and Z.M.Z. designed the study. H. L. and L. Y.

performed the experiments. H. L., L. Y., V.M., T. M., M. Y., P. J., M.C., D. R. L., and Z.M.Z.

analyzed data and participated in the discussion and interpretation of the results. H. L., L. Y., and

Z.M.Z drafted the manuscript. H. L., L. Y., and Z.M.Z. revised the manuscript. All authors read

and approved of the final manuscript.

DECLARATION OF INTERESTS: The authors declare no competing interests.

REFERENCES

1. Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, Jemal A: **Global cancer**
statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in
185 countries. *CA Cancer J Clin* 2024, **74:**229-263.
2. Bosch FX, Lorincz A, Munoz N, Meijer CJ, Shah KV: **The causal relation between human**
papillomavirus and cervical cancer. *J Clin Pathol* 2002, **55:**244-265.
3. Munoz N, Bosch FX, de Sanjose S, Herrero R, Castellsague X, Shah KV, Snijders PJ, Meijer CJ,
International Agency for Research on Cancer Multicenter Cervical Cancer Study G:
Epidemiologic classification of human papillomavirus types associated with cervical cancer. *N*
Engl J Med 2003, **348:**518-527.
4. Walboomers JM, Jacobs MV, Manos MM, Bosch FX, Kummer JA, Shah KV, Snijders PJ, Peto J,
Meijer CJ, Munoz N: **Human papillomavirus is a necessary cause of invasive cervical cancer**
worldwide. *J Pathol* 1999, **189:**12-19.
5. McCredie MR, Sharples KJ, Paul C, Baranyai J, Medley G, Jones RW, Skegg DC: **Natural history of**
cervical neoplasia and risk of invasive cancer in women with cervical intraepithelial neoplasia
3: a retrospective cohort study. *Lancet Oncol* 2008, **9:**425-434.
6. Niraj J, Farkkila A, D'Andrea AD: **The Fanconi Anemia Pathway in Cancer.** *Annu Rev Cancer Biol*
2019, **3:**457-478.
7. Hoskins EE, Morris TA, Higginbotham JM, Spardy N, Cha E, Kelly P, Williams DA, Wikenheiser-
Brokamp KA, Duensing S, Wells SI: **Fanconi anemia deficiency stimulates HPV-associated**
hyperplastic growth in organotypic epithelial raft culture. *Oncogene* 2009, **28:**674-685.
8. Liu H, Xu J, Yang Y, Wang X, Wu E, Majerciak V, Zhang T, Steenbergen RDM, Wang HK, Banerjee
NS, et al: **Oncogenic HPV promotes the expression of the long noncoding RNA lnc-FANCI-2**
through E7 and YY1. *Proc Natl Acad Sci U S A* 2021, **118:**e2014195118.

9.   Santegoets LA, van Baars R, Terlou A, Heijmans-Antonissen C, Swagemakers SM, van der Spek PJ,
Ewing PC, van Beurden M, van der Meijden WI, Helmerhorst TJ, Blok LJ: **Different DNA damage**
and cell cycle checkpoint control in low- and high-risk human papillomavirus infections of the
vulva. *Int J Cancer* 2012, **130:**2874-2885.
10.   Hoskins EE, Morreale RJ, Werner SP, Higginbotham JM, Laimins LA, Lambert PF, Brown DR,
Gillison ML, Nuovo GJ, Witte DP, et al: **The fanconi anemia pathway limits human**
papillomavirus replication. *J Virol* 2012, **86:**8131-8138.
11.   Park JW, Pitot HC, Strati K, Spardy N, Duensing S, Grompe M, Lambert PF: **Deficiencies in the**
Fanconi anemia DNA damage response pathway increase sensitivity to HPV-associated head
and neck cancer. *Cancer Res* 2010, **70:**9959-9968.
12.   Park JW, Shin MK, Lambert PF: **High incidence of female reproductive tract cancers in FA-**
deficient HPV16-transgenic mice correlates with E7's induction of DNA damage response, an
activity mediated by E7's inactivation of pocket proteins. *Oncogene* 2014, **33:**3383-3391.
13.   Hietanen S, Lain S, Krausz E, Blattner C, Lane DP: **Activation of p53 in cervical carcinoma cells by**
small molecules. *Proc Natl Acad Sci U S A* 2000, **97:**8501-8506.
14.   Tommasino M, Accardi R, Caldeira S, Dong W, Malanchi I, Smet A, Zehbe I: **The role of TP53 in**
Cervical carcinogenesis. *Hum Mutat* 2003, **21:**307-312.
15.   Prior IA, Lewis PD, Mattos C: **A comprehensive survey of Ras mutations in cancer.** *Cancer Res*
2012, **72:**2457-2467.
16.   Fernández-Medarde A, Santos E: **Ras in cancer and developmental diseases.** *Genes Cancer*
2011, **2:**344-358.
17.   Roman A, Munger K: **The papillomavirus E7 proteins.** *Virology* 2013, **445:**138-168.
18.   Vande Pol SB, Klingelhutz AJ: **Papillomavirus E6 oncoproteins.** *Virology* 2013, **445:**115-137.
19.   Liu X, Dakic A, Zhang Y, Dai Y, Chen R, Schlegel R: **HPV E6 protein interacts physically and**
functionally with the cellular telomerase complex. *Proc Natl Acad Sci U S A* 2009, **106:**18780-
18785.
20.   Klingelhutz AJ, Foster SA, McDougall JK: **Telomerase activation by the E6 gene product of**
human papillomavirus type 16. *Nature* 1996, **380:**79-82.
21.   Storey A, Pim D, Murray A, Osborn K, Banks L, Crawford L: **Comparison of the in vitro**
transforming activities of human papillomavirus types. *EMBO J* 1988, **7:**1815-1820.
22.   Barbosa MS, Schlegel R: **The E6 and E7 genes of HPV-18 are sufficient for inducing two-stage in**
vitro transformation of human keratinocytes. *Oncogene* 1989, **4:**1529-1532.
23.   Hawley-Nelson P, Vousden KH, Hubbert NL, Lowy DR, Schiller JT: **HPV16 E6 and E7 proteins**
cooperate to immortalize human foreskin keratinocytes. *EMBO J* 1989, **8:**3905-3910.
24.   Munger K, Phelps WC, Bubb V, Howley PM, Schlegel R: **The E6 and E7 genes of the human**
papillomavirus type 16 together are necessary and sufficient for transformation of primary
human keratinocytes. *J Virol* 1989, **63:**4417-4421.
25.   Halbert CL, Demers GW, Galloway DA: **The E7 gene of human papillomavirus type 16 is**
sufficient for immortalization of human epithelial cells. *J Virol* 1991, **65:**473-478.
26.   Phelps WC, Yee CL, Munger K, Howley PM: **The human papillomavirus type 16 E7 gene encodes**
transactivation and transformation functions similar to those of adenovirus E1A. *Cell* 1988,
53:539-547.
27.   DiPaolo JA, Woodworth CD, Popescu NC, Notario V, Doniger J: **Induction of human cervical**
squamous cell carcinoma by sequential transfection with human papillomavirus 16 DNA and
viral Harvey ras. *Oncogene* 1989, **4:**395-399.
28.   Schreiber K, Cannon RE, Karrison T, Beck-Engeser G, Huo D, Tennant RW, Jensen H, Kast WM,
Krausz T, Meredith SC, et al: **Strong synergy between mutant ras and HPV16 E6/E7 in the**
development of primary tumors. *Oncogene* 2004, **23:**3972-3979.

29.   Narisawa-Saito M, Yoshimatsu Y, Ohno S, Yugawa T, Egawa N, Fujita M, Hirohashi S, Kiyono T: **An**
888   **in vitro multistep carcinogenesis model for human cervical cancer.** *Cancer Res* 2008, **68:**5699-
889   5705.
890   30.   Conover CA, Bale LK, Durham SK, Powell DR: **Insulin-like growth factor (IGF) binding protein-3**
891   **potentiation of IGF action is mediated through the phosphatidylinositol-3-kinase pathway and**
892   **is associated with alteration in protein kinase B/AKT sensitivity.** *Endocrinology* 2000, **141:**3098-
893   3103.
894   31.   Allard JB, Duan C: **IGF-Binding Proteins: Why Do They Exist and Why Are There So Many?** *Front*
895   *Endocrinol (Lausanne)* 2018, **9:**117.
896   32.   Bach LA: **IGF-binding proteins.** *J Mol Endocrinol* 2018, **61:**T11-t28.
897   33.   Song S, Pitot HC, Lambert PF: **The human papillomavirus type 16 E6 gene alone is sufficient to**
898   **induce carcinomas in transgenic animals.** *J Virol* 1999, **73:**5887-5893.
899   34.   Herber R, Liem A, Pitot H, Lambert PF: **Squamous epithelial hyperplasia and carcinoma in mice**
900   **transgenic for the human papillomavirus type 16 E7 oncogene.** *J Virol* 1996, **70:**1873-1881.
901   35.   Song S, Liem A, Miller JA, Lambert PF: **Human papillomavirus types 16 E6 and E7 contribute**
902   **differently to carcinogenesis.** *Virology* 2000, **267:**141-150.
903   36.   Riley RR, Duensing S, Brake T, Munger K, Lambert PF, Arbeit JM: **Dissection of human**
904   **papillomavirus E6 and E7 function in transgenic mouse models of cervical carcinogenesis.**
905   *Cancer Res* 2003, **63:**4862-4871.
906   37.   Brake T, Lambert PF: **Estrogen contributes to the onset, persistence, and malignant**
907   **progression of cervical cancer in a human papillomavirus-transgenic mouse model.** *Proc Natl*
908   *Acad Sci U S A* 2005, **102:**2490-2495.
909   38.   Kato S, Endoh H, Masuhiro Y, Kitamoto T, Uchiyama S, Sasaki H, Masushige S, Gotoh Y, Nishida E,
910   Kawashima H, et al: **Activation of the estrogen receptor through phosphorylation by mitogen-**
911   **activated protein kinase.** *Science* 1995, **270:**1491-1494.
912   39.   Statello L, Guo CJ, Chen LL, Huarte M: **Gene regulation by long non-coding RNAs and its**
913   **biological functions.** *Nat Rev Mol Cell Biol* 2021, **22:**96-118.
914   40.   Liu H, Zheng ZM: **Linking a nuclear lncRNA to cytoplasmic lysosome integrity and cell death.**
915   *Proc Natl Acad Sci U S A* 2022, **119:**e2123082119.
916   41.   Scheffner M, Münger K, Byrne JC, Howley PM: **The state of the p53 and retinoblastoma genes**
917   **in human cervical carcinoma cell lines.** *Proceedings of the National Academy of Sciences* 1991,
918   **88:**5523-5527.
919   42.   BeltCappellino A, Majerciak V, Lobanov A, Lack J, Cam M, Zheng ZM: **CRISPR/Cas9-Mediated**
920   **Knockout and In Situ Inversion of the ORF57 Gene from All Copies of the Kaposi's Sarcoma-**
921   **Associated Herpesvirus Genome in BCBL-1 Cells.** *J Virol* 2019, **93:**e00628-00619.
922   43.   Schlomann U, Wildeboer D, Webster A, Antropova O, Zeuschner D, Knight CG, Docherty AJ,
923   Lambert M, Skelton L, Jockusch H, Bartsch JW: **The metalloprotease disintegrin ADAM8.**
924   **Processing by autocatalysis is required for proteolytic activity and cell adhesion.** *J Biol Chem*
925   2002, **277:**48210-48219.
926   44.   Conrad C, Yildiz D, Cleary SJ, Margraf A, Cook L, Schlomann U, Panaretou B, Bowser JL,
927   Karmouty-Quintana H, Li J, et al: **ADAM8 signaling drives neutrophil migration and ARDS**
928   **severity.** *JCI Insight* 2022, **7:**e149870.
929   45.   Liberzon A, Birger C, Thorvaldsdottir H, Ghandi M, Mesirov JP, Tamayo P: **The Molecular**
930   **Signatures Database (MSigDB) hallmark gene set collection.** *Cell Syst* 2015, **1:**417-425.
931   46.   Downward J: **Targeting RAS signalling pathways in cancer therapy.** *Nat Rev Cancer* 2003, **3:**11-
932   22.
933   47.   Simanshu DK, Nissley DV, McCormick F: **RAS Proteins and Their Regulators in Human Disease.**
934   *Cell* 2017, **170:**17-33.

48.   Castellano E, Downward J: **RAS Interaction with PI3K: More Than Just Another Effector**
Pathway. *Genes Cancer* 2011, **2:**261-274.
49.   Asati V, Mahapatra DK, Bharti SK: **PI3K/Akt/mTOR and Ras/Raf/MEK/ERK signaling pathways**
inhibitors as anticancer agents: Structural and pharmacological perspectives. *Eur J Med Chem*
2016, **109:**314-341.
50.   Cuesta C, Arevalo-Alameda C, Castellano E: **The Importance of Being PI3K in the RAS Signaling**
Network. *Genes (Basel)* 2021, **12:**1094.
51.   Vanhaesebroeck B, Perry MWD, Brown JR, Andre F, Okkenhaug K: **PI3K inhibitors are finally**
coming of age. *Nat Rev Drug Discov* 2021, **20:**741-769.
52.   Favata MF, Horiuchi KY, Manos EJ, Daulerio AJ, Stradley DA, Feeser WS, Van Dyk DE, Pitts WJ,
Earl RA, Hobbs F, et al: **Identification of a novel inhibitor of mitogen-activated protein kinase**
kinase. *J Biol Chem* 1998, **273:**18623-18632.
53.   Serrano ML, Sánchez-Gómez M, Bravo MM: **Insulin-like growth factor system gene expression**
in cervical scrapes from women with squamous intraepithelial lesions and cervical cancer.
Growth Horm IGF Res 2007, **17:**492-499.
54.   Wang Z, Xu Q, Zhang N, Du X, Xu G, Yan X: **CD146, from a melanoma cell adhesion molecule to**
a signaling receptor. *Signal Transduct Target Ther* 2020, **5:**148.
55.   Kebir A, Harhouri K, Guillet B, Liu JW, Foucault-Bertaud A, Lamy E, Kaspi E, Elganfoud N, Vely F,
Sabatier F, et al: **CD146 short isoform increases the proangiogenic potential of endothelial**
progenitor cells in vitro and in vivo. *Circ Res* 2010, **107:**66-75.
56.   Stalin J, Harhouri K, Hubert L, Garrigue P, Nollet M, Essaadi A, Muller A, Foucault-Bertaud A,
Bachelier R, Sabatier F, et al: **Soluble CD146 boosts therapeutic effect of endothelial**
progenitors through proteolytic processing of short CD146 isoform. *Cardiovasc Res* 2016,
111:240-251.
57.   Joshkon A, Heim X, Dubrou C, Bachelier R, Traboulsi W, Stalin J, Fayyad-Kazan H, Badran B,
Foucault-Bertaud A, Leroyer AS, et al: **Role of CD146 (MCAM) in Physiological and Pathological**
Angiogenesis-Contribution of New Antibodies for Therapy. *Biomedicines* 2020, **8:**633.
58.   Baxter RC: **Insulin-like growth factor binding protein-3 (IGFBP-3): Novel ligands mediate**
unexpected functions. *J Cell Commun Signal* 2013, **7:**179-189.
59.   Varma Shrivastav S, Bhardwaj A, Pathak KA, Shrivastav A: **Insulin-Like Growth Factor Binding**
Protein-3 (IGFBP-3): Unraveling the Role in Mediating IGF-Independent Effects Within the Cell.
Front Cell Dev Biol 2020, **8:**286.
60.   Chu C, Chang HY: **ChIRP-MS: RNA-Directed Proteomic Discovery.** *Methods Mol Biol* 2018,
1861:37-45.
61.   Gao X, Gao C, Liu G, Hu J: **MAP4K4: an emerging therapeutic target in cancer.** *Cell Biosci* 2016,
6:56.
62.   Gao X, Chen G, Gao C, Zhang DH, Kuan SF, Stabile LP, Liu G, Hu J: **MAP4K4 is a novel MAPK/ERK**
pathway regulator required for lung adenocarcinoma maintenance. *Mol Oncol* 2017, **11:**628-
639.
63.   Gonzalez-Montero J, Rojas CI, Burotto M: **MAP4K4 and cancer: ready for the main stage?** *Front*
Oncol 2023, **13:**1162835.
64.   Patterson V, Ullah F, Bryant L, Griffin JN, Sidhu A, Saliganan S, Blaile M, Saenz MS, Smith R,
Ellingwood S, et al: **Abrogation of MAP4K4 protein function causes congenital anomalies in**
humans and zebrafish. *Sci Adv* 2023, **9:**eade0631.
65.   Roth Flach RJ, Guo CA, Danai LV, Yawe JC, Gujja S, Edwards YJ, Czech MP: **Endothelial Mitogen-**
Activated Protein Kinase Kinase Kinase Kinase 4 Is Critical for Lymphatic Vascular
Development and Function. *Mol Cell Biol* 2016, **36:**1740-1749.

66. Castello A, Hentze MW, Preiss T: **Metabolic Enzymes Enjoying New Partnerships as RNA-**
Binding Proteins. *Trends Endocrinol Metab* 2015, **26:**746-757.
67. Hentze MW, Castello A, Schwarzl T, Preiss T: **A brave new world of RNA-binding proteins.** *Nat*
Rev Mol Cell Biol 2018, **19:**327-341.
68. Crook T, Storey A, Almond N, Osborn K, Crawford L: **Human papillomavirus type 16 cooperates**
with activated ras and fos oncogenes in the hormone-dependent transformation of primary
mouse cells. *Proc Natl Acad Sci U S A* 1988, **85:**8820-8824.
69. Matlashewski G, Schneider J, Banks L, Jones N, Murray A, Crawford L: **Human papillomavirus**
type 16 DNA cooperates with activated ras in transforming primary cells. *EMBO J* 1987,
6:1741-1746.
70. Greenhalgh DA, Wang XJ, Rothnagel JA, Eckhardt JN, Quintanilla MI, Barber JL, Bundman DS,
Longley MA, Schlegel R, Roop DR: **Transgenic mice expressing targeted HPV-18 E6 and E7**
oncogenes in the epidermis develop verrucous lesions and spontaneous, rasHa-activated
papillomas. *Cell Growth Differ* 1994, **5:**667-675.
71. Riou G, Barrois M, Sheng ZM, Duvillard P, Lhomme C: **Somatic deletions and mutations of c-Ha-**
ras gene in human cervical cancers. *Oncogene* 1988, **3:**329-333.
72. Eiben GL, Velders MP, Schreiber H, Cassetti MC, Pullen JK, Smith LR, Kast WM: **Establishment of**
an HLA-A*0201 human papillomavirus type 16 tumor model to determine the efficacy of
vaccination strategies in HLA-A*0201 transgenic mice. *Cancer Res* 2002, **62:**5792-5799.
73. Golijow CD, Mouron SA, Gomez MA, Dulout FN: **Differences in K-ras codon 12 mutation**
frequency between "high-risk" and "low-risk" HPV-infected samples. *Gynecol Oncol* 1999,
75:108-112.
74. Prokopakis P, Sourvinos G, Koumantaki Y, Koumantakis E, Spandidos DA: **K-ras mutations and**
HPV infection in cervicitis and intraepithelial neoplasias of the cervix. *Oncol Rep* 2002, **9:**129-
133.
75. Strickland SW, Vande Pol S: **The Human Papillomavirus 16 E7 Oncoprotein Attenuates AKT**
Signaling To Promote Internal Ribosome Entry Site-Dependent Translation and Expression of
c-MYC. *J Virol* 2016, **90:**5611-5621.
76. Nieto P, Ambrogio C, Esteban-Burgos L, Gómez-López G, Blasco MT, Yao Z, Marais R, Rosen N,
Chiarle R, Pisano DG, et al: **A Braf kinase-inactive mutant induces lung adenocarcinoma.** *Nature*
2017, **548:**239-243.
77. Yang J, Nie J, Ma X, Wei Y, Peng Y, Wei X: **Targeting PI3K in cancer: mechanisms and advances**
in clinical trials. *Mol Cancer* 2019, **18:**26.
78. Saliani M, Mirzaiebadizi A, Javadmanesh A, Siavoshi A, Ahmadian MR: **KRAS-related long**
noncoding RNAs in human cancers. *Cancer Gene Ther* 2022, **29:**418-427.
79. Martin JL, Weenink SM, Baxter RC: **Insulin-like growth factor-binding protein-3 potentiates**
epidermal growth factor action in MCF-10A mammary epithelial cells. Involvement of p44/42
and p38 mitogen-activated protein kinases. *J Biol Chem* 2003, **278:**2969-2976.
80. Józefiak A, Larska M, Pomorska-Mól M, Ruszkowski JJ: **The IGF-1 Signaling Pathway in Viral**
Infections. *Viruses* 2021, **13:**1488.
81. Jozefiak A, Pacholska-Bogalska J, Myga-Nowak M, Kedzia W, Kwasniewska A, Luczak M, Kedzia H,
Gozdzicka-Jozefiak A: **Serum and tissue levels of insulin-like growth factor-I in women with**
dysplasia and HPV-positive cervical cancer. *Mol Med Rep* 2008, **1:**231-237.
82. Serrano ML, Romero A, Cendales R, Sánchez-Gómez M, Bravo MM: **Serum levels of insulin-like**
growth factor-I and -II and insulin-like growth factor binding protein 3 in women with
squamous intraepithelial lesions and cervical cancer. *Biomedica* 2006, **26:**258-268.

83.   Schäfer B, Marg B, Gschwind A, Ullrich A: **Distinct ADAM metalloproteinases regulate G**
1029   **protein-coupled receptor-induced cell proliferation and survival.** *J Biol Chem* 2004, **279:**47929-
1030   47938.
1031   84.   Ohtsu H, Dempsey PJ, Eguchi S: **ADAMs as mediators of EGF receptor transactivation by G**
1032   **protein-coupled receptors.** *Am J Physiol Cell Physiol* 2006, **291:**C1-10.
1033   85.   Dang M, Dubbin K, D'Aiello A, Hartmann M, Lodish H, Herrlich A: **Epidermal growth factor (EGF)**
1034   **ligand release by substrate-specific a disintegrin and metalloproteases (ADAMs) involves**
1035   **different protein kinase C (PKC) isoenzymes depending on the stimulus.** *J Biol Chem* 2011,
1036   **286:**17704-17713.
1037   86.   Kleino I, Järviluoma A, Hepojoki J, Huovila AP, Saksela K: **Preferred SH3 domain partners of**
1038   **ADAM metalloproteases include shared and ADAM-specific SH3 interactions.** *PLoS One* 2015,
1039   **10:**e0121301.
1040   87.   Huang YK, Cheng WC, Kuo TT, Yang JC, Wu YC, Wu HH, Lo CC, Hsieh CY, Wong SC, Lu CH, et al:
1041   **Inhibition of ADAM9 promotes the selective degradation of KRAS and sensitizes pancreatic**
1042   **cancers to chemotherapy.** *Nat Cancer* 2024, **5:**400-419.
1043   88.   Sun C, Wu MH, Guo M, Day ML, Lee ES, Yuan SY: **ADAM15 regulates endothelial permeability**
1044   **and neutrophil migration via Src/ERK1/2 signalling.** *Cardiovasc Res* 2010, **87:**348-355.
1045   89.   Anfosso F, Bardin N, Frances V, Vivier E, Camoin-Jau L, Sampol J, Dignat-George F: **Activation of**
1046   **human endothelial cells via S-endo-1 antigen (CD146) stimulates the tyrosine phosphorylation**
1047   **of focal adhesion kinase p125(FAK).** *J Biol Chem* 1998, **273:**26852-26856.
1048   90.   Dye DE, Karlen S, Rohrbach B, Staub O, Braathen LR, Eidne KA, Coombe DR: **hShroom1 links a**
1049   **membrane bound protein to the actin cytoskeleton.** *Cell Mol Life Sci* 2009, **66:**681-696.
1050   91.   Bardin N, Frances V, Combes V, Sampol J, Dignat-George F: **CD146: biosynthesis and production**
1051   **of a soluble form in human cultured endothelial cells.** *FEBS Lett* 1998, **421:**12-14.
1052   92.   Xu J, Liu H, Yang Y, Wang X, Liu P, Li Y, Meyers C, Banerjee NS, Wang HK, Cam M, et al: **Genome-**
1053   **Wide Profiling of Cervical RNA-Binding Proteins Identifies Human Papillomavirus Regulation of**
1054   **RNASEH2A Expression by Viral E7 and E2F1.** *mBio* 2019, **10:**e02687-18.
1055   93.   Justus CR, Leffler N, Ruiz-Echevarria M, Yang LV: **In vitro cell migration and invasion assays.** *J Vis*
1056   *Exp* 2014, **88:**51046.
1057   94.   Yu L, Zheng ZM: **Human Papillomavirus Type 16 Circular RNA Is Barely Detectable for the**
1058   **Claimed Biological Activity.** *mBio* 2022, **13:**e0359421.
1059   95.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR:
1060   **STAR: ultrafast universal RNA-seq aligner.** *Bioinformatics* 2013, **29:**15-21.
1061   96.   Li B, Dewey CN: **RSEM: accurate transcript quantification from RNA-Seq data with or without a**
1062   **reference genome.** *BMC Bioinformatics* 2011, **12:**323.
1063   97.   Harrow J, Frankish A, Gonzalez JM, Tapanari E, Diekhans M, Kokocinski F, Aken BL, Barrell D,
1064   Zadissa A, Searle S, et al: **GENCODE: the reference human genome annotation for The ENCODE**
1065   **Project.** *Genome Res* 2012, **22:**1760-1774.
1066   98.   Law CW, Chen Y, Shi W, Smyth GK: **voom: Precision weights unlock linear model analysis tools**
1067   **for RNA-seq read counts.** *Genome Biol* 2014, **15:**R29.
1068   99.   Smyth GK: **Linear models and empirical bayes methods for assessing differential expression in**
1069   **microarray experiments.** *Stat Appl Genet Mol Biol* 2004, **3:**Article3.
1070   100.   Liberzon A, Subramanian A, Pinchback R, Thorvaldsdottir H, Tamayo P, Mesirov JP: **Molecular**
1071   **signatures database (MSigDB) 3.0.** *Bioinformatics* 2011, **27:**1739-1740.
1072   101.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A,
1073   Pomeroy SL, Golub TR, Lander ES, Mesirov JP: **Gene set enrichment analysis: a knowledge-**
1074   **based approach for interpreting genome-wide expression profiles.** *Proc Natl Acad Sci U S A*
1075   2005, **102:**15545-15550.

1076   102.   Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A: **Fast gene set**
1077   **enrichment analysis.** *bioRxiv* 2021.
1078   103.   Chu C, Quinn J, Chang HY: **Chromatin isolation by RNA purification (ChIRP).** *J Vis Exp* 2012:3912.

1079

 Figure legends

Fig. 1. Increased expression of lnc-FANCI-2 in HPV16-infected cervical squamous cell

carcinoma tissues and HPV16- and HPV18-infected raft culture tissues. (A and B) Expression

lnc-FANCI-2 (red) in HPV16$^+$ cervical squamous cell carcinoma tissues were examined by

RNAscope RNA-ISH analysis. Nuclei were stained with DAPI (blue). The corresponding

regions (A) in H&E staining are shown for the adjacent region from the tumor nest and

histotype. (C) Human foreskin keratinocyte (HFK)-derived raft cultures without (HFK) or with

HPV16 or HPV18 infections (HFK-HPV16 or HFK-HPV18) were examined at day 10 for lnc-

FANCI-2 RNA by RNAscope RNA-ISH analysis. Scale bars: 25 μm in the original figures and

10 μm in the zoomed insets.

Fig. 2. Knockout (KO) of lnc-FANCI-2 in CaSki cells affects cell proliferation, colony formation

and migration. (A) Diagram and KO strategies of lnc-FANCI-2 gene. On the figure top is the

lnc-FANCI-2 gene structure and its alternative transcription start sites (TSS) and polyadenylation

sites (pA). TSS2 or pA2 with a heavier arrow are predominately used for lnc-FANCI-2

expression. The lower figure part shows KO strategies. Red slashes represent gRNA-targeted

sites to create genomic DNA deletions by CRISPR-Cas9 technology. Deletion of two YY1-

binding motifs (ΔYY1) led to a deletion of 86-bp DNA fragment. Deletion of lnc-FANCI-2

promoter (ΔPr) led to delete a ~3.3-kb promoter region. Primers (P1-P4) used for PCR screening

were shown as black arrows. (B) PCR screening of single cell clones with homozygous lnc-

FANCI-2 KO, with indicated primer sets diagramed (A). Single cell clones selected from the

cells transfected with an empty vector served as control (Ctrl) cells. (C) Evaluation of lnc-

FANCI-2 KO efficiency in the selected single cell clones (B) by RT-qPCR. (D) Northern blot

validation of lnc-FANCI-2 KO efficiency from the individual single cell clones or the parental

(WT) CaSki cells on polyA$^+$ RNA (enriched from 100 µg of total cell RNA, lanes 1-4) or 10 µg

of total cell RNA (lanes 5-9). Total cell RNA (2 µg) of HEK293T cells with ectopic expression

of two isoforms (short, S or long, L) of lnc-FANCI-2 cDNA served as a control. Antisense oligo

probe P5 (A) labeled with ^{32}P was used for hybridization. GAPDH RNA served as a loading

control and hybridized with a GAPDH-specific oligo probe. (E-F) RNA in situ hybridization

(RNA-ISH) validation of lnc-FANCI-2 KO efficiency by RNAscope technology. Two single cell

clones were examined with red color for lnc-FANCI-2 and blue color for the nucleus (E). Bar

graphs show the copy number of lnc-FANCI-2 per cell in the WT, ΔYY1-D5 or ΔPr-A9 CaSki

cells (each averaged from 200 cells) (F). (G-I) Cell proliferation, colony formation and migration

of WT, ΔYY1-D5 or ΔPr-A9 cells show requirements of lnc-FANCI-2 expression for cell

proliferation, colony formation, and cell invasion of CaSki cells. Representative pictures are

shown from three separate experiments of cell proliferation by CCK8 assay (G), cell colony

formation under 1% methylcellulose solution (H), and cell migration by transwell assay (I) of

WT, ΔYY1-D5 or ΔPr-A9 cells. Cell colonies were stained with crystal violet as described in the

procedure. Numbers of the invaded cells crossed the transwell membrane were quantified and

shown as bar graphs (I). *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$ by two tailed Student t test.

Fig. 3. KO of lnc-FANCI-2 in CaSki cells affects expression of cellular soluble receptors. (A)

Dot blots show differentially expressed soluble receptors in ΔPr-A9 cells when compared to

those in the WT CaSki cells. A Proteome Profiler Human sReceptor Array was used to examine

105 cellular soluble receptors. Label in red, upregulated receptors; label in blue, downregulated

receptors. (B) Quantification of differentially expressed and/or released soluble receptors (A) in

ΔPr-A9 cells when compared to those in the WT CaSki cells. Error bar represents standard

deviation of replicates in each proteome array. The dash line indicates the threshold of fold

change (FC) with a score above or below the threshold (-/+ 0.3 FC) being determined as

differentially expressed and/or released receptors. (C) Validation of several differentially

expressed soluble receptors by immunoblot analysis using specific antibodies. Tubulin served as

an internal loading control. * non-specific band in MCAM immunoblot.

Fig. 4. Transcriptomic effect of lnc-FANCI-2 KO in CaSki cells by RNA-seq analysis. (A)

RNA-seq reads-coverage maps by IGV showing the expression levels of lnc-FANCI-2 from two

KO cell clones, ΔPr-A9 and ΔYY1-D5, to the WT CaSki cells. one representative coverage

profile of four in each type of cells is shown by IGV. FPKM, fragments per kilobase of transcript

per million. Red slashes represent the deleted genomic region, with validated fourteen lnc-

FANCI-2 RNA isoforms shown below [8]. (B) Similarity in heatmap comparison among ΔPr-

A9, ΔYY1-D5 and WT cells was generated using Limma-normalized counts from each sample

by Pearson complete linkage method. (C) Volcano plot visualization of differentially expressed

genes (DEGs) in ΔPr-A9 and ΔYY1-D5. The genes with the most significant change in the

increased (red) or decreased (blue) expression are indicated. FC, fold change. (D) Venn diagram

depicting the overlapped DEGs between ΔPr-A9 and ΔYY1-D5 cells over the WT CaSki cells.

(E) Heatmap shows overlapped DEGs with FPKM ≥7 in ΔPr-A9 and ΔYY1-D5 cells when

compared with the WT CaSki cells. (F) Validation of selective 12 upregulated or downregulated

DEGs shown in the heatmap by RT-PCR with gene-specific primers in the presence (+) or

absence (-) of reverse transcriptase (RT). GAPDH served as an internal RNA control. Relative

expression level of each gene was calculated based on the band density after normalizing to

GAPDH RNA band, with the expression level in the WT CaSki cells setting as 1. M, a 100-bp

DNA marker. (G) Validation of the upregulated expression of IGFBP3 in ΔPr-A9 cells over the

WT CaSki cells by immunoblot analysis.

Fig. 5. Pathway analyses of DEGs identified by RNA-seq. (A) Top 3 upregulated and top 4

downregulated pathways in ΔPr-A9 and ΔYY1-D5 cells over the WT CaSki cells. Data were

generated by Gene set enrichment analysis (GSEA) performed with Hallmark gene sets. NES

stands for normalized enrichment score. (B-E) Each GSEA Enrichment plot shows the

enrichment score and gene hits enriched in ΔPr-A9 cells using Hallmark gene sets,

KRAS_SIGNALING_UP (B), EPITHELIAL_MESENCHYMAL_TRANSITION (C),

INTERFERON_GAMMA_ RESPONSE (D) or INTERFERON_ALPHA_RESPONSE (E). padj,

adjusted p value; Zero crossed 7342, the middle number of total genes in the GSEA at ranking.

The heatmaps below the enrichment plots (B and C) visualize the genes enriched in respective

pathways of KRAS_SIGNALING_UP (B) and

EPITHELIAL_MESENCHYMAL_TRANSITION (C) in ΔPr-A9 cells when compared to the

WT CaSki cells.

Fig. 6. CaSki cells with lnc-FANCI-2 KO exhibit activation of RAS signaling pathway. (A)

CaSki cells with lnc-FANCI-2 KO (top, ΔPr-A9 cells) or knockdown (KD, lower) by lnc-

FANCI-2-specific siRNAs display increased RAS GTPase activity when compared to the WT

CaSki cells (top) or the WT cells treated with a nonspecific control siRNA (siNS). The relative

RAS GTPase activity was measured by a RAS GTPase Chemi ELISA assay. *, P< 0.05; ***,

P<0.001 by two tailed Student *t* test. (B and C) Selective validation of increased expression of

RAS signaling-related downstream genes in lnc-FANCI-2 KO ΔPr-A9 cells (B) over the WT

CaSki cells or lnc-FANCI-2 KD CaSki cells (C) over the WT cells treated with a nonspecific

control siRNA (siNS). Relative expression of the indicated genes in ΔPr-A9 cells in comparison

to the WT cells (B) or in the WT CaSki cells treated for 48 h and 96 h with a nonspecific control

siRNA (siNS) or a siRNA specifically targeting the lnc-FANCI-2 exon 3 (C) were

immunoblotted using the corresponding antibodies as indicated. Tubulin served as an internal

control for each blot. The level of p-Akt or p-Erk was calculated by normalizing to total Akt or

Erk protein and the other proteins by normalizing to tubulin. The protein level in the WT cells or

WT cells treated with siNS set as 1. The KD efficiency of lnc-FANCI-2 RNA (C, top bar graphs)

was examined by RT-qPCR. (D) Effect of blocking RAS signaling on the expression of MCAM

and VIM using PI3K inhibitor LY294002 (20 μM) and MEK inhibitor U0126 (10 μM). The

protein at each time point was immunoblotted with a corresponding antibody. The level of

MCAM or VIM was calculated after normalizing to tubulin. The protein level in ΔPr-A9 cells

without the inhibitor was set as 100%. *, nonspecific protein band. (E) The time dependent cell

viability of the WT CaSki and ΔPr-A9 cells in the presence of 20 μM LY294002 or 10 μM

U0126. Data was obtained in each time point after normalizing to the cells treated with DMSO.

The mean $\pm$ SD at each data point was calculated from 6 samples combined from two

independent experiments.

Fig. 7. lnc-FANCI-2 is suppressive to the expression of IGFBP3 and MCAM. (A and B)

Transient rescue expression of lnc-FANCI-2 in Δpr-A9 cells inhibits the expression of IGFBP3.

ΔPr-A9 cells were transfected with a major isoform lnc-FANCI-2a-PA2 (GenBank ACC. No.

MT669800.1) cDNA plasmid. lnc-FANCI-2 in red and IGFBP3 RNA in green were detected by

RNAscope RNA-ISH at 24h post transfection using each specific antisense RNA probe and

imaged by confocal microscopy (A). Expression levels of lnc-FANCI-2 RNA and IGFBP3 RNA

in four neighboring cells (B) were measured by signal intensity of a line crossing over the stained

cells in A (white line arrow). (C) Validation of differential expression of MCAM RNA in ΔPr-

A9 cells by RT-PCR in the presence (+) or absence (-) of reverse transcriptase (RT). One pair of

primers with one primer at the exons 11 and the other at exon 13 of MCAM RNA (NM_006500)

were used. GAPDH served as a loading control. (D) The major isoform lnc-FANCI-2a-PA2

repressed the expression of MCAM. ΔPr-A9 cells were transfected with lnc-FANCI-2a-PA2

cDNA plasmid. lnc-FANCI-2 RNA in red was detected by RNAscope ISH and MCAM protein

in green was detected by IF with an anti-MCAM antibody. (E) Expression levels of lnc-FANCI-

2 RNA and MCAM protein in three neighboring cells were measured by signal intensity of a line

crossing over the stained cells (D, white line arrow). (F) Calculation of the expression levels of

MCAM in lnc-FANCI-2 positive cells (n=35) and lnc-FANCI-2 negative cells (n=54) by

fluorescent intensity. (G) Subcellular MCAM distributions in the nucleus (N) and cytoplasm (C)

of WT CaSki cells and ΔPr-A9 by immunoblot analysis. Fractionation efficiency and sample

loading were controlled by cytoplasmic (represented by tubulin) and nuclear (represented by

hnRNP C1/C2) proteins. (H-I) Correlation and survival analysis of lnc-FANCI-2 and MCAM

expression with cervical squamous cell carcinoma (CESC) cases from the cancer genome atlas

(TCGA) datasets by GEPIA web server (http://gepia.cancer-pku.cn/). The negative correlation at

R= -0.321 (with p-value=1.02e-08) of lnc-FANCI-2 with MCAM in cervical cancer patients was

obtained from the RNA-seq data from the TCGA CESC tumors downloaded from the TCGA

data portal (https://portal.gdc.cancer.gov/). Only primary solid tumor samples (n=304 after

exclusion of 2 metastatic samples and 3 normal samples) were subjected to analysis, with the

data showing as a Scatter plot (H). Kaplan-Meier plot with log rank test p-value at 5.74e-05 (I)

shows MCAM as a biomarker for poor prognosis of cervical cancer survival with a lower

quartile group cutoff. RNA-seq and survival data are derived from the TCGA CESC cancer

patients. (J/K) AKT and ERK phosphorylation is partially regulated by MCAM and IGFBP3 in

CaSki cells. KD of MCAM (J) and IGFBP3 (K) expression in WT parental CaSki or ΔPr-A9

cells was treated with MCAM siRNA or IGFBP3 siRNA along with or without lnc-FANCI-2

siRNA for 48 h. Expression levels of individual proteins were immunoblotted using the

corresponding antibodies. The level of MCAM, p-Akt, or p-Erk1/2 was calculated after

normalizing to tubulin. The protein level in a non-targeting siRNA (siNS) control set as 1.

Fig. 8. lnc-FANCI-2 interacts with host factors to regulate RAS signaling pathway. (A) The lnc-

FANCI-2-associated proteins in the WT CaSki cells were identified by isolation of RNA-protein

complexes using RNA purification (IRPCRP)-mass spectrometry technology. (B) lnc-FANCI-2

RNA in the IRPCRP-1 and IRPCRP-2 pulldowns had the pooled antisense biotinylated oligos

(pool 1 with oligos in even numbers and pool 2 with oligos in odd numbers) immobilized to

avidin-beads first before mixed with cell lysates, while the IRPCRP-3 and IRPCRP-4 pulldowns

had the oligos pool 1 and 2 separately mixed with cell lysates first before addition to the avidin-

beads for the RNA pull-downs. RT-PCR in the absence (-) or presence (+) of reverse

transcriptase (RT) was carried out using the RNA isolated from the individual IRPCRP

experiments using a primer pair of oHBL5 and oHBL12 (Table S5) specific for lnc-FANCI-2

RNA detection. Beads only (no oligos) IRPCRP experiments served as a negative control. Total

RNA from the WT CaSki cells after sonication was used as an input control. The arrow indicates

the detected lnc-FANCI-2 RNA. (C) Proteins associated with lnc-FANCI-2 RNA identified from

lnc-FANCI-2 IRPCRP pulldowns. A total of 32 proteins were specifically pulled down from lnc-

FANCI-2 IRPCRP reactions 1-4 (PSM$\geq$2 from two separate pulldowns, Table S4), with top 10

proteins binding lnc-FANCI-2 shown in the order by the number of identified PSM (peptide

spectrum match) in LC-MS/MS. (D) Expression of p-Akt and p-Erk from CaSki WT cells 48 h

after siRNA KD of MAP4K4 or lnc-FANCI-2 was immunoblotted by the corresponding

antibodies. GAPDH served as a sample loading control. Fold change of the indicated proteins in

the cells with KD of MAP4K4 or lnc-FANCI-2 over the cells treated by a non-targeting siRNA

(siNS) was calculated after normalizing to GAPDH. (E) A proposed model illustrates how lnc-

FANCI-2-protein complexes inhibit the RAS signaling pathway to control Akt/Erk

phosphorylation and expression of host genes. RAS signaling can be regulated by integrating

external and internal factors. In HR-HPV infected cells, viral oncoprotein E7-YY1 complex

transactivates lnc-FANCI-2 expression. By interactions with cellular RBPs, the lnc-FANCI-2-

protein complex inhibits RAS signaling. In the absence of lnc-FANCI-2, increased RAS

signaling leads to phosphorylation of Akt and Erk and cascaded responses of transcription

factors (TFs) and thus regulates the expression of RAS signaling responder genes, such as

IGFBP3, MCAM, etc. Consequently, this brings fundamental biochemical and biological

processes under control by finetuning the RAS pathway.

1252

Supporting Tables

Table S1. Genome-wide Effect of lnc-FANCI-2 KO in ΔPr-A9 and ΔYY1 cells on gene

expression when compared with parental WT CaSki cells.

Table S2. The genes with altered expression found in both ΔPr-A9 and ΔYY1-D5 cells when

compared with parental WT CaSki cells.

Table S3. The genes with increased expression in KRAS signaling and EMT and with decreased

expression in IFN-gamma and IFN-alpha responses in lnc-FANCI-2 KO cells (ΔPr-A9 and

ΔYY1-D5) over the parental WT CaSki cells.

Table S4. CaSki proteins and their PSMs identified by lnc-FANCI-2 IRPCRP-LC-MS/MS

Table S5. Sequences of oligonucleotides used in this study.

Supporting Figures

Figure S1. Selective expression of lnc-FANCI-2 RNA in cervical cancer cell lines.

Figure S2. lnc-FANCI-2 KO on characteristic CaSki cell growth behavior and morphology

imaged at 24 h after spreading.

Figure S3. lnc-FANCI-2 KO on expression of viral oncoproteins and cell senescence.

Figure S4. GSEA Enrichment plots show enrichment scores and gene hits enriched in ΔYY1-D5

cells using Hallmark gene sets.

Figure S5. Pathway map of RAS Initiative with highlighted differentially expressed genes

(DEGs) in ΔPr-A9 cells vs parental wild type cells.

Figure S6. Knockout or Knockdown of lnc-FANCI-2 affects RAS signaling.

Figure S7. Expression of IGFBP3, MCAM, MAP4K4 and lnc-FANCI-2.

Figure S8. Expression levels of lnc-FANCI-2 and MCAM are significantly associated with

TCGA CESC (cervical squamous cell carcinoma) cancer patients' survival outcome.

A Cervical cancer tissue-tumor nest and its adjacent normal tissue area

Normal tissue

Tumor nest

Normal tissue

Tumor nest

H&E staining

B Cervical cancer tissue-Case 1

Cervical cancer tissue-Case 2

C

HFK HFK-HPV16 HFK-HPV18

Fig. 3

A

ΔPr-A9 VS WT

pathway	NES
KRAS_SIGNALING_UP	1.858008
EPITHELIAL_MESENCHYMAL_TRANSITION	1.771138
INFLAMMATORY_RESPONSE	1.490227
INTERFERON_GAMMA_RESPONSE	-1.47303
ADIPOGENESIS	-1.50073
INTERFERON_ALPHA_RESPONSE	-1.62585
BILE_ACID_METABOLISM	-1.64176

ΔYY1-D5 VS WT

pathway	NES
TNFA_SIGNALING_VIA_NFKB	1.911117
KRAS_SIGNALING_UP	1.746758
EPITHELIAL_MESENCHYMAL_TRANSITION	1.711204
ESTROGEN_RESPONSE_LATE	-1.63355
INTERFERON_GAMMA_RESPONSE	-1.69812
ESTROGEN_RESPONSE_EARLY	-1.71013
INTERFERON_ALPHA_RESPONSE	-2.22382

B KRAS_SIGNALING_UP
padj=0.0110, NES=1.86

C EPITHELIAL_MESENCHYMAL_TRANSITION
padj=0.0106, NES=1.77

D INTERFERON_GAMMA_RESPONSE
padj=0.045, NES=−1.47

E INTERFERON_ALPHA_RESPONSE
padj=0.040, NES=−1.63

A

UV crosslinking — Cell lysis — Biotinylated as-oligos to lnc-FANCI-2 / Magnetic avidin-beads — Hybridization & pulldown — RNA:protein complexes — Proteins (Mass Spec) / RNA (RT-PCR)

CaSki cells

Magnet

lnc-FANCI-2

B

IRPCRP

lnc-FANCI-2 RNA

bp	RT	input		Beads only		1		2		3		4	
		−	+	−	+	−	+	−	+	−	+	−	+

400-
300-
200-

% pulldown 100 0 70 65 115 132

C

1	H13
2	HNRH1
3	K1H1
4	MAP4K4
5	RNPS1
6	S27A3
7	TNR1B
8	TNC18
9	TBA3D
10	RL8

Top 10 proteins associated with lnc-FANCI-2 RNA

D

siNS | siMAP4K4 | silnc-FANCI-2 | kDa

MAP4K4 — -150
MCAM
1.0 1.7 2.0

IGFBP3 — -100 / -35
1.0 1.3 1.3

GAPDH — -37

p-Akt — -50
1.0 1.4 1.5

Akt — -50

p-Erk1/2 — -35
1.0 1.5 1.6

Erk1/2 — -35

GAPDH — -37

1 2 3

E

RBPs

lnc-FANCI-2 RNA

RAS — MCAM
PI3K — MAPK — IGFBP3
P Akt — P Erk
TFs

YY1 / E7

TFs — IGFBP3 / MCAM

Nucleus

HR-HPVs

Cytoplasm

Fig S1. Selective expression of lnc-FANCI-2 RNA in cervical cancer cell lines. (A)) RT-qPCR detection of lnc-FANCI-2 in HPV16+ cervical cancer cell lines SiHa, CaSki, and W12 20861 (integrated HPV16) and 20863 (episomal HPV16), HPV18+ cervical cancer cell lines HeLa and C4II, and HPV- cell lines C33A (cervical cancer cells with mutations of p53 and pRb), HCT116 (colorectal cancer cells), BCBL-1 (body cavity B lymphoma cells), HEK293 (Ad5 E1/E2-immortalized human kidney cells), and HaCaT (spontaneously immortalized human epidermal cells) in triplicates. (B) HeLa cells express no lnc-FANCI-2 when compared with C33A cells by Northern blot. (C) lnc-FANCI-2 is mainly cytoplasmic in CaSki but nuclear in SiHa and C33A cells. Cytoplasmic and nuclear fractionation efficiency was blotted for nuclear SRSF3 (serine- and arginine-rich splicing factor 3) and cytoplasmic GAPDH. Total fractionated cytoplasmic and nuclear RNAs were quantified for lnc-FANCI-2 by RT-qPCR in triplicates, with GAPDH RNA serving as an internal control for RNA fractionation efficiency. (D) Subcellular lnc-FANCI-2 (red) localization in CaSki, SiHa, and C33A cells determined by RNAscope RNA ISH analysis. Nuclei were stained with DAPI (blue). Scale bars: 25 μm in the top and 10 μm in the zoom.

A

B

Fig. S2. lnc-FANCI-2 KO on characteristic CaSki cell growth behavior and morphology imaged at 24 h after spreading. (A) Parental WT CaSki and ΔYY1-D5 CaSki cells. (B) Parental WT CaSki and ΔPr-A9 CaSki cells.

Fig. S3. lnc-FANCI-2 KO on the expression of viral oncoproteins and cell senescence. (A) Effect of lnc-FANCI-2 KO in CaSki cells on the expression of HPV16 E6 and E7 and their downstream targets. Total cell extracts from parental CaSki, ΔPr-A9, and ΔPr-B3 cells were immunobloted with the corresponding antibodies. Tubulin or GAPDH served as a protein loading control. The relative protein levels of E6, E7, p53, and E2F1 were calculated after normalizing to tubulin or GAPDH in the corresponding experiment. Expt I or II = experiment I or II. (B) KO of lnc-FANCI-2 expression enhances cell senescence in β-gal analysis.

Fig S4. GSEA Enrichment plots show enrichment scores and gene hits enriched in ΔYY1-D5 cells using Hallmark gene sets. (A) KRAS_SIGNALING_UP. (B) EPITHELIAL_MESENCHYMAL_TRANSITION. (C) INTERFERON_GAMMA_RESPONSE. (D) INTERFERON_ALPHA_RESPONSE. The heatmaps on the right side of Enrichment plot in the panels A and B visualize the genes with differentiated expression enriched in each pathway in ΔYY1-D5 cells.

Fig. S5. Pathway map of RAS Initiative with highlighted differentially expressed genes (DEGs) in ΔPr-A9 cells vs parental wild type cells. DEGs are mainly distributed and clustered in two main branches of RAS pathway: MAPK signaling branch and PI3K/AKT branches. DEGs were derived from RNA-seq data at cutoff of adjusted p-value <=0.01 and fold change>1.5 for both up- and down-regulated genes (indicated by red and blue arrows respectively) between ΔPr-A9 vs WT CaSki cells using three common analysis methods (DESeq2, edgeR, and limma-voom) and in-house BRB analysis pipeline from NCI. Highlighted genes in the pathway map are derived as DEGs in at least three of the analysis methods, although in most cases, these DEGs behaved consistently across all 4 methods. MAPK signaling branch is highlighted in transparent green box and PI3K/AKT branch is highlighted in transparent yellow box within the pathway map of RAS signaling pathway, which was collectively collated from community inputs that were organized and stimulated by RAS Initiative, and maintained as a common knowledge basis at URL below: https://www.cancer.gov/research/key-initiatives/ras/ras-central/blog/2015/ras-pathway-v2, which was also described in recent review (Figure 1 in Nissley and McCormick, 2022).

Reference (where the Ras pathway map has been described):
Nissley, D. V. and McCormick, F. (2022). RAS at 40: update from the RAS Initiative. *Cancer Discov.* https://doi.org/10.1158/2159-8290.CD-21-1554

Fig S6. Knockout or Knockdown of lnc-FANCI-2 affects RAS signaling. (A) Selective validation of increased expression of RAS signaling-related downstream genes in ΔPr-A9 and ΔYY1-D5 cells. (B) KD of lnc-FANCI-2 expression in SiHa cells enhanced phosphorylation of Akt and Erk1/2 but did not in HeLa cells expressing no lnc-FANCI-2.

Fig. S7. Expression of IGFBP3, MCAM, and MAP4K4 and lnc-FANCI-2. (A-B) RNA-seq reads-coverage of IGFBP3 and MCAM by IGV illustrates the increased expression of IGFBP3 and MCAM in lnc-FANCI-2 KO cells ΔPr-A9 and/or ΔYY1-D5 when compared to the parental WT CaSki cells. (C) KD of MAP4K4 expression in lnc-FANCI-2 KO ΔPr-A9 cells further enhances phosphorylation of Erk1/2 but little on MCAM and p-Akt and no effect on IGFBP3.

Fig. S8. Expression levels of lnc-FANCI-2 and MCAM are significantly associated with TCGA CESC (cervical squamous cell carcinoma) cancer patients' survival outcome. The expression of lnc-FANCI-2 (A) and of MCAM (B) along with survival of the same group of 304 cervical cancer patients were analyzed by our in-house GradientScanSurv pipeline (Yi et al 2018, PLoS One 13:e0207590). The **open grey circles** in each plot are ranked expression levels from high (left) to low (right) of lnc-FANCI-2 (A) and MCAM (B) from all patient samples as indicated by y-axis at the right side of the plot (in RSEM normalized value of RNAseq data). At each breaking point denoted by patient number (x-axis), the log rank test (y-axis at the left side of the plot) was performed between the higher expression group at the left side of the cut-point vs the lower expression group at the right side of the cut-point and the indicated log rank p-values (**blue dots**) are shown as at each cut-point at x-axis. The **horizontal red line** is the p=0.05 cutoff line for the log rank test p-values. If a log rank p-value (**blue dot**) at a cut-point is below the cutoff (p<0.05), there will be a **vertical green line** shown at the corresponding cut-point, indicative of a significant log rank test p-value, and also a **brown** diamond plotted either at the bottom part of the **vertical green line** (A) if higher expression of lnc-FANCI-2 led to less severe outcome (dying slower than the lower expressors), or at the top part of the **vertical green line** (B) if higher expression of MCAM led to more severe outcome (dying faster than the lower expressors).

The original GoodCount was defined as the number of significant log rank tests (the default significance level was set as 0.05) across all possible cut-points for the original dataset. We employed the bootstrap approach to permutate the data for 1000 times in this case. A similar procedure each time was performed like the original dataset and a permutated GoodCount was obtained for each permutated dataset. The GoodCount p-value was derived as the proportion of the times that the GoodCounts of permutated datasets were no less than that of the original dataset (Yi et al 2018, PLoS One 13:e0207590) and shown at the left top corner of each plot, 0.011 for lnc-FANCI-2 in panel A and 0.002 for MCAM in panel B. A significant GoodCount p-value (**blue dots** in both panels A and B) is indicative of significant association of the expression levels with the corresponding survival outcomes of the analyzed patient samples. Not shown in the panels A and B are the coxph p-value from univariant expression-based cox regression models (coxph p-value: derived by original gene expression values; or coxph p-value by rank: derived by corresponding ranks of gene expression values) and the cut-point-specific FDR and FDR2. FDR for each cut-point was defined as the portion of cases of permutated datasets that have log rank test p-values no higher than that of original dataset. FDR2 for each cut-point was defined as the portion of cases of permutated datasets that have log rank test p-values no higher than default setting of significance as 0.05.