The long noncoding RNA lnc-FANCI-2 restricts RAS signaling and phosphorylation of Akt

and Erk in HPV16-infected cervical cancer cells

Haibin Liu[1a,b], Lulu Yu[1a], Vladimir Majerciak[1], Thomas Meyer[2], Ming Yi[3], Peter F. Johnson[4],
Maggie Cam[2], Douglas R. Lowy[5], and Zhi-Ming Zheng[1]*

[1]Tumor Virus RNA Biology Section, HIV Dynamics and Replication Program, CCR/NCI/NIH

[2]Collaborative Bioinformatics Resource, CCR/NCI/NIH

[3]RAS Informatics Group, NCI/NIH

[4]Mouse Cancer Genetics Program, CCR/NCI/NIH

[5]Laboratory of Cellular Oncology, CCR/NCI/NIH

[a]Co-first authors

Running title: lnc-FANCI-2 regulates RAS signaling

[b]Present address: Center for Emerging Infectious Diseases, Wuhan Institute of Virology, and

Center for Biosafety Mega-Science, Chinese Academy of Sciences, Wuhan, Hubei, 430207,

China

*Corresponding: zhengt@exchange.nih.gov

ABSTRACT

We recently discovered increased expression of a cytoplasmic long noncoding RNA, lnc-FANCI-2, coinciding with cervical lesion progression from CIN1, CIN2-3 to cervical cancer. Viral E7 of high-risk HPVs and host transcription factor YY1 are two major factors promoting lnc-FANCI-2 expression. To explore possible roles of lnc-FANCI-2 in HPV-induced cervical carcinogenesis, we ablated the expression of *lnc-FANCI-2* in the HPV16-positive cervical cancer cell line, CaSki. Knock-out (KO) single cell clones expressed HPV16 oncogenes normally but displayed altered cell morphology and reduced proliferation and migration when compared with the parental cells. Proteomic profiling of cytosolic and secreted proteins from the parental and KO cells showed that lnc-FANCI-2 regulates expression of a subset of cell surface and adhesion-related proteins, including inhibition of MCAM, PODXL2 and ECM1 and increased levels of ADAM8 and TIMP2. RNA-seq analyses revealed that, relative to the parental cells, KO cells exhibited significantly increased RAS signaling but decreased IFN pathways. In KO cells, phosphorylated Akt and Erk1/2, two important RAS pathway effectors, were increased more than 3-fold, accompanied by increase of MCAM, VIM, and CCND2 (cyclin D2) and decrease of RAC3. Accordingly, high levels of lnc-FANCI-2 and lower levels of MCAM in cervical cancer patients are associated with improved survival. We found that lnc-FANCI-2 in CaSki cells interacts specifically with 32 host proteins, including H13, HNRH1, K1H1, MAP4K4, and RNPS1, and knockdown of MAP4K4 led to decreased phosphorylation of Akt and Erk1/2. In summary, a key function of lnc-FANCI-2 is to regulate RAS signaling, thereby affecting cervical cancer outcome.

INTRODUCTION

Cervical cancer is a leading cause of mortality in women worldwide. It is the second most diagnosed cancer among women, with an estimated 530,000 new cases worldwide each year, and the third most frequent cause of cancer-related death, accounting for 270,000 deaths annually[1]. Persistent infections in the cervical epithelium with the cancer-associated alpha human papillomaviruses (high-risk HPVs), in particular HPV16 and HPV18 with the sustained expression of viral oncoproteins, E6 and E7, contribute to squamous epithelial cell tumorigenesis [2-4]. Approximate 1.4% of high-grade neoplastic cervical lesions progress to invasive cervical cancer [5], while most lesions resolve spontaneously. The progression to cervical cancer typically takes 15 to 20 years after initial infection (https://www.who.int/news-room/fact-sheets/detail/cervical-cancer). These observations suggest that active host defense systems play important roles in preventing malignant transformation of neoplastic cervical lesions, and additional factors besides E6 and E7 are necessary for cervical cancer development. Indeed, one tumor suppressor network, the Fanconi anemia (FA) pathway [6], can be activated by HPV-induced DNA damage. FA proteins including FANCA, FANCD2 and FANCI are elevated upon HPV infection [7-9] and the activated FA pathway restricts HPV replication and guards host genome stability [10]. FANCD2 mutation stimulates HPV16 and HPV31 genome amplification and also promotes cervical and vaginal cancer development in HPV16 E7 transgenic mouse [11, 12].

Viral oncoprotein E7 plays a major role in the immortalization of primary epithelial cells mainly by inactivation of pRb family of proteins [13], whereas E6 mediates degradation of tumor suppressor p53 and activation of human telomerase reverse transcriptase (hTERT) transcription [14-16]. However, high-risk E6 and E7 are necessary but not sufficient to transform fully the immortalized cells into malignant cells [17-21] and additional oncogenic stress is needed. For

example, the HRAS[G12V], a constitutively activated RAS GTPase mutation, triggers malignant

transformation of E6/E7-expressing keratinocytes[22-25]. Normal RAS regulates cell proliferation,

cell differentiation and cell adhesion through cellular signal transduction from growth-factor

receptors. Thus, overactive RAS signaling might facilitate transformation of E6/E7-expressing

keratinocytes. In transgenic mice, HPV16 E6 and E7 can promote the development of

spontaneous epithelial skin tumors, but not spontaneous tumors of the reproductive tract [26-29].

Prolonged estrogen treatment is required [30]. Estrogen receptor (ER) activation stimulates the

mitogen-activated protein kinase (MAPK/Erk) and phosphoinositide 3-kinase (Pl3K/Akt)

pathways, both of which are mediated by RAS [31]. However, the molecular mechanism

underlying estrogen-induced cervical cancer in E6 and E7 transgene mice is not fully understood.

Long non-coding RNAs (lncRNAs) are RNAs over 200 nucleotides (nt) in length lacking

a coding capacity for translation into functional proteins. Although several lncRNAs have been

proposed to function in diverse biological processes, evidence is still lacking to support the

functionality of the majority of lncRNAs [32]. Nevertheless, the identified lncRNA functions

include regulating: (1) transcription of neighboring or distant genes by recruiting histone

modifiers, as shown for Xist [33] and ANRIL [34], or by causing chromosome looping, as proposed

for PVT1 [35] and CCAT1-L [36]. (2) RNA splicing by acting as decoy of splicing factors, as

demonstrated for MALAT1 [37] and PNCTR [38]. (3) mRNA stability by directly base pairing as

antisense transcripts [39], by acting as RNA binding protein (RBP) decoy to prevent mRNA-RBP

interaction, such is the case for NORAD [40], or by functioning as miRNA sponges to prevent

miRNA-induced degradation of mRNAs, as shown for PTEN [41] and PNUTS [42]. And (4) protein

stability by directly binding, such as MaIL1 [43], or by acting as protein decoy to prevent ubiquitin-

mediated proteasomal degradation, such as FAST [44]. Thus, lncRNA genomic position,

subcellular localization and their interactions with DNA, RNA and proteins are crucial

determinants to the functionality of individual lncRNAs.

We recently reported an increased level of lnc-FANCI-2 expression in high-risk HPV-

positive cervical intraepithelial neoplasia (CIN) and invasive cervical cancer (ICC) tissues [8]. At

least 14 RNA isoforms of lnc-FANCI-2 are expressed by alternative promoter usage, alternative

RNA splicing, and alternative selection of RNA polyadenylation sites from the genomic locus

adjacent to the FANCI. Both lnc-FANCI-2 and FANCI are up-regulated simultaneously in

neoplastic cervical lesions and cervical cancer by high-risk HPV infections. E6 and in particular,

E7 are responsible for the enhanced expression of lnc-FANCI-2, the transcription of which is

also regulated by YY1[8]. HPV infection increases YY1 levels but decreases the expression of

p53-dependent miR-29a which targets the YY1 3' UTR. Viral E7 interacts with YY1 and

facilitates YY1 transactivation of lnc-FANCI-2 promoter [8]. In situ hybridization has revealed

that lnc-FANCI-2 is preferentially cytoplasmic, but its function in HPV infected cells has not

been characterized. In this report, we demonstrate that knockout (KO) or knockdown (KD) of

lnc-FANCI-2 promotes RAS signaling and phosphorylation of Akt and Erk.

RESULTS

lnc-FANCI-2 regulates proliferation of HPV-transformed cervical cancer cells.

lnc-FANCI-2 RNA is transcribed mainly from a proximal promoter TSS2, but also from

an alternative minor, distal promoter TSS1. Two highly conserved YY1 binding motifs upstream

of the TSS2 are essential for the TSS2 transcriptional activity [8] (Fig.1A). To elucidate the

function of lnc-FANCI-2 in high-risk HPV-infected cervical cancer cells, we knocked out (KO)

lnc-FANCI-2 expression in HPV16-positive CaSki cells using CRISPR/Cas9 deletion of either a

3-kb promoter region encompassing both TSS1 and TSS2 or an 86-bp region containing two

YY1-binding motifs in the TSS2 promoter (Fig.1A). Two tested gRNAs with high KO efficiency

were selected and cloned into a modified CRISPR/Cas9 expression vector to express both 5'-

specific gRNA and 3'-specific gRNA simultaneously for efficient genome editing [45]. CaSki cells

stably transfected with the dual gRNA expression vector were generated. Through a serial

dilution, several single cell clones were isolated and verified for homozygous deletion by PCR

screening. PCR genotyping indicated successful homozygous deletion of the promoter or YY1-

binding motifs, respectively (Fig.1B).

Loss of lnc-FANCI-2 expression in the single-cell clones was examined by RT-qPCR

(Fig.1C). When compared to all control clones transfected and selected from an empty vector

containing no gRNA, all single cell clones with deletion of YY1-binding motifs (ΔYY1) showed

60-70% reduction of lnc-FANCI-2 expression, whereas the single cell clones with deletion of

both TSS1 and TSS2 promoters (ΔPr) showed at least 90% reduction (Fig. 1C). Northern blot

analysis confirmed the decrease in levels of the two major lnc-FANCI-2 isoforms (Fig. 1D), with

an abundant 4-kb lnc-FANCI-2 RNA derived from a distal polyadenylation site PA2 and a less

abundant 2-kb of lnc-FANCI-2 RNA derived from a proximal polyadenylation site PA1[8] (Fig.

1A and 1D). Reduced lnc-FANCI-2 RNA expression was also confirmed by RNAscope single

molecule RNA in situ hybridization (RNA_ISH) (Fig. 1E and 1F). The parental (WT) CaSki

cells expressed ~10 copies of lnc-FANCI-2 RNA per cell [8], whereas the copy number of lnc-

FANCI-2 RNA in the ΔYY1-D5 cells dropped to ~5 copies and to ~2 copies per cell in the ΔPr-

A9 cells (each averaged from 200 cells).

Although the WT CaSki cells grow as cell islands, the lnc-FANCI-2 KO cells displayed a

dispersed cell growth pattern and an irregular or spindle-like cell morphology which was more

obvious in the ΔPr-A9 cells (Fig.S1A). The KO cells also grew significantly slower than the WT

CaSki cells (Fig. 1G), formed fewer colonies with reduced size (Fig. 1H), and showed greatly

decreased migration capacities (Fig. 1I), with ΔPr-A9 cells having a more severe phenotype than

ΔYY1-D5 cells. The ΔPr-A9 cells were further examined for effects on HPV16 E6 and E7

expression and cell senescence. When compared with WT CaSki cells, the ΔPr-A9 cells showed

little change in E6, E7, p53 (E6 downstream target) and E2F1 (E7 downstream target) protein

levels (Fig. S1B) but a significant increase in cell senescence (Fig. S1C). Altogether, the data

indicate that lnc-FANCI-2 RNA is important for cell growth, colony formation, and cell

migration.

lnc-FANCI-2 regulates the expression and secretion of cell soluble receptors.

We considered that altered expression of cell membrane proteins and secreted factors in

the lnc-FANCI-2 KO cells might contribute to the observed different cell morphology and

growth properties. Therefore, we performed a Proteome Profiler Human sReceptor Array

analysis to examine possible changes in expression of 105 well-characterized soluble protein

receptors using total cell lysates and cell culture supernatants from ΔPr-A9 and WT CaSki cells.

We identified from the ΔPr-A9 cell lysate nine proteins with increased expression

compared to the WT CaSki cells, including PODXL2, ECM1, NECTIN2, MCAM, ADAM9,

ADAM10, CDH5, ITGA5 and NOTCH1, and six proteins with decreased expression including

ITGB6, CDH13, LGALS3BP, TIMP2, ADMA8 and SCARF2 (Fig. 2A and 2B). We also found

from the ΔPr-A9 cell culture supernatant five proteins with increased expression, including

ADAM9, NECTIN2, ADAM10, CDH5 and ECM1, and the decreased expression of four

proteins, including CRELD2, SDC1, SDC4 and TIMP2 (Fig. 2A and 2B). By immunoblot

assays, we verified selectively the increase in PODXL2, MCAM, and ECM1 and the decrease in

ADAM8 and TIMP2 from the ΔPr-A9 cell lysates and the increase in ECM1 and the decrease in

TIMP2 in the ΔPr-A9 cell culture supernatant (Fig. 2C). As ADAM8 is proteolytically processed

into two protein isoforms [46], we confirmed by immunoblot the decreased expression of all three

sizes of ADAM8 protein in the ΔPr-A9 cell lysate over the WT CaSki cells (Fig. 2C).

lnc-FANCI-2 regulates the expression of genes involved in RAS signaling.

We next conducted genome-wide RNA-seq analyses of WT and lnc-FANCI-2 KO cells

(four samples/group) to determine the transcriptomic consequences of lnc-FANCI-2 deficiency.

We obtained ~120 million mappable RNA reads to the human reference genome hg38 from each

sample. Analysis of RNA-seq reads-coverage map by Integrative Genomics Viewer (IGV)

showed ~90% reduction of lnc-FANCI-2 expression in ΔPr-A9 cells and ~72% decrease in

ΔYY1-D5 cells compared to the WT CaSki cells (Fig. 3A), consistent with the results shown in

Fig. 1C-F.

Hierarchical clustering analysis showed more global transcriptional similarity between

ΔYY1-D5 and ΔPr-A9 cells (Fig. 3B). By applying a threshold of fold change (FC) $\geq$ 1.8 or FC

$\leq$ -1.8 with a p value of FDR $\leq$ 0.01, we found 1230 genes in ΔPr-A9 and 797 genes in ΔYY1-

D5 cells with significantly differential expression relative to the WT CaSki cells expressing

~15890 genes. The most significantly affected genes are shown in Volcano plots (Fig. 3C and

Table S1). Among these, 211 were upregulated and 189 were downregulated in both ΔPr-A9 and

Δ YY1-D5 cells (Fig. 3D and Table S2). By applying more stringent criteria with FPKM $\geq$7 in at

least one of four RNA-seq samples as a cutoff, we profiled the genes with large expression

differences in both ΔPr-A9 and ΔYY1-D5 cells. The results displayed in the heatmap of Fig. 3E

included 51 upregulated with only one outlier, IGFBP3, and 47 downregulated genes excluding

lnc-FANCI-2. Subsequently, we selectively verified by RT-PCR the decreased expression of

ITGB6, NLRP2, PLAC8, PSG4, PSMB9, and SERPINB1 and increased expression of CFH,

CNTN5, EMP3, IGFBP3, PTGS2, and SERPINB2 in ΔPr-A9 cells (Fig. 3F), as well as the

increased expression of IGFBP3 protein in ΔPr-A9 cells by immunoblot (Fig. 3G).

By performing the Gene Set Enrichment Analysis (GSEA) on the Hallmark gene sets,

which provide more refined and concise inputs for GSEA [47], we found the most significantly

upregulated pathways in both ΔPr-A9 and ΔYY1-D5 cells, when compared with the WT CaSki

cells, were KRAS signaling and epithelial mesenchymal transition (EMT). The most

significantly downregulated pathways in both ΔPr-A9 and ΔYY1-D5 cells over the WT CaSki

cells were interferon gamma (IFN-γ) and interferon alpha (IFN-α) responses (Fig. 4A). GSEA

plots for ΔPr-A9 shows 29 of 111 genes in RAS signaling and 39 of 130 genes in EMT (Fig. 4B

and 4C, Table S3) were significantly upregulated, whereas 37 of 139 genes in IFN-γ response

and 24 of 71 genes in IFN-α response were significantly downregulated (Fig. 4D and 4E, Table

S3). Similar enriched gene sets with increased RAS signaling and EMT and decreased IFN-γ and

IFN-α responses were observed in ΔYY1-D5 cells (Fig. S2, Table S3).

Since ΔPr-A9 cells exhibit higher KO efficiency of lnc-FANCI-2 and a higher number of

differentially expressed genes (DEGs) and show more severe phenotypic changes in cell

proliferation, migration, and colony formation than ΔYY1-D5 cells, the ΔPr-A9 cells were

primarily used for further focused studies on RAS signaling in this report.

lnc-FANCI-2 regulates RAS GTPase activities to phosphorylate RAS signaling effectors

Akt and Erk.

RAS activation triggers two major downstream signal transduction pathways,

Raf/Mek/Erk and PI3K/Akt (Fig. S3), to transduce signals from extracellular stimuli to the cell

nucleus where specific effector genes are activated for their corresponding functions [48, 49]. To

further confirm the GSEA data and verify that two major signaling pathways of RAS are

activated, we first examined and compared the RAS GTPase activities in ΔPr-A9 cells and the

WT CaSki cells. We found a significant increase in RAS GTPase activity in ΔPr-A9 cells (Fig.

5A, top bar graphs), indicating that the endogenous lnc-FANCI-2 RNA in CaSki cells suppresses

RAS activation. Since ΔPr-A9 cells had undergone long term selection and adaption during

single cell screening, the increased RAS GTPase activity might result from selection pressure for

cell survival. However, by transient siRNA knockdown (KD) of lnc-FANCI-2 expression in the

WT CaSki cells, we obtained a similar, albeit weaker, increase of RAS GTPase activity (Fig. 5A,

lower bar graphs). This result suggests that the increased RAS GTPase activity in ΔPr-A9 cells

was unrelated to the persistent selection pressure, but rather the deficiency of lnc-FANCI-2.

Next, we examined activation of the PI3K/Akt and Raf/Mek/Erk pathways [50-52]. We

observed a 3-fold increase in phosphorylated Akt (p-Akt) and 2.5-fold increase of p-Erk1/2

(p44/p42 MAPK), after being normalized to total Akt and Erk, in ΔPr-A9 cells over the WT

CaSki cells (Fig. 5B). These results were confirmed in ΔYY1-D5 cells (Fig. S4A). The increased

p-Akt and p-Erk (mostly p-Erk2/p42) was accompanied by elevated expression of MCAM, VIM,

and CCND2, and decreased expression of RAC3 (Fig. 5B). These randomly chosen potential

RAS effector proteins facilitate membrane receptor function, cell proliferation, and EMT.

Moreover, transient siRNA knockdown of lnc-FANCI-2 in the WT CaSki cells also led to the

increased levels of p-Akt and p-Erk (Fig. 5C) and increased expression of MCAM and VIM at

48 h and 96 h post-transfection (Fig. 5C). These observations indicate that a transient loss of lnc-

FANCI-2 in CaSki cells is sufficient to trigger RAS signaling. The same siRNA transfection in

HeLa cells, a lnc-FANCI-2 negative cell line (Fig. S4B and S4C), exhibited no effect on p-Akt

or p-Erk1/2 levels (Fig. S4F), but in SiHa cells containing mainly nuclear lnc-FANCI-2 (Fig.

S4D and S4E) enhanced the expression of p-Akt and p-Erk1/2 (Fig. S4F), verifying the specific

effect of lnc-FANCI-2 depletion on RAS signaling pathways no matter where a nuclear or

cytoplasmic lnc-FANCI-2 is predominant in the cancer cells. We found both cytoplasmic and

nuclear locations of the increased lnc-FANCI-2 in HPV-infected human foreskin keratinocytes

and cervical cancer tissues (Fig. S5). The lnc-FANCI-2 KO-mediated increase in p-Akt and p-

Erk was sensitive to PI3K inhibitor LY294002 [53] and MEK1/2 inhibitor U0126 [54] (Fig. 5D),

respectively. The inhibitors also decreased the expression of VIM and MCAM and cell

proliferation, in particular by PI3K inhibitor LY294002 (Fig. 5D and 5E). These effects on cell

proliferation were similar from ΔPr-A9 to the WT CaSki cells.

lnc-FANCI-2 inhibits MCAM expression to promote cervical cancer survival.

As shown in Fig. 3E and Fig. S6A, the increased RNA expression of MCAM appears to

be transcriptional or posttranscriptional in both ΔPr-A9 and ΔYY1-D5 cells. We confirmed by

RT-PCR the lnc-FANCI-2 KO-mediated increase of MCAM RNA expression in ΔPr-A9 cells

(Fig. 6A). A rescue experiment was further performed by transient expression of one major

isoform of lnc-FANCI-2 RNA (a-PA2, GenBank: MT669800.1) [8] in ΔPr-A9 cells to determine

both nuclear and cytoplasmic MCAM expression [55, 56] in individual lnc-FANCI-2 expressing

cells. ΔPr-A9 cells ectopically expressing diffused cytoplasmic lnc-FANCI-2 RNA, as detected

by RNAscope RNA-ISH, exhibited a marked reduction in nuclear MCAM protein (Fig. 6B-C)

and cytoplasmic IGFBP3 RNA, another lnc-FANCI-2 effector gene (Fig. S6B-D). Quantitative

analyses of the ΔPr-A9 cells with or without transient lnc-FANCI-2 RNA expression showed

significant reduction of nuclear MCAM protein in the 35 cells with rescued lnc-FANCI-2

expression when compared to the 54 cells expressing no lnc-FANCI-2 (Fig. 6D). Due to

RNAscope procedures, the membrane-bound and cytoplasmic MCAM protein was mostly

removed by protease III treatment when RNA-ISH was performed and thus only the nuclear

signal of MCAM protein remained was detectable by IF staining. Using cell fractionation, we

detected a cleaved MCAM of ~46 kDa mainly in the nuclear fraction by Western blot (Fig. 6E).

The increased nuclear MCAM expression in ΔPr-A9 cells was proportional to the cytoplasmic

and total MCAM levels when compared with the WT CaSki cells (Fig. 6E). The nuclear MCAM

signal appeared as a proteolytic cleavage product presumably by the increased metalloprotease

ADAM9 and ADAM10 proteins and decreased metalloprotease inhibitor TIMP2 (Fig. 2), as

observed in other studies [56]. There was no secreted soluble MCAM detectable in the CaSki cell

culture supernatant (Fig. 6F).

The significance of the inverse correlation of lnc-FANCI-2 and MCAM in CaSki cells

was further investigated by analyzing their expression in 304 cervical cancer samples from the

TCGA dataset. There was a significant negative correlation in lnc-FANCI-2 (LINC00925 being

wrongly assigned by NCBI)[8] and MCAM RNA levels (Fig. 6G). The opposite effects of lnc-

FANCI-2 and MCAM on cervical cancer survival (Fig. 6H and Fig. S7) show that the patients

with a higher level of lnc-FANCI-2 [8] but a lower level of MCAM in the cervical tissue (Fig. S7

and Fig. 6H) exhibited a better survival prognosis for cervical cancer.

MAP4K4 association with lnc-FANCI-2 RNA in CaSki cells regulates RAS signaling and

phosphorylation of Akt and Erk.

We speculated lnc-FANCI-2 restriction on RAS signaling through interactions with

cellular proteins. Subsequently, comprehensive identification of lnc-FANCI-2-binding proteins

in the WT CaSki cells was performed by an IRPCRP protocol (modified from published ChIRP

[57]) in combination of mass spectrometry (Fig. 7A). In this protocol, the lnc-FANCI-2-binding

proteins in the WT CaSki cells were covalently crosslinked to lnc-FANCI-2 RNA via UV

irradiation and pulled down by pooled 30 antisense oligos crossover the entire lnc-FANCI-2. The

RNA extracted from the pulldowns were verified to be lnc-FANCI-2 specific by RT-PCR in the

absence (-) or presence (+) of reverse transcriptase (RT) (Fig. 7B). The lnc-FANCI-2-binding

proteins in the pulldowns were then subjected to LC-MS/MS analyses. We identified 32 specific

lnc-FANCI-2-binding proteins, including H13, HNRH1, K1H1, MAP4K4, and RNPS1(Fig. 7C,

Table S4).

MAP4K4, a serine/threonine protein kinase, stimulates cancer cell proliferation, invasion

and migration, and is recently characterized as a novel MAPK/ERK pathway regulator [58-60] and

regulates RAS signaling in the context of the early embryo [61]. Thus, whether MAP4K4 could

upregulate PI3K/Akt signal transduction was investigated in the WT CaSki cells. We

demonstrated that siRNA knockdown of MAP4K4 expression in the WT CaSki cells led to

~40% reduction of phosphorylation of both Akt and Erk1/2 (Fig. 7D). The data suggest that,

through RNA-protein interactions, the increased lnc-FANCI-2 RNA in cells might titrate

MAP4K4 and other cellular proteins to control RAS signaling and gene expression of its

effectors (Fig. 7E).

DISCUSSION

HPV oncoproteins E6 and E7 are necessary but not sufficient for development of cervical

cancer. Indeed, early studies indicated that other oncogenic stress, such as activated RAS mutant,

is needed to trigger malignant transformation of E6 and E7 expressing cells [17, 22, 62, 63] and

tumorigenesis [23, 24, 64-68]. Moreover, E7 has been also reported to repress phosphorylation of Akt

and Akt mediated signaling[69]. In this study, we show that lnc-FANCI-2, whose expression is

highly dependent on E7 and YY1[8], restricts RAS GTPase activities and Akt and Erk

phosphorylation presumably by interacting with one of cellular factors, MAP4K4, in HPV16-

positive CaSki cells. Interestingly, early reports indicated that p-Akt attenuation by E7 could be

abolished by introduction of H73E mutation in the E7 CR3 domain[69]. This E7 CR3 domain is

also essential to interact with transcription factor YY1 to activate lnc-FANCI-2 transcription [8].

Thus, it would be presumable that the reported p-Akt attenuation by E7 might be mediated by the

increased expression of lnc-FANCI-2. More importantly, these findings also suggest that

additional oncogenic stress, such as activated RAS mutant, is required to overcome the lnc-

FANCI-2 restriction on RAS signaling for the early observed malignant transformation [17, 22, 62, 63]

and tumorigenesis [23, 24, 64-68] of E6 and E7 expressing cells. In CaSki cells, loss of lnc-FANCI-2

in this report was found not only to promote RAS signaling but also retard cell proliferation and

induce cell senescence. This is consistent with previous observations showing that high levels of

mutant HRAS[G12V] induce cell cycle arrest and cell death [70, 71].

Some lncRNAs function as RAS regulators [72, 73] by their ability to sequester RAS-

targeting miRNAs, including MALAT1/ miR-217, RMRP/miR-206, and KRAS1P/miR-143/let-

7 [73]. Conceivably, lncRNAs may also regulate RAS signaling through other mechanisms beyond

miRNAs. In this report, loss of lnc-FANCI-2 in HPV16-positive CaSki and SiHa cells stimulates

RAS signaling and expression of RAS signaling effectors (Fig. 7E). By profiling the proteins

associated with lnc-FANCI-2 in CaSki cells, we identified MAP4K4, a serine/threonine protein

kinase and a negative RAS signaling regulator in the context of the early embryo [59, 61], as one of

major lnc-FANCI-2-binding proteins that may partially mediate the lnc-FANCI-2 restriction on

RAS signaling. MAP4K4 is thought to regulate the c-Jun NH2-terminal kinase (JNK) signaling

[74, 75] and the MAPK/ERK pathway [59]. Indeed, silencing of MAP4K4 in CaSki cells led to

reduction of both p-Akt and p-Erk1/2, rather than an expected enhancement of p-Akt and p-

Erk1/2 expression according to the observations in the context of the early embryo[61].

Although all HPV16-positive cervical cancer and pre-cancer cell lines examined, such as

CaSki, SiHa, W12 subclone cells, produce predominantly cytoplasmic or nuclear lnc-FANCI-2,

unexpectedly, we found not all cervical cancer cell lines, including HPV18-positive HeLa and

C4II cells, express lnc-FANCI-2, nor colorectal cancer cell line HCT116, adenovirus 5 DNA-

transformed human embryonic kidney cell line HEK293, spontaneously transformed aneuploid

immortal keratinocyte cell line HaCaT, and KSHV-infected B cell lymphoma cell line BCBL-1

cells which are all HPV-negative cell lines. It remains to know what negatively regulates lnc-

FANCI-2 expression in those cell lines. However, we did find that HPV-negative cervical cancer

cell line C33A cells expressing mutant p53 and mutant pRB produce a high level of lnc-FANCI-

2 (Fig. S4B and S4C). In the lnc-FANCI-2 producing cells, we noticed that cellular location of

lnc-FANCI-2 also varies from cell types, with lnc-FANCI-2 predominantly in the nucleus in

SiHa cells, but in the cytoplasm in CaSki cells, although both contribute to regulate RAS

signaling. This suggests that a variety of nuclear and cytoplasmic functions of lnc-FANCI-2

remains to be explored.

It is obvious from our RNA-seq and human soluble receptor array analysis that KO of

lnc-FANCI-2 in CaSki cells affects the expression of a large set of genes responsible for RAS

signaling, EMT pathway, and IFN responses. In addition to the notified activation of PI3K/Akt

and Raf/Mek/Erk pathways, one of the major RAS signaling effectors in lnc-FANCI-2 KO or

KD cells was MCAM. MCAM binds various cellular surface receptors or co-receptors to trigger

signal transduction, cell proliferation and motility, tumor angiogenesis, and tumor cell metastasis

[76]. It is an integral, highly glycosylated membrane protein with three common protein isoforms

[77-79]. Two membrane-anchored forms with a long (p113) or short (p76) cytoplasmic tail produced

by alternative RNA splicing and a proteolytically cleaved soluble form without transmembrane

and cytoplasmic regions [78-80]. In addition to its roles in cell signal transduction [76-79], the short

isoform produced by an intramembrane cleavage may be directed towards the nucleus to regulate

gene transcription [56]. We found that CaSki cells express mainly a cytoplasmic form of full-length

MCAM (~113 kDa) and a small nuclear form (~46 kDa), but not the short ~78 kDa isoform nor

the soluble product. Moreover, lnc-FANCI-2 KO cells exhibit high MCAM levels, which was

most likely a result of the increased RAS signaling. Consistently, lnc-FANCI-2 and MCAM

levels are negatively correlated in cervical cancer patients, and low levels of MCAM and high

levels of lnc-FANCI-2 indicate better prognosis. This suggests that increased MCAM might

promote cervical tumorigenesis in vivo.

In conclusion, this study provides the first evidence that one of lnc-FANCI-2 functions is

to maintain epithelial functional integrity by repressing RAS signaling in preventing malignant

transformation of neoplastic cervical lesions. In addition, more extensive studies are undergoing

to elucidate the role of lnc-FANCI-2 in regulation of IFN responses, which are not covered in

this report. Since transactivation of lnc-FANCI-2 expression is largely dependent on host

transcription factor YY1 of which expression is regulated by HPV oncoproteins and host miR-

29a, dissecting their transcription network on lnc-FANCI-2 and its effectors may further provide

novel insights involved in host homeostasis against HPV infection-induced carcinogenesis. Our

observations also highlight additional, unexpected players beyond viral E6 and E7 that drive

cells toward malignant transformation.

MATERIALS AND METHODS

Cell cultures and treatment

CaSki, a HPV16-positive human cervical cancer cell line with wild-type p53, pRb, HRAS, KRAS, and NRAS (https://depmap.org/portal/cell_line/ACH-001336?tab=mutation) was obtained from the American Type Culture Collection (ATCC, Manassas, VA), maintained in Dulbecco's modified Eagle medium (DMEM) (Thermo Fisher Scientific) supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin, and grown in a cell culture incubator at 37°C with 5% CO_2.

SiHa (HPV16+), HeLa (HPV18+), C4II (HPV18+), C33A (no HPV), and other HPV-negative cell lines, HEK293 (Ad5 E1/E2-immortalized human kidney cells), and HaCaT (spontaneously immortalized human epidermal cell), were all obtained from ATCC and grown in Dulbecco's modified Eagle's medium (DMEM) with 10% FBS at 37°C and 5% CO_2. HCT116 cell line, a colorectal cancer cell line, was a gift from Dr. Bert Vogelstein of Johns Hopkins University and grown in McCoy's 5A medium with 10% FBS at 37°C and 5% CO_2. W12 subclones 20861 (HPV16+, integrated) and 20863 (HPV16+, episomal) were gift cells from Paul Lambert of University of Wisconsin and grown on pretreated NIH3T3 feeder cells in F12 medium at 37°C and 5% CO_2. BCBL-1 (body cavity B lymphoma cells with KSHV infection) was obtained from the AIDS Research and Reference Reagent Program, Division of AIDS, NIAID, NIH and cultured in RPMI 1640 containing 10% FBS at 37°C and 5% CO_2.

CaSki cells were treated with 20 µM phosphoinositide 3-kinase (PI3K) inhibitor, LY294002 (Cell Signaling Technology, Danvers, MA, #9901) or 10 µM mitogen-activated protein kinase kinase (MEK) inhibitor, U0126 (Cell Signaling Technology, Danvers, MA, #9903) and

examined for their proliferation and viability by CCK-8 assay and RAS effector expression by

Western blot.

siRNAs, plasmid constructions and transfections

Synthetic double-stranded siRNA targeting the lnc-FANCI-2 transcript is listed in a

supplementary Table S5. Human MAP4K4 siRNA (#M-003971) and non-targeting control

siRNA(#D-001210-01) were purchased from Dharmacon, Inc (Lafayette, CO). Each siRNA

transfection into CaSki or other type of cells was carried out by LipoJet in vitro transfection

reagent (SignaGen, MD, USA). Plasmid pHBL03 has a short isoform of lnc-FANCI-2 cDNA

(lnc-FANCI-2a-PA1, GenBank MT669801). Overlapping PCR products generated from 5' and

3' RACE PCR products with a primer pair of oHBL 17 and oHBL 18 were digested and inserted

into pcDNA3 at Hind III and Xba I sites. The insertion was verified by sequencing. Plasmid

pHBL05 has a long isoform of lnc-FANCI-2 (lnc-FANCI-2a-PA2, GenBank MT669800). PCR

products generated from genomic DNA with oHBL 19 and oHBL 23 were digested and swapped

into pHBL03 to replace lnc-FANCI-2a-PA1 at Bsmb I and Xba I sites.

CRISPR/Cas9 system was used to knockout lnc-FANCI-2 gene from CaSki cells. Two lnc-

FANCI-2 specific guide RNAs (gRNAs) were designed by an on-line CRISPR design tool

developed by the Zhang lab (http://crispr.mit.edu). A modified cloning strategy was used to

clone the two gRNA into pSpCas9(BB)-2A-Puro vector (Addgene, Watertown, MA, #62988)

and described in our previous publication [45]. This strategy allows all three components (two

gRNAs and Cas9) express from the same plasmid to ensure the transfected cells expressing two

gRNAs simultaneously knocking out the targeted gene. The sequences of primers to make guide

RNAs were listed in Supplementary Table S5.

Knockout of lnc-FANCI-2 gene in CaSki cells, PCR screening and genotyping

Two gRNAs each under control by a U6 promoter in a pSpCas9(BB)-2A-Puro vector were

constructed as described [45] and used for transfection of CaSki cells. Plasmid pHBL25 which

contains gRNA 1 and 6 was used to delete the entire promoter region of lnc-FANCI-2 and

plasmid pHBL27 which contains gRNA 3 and 4 was used to delete only two YY1-binding motifs

spanning over an 87-bp region in the lnc-FANCI-2 promoter. CaSki cells transfected with the

gRNA-expressing vectors were then under 2 weeks of puromycin selection (0.1 µg/ml). The

puromycin-resistant CaSki cells were diluted to a final concentration of 0.5 cells per 100 µl,

plated 100 µl of diluted cells into each well of a 96-well plate, and grew under purymycin

selection. The colonies were inspected to ensure the clones were grown from one cell, and refed

or re-plated as needed in 3 weeks of expansion.

A direct PCR was performed for number of single cell clones as described [45]. Briefly, several

hundreds of cells expanded from a single cell clone were resuspended in 50 µl of phosphate-

buffered saline (PBS) and then frozen-thawed on dry ice for three times. Cell lysates were treated

with 4 µl of Qiagen protease (QIAGEN, Hilden, Germany, #1017782,) at 56°C for 10 min, and

the protease was inactivated at 95°C for 5 mins. The resulting product was directly used for PCR

screening. P1 + P5 were used for verification of promoter deletion and P3 + P4 for YY1 motifs

deletion. To ensure the selected single cell clones displaying homozygous knockout of lnc-

FANCI-2, total cell DNA was isolated using a QIAamp DNA Blood Minikit (QIAGEN, Hilden,

Germany, #51106) and then the homozygous deletion was screened and the cells with

homozygous KO were selected and confirmed by PCR. The primers used for genotyping are

listed in supplementary Table S5. At least two single cell clones either with lnc-FANCI-2

promoter deletion or with YY1 motif deletion were selected. Three single cell clones from the

428   cells transfected with an empty vector, pSpCas9(BB)-2A-Puro, were also selected and served as

429   controls for the genotyping screening.

Cell proliferation and viability assay

Cell proliferation and viability were determined by a CCK-8 assay (Dojindo Molecular

Technologies, Rockville, MD). For proliferation assay, 500 µl of cell suspension (50,000

cells/well) were dispensed into individual wells in a 24-well plate. Cells were then treated with

inhibitors (20 µM LY294002, or 10 µM U0126 respectively) after 24 h culture. At each time

point of 24 h, 48 h and 72 h, 50 µl of CCK-8 solution was added in three wells of each group and

incubated for 1 h in a cell culture incubator at 37°C. Cell proliferation was determined by

absorbance at 450 nm viable cells, and normal medium was used to subtract background.

Percentage of cell viability was calculated by the following formula: cell viability % = OD450 of

inhibitors treated sample / OD450 of untreated sample × 100%.

Colony formation, cell migration, and senescence analysis

CaSki Cells were seeded at 1×10^4, 5×10^3 or 1×10^3 cells/well in 6-well plates. After 24 h of

cell growth, the culture medium in each well was replaced by DMEM containing 1%

methylcellulose and 10% FBS. The plates were fixed with 3.7% formaldehyde and stained with

1% crystal violet in 2 weeks. Transwell cell migration assays were performed as previously

described [81]. Briefly, CaSki cells were re-suspended in serum free DMEM containing 0.1% BSA.

1×10^5 cells in 100 µl were then plated on top of the filter membrane in a transwell insert

(SARSTEDT, Nümbrecht, Germany, # 83.3932.800) and incubated for 10 min at 37 °C to allow

the cells to settle down, then 600 µl of complete media containing 10% FBS were added into the

bottom of the lower chamber in a 24-well plate. After 24 h in culture, the cells on the top of the

insert membrane were removed, and cells on the bottom were fixed, stained by 0.2% crystal

violet, and counted.

Cell senescence was determined by a Cellular Senescence Assay (MilliporeSigma, Burlington,

MA, # QIA117-1KIT). Briefly, CaSki cells were plated at 2×10^5 cells/well in 6-well plates and

cultured for 2 days to 50% confluence. Cells were washed once with 2 ml PBS and fixed with

0.5 ml of fixative solution at room temperature for 10 to 15 min. Then, cells were washed twice

with 2 ml PBS, and 1 ml staining solution was added to each well. Cells were incubated at 37°C

overnight without CO_2 and protected from the light. Blue stained cells were then counted under

light microscopy.

Nuclear and cytoplasmic fractionation

CaSki and SiHa cells were fractionated by using Nuclei EZ Prep Kit (Sigma-Aldrich, #NUC-

101) following the manufacturer's protocol. The details of the method could be found in our

previous publication [82]. Western blot analysis for nuclear protein SRSF3 and cytoplasmic protein

GAPDH was used to determine the fractionation efficiency. The fractionated cytoplasmic and

nuclear RNAs were used for detection of lnc-FANCI-2 RNA with human GAPDH RNA serving

as a control for RNA fractionation efficiency of the cytoplasmic RNAs.

RT-qPCR

Detection of lnc-FANCI-2 by RT-qPCR was performed as described [8]. Pre-designed primers for

lnc-FANCI-2 are listed in Table S5. Briefly, 2 µg total RNA was converted to cDNA using

Superscript First-stand Synthesis kit (Thermo Fisher Scientific, Waltham, MA, #11904018).

qPCR was performed using TaqMan gene expression Master Mix (Applied Biosystems,

Waltham, MA, #4369016) on a StepOne Plus Real-Time PCR system (Applied Biosystems).

TaqMan Gene Expression Assay (Applied Biosystems, #4331182) for GADPH

(Hs02758991_g1) was served as an internal control. Data were plotted as fold change over the

control group using the $2^{-\Delta\Delta Ct}$ method by which the data was normalized first to the values for

GAPDH and then to the median value for control samples. Data are presented as a bar graph with

mean ± SD for each group.

Northern blot

To validate lnc-FANCI-2 KO in CaSki cells, Northern blot was performed using poly(A)$^+$

mRNA isolated from 100 µg of total CaSki RNA with PolyATtract mRNA Isolation Systems

(Promega, #Z5310) or 10 µg of total CaSki RNA as described previously [8]. Briefly, RNA

samples were denatured in NorthernMax Formaldehyde loading dye (Thermo Fisher Scientific,

#AM8552) at 75°C for 15 min, then separated in an 1% formaldehyde-containing agarose gel

and transferred onto a GeneScreen Plus hybridization transfer membrane (Perkin Elmer,

Waltham, MA, #NEF987001PK). RNAs on the membrane were crosslinked by exposing to UV

light, and then prehybridized with PerfectHyb Plus hybridization buffer (Sigma-Aldrich, St.

Louis, MO, #H7003) for 2 h at 42°C. Specific oligos against lnc-FANCI-2 or GAPDH listed in

Table S5 were labeled with [γ-^{32}P]-ATP using T4 PNK (Thermo Fisher Scientific, Waltham,

MA, #18004–010) and added into hybridization buffer for overnight incubation at 42°C. The

membrane was then washed with 2× SSPE-0.5% SDS solution for 5 mins, followed by twice

washes with 0.5× SSPE-0.5% SDS each for 15 min at 42°C and then exposed to a

PhosphorImager screen.

RNA-seq and data analysis

Total RNA was from WT CaSki cells, ΔYY1-D5 and ΔPr-A9 cells were extracted using TRIzol

reagent (Thermo Fisher Scientific, Waltham, MA, #15596018). Total ribo-minus RNA-seq

libraries, four samples in each group, were prepared with TruSeq Stranded Total RNA Library

Kit and then subjected to pair-end sequencing using Illumina-HiSeq3000/4000 platform. The

samples had 123 to 140 million pass filter reads with more than 89% of bases above the quality

score of Q30. Reads of the samples were trimmed from adapters and low-quality bases using

Cutadapt before alignment with the reference genome (Human – hg38) and the annotated

transcripts using STAR. The average mapping rate of all samples was 95%. Unique alignment

was above 84%. There were 4.38 to 5.72% unmapped reads. The mapping statistics were

calculated using Picard software. Percent coding bases were between 33-39%. Percent UTR

bases were 25-31%, and mRNA bases were between 59-70% for all the samples. Library

complexity was measured in terms of unique fragments in the mapped reads using Picard's

MarkDuplicate utility. The samples had 71-76% non-duplicate reads. Raw data and the analyzed

RNA-seq data supporting the findings in this study have been deposited in the NCBI GEO

database (the accession#: GSE190904).

For RNA sequencing, sample reads were processed using the CCBR Pipeliner utility

(https://github.com/CCBR/Pipeliner). Briefly, reads were trimmed from adapters and low-quality

bases using Cutadapt (version 1.18) (https://bioweb.pasteur.fr/packages/pack@cutadapt@1.18)

before alignment to the custom reference genome described below. The transcripts were aligned

using STAR v2.5.2b in two-pass mode [83]. Expression levels were quantified using RSEM

(version 1.3.0) [84] with a custom gene annotation described below.

The custom reference genome allowing quantification of both viral and host expression used in

this alignment consisted of the human reference genome (hg38/Dec. 2013/GRCh38) FASTA

with a HPV16 FASTA sequence added as an additional pseudochromosome. The custom gene

annotation used for gene expression quantification consisted of a concatenation of the hg38

GENCODE annotation version 30 [85] and the HPV16 genome, with one other notable alteration.

The hg38 v30 gene annotation is incorrect at the location of the lnc-FANCI-2 locus in the hg38

genome at chr15:89378104-89398487, which affects quantification of this gene. To correct this

annotation and provide accurate quantification, therefore, we first performed BLAST against

hg38 using the sequences of the 14 known isoform transcripts of lnc-FANCI-2 (MT669800 for

lnc-FANCI-2a-PA2, MT669801 for lnc-FANCI-2a-PA1, MT669802 for lnc-FANCI-2b-PA2,

MT669803 for lnc-FANCI-2b-PA1, MT669804 for lnc-FANCI-2c-PA2, MT669805 for lnc-

FANCI-2c-PA1, MT669806 for lnc-FANCI-2d-PA2, MT669807 for lnc-FANCI-2d-PA1,

MT669808 for lnc-FANCI-2e-PA2, MT669809 for lnc-FANCI-2e-PA1, MT669810 for lnc-

FANCI-2f-PA2, MT669811 for lnc-FANCI-2f-PA1, MT669812 for lnc-FANCI-2g-PA2, and

MT669813 for lnc-FANCI-2g-PA1). The results were used to determine the precise exon start

and stop sites for each of the isoforms. These were then used to manually adjust the default hg38

v30 gene annotations at the locus indicated above to account for the correct structure of the lnc-

FANCI-2 locus and its 14 isoforms.

Downstream analysis and visualization were performed within the NIH Integrated Analysis

Portal (NIDAP) using R programs developed on the Foundry platform (Palantir Technologies).

Briefly, raw counts data produced by RSEM were imported into the NIDAP platform, genes

were filtered for low counts (<1 cpm) and the voom algorithm [86] from the Limma R package

(version 3.40.6) [87] was used for quantile normalization and calculation of differentially expressed

genes. Pre-ranked GSEA was performed using the Molecular Signatures Database version 6.2 [88,

89] and the fgsea package [90].

Human Soluble Receptor Array

To detect the changes of soluble receptors expressed and released from parental WT CaSki cells

and lnc-FANCI-2 KO cells, Human Soluble Receptor Array (R&D Systems, Minneapolis, MN, #

ARY012) was performed according to a manufacturer protocol. Briefly, 7×10^6 WT CaSki cells

and ΔPr-A9 cells were cultured in 15 ml of DMEM supplemented with 10% FBS in a T75 flask

for 24 h. Cell culture supernatants were then collected and centrifuged at 4,000 x g for 15 min to

remove cell debris. Cells were rinsed by 10 ml of PBS and solubilized in Lysis Buffer 17 at 4°C

for 30 min. The cell lysis was then centrifuged at 14,000 x g for 5 min to remove cell debris.

Protein concentration of cell lysis was measured by Micro BCA Protein Assay (PIERCE,

Waltham, MA, #23235). Next, two sets of N and C membranes were blocking by Buffer 8/1 in

4-Well Multi-dish for 1 h on a rocking platform. 500 μl of culture supernatant or 100 μg cell

lysis generated from the WT CaSki cells or ΔPr-A9 cells were diluted in 3 ml of Buffer 8/1, and

then applied to each membrane after three washes with 1X Wash Buffer. The membranes were

incubated overnight at 4 °C on a rocking platform and washed three times to remove unbound

materials followed by incubation with their specific cocktail of biotinylated detection antibodies

for 2 h at room temperature on a rocking platform. 2 ml of diluted Streptavidin-HRP was then

applied to each membrane for 30 min at room temperature on a rocking platform after three

times of wash. 1 ml of the prepared Chemi Reagent Mix was added onto the membranes after

three times of wash. The signal was then captured by ChemiDoc™ Touch Imaging System (Bio-

Rad). Quantification of the relative protein expression in ΔPr-A9 cells compared to the WT cells

was determined in Image Lab software (Bio-Rad). The average signal of duplicate spots

subtracting background value from negative control spots represented the levels of each protein.

The relative change in protein levels was then determined by comparing ΔPr-A9 cells to the WT

cells.

Antibodies and Immunoblot

Immunoblot analysis was performed for individual proteins by using the following antibodies:

Anti-MCAM (#17564-1-AP), anti-E2F1 (#12171-1-AP), anti-MAP4K4 (#55247-1-AP), and

anti-p53 (#10442-1-AP) antibodies were from ProteinTech (Rosemont, IL). Anti-tubulin

(#T5201) antibodies were from Sigma-Aldrich (St. Louis, MO). Anti-hnRNP C1/C2 (#Ab10294)

and anti-RAC3 (#ab129062) antibodies were from Abcam (Cambridge, United Kingdom). Anti-

CCND2 (Cyclin D2, #3741), anti-Akt (#9272), anti-phospho-Akt (#9271), anti-Erk1/2 (p44/42

MAPK, #4695), anti-phospho-Erk1/2 (#4370), and anti-IGFBP3 (D1U9C, #25864) were from

Cell Signaling Technology (Danvers, MA). Anti-VIM (#MA5-11883) was from Thermo Fisher

Scientific (Waltham, MA). Anti- PODXL2 (#AF1524), anti-ECM1 (#MAB39371), anti-TIMP2

(#AF971) and anti-ADAM8 (#AF1031) were from R&D Systems (Minneapolis, MN). Anti-

SRSF3 (#NBP2-76892) was from Novus Biologicals (Centennial, CO). Rabbit polyclonal anti-

HPV16 E7 (#GTX133411) was from GeneTex (Irvine, CA).

RAS GTPase Chemi ELISA

RAS GTPase activity in cell lysate of the WT CaSki or ΔPr-A9 cells was determined by RAS

GTPase Chemi ELISA Kit (Active Motif, Carlsbad, CA, # 52097) according to the manufacturer

protocol. Briefly, 2×10^7 cells were solubilized in 500 µl of Complete Lysis/Binding buffer at

4°C for 15 min. Protein concentration of cell extract was then measured by Micro BCA Protein

Assay (PIERCE, #23235) after cell debris was removed by centrifuged at 14,000 x g for 10 min.

2 µg of GST-Raf-RBD in 50 µl of Complete Lysis/Binding buffer was coated in each well of a

96-well plate by incubating for 1 h at 4°C with 100 rpm agitation. After three times of wash, 50

µg of extract in 50 µl Complete Lysis/Binding buffer was then added in each well. HeLa (EGF

treated) extract was served as a positive control. The plate was incubated for 1 h at room

temperature with 100 rpm agitation. The active RAS protein (GST-Raf-RBD binding) in each

well was then detected by incubating with primary RAS antibody and three times of wash, and

then, HRP secondary antibody and three times of wash. Chemiluminescence in each well was

read in luminometer by adding 50 µl room-temperature Chemiluminescent Working Solution.

RNA *in situ* hybridization (RNA-ISH)

Endogenous lnc-FANCI-2 in CaSki cells was examined by RNAScope Multiplex Fluorescent V2

Assay (Advanced Cell Diagnostics, Minneapolis, MN, #323100) as described previously [8]. A

custom-designed probe targeting to nt 359-1713 of GenBank MT669800.1 transcript for lnc-

FANCI-2a-PA2 was utilized. Dual staining of lnc-FANCI-2 and IGFBP RNA was performed

according to RNAscope Multplex Fluorescent v2 Manual part 2 (#323100), and the channel one

probe for IGFBP3 (#310351) and channel three probe for lnc-FANCI-2 (#509061-C3) were

applied.

For RNA-ISH combined IF staining of MCAM protein, 3 x 10^5 of CaSki cells were grown on a

glass coverslip in a 6-well plate for 24 h before transfection. Plasmid pHBL05 containing a long

isoform of lnc-FANCI-2a-PA2 (GenBank, MT669800.1) were transfected into cells with LipoD

293 DNA in vitro transfection reagent. Cultured Adherent Cell Sample Preparation for the

RNAscope Multplex Fluorescent v2 was performed according to a manufacturer protocol

(Advanced Cell Diagnostics, Minneapolis, MN, MK-50-010). Briefly, the cells were washed

with PBS and fixed by 10% neutral buffered formalin at room temperature for 30 min. The cells

were washed with PBS three times and dehydrated by 50% ethanol for 5 min, 70% ethanol for 5

min, 100% ethanol for 5min, and 100% ethanol for 10 min. The cells were then rehydrated by

70% ethanol for 2 min, 50% ethanol for 2min, and PBS for 10 min. The cells were then applied

to hydrogen peroxide at RT for 10 min and washed with distilled water for three times. The cells

were then applied to Protease III (1:15 dilution with PBS) at RT for 10 min. The cells were ready

for RNAscope Multplex Fluorescent v2 Manual part 2 (#323100). IF was performed after RNA-

ISH but before counterstaining with DAPI. The cells were washed with PBS, and then were

blocked with 2% bovine serum albumin (BSA) in PBS for 1 h at 37°C or overnight at 4°C. The

cells were incubated with the MCAM primary antibody (ProteinTech, Rosemont, IL, #17564-1-

AP, diluted 1:200 in 2% BSA blocking buffer) for 2 h at 37°C. An alexaFluor-conjugated

secondary antibody (1: 500, ThermoFisher Scientific) was diluted in a blocking solution and

incubated for 1 h at 37˚C. The slides were washed with PBS for three times. DAPI was used for

nuclei counterstaining before mounting in a Prolong Gold Antifade mounting medium (Thermo

Fisher Scientific, #P36934).

Confocal images were collected using a Zeiss LSM710 laser-scanning microscope

equipped with a 63 × Plan-Apochromat (N.A. 1.4) objective lens. Three dimensional

distributions of lnc-FANCI-2 were generated by Z stacks using ZEN 2.3 software (Zeiss).

Isolation of lnc-FANCI-2 <u>R</u>NA-<u>p</u>rotein <u>c</u>omplex by <u>R</u>NA <u>P</u>urification (IRPCRP) and LC-

MS/MS analysis

To identify the lnc-FANCI-2 RNA-associated proteins, we pulled down lnc-FANCI-2

RNA from WT CaSki cells using IRPCRP, a modified ChIRP protocol from the published

method [91], followed by mass spectrometry [57]. Briefly, a total of 30 lnc-FANCI-2 anti-sense oligo

probes (oLLY496-oLLY525), each with biotinTEG at 3' end, across entire lnc-FANCI-2 RNA

(GenBank: MT669800.1) were designed using the online probe designer

(singlemoleculefish.com). Two probe pools were used in IRPCRP assays: the probe pool 1 was

the mixed oligos in even numbers (oLLY496

/498/500/502/504/506/508/510/512/514/516/518/520/522/524) and the probe pool 2 mixed with

the oligos in odd numbers

(oLLY497/499/501/503/505/507/509/511/513/515/517/519/521/523/525) (Table S5). CaSki

cells were seeded and harvested at 24 h and the cell lysates from 200 million cells were used for

each IRPCRP reaction. Instead of formaldehyde crosslink in the standard protocol [91], UV cross-

linking (254 nm, energy at 480 mJ/cm^2) was applied in our study to minimize non-specific

binding. After UV cross-linking, the cells were lysed on ice in 1× radioimmunoprecipitation

assay (RIPA) buffer (Boston Bioproducts, Ashland, MA, #BP-115) (50 mM Tris base/Tris-HCl

[pH 7.4], 150 mM NaCl, 0.5% sodium deoxycholate, 0.1% sodium dodecyl sulfate [SDS], 1%

NP-40) supplemented with protease inhibitors (complete mini-EDTA-free protease inhibitor

cocktail, Millipore Sigma, Burlington, MA, #469315900) and RNase inhibitor (Thermo Fisher

Scientific, #AM2694) for 30 min, followed by a brief sonication. The obtained cell lysates were

spin for 10 min of centrifugation at 20,000 × g at 4°C and the collected cell lysate supernatants

were pre-absorbed with C-1 magnetic beads (Thermo Fisher Scientific, #65002) for 1 h at room

temperature before proceeding to oligo probe hybridization. Two different hybridization

protocols were used to pull down lnc-FANCI-2 RNA. One had the pre-absorbed cell lysate

incubated with two separate probe pools in hybridization buffer (750 mM NaCl, 0.1% SDS, 50

mM Tris-Cl pH 7.0, 1 mM EDTA, 15% formamide) at room temperature overnight and the pre-

washed C-1 magnetic beads was then added and incubated for additional 4 h; the other had two

separate oligo probe pools immobilized onto pre-washed C-1 magnetic beads first at room

temperature for 1 h and then mixed with the pre-absorbed cell lysates in hybridization buffer

overnight at room temperature, along with the negative control-beads only IRPCRP reaction

without oligo probes serving as a control in the modified protocol. After hybridization, the beads

were washed with 1x wash buffer (2x NaCl and sodium citrate, 0.1% SDS) for 5 times and

resuspend in 1 ml lysis buffer. 100 µl beads from each IRPCRP were set aside for RNA isolation

of lnc-FANCI-2 pulldown efficiency and the leftover beads were sent for mass spectrometry

analysis. The input and IRPCRP RNA were isolated using a MIRNeasy mini kit (Qiagen,

#217004) following the standard protocol after proteinase K (Millipore Sigma, #71049)

digestion. For mass spectrometry analysis, the IRPCRP beads were processed for trypsin/LysC

digestion before submitted to Thermo Scientific Orbitrap Exploris 240 Mass Spectrometer and a

Thermo Dionex UltiMate 3000 RSLCnano System for Proteinomics. Peptides from trypsin

digestion were loaded onto a peptide trap cartridge at a flow rate of 5 µl /min. The trapped

peptides were eluted onto a reversed-phase Easy-Spray Column PepMap RSLC, C18, 2 µM,

100A, 75 µm x 250 mm (Thermo Scientific) using a linear gradient of acetonitrile (3-36%) in

0.1% formic acid. The elution duration was 110 min at a flow rate of 0.3 µl/min. Eluted peptides

from the Easy-Spray column were ionized and sprayed into the mass spectrometer, using a Nano

Easy-Spray Ion Source (Thermo Fisher Scientific) under the following settings: spray voltage,

1.6 kV, Capillary temperature, 275 °C. Other settings were empirically determined. Raw data

files were searched against human protein sequences database using the Proteome Discoverer 2.4

software (Thermo, San Jose, CA) based on the SEQUEST algorithm. All the protein peptides

identified in the IRPCRP pulldowns were summarized in Table S4.

ACKNOWLEDGEMENTS

We thank Louise T. Chow of the University of Alabama at Birmingham for her critical reading

of the manuscript. We thank Craig Meyers of Penn State University Hershey Medical Center for

providing HPV16- and HPV18- infected raft cultures, Xing Xie and Yang Li of Zhejiang

University Women's Hospital of China for cervical tissue sections, and Johannes G. Schweizer

from Arbor Vita corporation for anti-HPV16 E6-specific antibodies. This study was supported

by the Intramural Research Program of the NIH, National Cancer Institute, Center for Cancer

Research.

DATA AVAILABILITY

NCBI GEO database Accession# GSE190904 for CaSki cell RNA-seq.

REFERENCES

1. Torre, L.A., Bray, F., Siegel, R.L., Ferlay, J., Lortet-Tieulent, J. & Jemal, A. Global cancer statistics, 2012. *CA Cancer J Clin* **65**, 87-108 (2015).
2. Bosch, F.X., Lorincz, A., Munoz, N., Meijer, C.J. & Shah, K.V. The causal relation between human papillomavirus and cervical cancer. *J Clin Pathol* **55**, 244-265 (2002).
3. Munoz, N. *et al.* Epidemiologic classification of human papillomavirus types associated with cervical cancer. *N Engl J Med* **348**, 518-527 (2003).
4. Walboomers, J.M. *et al.* Human papillomavirus is a necessary cause of invasive cervical cancer worldwide. *J Pathol* **189**, 12-19 (1999).
5. McCredie, M.R. *et al.* Natural history of cervical neoplasia and risk of invasive cancer in women with cervical intraepithelial neoplasia 3: a retrospective cohort study. *Lancet Oncol* **9**, 425-434 (2008).
6. Niraj, J., Farkkila, A. & D'Andrea, A.D. The Fanconi Anemia Pathway in Cancer. *Annu Rev Cancer Biol* **3**, 457-478 (2019).
7. Hoskins, E.E. *et al.* Fanconi anemia deficiency stimulates HPV-associated hyperplastic growth in organotypic epithelial raft culture. *Oncogene* **28**, 674-685 (2009).
8. Liu, H. *et al.* Oncogenic HPV promotes the expression of the long noncoding RNA lnc-FANCI-2 through E7 and YY1. *Proc Natl Acad Sci U S A* **118** (2021).
9. Santegoets, L.A. *et al.* Different DNA damage and cell cycle checkpoint control in low- and high-risk human papillomavirus infections of the vulva. *Int J Cancer* **130**, 2874-2885 (2012).

10.   Hoskins, E.E. *et al.* The fanconi anemia pathway limits human papillomavirus replication. *J Virol*
86, 8131-8138 (2012).
11.   Park, J.W. *et al.* Deficiencies in the Fanconi anemia DNA damage response pathway increase
sensitivity to HPV-associated head and neck cancer. *Cancer Res* **70**, 9959-9968 (2010).
12.   Park, J.W., Shin, M.K. & Lambert, P.F. High incidence of female reproductive tract cancers in FA-
deficient HPV16-transgenic mice correlates with E7's induction of DNA damage response, an
activity mediated by E7's inactivation of pocket proteins. *Oncogene* **33**, 3383-3391 (2014).
13.   Roman, A. & Munger, K. The papillomavirus E7 proteins. *Virology* **445**, 138-168 (2013).
14.   Vande Pol, S.B. & Klingelhutz, A.J. Papillomavirus E6 oncoproteins. *Virology* **445**, 115-137 (2013).
15.   Liu, X., Dakic, A., Zhang, Y., Dai, Y., Chen, R. & Schlegel, R. HPV E6 protein interacts physically and
functionally with the cellular telomerase complex. *Proc Natl Acad Sci U S A* **106**, 18780-18785
(2009).
16.   Klingelhutz, A.J., Foster, S.A. & McDougall, J.K. Telomerase activation by the E6 gene product of
human papillomavirus type 16. *Nature* **380**, 79-82 (1996).
17.   Storey, A., Pim, D., Murray, A., Osborn, K., Banks, L. & Crawford, L. Comparison of the in vitro
transforming activities of human papillomavirus types. *EMBO J* **7**, 1815-1820 (1988).
18.   Barbosa, M.S. & Schlegel, R. The E6 and E7 genes of HPV-18 are sufficient for inducing two-stage
in vitro transformation of human keratinocytes. *Oncogene* **4**, 1529-1532 (1989).
19.   Hawley-Nelson, P., Vousden, K.H., Hubbert, N.L., Lowy, D.R. & Schiller, J.T. HPV16 E6 and E7
proteins cooperate to immortalize human foreskin keratinocytes. *EMBO J* **8**, 3905-3910 (1989).
20.   Munger, K., Phelps, W.C., Bubb, V., Howley, P.M. & Schlegel, R. The E6 and E7 genes of the
human papillomavirus type 16 together are necessary and sufficient for transformation of
primary human keratinocytes. *J Virol* **63**, 4417-4421 (1989).
21.   Halbert, C.L., Demers, G.W. & Galloway, D.A. The E7 gene of human papillomavirus type 16 is
sufficient for immortalization of human epithelial cells. *J Virol* **65**, 473-478 (1991).
22.   Phelps, W.C., Yee, C.L., Munger, K. & Howley, P.M. The human papillomavirus type 16 E7 gene
encodes transactivation and transformation functions similar to those of adenovirus E1A. *Cell*
53, 539-547 (1988).
23.   DiPaolo, J.A., Woodworth, C.D., Popescu, N.C., Notario, V. & Doniger, J. Induction of human
cervical squamous cell carcinoma by sequential transfection with human papillomavirus 16 DNA
and viral Harvey ras. *Oncogene* **4**, 395-399 (1989).
24.   Schreiber, K. *et al.* Strong synergy between mutant ras and HPV16 E6/E7 in the development of
primary tumors. *Oncogene* **23**, 3972-3979 (2004).
25.   Narisawa-Saito, M. *et al.* An in vitro multistep carcinogenesis model for human cervical cancer.
Cancer Res **68**, 5699-5705 (2008).
26.   Song, S., Pitot, H.C. & Lambert, P.F. The human papillomavirus type 16 E6 gene alone is
sufficient to induce carcinomas in transgenic animals. *J Virol* **73**, 5887-5893 (1999).
27.   Herber, R., Liem, A., Pitot, H. & Lambert, P.F. Squamous epithelial hyperplasia and carcinoma in
mice transgenic for the human papillomavirus type 16 E7 oncogene. *J Virol* **70**, 1873-1881
(1996).
28.   Song, S., Liem, A., Miller, J.A. & Lambert, P.F. Human papillomavirus types 16 E6 and E7
contribute differently to carcinogenesis. *Virology* **267**, 141-150 (2000).
29.   Riley, R.R., Duensing, S., Brake, T., Munger, K., Lambert, P.F. & Arbeit, J.M. Dissection of human
papillomavirus E6 and E7 function in transgenic mouse models of cervical carcinogenesis. *Cancer*
Res **63**, 4862-4871 (2003).
30.   Brake, T. & Lambert, P.F. Estrogen contributes to the onset, persistence, and malignant
progression of cervical cancer in a human papillomavirus-transgenic mouse model. *Proc Natl*
Acad Sci U S A **102**, 2490-2495 (2005).

31.   Kato, S. *et al.* Activation of the estrogen receptor through phosphorylation by mitogen-activated
protein kinase. *Science* **270**, 1491-1494 (1995).

32.   Statello, L., Guo, C.J., Chen, L.L. & Huarte, M. Gene regulation by long non-coding RNAs and its
biological functions. *Nat Rev Mol Cell Biol* **22**, 96-118 (2021).

33.   Zhao, J., Sun, B.K., Erwin, J.A., Song, J.J. & Lee, J.T. Polycomb proteins targeted by a short repeat
RNA to the mouse X chromosome. *Science* **322**, 750-756 (2008).

34.   Yap, K.L. *et al.* Molecular interplay of the noncoding RNA ANRIL and methylated histone H3
lysine 27 by polycomb CBX7 in transcriptional silencing of INK4a. *Mol Cell* **38**, 662-674 (2010).

35.   Cho, S.W. *et al.* Promoter of lncRNA Gene PVT1 Is a Tumor-Suppressor DNA Boundary Element.
Cell **173**, 1398-1412 e1322 (2018).

36.   Xiang, J.F. *et al.* Human colorectal cancer-specific CCAT1-L lncRNA regulates long-range
chromatin interactions at the MYC locus. *Cell Res* **24**, 513-531 (2014).

37.   Tripathi, V. *et al.* The nuclear-retained noncoding RNA MALAT1 regulates alternative splicing by
modulating SR splicing factor phosphorylation. *Mol Cell* **39**, 925-938 (2010).

38.   Yap, K., Mukhina, S., Zhang, G., Tan, J.S.C., Ong, H.S. & Makeyev, E.V. A Short Tandem Repeat-
Enriched RNA Assembles a Nuclear Compartment to Control Alternative Splicing and Promote
Cell Survival. *Mol Cell* **72**, 525-540 e513 (2018).

39.   Nishizawa, M., Ikeya, Y., Okumura, T. & Kimura, T. Post-transcriptional inducible gene regulation
by natural antisense RNA. *Front Biosci (Landmark Ed)* **20**, 1-36 (2015).

40.   Lee, S. *et al.* Noncoding RNA NORAD Regulates Genomic Stability by Sequestering PUMILIO
Proteins. *Cell* **164**, 69-80 (2016).

41.   Poliseno, L., Salmena, L., Zhang, J., Carver, B., Haveman, W.J. & Pandolfi, P.P. A coding-
independent function of gene and pseudogene mRNAs regulates tumour biology. *Nature* **465**,
1033-1038 (2010).

42.   Grelet, S. *et al.* A regulated PNUTS mRNA to lncRNA splice switch mediates EMT and tumour
progression. *Nat Cell Biol* **19**, 1105-1115 (2017).

43.   Aznaourova, M. *et al.* Noncoding RNA MaIL1 is an integral component of the TLR4-TRIF pathway.
Proc Natl Acad Sci U S A **117**, 9042-9053 (2020).

44.   Guo, C.J. *et al.* Distinct Processing of lncRNAs Contributes to Non-conserved Functions in Stem
Cells. *Cell* **181**, 621-636 e622 (2020).

45.   BeltCappellino, A., Majerciak, V., Lobanov, A., Lack, J., Cam, M. & Zheng, Z.M. CRISPR/Cas9-
Mediated Knockout and In Situ Inversion of the ORF57 Gene from All Copies of the Kaposi's
Sarcoma-Associated Herpesvirus Genome in BCBL-1 Cells. *J Virol* **93** (2019).

46.   Schlomann, U. *et al.* The metalloprotease disintegrin ADAM8. Processing by autocatalysis is
required for proteolytic activity and cell adhesion. *J Biol Chem* **277**, 48210-48219 (2002).

47.   Liberzon, A., Birger, C., Thorvaldsdottir, H., Ghandi, M., Mesirov, J.P. & Tamayo, P. The
Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* **1**, 417-425
(2015).

48.   Downward, J. Targeting RAS signalling pathways in cancer therapy. *Nat Rev Cancer* **3**, 11-22
(2003).

49.   Simanshu, D.K., Nissley, D.V. & McCormick, F. RAS Proteins and Their Regulators in Human
Disease. *Cell* **170**, 17-33 (2017).

50.   Castellano, E. & Downward, J. RAS Interaction with PI3K: More Than Just Another Effector
Pathway. *Genes Cancer* **2**, 261-274 (2011).

51.   Asati, V., Mahapatra, D.K. & Bharti, S.K. PI3K/Akt/mTOR and Ras/Raf/MEK/ERK signaling
pathways inhibitors as anticancer agents: Structural and pharmacological perspectives. *Eur J*
Med Chem **109**, 314-341 (2016).

52. Cuesta, C., Arevalo-Alameda, C. & Castellano, E. The Importance of Being PI3K in the RAS
Signaling Network. *Genes (Basel)* **12** (2021).
53. Vanhaesebroeck, B., Perry, M.W.D., Brown, J.R., Andre, F. & Okkenhaug, K. PI3K inhibitors are
finally coming of age. *Nat Rev Drug Discov* **20**, 741-769 (2021).
54. Favata, M.F. *et al.* Identification of a novel inhibitor of mitogen-activated protein kinase kinase. *J*
Biol Chem **273**, 18623-18632 (1998).
55. Kebir, A. *et al.* CD146 short isoform increases the proangiogenic potential of endothelial
progenitor cells in vitro and in vivo. *Circ Res* **107**, 66-75 (2010).
56. Stalin, J. *et al.* Soluble CD146 boosts therapeutic effect of endothelial progenitors through
proteolytic processing of short CD146 isoform. *Cardiovasc Res* **111**, 240-251 (2016).
57. Chu, C. & Chang, H.Y. ChIRP-MS: RNA-Directed Proteomic Discovery. *Methods Mol Biol* **1861**, 37-
45 (2018).
58. Gao, X., Gao, C., Liu, G. & Hu, J. MAP4K4: an emerging therapeutic target in cancer. *Cell Biosci* **6**,
56 (2016).
59. Gao, X. *et al.* MAP4K4 is a novel MAPK/ERK pathway regulator required for lung
adenocarcinoma maintenance. *Mol Oncol* **11**, 628-639 (2017).
60. Gonzalez-Montero, J., Rojas, C.I. & Burotto, M. MAP4K4 and cancer: ready for the main stage?
Front Oncol **13**, 1162835 (2023).
61. Patterson, V. *et al.* Abrogation of MAP4K4 protein function causes congenital anomalies in
humans and zebrafish. *Sci Adv* **9**, eade0631 (2023).
62. Crook, T., Storey, A., Almond, N., Osborn, K. & Crawford, L. Human papillomavirus type 16
cooperates with activated ras and fos oncogenes in the hormone-dependent transformation of
primary mouse cells. *Proc Natl Acad Sci U S A* **85**, 8820-8824 (1988).
63. Matlashewski, G., Schneider, J., Banks, L., Jones, N., Murray, A. & Crawford, L. Human
papillomavirus type 16 DNA cooperates with activated ras in transforming primary cells. *EMBO J*
6, 1741-1746 (1987).
64. Greenhalgh, D.A. *et al.* Transgenic mice expressing targeted HPV-18 E6 and E7 oncogenes in the
epidermis develop verrucous lesions and spontaneous, rasHa-activated papillomas. *Cell Growth*
Differ **5**, 667-675 (1994).
65. Riou, G., Barrois, M., Sheng, Z.M., Duvillard, P. & Lhomme, C. Somatic deletions and mutations
of c-Ha-ras gene in human cervical cancers. *Oncogene* **3**, 329-333 (1988).
66. Eiben, G.L. *et al.* Establishment of an HLA-A*0201 human papillomavirus type 16 tumor model
to determine the efficacy of vaccination strategies in HLA-A*0201 transgenic mice. *Cancer Res*
62, 5792-5799 (2002).
67. Golijow, C.D., Mouron, S.A., Gomez, M.A. & Dulout, F.N. Differences in K-ras codon 12 mutation
frequency between "high-risk" and "low-risk" HPV-infected samples. *Gynecol Oncol* **75**, 108-112
(1999).
68. Prokopakis, P., Sourvinos, G., Koumantaki, Y., Koumantakis, E. & Spandidos, D.A. K-ras mutations
and HPV infection in cervicitis and intraepithelial neoplasias of the cervix. *Oncol Rep* **9**, 129-133
(2002).
69. Strickland, S.W. & Vande Pol, S. The Human Papillomavirus 16 E7 Oncoprotein Attenuates AKT
Signaling To Promote Internal Ribosome Entry Site-Dependent Translation and Expression of c-
MYC. *J Virol* **90**, 5611-5621 (2016).
70. Cararo-Lopes, E. *et al.* Autophagy buffers Ras-induced genotoxic stress enabling malignant
transformation in keratinocytes primed by human papillomavirus. *Cell Death Dis* **12**, 194 (2021).
71. Singh, A. *et al.* A gene expression signature associated with "K-Ras addiction" reveals regulators
of EMT and tumor cell survival. *Cancer Cell* **15**, 489-500 (2009).

72. Yang, J., Nie, J., Ma, X., Wei, Y., Peng, Y. & Wei, X. Targeting PI3K in cancer: mechanisms and
advances in clinical trials. *Mol Cancer* **18**, 26 (2019).

73. Saliani, M., Mirzaiebadizi, A., Javadmanesh, A., Siavoshi, A. & Ahmadian, M.R. KRAS-related long
noncoding RNAs in human cancers. *Cancer Gene Ther* (2021).

74. Su, Y.C., Han, J., Xu, S., Cobb, M. & Skolnik, E.Y. NIK is a new Ste20-related kinase that binds NCK
and MEKK1 and activates the SAPK/JNK cascade via a conserved regulatory domain. *EMBO J* **16**,
1279-1290 (1997).

75. Collins, C.S. *et al.* A small interfering RNA screen for modulators of tumor cell motility identifies
MAP4K4 as a promigratory kinase. *Proc Natl Acad Sci U S A* **103**, 3775-3780 (2006).

76. Joshkon, A. *et al.* Role of CD146 (MCAM) in Physiological and Pathological Angiogenesis-
Contribution of New Antibodies for Therapy. *Biomedicines* **8** (2020).

77. Anfosso, F. *et al.* Activation of human endothelial cells via S-endo-1 antigen (CD146) stimulates
the tyrosine phosphorylation of focal adhesion kinase p125(FAK). *J Biol Chem* **273**, 26852-26856
(1998).

78. Dye, D.E. *et al.* hShroom1 links a membrane bound protein to the actin cytoskeleton. *Cell Mol*
Life Sci **66**, 681-696 (2009).

79. Wang, Z., Xu, Q., Zhang, N., Du, X., Xu, G. & Yan, X. CD146, from a melanoma cell adhesion
molecule to a signaling receptor. *Signal Transduct Target Ther* **5**, 148 (2020).

80. Bardin, N., Frances, V., Combes, V., Sampol, J. & Dignat-George, F. CD146: biosynthesis and
production of a soluble form in human cultured endothelial cells. *FEBS Lett* **421**, 12-14 (1998).

81. Justus, C.R., Leffler, N., Ruiz-Echevarria, M. & Yang, L.V. In vitro cell migration and invasion
assays. *J Vis Exp* (2014).

82. Yu, L. & Zheng, Z.M. Human Papillomavirus Type 16 Circular RNA Is Barely Detectable for the
Claimed Biological Activity. *mBio* **13**, e0359421 (2022).

83. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

84. Li, B. & Dewey, C.N. RSEM: accurate transcript quantification from RNA-Seq data with or without
a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

85. Harrow, J. *et al.* GENCODE: the reference human genome annotation for The ENCODE Project.
Genome Res **22**, 1760-1774 (2012).

86. Law, C.W., Chen, Y., Shi, W. & Smyth, G.K. voom: Precision weights unlock linear model analysis
tools for RNA-seq read counts. *Genome Biol* **15**, R29 (2014).

87. Smyth, G.K. Linear models and empirical bayes methods for assessing differential expression in
microarray experiments. *Stat Appl Genet Mol Biol* **3**, Article3 (2004).

88. Liberzon, A., Subramanian, A., Pinchback, R., Thorvaldsdottir, H., Tamayo, P. & Mesirov, J.P.
Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**, 1739-1740 (2011).

89. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550
(2005).

90. Korotkevich, G., Sukhov, V., Budin, N., Shpak, B., Artyomov, M.N. & Sergushichev, A. Fast gene
set enrichment analysis. *bioRxiv* (2021).

91. Chu, C., Quinn, J. & Chang, H.Y. Chromatin isolation by RNA purification (ChIRP). *J Vis Exp* (2012).

 Figure legends

Fig. 1. Knockout (KO) of lnc-FANCI-2 in CaSki cells affects cell proliferation, colony formation

and migration. (A) Diagram and KO strategies of lnc-FANCI-2 gene. On the figure top is the

lnc-FANCI-2 gene structure and its alternative transcription start sites (TSS) and polyadenylation

sites (pA). TSS2 or pA2 with a heavier arrow are predominately used for lnc-FANCI-2

expression. The lower figure part shows KO strategies. Red slashes represent gRNA-targeted

sites to create genomic DNA deletions by CRISPR-Cas9 technology. Deletion of YY1-binding

motifs (ΔYY1 motifs or ΔYY1) led to a deletion of 86-bp DNA fragment containing two YY1-

binding motifs. Deletion of lnc-FANCI-2 promoter (ΔPromoter or ΔPr) led to delete a ~3.3-kb

promoter region. Primers (P1-P4) used for deletion screening were shown as black arrows. (B)

PCR screening of single cell clones with homozygous lnc-FANCI-2 KO, with indicated primer

sets diagramed (A). Single cell clones selected from the cells transfected with an empty vector

served as control cells. (C) Evaluation of lnc-FANCI-2 KO efficiency in the selected single cell

clones (B) by RT-qPCR. (D) Northern blot validation of lnc-FANCI-2 KO efficiency from the

individual single cell clones on polyA$^+$ RNA (enriched from 100 μg of total cell RNA, Lanes 1-

4) or 10 μg of total cell RNA (lanes 5-9). Total cell RNA (2 μg) of HEK293T cells with ectopic

expression of two isoforms (short, S or long, L) of lnc-FANCI-2 cDNA served as a control.

Oligo probe P5 (A) labeled with ^{32}P was used for hybridization. GAPDH RNA served as a

loading control and hybridized with a GAPDH-specific oligo probe. (E-F) RNA in situ

hybridization (RNA-ISH) validation of lnc-FANCI-2 KO efficiency by RNAscope technology.

Two single cell clones were examined with red color for lnc-FANCI-2 and blue color for the

nucleus (E). Bar graphs show the copy number of lnc-FANCI-2 per cell in the parental (WT),

ΔYY1-D5 or ΔPr-A9 cells (each averaged from 200 cells) (F). (G-I) lnc-FANCI-2 expression is

required for cell proliferation, colony formation, and cell invasion of CaSki cells. Representative

pictures are shown from three separate experiments of cell proliferation by CCK8 assay (G), cell

colony formation under 1% methylcellulose solution (H), and cell migration by transwell assay

(I) of WT, ΔYY1-D5 or ΔPr-A9 cells. Cell colonies were stained with crystal violet as described

in the procedure. Numbers of the invaded cells crossed the transwell membrane were quantified

and shown as bar graphs (I). *, P< 0.05; **, P<0.01; ***, P<0.001 by two tailed Student *t* test.

Fig. 2. KO of lnc-FANCI-2 in CaSki cells affects expression of cellular soluble receptors. (A)

Dot blots shows differentially expressed soluble receptors in ΔPr-A9 cells when compared to

those in the WT CaSki cells. A Proteome Profiler Human sReceptor Array was used to examine

105 cellular soluble receptors. Label in red, upregulated soluble receptors; label in blue,

downregulated soluble receptors. (B) Quantification of differentially expressed and/or released

soluble receptors (A) in ΔPr-A9 cells when compared to those in the WT CaSki cells. Error bar

represents standard deviation of replicates in each proteome array. The dash line indicates the

threshold of fold change-1 with a score above or below the threshold being determined as

differentially expressed and/or released soluble receptors. (C) Validation of several differentially

expressed soluble receptors by immunoblot analysis using specific antibodies. Tubulin served as

an internal loading control. * non-specific band in MCAM immunoblot.

Fig. 3. Transcriptomic effect of lnc-FANCI-2 KO in CaSki cells by RNA-seq analysis. (A)

RNA-seq reads-coverage maps by IGV showing the expression levels of lnc-FANCI-2 from two

KO cell clones, ΔPr-A9 and ΔYY1-D5, to the WT CaSki cells. one representative coverage

profile of four in each type of cells is shown by IGV. Red slashes represent the deleted genomic

region, with validated fourteen lnc-FANCI-2 RNA isoforms shown below [8]. (B) Similarity in

heatmap comparison among ΔPr-A9, ΔYY1-D5 and WT cells was generated using Limma-

normalized counts from each sample by Pearson complete linkage method. (C) Volcano plot

visualization of differentially expressed genes (DEGs) in ΔPr-A9 and ΔYY1-D5. The genes with

the most significant change in the increased (red) or decreased (blue) expression are indicated.

FC, fold change. (D) Venn diagram depicting the overlapped DEGs between ΔPr-A9 and ΔYY1-

D5 cells over the WT CaSki cells. (E) Heatmap shows overlapped DEGs with FPKM equal or

more than 7 in ΔPr-A9 and ΔYY1-D5 cells when compared with the WT CaSki cells. (F)

Validation of selective 12 upregulated or downregulated DEGs shown in the heatmap by RT-

PCR with gene-specific primers in the presence (+) or absence (-) of reverse transcriptase (RT).

GAPDH served as an internal RNA control. Relative expression level of each gene was

calculated based on band density after normalizing to GAPDH RNA band, with the expression

level in the WT CaSki cells setting as 1. (G) Validation of the upregulated expression of IGFBP3

in ΔPr-A9 cells over the WT CaSki cells by immunoblot analysis.

Fig. 4. Pathway analyses of DEGs identified by RNA-seq. (A) Top 3 upregulated and top 4

downregulated pathways in ΔPr-A9 and ΔYY1-D5 cells over the WT CaSki cells. Data were

generated by Gene set enrichment analysis (GSEA) performed with Hallmark gene sets. NES

stands for normalized enrichment score. (B-E) Each GSEA Enrichment plot shows the

enrichment score and gene hits enriched in ΔPr-A9 cells using Hallmark gene sets,

KRAS_SIGNALING_UP (B), EPITHELIAL_MESENCHYMAL_TRANSITION (C),

INTERFERON_GAMMA_ RESPONSE (D) or INTERFERON_ALPHA_RESPONSE (E). padj,

adjusted p value; Zero crossed 7342, the middle number of total genes in the GSEA at ranking.

The heatmaps below the enrichment plots (B and C) visualize the genes enriched in respective

pathways of KRAS_SIGNALING_UP (B) and

EPITHELIAL_MESENCHYMAL_TRANSITION (C) in ΔPr-A9 cells when compared to the

WT CaSki cells.

Fig. 5. CaSki cells with lnc-FANCI-2 KO exhibit activation of RAS signaling pathway. (A)

CaSki cells with lnc-FANCI-2 KO (top, ΔPr-A9 cells) or knockdown (KD, lower) by lnc-

FANCI-2-specific siRNAs display increased RAS GTPase activity when compared to the WT

CaSki cells (top) or the WT cells treated with a nonspecific control siRNA (siNS). The relative

RAS GTPase activity was measured by a RAS GTPase Chemi ELISA assay. *, P< 0.05; ***,

P<0.001 by two tailed Student *t* test. (B and C) Selective validation of increased expression of

RAS signaling-related downstream genes in lnc-FANCI-2 KO ΔPr-A9 cells (B) or lnc-FANCI-2

KD CaSki cells (C) over the WT CaSki cells. Relative expression of the indicated genes in ΔPr-

A9 cells in comparison to the WT cells (B) or in the WT CaSki cells treated for 48 h and 96 h

with a nonspecific control siRNA (siNS) or a siRNA specifically targeting the lnc-FANCI-2

exon 3 (C) were examined by immunoblot analyses using corresponding antibodies as indicated.

Tubulin served as an internal control for each blot. The level of p-Akt or p-Erk was calculated by

normalizing to total Akt or Erk protein and the other proteins by normalizing to tubulin. The

protein level in the WT cells or WT cells treated with siNS set as 1. The KD efficiency of lnc-

FANCI-2 RNA (C, top bar graphs) was examined by RT-qPCR. (D) The effect of blocking RAS

signaling on the expression of MCAM and VIM using PI3K inhibitor LY294002 (20 μM) and

MEK inhibitor U0126 (10 μM). The protein at each time point was examined by immunoblot

analysis with a corresponding antibody. The level of MCAM or VIM was calculated after

normalizing to tubulin. The protein level in ΔPr-A9 cells without the inhibitor was set as 100%.

*, nonspecific protein band. (E) The time dependent cell viability of the WT CaSki and ΔPr-A9

cells in the presence of 20 μM LY294002 or 10 μM U0126. Data was obtained in each time point

after normalizing to the cells treated with DMSO. The mean $\pm$ SD at each data point was

calculated from 6 samples combined from two independent experiments.

Fig. 6. Negative correlation of lnc-FANCI-2 and MCAM predicts cervical cancer patient

prognosis. (A) Validation of differential expression of MCAM RNA in Δpr-A9 cells by RT-PCR

in the presence (+) or absence (-) of reverse transcriptase (RT). One pair of primers with one

primer at the exons 11 and the other at exon 13 of MCAM RNA (NM_006500) were used.

GAPDH served as a loading control. (B) The major isofrom lnc-FANCI-2a-PA2 (GenBank,

MT669800.1) repressed the expression of MCAM. Δpr-A9 cells were transfected with lnc-

FANCI-2a-PA2 cDNA plasmid. lnc-FANCI-2 RNA in red was detected by RNAscope ISH and

MCAM protein in green was detected by IF with an anti-MCAM antibody. (C) Expression levels

of lnc-FANCI-2 RNA and MCAM protein in three neighboring cells were measured by signal

intensity of a line crossing over the stained cells (B, white line arrow). (D) Calculation of the

expression levels of MCAM in lnc-FANCI-2 positive cells (n=35) and lnc-FANCI-2 negative

cells (n=54) by fluorescent intensity. (E) Subcellular MCAM distributions in the nucleus (N) and

cytoplasm (C) of WT CaSki cells and Δpr-A9 by immunoblot analysis. Fractionation efficiency

and sample loading were controlled by cytoplasmic (represented by tubulin) and nuclear

(represented by hnRNP C1/C2) proteins. (F). No detectable MCAM in the culture supernatant by

immunoblot. (G-H) Correlation and survival analysis of lnc-FANCI-2 and MCAM expression

with cervical squamous cell carcinoma (CESC) cases from the cancer genome atlas (TCGA)

datasets by GEPIA web server (http://gepia.cancer-pku.cn/). The negative correlation at r=-0.321

(with p-value=1.02e-08) of Inc-FANCI-2 with MCAM in cervical cancer patients was obtained

from the RNA-seq data from the TCGA for CESC tumor type downloaded from the TCGA data

portal (https://portal.gdc.cancer.gov/). Only primary solid tumor samples (n=304 after exclusion

of 2 metastatic samples and 3 normal samples) were subjected to analysis, with the data showing

as a Scatter plot (G). Kaplan-Meier plot with log rank test p-value at 5.74e-05 (H) shows MCAM

a biomarker for poor prognosis of cervical cancer survival with a lower quartile group cutoff.

RNA-seq and survival data are derived from the TCGA CESC cancer patients.

Fig. 7. lnc-FANCI-2 interacts with host factors to regulate RAS signaling pathway. (A) The lnc-

FANCI-2-associated proteins in the WT CaSki cells were identified by isolation of RNA-protein

complexes using RNA purification (IRPCRP)-Mass spectrometry technology. (B) lnc-FANCI-2

RNA in the IRPCRP-1 and IRPCRP-2 pulldowns had the pooled antisense biotinylated oligos

(pool 1 with oligos in even numbers and pool 2 with oligos in odd numbers) immobilized to

avidin-beads first before mixed with cell lysates, while the IRPCRP-3 and IRPCRP-4 pulldowns

had the oligos pool 1 and 2 separately mixed with cell lysates first before addition to the avidin-

beads for the RNA pull-downs. RT-PCR in the absence (-) or presence (+) of reverse

transcriptase (RT) was carried out using the RNA isolated from the individual IRPCRP

experiments using a primer pair of oHBL5 and oHBL12 (Table S5) specific for lnc-FANCI-2

RNA detection. Beads only (no oligos) IRPCRP experiments served as a negative control. Total

RNA from the WT CaSki cells after sonication was used as an input control. Arrow indicates the

detected lnc-FANCI-2 RNA. (C) Proteins associated with lnc-FANCI-2 RNA identified from

lnc-FANCI-2 IRPCRP pulldowns. A Venn diagram circle in orange shows the proteins pulled

down specifically from lnc-FANCI-2 IRPCRP reactions 1-4 (PSM>=2 from two separate

reactions), and in blue the proteins non-specifically pulled down by beads only (oligo-free) in the

IRPCRP reactions (PSM >=1). Top 10 proteins binding lnc-FANCI-2 are shown on the left in

the order by the number of identified PSM. (D) Expression of p-Akt and p-Erk from CaSki WT

cells 48 h after siRNA-mediated MAP4K4 knockdown (KD) was immunoblotted by the

corresponding antibodies. GAPDH served as a sample loading control. Fold change of the

indicated proteins in the cells with MAP4K4 KD over the cells treated by non-targeting siRNA

(siRNA-NS) was calculated after normalizing to GAPDH. (E) A proposed model illustrates how

lnc-FANCI-2-protein complexes regulate the RAS signaling pathway to control Akt/ErK

phosphorylation and expression of host genes.

1037

A

ΔPr-A9 VS WT

pathway	NES
KRAS_SIGNALING_UP	1.858008
EPITHELIAL_MESENCHYMAL_TRANSITION	1.771138
INFLAMMATORY_RESPONSE	1.490227
INTERFERON_GAMMA_RESPONSE	-1.47303
ADIPOGENESIS	-1.50073
INTERFERON_ALPHA_RESPONSE	-1.62585
BILE_ACID_METABOLISM	-1.64176

ΔYY1-D5 VS WT

pathway	NES
TNFA_SIGNALING_VIA_NFKB	1.911117
KRAS_SIGNALING_UP	1.746758
EPITHELIAL_MESENCHYMAL_TRANSITION	1.711204
ESTROGEN_RESPONSE_LATE	-1.63355
INTERFERON_GAMMA_RESPONSE	-1.69812
ESTROGEN_RESPONSE_EARLY	-1.71013
INTERFERON_ALPHA_RESPONSE	-2.22382

B KRAS_SIGNALING_UP
padj=0.0110, NES=1.86

C EPITHELIAL_MESENCHYMAL_TRANSITION
padj=0.0106, NES=1.77

D INTERFERON_GAMMA_RESPONSE
padj=0.045, NES=−1.47

E INTERFERON_ALPHA_RESPONSE
padj=0.040, NES=−1.63

Fig. 5

Fig. S1. Characterization of lnc-FANCI-2 KO on CaSki cell growth, viral oncoprotein expression, and cell senescence. (A) The characteristic cell growth behavior and morphology was imaged at 24 h after spreading. (B) Effect of lnc-FANCI-2 KO in CaSki cells on the expression of HPV16 E6 and E7 and their downstream targets. Total cell extracts from parental CaSki cells and ΔPr-A9 cells were examined by immunoblotting with corresponding antibodies. Tubulin served as a protein loading control. The relative protein levels of E6, E7, p53, and E2F1 were calculated after normalizing to tubulin. (C) KO of lnc-FANCI-2 expression enhances cell senescence in b-gal analysis.

Fig S2. GSEA Enrichment plot shows enrichment score and gene hits enriched in ΔYY1-D5 cells using Hallmark gene sets, (A) KRAS_SIGNALING_UP. (B) EPITHELIAL_MESENCHYMAL_TRANSITION. (C) INTERFERON_ALPHA_RESPONSE. (D) INTERFERON_GAMMA_RESPONSE. The heatmap on the right side of Enrichment plot in each panel visualize the genes with differentiated expression enriched in each pathway in ΔYY1-D5 cells.

Fig. S3. Pathway map of RAS Initiative with highlighted differentially expressed genes (DEGs) between A9 cells vs wild type cells. DEGs are mainly distributed and clustered in two main branches of RAS pathway: MAPK signaling branch and PI3K/AKT branches. DEGs were derived from RNA-seq data at cutoff of adjusted p-value <=0.01 and fold change>1.5 for both up- and down-regulated genes (indicated by red and blue arrows respectively) between A9 cells vs WT CaSki cells using three common analysis methods (DESeq2, edgeR, and limma-voom) and in-house BRB analysis pipeline from NCI. Highlighted genes in the pathway map are derived as DEGs in at least three of the analysis methods, although in most cases, these DEGs behaved consistently across all 4 methods. MAPK signaling branch is highlighted in transparent green box and PI3K/AKT branch is highlighted in transparent yellow box within the pathway map of RAS signaling pathway, which was collectively collated from community inputs that were organized and stimulated by RAS Initiative, and maintained as a common knowledge basis at URL below: https://www.cancer.gov/research/key-initiatives/ras/ras-central/blog/2015/ras-pathway-v2, which was also described in recent review (Figure 1 in Nissley and McCormick, 2022).

Reference (where the Ras pathway map has been described):
Nissley, D. V. and McCormick, F. (2022). RAS at 40: update from the RAS Initiative. *Cancer Discov.* https://doi.org/10.1158/2159-8290.CD-21-1554

Fig S4. (A) Selective validation of increased expression of RAS signaling-related downstream genes in ΔPr-A9 and ΔYY1-D5 cells. (B) Selective expression of lnc-FANCI-2 RNA in a subset of cervical cancer cell lines as determined by RT-qPCR. Cervical cancer cell lines: SiHa (HPV16+), CaSki (HPV16+), HeLa (HPV18+), C33A (no HPV), C4II (HPV18+), and W12 subclones 20861 (HPV16+, integrated) and 20863 (HPV16+, episomal); other HPV negative cell lines: HCT116 (colorectal cancer cells), BCBL-1 (body cavity B lymphoma cells), HEK293 (Ad5 E1/E2-immortalized human kidney cells), HaCaT (spontaneously immortalized human epidermal cells). (C) HeLa cells express no lnc-FANCI-2 when compared with C33A cells as determined by Northern blot (C). (D) Subcellular determination of lnc-FANCI-2 as being predominantly cytoplasmic in CaSki but nuclear in SiHa cells. Cytoplasmic and nuclear fractionation efficiency was determined by Western blot analyses of nuclear protein SRSF3 (serine- and arginine-rich splicing factor 3) and cytoplasmic GAPDH protein. Total fractionated cytoplasmic and nuclear RNAs from CaSki and SiHa cells were quantified for lnc-FANCI-2 by RT-qPCR, with cytoplasmic GAPDH RNA serving as internal controls for RNA fractionation efficiency. (E) Subcellular lnc-FANCI-2 (red) localization in SiHa cells was examined by RNAscope RNA ISH analysis. Nuclei were stained with DAPI (blue). Scale bars: 25 µm in the left and 10 µm in the zoom. (F) KD of lnc-FANCI-2 expression in SiHa cells enhanced phosphorylation of Akt and Erk1/2 but did not in HeLa cells expressing no lnc-FANCI-2 (B, C).

A

HFK HFK-HPV16 HFK-HPV18

B

Cervical cancer tissue-adjacent normal Cervical cancer tissue-tumor area 1 Cervical cancer tissue-tumor area 2

Fig. S5. Increased expression of lnc-FANCI-2 in HPV 16- and HPV18-infected raft culture tissues and cervical cancer tissues. (A) HFK-derived raft cultures without (HFK) or with HPV16 or HPV18 infections (HFK-HPV16 or HFK-HPV18) was examined at day 10 for lnc-FANCI-2 RNA by RNAscope RNA-ISH analysis. (B) lnc-FANCI-2 expression in HPV+ cervical cancer tissues was examined by RNAscope RNA-ISH analysis. Nuclei were stained with DAPI (blue). Scale bars: 25 μm in the original figures and 10 μm in the zoomed insets.

Fig. S6. MCAM expression and lnc-FANCI-2. (A-B) RNA-seq reads-coverage of MCAM and IGFBP3 by IGV illustrates the increased expression of MCAM and IGFBP3 in lnc-FANCI-2 KO cells ΔPr-A9 and/or ΔYY1-D5 when compared to the parental WT CaSki cells. (C-D) Transient rescue expression of lnc-FANCI2 in Δpr-A9 cells inhibits expression of IGFBP3. Δpr-A9 cells were transfected with a lnc-FANCI-2a-PA2 cDNA plasmid. lnc-FANCI-2 in red and IGFBP3 RNA in green were detected by RNA-ISH at 24h post transfection using each specific antisense RNA probe and imaged by confocal microscopy (C). Expression levels of lnc-FANCI-2 RNA and IGFBP3 RNA in four neighboring cells (D) were measured by signal intensity of a line crossing over the stained cells in C (white line arrow).

A

lnc-FANCI-2:CESC:log rank test p-values vs gene exp gradient with days, 2followup perm No:1000

B

MCAM:CESC:log rank test p-values vs gene exp gradient with days, 2followup perm No:1000

Fig. S7. Expression levels of lnc-FANCI-2 and MCAM are significantly associated with TCGA CESC cancer patients' survival outcome. The expression of lnc-FANCI-2 (A) and of MCAM (B) along with survival of the same group of 304 cervical cancer patients were analyzed by our in-house GradientScanSurv pipeline as described (Yi et al 2018, PLoS One 13(12):e0207590). The **open grey circles** in each plot are ranked expression levels from high (left) to low (right) of lnc-FANCI-2 (A) and MCAM (B) from all patient samples as indicated by y-axis at the right side of the plot (in RSEM normalized value of RNAseq data). At each breaking point denoted by patient number (x-axis), the log rank test (y-axis at the left side of the plot) was performed between the higher expression group at the left side of the cut-point vs the lower expression group at the right side of the cut-point and the indicated log rank p-values (**blue dots**) are shown as at each cut-point at x-axis. The **horizontal red line** is the p=0.05 cutoff line for the log rank test p-values. If a log rank p-value (**blue dot**) at a cut-point is below the cutoff (p<0.05), there will be a **vertical green line** shown at the corresponding cut-point, indicative of a significant log rank test p-value, and also a **brown** diamond plotted either at the bottom part of the **vertical green line** (A) if higher expression of lnc-FANCI-2 led to less severe outcome (dying slower than the lower expressors), or at the top part of the **vertical green line** (B) if higher expression of MCAM led to more severe outcome (dying faster than the lower expressors).

The original GoodCount was defined as the number of significant log rank tests (the default significance level was set as 0.05) across all possible cut-points for the original dataset. We employed the bootstrap approach to permutate the data for 1000 times in this case, and for each time, a similar procedure was performed like the original dataset, and a permutated GoodCount was obtained for each permutated dataset. The GoodCount p-value was derived as the proportion of the times that the GoodCounts of permutated datasets were no less than that of the original dataset (see Yi et al 2018, PLoS One 13(12):e0207590) and shown at the left top corner of each plot, 0.014 for lnc-FANCI-2 in panel A and 0.004 for MCAM in panel B. A significant GoodCount p-value (**blue dots** in both panels A and B) is indicative of significant association of the expression levels with the corresponding survival outcomes of the analyzed patient samples. Derived p-values from univariant expression-based cox regression models (coxph p-value: derived by original gene expression values; or coxph p-value By Rank: derived by corresponding ranks of gene expression values) were shown at the top left corner of each plot. The cut-point-specific FDR (**green dots**) and FDR2 (**red dots**) were shown at the top right corner of each plot. FDR for each cut-point was defined as the portion of cases of permutated datasets that have log rank test p-values no higher than that of original dataset. FDR2 for each cut-point was defined as the portion of cases of permutated datasets that have log rank test p-values no higher than default setting of significance as 0.05.