The role of NMDA receptors in memory and prediction in cultured neural networks

Martina Lamberti[1,*], Michel J.A.M. van Putten[1], Sarah Marzen[2], Joost le Feber[1,*],

1 Department of Clinical Neurophysiology, University of Twente, Enschede, PO Box 217 7500AE, The Netherlands
2 W. M. Keck Science Department, Pitzer, Scripps and Claremont McKenna College, Claremont, 91711, USA

* Martina Lamberti m.lamberti@utwente.nl or Joost le Feber j.lefeber@utwente.nl

Abstract

Memory has been extensively studied at the behavioural as well as the cellular level. Spike timing dependent plasticity (STDP) is associated with N-methyl-D-aspartate (NMDA) receptor activation and is widely accepted to be essential for long-term memory. However, experimental evidence remains sparse, probably due to the required complex combination of cellular and functional readouts. Recent work showed that in-vitro cortical networks memorize and predict inputs. The initial dependency of prediction on short-term memory decreased during the formation of long-term memory traces. Here, we stimulated in-vitro networks to investigate memory and prediction under control conditions, or using the NMDA antagonist APV. APV did not significantly reduce network excitability, but did impede long-term memory trace formation. In APV-treated cultures short-term memory of stimuli persisted, and they were still able to predict. In contrast to control cultures, prediction remained fully dependent on short-term memory. This confirms that NMDA receptor activation is essential for the formation of long-term memory traces and supports the notion that, as control cultures learn to memorize the stimulus, long-term memory starts to contribute to their predictive capability.

Keywords:

NMDA receptors, Synaptic plasticity, Memory and prediction, In-vitro neural networks, Multi electrodes array

Introduction

Memory is a crucial cognitive function for daily life [1,2] that has been the subject of extensive investigation at various levels, ranging from cognitive psychological to neurobiological approaches. Cognitive psychology has mainly focused on abstract explanations of memory [3–5], and discriminates between short-term memory, acting on time scales of seconds, and long-term memory, on time scales of minutes to days [6–8]. New experiences are first stored in the hippocampus in a process called synaptic consolidation [9,10]. Then, memories are transferred to the neocortex for long-term storage by repeated replay in the hippocampus, particularly during slow-wave sleep (system consolidation) [10,11]. On the other hand, neurobiology has revealed detailed

insights into neuronal functioning and synaptic plasticity that are likely to underlie memory [12]. Bridging the gap between cognitive and neurobiological studies has been challenging, at least in part because experimental paradigms usually do not facilitate assessment of parameters that cover the entire range from the molecular or cellular level to cognitive functioning.

It is widely accepted that plasticity of neuronal connectivity is crucial for memory. Connection strengths between neurons can be determined in patch clamp experiments [13], but this approach reveals only one, or at the best a few, connection strengths at a time. In past decades, micro electrode arrays (MEAs) have been used to record activity from networks of cultured neurons [14], and recorded activity patterns can be used to infer functional connectivity and study memory [15–17].Connectivity and activity in neuronal networks mutually affect each other. The finding that input deprived networks develop quasi stable activity patterns [18,19] and connectivity [20] suggests that activity and connectivity are in equilibrium [21]. External stimuli induce new activity patterns, and disturb the equilibrium between activity and connectivity [22,23]. After a while the network reaches a new equilibrium with spontaneous activity patterns that include the stimulus response, which is interpreted as the formation of long-term memory of the input. Consequently, repeated exposure to this stimulus induces no further connectivity changes [22,23]. Recent work showed that cultured networks are able to form memory traces when repeatedly exposed to a stimulus during a few hours [22], and that this ability depends on slow-wave sleep-like conditions [23]. These preparations facilitate measurement and manipulation of cellular and synaptic properties, which provides the opportunity to bridge part of the gap between neurobiological and cognitive studies.

In addition to memory, prediction is also crucial to successfully conclude every-day actions. Prediction can be defined as the ability to reduce the uncertainty on upcoming external inputs, and is strongly related to memory [24–26]. In principle there can be no prediction without memory [25]. Most evidence to support the notion that neural networks are able to predict comes from theoretical work [1,2,24], and a few pioneering experimental studies using retinal [26–30] or hippocampal preparations [31–33].

Recent work showed that also random networks of dissociated cortical neurons are able to predict, suggesting that prediction is independent of specific network architecture [34]. In that study, prediction was shown to initially depend on short-term memory of given stimuli, which typically lasted several hundreds of milliseconds. During multiple hours of repeated focal stimulation, long-term memory traces were formed and prediction became less dependent on short-term memory. This suggests that focal stimulation-induced long-term memory traces also play a role in prediction. It remains unclear which mechanisms underlie prediction and to what extent long-term memory is involved.

Several studies have attributed the formation of long-term memory traces to spike timing dependent plasticity (STDP). [6,35–42]. STDP governs changes in synaptic weights based on the timing of pre- and postsynaptic action potentials, and may lead to either long-term potentiation (LTP) or depression (LTD) of synaptic strengths [43–46]. Calcium influx mediated by NMDA receptor activation plays an important role in STDP [36,43,45,47], and has been shown to affect the density of AMPA receptors in the synaptic region [36,47,48]. Depending on the amplitude and kinetics of the calcium influx, synapses can be either potentiated or depressed [36,47,49]. The strong association between memory and prediction suggests that STDP and NMDA receptor activation may also play a role in prediction.

In the current work, we present an in-vitro model to investigate memory and prediction. In particular we aimed to determine the involvement of NMDA receptor activation in memory and prediction. We applied focal electrical stimulation to

networks of rat cortical neurons plated on MEAs, with and without pharmacological blockade of NMDA receptors, and investigated whether networks were able to i) form long-term memory traces, and ii) predict future stimuli. In addition, we determined to what extent prediction depended on (short-term or long-term) memory.

Materials and methods

Culture preparation

Cortical cells, obtained from newborn rats were dissociated by trypsin treatment and trituration, and then 60 µl droplets of cell suspension (about $50,000$ cells) were plated on multi electrode arrays (MEAs; Multi Channel Systems, Reutlingen, Germany), precoated with poly ethylene imine (PEI). The glass substrate contained 60 titanium nitride electrodes (diameter 30 µm and pitch 200 µm), on which a Plexiglas ring (diameter 20 mm) was glued to create a culture well (Figure 1A). The well was filled with 1 ml of Neurobasal commercial medium (Neurobasal-A medium -gluc -pyr, Thermo Fisher), supplemented with B27, 1.24 g/L glucose, Pen/strep/glutamine, 100 mg/L vitamin C and 10 ng/L NGF. MEAs were stored in an incubator, under standard conditions of 36°C , high humidity, and 5% CO2 in air. The culture medium was refreshed twice a week by removing 300 µl and adding 400 µl of fresh medium, thus compensating for evaporation. All cultures were grown for at least 3 weeks before experiments started, to allow for culture maturation [20, 50, 51]. Mature cultures contained excitatory and inhibitory neurons [52], and astrocytes (ratio astrocytes neurons: 4:1) [53] (see Figure S3 in Supplementary Information). Before experiments, culture chambers were firmly sealed with watertight but O_2 and CO_2 permeable foil (MCS; ALA scientific). Then, MEAs were placed in a measurement setup outside the incubator. In this setup, standard conditions of 36°C, high humidity, and 5% CO_2, were maintained. Every recording began after a 15 minute accommodation period. At the end of each experiment, cultures were returned to the incubator. All surgical and experimental procedures were approved by the Dutch committee on animal use (Centrale Commissie Dierproeven; AVD110002016802), and complied with Dutch and European laws and guidelines. Results are presented in compliance with the ARRIVE guidelines.

Recording set-up

To record activity, we placed MEAs into a setup outside the incubator, consisting of a MC1060BC preamplifier and FA60s filter amplifier (both MultiChannelSystems GmbH, Reut-lingen, Germany). Signals from the network were recorded by 59 electrodes using a custom-made Lab-View program, at a sampling frequency of 16 kHz per electrode. Acquired analogue signals were band-pass filtered (2^{nd} order Butterworth $0.1 – 6$ kHz) before sampling [17, 34]. Action potentials were detected whenever recorded voltages exceeded a threshold, set at 5.5 times the estimated root- mean-square noise level (ranging $3 – 5$ µV). During recordings this noise estimation was continuously updated for each electrode. For each detected event the setup stores a time stamp, the electrode number, and the wave shape (6 ms). For off-line artifact detection and removal we used a wave shape based algorithm adapted from [54].

Focal stimulation was applied by electrical stimulation through one of the electrodes, using biphasic rectangular current pulses of 200 µs per phase [23]. After probing all electrodes at various amplitudes ($16 − 24$ µA), one electrode was selected for stimulation, using the lowest amplitude that induced a network response after at least 50% of all stimuli. Amplitudes were low enough to avoid electrolysis.

Experimental Design

We used 20 mM 2-amino-5-phosphonovaleric acid (APV) to block NMDA receptors. This concentration clearly affected firing patterns and stimulation induced connectivity changes, but had no significant effect on network excitability (see Supplementary Information). Addition of APV was done under a laminar flow hood to maintain sterility. We performed experiments under control conditions (CTRL cultures) or using an NMDA antagonist (APV cultures).

Memory

We acquired 1 h of spontaneous activity at the beginning of the experiment ($Baseline$). After APV administration we recorded a second hour of spontaneous activity ($Baseline_{APV}$). Then both CTRL and APV cultures underwent focal stimulation applied to two different electrodes (A and B). For each electrode we applied 5 stimulation periods of ten minutes each at a constant frequency of 0.2 Hz, alternated by 1 h spontaneous activity recordings. First, electrode A was used, then electrode B, and then again electrode A, leading to a total of 15 stimulation periods of 10 minutes. The total duration of the experimental protocol was 17 h and 30 min (Figure 1B).

Prediction

To study prediction we adapted the protocol developed in earlier work [34]. In short, we first acquired 1 h of spontaneous activity under control conditions. After APV administration we recorded another hour of spontaneous activity. Then, cultures were subjected to 20 h of focal electrical stimulation (Figure 1B). For these experiments we used inter stimulus intervals (ISIs) drawn independently and identically from a density distribution designed to produce long-range temporal correlations, which were read from a pre-generated list (see Figure 4C). The long and intensive stimulation protocol adopted for the prediction experiments might be too stressful for the cultures to be applied twice. We therefore compared current results under NMDA blockade to control experiments from a previous study [34].

Data analysis

Memory trace formation

Functional connectivity was inferred from spontaneous activity recordings using conditional firing probability (CFP) [20]. In short, CFP calculates the probability that electrode j spikes at time $t = \tau$, given that electrode i spiked at $t = 0$. The maximum of this probability curve $M_{i,j}$ was interpreted as the strength of the connection from electrode i to j (see Supplementary Information for details). This analysis was performed on chunks of data that contained 2^{13} action potentials. Typically, an hour of spontaneous activity could be divided into multiple chunks. This yielded a 60 x 60 connectivity matrix M for each chunk, that contained the strengths of connections between all possible pairs of electrodes. To assess the formation of memory traces, we quantified connectivity changes induced by each stimulation period by computation of Euclidean distances (EDs) between connectivity matrices [17, 23] (equation S2 in Supplementary Information). First we calculated EDs between all chunks within baseline, which were then averaged to obtain a measure of spontaneous connectivity fluctuations: ED_{baseline}. Then we computed EDs between the last chunk of baseline and each chunk contained in the (5) spontaneous periods that followed stimulation of electrode A. These distances are referred to as ED_0. This procedure was repeated for electrode B, but now the last chunks before the first stimulation at electrode B was used

Fig 1. Example of MEA, experimental protocols and recorded activity. A) shows an example of a 60 electrodes single well MEA used in this study. B) shows the protocols for memory experiments (top) and prediction experiments (bottom). C) shows an example of recorded spontaneous activity under control conditions. Each tick represents a recorded spike at the indicated electrode. Bottom trace shows summed activity (array wide firing rate; AWFR). D) represents an example of spontaneous activity recorded after APV administration. C) and D) show that APV administration reduced the spontaneous firing rates, but firing patterns still contained synchronous activity (networks bursts)

as baseline. Similarly, after return to electrode A, the last chunk before the first (second round) stimulation at electrode A was used as baseline (see Figure 2). In the computation we considered only the nonzero elements of the M matrices. These analyses were applied to both controls and APV experimental data.

Fig 2. Calculation of Euclidean distances (ED). Shown is a visual explanation on how Euclidean distances were calculated in regards to the memory experiments. Black vertical lines indicates the division in blocks applied to calculate CFP and get the connectivity matrices M. Red dashed arrow lines indicates ED0 calculation between different connectivity matrices and their corresponding baseline. Black circles shows that we took in consideration the final average ED_0 per each spontaneous period recorded.

Network excitability

Network excitability, defined as the average network response to a spike in one neuron, was calculated with SPRs as developed in [55]. This allows estimation of the average response at electrode j to a single spike at electrode i under widely varying dynamic regimes. The computation of SPRs requires deconvolving the auto-correlation of electrode i from $CFP_{i,j}$. Also in this case spontaneous recordings were subdivided in blocks of 2^{13} recorded spikes. SPR strengths of all pairs of active electrodes were calculated for each period of spontaneous activity in both control and APV groups, and averaged to obtain a measure of network excitability.

Quantification of prediction

We first verified that networks remained responsive to stimulation throughout experiments (see Supplementary Information). We used mutual information to quantify prediction [26, 34]. Mutual information estimates how much one signal reduces the uncertainty of another one. Then we first binarized the recorded neural activity (X) in bins of $\Delta t = 100$ ms by setting $X_i = 1$ if electrode i recorded activity in that time bin, and $X_i = 0$ if not. The same binarization was applied to the stimulation signal (S). After binarizing the activity X and the stimulation signal S, for each hour, we calculated MI between S and X, with X shifted in time by $n \times \Delta t$, with $-10 < n < 20$. Negative shifts reveal memory, as quantified by MI_{past}:

$$MI_{\mathrm{past}}(S; X) = MI(S_{\mathrm{t}}; X_{\mathrm{t}+\Delta \mathrm{t}})) \ \forall \ -1000 \leq \Delta t \leq 0 \text{ ms}, \tag{1}$$

where S_{t} and $X_{\mathrm{t}+\Delta \mathrm{t}}$ represent the unshifted stimulation signal and the time-shifted binarized activity. Activity was shifted forward up until a maximum of 2 seconds. Positive shifts are used to quantify prediction by computing MI_{future}:

$$MI_{\mathrm{future}}(S; X) = MI(S_{\mathrm{t}}; X_{\mathrm{t}+\Delta \mathrm{t}})) \ \forall \ 100 \leq \Delta t \leq 2000 \text{ ms}. \tag{2}$$

Consequently from Eqs. 1 and 2 MI_{past} and MI_{future} are functions of the shift Δt. We applied this analysis to the APV prediction experiments and compared our results to control experiments obtained in earlier work [34]. In APV experiments, stimulus responses significantly decreased after 13 hours (see Supplementary Information, Fig S2). Therefore, we took into account only the first 10h of stimulation (instead of all 20h) for APV as well as control cultures.

Relationship between prediction and short-term memory

To study possible effects of APV on the relationship between short-term memory and prediction we analyzed the area under the curve of MI_{future} ($\Sigma MI_{\mathrm{future}}$) and MI_{past} ($\Sigma MI_{\mathrm{future}}$) for each analyzed hour, for each experiment (see Eq. S3 and S4 Supplementary Information) [34]. Then we plotted $\Sigma MI_{\mathrm{future}}$ against $\Sigma MI_{\mathrm{past}}$, and fitted a linear equation to determine the slope and offset of the relation, and the correlation coefficient. Finally, we investigated whether and how slope and offset changed during the considered 10 hours of stimulation.

Efficiency of predictions

We used the predictive information bottleneck concept as shown in Lamberti et al. [34] to determine whether APV affected the efficiency of prediction. This allows us to determine the minimum information about the past (I_{mem}) necessary to make predictions on the future (I_{pred}). The entire past and future were replaced by the time

since the last stimulus (past; S^+), and the time to the next stimulus (future; S^-) [34, 56]:

$$I_{\mathrm{mem}} = I[S^+; X], \ I_{\mathrm{pred}} = I[S^-; X]. \tag{3}$$

Prediction as quantified in experimental data under APV or control conditions was then compared to a theoretical optimum ($R(D)$). The theoretical optimum was obtained calculating the minimum memory required to enable a certain prediction. This resulted in a line that divides between achievable and unachievable combinations of information about the past (I_{mem}) and the future (I_{pred}).

$$R(D) = \min_{p(x|s^+):I[X;S^-]\geq D} I[X; S^+]. \tag{4}$$

Here D corresponds to the required predictive power, $R(D)$ is the memory necessary to reach that predictive power, $p(x|s^+)$ is the conditional probability of neural activity given time since the last stimulus, x is a realization of the activity X, s^+ is a realization of S^+ [34]. For both controls and APV treated cultures we computed I_{mem} and I_{pred} considering the first 10 h of stimulation only.

Statistical Analysis

All analyses were performed using SPSS statistics for Windows (IBM, Inc., Chicago, IL) or Matlab (The Mathworks, Inc., Natick, MA, USA). Normality of data distribution was verified using Shapiro-Wilk tests. Effects of APV concentrations was testes with one sample t-test, in case of normality, or one sample Wilcoxon test otherwise. Significance of connectivity changes was assessed using repeated measurements ANOVA, in case of normality, or repeated measurement Friedman test otherwise. Assessment of possible effects of APV on network excitability was performed with two way repeated measurement ANOVA, with condition and time as independent variables. In both memory and prediction experiments significance of temporal differences in effectiveness of stimulation were analyzed by one way ANOVA (if normally distributed) or Kruskal-Wallis (if non-normal). In the end t-statistics measure was applied to check trends of slope and offset of the relationship between memory and prediction. P-values < 0.05 were considered to indicate significance.

Results

Long-term memory

In total 12 cultures were used, first under control conditions (CTRL experiments), and then under NMDA blockade (APV experiments). All 12 control recordings were included for analysis, and 7 of the APV experiments. In the other 5 cultures APV administration reduced network activity to such low levels that CFP analysis could not be done to estimate functional connectivity.

Network excitability

Network excitability was quantified by the average magnitude of SPRs during each period of spontaneous activity. There was no significant difference in network excitability between APV treated cultures and controls (see Figure 3B; $p = 0.11$), and network excitability in APV treated cultures did not significantly change during experiments ($p = 0.13$). In the control group network excitability slightly decreased with time ($p = 0.045$).

Memory trace formation

To study memory trace formation we first computed baseline functional connectivity in the networks and then quantified connectivity changes induced by 10 min periods of low frequency stimulation at one of the electrodes (Eq. S2 Supplementary Information). Figure 3A shows that in the control group the first sets of stimuli at electrode A and the first sets of stimuli at electrode B induced significant connectivity changes (respectively $p = 0.002$ and $p = 0.003$). Interestingly, second sets of stimuli to electrode A did not induce further significant connectivity changes in the network (repeated measurement Friedman test: $p = 0.08$). In the APV group, stimulation at either electrode did not lead to significant connectivity changes (A first stimulation set $p = 0.12$, B stimulation $p = 0.88$, A second stimulation set $p = 0.61$).

Fig 3. Memory formation and time course of network excitability. A) shows stimulation-induced connectivity changes for controls (orange) and APV treated (green) cultures. First series of stimulation at electrode A induced connectivity changes in controls ($p = 0.002$) but not in APV experiments ($p = 0.12$). Stimulation at electrode B yielded similar effects (control $p = 0.003$, APV $p = 0.88$). Return to electrode A did not induce connectivity changes in either group (control $p = 0.08$, APV $p = 0.61$). B) shows that network excitability was not affected by APV treatment ($p = 0.11$). In addition time was not affecting either group (control $p = 0.13$, APV $p = 0.045$). In both panels the red arrow indicates the addition of APV. Shaded areas show SEM, and indicate differences between experiments.

Quantification of prediction

We computed MI_{past} (Eq 1) and MI_{future} (Eq 2) to quantify short term memory and prediction. Figures 4A and B show MI_{past} and MI_{future} for control experiments (N=8) and APV experiments (N=10). Data from the control group has been used in an earlier study [34]. Both MI_{past} curves show a clear peak that corresponds to induced stimulus responses, which lasted up to ≈ 300 ms (Figure 4A). Both MI_{future} curves show an initial plateau until $\Delta t = 1000$ ms, followed by a peak between $\Delta t = 1000$ ms and $\Delta t = 1100$ ms. These correspond to the most probable inter stimulus interval (see ISI distribution figure 4C). These peaks were followed by a decrease in MI_{future} for $1100 < \Delta t < 2000$ ms, reflecting the probability distribution of ISIs . The narrow shaded areas, that indicate SEM and represent differences between different hours of analysis across experiments, show that curves hardly changed during the 10 analyzed hours.

Fig 4. MI curves obtained in CTRL (orange) and APV experiments (green). A) MI_{past} shows a peak at $\Delta t = -200 - 0$ ms corresponding to recorded stimulus responses. B) MI_{future} shows a peak at $\Delta t = 1100$ ms for CTRL experiments and at $\Delta t = 1000$ ms for APV experiments. Both follow the distribution of interstimulus intervals (ISIs) used and shown in (C). Shaded areas represents SEM.

Relationship between prediction and memory

Figure 5 shows how prediction, as quantified by $\Sigma MI_{\text{future}}$ (Eq. S3 Supplementary Information), depended on short-term memory, as assessed by ΣMI_{past} (Eq. S4 Supplementary Information). Prediction was strongly correlated with short-term memory in CTRL ($R = 0.8 \pm 0.07$) as well as APV experiments ($R = 0.87 \pm 0.09$). Under control conditions the slope of this relationship decreased with time ($p = 0.02$) and the offset increased ($p = 0.004$). In APV experiments slope and offset did not significantly change in the considered 10h of stimulation ($p = 0.56$ and $p = 0.79$). See Figures 5C and D.

Efficiency of predictions

To study how efficiently APV-treated cultures predict compared to controls, we computed I_{mem} and I_{pred} (Eq. 3) for each culture, using the first 10 h of stimulation. Under control conditions cultures reached a predictive power (I_{pred}) that was on average $62.7 \pm 7\%$ of the theoretical maximum ($R(D)$), (Figure 5F). When treated with APV cultures reached $69.6 \pm 9\%$ (Figure 5E). APV treated cultures were as efficient as control ones ($p = 0.1$).

Discussion

The current study confirms that *in-vitro* networks of dissociated cortical neurons form memory traces when focally stimulated as shown in earlier work [22, 23]. Blockade of NMDA receptors impeded the formation of memory traces, demonstrating the pivotal role of NMDA receptor activation in the formation of memory traces. Networks under NMDA blockade were still able to predict upcoming stimuli. However, whereas prediction became less dependent on short-term memory with the formation of long-term memory traces in control cultures, this dependency did not decrease under NMDA blockade.

NMDA receptor activation plays an important role in STDP. The current finding that NMDA receptor blockade hampered long-term memory formation in cortical networks is in line with the notion that STDP is critically involved in this process. However, blocking a subset of excitatory receptors may also reduce network excitability, which has been reported to hamper memory formation [23]. We therefore verified that network excitability did not decrease significantly (Figure 3), which suggests that other mechanisms than reduced network excitability lead to impeded memory trace formation. Although NMDA block has been shown to affect the dynamics of network bursts [57], it

Fig 5. Relationship between prediction ($\Sigma MI_{\text{future}}$) and short term memory (ΣMI_{past}), and efficiency of prediction. A,B) examples (hour 1, 6 and 10) of the relationship between $\Sigma MI_{\text{future}}$ and ΣMI_{past} and fitted trend lines in APV-treated cultures (A) and controls (B). C) time course of slope and offset of fitted lines in (A). In APV experiments slope and offset did not significantly change with time ($p = 0.82$ and $p = 0.97$). D) Time course of slope and offset of fitted lines in (B). In control experiments slope significantly decreased ($p = 0.02$) and offset significantly increased ($p = 0.004$) with time. E) and F) Efficiency of prediction in APV (E) and CTRL (F) cultures. I_{pred} and I_{mem} were calculated using Eq 3, dashed lines indicate the theoretical optimum as derived in Eq. 4. This shows us the minimum memory necessary to enable a certain prediction.

did not impede network responses to electrical stimulation or the occurrence of spontaneous network bursts, which are the most plausible driving forces to reach a connectivity <=> activity equilibrium. Taken together, these observations are in agreement with a crucial role of STDP in memory trace formation [35, 45, 58, 59].

Prediction strongly depended on the presence of clearly recognizable stimulus responses [34]. These stimulus responses are often interpreted as short-term memory of a stimulus, which has been suggested to depend mainly on sustained spiking in recurrent networks, creating attractor states [60]. Blocking NMDA receptors resulted in increasing stimulus responses during the first 5 hours after APV administration, followed by a decrease towards baseline values during the next hours (Figure S2). This suggests increasing excitability during the first 5 hours, which may reflect homeostatic mechanisms aiming to compensate for the blockade of NMDA receptors. On average, the NMDA antagonist did not reduce network excitability throughout the 10h analysis period. Networks remained responsive to focal electrical stimulation and they were still able to predict (Figure 4). This is in agreement with the notion that prediction, at least initially, depends on short-term memory, which is not blocked by the NMDA antagonist. The slightly higher values of MI_{future} and the slightly higher prediction efficiency (Figure 4B and Figure 5E,F) might be due to the reduction of spontaneous activity and network bursts in APV-treated cultures, which yielded more recognizable stimulus

responses in the recorded patterns (Figure 1C and D). Whereas prediction initially seemed to fully depend on short-term memory, this dependency decreased with time in focally stimulated networks under control conditions [34]. This type of stimulation induces long term memory traces in the network, suggesting that long-term memory traces are also involved in prediction under control conditions. This confirms earlier theoretical work on the role of long-term memory in prediction [61]. Interestingly, even though APV-treated networks were still able to predict, short-term memory dependency of prediction did not decrease with time (Figure 5D). Persistent short-term memory dependency was also observed in globally stimulated networks, where no long-term memory traces were formed [17,34]. This further supports the putative role of long-term memory in prediction.

The current study had a few limitations. The definition of memory differs from most other studies. This had the advantage that we were able to observe the formation of memory traces independently from recall. The mechanism of memory recall in our model is not clear yet, but the findings that stimulus responses stabilize during memory experiments, and start to appear in spontaneous firing patterns [22] support the concept of pattern completion [62,63]. Further understanding of recall, preferably a quantitative measure of memory recall, could further confirm memory trace formation. It is not well-known how long the NMDA antagonist APV remains functional, and we could not directly estimate the half life of APV in the culture medium. However, indirect measures suggest that APV remained functional throughout experiments. No long-term memory traces were formed, even > 10 h after APV administration. Network excitability could not be measured in the prediction experimental protocol due to ongoing stimulation. However, the period of analysis in this protocol was limited to 10 h, and it seems unlikely that the effects of APV wore off more rapidly in this protocol than in the memory protocol. It would be interesting to see whether network excitability was affected by APV under continuous stimulation, as used in prediction experiments, but the method used to estimate network connectivity [55] requires longer periods of spontaneous activity between stimuli, containing a sufficient number of action potentials for statistical analysis. Controls and APV experiments were not performed on the same cultures, which may have introduced a bias related to possible culture differences.

In conclusion, We developed an in-vitro model that allows us to investigate cellular mechanisms that underlie memory and prediction. Blocking NMDA receptors in this model impedes long-term memory formation. Short-term memory of stimuli persisted under NMDA block, and networks were still able to predict. However, in contrast to control conditions, the dependency on short-term memory persisted throughout time. This further sustains the hypothesis that long-term memory becomes involved in prediction after extended exposure to a focal stimulus.

Data availability

Part of the data (prediction control experiments) are available on Dryad (doi:10.5061/dryad.18931zd2t). All other data will be made available on Dryad upon acceptance.

Acknowledgments

The authors would like to thank Gerco Hassink and Marloes Levers for the technical assistance, and Stephanie Palmer and Chris Hillar for valuable comments. This study was supported by the US Air Force Office for Scientific Research, Grant Number FA9550-19-1-0411.

Author contributions statement

J.lF, S.M and M.L. conceived the study design, M.L. performed the experiments, M.L. J.lF. and analysed the data with the contribution of S.M., M.L. J. lF. developed the analysis scripts with a contribution from S.M., M.L. M.vP. J.lF. and S.M. wrote the manuscript draft. All authors reviewed the manuscript.

Competing interests

All authors declare no competing interests.

References

1. Bubic A, Von Cramon DY, Schubotz RI. Prediction, cognition and the brain. Frontiers in human neuroscience. 2010; p. 25.

2. Pitkow X, Liu S, Angelaki DE, DeAngelis GC, Pouget A. How can single sensory neurons predict behavior? Neuron. 2015;87(2):411–423.

3. Baddeley A. Cognitive psychology and human memory. Trends in neurosciences. 1988;11(4):176–181.

4. Roediger HL. Memory metaphors in cognitive psychology. Memory & Cognition. 1980;8:231–246.

5. Squire LR. Memory and brain systems: 1969–2009. Journal of Neuroscience. 2009;29(41):12711–12716.

6. Kukushkin NV, Carew TJ. Memory takes time. Neuron. 2017;95(2):259–279.

7. Kelley P, Whatson T. Making long-term memories in minutes: a spaced learning pattern from memory research in education. Frontiers in human neuroscience. 2013;7:589.

8. Tetzlaff C, Kolodziejski C, Markelic I, Wörgötter F. Time scales of memory, learning, and plasticity. Biological cybernetics. 2012;106:715–726.

9. Squire LR, Cohen NJ, Nadel L. The medial temporal region and memory consolidation: A new hypothesis. Memory consolidation: Psychobiology of cognition. 1984; p. 185–210.

10. Born J, Wilhelm I. System consolidation of memory during sleep. Psychological research. 2012;76:192–203.

11. Frankland PW, Bontempi B. The organization of recent and remote memories. Nature reviews neuroscience. 2005;6(2):119–130.

12. Kandel ER, Dudai Y, Mayford MR. The molecular and systems biology of memory. Cell. 2014;157(1):163–186.

13. Linders LE, Supiot LF, Du W, D'Angelo R, Adan RA, Riga D, et al. Studying synaptic connectivity and strength with optogenetics and patch-clamp electrophysiology. International Journal of Molecular Sciences. 2022;23(19):11612.

14. Cerina M, Piastra MC, Frega M. The potential of in vitro neuronal networks cultured on Micro Electrode Arrays for biomedical research. Progress in Biomedical Engineering. 2023;.

15. Morin FO, Takamura Y, Tamiya E. Investigating neuronal activity with planar microelectrode arrays: achievements and new perspectives. Journal of bioscience and bioengineering. 2005;100(2):131–143.

16. Marom S, Shahaf G. Development, learning and memory in large random networks of cortical neurons: lessons beyond anatomy. Quarterly reviews of biophysics. 2002;35(1):63–87.

17. Lamberti M, Hess M, Dias I, van Putten M, le Feber J, Marzen S. Maximum entropy models provide functional connectivity estimates in neural networks. Scientific reports. 2022;12(1):9656.

18. Stegenga J, Le Feber J, Marani E, Rutten WL. Analysis of cultured neuronal networks using intraburst firing characteristics. IEEE transactions on biomedical engineering. 2008;55(4):1382–1390.

19. Van Pelt J, Wolters PS, Corner MA, Rutten WL, Ramakers GJ. Long-term characterization of firing dynamics of spontaneous bursts in cultured neural networks. IEEE Transactions on Biomedical Engineering. 2004;51(11):2051–2062.

20. le Feber J, Rutten W, Stegenga J, Wolters P, Ramakers G, Van Pelt J. Conditional firing probabilities in cultured neuronal networks: a stable underlying structure in widely varying spontaneous activity patterns. Journal of neural engineering. 2007;4(2):54.

21. Le Feber J, Stegenga J, Rutten WL. The effect of slow electrical stimuli to achieve learning in cultured networks of rat cortical neurons. PloS one. 2010;5(1):e8871.

22. le Feber J, Witteveen T, van Veenendaal TM, Dijkstra J. Repeated stimulation of cultured networks of rat cortical neurons induces parallel memory traces. Learning & memory. 2015;22(12):594–603.

23. Dias I, Levers MR, Lamberti M, Hassink GC, Van Wezel R, Le Feber J. Consolidation of memory traces in cultured cortical networks requires low cholinergic tone, synchronized activity and high network excitability. Journal of neural engineering. 2021;18(4):046051.

24. Barron HC, Auksztulewicz R, Friston K. Prediction and memory: A predictive coding account. Progress in neurobiology. 2020;192:101821.

25. Marzen S. Difference between memory and prediction in linear recurrent networks. Physical Review E. 2017;96(3):032308.

26. Palmer SE, Marre O, Berry MJ, Bialek W. Predictive information in a sensory population. Proceedings of the National Academy of Sciences. 2015;112(22):6908–6913.

27. Berry MJ, Brivanlou IH, Jordan TA, Meister M. Anticipation of moving stimuli by the retina. Nature. 1999;398(6725):334–338.

28. Schwartz G, Harris R, Shrom D, Berry MJ. Detection and prediction of periodic patterns by the retina. Nature neuroscience. 2007;10(5):552–554.

29. Kastner DB, Baccus SA. Insights from the retina into the diverse and general computations of adaptation, detection, and prediction. Current opinion in neurobiology. 2014;25:63–69.

30. Körding KP, Wolpert DM. Bayesian decision theory in sensorimotor control. Trends in cognitive sciences. 2006;10(7):319–326.

31. Stachenfeld KL, Botvinick MM, Gershman SJ. The hippocampus as a predictive map. Nature neuroscience. 2017;20(11):1643–1653.

32. Hsu A, Marzen SE. Time cells might be optimized for predictive capacity, not redundancy reduction or memory capacity. Physical Review E. 2020;102(6):062404.

33. Guo C, Huson V, Macosko EZ, Regehr WG. Graded heterogeneity of metabotropic signaling underlies a continuum of cell-intrinsic temporal responses in unipolar brush cells. Nature Communications. 2021;12(1):5491.

34. Lamberti M, Tripathi S, van Putten MJAM, Marzen S, le Feber J. Prediction in cultured cortical neural networks. PNAS Nexus. 2023;2(6). doi:10.1093/pnasnexus/pgad188.

35. Debanne D, Inglebert Y. Spike timing-dependent plasticity and memory. Current Opinion in Neurobiology. 2023;80:102707.

36. Bruel-Jungerman E, Davis S, Laroche S. Brain plasticity mechanisms and memory: a party of four. The Neuroscientist. 2007;13(5):492–505.

37. Pokorny C, Ison MJ, Rao A, Legenstein R, Papadimitriou C, Maass W. STDP forms associations between memory traces in networks of spiking neurons. Cerebral Cortex. 2020;30(3):952–968.

38. le Feber J, Van Pelt J, Rutten W. Latency-related development of functional connections in cultured cortical networks. Biophysical journal. 2009;96(8):3443–3450.

39. Geinisman Y. Structural synaptic modifications associated with hippocampal LTP and behavioral learning. Cerebral Cortex. 2000;10(10):952–962.

40. Johnston D. A missing link? LTP and learning. Science. 1997;278(5337):401–402.

41. Rioult-Pedotti MS, Friedman D, Donoghue JP. Learning-induced LTP in neocortex. science. 2000;290(5491):533–536.

42. Pinsker H, Kupfermann I, Castellucci V, Kandel E. Habituation and dishabituation of the gill-withdrawal reflex in Aplysia. Science. 1970;167(3926):1740–1742.

43. Haas J, Nowotny T, Abarbanel HDI. Spike-Timing-Dependent Plasticity of Inhibitory Synapses in the Entorhinal Cortex. Journal of neurophysiology. 2006;96:3305–13. doi:10.1152/jn.00551.2006.

44. Frémaux N, Gerstner W. Neuromodulated spike-timing-dependent plasticity, and theory of three-factor learning rules. Frontiers in neural circuits. 2016;9:85.

45. Markram H, Gerstner W, Sjöström PJ. Spike-timing-dependent plasticity: a comprehensive overview. Frontiers in synaptic neuroscience. 2012;4:2.

46. Feldman DE. The spike-timing dependence of plasticity. Neuron. 2012;75(4):556–571.

47. Lu WY, Man HY, Ju W, Trimble WS, MacDonald JF, Wang YT. Activation of synaptic NMDA receptors induces membrane insertion of new AMPA receptors and LTP in cultured hippocampal neurons. Neuron. 2001;29(1):243–254.

48. Song I, Huganir RL. Regulation of AMPA receptors during synaptic plasticity. Trends in neurosciences. 2002;25(11):578–588.

49. Yang SN, Tang YG, Zucker RS. Selective induction of LTP and LTD by postsynaptic [Ca2+] i elevation. Journal of neurophysiology. 1999;81(2):781–787.

50. Kamioka H, Maeda E, Jimbo Y, Robinson HP, Kawana A. Spontaneous periodic synchronized bursting during formation of mature patterns of connections in cortical cultures. Neuroscience letters. 1996;206(2-3):109–112.

51. Chiappalone M, Bove M, Vato A, Tedesco M, Martinoia S. Dissociated cortical networks show spontaneously correlated activity patterns during in vitro development. Brain research. 2006;1093(1):41–53.

52. Le Feber J, Dummer A, Hassink GC, Van Putten MJ, Hofmeijer J. Evolution of excitation–inhibition ratio in cortical cultures exposed to hypoxia. Frontiers in cellular neuroscience. 2018;12:183.

53. le Feber J, Erkamp N, Van Putten MJ, Hofmeijer J. Loss and recovery of functional connectivity in cultured cortical networks exposed to hypoxia. Journal of neurophysiology. 2017;118(1):394–403.

54. Wagenaar DA, Madhavan R, Pine J, Potter SM. Controlling bursting in cortical cultures with closed-loop multi-electrode stimulation. Journal of Neuroscience. 2005;25(3):680–688.

55. Le Feber J, Stoyanova II, Chiappalone M. Connectivity, excitability and activity patterns in neuronal networks. Physical biology. 2014;11(3):036005.

56. Marzen SE, Crutchfield JP. Predictive rate-distortion for infinite-order Markov processes. Journal of Statistical Physics. 2016;163:1312–1338.

57. Teppola H, Aćimović J, Linne ML. Unique features of network bursts emerge from the complex interplay of excitatory and inhibitory receptors in rat neocortical networks. Frontiers in Cellular Neuroscience. 2019;13:377.

58. Li F, Tsien JZ. Memory and the NMDA receptors. The New England journal of medicine. 2009;361(3):302.

59. Mansvelder HD, Verhoog MB, Goriounova NA. Synaptic plasticity in human cortical circuits: cellular mechanisms of learning and memory in the human brain? Current opinion in neurobiology. 2019;54:186–193.

60. Hopfield JJ. Neural networks and physical systems with emergent collective computational abilities. Proceedings of the national academy of sciences. 1982;79(8):2554–2558.

61. Luczak A, McNaughton BL, Kubo Y. Neurons learn by predicting future activity. Nature Machine Intelligence. 2022;4:62–72.

62. Hunsaker MR, Kesner RP. The operation of pattern separation and pattern completion processes associated with different attributes or domains of memory. Neuroscience & Biobehavioral Reviews. 2013;37(1):36–58.

63. Knierim JJ, Neunuebel JP. Tracking the flow of hippocampal computation: Pattern separation, pattern completion, and attractor dynamics. Neurobiology of learning and memory. 2016;129:38–49.

64. Bologna LL, Pasquale V, Garofalo M, Gandolfo M, Baljon PL, Maccione A, et al. Investigating neuronal activity by SPYCODE multi-channel data analyzer. Neural Networks. 2010;23(6):685–697.

Supplementary Information

Additional analysis

APV concentration

We blocked NMDA receptors aiming to impede synaptic plasticity induced by NMDA receptor activation. Blocking a subgroup of excitatory receptors may also affect network excitability, which has been suggested as a possible determinant for memory functioning [23], and might thus bias results. We tested three APV concentrations: 15, 20 and 25 µM, in 5 cultures each. We first acquired 1 h of spontaneous activity ($Baseline$), and after APV administration we acquired an other hour of spontaneous activity ($Baseline_{APV}$) (see Figure S1). Periods of spontaneous activity ($Baseline$ and $Baseline_{APV}$) were used to determine Mean firing rate (MFR), burstiness index (BI), and mean connectivity strength, which are related to network excitability. The MFR was computed as defined by Bologna et al and following Dias et al [23, 64]. BI was used to quantify the synchronicity of each culture following Wagenaar et al. and Dias et al. [23, 54]. Mean connectivity strength was determined applying CFP [20]. We computed the ratio (baselineAPV/baseline) of MFR, BI, and mean strength of functional connections to quantify the effects of APV on these network parameters.

Responsiveness to stimulation

All memory experiments were performed using a constant stimulation frequency of 0.2 Hz, while in all prediction experiments inter-stimulus intervals were taken from a distribution with average stimulation frequency around 0.2 Hz. We chose this relatively low frequency to facilitate long lasting effectiveness of stimulation. We computed post stimulus time histograms (PSTHs) to assess the responsiveness of cultures to stimulation throughout the entire experimental protocol (memory: 17.5 h; prediction: 20 h). PSTHs were generated by summing all recorded activity in 5 ms bins from 300 ms before the stimulus until 500 ms after. In memory experiments PSTHs were computed for all 10 min stimulation periods, and in prediction experiments PSTHs were averaged per hour. To quantify network responsiveness to stimulation, we calculated the area under the curve (AUC), starting 15 ms after the stimulus (thus excluding stimulation artifacts and directly induced activity) and ending 500 ms after the stimulus. In memory experiments AUCs were normalized to the first 10 minutes of stimulation on that specific electrode, and in prediction experiments AUCs were normalized to the first hour. The duration of stimulus responses was determined from PSTHs as the maximum time after the stimulus where the average PSTH was still higher than the averaged activity before stimulation plus 5 times its standard deviation.

Memory formation

To investigate memory formation we estimated changes in functional connectivity of the network. Functional connectivity was inferred from spontaneous activity recordings

using conditional firing probability (CFP) [20]. This allows us to estimate strengths of connections between electrodes (groups of neurons), using the following equation:

$$CFP_{ij}(\tau) \approx o_{i,j} + \frac{M_{i,j}}{1 + \left(\frac{\tau - T_{i,j}}{w_{i,j}}\right)^2}. \tag{S1}$$

Here $M_{i,j}$ is interpreted as the strength of the connection, $T_{i,j}$ as the latency, $o_{i,j}$ represents uncorrelated background activity, and $w_{i,j}$ accounts for the width of the peak. For this analysis data were divided into chunks containing 2^{13} action potentials. This yielded a $60x60$ connectivity matrix M for each chunk, that contained the strengths of connections between all possible pairs of electrodes. Formation of memory traces was investigated quantifying connectivity changes induced by each stimulation period by computation of Euclidean distances (EDs) between connectivity matrices M [17, 23] using

$$ED(t) = \sqrt{\sum_{i=1}^{n}\sum_{j=1}^{n}(M_{ij}(t) - M_{ij}(t_0))^2} \tag{S2}$$

with $t > t_0$.

Relationship between memory and prediction

Estimates on the relationship between short-term memory and prediction were studied by computing the area under the curve of MI_{future} ($\Sigma MI_{\text{future}}$) and MI_{past} ($\Sigma MI_{\text{future}}$) for each hour of interest, for each experiment, using the following equations [34]:

$$\Sigma MI_{\text{future}} = \sum_{\Delta t=100}^{2000 \text{ ms}} MI_{\text{future}}(S, X)) \tag{S3}$$

$$\Sigma MI_{\text{past}} = \sum_{\Delta t=-1000}^{0 \text{ ms}} MI_{\text{past}}(S, X)). \tag{S4}$$

Here S indicates the binarized stimulation vector, and X refers to the binarized neuronal activity.

Cultured neural network composition

We used cultures on coverslips with the same density as cultures on MEA to visualize the presence of both neurons and astrocytes under control conditions [53]. They were fixated after 19–26 days in vitro using 4% paraformaldahyde, permeabilized with 0.2% Triton X−100 and blocked with 1% BSA. Then they were stained for mouse anti-microtubuleassociated protein 2 (MAP2; M9942, 1:200; Sigma-Aldrich, St. Louis, MO) or goat anti-glial fibrillary acidic protein (GFAP; SAB250046, 1:200; Sigma-Aldrich) [53]. Secondary antibodies used were donkey anti mouse-CF555 (SAB4600060, 1:500; Sigma-Aldrich) and donkey anti goat-488 (SAB4600032, 1:500; Sigma-Aldrich). We also stained excitatory and inhibitory (pre) synapses. Fixation was the same as mention above but NGS was used as blocking buffer [52]. Primary antibodies used were: mouse anti vGLUT; Sigma-Aldrich (Cat. Nr. SAB5200262-100UG), and rabbit anti vGAT; Sigma-Aldrich (Cat. Nr. V5764-200UL). Secondary antibodies used were donkey anti mouse 555; Sigma–Aldrich (SAB4600060-250UL), goat anti mouse TR; Santa Cruz (sc-2983), and donkey anti rabbit FITC; Jackson ImmunoResearch (711-095-152). In addition, DAPI staining was applied(1:1000; Sigma-Aldrich, Cat. Nr. D9542-1MG) [52, 53]. All pictures were taken using a Nikon Eclipse 50i fluorescence microscope.

Additional results

APV concentration

We tested 3 different concentrations of APV (15, 20 and 25 µM). The averaged mean firing rate significantly decreased by about 50% for 15 ($p = 0.026$) and 20 µM of APV ($p = 0.024$), and remained unaffected by 25 µM ($p = 0.34$). The burstiness index tended to decrease (by $\approx 50\%$) for all concentrations, but changes were not significant ($p = 0.19$, $p = 0.05$ and $p = 0.34$, respectively). The mean connectivity strength tended to decrease for all concentrations, but changes were not significant(15 µM $p = 0.08$, 20 µM $p = 0.34$ and 25 µM $p = 0.08$). We chose to use 20 µM of APV because it showed the best trade-off between the effect on plasticity and reduction of network excitability.

Responsiveness to stimulation

To assess responsiveness to stimulation throughout experiments, we calculated the area under the curve of post-stimulus time histograms as shown in Figure S2A and C. and the duration of stimulus responses as shown in Figure S2B and D. Two-way repeated measurement ANOVA showed that responsiveness did not significantly depend on time in the memory protocol ($p = 0.29$). Prediction data showed a significant interaction between time and APV application ($p < 0.05$), thus we performed one-way ANOVA to test the effect of time in controls and APV experiments separately. Stimulus response were not affected by time under control conditions ($p = 0.68$). However under APV administration responsiveness significantly depended on time ($p < 0.001$), with increasing responsiveness during the first 7 hours, followed by a decrease (Figure S2C). To avoid a possible bias of the quantification of prediction by the decreasing responsiveness, we chose to analyze only the first 10 hours. In both protocols, responsiveness did not significantly differ between conditions (control vs APV); memory protocol ($p = 0.53$), prediction protocol: $p = 0.54$), or on the two-way interaction between time and condition ($p = 0.63$). Figure S2B and D show that the duration of stimulus responses did not change with time (Memory protocol: $p = 0.3$; prediction protocol: $p = 0.49$), and did not differ between conditions (Memory protocol: $p = 0.51$; prediction protocol: $p = 0.56$), and there was no significant interaction ($p = 0.71$).

Additional figures

Fig S1. Effect of different APV concentrations on network excitability and plasticity. A) experimental protocol used to test each concentration. B) Effect of 15 (light blue), 20 (oblique stripes) and 25 µM (light green) of APV on mean firing rate (MFR). C) Effect of different APV concentrations on burstiness index (BI). D) Effect of the different APV concentrations on the mean connectivity strength. Significant changes are indicated by *.

Fig S2. Effectiveness of stimulation. In all panels control experiments are shown in orange and APV ones in green. A) and B) are analysis related to the memory experiments. A) shows that there was no effect on the AUC of the PSTHs of either time ($p = 0.29$) or APV addition ($p = 0.53$). B) shows that there was no effect on the end of the PSTHs of either time or APV addition (respectively $p = 0.3$ and $p = 0.51$). C) and D) regard prediction experiments. C) shows that there was no effect on the AUC of the PSTHs induced by APV addition ($p = 0.54$), but in this case AUC was influenced by time ($p < 0.001$), while this did not happen under control conditions ($p = 0.68$). D) shows that there was no effect on the end of the PSTHs of either time or APV addition ($p = 0.49$ and $p = 0.56$ respectively). Shaded areas represent SEM.

Fig S3. Cultured neural networks contained neurons and astrocytes, and excitatory and inhibitory synapses. A) Example of a culture, stained for MAP2 (neurons; red) and GFAP (astrocytes; green). B) Example of a culture stained for vGLUT (excitatory synapses; red) and vGAT (inhibitory synapses; green). In both panels cell nuclei were stained with DAPI (blue).