Reduction in neurons immunoreactive for parvalbumin, calretinin and calbindin in the anteroventral

thalamic nuclei of individuals with Down syndrome.

James C. Perry & Seralynne D. Vann

School of Psychology & Neuroscience and Mental Health Innovation Institute, Cardiff University,

Cardiff, UK

Corresponding authors:

Professor Seralynne Vann

School of Psychology

70 Park Place

Cardiff, CF10 3AT, UK

Abstract

The anterior thalamic nuclei are important for cognition, and memory in particular. However, little is known about how the anterior thalamic nuclei are affected in many neurological disorders partly due to difficulties in selective segmentation in *in vivo* scans, due to their size and location. Post-mortem studies, therefore, remain a valuable source of information about the status of the anterior thalamic nuclei. We used post-mortem tissue to assess the status of the anteroventral thalamic nucleus in Down syndrome using samples from males and females ranging from 22-65 years in age and comparing to tissue from age matched controls. As expected, there was increased beta-amyloid plaque expression in the Down syndrome group. While there was a significant increase in neuronal density in the Down syndrome group, the values showed more variation consistent with a heterogeneous population. The surface area of the anteroventral thalamic nucleus was smaller in the Down syndrome group suggesting the increased neuronal density was due to greater neuronal packing but likely fewer overall neurons. There was a marked reduction in the proportion of neurons immunoreactive for the calcium-binding proteins calbindin, calretinin, and parvalbumin in individuals with Down syndrome across all ages. These findings highlight the vulnerability of calcium-binding proteins in the anteroventral nucleus in Down syndrome, which could both be driven by, and exacerbate, Alzheimer-related pathology in this region.

Introduction

Down syndrome is the most common chromosomal disorder, affecting 1 in 1000 live births in the UK. This highlights the need to have a better understanding of how the brain can be affected with the long-term goal of better supporting cognitive function and reducing age-related decline. Memory is often affected in individuals with Down syndrome and this has typically been attributed to structural changes in the hippocampal formation (Carducci et al., 2013, Kemper 1991, Pinter et al., 2001). However, other brain regions that are important for memory, including the anterior thalamic nuclei, also appear to be compromised in Down syndrome; for example, we previously found a marked reduction of neurons in the anterior thalamic nuclei in individuals with Down syndrome (Perry et al., 2019). While these findings implicate the anterior thalamic nuclei in Down syndrome, a limitation of that study was that the tissue was exclusively from older females, so it is unclear whether these changes reflect the presence of advanced Alzheimer pathology in this group, rather than being more specifically related to Down syndrome. Alzheimer-related pathology is a common feature in individuals with Down syndrome with the onset of amyloid pathology commencing as early as the late teens and dementia often occuring by the mid 50s (Davidson et al., 2018, Head et al., 2012).

One aim of the current study was to determine whether the anterior thalamic nuclei exhibit changes in Down syndrome at younger ages. We, therefore, analysed post-mortem tissue from 15 individuals with Down syndrome (6 female) ranging from 22 to 62 years and compared this to tissue from 12 control individuals (6 female) ranging in age from 38 to 70 years. We measured cell density in the anteroventral thalamic nucleus, including neurons and glial cells, as well as the numbers of cells that stained positive for the calcium-binding proteins, calretinin, parvalbumin and calbindin, to determine whether any of these cell populations appeared to be affected in individuals with Down syndrome. The number of neurons immunopositive for calcium-binding proteins is generally reduced in Alzheimer's disease (Satoh et al., 1991). A reduction in calbindin and parvalbumin stained neurons

has also been reported in the cortex of older participants with Down syndrome (Kobayashi et al.,

1990) but there is relatively little known about the expression of calcium-binding proteins in the

human anterior thalamic nuclei in general, and even less knowledge in the case of Down syndrome.

Calcium-binding proteins are important for calcium homeostasis and intracellular signalling (Fairless

et al., 2019), highlighting the importance of understanding their expression in Down syndrome.

A subsequent aim was to assess whether Alzheimer's disease related pathology contributed to

neuronal calcium-binding protein expression given the links between calcium homeostasis and both

beta-amyloid (Aβ) and tau (Guan et al., 2021). Diffuse Aβ plaques are comprised primarily of Aβ42

and develop at a younger age, with the earliest cortical deposits being found in Down syndrome

around age 12 (Lemere et al., 1996). In contrast, the more mature plaques containing Aβ40

(Iwatsubo et al., 1995), and neurofibrillary tangles, have been more typically reported in cortical and

parahippocampal regions from age 30-35 in individuals with Down syndrome (Lemere et al., 1996,

Head et al., 2003). To determine how the differential development of plaques and tangles relate to

expression of calcium-binding proteins, we processed the tissue for tau, Aβ42, Aβ40, and Aβ-4G8 to

visualize total amyloid burden (Mandler et al., 2014). A more comprehensive understanding of the

vulnerability of cells within the anteroventral thalamic nucleus will help identify possible

contributing factors underlying structural changes and the extent to which these are driven by

Alzheimer-related pathology.

Methods

Tissue samples

Samples originated from formalin fixed paraffin embedded tissue blocks from the brains of 15 people

with Down syndrome (age range 22 – 65 years) and 12 controls (age range 38 – 70). Slide-mounted

consecutive serial coronal sections cut at 6μm thickness were provided by the UK Brain Banks

Network (UKBBN). The UKBBN adheres to agreed common standards, therefore, handling of tissue

was comparable across sites. Details of each case are provided in Additional File 1: Supplementary

Table 1. This study was conducted in accordance with the UK Human Tissue Act (2004). Approval for

the study was given by the School of Psychology Research Ethics Committee, Cardiff University, and

use of the tissue was covered by the generic approval from each brain bank (Manchester, REC

reference 19/NE/0242; London, 23/WA/0124; Cambridge, 20/EE/0283; Oxford, 23/SC/0241).

Immunohistochemistry

For each immunohistochemistry procedure, all sections were processed as a single batch with

staining carried out within a single session. Sections were deparaffinized in xylene then rehydrated in

a descending series of alcohols. Endogenous enzyme activity was quenched in either methanol

containing 3% H_2O_2 or Bloxall solution (SP-6000-100, Vector Laboratories, US) to block both

peroxidase and alkaline phosphate activity. Epitope retrieval was performed by incubating sections in

98% formic acid for 5 mins. Sections were then washed three times in 0.1M PBS containing 0.2%

triton-X 100 followed by blocking in 10% normal goat or horse serum (Vector Laboratories, US) for 1

h at room temperature. Sections were incubated in primary antibody solution overnight at 4°C in a

humidified chamber (Table 1). The Aβ-4G8, AT8, parvalbumin, and calbindin samples were incubated

in an appropriate biotinylated secondary antibody solution (1:200 dilution) for 1 h at room

temperature followed by streptavidin horseradish peroxidase solution for 30 mins. The calretinin

samples were incubated in ImmPRESS horseradish peroxidase polymer detection kit for 30mins

(Vector Laboratories, US). Samples were developed in a 3,3'-diaminobenzidine (DAB) substrate kit

(Vector Laboratories, US), counterstained using QS haematoxylin (Vector Laboratories, US), then

dehydrated in an ascending series of alcohols, cleared in xylene and coverslipped using DPX

mounting media. The Aβ 42/40 double-labelled samples were incubated in ImmPRESS horseradish

peroxidase polymer then ImmPRESS alkaline phosphate polymer kits for 30 mins each (Vector

Laboratories, US). For these sections, Aβ40 was visualized using StayYellow HRP chromogen

(ab169561, Abcam,UK) followed by Aβ42 visualization using StayBlue AP chromogen (ab176915,

Abcam, UK), counterstained with haematoxylin, air dried and coverslipped using aqueous mounting

media. For each case, one section was processed for each immunohistochemical marker except for

one 70-year-old control case where tissue was not stained for calbindin and Tau AT8 due to

insufficient samples. Primary antibody specificity was assessed by omitting the primary from one

sample per marker. In addition, the expected regional specificity of the neuropil stain for calretinin,

parvalbumin, and calbindin at low magnification along with neuronal, but not glial cell, label at

higher magnification was confirmed.

Table 1. Primary antibodies used for immunohistochemistry.

Antigen	Antibody	Dilution	Detection	Source
Aβ, 17-24	Mouse anti-4G8 (monoclonal)	1:1000	Streptavidin/DAB	BioLegend 800708
Aβ, 1-40	Rabbit anti- β1-40 (polyclonal)	1:200	ImmPress polymer horse-anti-rabbit HRP/StayYellow	Merck, AB5074P
Aβ, 1-42	Mouse anti-124F (monoclonal)	1:500	ImmPress polymer horse-anti-mouse AP/StayBlue	BioLegend, 805509
Tau (pSer202/ PThr205)	Rabbit anti-AT8/AH36 (monoclonal)	1:500	Streptavidin/DAB	StressMarq Biosciences, SMC-601.
Parvalbumin	Recombinant rabbit anti-parvalbumin (monoclonal)	1:500	Streptavidin/DAB	Abcam, ab181086
Calretinin	Goat anti-calretinin (polyclonal)	1:100	ImmPress polymer horse-anti-goat HRP/DAB	Novus, AF5065
Calbindin	Chicken anti-calbindin D-18K (polyclonal)	1:250	Streptavidin/DAB	Novus, NBP2-50028

Imaging and analysis

All analyses of the anterior thalamus in this study employed digital pathology methods. This allowed

objective assessments of the whole region on each section. To do this, images of the anterior

thalamus were obtained using an automated slide scanner (Olympus VS200, Olympus Life Science,

Japan) equipped with a color iDS camera (Olympus VS-264C). The anterior thalamus was imaged

using a 40x objective (NA 0.95) for Aβ-4G8, Tau AT8, hematoxylin, calretinin, parvalbumin and

calbindin and a 20x objective (NA 0.8) for Aβ 40/42. The images were loaded into QuPath (v0.4.3,

Bankhead et al., 2017) and the anterior ventral thalamic nuclei was annotated as previously

described (Perry et al., 2019). The percentage area of the anteroventral thalamic nucleus occupied by

Aβ-4G8, Aβ 40/42, and Tau AT8 was calculated by training a pixel classification model to identify the

relative area occupied by each marker.

For neurons and glia cells, the StarDist deep-learning model (Schmidt et al., 2018) was used to

identify each stained object in sections stained with hematoxylin alone. An object classifier was

trained to identify each object as either a neuron, astrocyte, oligodendrocyte, or microglia. The

classifier employed several measures including area, length, circularity, diameter, hematoxylin

intensity and a 100μm weighted smoothing for each measure. Similarly, StarDist was used to detect

calretinin, parvalbumin, and calbindin positive neurons in DAB and hematoxylin counter-stained

sections. To do this, the DAB and hematoxylin channels were extracted and summed into a single

detection channel. Positive neurons were identified by filtering detections to a minimum nucleus

area, circularity, DAB and hematoxylin intensity. To provide an indication of the performance of each

cell classifier, one expert manually annotated a 1,000,000 μm^2 area on one section for hematoxylin

stained neurons and glia and a 4,000,000μm^2 area on one section per calcium binding protein marker

for samples from both the controls and people with Down syndrome. The manually annotated

sections were regarded as the ground truth and were used to compare the accuracy, precision, and

recall of each cell classifier. Each cell classifier performed well, with high accuracy (range: 89-99%),

precision (range 89-98%), and recall (range: 89-97%) achieved, with comparable performance for

samples from both the control and Down syndrome groups (see Additional File 1: Supplementary

Table 2).

Neuron and glia counts could not be obtained from DAB-stained sections that were counterstained

with hematoxylin due to the background staining from DAB. As such, neuron and glia measurements

were only acquired from the hemaotxylin-only stained sections (one section per case). The numbers

of microglia were added to the total glia density and not analysed alone due to low numbers.

Calbindin, calretinin and parvalbumin counts were all derived from separate brain DAB-stained

sections, and again from one section per case. For neuron and glial counts, a measure of the total

number of items was derived within the boundaries of the anteroventral thalamic nuclei on each

section, these measures were then adjusted for the area of the anteroventral thalamic nucleus on

the section, providing a measure of objects per mm^2. Calbindin, calretinin and parvalbumin counts

were further adjusted for neuronal density for each case, to provide an approximate proportion of

immunopositive neurons.

One 40x annotated image, selected from the middle of the available consecutive series of sections

(i.e., the largest available region for each case) was used to provide an approximate estimate of the

cross-sectional area of the anteroventral thalamic nucleus (AV). Each section was coded as either

anterior or posterior AV, as the anteroposterior levels varied across samples.

Data Analysis

Statistical analyses were performed in JASP version 0.18.2 (JASP Team, 2024). Independent means *t*-

tests were used to compare the age and post-mortem delay of controls and individuals with Down

syndrome. The distribution of gender across the groups was assessed using a Chi-squared test.

Pearson correlation was used to assess whether post-mortem delay, i.e. the time between death and

tissue fixation, independent of group, was correlated with any measure. A 2 x 2 factorial ANOVA was

used to assess the cross-sectional area of the AV with Diagnosis (Control vs Down syndrome) as the

between groups factor and the Anterior-posterior level (anterior vs posterior) of the AV as the within

groups factor. Comparisons between histological markers across groups were carried out using

independent means t-tests. One-tailed tests were used for Aβ pathology markers while two-tailed

test were used for all other measures. Welch's t-test for unequal variances was used wherever

appropriate. A Mann-Whitney U test was used when normality was not assumed. Wherever relevant,

effect sizes between groups were expressed as Hedges g when equal variances were assumed,

Glass's Δ when equal variances were not assumed, rank-biserial correlation following a Mann-

Whitney U test, or η_p^2 following factorial ANOVA. Separate multiple regressions with backward

elimination (p <0.1 retention) were modelled with neuronal density, calretinin, parvalbumin, and

calbindin proportions as the dependent variables, and Aβ42, diagnosis, and age as the co-variates.

Among the Aβ pathology markers stained in this study, Aβ42 was selected as a predictor for the

regression model as it is thought to be the more pathogenic form of Aβ (Bodani et al., 2015,

Iwatsubo et al., 1995). Age was included in the model due to the association between advancing age

and AD pathology in Down syndrome (Davidson et al., 2018). Tau was not included due low

expression. The most parsimonious model with the best fit is reported. Additional details of

regression summaries, ANOVAs, and coefficients for each model are included in Additional File 2:

Supplementary tables 3 -14. Multiple comparisons were corrected with the Benjamini-Hochberg

procedure (Q = 0.05). The corrected p values (p_c) of less than 0.05 were considered significant. Given

the small numbers, correlations with age were not carried out but the data are presented in

scatterplots so that age and gender distribution across groups can be visualized.

In addition to significance testing, the Bayesian t-test framework was applied to the key outcome

measures: proportion of calretinin, parvalbumin, and calbindin positive neurons. Bayesian Mann-

Whitney U test, with Markov Monte Carlo set to five chains of 10, 000 iterations, was used when

normality was not assumed. The standardised effect size (δ) was assigned a Cauchy prior distribution

with $r = 1/\sqrt{2}$. For each test, the Bayes factor (BF_{10}), was calculated to quantify the relative evidence

for the null hypothesis (H_0) compared to the alternative hypothesis (H_1). BF_{10} provides a continuous

measure of evidence where a BF_{10} between 1 and 3 is considered anecdotal evidence, between 3

and 10 is considered moderate evidence and BF_{10} greater than 10 is considered strong evidence, in

favour of the alternative hypothesis, respectively. In comparison, a BF_{10} between 1 and 0.33 is

considered anecdotal evidence, between 0.33 and 0.1 is considered moderate evidence and lower

than 0.1 can be interpreted as strong evidence, in favour of the null hypothesis, respectively (van

Doorn et al., 2021). Here it is important to note that verbal labels used to categorise different Bayes

factors can be useful to facilitate scientific communication, but caution is needed due to the arbitrary

nature of these labels and the continuous nature of the Bayes factor. The Bayesian *t* test and

Bayesian Mann-Whitney *U* test posterior and prior plots, along with robustness checks where

applicable are included in Additional File 3: Supplementary figures 1 -3. Finally, we applied the

Bayesian framework to the calcium binding protein regression analyses as described above. In

addition to providing the BF_{10} for the most parsimonious model, this framework also quantifies the

relative evidence for the inclusion ($BF_{inclusion}$) of each co-variate in the model. A beta prior probability

(a = b =1) was applied to the models and a JZS prior (r scale = 0.354) was applied to the regression

coefficients. Additional details of Bayesian regression model comparisons, posterior summaries of

coefficients, and descriptive statistics are included in Additional File 4: Supplementary tables 15 – 23.

Figures were produced in R using ggplot2 (Wickham, 2016).

Results

Group demographics

The control group tissue came from a slightly older population (range 38-70 years) than the Down

syndrome (DS) group tissue (range 22-62 years), although the difference did not reach significance

(Table 2). There was no difference in gender distribution across the two groups (see Table 2).

Table 2. Demographic data

	Control group n=12	Down syndrome group n=15	t value/χ^2	p value
Mean age ± S.D.	57.4 ± 11.2 years	47.8 ± 13.1 years	2.02	0.06
Sex (M:F)	6:6	9:6	0.27	0.60

Given the sample size and distribution of data, we did not look at specific correlations between

histological measures and age. While there were no effects of gender, due to the small numbers it is

possible that more subtle gender effects may not be detected, therefore, age and gender

distributions are presented graphically for the histological measures. Post-mortem delay data were

available for all 12 controls and 7 of the 15 Down syndrome samples. While there was a trend for

individuals with Down syndrome (M = 70.7h, SD = 28.6) to have a longer post-mortem delay than

controls (M = 54.2h, SD = 21.7), this difference was not significant $t(17)$ = -1.43, p = 0.17,

uncorrected, g = 0.47, 95% CI: [-1.60, 0.32]. Pearson correlation was used to assess whether the

post-mortem delay, independent of group, was correlated with each measure. While there was no

significant correlation between post-mortem delay and any measure, there was a moderate

relationship between post-mortem delay and calretinin density (r = -0.45, p = 0.06, uncorrected; see

Additional File 1: Supplementary Table 3).

Regional area of anteroventral thalamic nucleus

While we did not have the tissue necessary to derive volumetric measurements for the anteroventral

thalamic nucleus (AV), we did estimate the cross-sectional regional area from one mid-section per

case (Fig. 1a - d). We found that the area of AV was approximately 37% smaller in the DS group

compared to controls and this difference was significant as shown by a main effect of Diagnosis ($F(1,$

$23)$ = 7.06, p_c = 0.023, η_p^2 = 0.24, Fig. 1e). The smaller AV area in the DS group was not due to

differences in the anterior-posterior levels between the samples as neither the main effect of

Anterior-posterior level ($F(1, 23)$ = 0.59, p_c = 0.45, η_p^2 = 0.03) nor the Diagnosis x Anterior-posterior

level interaction were significant ($F(1, 23)$ = 4.01, p_c = 0.07, η_p^2 = 0.15).

Fig 1. Cross-sectional area of the anteroventral thalamic nucleus. Representative examples of the

anteroventral thalamic nucleus from similar middle anterior – posterior levels in (a) 63yo male

control (case# BBN110.35427), (b) 61yo female control (case# BBN110.28899), (c) 47yo male with

Down syndrome (case# BBN_17189), (d) 60yo female with Down syndrome (case# BBN_2990) and

(e) box plot of the cross-sectional area of the anteroventral thalamic nuclei. The box indicates the

inter-quartile range (IQR), the whiskers show the range of values that are within 1.5 x IQR, the

horizontal line indicates the median, and the cross indicates the mean. The notches represent the

95% confidence interval for each median (1.58 x IQR/√(n)). Control, control group; DS, Down

syndrome group, F, females, M, males; *p<0.05, **p<0.01, ***p<0.001. Scale bar = 1mm. Note: The

examples of the anteroventral thalamic nucleus in (a) to (d) are all oriented with dorsal towards the

top of the image. The orientation of the anteroventral nucleus can vary between individuals at

similar anterior – posterior levels.

Neurons and glia

There was increased neuron density in the DS group ($t(16.482) = -2.27$, $p_c = 0.049$, $\Delta = -0.61$, 95% CI:

[-1.39, 0.19], Fig. 2a, e, i), however, the range in neuron density was much greater in the DS group

(44-288 cells/mm^2) than in the control group (74-129 cells/mm^2) suggesting a more heterogeneous

population. While the density of total glia was slightly higher in the DS group, this difference was not

significant ($t(25) = -1.13$, $p_c = 0.30$, $g = -0.423$, 95% CI: [-1.19, 0.35], Fig. 2b, f, j). The numerical

increase in total glia density was driven by a significant increase in astroglia density ($t(25) = -2.33$, p_c

= 0.04, *g* = -0.88, 95% CI: [-1.67, -0.07], Fig. 2c, g, k), but not, oligodendrocyte density (*t*(25) = 0.81, *p_c*

= 0.45, *g* = 0.30, 95% CI: [-0.46, 1.06], Fig. 2d, h, l), which was comparable across the groups.

Fig 2. Neuron and glia density in the anteroventral thalamic nucleus. Top panel, hematoxylin stained

examples of (a) a neuron with clearly stained nucleolus, nucleus, and surrounded by cytoplasm, (b)

examples of astroglia (yellow arrows), oligodendrocytes (black arrows), comma shaped microglia (red

arrow), and neurons (white arrow) and zoomed in examples of (c) astroglia showing a typical pale

nucleus with heterochromatin staining concentrated in granules below the nuclear membrane and

(d) oligodendrocytes with a classic rounded nucleus and dense chromatin staining. Middle panel, box

plots of the densities of (e) neurons, (f) total glia, (g) astroglia and (h) oligodendrocytes. The box

indicates the inter-quartile range (IQR), the whiskers show the range of values that are within 1.5 x

IQR, the horizontal line indicates the median, and the cross indicates the mean. The notches

represent the 95% confidence interval for each median (1.58xIQR/√(n)). Bottom panel, scatterplot

showing the relationship between age and density of (i) neurons, (j) total glia, (k) astroglia and (l)

oligodendrocytes. The lines are fitted with the general linear model and the shaded areas are the

95% confidence interval. Control, control group; DS, Down syndrome group, F, females, M, males.

*p<0.05, **p<0.01, ***p<0.001. Note: All neuron and glia examples come from the same 61yo male

control (case# BBN110.28899). Samples were imaged at 40x magnification and zoomed in using

QuPath to show the features of each cell type.

Amyloid and tau

As expected, the area fraction of AV occupied by Aβ was significantly higher in the DS group than the

control group for all three markers: Aβ-4G8 (Fig. 3a, d), Aβ42 (Fig. 3b, e) and Aβ40 (Fig. 3b, e) [Aβ-

4G8: Mann-Whitney U = 15.00, p_c < 0.001, r = -0.83, 95% CI: [-∞, -0.68], Fig. 3g, i; Aβ42: t(14.03) = -

4.32, p_c = 0.002, Δ = -1.12, 95% CI: [-∞, -0.38], Fig. 3h, j; Aβ40: Mann-Whitney U = 49.50, p_c = 0.02, r

= -0.45, 95% CI: [-∞, -0.11]]. While there was a significant group difference for Aβ40, expression was

generally low with levels near zero for 13 of the 15 tissue samples from the DS group. However,

moderate expression was found in one case (47-year-old female, 1.14% coverage), and high

expression was found in one case (47-year-old male, 7.24% coverage). Tau AT8 expression in the AV

was generally low and did not differ between the groups (p = 0.36, uncorrected). Tau AT8 expression

was near zero in 12 of the 15 tissue samples from individuals with DS. However, moderate expression

was found in one case (65-year-old male: 1.7% coverage), and high expression was found in two

cases (47-year-old female, 9.1% coverage; 60-year-old male, 6.5% coverage; Fig. 3c, f).

 Fig 3. Alzheimer's disease pathology in the anteroventral thalamic nucleus. Top panel, examples of

 (a) Aβ-4G8 plaques, (b) blue stained Aβ42 (white arrow) and yellow stained Aβ40 (black arrow)

 plaques taken from the same area of the anteroventral thalamic nucleus from a 47yo female with

Down syndrome (case# BBN_17186) and (c) Tau AT8 positive neurofibrillary tangles (white arrows)

and neuropil threads (black arrows) taken from a 65yo male with Down syndrome (case# BBN_2975).

Second panel, no expression of (d) Aβ-4G8, (e) Aβ42 and Aβ40, and (f) Tau AT8 in a representative

control taken from a 61yo female control (case# BBN110.28899). Third panel, box plots of the

percent area fraction occupied by (g) Aβ 4G8 and (h) Aβ42. The box indicates the inter-quartile range

(IQR), the whiskers show the range of values that are within 1.5xIQR, the horizontal line indicates the

median, and the cross indicates the mean. The notches represent the 95% confidence interval for

each median (1.58xIQR/√(n)). Bottom panel, scatterplot showing the relationship between age and

the percent area fraction occupied by (i) Aβ 4G8 and (j) Aβ42. The lines are fitted with the general

linear model and the shaded areas are the 95% confidence interval, truncated at zero. Control,

control group, DS, Down syndrome group; F, females, M, males. *p<0.05, **p<0.01, ***p<0.001.

Calcium-binding proteins (calretinin, calbindin, parvalbumin)

Counts of immunopositive neurons for each of the calcium-binding proteins were normalized by

dividing each calcium-binding protein marker density by the total neuronal density for each

individual case, to provide a relative proportion of immunopositive neurons. The proportion of

neurons immunopositive for all three calcium-binding proteins in AV was significantly reduced in the

DS group [calretinin, $t(25)$ = 3.23, p_c = 0.008, g = 1.21, 95% CI: [0.37, 2.03], Fig. 4a, d, g, j;

parvalbumin, Mann-Whitney U = 149, p_c = 0.009, r = 0.66, 95% CI: [0.33, 0.84], Fig. 4b, e, h, k;

calbindin, Mann-Whitney U = 156, p_c < 0.001, r = 0.89, 95% CI: [0.75, 0.96], Fig. 4c, f, i, l]. The same

pattern of results was found when using the density of neurons immunopositive for calcium-binding

proteins in the AV, rather than proportion [calretinin, $t(25)$ = 2.61, p_c = 0.023, g = 0.98, 95% CI: [0.16,

1.78]; parvalbumin, Mann-Whitney U = 144, p_c = 0.018, r = 0.60, 95% CI: [0.24, 0.81]; calbindin,

Mann-Whitney U = 140, p_c = 0.006, r = 0.70, 95% CI: [0.38, 0.87]].

Subsequent Bayesian analyses indicated strong evidence in favour of reduced calretinin and calbindin

proportions (BF_{10} = 11.41 and 19.43, respectively) and moderate evidence in favour of reduced

parvalbumin proportion (BF_{10} = 5.36), over a null hypothesis of no difference in proportions between

controls and individuals with Down syndrome.

Fig 4. Calcium-binding proteins in the anteroventral thalamic nucleus. Top panel, representative

examples of (a) calretinin from a 66yo female control (case# BBN_24381), (b) parvalbumin from a

60yo female control (case# BBN110.28668), and (c) calbindin positive neurons from a 61yo female

control (case# BBN110.28899). Second panel, representative examples of (d) calretinin from a 37yo

male with DS (case# BBN002.28703), (e) parvalbumin from a 60yo female with DS (case# BBN_2990),

and (f) calbindin from a 37yo male with DS (case# BBN002.28703). Third panel, box plots of

normalized densities of (g) calretinin, (h) parvalbumin, and (i) calbindin positive neurons. The box

indicates the inter-quartile range (IQR), the whiskers show the range of values that are within

1.5xIQR, the horizontal line indicates the median, and the cross indicates the mean. The notches

represent the 95% confidence interval for each median (1.58xIQR/√(n)). Bottom panel, scatterplot

showing the relationship between age and normalized densities of (j) calretinin, (k) parvalbumin, and

(l) calbindin positive neurons. The lines are fitted with the general linear model and the shaded areas

are the 95% confidence interval, truncated at zero. Control, control group; DS, Down syndrome

group; F, females, M, males. $*p<0.05$, $**p<0.01$, $***p<0.001$.

Multiple Regression

Backward regression analysis identified a model containing diagnosis and age as the best model

predicting normalised calretinin expression (adjusted $R^2 = 0.36$, $F(2,24) = 8.20$, $p_c = 0.006$). In

contrast, models containing only diagnosis were the best at predicting parvalbumin (adjusted $R^2 =$

0.29, $F(1, 25) = 11.71$, $p_c = 0.006$) and calbindin expression (adjusted $R^2 = 0.47$, $F(1, 24) = 23.16$, $p_c <$

0.001). A model containing Aβ42 neared significance as the best predictor of increasing neuronal

density (adjusted $R^2 = 0.12$, $F(1,25) = 4.45$, $p_c = 0.056$).

Bayesian regression analyses were carried out using diagnosis, age, and Aβ42 as predictors of

normalized calretinin, parvalbumin, and calbindin expression. For calretinin, there was strong

evidence that the model with diagnosis and age explained a large amount of the variance, $R^2 = 0.41$,

(BF_{10} = 21.47). Diagnosis provided moderate evidence (β_1 =-0.17, $BF_{inclusion}$ = 6.25) while Age (β_2 = -

0.003, $BF_{inclusion}$ = 1.84) and Aβ42 (β_3 = -0.012, $BF_{inclusion}$ = 1.84) provided only anecdotal evidence for

inclusion in the model. For parvalbumin, there was strong evidence that the model with diagnosis

alone explained a large amount of variance R^2 = 0.32 (BF_{10} = 16.70). Diagnosis provided moderate

evidence for inclusion in the model (β_1 =-0.21, $BF_{inclusion}$ = 7.11). In contrast, both age (β_2 =-0.00036,

$BF_{inclusion}$ = 0.57) and Aβ42 (β_3 =0.003, $BF_{inclusion}$ = 0.59) provided anecdotal evidence for exclusion

from the model. Finally, for calbindin, there was strong evidence that the model with diagnosis alone

explained a large amount of variance R^2 = 0.49 (BF_{10} = 290). Diagnosis provided strong evidence for

inclusion in the model (β_1 =-0.29, $BF_{inclusion}$ = 25.85). In contrast, both age (β_2 =-0.00094, $BF_{inclusion}$ =

0.57) and Aβ42 (β_3 =-0.014, $BF_{inclusion}$ = 0.57) provided anecdotal evidence for exclusion from the

model.

Discussion

Despite the longstanding association between the medial diencephalon and memory, there has been

surprisingly little research into the status of this brain region in individuals with Down syndrome. In

the current study we assessed the anteroventral thalamic nucleus (AV) in post-mortem tissue from

15 individuals with Down syndrome (DS group). In addition to measuring surface area and neuronal

density, we also measured calcium-binding protein immunoreactivity; calcium-binding proteins play

an important role in calcium homeostasis and intracellular signalling and have been linked to

neurodegeneration (Kelemen and Szilágyi, 2021).

As expected, we found higher levels of amyloid markers in the DS group, most notably in total Aβ

burden (as measured by Aβ-4G8) and diffuse Aβ plaques (as measured by Aβ42 expression).

Expression of Aβ40-positive mature plaques was generally low with near zero expression in 13 of 15

samples from individuals with Down syndrome. These findings are in line with previous reports

showing that Aβ42 plaques are more prevalent than Aβ40 plaques across all ages in Down syndrome

(Head et al., 2016). Previous studies have found less pronounced tau expression in the anterior

thalamic nuclei in Alzheimer's disease than other regions (Rüb et al., 2016, Braak and Braak, 1991,

Viney et al., 2022). Consistent with this, there was very little tau pathology in most of the cases we

examined across groups with near zero levels of tau in 12 of the 15 cases from the DS group. While

there were too few cases with high tau expression to draw any clear conclusions, there did not seem

to be an association with age as the individual with the highest tau levels was 49 years old, but that

case also had the lowest neuron density, suggesting neurons may be particularly vulnerable to high

levels of tau in the AV. The heterogeneity in both tau and amyloid expression in younger cases

suggests some individuals may have better protection against earlier Alzheimer's pathology (Maxwell

et al., 2021).

Astroglial, but not oligodendrocyte, density was significantly increased in the AV in the DS group. In

addition, there was an increase in neuron density in the AV of individuals with Down syndrome,

although there was far greater variability in the DS group with some showing very low neuron

density and others showing increased neuron density. The overall pattern of increased neuronal and

glial density may appear to contradict our previous findings (Perry et al., 2019), where we found

reduced total neuron and glia counts. However, the increased neuron density in the current study

likely reflects increased packing due to a reduction in AV volume. Due to sample availability, we were

not able to acquire total volume measures, as we were in our previous study (Perry et al., 2019), but

we acquired a surface area measure for AV for each case to provide a relative approximation of size.

The surface area of AV was approximately 37% smaller in the DS group than controls, therefore, even

with preserved or increased neuronal density, there are likely fewer neurons overall in the AV in the

DS group. Only a few studies have quantitatively assessed the anterior thalamic nuclei in Alzheimer's

disease (De Simone et al., 2024) and even fewer have carried out neuronal counts but, from the

studies that are available, there does not seem to be a significant reduction in either volume or

neuronal counts in this region using post-mortem tissue (Hornberger et al., 2012, Xuereb et al.,

1991), suggesting that the AV may be more impacted in Down syndrome than Alzheimer's disease

alone.

A further aim of the current study was to assess calcium-binding proteins in the AV in Down

syndrome. The proportion of AV neurons immunopositive for calbindin, calretinin and parvalbumin

was significantly reduced in the DS group, by 70%, 49% and 61%, respectively. Furthermore, we

found strong evidence in favour of reduced proportions of calretinin and calbindin positive cells and

moderate evidence in favour of reduced proportions of parvalbumin positive cells, according to the

Bayesian framework. A previous study had reported a reduction in both calbindin and parvalbumin

in the cortices of four older individuals with Down syndrome (Kobayashi et al., 1990). For calbindin,

they found a 38% and 28% reduction in immunoreactive neurons in frontal and temporal cortices,

respectively. The reported reduction in parvalbumin was slightly less than calbindin, with a 27%

decrease in frontal cortex and 22% decrease in temporal cortex. Consistent with these cortical

findings (Kobayashi et al., 1990), we also found a greater decrease in calbindin than parvalbumin.

However, the relative reduction for both parvalbumin and calbindin appears greater in AV than in

the cortex, especially considering our sample population also included younger individuals. In our

data set, there was increased neuronal density in the DS group but reduced calcium-binding protein

immunoreactivity; this would suggest a downregulation in expression rather than a loss of neurons.

However, we cannot tell from this data set whether there are inherent differences in these neuron

populations in Down syndrome, i.e., the expression profile is different from birth, or whether

downregulation occurs due to pathophysiological conditions or, for example, changes in BDNF-

levels, which are altered in Down syndrome and affect calcium-binding protein expression (Fairless

et al., 2019).

Parvalbumin immunoreactivity has been previously analysed in human AV. One study found 53-58%

of total neurons to be immunoreactive for parvalbumin (Dixon et al., 2000) while another found the

percentage of parvalbumin positive cells to range from 55-63% (Danos et al., 1998). In our study, we

found approximately 45% of cells to be immunoreactive for parvalbumin in the control group. The

methodology and samples are very different across studies, but it seems a consistent finding that

approximately half the neurons in AV are immunoreactive for parvalbumin.

Reductions in parvalbumin, calretinin, and calbindin immunoreactivity have been reported in

Alzheimer's disease (Zallo et al., 2018, Brady and Mufson, 1997, Kook et al., 2014b), however, these

studies have typically focused on the hippocampus and frontal cortex; to our knowledge, no study

has specifically assessed these markers in the anterior thalamic nuclei in Alzheimer's disease. It is,

therefore, not clear whether the reduction in calcium-binding proteins in the DS group simply

reflects a sensitivity to Alzheimer-related pathology or whether there are underlying changes to the

cell populations in Down syndrome that co-occur with, or even pre-date, amyloid changes. There did

not seem to be a clear relationship between amyloid or tau and calcium-binding protein expression

but there were cases with very low amyloid and tau expression that also had very low levels of

calbindin, parvalbumin and calretinin expression, consistent with the calcium-binding protein

changes being found irrespective of amyloid status. In support of this, according to our most

parsimonious frequentist and Bayesian regression models, $A\beta42$, the most pathological form of these

$A\beta$ species (Bodani et al., 2015, Iwatsubo et al., 1995), did not predict expression of any calcium-

binding protein. Down syndrome alone was the best predictor of calcium-binding protein expression.

In comparison, advancing age and $A\beta42$ deposition provided only some evidence in favour of

predicting calretinin expression, but some evidence to be excluded as predictors of parvalbumin and

calbindin expression. It is possible that biological age could be a stronger predictor than the

chronological age used in this study. Recently, others have proposed mathematical models to predict

a brain's biological age based on MRI derived measures of cortical thickness, cortical surface area,

and subcortical volumes (Liem et al., 2017, Salih et al., 2023). Future human tissue studies could

estimate brain age using stereological assessment of cortical and subcortical volumes. While there

does not seem to be a clear association between Alzheimer markers and calcium-binding protein

expression in this current data set, it is possible that earlier changes, for example Aβ and/or tau

seeding (Ye et al., 2017, Condello et al., 2022), may show clearer links. A reduction in calcium-binding

proteins would disrupt calcium homeostasis, and a loss of calbindin in a rodent model of Alzheimer's

disease was linked to increased pathogenesis (Kook et al., 2014a). It is, therefore, possible that an

early loss of calbindin could exacerbate Alzheimer-related pathology in Down syndrome, given

calbindin shows the greatest reduction in the present study.

While there are few studies on the thalamus in Down syndrome, a recent study assessed three

thalamic nuclei in tissue from foetuses with Down syndrome: mediodorsal nucleus, centromedian

nucleus and parafasciular nucleus (Stagni et al., 2020). A similar pattern was found across all nuclei,

with a reduction in neuron and glia density but no difference in surface area, consistent with an

overall reduction in total neurons in these thalamic regions. There was high expression of calretinin

in the mediodorsal thalamic nucleus, with approximately 80% of cells staining positive for calretinin

but there was no difference in expression between the control and Down syndrome group (Stagni et

al., 2020). This might suggest that expression levels are generally comparable across individuals with

Down syndrome and controls in early life, with downregulation only occurring with increasing age.

Alternatively, there may be variability across thalamic nuclei and the age at which changes occur,

which might be expected given the different expression profiles and functions across thalamic nuclei

(Stagni et al., 2020, Morel et al., 1997, Münkle et al., 2000, Wolff and Vann, 2019). Similarly, a

stereological study of the superior temporal gyrus of four males with Down syndrome aged 19-33

years found reduced neurons but preserved parvalbumin and a non-significant reduction in

calretinin-positive neurons (Giffin-Rao et al., 2022). This reinforces the importance of future studies

into the anterior thalamic nuclei in earlier age groups to identify whether there is downregulation in

calcium-binding protein in early life or if changes only occur after adolescence. If there is a

downregulation of calcium-binding protein expression after adolescence, then finding ways to

preserve the expression of these proteins may help protect against subsequent pathogenesis (Hijazi

et al., 2020).

Unlike other cortical and allocoritcal regions, the majority of calcium-binding positive neurons in the

primate anterior thalamus are projection neurons rather than interneurons (Jones and Hendry, 1989,

Dixon et al., 2000). Although calcium-binding protein expressing neurons are thought to largely

comprise distinct neuronal populations, it is possible that these proteins are co-expressed in some

anterior thalamic neurons. In support of this, neuronal co-expression of these proteins has been

reported in the human visual and temporal cortex regions (del Río and DeFelipe, 1997, Leuba and

Saini, 1997) and in the anterior thalamus of guinea pigs (Zakowski et al., 2013). We were not able to

assess colocalization in the current study, however, the total proportion of calcium-binding protein

positive neurons was greater than one among control samples but less than one among samples

from people with Down syndrome. This suggests some colocalization of neuronal calcium-binding

protein expression may be occurring, especially among control samples, highlighting the need for

future studies to examine the co-expression of calcium-binding proteins in the human anterior

thalamus.

The current study used digital pathology methods. This enabled objective assessment of all the

stained objects in the AV on each section. While this method to identify and quantify cells and for

measurements of surface area is suitable for identifying relative numbers between groups, it cannot

estimate the true number of cells or surface area in these samples. This can only be achieved with

stereological assessment which requires access to all the tissue from the region of interest. Much of

the tissue from current study is historical, with portions of the anterior thalamus having been used in

other studies. However, despite the lack of stereological assessments, the data still provides valuable

insights to changes in the anterior thalamus of younger and older adults with Down syndrome given

the rarity of this tissue and the lack of knowledge of the anterior thalamus in Down syndrome.

Although, there was variation in the anteroposterior level assessed across cases, the sections for

both the DS and control groups came from a similar range of anteroposterior levels. Furthermore,

cell density and calretinin immunoreactivity in the human anterior thalamus is uniform across the

anteroposterior extent of the nuclei (Braak and Weinel, 1985, Morel et al., 1997, Fortin et al., 1998),

giving additional weight to the findings. In immunohistochemistry studies, the reliability of findings

can be influenced by variations in tissue handling and staining and this is especially important when

few sections per case are stained. This source of variation was reduced in the current study as

samples were sourced from UKBBN banks that collected tissue according to an agreed common

standard and all sections for each immunohistochemistry procedure were processed in a single

session.

The present study adds to the growing literature linking neuropathology in the medial diencephalon,

including anterior thalamic nuclei and mammillary bodies, to various neurological disorders

(Milczarek et al., 2023a, Lequin et al., 2021, Meys et al., 2022, Clarke et al., 1994, Connaughton et al.,

2023, Danos et al., 1998). Pathology in these brain areas would contribute to memory impairments

often accompanying these conditions by affecting both the encoding and consolidation of memories

(Milczarek et al., 2023b, Aggleton et al., 2023). In the present study, we found changes in the AV in

Down syndrome at a younger age than previously reported and showed for the first time that there

is a marked loss of calcium-binding protein immunoreactivity. Noticeably, this reduction in

parvalbumin, calretinin and calbindin immunoreactivity was not simply driven by overall neuronal

loss but likely reflects a downregulation in expression. These reductions in calcium-binding proteins

may increase the vulnerability of cells within this region and potentially exacerbate expression of

Alzheimer-related pathology.

Acknowledgements

This research was funded by a grant from the Jerome Lejeune Foundation (# 1899) and a Wellcome

Trust Senior Research Fellowship (WT212273/Z/18) awarded to SDV.

Tissue samples were sourced from brain banks in the UK Brain Banks Network. The Manchester Brain

Bank, which is part of the Brains for Dementia Research Initiative, jointly funded by Alzheimer's

Society and Alzheimer's Research UK. The Cambridge Brain Bank is supported by the NIHR

Cambridge Biomedical Research Centre (NIHR203312). We acknowledge the Oxford Brain Bank,

supported by Brains Dementia Research (BDR) (Alzheimer Society and Alzheimer Research UK) and

the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC). Tissue

samples were provided by The London Neurodegenerative Diseases Brain Bank at King's College

London. The brain bank receives funding as part of the Brains for Dementia Research programme,

jointly funded by Alzheimer's Research UK and the Alzheimer's Society.

List of abbreviations

AV, anteroventral thalamic nucleus; Aβ, beta-amyloid; BF_{10}, Bayes Factor; DAB, 3,3'-

diaminobenzidine; DS, Down syndrome.

References

Aggleton, J. P., Vann, S. D. & O'mara, S. M. 2023. Converging diencephalic and hippocampal supports
for episodic memory. *Neuropsychologia,* 191**,** 108728.
Bankhead, P., Loughrey, M. B., Fernández, J. A., Dombrowski, Y., Mcart, D. G., Dunne, P. D., Mcquaid,
S., Gray, R. T., Murray, L. J., Coleman, H. G., James, J. A., Salto-Tellez, M. & Hamilton, P. W.
2017. QuPath: Open source software for digital pathology image analysis. *Scientific Reports,*
7**,** 16878.
Bodani, R. U., Sengupta, U., Castillo-Carranza, D. L., Guerrero-Muñoz, M. J., Gerson, J. E., Rudra, J. &
Kayed, R. 2015. Antibody against Small Aggregated Peptide Specifically Recognizes Toxic Aβ-
42 Oligomers in Alzheimer's Disease. *ACS Chem Neurosci,* 6**,** 1981-9.
Braak, H. & Braak, E. 1991. Alzheimer's disease affects limbic nuclei of the thalamus. *Acta*
Neuropathol, 81**,** 261-8.
Braak, H. & Weinel, U. 1985. The percentage of projection neurons and local circuit neurons in
different nuclei of the human thalamus. *J Hirnforsch,* 26**,** 525-30.
Brady, D. R. & Mufson, E. J. 1997. Parvalbumin-immunoreactive neurons in the hippocampal
formation of Alzheimer's diseased brain. *Neuroscience,* 80**,** 1113-1125.
Carducci, F., Onorati, P., Condoluci, C., Di Gennaro, G., Quarato, P. P., Pierallini, A., Sara, M., Miano, S.,
Cornia, R. & Albertini, G. 2013. Whole-brain voxel-based morphometry study of children and
adolescents with Down syndrome. *Funct Neurol,* 28**,** 19-28.
Clarke, S., Assal, G., Bogousslavsky, J., Regli, F., Townsend, D. W., Leenders, K. L. & Blecic, S. 1994.
Pure amnesia after unilateral left polar thalamic infarct: topographic and sequential

596   neuropsychological and metabolic (PET) correlations. *J Neurol Neurosurg Psychiatry,* 57, 27-
597   34.

598   Condello, C., Maxwell, A. M., Castillo, E., Aoyagi, A., Graff, C., Ingelsson, M., Lannfelt, L., Bird, T. D.,
599   Keene, C. D., Seeley, W. W., Perl, D. P., Head, E. & Prusiner, S. B. 2022. Aβ and tau prions
600   feature in the neuropathogenesis of Down syndrome. *Proc Natl Acad Sci U S A,* 119,
601   e2212954119.

602   Connaughton, M., O'hanlon, E., Silk, T. J., Paterson, J., O'neill, A., Anderson, V., Whelan, R. & Mcgrath,
603   J. 2023. The Limbic System in Children and Adolescents with Attention-Deficit/Hyperactivity
604   Disorder: A Longitudinal Structural MRI Analysis. *Biological Psychiatry Global Open Science.*

605   Danos, P., Baumann, B., Bernstein, H.-G., Franz, M., Stauch, R., Northoff, G., Krell, D., Falkai, P. &
606   Bogerts, B. 1998. Schizophrenia and anteroventral thalamic nucleus: selective decrease of
607   parvalbumin-immunoreactive thalamocortical projection neurons. *Psychiatry Research:*
608   *Neuroimaging,* 82, 1-10.

609   Davidson, Y. S., Robinson, A., Prasher, V. P. & Mann, D. M. A. 2018. The age of onset and evolution of
610   Braak tangle stage and Thal amyloid pathology of Alzheimer's disease in individuals with
611   Down syndrome. *Acta Neuropathol Commun,* 6, 56.

612   De Simone, M. S., Spalletta, G., Vecchio, D., Bassi, A., Carlesimo, G. A. & Piras, F. 2024. The Role of the
613   Anterior Thalamic Nuclei in the Genesis of Memory Disorders in Alzheimer's Disease: An
614   Exploratory Study. *J Alzheimers Dis,* 97, 507-519.

615   Del Río, M. R. & Defelipe, J. 1997. Colocalization of parvalbumin and calbindin D-28k in neurons
616   including chandelier cells of the human temporal neocortex. *J Chem Neuroanat,* 12, 165-73.

617   Dixon, G., Dissanaike, S. & Harper, C. G. 2000. Parvalbumin-immunoreactive neurons in the human
618   anteroventral thalamic nucleus. *Neuroreport,* 11, 97-101.

619   Fairless, R., Williams, S. K. & Diem, R. 2019. Calcium-Binding Proteins as Determinants of Central
620   Nervous System Neuronal Vulnerability to Disease. *Int J Mol Sci,* 20.

621   Fortin, M., Asselin, M. C., Gould, P. V. & Parent, A. 1998. Calretinin-immunoreactive neurons in the
622   human thalamus. *Neuroscience,* 84, 537-548.

623   Giffin-Rao, Y., Sheng, J., Strand, B., Xu, K., Huang, L., Medo, M., Risgaard, K. A., Dantinne, S., Mohan,
624   S., Keshan, A., Daley, R. A., Jr., Levesque, B., Amundson, L., Reese, R., Sousa, A. M. M., Tao, Y.,
625   Wang, D., Zhang, S. C. & Bhattacharyya, A. 2022. Altered patterning of trisomy 21
626   interneuron progenitors. *Stem Cell Reports,* 17, 1366-1379.

627   Guan, P. P., Cao, L. L. & Wang, P. 2021. Elevating the Levels of Calcium Ions Exacerbate Alzheimer's
628   Disease via Inducing the Production and Aggregation of β-Amyloid Protein and
629   Phosphorylated Tau. *Int J Mol Sci,* 22.

630   Head, E., Lott, I. T., Hof, P. R., Bouras, C., Su, J. H., Kim, R., Haier, R. & Cotman, C. W. 2003. Parallel
631   Compensatory and Pathological Events Associated with Tau Pathology in Middle Aged
632   Individuals with Down Syndrome. *Journal of Neuropathology & Experimental Neurology,* 62,
633   917-926.

634   Head, E., Lott, I. T., Wilcock, D. M. & Lemere, C. A. 2016. Aging in Down Syndrome and the
635   Development of Alzheimer's Disease Neuropathology. *Curr Alzheimer Res,* 13, 18-29.

636   Head, E., Powell, D., Gold, B. T. & Schmitt, F. A. 2012. Alzheimer's Disease in Down Syndrome. *Eur J*
637   *Neurodegener Dis,* 1, 353-364.

638   Hijazi, S., Heistek, T. S., Scheltens, P., Neumann, U., Shimshek, D. R., Mansvelder, H. D., Smit, A. B. &
639   Van Kesteren, R. E. 2020. Early restoration of parvalbumin interneuron activity prevents
640   memory loss and network hyperexcitability in a mouse model of Alzheimer's disease. *Mol*
641   *Psychiatry,* 25, 3380-3398.

642   Hornberger, M., Wong, S., Tan, R., Irish, M., Piguet, O., Kril, J., Hodges, J. R. & Halliday, G. 2012. In
643   vivo and post-mortem memory circuit integrity in frontotemporal dementia and Alzheimer's
644   disease. *Brain,* 135, 3015-25.

645   Iwatsubo, T., Mann, D. M., Odaka, A., Suzuki, N. & Ihara, Y. 1995. Amyloid beta protein (A beta)
646   deposition: A beta 42(43) precedes A beta 40 in Down syndrome. *Ann Neurol,* 37, 294-9.

Jones, E. G. & Hendry, S. H. 1989. Differential Calcium Binding Protein Immunoreactivity
Distinguishes Classes of Relay Neurons in Monkey Thalamic Nuclei. *Eur J Neurosci,* 1, 222-
246.
Kelemen, K. & Szilágyi, T. 2021. New Approach for Untangling the Role of Uncommon Calcium-
Binding Proteins in the Central Nervous System. *Brain Sci,* 11.
Kemper , T. L. 1991. Down Syndrome. *In:* PETERS, A. & JONES, E. G. (eds.) *Cerebral Cortex: Normal*
and altered states of function. Plenum Press.
Kobayashi, K., Emson, P. C., Mountjoy, C. Q., Thornton, S. N., Lawson, D. E. & Mann, D. M. 1990.
Cerebral cortical calbindin D28K and parvalbumin neurones in Down's syndrome. *Neurosci*
Lett, 113, 17-22.
Kook, S. Y., Jeong, H., Kang, M. J., Park, R., Shin, H. J., Han, S. H., Son, S. M., Song, H., Baik, S. H.,
Moon, M., Yi, E. C., Hwang, D. & Mook-Jung, I. 2014a. Crucial role of calbindin-D28k in the
pathogenesis of Alzheimer's disease mouse model. *Cell Death Differ,* 21, 1575-87.
Kook, S. Y., Jeong, H., Kang, M. J., Park, R., Shin, H. J., Han, S. H., Son, S. M., Song, H., Baik, S. H.,
Moon, M., Yi, E. C., Hwang, D. & Mook-Jung, I. 2014b. Crucial role of calbindin-D28k in the
pathogenesis of Alzheimer's disease mouse model. *Cell Death & Differentiation,* 21, 1575-
1587.
Lemere, C. A., Blusztajn, J. K., Yamaguchi, H., Wisniewski, T., Saido, T. C. & Selkoe, D. J. 1996.
Sequence of deposition of heterogeneous amyloid beta-peptides and APO E in Down
syndrome: implications for initial events in amyloid plaque formation. *Neurobiol Dis,* 3, 16-
32.
Lequin, M. H., Steggerda, S. J., Severino, M., Tortora, D., Parodi, A., Ramenghi, L. A., Groenendaal, F.,
Meys, K. M. E., Benders, M., De Vries, L. S. & Vann, S. D. 2021. Mammillary body injury in
neonatal encephalopathy: a multicentre, retrospective study. *Pediatr Res.*
Leuba, G. & Saini, K. 1997. Colocalization of parvalbumin, calretinin and calbindin D-28k in human
cortical and subcortical visual structures. *J Chem Neuroanat,* 13, 41-52.
Liem, F., Varoquaux, G., Kynast, J., Beyer, F., Kharabian Masouleh, S., Huntenburg, J. M., Lampe, L.,
Rahim, M., Abraham, A., Craddock, R. C., Riedel-Heller, S., Luck, T., Loeffler, M., Schroeter, M.
L., Witte, A. V., Villringer, A. & Margulies, D. S. 2017. Predicting brain-age from multimodal
imaging data captures cognitive impairment. *Neuroimage,* 148, 179-188.
Mandler, M., Walker, L., Santic, R., Hanson, P., Upadhaya, A. R., Colloby, S. J., Morris, C. M., Thal, D.
R., Thomas, A. J., Schneeberger, A. & Attems, J. 2014. Pyroglutamylated amyloid-β is
associated with hyperphosphorylated tau and severity of Alzheimer's disease. *Acta*
Neuropathol, 128, 67-79.
Maxwell, A. M., Yuan, P., Rivera, B. M., Schaaf, W., Mladinov, M., Prasher, V. P., Robinson, A. C.,
Degrado, W. F. & Condello, C. 2021. Emergence of distinct and heterogeneous strains of
amyloid beta with advanced Alzheimer's disease pathology in Down syndrome. *Acta*
Neuropathologica Communications, 9, 201.
Meys, K. M. E., De Vries, E., Groenendaal, F., Vann, S. & Lequin, M. H. 2022. The mammillary bodies:
a review of causes of injury in infants and children. *AJNR Am J Neuroradiol.*
Milczarek, M. M., Gilani, S. I. A., Lequin, M. H. & Vann, S. D. 2023a. Reduced mammillary body
volume in individuals with a schizophrenia diagnosis: an analysis of the COBRE data set.
Schizophrenia (Heidelb), 9, 48.
Milczarek, M. M., Perry, J. C., Amin, E., Haniffa, S., Hathaway, T. & Vann, S. D. 2023b. Impairments in
the early consolidation of spatial memories via group II mGluR agonism in the mammillary
bodies. *bioRxiv,* 2023.12.13.571542.
Morel, A., Magnin, M. & Jeanmonod, D. 1997. Multiarchitectonic and stereotactic atlas of the human
thalamus. *Journal of Comparative Neurology,* 387, 588-630.
Münkle, M. C., Waldvogel, H. J. & Faull, R. L. M. 2000. The distribution of calbindin, calretinin and
parvalbumin immunoreactivity in the human thalamus. *Journal of Chemical Neuroanatomy,*
19, 155-173.

Perry, J. C., Pakkenberg, B. & Vann, S. D. 2019. Striking reduction in neurons and glial cells in anterior
thalamic nuclei of older patients with Down syndrome. *Neurobiol Aging,* 75**,** 54-61.
Pinter, J. D., Brown, W. E., Eliez, S., Schmitt, J. E., Capone, G. T. & Reiss, A. L. 2001. Amygdala and
hippocampal volumes in children with Down syndrome: a high-resolution MRI study.
Neurology, 56**,** 972-4.
Rüb, U., Stratmann, K., Heinsen, H., Del Turco, D., Ghebremedhin, E., Seidel, K., Den Dunnen, W. &
Korf, H. W. 2016. Hierarchical Distribution of the Tau Cytoskeletal Pathology in the Thalamus
of Alzheimer's Disease Patients. *J Alzheimers Dis,* 49**,** 905-15.
Salih, A., Nichols, T., Szabo, L., Petersen, S. E. & Raisi-Estabragh, Z. 2023. Conceptual Overview of
Biological Age Estimation. *Aging Dis,* 14**,** 583-588.
Satoh, J., Tabira, T., Sano, M., Nakayama, H. & Tateishi, J. 1991. Parvalbumin-immunoreactive neurons
in the human central nervous system are decreased in Alzheimer's disease. *Acta*
Neuropathologica, 81**,** 388-395.
Schmidt, U., Weigert, M., Broaddus, C. & Myers, G. Cell Detection with Star-Convex Polygons. *In:*
FRANGI, A. F., SCHNABEL, J. A., DAVATZIKOS, C., ALBEROLA-LÓPEZ, C. & FICHTINGER, G., eds.
Medical Image Computing and Computer Assisted Intervention – MICCAI 2018, 2018// 2018
Cham. Springer International Publishing, 265-273.
Stagni, F., Giacomini, A., Emili, M., Uguagliati, B., Bonasoni, M. P., Bartesaghi, R. & Guidi, S. 2020.
Neuroanatomical alterations in higher-order thalamic nuclei of fetuses with Down syndrome.
Clinical Neurology and Neurosurgery, 194**,** 105870.
Team, J. 2024. JASP (Version 0.18.2)[Computer software]. https:// jasp-stats.org/
Van Doorn, J., Van Den Bergh, D., Böhm, U., Dablander, F., Derks, K., Draws, T., Etz, A., Evans, N. J.,
Gronau, Q. F., Haaf, J. M., Hinne, M., Kucharský, Š., Ly, A., Marsman, M., Matzke, D., Gupta, A.
R. K. N., Sarafoglou, A., Stefan, A., Voelkel, J. G. & Wagenmakers, E.-J. 2021. The JASP
guidelines for conducting and reporting a Bayesian analysis. *Psychonomic Bulletin & Review,*
28**,** 813-826.
Viney, T. J., Sarkany, B., Ozdemir, A. T., Hartwich, K., Schweimer, J., Bannerman, D. & Somogyi, P. 2022.
Spread of pathological human Tau from neurons to oligodendrocytes and loss of high-firing
pyramidal neurons in aging mice. *Cell Rep,* 41**,** 111646.
Wickham, H. 2016. *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag New York.
Wolff, M. & Vann, S. D. 2019. The Cognitive Thalamus as a Gateway to Mental Representations. *J*
Neurosci, 39**,** 3-14.
Xuereb, J. H., Perry, R. H., Candy, J. M., Perry, E. K., Marshall, E. & Bonham, J. R. 1991. Nerve cell loss
in the thalamus in Alzheimer's disease and Parkinson's disease. *Brain,* 114 (Pt 3)**,** 1363-79.
Ye, L., Rasmussen, J., Kaeser, S. A., Marzesco, A. M., Obermüller, U., Mahler, J., Schelle, J., Odenthal,
J., Krüger, C., Fritschi, S. K., Walker, L. C., Staufenbiel, M., Baumann, F. & Jucker, M. 2017. Aβ
seeding potency peaks in the early stages of cerebral β-amyloidosis. *EMBO reports,* 18**,** 1536-
1544-1544.
Zakowski, W., Bogus-Nowakowska, K. & Robak, A. 2013. Embryonic and postnatal development of
calcium-binding proteins immunoreactivity in the anterior thalamus of the guinea pig. *J*
Chem Neuroanat, 53**,** 25-32.
Zallo, F., Gardenal, E., Verkhratsky, A. & Rodríguez, J. J. 2018. Loss of calretinin and parvalbumin
positive interneurones in the hippocampal CA1 of aged Alzheimer's disease mice.
Neuroscience Letters, 681**,** 19-25.
742