1   Reduced expression of calcium-binding proteins in the anteroventral thalamic nucleus of individuals

2   with Down syndrome

James C. Perry & Seralynne D. Vann

School of Psychology & Neuroscience and Mental Health Innovation Institute, Cardiff University,

Cardiff, UK

Corresponding authors:

Professor Seralynne Vann

School of Psychology

70 Park Place

Cardiff, CF10 3AT, UK

Abstract

The anterior thalamic nuclei are important for cognition, and memory in particular. However, little is known about how the anterior thalamic nuclei are affected in many neurological disorders partly due to difficulties in selective segmentation in *in vivo* scans, due to their size and location. Post-mortem studies, therefore, remain a valuable source of information about the status of the anterior thalamic nuclei. We used post-mortem tissue to assess the status of the anteroventral thalamic nucleus in Down syndrome using samples from males and females ranging from 22-65 years in age and comparing to tissue from age matched controls. As expected, there was increased beta-amyloid plaque expression in the Down syndrome group. While there was a significant increase in neuronal density in the Down syndrome group, the values showed more variation consistent with a heterogeneous population. The surface area of the anteroventral thalamic nucleus was smaller in the Down syndrome group suggesting the increased neuronal density was due to greater neuronal packing but likely fewer overall neurons. There was a marked reduction in the proportion of neurons immunoreactive for the calcium-binding proteins calbindin, calretinin, and parvalbumin in individuals with Down syndrome across all ages. These findings highlight the vulnerability of calcium-binding proteins in the anteroventral nucleus in Down syndrome, which could both be driven by, and exacerbate, Alzheimer-related pathology in this region.

Introduction

Down syndrome is the most common chromosomal disorder, affecting 1 in 1000 live births in the UK. This highlights the need to have a better understanding of how the brain can be affected with the long-term goal of better supporting cognitive function and reducing age-related decline. Memory is often affected in individuals with Down syndrome and this has typically been attributed to structural changes in the hippocampal formation (Carducci et al., 2013, Kemper 1991, Pinter et al., 2001). However, other brain regions that are important for memory, including the anterior thalamic nuclei, also appear to be compromised in Down syndrome; for example, we previously found a marked reduction of neurons in the anterior thalamic nuclei in individuals with Down syndrome (Perry et al., 2019). While these findings implicate the anterior thalamic nuclei in Down syndrome, a limitation of that study was that the tissue was exclusively from older females, so it is unclear whether these changes reflect the presence of advanced Alzheimer pathology in this group, rather than being more specifically related to Down syndrome. Alzheimer-related pathology is a common feature in individuals with Down syndrome with the onset of amyloid pathology commencing as early as the late teens and dementia often occuring by the mid 50s (Davidson et al., 2018, Head et al., 2012).

One aim of the current study was to determine whether the anterior thalamic nuclei exhibit changes in Down syndrome at younger ages. We, therefore, analysed post-mortem tissue from 15 individuals with Down syndrome (6 female) ranging from 22 to 62 years and compared this to tissue from 12 control individuals (6 female) ranging in age from 38 to 70 years. We measured cell density in the anteroventral thalamic nucleus, including neurons and glial cells, as well as the numbers of cells that stained positive for the calcium-binding proteins, calretinin, parvalbumin and calbindin, to determine whether any of these cell populations appeared to be affected in individuals with Down syndrome. The number of neurons immunopositive for calcium-binding proteins is generally reduced in Alzheimer's disease (Satoh et al., 1991). A reduction in calbindin and parvalbumin stained neurons

has also been reported in the cortex of older participants with Down syndrome (Kobayashi et al.,

1990) but there is relatively little known about the expression of calcium-binding proteins in the

human anterior thalamic nuclei in general, and even less knowledge in the case of Down syndrome.

Calcium-binding proteins are important for calcium homeostasis and intracellular signalling (Fairless

et al., 2019), highlighting the importance of understanding their expression in Down syndrome.

A subsequent aim was to assess whether Alzheimer's disease related pathology contributed to

neuronal calcium-binding protein expression given the links between calcium homeostasis and both

beta-amyloid (Aβ) and tau (Guan et al., 2021). Diffuse Aβ plaques are comprised primarily of Aβ42

and develop at a younger age, with the earliest cortical deposits being found in Down syndrome

around age 12 (Lemere et al., 1996). In contrast, the more mature plaques containing Aβ40

(Iwatsubo et al., 1995), and neurofibrillary tangles, have been more typically reported in cortical and

parahippocampal regions from age 30-35 in individuals with Down syndrome (Lemere et al., 1996,

Head et al., 2003). To determine how the differential development of plaques and tangles relate to

expression of calcium-binding proteins, we processed the tissue for tau, Aβ42, Aβ40, and Aβ-4G8 to

visualize total amyloid burden (Mandler et al., 2014). A more comprehensive understanding of the

vulnerability of cells within the anteroventral thalamic nucleus will help identify possible

contributing factors underlying structural changes and the extent to which these are driven by

Alzheimer-related pathology.

Methods

Tissue samples

Samples originated from formalin fixed paraffin embedded tissue blocks from the brains of 15 people

with Down syndrome (age range 22 – 65 years) and 12 controls (age range 38 – 70). Slide-mounted

consecutive serial coronal sections cut at 6μm thickness were provided by the UK Brain Bank

Network. Details of each case are provided in Additional File 1: Supplementary Table 1. This study

was conducted in accordance with the UK Human Tissue Act (2004). Approval for the study was given

by the School of Psychology Research Ethics Committee, Cardiff University, and use of the tissue was

covered by the generic approval from each brain bank (Manchester, REC reference 19/NE/0242;

London, 23/WA/0124; Cambridge, 20/EE/0283; Oxford, 23/SC/0241).

Immunohistochemistry

Sections were deparaffinized in xylene then rehydrated in a descending series of alcohols.

Endogenous enzyme activity was quenched in either methanol containing 3% H_2O_2 or Bloxall solution

(SP-6000-100, Vector Laboratories, US) to block both peroxidase and alkaline phosphate activity.

Epitope retrieval was performed by incubating sections in 98% formic acid for 5 mins. Sections were

then washed three times in 0.1M PBS containing 0.2% triton-X 100 followed by blocking in 10%

normal goat or horse serum (Vector Laboratories, US) for 1 h at room temperature. Sections were

incubated in primary antibody solution overnight at 4°C in a humidified chamber (Table 1). The Aβ-

4G8, AT8, parvalbumin, and calbindin samples were incubated in an appropriate biotinylated

secondary antibody solution (1:200 dilution) for 1 h at room temperature followed by streptavidin

horseradish peroxidase solution for 30 mins. The calretinin samples were incubated in ImmPRESS

horseradish peroxidase polymer detection kit for 30mins (Vector Laboratories, US). Samples were

developed in a 3,3'-diaminobenzidine (DAB) substrate kit (Vector Laboratories, US), counterstained

using QS haematoxylin (Vector Laboratories, US), then dehydrated in an ascending series of alcohols,

cleared in xylene and coverslipped using DPX mounting media. The Aβ 42/40 double-labelled

samples were incubated in ImmPRESS horseradish peroxidase polymer then ImmPRESS alkaline

phosphate polymer kits for 30 mins each (Vector Laboratories, US). For these sections, Aβ40 was

visualized using StayYellow HRP chromogen (ab169561, Abcam,UK) followed by Aβ42 visualization

using StayBlue AP chromogen (ab176915, Abcam, UK), counterstained with haematoxylin, air dried

and coverslipped using aqueous mounting media. For each case, one section was processed for each

immunohistochemical marker except for one 70-year-old control case where tissue was not stained

for calbindin and Tau AT8 due to insufficient samples.

Table 1. Primary antibodies used for immunohistochemistry.

Antigen	Antibody	Dilution	Detection	Source
Aβ, 17-24	Mouse anti-4G8 (monoclonal)	1:1000	Streptavidin/DAB	BioLegend 800708
Aβ, 1-40	Rabbit anti-β1-40 (polyclonal)	1:200	ImmPress polymer horse-anti-rabbit HRP/StayYellow	Merck, AB5074P
Aβ, 1-42	Mouse anti-124F (monoclonal)	1:500	ImmPress polymer horse-anti-mouse AP/StayBlue	BioLegend, 805509
Tau (pSer202/ PThr205)	Rabbit anti-AT8/AH36 (monoclonal)	1:500	Streptavidin/DAB	StressMarq Biosciences, SMC-601.
Parvalbumin	Recombinant rabbit anti-parvalbumin (monoclonal)	1:500	Streptavidin/DAB	Abcam, ab181086
Calretinin	Goat anti-calretinin (polyclonal)	1:100	ImmPress polymer horse-anti-goat HRP/DAB	Novus, AF5065
Calbindin	Chicken anti-calbindin D-18K (polyclonal)	1:250	Streptavidin/DAB	Novus, NBP2-50028

Imaging and analysis

Images of the anterior thalamus were obtained using an automated slide scanner (Olympus VS200,

Olympus Life Science, Japan) equipped with a color iDS camera (Olympus VS-264C). The anterior

thalamus was imaged using a 40x objective (NA 0.95) for Aβ-4G8, Tau AT8, hematoxylin, calretinin,

parvalbumin and calbindin and a 20x objective (NA 0.8) for Aβ 40/42. The images were loaded into

QuPath (v0.4.3, Bankhead et al., 2017) and the anterior ventral thalamic nuclei was annotated as

previously described (Perry et al., 2019). The percentage area of the anteroventral thalamic nucleus

occupied by Aβ-4G8, Aβ 40/42, and Tau AT8 was calculated by training a pixel classification model to

identify the relative area occupied by each marker.

For neurons and glia cells, the StarDist deep-learning model (Schmidt et al., 2018) was used to

identify the nuclei of each object in sections stained with hematoxylin alone. An object classifier was

trained to identify each object as either a neuron, astrocyte, oligodendrocyte, or microglia. The

classifier employed several measures including area, length, circularity, diameter, hematoxylin

intensity and a 100µm weighted smoothing for each measure. Similarly, StarDist was used to detect

calretinin, parvalbumin, and calbindin positive neurons in DAB and hematoxylin counter-stained

sections. To do this, the DAB and hematoxylin channels were extracted and summed into a single

detection channel. Positive neurons were identified by filtering detections to a minimum nucleus

area, circularity, DAB and hematoxylin intensity. To provide an indication of the performance of each

cell classifier, one expert manually annotated a 1,000,000 $µm^2$ area on one section for hematoxylin

stained neurons and glia and a 4,000,000$µm^2$ area on one section per calcium binding protein

marker. The manually annotated sections were regarded as the ground truth and were used to

compare the accuracy, precision, and recall of each cell classifier. Each cell classifier performed well,

with high accuracy (range: 89-98%), precision (range 89-97%), and recall (range: 89-97%) achieved

(see Additional File 1: Supplementary Table 2).

Neuron and glia counts could not be obtained from DAB-stained sections that were counterstained

with hematoxylin due to the background staining from DAB. As such, neuron and glia measurements

were only acquired from the hemaotxylin-only stained sections (one section per case). The numbers

of microglia were added to the total glia density and not analysed alone due to low numbers.

Calbindin, calretinin and parvalbumin counts were all derived from separate brain DAB-stained

sections, and again from one section per case. For neuron and glial counts, a measure of the total

number of items was derived within the boundaries of the anteroventral thalamic nuclei on each

section, these measures were then adjusted for the area of the anteroventral thalamic nucleus on

the section, providing a measure of objects per mm^2. Calbindin, calretinin and parvalbumin counts

were further adjusted for neuronal density for each case, to provide an approximate proportion of

immunopositive neurons.

One 40x annotated image, selected from the middle of the available consecutive series of sections

(i.e., the largest available region for each case) was used to provide an approximate estimate of the

cross-sectional area of the anteroventral thalamic nucleus (AV). Each section was coded as either

anterior or posterior AV, as the anteroposterior levels varied across samples.

Data Analysis

Statistical analyses were performed in JASP version 0.18.2 (JASP Team, 2024). An independent means

t-test was used to compare the age of controls and individuals with Down syndrome. The distribution

of gender across the groups was assessed using a Chi-squared test. A 2 x 2 factorial ANOVA was used

to assess the cross-sectional area of the AV with Diagnosis (Control vs Down syndrome) as the

between groups factor and the Anterior-posterior level (anterior vs posterior) of the AV as the within

groups factor. Comparisons between histological markers across groups were carried out using

independent means *t*-tests. One-tailed tests were used for Aβ pathology markers while two-tailed

test were used for all other measures. Welch's *t*-test for unequal variances was used wherever

appropriate. A Mann-Whitney *U* test was used when normality tests failed. Wherever relevant, effect

sizes between groups were expressed as Hedges *g* when equal variances were assumed, Glass's Δ

when equal variances were not assumed, rank-biserial correlation following a Mann-Whitney *U* test,

or η_p^2 following factorial ANOVA. Separate multiple regressions with backward elimination ($p < 0.1$

retention) were modelled with neuronal density, calretinin, parvalbumin, and calbindin proportions

as the dependent variables, and Aβ42, diagnosis, and age as the co-variates. Among the Aβ

pathology markers stained in this study, Aβ42 was selected as a predictor for the regression model as

it is thought to be the more pathogenic form of Aβ (Bodani et al., 2015, Iwatsubo et al., 1995). Age

was included in the model due to the association between advancing age and AD pathology in Down

syndrome (Davidson et al., 2018). Tau was not included due low expression. The most parsimonious

model with the best fit is reported. Additional details of regression summaries, ANOVAs, and

coefficients for each model are included in Additional File 2: Supplementary tables 3 -14. Multiple

comparisons were corrected with the Benjamini-Hochberg procedure (Q = 0.05). The corrected p

values (p_c) of less than 0.05 were considered significant. Given the small numbers, correlations with

age were not carried out but the data are presented in scatterplots so that age and gender

distribution across groups can be visualized.

In addition to significance testing, the Bayesian t-test framework was applied to the key outcome

measures: proportion of calretinin, parvalbumin, and calbindin positive neurons. Bayesian Mann-

Whitney U test, with Markov Monte Carlo set to five chains of 10, 000 iterations, was used when

normality was not assumed. The standardised effect size (δ) was assigned a Cauchy prior distribution

with r = 1/√2. For each test, the Bayes factor (BF_{10}), was calculated to quantify the relative evidence

for the null hypothesis (H_0) compared to the alternative hypothesis (H_1). BF_{10} provides a continuous

measure of evidence where a BF_{10} between 1 and 3 is considered anecdotal evidence, between 3

and 10 is considered moderate evidence and BF_{10} greater than 10 is considered strong evidence, in

favour of the alternative hypothesis, respectively. In comparison, a BF_{10} between 1 and 0.33 is

considered anecdotal evidence, between 0.33 and 0.1 is considered moderate evidence and lower

than 0.1 can be interpreted as strong evidence, in favour of the null hypothesis, respectively (van

Doorn et al., 2021). Here it is important to note that verbal labels used to categorise different Bayes

factors can be useful to facilitate scientific communication, but caution is needed due to the arbitrary

nature of these labels and the continuous nature of the Bayes factor. The Bayesian t test and

Bayesian Mann-Whitney U test posterior and prior plots, along with robustness checks where

applicable are included in Additional File 3: Supplementary figures 1 -3. Finally, we applied the

Bayesian framework to the calcium binding protein regression analyses as described above. In

addition to providing the BF_{10} for the most parsimonious model, this framework also quantifies the

relative evidence for the inclusion ($BF_{inclusion}$) of each co-variate in the model. A beta prior probability

(a = b =1) was applied to the models and a JZS prior (r scale = 0.354) was applied to the regression

coefficients. Additional details of Bayesian regression model comparisons, posterior summaries of

coefficients, and descriptive statistics are included in Additional File 4: Supplementary tables 15 – 23.

Figures were produced in R using ggplot2 (Wickham, 2016).

Results

Group demographics

The control group tissue came from a slightly older population (range 38-70 years) than the Down

syndrome (DS) group tissue (range 22-62 years), although the difference did not reach significance

(Table 2). There was no difference in gender distribution across the two groups (see Table 2).

Table 2. Demographic data

	Control group n=12	Down syndrome group n=15	t value/χ^2	p value
Mean age ± S.D.	57.4 ± 11.2 years	47.8 ± 13.1 years	2.02	0.06
Sex (M:F)	6:6	9:6	0.27	0.60

Given the sample size and distribution of data, we did not look at specific correlations between

histological measures and age. While there were no effects of gender, due to the small numbers it is

possible that more subtle gender effects may not be detected, therefore, age and gender

distributions are presented graphically for the following histological measures.

Regional area of anteroventral thalamic nucleus

While we did not have the tissue necessary to derive volumetric measurements for the anteroventral

thalamic nucleus (AV), we did estimate the cross-sectional regional area from one mid-section per

case (Fig. 1a, b). We found that the area of AV was approximately 37% smaller in the DS group

compared to controls and this difference was significant as shown by a main effect of Diagnosis (F(1,

23) = 7.06, p_c = 0.023, η_p^2 = 0.24, Fig. 1c). The smaller AV area in the DS group was not due to

differences in the anterior-posterior levels between the samples as neither the main effect of

Anterior-posterior level (F(1, 23) = 0.59, p_c = 0.45, η_p^2 = 0.03) nor the Diagnosis x Anterior-posterior

level interaction were significant (F(1, 23) = 4.01, p_c = 0.07, η_p^2 = 0.15).

Fig 1. Cross-sectional area of the anteroventral thalamic nucleus. Representative examples of the

anteroventral thalamic nucleus from similar middle anterior – posterior levels in (a) 63yo male

control (case# BBN110.35427) and (b) 60yo female with Down syndrome (case# BBN_2990), and (c)

box plot of the cross-sectional area of the anteroventral thalamic nuclei. The box indicates the inter-

quartile range (IQR), the whiskers show the range of values that are within 1.5 x IQR, the horizontal

line indicates the median, and the cross indicates the mean. The notches represent the 95%

confidence interval for each median (1.58 x IQR/√(n)). Control, control group; DS, Down syndrome

group, F, females, M, males; *p<0.05, **p<0.01, ***p<0.001. Scale bar = 1mm. Note: The examples

of the anteroventral thalamic nucleus in (a) and (b) are both oriented with dorsal towards the top of

the image. The orientation of the anteroventral nucleus can vary between individuals at similar

anterior – posterior levels.

Neurons and glia

There was increased neuron density in the DS group (t(16.482) = -2.27, p_c = 0.049, Δ = -0.61, 95% CI:

[-1.39, 0.19], Fig. 2a, e, I), however, the range in neuron density was much greater in the DS group

(44-288 cells/mm^2) than in the control group (74-129 cells/mm^2) suggesting a more heterogeneous

population. While the density of total glia was slightly higher in the DS group, this difference was not

significant (t(25) = -1.13, p_c = 0.30, g = -0.423, 95% CI: [-1.19, 0.35], Fig. 2b, f, j). The numerical

increase in total glia density was driven by a significant increase in astroglia density (t(25) = -2.33, p_c

= 0.04, g = -0.88, 95% CI: [-1.67, -0.07], Fig. 2c, g, k), but not, oligodendrocyte density (t(25) = 0.81, p_c

= 0.45, g = 0.30, 95% CI: [-0.46, 1.06], Fig. 2d, h, l), which was comparable across the groups.

Fig 2. Neuron and glia density in the anteroventral thalamic nucleus. Top panel, hematoxylin stained examples of (a) a neuron with clearly stained nucleolus, nucleus, and surrounded by cytoplasm, (b) examples of astroglia (yellow arrows), oligodendrocytes (black arrows), comma shaped microglia (red arrow), and neurons (white arrow) and zoomed in examples of (c) astroglia showing a typical pale nucleus with heterochromatin staining concentrated in granules below the nuclear membrane and (d) oligodendrocytes with a classic rounded nucleus and dense chromatin staining. Middle panel, box plots of the densities of (e) neurons, (f) total glia, (g) astroglia and (h) oligodendrocytes. The box indicates the inter-quartile range (IQR), the whiskers show the range of values that are within 1.5 x IQR, the horizontal line indicates the median, and the cross indicates the mean. The notches represent the 95% confidence interval for each median (1.58xIQR/√(n)). Bottom panel, scatterplot showing the relationship between age and density of (i) neurons, (j) total glia, (k) astroglia and (l) oligodendrocytes. The lines are fitted with the general linear model and the shaded areas are the

95% confidence interval. Control, control group; DS, Down syndrome group, F, females, M, males.

*p<0.05, **p<0.01, ***p<0.001. Note: All neuron and glia examples come from the same 61yo male

control (case# BBN110.28899). Samples were imaged at 40x magnification and zoomed in using

QuPath to show the features of each cell type.

Amyloid and tau

As expected, the area fraction of AV occupied by Aβ was significantly higher in the DS group than the

control group for all three markers: Aβ-4G8 (Fig. 3a), Aβ42 (Fig. 3b) and Aβ40 (Fig. 3b) [Aβ-4G8:

Mann-Whitney U = 15.00, p_c < 0.001, r = -0.83, 95% CI: [-∞, -0.68], Fig. 3d,f; Aβ42: t(14.03) = -4.32,

p_c = 0.002, Δ = -1.12, 95% CI: [-∞, -0.38], Fig. 3e, f; Aβ40: Mann-Whitney U = 49.50, p_c = 0.02, r = -

0.45, 95% CI: [-∞, -0.11]]. While there was a significant group difference for Aβ40, expression was

generally low with levels near zero for 13 of the 15 tissue samples from the DS group. However,

moderate expression was found in one case (47-year-old female, 1.14% coverage), and high

expression was found in one case (47-year-old male, 7.24% coverage).

Tau AT8 expression in the AV was generally low and did not differ between the groups (p = 0.36,

uncorrected). Tau AT8 expression was near zero in 12 of the 15 tissue samples from individuals with

DS. However, moderate expression was found in one case (65-year-old male: 1.7% coverage), and

high expression was found in two cases (47-year-old female, 9.1% coverage; 60-year-old male, 6.5%

coverage; Fig. 3c).

Fig 3. Alzheimer's disease pathology in the anteroventral thalamic nucleus. Top panel, examples of

(a) Aβ-4G8 plaques, (b) blue stained Aβ42 (white arrow) and yellow stained Aβ40 (black arrow)

plaques taken from the same area of the anteroventral thalamic nucleus from a 47yo female with

Down syndrome (case# BBN_17186) and (c) Tau AT8 positive neurofibrillary tangles (white arrows)

and neuropil threads (black arrows) taken from a 65yo male with Down syndrome (case# BBN_2975).

Middle panel, box plots of the percent area fraction occupied by (d) Aβ 4G8 and (e) Aβ42. The box

indicates the inter-quartile range (IQR), the whiskers show the range of values that are within

1.5xIQR, the horizontal line indicates the median, and the cross indicates the mean. The notches

represent the 95% confidence interval for each median (1.58xIQR/√(n)). Bottom panel, scatterplot

showing the relationship between age and the percent area fraction occupied by (f) Aβ 4G8 and (g)

Aβ42. The lines are fitted with the general linear model and the shaded areas are the 95%

confidence interval. Control, control group, DS, Down syndrome group; F, females, M, males. Scale

bar = 20 μm. *p<0.05, **p<0.01, ***p<0.001.

Calcium-binding proteins (calretinin, calbindin, parvalbumin)

Expression of each calcium-binding protein was normalized against the neuronal density for each

individual case, to provide a relative proportion of immunopositive neurons. The proportion of all

three calcium-binding proteins in AV was significantly reduced in the DS group [calretinin, $t(25)$ =

3.23, p_c = 0.008, g = 1.21, 95% CI: [0.37, 2.03], Fig. 4a, d, g; parvalbumin, Mann-Whitney U = 149, p_c =

0.009, r = 0.66, 95% CI: [0.33, 0.84], Fig. 4b, e, h; calbindin, Mann-Whitney U = 156, p_c < 0.001, r =

0.89, 95% CI: [0.75, 0.96], Fig. 4c, f, l]. The same pattern of results was found when using the density

of neurons immunopositive for calcium-binding proteins in the AV, rather than proportion [calretinin,

$t(25)$ = 2.61, p_c = 0.023, g = 0.98, 95% CI: [0.16, 1.78]; parvalbumin, Mann-Whitney U = 144, p_c =

0.018, r = 0.60, 95% CI: [0.24, 0.81]; calbindin, Mann-Whitney U = 140, p_c = 0.006, r = 0.70, 95% CI:

[0.38, 0.87]].

Subsequent Bayesian analyses indicated strong evidence in favour of reduced calretinin and calbindin

proportions (BF_{10} = 11.41 and 19.43, respectively) and moderate evidence in favour of reduced

parvalbumin proportion (BF_{10} = 5.36), over a null hypothesis of no difference in proportions between

controls and individuals with Down syndrome.

Fig 4. Calcium-binding proteins in the anteroventral thalamic nucleus. Top panel, representative

examples of (a) calretinin from a 66yo female control (case# BBN_24381), (b) parvalbumin from a

60yo female control (case# BBN110.28668), and (c) calbindin positive neurons from a 61yo female

control (case# BBN110.28899). Middle panel, box plots of normalized densities of (d) calretinin, (e)

parvalbumin, and (f) calbindin positive neurons. The box indicates the inter-quartile range (IQR), the

whiskers show the range of values that are within 1.5xIQR, the horizontal line indicates the median,

and the cross indicates the mean. The notches represent the 95% confidence interval for each

median (1.58xIQR/√(n)). Bottom panel, scatterplot showing the relationship between age and

normalized densities of (g) calretinin, (h) parvalbumin, and (i) calbindin positive neurons. The lines

are fitted with the general linear model and the shaded areas are the 95% confidence interval.

Control, control group; DS, Down syndrome group; F, females, M, males. Scale bar = 20 μm. *p<0.05,

**p<0.01, ***p<0.001.

Multiple Regression

Backward regression analysis identified a model containing diagnosis and age as the best model

predicting normalised calretinin expression (adjusted R^2 = 0.36, $F(2,24)$ = 8.20, p_c = 0.006). In

contrast, models containing only diagnosis were the best at predicting parvalbumin (adjusted R^2 =

0.29, $F(1, 25)$ = 11.71, p_c = 0.006) and calbindin expression (adjusted R^2 = 0.47, $F(1, 24)$ = 23.16, p_c <

0.001). A model containing Aβ42 neared significance as the best predictor of increasing neuronal

density (adjusted R^2 = 0.12, $F(1,25)$ = 4.45, p_c = 0.056).

Bayesian regression analyses were carried out using diagnosis, age, and Aβ42 as predictors of

normalized calretinin, parvalbumin, and calbindin expression. For calretinin, there was strong

evidence that the model with diagnosis and age explained a large amount of the variance, R^2 = 0.41,

(BF$_{10}$ = 21.47). Diagnosis provided moderate evidence ($β_1$ =-0.17, BF$_{inclusion}$ = 6.25) while Age ($β_2$ = -

0.003, BF$_{inclusion}$ = 1.84) and Aβ42 ($β_3$ = -0.012, BF$_{inclusion}$ = 1.84) provided only anecdotal evidence for

inclusion in the model. For parvalbumin, there was strong evidence that the model with diagnosis

alone explained a large amount of variance R^2 = 0.32 (BF$_{10}$ = 16.70). Diagnosis provided moderate

evidence for inclusion in the model ($β_1$ =-0.21, BF$_{inclusion}$ = 7.11). In contrast, both age ($β_2$ =-0.00036,

BF$_{inclusion}$ = 0.57) and Aβ42 ($β_3$ =0.003, BF$_{inclusion}$ = 0.59) provided anecdotal evidence for exclusion

from the model. Finally, for calbindin, there was strong evidence that the model with diagnosis alone

explained a large amount of variance R^2 = 0.49 (BF$_{10}$ = 290). Diagnosis provided strong evidence for

inclusion in the model ($β_1$ =-0.29, BF$_{inclusion}$ = 25.85). In contrast, both age ($β_2$ =-0.00094, BF$_{inclusion}$ =

0.57) and Aβ42 (β_3 =-0.014, $BF_{inclusion}$ = 0.57) provided anecdotal evidence for exclusion from the

model.

Discussion

Despite the longstanding association between the medial diencephalon and memory, there has been

surprisingly little research into the status of this brain region in individuals with Down syndrome. In

the current study we assessed the anteroventral thalamic nucleus (AV) in post-mortem tissue from

15 individuals with Down syndrome (DS group). In addition to measuring surface area and neuronal

density, we also measured the expression of calcium-binding proteins which play an important role

in calcium homeostasis and intracellular signalling and have been linked to neurodegeneration

(Kelemen and Szilágyi, 2021).

As expected, we found higher levels of amyloid markers in the DS group, most notably in total Aβ

burden (as measured by Aβ-4G8) and diffuse Aβ plaques (as measured by Aβ42 expression).

Expression of Aβ40-positive mature plaques was generally low with near zero expression in 13 of 15

samples from individuals with Down syndrome. These findings are in line with previous reports

showing that Aβ42 plaques are more prevalent than Aβ40 plaques across all ages in Down syndrome

(Head et al., 2016). Previous studies have found less pronounced tau expression in the anterior

thalamic nuclei in Alzheimer's disease than other regions (Rüb et al., 2016, Braak and Braak, 1991,

Viney et al., 2022). Consistent with this, there was very little tau pathology in most of the cases we

examined across groups with were near zero levels of tau in 12 of the 15 cases from the DS group.

While there were too few cases with high tau expression to draw any clear conclusions, there did not

seem to be an association with age as the individual with the highest tau levels was 49 years old, but

that case also had the lowest neuron density, suggesting neurons may be particularly vulnerable to

high levels of tau in the AV. The heterogeneity in both tau and amyloid expression in younger cases

suggests some individuals may have better protection against earlier Alzheimer's pathology (Maxwell

et al., 2021).

Astroglial, but not oligodendrocyte, density was significantly increased in the AV in the DS group. In

addition, there was an increase in neuron density in the AV of individuals with Down syndrome,

although there was far greater variability in the DS group with some showing very low neuron

density and others showing increased neuron density. The overall pattern of increased neuronal and

glial density may appear to contradict our previous findings (Perry et al., 2019), where we found

reduced total neuron and glia counts. However, the increased neuron density in the current study

likely reflects increased packing due to a reduction in AV volume. Due to sample availability, we were

not able to acquire total volume measures, as we were in our previous study (Perry et al., 2019), but

we acquired a surface area measure for AV for each case to provide a relative approximation of size.

The surface area of AV was approximately 37% smaller in the DS group than controls, therefore, even

with preserved or increased neuronal density, there are likely fewer neurons overall in the AV in the

DS group. Only a few studies have quantitatively assessed the anterior thalamic nuclei in Alzheimer's

disease (De Simone et al., 2024) and even fewer have carried out neuronal counts but, from the

studies that are available, there does not seem to be a significant reduction in either volume or

neuronal counts in this region using post-mortem tissue (Hornberger et al., 2012, Xuereb et al.,

1991), suggesting that the AV may be more impacted in Down syndrome than Alzheimer's disease

alone.

A further aim of the current study was to assess expression of calcium-binding proteins in the AV in

Down syndrome. The proportion of AV neurons immunopositive for calbindin, calretinin and

parvalbumin was significantly reduced in the DS group, by 70%, 49% and 61%, respectively.

Furthermore, we found strong evidence in favour of reduced proportions of calretinin and calbindin

positive cells and moderate evidence in favour of reduced proportions of parvalbumin positive cells,

according to the Bayesian framework. A previous study had reported a reduction in both calbindin

and parvalbumin in the cortices of four older individuals with Down syndrome (Kobayashi et al.,

1990). For calbindin, they found a 38% and 28% reduction in immunoreactive neurons in frontal and

temporal cortices, respectively. The reported reduction in parvalbumin was slightly less than

calbindin, with a 27% decrease in frontal cortex and 22% decrease in temporal cortex. Consistent

with these cortical findings (Kobayashi et al., 1990), we also found a greater decrease in calbindin

than parvalbumin. However, the relative reduction for both parvalbumin and calbindin appears

greater in AV than in the cortex, especially considering our sample population also included younger

individuals.

Parvalbumin expression has been previously analysed in human AV. One study found 53-58% of total

neurons to be immunoreactive for parvalbumin (Dixon et al., 2000) while another found the

percentage of parvalbumin positive cells to range from 55-63% (Danos et al., 1998). In our study, we

found approximately 45% of cells to be immunoreactive for parvalbumin in the control group. The

methodology and samples are very different across studies but in seems a consistent finding that

approximately half the neurons in AV are immunoreactive for parvalbumin.

Reductions in parvalbumin, calretinin, and calbindin expression have been reported in Alzheimer's

disease (Zallo et al., 2018, Brady and Mufson, 1997, Kook et al., 2014b), however, these studies have

typically focused on the hippocampus and frontal cortex; to our knowledge, no study has specifically

assessed these markers in the anterior thalamic nuclei in Alzheimer's disease. It is, therefore, not

clear whether the reduction in calcium-binding proteins in the DS group simply reflects a sensitivity

to Alzheimer-related pathology or whether there are underlying changes to the cell populations in

Down syndrome that co-occur with, or even pre-date, amyloid changes. There did not seem to be a

clear relationship between amyloid or tau and calcium-binding protein expression but there were

cases with very low amyloid and tau expression that also had very low levels of calbindin,

parvalbumin and calretinin expression, consistent with the calcium-binding protein changes being

found irrespective of amyloid status. In support of this, according to our most parsimonious

frequentist and Bayesian regression models, Aβ42, the most pathological form of these Aβ species

(Bodani et al., 2015, Iwatsubo et al., 1995), did not predict expression of any calcium-binding protein.

Down syndrome alone was the best predictor of calcium-binding protein expression. In comparison,

advancing age and Aβ42 deposition provided only some evidence in favour of predicting calretinin

expression, but some evidence to be excluded as predictors of parvalbumin and calbindin expression.

While there does not seem to be a clear association between Alzheimer markers and calcium-binding

protein expression in this current data set, it is possible that earlier changes, for example Aβ and/or

tau seeding (Ye et al., 2017, Condello et al., 2022), may show clearer links. A reduction in calcium-

binding proteins would disrupt calcium homeostasis, and a loss of calbindin in a rodent model of

Alzheimer's disease was linked to increased pathogenesis (Kook et al., 2014a). It is, therefore,

possible that an early loss of calbindin could exacerbate Alzheimer-related pathology in Down

syndrome, given calbindin shows the greatest reduction in the present study.

While there are few studies on the thalamus in Down syndrome, a recent study assessed three

thalamic nuclei in tissue from foetuses with Down syndrome: mediodorsal nucleus, centromedian

nucleus and parafasciular nucleus (Stagni et al., 2020). A similar pattern was found across all nuclei,

with a reduction in neuron and glia density but no difference in surface area, consistent with an

overall reduction in total neurons in these thalamic regions. There was high expression of calretinin

in the mediodorsal thalamic nucleus, with approximately 80% of cells staining positive for calretinin

but there was no difference in expression between the control and Down syndrome group (Stagni et

al., 2020). This might suggest that expression levels are generally comparable across individuals with

Down syndrome and controls in early life, with a loss only occurring with increasing age.

Alternatively, there may be variability across thalamic nuclei and the age at which changes occur,

which might be expected given the different expression profiles and functions across thalamic nuclei

(Stagni et al., 2020, Morel et al., 1997, Münkle et al., 2000, Wolff and Vann, 2019). This reinforces the

importance of future studies into the anterior thalamic nuclei in earlier age groups to identify

whether there is a reduction in calcium-binding protein in early life or if a loss of expression only

occurs after adolescence. If calcium-binding protein expression reduces after adolescence, then

finding ways to preserve the expression of these proteins may help protect against subsequent

pathogenesis (Hijazi et al., 2020).

One of the limitations of the study is that we used digital pathology methods to identify and quantify

cells. This method is suitable for identifying relative numbers between groups but cannot estimate

the true number of cells in these samples. This can only be achieved with stereological assessment

which requires all of tissue from the region of interest. Much of the tissue from current study is

historical, with portions of the anterior thalamus having been used in other studies. However,

despite the lack of stereological assessments, the data still provides valuable insights to changes in

the anterior thalamus of younger and older adults with Down syndrome given the rarity of this tissue

and the lack of knowledge of the anterior thalamus in Down syndrome.

An additional limitation is that the different stains were carried out on adjacent tissue sections for

each individual which could add to some natural variation in staining levels and tissue quality and we

were not able to assess colocalization of staining. Furthermore, there was also variation in the

anteroposterior level assessed across cases. However, the sections for both the DS and control

groups both came from a similar range of anteroposterior levels giving additional weight to the

findings. But this again highlights the importance of future stereological studies into calcium-binding

protein expression within the AV nucleus both in control and patient populations.

The present study adds to the growing literature linking neuropathology in the medial diencephalon,

including anterior thalamic nuclei and mammillary bodies, to various neurological disorders

(Milczarek et al., 2023a, Lequin et al., 2021, Meys et al., 2022, Clarke et al., 1994, Connaughton et al.,

2023, Danos et al., 1998). Pathology in these brain areas would contribute to memory impairments

often accompanying these conditions by affecting both the encoding and consolidation of memories

(Milczarek et al., 2023b, Aggleton et al., 2023). In the present study, we found changes in the AV in

Down syndrome at a younger age than previously reported and showed for the first time that there

is a marked loss of expression of calcium-binding proteins. Noticeably, this reduction in parvalbumin,

calretinin and calbindin expression was not simply driven by overall neuronal loss. These reductions

in calcium-binding proteins may increase the vulnerability of cells within this region and potentially

exacerbate expression of Alzheimer-related pathology.

References

Aggleton, J. P., Vann, S. D. & O'mara, S. M. 2023. Converging diencephalic and hippocampal supports
for episodic memory. *Neuropsychologia,* 191**,** 108728.
Bankhead, P., Loughrey, M. B., Fernández, J. A., Dombrowski, Y., Mcart, D. G., Dunne, P. D., Mcquaid,
S., Gray, R. T., Murray, L. J., Coleman, H. G., James, J. A., Salto-Tellez, M. & Hamilton, P. W.
2017. QuPath: Open source software for digital pathology image analysis. *Scientific Reports,*
7**,** 16878.
Bodani, R. U., Sengupta, U., Castillo-Carranza, D. L., Guerrero-Muñoz, M. J., Gerson, J. E., Rudra, J. &
Kayed, R. 2015. Antibody against Small Aggregated Peptide Specifically Recognizes Toxic Aβ-
42 Oligomers in Alzheimer's Disease. *ACS Chem Neurosci,* 6**,** 1981-9.
Braak, H. & Braak, E. 1991. Alzheimer's disease affects limbic nuclei of the thalamus. *Acta*
Neuropathol, 81**,** 261-8.
Brady, D. R. & Mufson, E. J. 1997. Parvalbumin-immunoreactive neurons in the hippocampal
formation of Alzheimer's diseased brain. *Neuroscience,* 80**,** 1113-1125.
Carducci, F., Onorati, P., Condoluci, C., Di Gennaro, G., Quarato, P. P., Pierallini, A., Sara, M., Miano, S.,
Cornia, R. & Albertini, G. 2013. Whole-brain voxel-based morphometry study of children and
adolescents with Down syndrome. *Funct Neurol,* 28**,** 19-28.
Clarke, S., Assal, G., Bogousslavsky, J., Regli, F., Townsend, D. W., Leenders, K. L. & Blecic, S. 1994.
Pure amnesia after unilateral left polar thalamic infarct: topographic and sequential
neuropsychological and metabolic (PET) correlations. *J Neurol Neurosurg Psychiatry,* 57**,** 27-
34.
Condello, C., Maxwell, A. M., Castillo, E., Aoyagi, A., Graff, C., Ingelsson, M., Lannfelt, L., Bird, T. D.,
Keene, C. D., Seeley, W. W., Perl, D. P., Head, E. & Prusiner, S. B. 2022. Aβ and tau prions
feature in the neuropathogenesis of Down syndrome. *Proc Natl Acad Sci U S A,* 119**,**
e2212954119.
Connaughton, M., O'hanlon, E., Silk, T. J., Paterson, J., O'neill, A., Anderson, V., Whelan, R. & Mcgrath,
J. 2023. The Limbic System in Children and Adolescents with Attention-Deficit/Hyperactivity
Disorder: A Longitudinal Structural MRI Analysis. *Biological Psychiatry Global Open Science*.
Danos, P., Baumann, B., Bernstein, H.-G., Franz, M., Stauch, R., Northoff, G., Krell, D., Falkai, P. &
Bogerts, B. 1998. Schizophrenia and anteroventral thalamic nucleus: selective decrease of
parvalbumin-immunoreactive thalamocortical projection neurons. *Psychiatry Research:*
Neuroimaging, 82**,** 1-10.

Davidson, Y. S., Robinson, A., Prasher, V. P. & Mann, D. M. A. 2018. The age of onset and evolution of
Braak tangle stage and Thal amyloid pathology of Alzheimer's disease in individuals with
Down syndrome. *Acta Neuropathol Commun,* 6**,** 56.
De Simone, M. S., Spalletta, G., Vecchio, D., Bassi, A., Carlesimo, G. A. & Piras, F. 2024. The Role of the
Anterior Thalamic Nuclei in the Genesis of Memory Disorders in Alzheimer's Disease: An
Exploratory Study. *J Alzheimers Dis,* 97**,** 507-519.
Dixon, G., Dissanaike, S. & Harper, C. G. 2000. Parvalbumin-immunoreactive neurons in the human
anteroventral thalamic nucleus. *Neuroreport,* 11**,** 97-101.
Fairless, R., Williams, S. K. & Diem, R. 2019. Calcium-Binding Proteins as Determinants of Central
Nervous System Neuronal Vulnerability to Disease. *Int J Mol Sci,* 20.
Guan, P. P., Cao, L. L. & Wang, P. 2021. Elevating the Levels of Calcium Ions Exacerbate Alzheimer's
Disease via Inducing the Production and Aggregation of β-Amyloid Protein and
Phosphorylated Tau. *Int J Mol Sci,* 22.
Head, E., Lott, I. T., Hof, P. R., Bouras, C., Su, J. H., Kim, R., Haier, R. & Cotman, C. W. 2003. Parallel
Compensatory and Pathological Events Associated with Tau Pathology in Middle Aged
Individuals with Down Syndrome. *Journal of Neuropathology & Experimental Neurology,* 62**,**
917-926.
Head, E., Lott, I. T., Wilcock, D. M. & Lemere, C. A. 2016. Aging in Down Syndrome and the
Development of Alzheimer's Disease Neuropathology. *Curr Alzheimer Res,* 13**,** 18-29.
Head, E., Powell, D., Gold, B. T. & Schmitt, F. A. 2012. Alzheimer's Disease in Down Syndrome. *Eur J*
Neurodegener Dis, 1**,** 353-364.
Hijazi, S., Heistek, T. S., Scheltens, P., Neumann, U., Shimshek, D. R., Mansvelder, H. D., Smit, A. B. &
Van Kesteren, R. E. 2020. Early restoration of parvalbumin interneuron activity prevents
memory loss and network hyperexcitability in a mouse model of Alzheimer's disease. *Mol*
Psychiatry, 25**,** 3380-3398.
Hornberger, M., Wong, S., Tan, R., Irish, M., Piguet, O., Kril, J., Hodges, J. R. & Halliday, G. 2012. In
vivo and post-mortem memory circuit integrity in frontotemporal dementia and Alzheimer's
disease. *Brain,* 135**,** 3015-25.
Iwatsubo, T., Mann, D. M., Odaka, A., Suzuki, N. & Ihara, Y. 1995. Amyloid beta protein (A beta)
deposition: A beta 42(43) precedes A beta 40 in Down syndrome. *Ann Neurol,* 37**,** 294-9.
Kelemen, K. & Szilágyi, T. 2021. New Approach for Untangling the Role of Uncommon Calcium-
Binding Proteins in the Central Nervous System. *Brain Sci,* 11.
Kemper , T. L. 1991. Down Syndrome. *In:* PETERS, A. & JONES, E. G. (eds.) *Cerebral Cortex: Normal*
and altered states of function. Plenum Press.
Kobayashi, K., Emson, P. C., Mountjoy, C. Q., Thornton, S. N., Lawson, D. E. & Mann, D. M. 1990.
Cerebral cortical calbindin D28K and parvalbumin neurones in Down's syndrome. *Neurosci*
Lett, 113**,** 17-22.
Kook, S. Y., Jeong, H., Kang, M. J., Park, R., Shin, H. J., Han, S. H., Son, S. M., Song, H., Baik, S. H.,
Moon, M., Yi, E. C., Hwang, D. & Mook-Jung, I. 2014a. Crucial role of calbindin-D28k in the
pathogenesis of Alzheimer's disease mouse model. *Cell Death Differ,* 21**,** 1575-87.
Kook, S. Y., Jeong, H., Kang, M. J., Park, R., Shin, H. J., Han, S. H., Son, S. M., Song, H., Baik, S. H.,
Moon, M., Yi, E. C., Hwang, D. & Mook-Jung, I. 2014b. Crucial role of calbindin-D28k in the
pathogenesis of Alzheimer's disease mouse model. *Cell Death & Differentiation,* 21**,** 1575-
1587.
Lemere, C. A., Blusztajn, J. K., Yamaguchi, H., Wisniewski, T., Saido, T. C. & Selkoe, D. J. 1996.
Sequence of deposition of heterogeneous amyloid beta-peptides and APO E in Down
syndrome: implications for initial events in amyloid plaque formation. *Neurobiol Dis,* 3**,** 16-
32.
Lequin, M. H., Steggerda, S. J., Severino, M., Tortora, D., Parodi, A., Ramenghi, L. A., Groenendaal, F.,
Meys, K. M. E., Benders, M., De Vries, L. S. & Vann, S. D. 2021. Mammillary body injury in
neonatal encephalopathy: a multicentre, retrospective study. *Pediatr Res.*

Mandler, M., Walker, L., Santic, R., Hanson, P., Upadhaya, A. R., Colloby, S. J., Morris, C. M., Thal, D.
R., Thomas, A. J., Schneeberger, A. & Attems, J. 2014. Pyroglutamylated amyloid-β is
associated with hyperphosphorylated tau and severity of Alzheimer's disease. *Acta*
Neuropathol, 128**,** 67-79.
Maxwell, A. M., Yuan, P., Rivera, B. M., Schaaf, W., Mladinov, M., Prasher, V. P., Robinson, A. C.,
Degrado, W. F. & Condello, C. 2021. Emergence of distinct and heterogeneous strains of
amyloid beta with advanced Alzheimer's disease pathology in Down syndrome. *Acta*
Neuropathologica Communications, 9**,** 201.
Meys, K. M. E., De Vries, E., Groenendaal, F., Vann, S. & Lequin, M. H. 2022. The mammillary bodies:
a review of causes of injury in infants and children. *AJNR Am J Neuroradiol.*
Milczarek, M. M., Gilani, S. I. A., Lequin, M. H. & Vann, S. D. 2023a. Reduced mammillary body
volume in individuals with a schizophrenia diagnosis: an analysis of the COBRE data set.
Schizophrenia (Heidelb), 9**,** 48.
Milczarek, M. M., Perry, J. C., Amin, E., Haniffa, S., Hathaway, T. & Vann, S. D. 2023b. Impairments in
the early consolidation of spatial memories via group II mGluR agonism in the mammillary
bodies. *bioRxiv***,** 2023.12.13.571542.
Morel, A., Magnin, M. & Jeanmonod, D. 1997. Multiarchitectonic and stereotactic atlas of the human
thalamus. *Journal of Comparative Neurology,* 387**,** 588-630.
Münkle, M. C., Waldvogel, H. J. & Faull, R. L. M. 2000. The distribution of calbindin, calretinin and
parvalbumin immunoreactivity in the human thalamus. *Journal of Chemical Neuroanatomy,*
19**,** 155-173.
Perry, J. C., Pakkenberg, B. & Vann, S. D. 2019. Striking reduction in neurons and glial cells in anterior
thalamic nuclei of older patients with Down syndrome. *Neurobiol Aging,* 75**,** 54-61.
Pinter, J. D., Brown, W. E., Eliez, S., Schmitt, J. E., Capone, G. T. & Reiss, A. L. 2001. Amygdala and
hippocampal volumes in children with Down syndrome: a high-resolution MRI study.
Neurology, 56**,** 972-4.
Rüb, U., Stratmann, K., Heinsen, H., Del Turco, D., Ghebremedhin, E., Seidel, K., Den Dunnen, W. &
Korf, H. W. 2016. Hierarchical Distribution of the Tau Cytoskeletal Pathology in the Thalamus
of Alzheimer's Disease Patients. *J Alzheimers Dis,* 49**,** 905-15.
Satoh, J., Tabira, T., Sano, M., Nakayama, H. & Tateishi, J. 1991. Parvalbumin-immunoreactive neurons
in the human central nervous system are decreased in Alzheimer's disease. *Acta*
Neuropathologica, 81**,** 388-395.
Schmidt, U., Weigert, M., Broaddus, C. & Myers, G. Cell Detection with Star-Convex Polygons. *In:*
FRANGI, A. F., SCHNABEL, J. A., DAVATZIKOS, C., ALBEROLA-LÓPEZ, C. & FICHTINGER, G., eds.
Medical Image Computing and Computer Assisted Intervention – MICCAI 2018, 2018// 2018
Cham. Springer International Publishing, 265-273.
Stagni, F., Giacomini, A., Emili, M., Uguagliati, B., Bonasoni, M. P., Bartesaghi, R. & Guidi, S. 2020.
Neuroanatomical alterations in higher-order thalamic nuclei of fetuses with Down syndrome.
Clinical Neurology and Neurosurgery, 194**,** 105870.
Team, J. 2024. JASP (Version 0.18.2)[Computer software]. https:// jasp-stats.org/
Van Doorn, J., Van Den Bergh, D., Böhm, U., Dablander, F., Derks, K., Draws, T., Etz, A., Evans, N. J.,
Gronau, Q. F., Haaf, J. M., Hinne, M., Kucharský, Š., Ly, A., Marsman, M., Matzke, D., Gupta, A.
R. K. N., Sarafoglou, A., Stefan, A., Voelkel, J. G. & Wagenmakers, E.-J. 2021. The JASP
guidelines for conducting and reporting a Bayesian analysis. *Psychonomic Bulletin & Review,*
28**,** 813-826.
Viney, T. J., Sarkany, B., Ozdemir, A. T., Hartwich, K., Schweimer, J., Bannerman, D. & Somogyi, P. 2022.
Spread of pathological human Tau from neurons to oligodendrocytes and loss of high-firing
pyramidal neurons in aging mice. *Cell Rep,* 41**,** 111646.
Wickham, H. 2016. *ggplot2: Elegant Graphics for Data Analysis,* Springer-Verlag New York.
Wolff, M. & Vann, S. D. 2019. The Cognitive Thalamus as a Gateway to Mental Representations. *J*
Neurosci, 39**,** 3-14.

Xuereb, J. H., Perry, R. H., Candy, J. M., Perry, E. K., Marshall, E. & Bonham, J. R. 1991. Nerve cell loss
in the thalamus in Alzheimer's disease and Parkinson's disease. *Brain,* 114 (Pt 3)**,** 1363-79.
Ye, L., Rasmussen, J., Kaeser, S. A., Marzesco, A. M., Obermüller, U., Mahler, J., Schelle, J., Odenthal,
J., Krüger, C., Fritschi, S. K., Walker, L. C., Staufenbiel, M., Baumann, F. & Jucker, M. 2017. Aβ
seeding potency peaks in the early stages of cerebral β-amyloidosis. *EMBO reports,* 18**,** 1536-
1544-1544.
Zallo, F., Gardenal, E., Verkhratsky, A. & Rodríguez, J. J. 2018. Loss of calretinin and parvalbumin
positive interneurones in the hippocampal CA1 of aged Alzheimer's disease mice.
Neuroscience Letters, 681**,** 19-25.
650
651

Declarations

Ethics approval and consent to participate

Not applicable. Included in methods section.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analysed during the current study available from the corresponding author

on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This research was funded by a grant from the Jerome Lejeune Foundation (# 1899) and a Wellcome

Trust Senior Research Fellowship (WT212273/Z/18) awarded to SDV.

Contributions

JCP and SDV conceived and designed the study. JCP performed the experiments, analysed the data,

and produced the figures. JCP and SDV interpreted the data. JCP and SDV wrote the manuscript.

Acknowledgements

This research was funded by a grant from the Jerome Lejeune Foundation (# 1899) and a Wellcome

Trust Senior Research Fellowship (WT212273/Z/18) awarded to SDV.

Tissue samples were sourced from brain banks in the UK Brain Banks Network. The Manchester Brain

Bank, which is part of the Brains for Dementia Research Initiative, jointly funded by Alzheimer's

Society and Alzheimer's Research UK. The Cambridge Brain Bank is supported by the NIHR

Cambridge Biomedical Research Centre (NIHR203312). We acknowledge the Oxford Brain Bank,

supported by Brains Dementia Research (BDR) (Alzheimer Society and Alzheimer Research UK) and

the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC). Tissue

samples were provided by The London Neurodegenerative Diseases Brain Bank at King's College

London. The brain bank receives funding as part of the Brains for Dementia Research programme,

jointly funded by Alzheimer's Research UK and the Alzheimer's Society.

Authors' information

School of Psychology & Neuroscience and Mental Health Innovation Institute, Cardiff University,

Cardiff, UK.

JCP & SDV

List of abbreviations

AV, anteroventral thalamic nucleus; Aβ, beta-amyloid; BF_{10}, Bayes Factor; DAB, 3,3'-

diaminobenzidine; DS, Down syndrome.