In vitro antimicrobial activity of non-*aureus Staphylococci* and *Mammaliicocci* against *Staphylococcus aureus* and *Streptococcus uberis* and their relationship with the presence of intramammary infections in organic dairy cows

F. Peña-Mosca [1], T. N. Gaire [1], C. Dean [1], D. Manriquez [2], P. Pinedo [3], N. Noyes [1], L. Caixeta [*] [1]

[1] Department of Veterinary Population Medicine, University of Minnesota, St. Paul, 55108

[2] AgNext, Colorado State University, Fort Collins, 80523

[2] Department of Animal Science, Colorado State University, Fort Collins, 80523

* Corresponding author: lcaixeta@umn.edu

ABSTRACT

This study aimed to: (1) Assess the *in vitro* antimicrobial activity of non-*aureus*

Staphylococci and *Mammaliicocci* (**NASM**) isolates from organic cows teat apices with and

without intramammary infections (**IMI**) by *Staphylococcus aureus* (**SAU**) or *Streptococcus* spp.

and *Streptococcus*-like organisms (**SSLO**); (2) Investigate the relationship between IMI status

and NASM counts on teat apex; (3) Explore the relationship between NASM species assigned

taxonomy and the inhibitory activity against SAU and *Streptococcus uberis* (**SUB**). A total of

114 cows from 2 farms were sampled and followed from 8 weeks before to 5 weeks after

calving. Weekly milk samples were cultured and used to design a case control study. Cases were

defined as cows that had an IMI by SAU or SSLO, and controls were randomly chosen from

cows without an IMI by these microorganisms. One teat apex sample from the week prior to IMI

diagnosis was aerobically cultured. Taxonomy of recovered isolates was confirmed using mass

spectrometry and whole genome sequencing. The inhibitory activity against SAU and SUB was

investigated using agar dilution methods. Isolates were ranked based on in vitro inhibitory

activity against SAU and SUB and categorized as "top 10" with the lowest minimum inhibitory

concentration or not. The presence of NASM isolates classified as "top 10" against SAU was

highest in controls (19.4%) compared to cases (5.8%). No difference in NASM counts on teat

apex was observed between cases and controls. The inhibitory activity varied across NASM,

with *Staphylococcus succinus* showing the highest inhibitory activity against both SAU and

SUB.

 KEY WORDS

 Coagulase negative *Staphylococci*. Organic heifers. Antimicrobial activity.

 INTRODUCTION

 Mastitis is the leading cause of morbidity (USDA, 2007) and milk production loss in

 dairy farms (Heikkila et al., 2018). Although preventive measures are important for the control

 of mastitis, antimicrobials are still a major component of on-farm contagious mastitis control

 (Ruegg, 2017). However, the availability of antimicrobials for dairy farmers and veterinarians is

 at-risk due to concerns about antimicrobial resistance to medically important antibiotics and

 policy restrictions on antimicrobial use in livestock (Rajala-Schultz et al., 2021; McCubbin et al.,

 2022; Ruegg, 2022). This limitation is especially remarkable on United States organic-certified

 dairy farms, where regulations for production of organic milk include a strict prohibition against

 the use of antimicrobials and other synthetic substances (NMC, 2019).

 Prior data from our group showed that *Staphylococcus aureus* (**SAU**) and *Streptococcus*

 spp. had a high prevalence and persistence in the mammary gland of dairy cows in organic farms

 in the first 35 days of lactation (Peña-Mosca et al., 2023b). Moreover, this study showed that

 SAU and *Streptococcus* spp. in early lactation were associated with a higher risk of high somatic

 cell count (**SCC**) during the first 180 days of lactation in organic dairy cows (Peña-Mosca et al.,

 2023a). Non-*aureus Staphylococci* and *Mammaliicocci* (**NASM**) are also highly prevalent in

 milk samples (De Buck et al., 2021; Peña-Mosca et al., 2023b), but unlike SAU and

 Streptococcus, NASM have been associated with increased milk production (Piepers et al.,

 2010). Indeed, the infusion into the mammary gland of NASM has been associated with

 decreased risk for intramammary infection (**IMI**) with SAU and *Streptococcus* spp., suggesting a

 potential protective effect of NASM against these two pathogens (Reyher et al., 2012). *In vitro*

studies using isolates obtained from the mammary gland of dairy cows have shown that NASM

are capable of inhibiting the growth of SAU (De Vliegher et al., 2004a; Carson et al., 2017;

Toledo-Silva et al., 2022). Using the cross-streak method for inhibitory testing, these studies

reported that between 9.1% and 14.3% of the isolates were able to inhibit the growth of SAU

(Carson et al., 2017; Toledo-Silva et al., 2022). Recently, agar dilution methods have also been

utilized to investigate the inhibitory capabilities of NASM species on mastitis pathogens

(Toledo-Silva et al., 2022). Moreover, studies in humans have also reported that specific NASM

strains were associated with a decreased risk of skin colonization by SAU (Nakatsuji et al.,

2017).

These findings have raised the hypothesis that certain NASM species could be used for

the development of products to prevent or treat mastitis in dairy farms (De Buck et al., 2021).

However, despite the *in vitro* antimicrobial activity demonstrated by NASM, it remains uncertain

whether any species or strains within the NASM group would yield *in vivo* protective effects

against IMI by major mastitis pathogens. Lastly, the use of the agar dilution method has only

been investigated using a few isolates, and a more comprehensive investigation needs to be

performed to assess the inhibitory activity of different NASM species.

The first objective of this study was to investigate the phenotypic *in vitro* antimicrobial

activity of NASM isolated from the teat apex of organic cows with and without an IMI caused by

SAU or *Streptococcus* spp. and *Streptococcus*-like organisms (**SSLO**) on the week preceding

IMI diagnosis. We hypothesized that NASM isolates from cows without an IMI caused by SAU

or SSLO (i.e., controls) would exhibit higher *in vitro* inhibitory activity against SAU and

Streptococcus uberis (**SUB**) compared to NASM isolates from cows with SAU or SSLO IMI

(i.e., cases). Our second objective was to investigate the relationship between the presence of an

IMI caused by SAU or SSLO and NASM counts on teat-apex swab samples collected the week

prior to the diagnosis of SAU or SSLO IMI. We hypothesized that the NASM counts from the

teat apex samples would be higher in organic cows without an IMI compared to cows with an

IMI. Lastly, our third objective was to explore the relationship between the assigned taxonomy

of NASM species and the presence of inhibitory activity against SAU and SUB. We

hypothesized that there will be differences in the *in vitro* inhibitory across different NASM

species, with a few specific species showing high *in vitro* antimicrobial activity.

74   **MATERIALS AND METHODS**

The Strengthening the Reporting of Observational Studies in Epidemiology - Veterinary

(STROBE-Vet) Statement guidelines were followed for the preparation of this manuscript

(Sargeant et al., 2016). All study activities were approved by the University of Minnesota

Institutional Animal Care and Use Committee (**IACUC**) (Protocol Number: 1807: 36109A),

Colorado State University IACUC (Protocol number: 1442), and Texas Tech University IACUC

(Protocol number: 18068-10). Laboratory activities were approved by the Institutional Biosafety

Committee protocol: 2205-40001H.

Inclusion criteria

Only cows from USDA organic certified dairy farms were enrolled in this study. Farms,

from different regions of the United States and with different herd sizes, were selected for the

study. Enrollment was based on willingness to participate, and proximity to the Universities

involved in the study. This study was part of a larger research initiative to investigate potential

associations between the udder microbiome and udder health (Dean et al., 2021). For this

purpose, cows from 5 organic dairy farms were enrolled 8 weeks before calving and followed up

during the first 5 weeks of lactation (Peña-Mosca et al., 2023b). In this manuscript, data from

only two of these farms were included, as teat apex samples for bacteriological culture were only

collected in these two farms due to limited research labor availability. Specifically, this

manuscript used samples collected from a subset of 114 cows (Farm A [n =21] and Farm B [n

=93]) out of the 596 cows enrolled in the main research project. Samples from Farm A were

collected between August 2019 and January 2020, and included all first-lactation cows that

calved during that period. Samples from Farm B were collected between February and July 2021

and included both first and second lactation cows that calved throughout that interval.

Milk sample collection

After calving, quarter milk samples were collected weekly during the first 5 postpartum

weeks prior to the morning milking by members of the research team, following guidelines from

the National Mastitis Council (NMC, 2017). Upon removal of the pre-dipping disinfectant, 3 to 4

streams of milk were discarded from each quarter, and the teat apex of each quarter was

scrubbed with a gauze soaked in 70% ethanol. Milk samples were collected in a clockwise

direction, beginning with the right rear quarter, and ending with the right front quarter.

Approximately 10 mL of milk from each quarter were collected into separate 60 mL plastic vials

(Capitol Vials, Thermo Fisher Scientific). Samples were placed on ice immediately after

collection and stored in a freezer at -20 °C until further processing.

Milk pooling

Quarter milk samples were combined into composite samples using the following

protocol (Peña-Mosca et al., 2023b). First, milk samples were thawed at 4 °C overnight. Next,

quarter milk samples were homogenized by inverting them upside down twice. Then, 2 mL was

extracted from each vial and dispensed into a single plastic vial inside a laminar hood. Lab-

pooled composite samples were then submitted to the Laboratory for Udder Health at the

University of Minnesota (St. Paul, MN, USA) for milk culture.

Milk culture

Using a cotton swab, approximately 100 μL of milk were plated onto Columbia CNA

agar with 5% sheep blood and MacConkey agar. Agar plates were incubated in aerobic

conditions at 37 °C for a total of 42 to 48 h. Plates were examined by a trained technician to

evaluate the presence of growth of distinct isolates at 18 to 24 hours and 42 to 48 hours. Samples

were defined as contaminated if more than 3 different isolates were identified and omitted from

further analysis (Dean et al., 2022). Taxonomic assignments of cultured isolates were made using

a Matrix-Assisted Laser Desorption/Ionization-Time of Flight (**MALDI-TOF**) mass

spectrometer (**MS**) (Microflex; BrukerDaltonics Inc.) (Jahan et al., 2021). Mass spectra profiles

produced from each isolate were matched against the Biotyper reference library. Confidence

scores were used to assign genus- and species-level classifications (≥ 1.8 and ≥ 2.0 for genus- and

species- classification, respectively) (Jahan et al., 2021). An IMI was defined as a composite

sample containing one or more than one colony forming units (10 CFU/mL) of any cultured

isolate.

Implementation of case control study

Milk culture results were used for the implementation of a case control study. Cases were

defined as cows that had an IMI caused by SAU or SSLO during the first 35 DIM. Controls, with

1 control chosen for each case, were randomly chosen from the list of sampled cows that did not

have an IMI caused by these microorganisms during the first 35 DIM. Cases and controls were

individually matched by farm, lactation number, and DIM of when SAU or SSLO IMI were

diagnosed (±7 DIM). From each selected cow, one teat apex sample from the week prior to the

diagnosis of a SAU or SSLO IMI was processed using the protocol outlined below. Because 3

cows did not have a teat apex sample available in the week prior to first diagnosis of SAU or

SSLO IMI, 2 samples were processed for these animals from the 2 closest available samples

before IMI diagnosis, resulting in a total of 71 samples from 68 cows (34 cases and 34 controls).

Teat apex sample collection

Cows enrolled from two of the participating farms, where labor was available, were

sampled weekly from 8 weeks before calving to 5 weeks after calving in the milking parlor prior

to the morning milking. First, teats were visually inspected, and any visible debris was removed

using a paper towel. Then, using a new pair of disposable gloves, a single sterile pre-moistened

gauze was used to thoroughly swab the distal third of all four teats. After sampling, gauze was

placed in individual tubes containing a pre-loaded solution of 50% glycerol. Gauze samples were

transported in a cooler with ice and stored in -80 °C after arrival to research facilities.

Postpartum samples were collected before pre-dipping disinfectant solutions were applied to the

teat and before milk samples were collected.

Teat apex sample culture

The methods for culturing gauze samples collected from teat apex were adapted from

protocols utilized in the Laboratory for Udder Health at the University of Minnesota to process

towel samples (Rowe et al., 2019). Before sample processing and to reduce the possibility of

batch effects, sample IDs for each gauze sample were randomly sorted using Excel (Microsoft

Corporation). In accordance with this order, isolates were selected for each sampling processing

date in batches of 10 isolates. Gauze samples were thawed at 4 °C overnight and then transferred

into 50 mL round-bottom falcon tubes (Corning Inc.). Afterwards, 5 mL of Phosphate-Buffered

Saline (**PBS**; Gibco, ThermoFisher Scientific) was added to reduce viscosity. Samples were

homogenized carefully by turning tubes upside down twice and vortexing for 10 seconds at low

speed. Samples were allowed to stand for 10 minutes, and the homogenization process was

repeated once. Following homogenization, a single 1 in 10 serial dilution was done using PBS. A

100 uL aliquot from each sample (i.e. undiluted and 10^{-1}) was inoculated onto Mannitol salt agar

(Oxoid). Plates were then incubated in aerobic conditions for 24 hours at 37 °C. Following

incubation, bacterial groups were visually assessed and counted from the dilution plate with the

optimal number of colonies (25 to 250 per plate). Isolates of visually distinct colonies were then

picked from the plate using a sterile calibrated loop (ThermoFisher Scientific), streaked onto

blood agar (Hardy Diagnostics) and incubated for 24 hours at 37 °C.

Species-level identification of isolates

After reaching pure growth, the identity of isolates was investigated using MALDI-TOF

MS, using similar protocols to those outlined above for isolates from milk samples. Non-*aureus*

Staphylococci and *Mammaliicocci* isolates were then stored for further investigation in -80 °C

using commercial cryovials (Hardy Diagnostics). Isolates identified as NASM using MALDI-

TOF MS were further investigated using whole genome sequencing. Isolates were first grown on

brain heart infusion for 24 hours, centrifuged at 10,000 rpm for 5 minutes, and the pellet was

collected in 2 mL sterile polypropylene microcentrifuge tubes (Thermo Fisher Scientific).

Following this procedure, DNA extraction was performed utilizing commercial kits and

following manufacturer's protocol (Qiagen's DNeasy 96 Blood & Tissue Kit, Qiagen). The

amount and quality of extracted DNA was evaluated using PicoGreen fluorometer and

NanoDrop spectrophotometer (ThermoFisher Scientific). Subsequently, library preparation was

performed using the DNA Prep Sample Preparation Kit (Illumina Inc.), and libraries were pooled

for 2x300 paired-end sequencing on the Illumina MISEQ platform, with a targeted depth of 50x

coverage across the *Staphylococcus* spp. genome for each library. Quality control of raw

sequencing data was performed using FastQC (Andrews, 2010) and MultiQC (Ewels et al.,

2016). Trimmomatic was utilized to trim low quality bases and adapters from raw reads (Bolger

et al., 2014). Genomes were assembled utilizing SPAdes version 3.15.5 (Bankevich et al., 2012),

and assembly quality was evaluated using Quast version 5.2.0 (Gurevich et al., 2013). Species

identification of isolates was performed utilizing GTDB-Tk version 1.7.0 (Chaumeil et al., 2020)

and GTDB release 202 (Parks et al., 2022) as implemented in KBase (Arkin et al., 2018).

Determination of in vitro antimicrobial activity of NASM isolates using agar dilution methods

The methods utilized in this study were based on procedures previously described by

others (Toledo-Silva et al., 2022). In each batch of samples, on day 1, isolates that were

previously identified using MALDI-TOF MS as NASM were randomly sorted using Excel

(Microsoft Corporation). Following this procedure, isolates were grown for 24 hours in blood

agar. If evidence of mixed growth (more than 1 colony type) was present, samples were replated.

On day 2, one colony was picked with a 10 uL disposable loop and was grown on brain heart

infusion (**BHI**; Research Products International Corporation) media for 18 to 20 hours. The next

day, 1 in 10 serial dilutions solutions were created using sterile PBS in a 96 well plate

(ThermoFisher Scientific). Bacterial counts (CFU/mL) were estimated by plating 100 uL of the

serial dilutions on blood agar plates that were then aerobically incubated for 18 to 20 h at 37 °C.

One hundred microliters of bacterial suspensions containing NASM isolates at different dilutions

(10^{-2}, 10^{-3}, 10^{-4}, 10^{-5}, 10^{-6}, 10^{-7}) were added to 10 mL of 2% Tryptic soy agar (**TSA**; Oxoid)

inside Falcon round-bottom polypropylene tubes (Corning Inc.). The temperature was examined

using a laboratory digital thermometer (Thermco) before the inclusion of the 2% TSA. The

media were not used until the temperature reached 50 °C (Toledo-Silva et al., 2022). Tubes were

then homogenized and poured into pre-heated (37 °C) TSA plates. Following aerobic incubation

for 18 to 20 h, 10 uL of the bacterial suspensions containing approximately 10,000 CFU of SAU

(American Type Culture Collection (**ATCC**) strain Seattle 1945; *Staphylococcus aureus* subsp.

aureus Rosenbach 25923^{TM}) and SUB (ATCC strain 01040 J; *Streptococcus uberis* Diernhofer

BAA-854 ^{TM}) were inoculated onto the top of plates containing different concentrations of NASM

isolates and incubated for 18 to 20 h. For each batch of samples, negative controls were included

by incubating a 2% TSA agar plate with the inclusion of 100 uL of BHI media only (instead of

NASM isolates). Additionally, a 10 μL droplet of PBS was added on top of the media on each

plate to serve as an additional negative control, ensuring the absence of PBS contamination.

Staphylococcus aureus and SUB were incorporated into the media as positive controls, and their

self-inhibition was tested to explore the determination of MIC driven by nutrient competition

rather than *in vitro* inhibitory activity. Growth of SAU and SUB (i.e., yes vs no) was evaluated

by visual examination of each of the plates and comparison with negative control plates within

each batch. Minimum inhibitory concentration (**MIC**) was defined as the minimum

concentration of NASM (CFU/mL) on the 2% TSA top layer able to inhibit the growth of SAU

or SUB.

Statistical analysis

All analyses were performed in R 4.3.2 (https://www.r-project.org/). Code and output can

be found online: https://fepenamosca.github.io/NASM_in_vitro_inhibitory_activity.io/ . No

priori sample size calculation was performed for this exploratory study. Before modeling, descriptive

characteristics for numerical variables and categorical outcomes were evaluated using the "table

1" package in R (Rich et al., 2023). Data distribution was evaluated using density plots as

implemented in the "car" package in R (Fox et al., 2022). The relationship between IMI status

(i.e., cases versus controls) and $\log_{10}$ -transformed bacterial counts from teat apex samples was

investigated using linear regression models. The relationship between IMI status and presence of

NASM on the teat apex samples (i.e., yes vs no) was investigated using a generalized linear

model with a "binomial" family and "logit" link (logistic regression). For analysis, NASM

taxonomy was determined based on the assigned taxonomy using MALDI-TOF MS and whole

genome sequencing. It was represented as a six-level multi-level variable (*Staphylococcus*

chromogenes, *Staphylococcus haemolyticus*, *Staphylococcus sciuri*, Staphylococcus *succinus*,

Staphylococcus xylosus/pseudoxylosus, and other rare NASM [*Staphylococcus cohnii*,

Staphylococcus devrieseii, *Staphylococcus equorum*]). Cox proportional hazards regression was

used to assess the association between presence of IMI, NASM taxonomy and the concentration

to NASM *in vitro* growth inhibition against SAU and SUB. Observations were right censored

when inhibition was still not present at maximum NASM concentration analyzed (i.e., 10^{-7}

dilution). Proportional hazards assumption was tested by evaluating the Schoenfeld residuals.

Since multiple isolates were harbored from each cow, the non-independence of observations

within each enrolled animal was accounted for using the robust sandwich estimator. The NASM

MIC ($\log_{10}$, CFU/mL) against SAU and SUB was further explored and compared across cases

and controls and different NASM species using mixed linear regression. These models

incorporated cow-ID as a random effect to address the non-independence of observations for

isolates harbored from the same cow. Subsequently, isolates were ranked according to their MIC

(e.g., 1 to 79) and classified as being in the top 10 isolates with lowest MIC or not ("top 10").

The relationship between IMI status and presence of a "top 10" NASM on the teat apex was

investigated using a generalized linear model (logistic regression) with "binomial" family and

"logit" link. When performing this analysis by NASM species, we used the Fisher exact test due

to the absence of "top 10" isolates for some NASM species. The intraclass correlation coefficient

was estimated using the "performance" package (Lüdecke et al., 2021). Estimated marginal

means were estimated using the "emmeans" package in R (Lenth et al., 2024) . Normality

assumption was investigated by evaluation of model residuals and the use of quantile-quantile

plots. Multiple comparisons were accounted for with Tukey adjustment as implemented in the

"emmeans" package (Lenth et al., 2024). Farm-ID was forced into all models that investigated

the relationship between IMI status and NASM inhibitory activity against SAU and SUB.

Because farm ID was strongly associated with the prevalence of different NASM species (i.e.,

each of the NASM species were in most cases highly prevalent in one farm but rarely found in

the other one), this variable was not included in the models comparing *in vitro* inhibitory activity

across different NASM species.

For all models, processing batch (i.e., the date at which gauze samples were processed or

antimicrobial activity of NASM was evaluated) and parity of the cow (i.e., first vs second

lactation) were offered as potential confounders. Presence of confounding was evaluated by

examining the change in the estimates after adjustment by each confounder ([Crude estimate –

Adjusted estimate] / Adjusted estimate). Confounders that changed the estimates by 10% or more

were kept in the models.

RESULTS

Descriptive characteristics

Farms enrolled in this study were located in Minnesota (farm A) and Colorado (farm B)

with herd sizes of 100 and 1,700 milking cows, respectively. Both farms utilized a free-stall

housing system and cows consumed at least 30% of dry matter intake from pasture, when

possible, following organic regulations. All enrolled cows from farm A (21/21) were starting

their first lactation, while in farm B 46.2% (43/93) were starting their first lactation and 53.8%

(50/93) were transitioning to their second lactation. A total of 68 animals (34 cases and 34

controls) were included in this case control study (**Table 1**). Fifteen of these cases had a SAU-

IMI, 15 had a SSLO-IMI and 4 had an IMI caused by both microorganisms. In the first

postpartum sample, 64.7% (22/34) of the cases had the presence of a SAU or SSLO IMI. Teat

apex samples processed from cases and controls showed similar days relative to calving (mean

[SD]; Controls: -2.8 [15.2] vs Cases: -1.2 [16.5]). A flowchart describing the *in vitro* study as

well as the number of observations lost at each step is presented in **Figure 1**.

In vitro inhibitory activity of* NASM *isolated from case and control cows against

Staphylococcus aureus and Streptococcus uberis

283   A total of 275 isolates were submitted for MALDI-TOF MS as shown in **Table 2**.

Surprisingly, *Bacillus* spp. were the most frequently identified genus in controls (61.9%

[83/134]) and cases (61.7% [87/141]). Non-*aureus Staphylococci Mammaliicocci* were identified

in 32.8% (44/134) and 27.7% (39/141) of the isolates obtained from the teat apex samples from

controls and cases, respectively. The most frequent species identified within this group included

Staphylococcus chromogenes (8.0% [22/275]) and *Staphylococcus haemolyticus* (7.6%

[21/275]). The overall inhibitory activity of NASM on SAU was not different between isolates

obtained from cases and controls (HR (95%CI): 0.92 (0.56, 1.49), *P* = 0.72; **Figure 2**). The

adjusted MIC ($\log_{10}$, CFU/mL) for SAU was 3.26 and 3.37 for isolates recovered from controls

and cases, respectively (Estimate [95%CI]: 0.11 [-0.21, 0.44]; *P* = 0.49). Similarly, we did not

find evidence of a difference in the inhibitory activity of NASM on SUB between isolates

obtained from cases and controls (HR (95%CI): 1.23 (0.74, 2.08), *P*=0.45; **Figure 2**). The

adjusted MIC ($\log_{10}$, CFU/ml) of NASM against SUB was 3.17 in isolates from controls and

3.08 in isolates from cases (Estimate [95%CI]: -0.10 [-0.45, 0.25]; $P = 0.58$).

During the study, we evaluated the inhibitory activity of 79 NASM isolates. We were not

able to determine the MIC against SAU and SUB for 5 and 14 NASM isolates, respectively

(**Figure 3**). Among these isolates, the average (SD) highest concentration with an undetermined

MIC was 3.98 (0.45) CFU/mL for SAU and 4.30 (0.52) CFU/mL for SUB.

When SAU was included in the media as a positive control, the MIC against SAU was

3.09. Among the studied NASM isolates, 34.2% (27/52) displayed an MIC below this value

(38.1% [16/42] and 30.0% [11/37] for isolates recovered from controls and cases, respectively).

Upon the inclusion of SUB in the media as a positive control, its MIC against SUB was 3.61.

Twenty percent of the isolates (16/79) displayed a MIC below this value (21.4% [9/42] for

isolates from controls and 18.9% [7/37] for those from cases).

To explore isolates with high inhibitory activity, we arranged the isolates based on their

MIC values against SAU and SUB (**Figure 3A and 3B**). The 10 isolates with lowest MICs

displayed a mean (SD) MIC ($\log_{10}$, CFU/ml) of 2.41 (0.26) and 2.16 (0.32) for SAU and SUB,

respectively. These isolates showed MICs that ranged between 1.92 to 2.66 $\log_{10}$ CFU/mL

against SAU growth and 1.65 to 2.56 $\log_{10}$ CFU/mL against SUB growth. Top 10 isolates with

lowest MIC against SAU were more prevalent in controls (19.4% (7/36)) than in cases (5.8%

(2/35)) (OR (95%CI): 0.25 (0.04, 1.13), $P = 0.07$). However, no evidence of an association was

found between IMI status and the likelihood of harboring a 10 top isolate for SUB (controls:

11.1% (4/35) vs cases: 14.7% (5/36); (OR (95%CI): 1.33 (0.32, 5.83), $P = 0.69$).

Relationship between case control status and bacterial counts on teat apex

The association between IMI status and bacterial count in the teat apex samples is

presented in **Table 3**. No evidence of a difference in NASM counts was found between cases and

controls (Estimate [95%CI]: -0.09 [-0.51, 0.36], *P* = 0.72). The likelihood of harboring NASM

on the teat apex was not different between controls (66.7% [22/36]) and cases (62.9% [24/35])

(OR (95%CI): 0.85 (0.32, 2.25), *P* = 0.74) (**Table 4**). The distribution of NASM species differed

by farm. *Staphylococcus chromogenes, Staphylococcus haemolyticus* and *Staphylococcus*

succinus showed a high prevalence in farm A (28.1% [16/57], 28.1% [16/57] and 14.1% [8/57],

respectively), but were rarely found on farm B (7.1% [1/14], 7.1% [1/14] and 0% [0/14],

respectively). On the other hand, *Staphylococcus xylosus* or *Staphylococcus pseudoxylosus* were

found in half of the samples from farm B (50.0% [7/14]) but showed a low prevalence in farm A

(1.8% [1/57]).

Relationship between NASM taxonomy and inhibitory activity against Staphylococcus aureus

and Streptococcus uberis

The inhibitory activity of NASM on SAU and SUB differed across different NASM

species (*P* < 0.001; **Figure 3, Figure 4**). *Staphylococcus haemolyticus* and other rare NASM

species (i.e., *Staphylococcus cohni*, *Staphylococcus devrieseii*, *Staphylococcus equorum*) showed

lower inhibitory activity against SAU compared to *Staphylococcus chromogenes* (HR [95%CI]:

0.28 [0.09, 0.90], *P*=0.02; and HR [95%CI]: 0.15 [0.03, 0.71], *P*=0.006, respectively),

Staphylococcus sciuri (HR [95%CI]: 0.21 [0.08, 0.58]; *P* = 0.001, and HR [95%CI]: 0.11 [0.02,

0.50], *P*<0.001, *respectively*), *Staphylococcus succinus* (HR [95%CI]: 0.15 [0.05, 0.42],

P<0.001; and HR [95%CI]: 0.08 [0.02, 0.36], *P*<0.001, respectively) and *Staphylococcus*

xylosus/*pseudoxylosus* (HR [95%CI]: 0.14 [0.05, 0.38], *P*<0.001; and HR [95%CI]: 0.07 [0.02,

0.28], $P<0.001$, respectively). The adjusted MIC ($\log_{10}$ CFU/ml) of NASM on SAU for different

NASM species ranged between 2.92 and 3.99 (**Table 5)**. Similarly, when inhibition of SUB was

the outcome of interest, *Staphylococcus haemolyticus* displayed a reduced ability to inhibit SUB

compared to *Staphylococcus chromogenes* (HR [95%CI]: 0.11 [0.02, 0.49]), *Staphylococcus*

sciuri (HR (95%CI): 0.17 (0.06, 0.48)), *Staphylococcus succinus* (HR [95%CI]: 0.13 [0.03,

0.50]) and *Staphylococcus xylosus/pseudoxylosus* (HR [95%CI]: 0.11 [0.03, 0.43]), all $P<0.001$).

However, the inhibitory activity of *Staphylococcus haemolyticus* against SUB was not different

from other rare NASM species (e.g., *Staphylococcus cohnii*, *Staphylococcus devriesei*,

Staphylococcus equorum) (HR [95%CI]: 0.62 [0.06, 6.31], P=0.99). The adjusted MIC ($\log_{10}$

CFU/ml) of NASM against SUB in different NASM species ranged between 2.92 and 3.63

(**Table 5)**. The distribution of isolates ranked by their MIC on SAU and SUB growth is shown in

Figure 3C and 3D. The proportion of isolates classified as "top 10" with respect to SAU and

SUB was different between the NASM species ($P = 0.04$ and 0.06, respectively). Only 4.8%

(1/21) and 0% (0/5) of the isolates identified as *Staphylococcus haemolyticus* and other rare

NASM species, respectively, were classified as being "top 10" against both SAU and SUB. By

comparison, 45.5% (5/11) of *Staphylococcus succinus* isolates were classified as "top 10" against

both SAU and SUB (**Table 6**).

356   **DISCUSSION**

This study aimed to explore the association of naturally occurring NASM isolates and

their ability to inhibit the growth of SAU and SUB, two major mastitis pathogens that are

difficult to control and eradicate on dairy farms (Barkema et al., 2006; Peña-Mosca et al., 2023b;

Woudstra et al., 2023). In our study, we observed an association between the presence of isolates

with high *in vitro* antimicrobial activity against SAU and absence of IMI by SAU or SSLO. In

addition, the inhibitory activity of NASM showed substantial variation across different NASM

species.

Relationship between in vitro antimicrobial activity of non-aureus Staphylococcus and

Mammaliicocci isolates and presence of intramammary infections

Our primary objective was to investigate the *in vitro* antimicrobial activity of NASM

isolates and their association with the presence of IMI. We discovered a wide variation in the

inhibitory activity across NASM isolates. A considerable number of the NASM isolates did not

show inhibitory activity at the maximum concentration studied suggesting the absence of

inhibitory activity. Moreover, we did not find evidence of a difference in the average inhibitory

activity of NASM on SAU or SUB across cases and controls. One possible explanation for these

findings is that average antimicrobial activity may not be an important predictor of the risk of

IMI, and only high antimicrobial activity against a specific pathogen may be effective to prevent

the risk of IMI (Nakatsuji et al., 2017). In this study, we discovered that control cows had the

majority of NASM isolates that were within the "top 10" with the lowest MIC against SAU. In

addition, cows in the control group had a higher chance of having these highly inhibitory isolates

on their teat apices than cows in the case group. Although our results need to be interpreted with

caution, given the small sample size and imprecision of estimates, these results suggest the

presence of a protective effect of NASM with high *in vitro* antimicrobial activity against SAU on

the presence of IMI by SAU or SSLO. These findings are in line with studies performed in

humans in which isolates with high phenotypic antimicrobial activity showed a high prevalence

on the skin on controls, but were rarely identified on people with SAU skin infections (Nakatsuji

et al., 2017). As whole, our results suggest that commensal NASM may provide defense against

mammary gland pathogen invasion, albeit these research discoveries require more study.

Relationship between teat apex colonization by non-aureus Staphylococcus and

Mammaliicocci and presence of intramammary infections

The prevalence of NASM on teat apex (64.8%) was similar to that reported by other

studies in conventional dairy farms in which 70.1% (Mahmmod et al., 2018) and 71.8% (De

Visscher et al., 2016) of the teat-apex samples showed presence of NASM. Consistently, we

found a high prevalence of *Staphylococcus chromogenes* and differences in the prevalence of the

different NASM species across the enrolled farms, which concurs with these previous reports

(De Visscher et al., 2016; Mahmmod et al., 2018). These differences have been previously

attributed to differences in management across the enrolled farms (De Buck et al., 2021) and are

not surprising, given the differences in the prevalence and distribution of different

microorganisms that cause IMI on organic dairy farms (Peña-Mosca et al., 2023b).

Contrary to our hypothesis, the counts or the presence of NASM on teat apex were not

associated with the presence of SAU or SSLO IMI. These results are inconsistent to those

reported by prior studies, in which prepartum colonization of teat apices by NASM was related

to a lower risk of postpartum IMI (Piepers et al., 2011) and subclinical mastitis (De Vliegher et

al., 2003). These findings suggest that, in our study population, the mere presence of NASM on

the teat apex was not a significant factor driving the risk of IMI in early lactation cows. Other

factors, which could potentially account for NASM's ability to prevent major pathogens IMI, as

has been proposed in previous research (Reyher et al., 2012), including the presence of *in vitro*

antimicrobial activity (Braem et al., 2014; Carson et al., 2017; Toledo-Silva et al., 2022).

Relationship between non-aureus Staphylococci and Mammaliicocci taxonomy and presence

of in vitro antimicrobial activity

In our study, nine different NASM species isolated were identified from teat apices of

organic dairy cows. This findings align with previous research (Braem et al., 2014; Carson et al.,

2017), highlighting important differences among various NASM species. In these studies, the

assessment of NASM's antimicrobial activity relied on the cross-streak method. This

methodology revealed that between 9.1% and 13% of the isolates showed signs of antimicrobial

activity (Braem et al., 2014; Carson et al., 2017). In our study, agar dilution methods were used

(Wiegand et al., 2008). This methodology not only aligns with the cross-streak technique but

also provides a quantitative measurement of the antimicrobial activity of bacterial isolates

(Toledo-Silva et al., 2022).

To the best of our knowledge, the quantitative method has only been investigated in 3

different NASM species isolates from the udder of dairy cows, which included *Staphylococcus*

chromogenes, *Staphylococcus epidermidis* and *Staphylococcus simulans*. The MIC of these

species against *Staphylococcus aureus* ranged between 5.10 and 8.24 CFU/mL (log$_{10}$) (Toledo-

Silva et al., 2022). In our study, variations were observed in the inhibitory activity of NASM on

SAU or SUB. Our results suggested that *Staphylococcus haemolyticus* possesses little to no

inhibitory activity on SAU or SUB. This finding is consistent with earlier research that showed

that SAU was not inhibited by *Staphylococcus haemolyticus* (Carson et al., 2017). In contrast,

Staphylococcus succinus exhibited the lowest MIC values, with almost fifty percent of these

isolates falling within the "top 10" isolates possessing the lowest MIC. This differs from other

species, where this proportion did not exceed 20%. In addition, this contradicts prior research

showing no inhibitory activity of *Staphylococcus succinus* against SAU (Carson et al., 2017).

These results may be explained by the observation that antibacterial activity in earlier research in

humans appeared to be strain-specific (Nakatsuji et al., 2017). Hence, it is plausible that

antimicrobial activity of NASM could be attributed more to specific strains rather than distinct

species. Investigating the antimicrobial activity across different strains within each of the NASM

species was beyond the scope of this project and should be further investigated in the future.

Study validity and limitations

This study investigated the quantitative antimicrobial activity of NASM isolates against

SAU and SUB. While this method represents a novel approach that has only recently been

described for its use on mastitis pathogens (Toledo-Silva et al., 2022), it also has certain

limitations. For instance, the novelty of these methods in mastitis research, coupled with the

absence of predetermined MIC cut-off points, complicates the interpretation of results. This

makes it challenging to discern whether the absence or presence of inhibition is due to

antimicrobial activity or merely a competition for nutrients that creates unfavorable conditions

for the growth of target microorganisms in the media (De Vliegher et al., 2004b; Toledo-Silva et

al., 2022). To overcome this limitation, we tested the self-inhibition of SAU and SUB, and we

found that a considerable number of NASM isolates were able to inhibit the growth of these

mastitis pathogens at lower concentrations than those necessary for self-inhibition of SAU and

SUB, suggesting the presence of inhibitory activity.

In this investigation, steps to control for potential technical and biological confounders

were taken. This included the randomization of samples for processing, matching by important

confounders, and assessing the magnitude of confounding by including these variables in the

models used in this study. Nonetheless, it is vital to understand that some of our results could be

still biased by other unmeasured confounders. For instance, cow-level factors that increased the

likelihood of presence of IMI such as abnormally high or low body condition score, milk leakage

(Fernandes et al., 2022), dysfunctional immune response (Sordillo, 2018), or dirty udders

(Schreiner and Ruegg, 2003; Compton et al., 2007) could have influenced our results. Moreover,

the small sample size and resulting imprecision of estimates could hinder our ability to make

inference from the results in this exploratory study. The use of composite milk samples

represents a potential limitation of this observational study, considering the lower sensitivity of

milk culture to detect IMI (Reyher and Dohoo, 2011; Toft et al., 2019). However, repeated

sampling within the first 35 DIM addressed this limitation, improving milk culture sensitivity for

IMI detection (Buelow et al., 1996).

Lastly, one major limitation of our study was the fact that almost two thirds of SAU and

SSLO IMI were detected in the first postpartum milk sample. This suggests that IMI may have

been acquired before calving. This finding is not surprising given the challenges to control

mastitis without antibiotics (NMC, 2019) and the increased prevalence of SAU on organic dairy

farms (Pol and Ruegg, 2007; Cicconi-Hogan et al., 2013). All the above limits our ability to

determine if colonization of NASM with antimicrobial activity preceded the acquisition of SAU

or SSLO IMI. Therefore, it is conceivable that other factors not explored in this study, such as

the presence of a close-up pen with poor bedding management, high stocking density, high fly

presence, and presence of IMI carried across lactations for multiparous cows, may be influencing

the results and explaining why some of these animals calved with IMI (Green et al., 2008; De

Vliegher et al., 2012; Fernandes et al., 2022).

External validity

This study was conducted in two organic dairy farms located in Minnesota and Colorado.

Considering the substantial differences in the prevalence NASM species across different dairy

farms in both this and previous studies (De Visscher et al., 2016; Mahmmod et al., 2018), it is

prudent to restrict the generalization of results to dairy farms managed under similar conditions.

In addition, because antimicrobial activity differed across different NASM species and even

within the species, results should be confirmed including a larger number of isolates

representative of different species that can be found on the mammary gland of dairy cows.

Finally, to address the wide heterogeneity of NASM antimicrobial activity, strain-level

identification is required to identify clades of bacteria and antimicrobial peptides connected to

the presence of *in vitro* antimicrobial activity.

CONCLUSIONS

The presence of NASM isolates classified as "top 10" with lowest MIC against SAU was

higher in cows without SAU or SSLO IMI (controls), compared to cows that had an IMI by these

microorganisms (cases) in early lactation. However, no evidence of an association was found

between presence of isolates with "top 10" lowest MIC against SUB and IMI status. In this

study, no evidence of an association was found between NASM counts on teat apex and presence

of IMI by SAU or SSLO. The inhibitory activity of NASM was different across different NASM

species. This study provides initial information about the inhibitory capabilities of naturally

occurring NASM isolated against SAU and SUB and contributes to our understanding of the

relationship between NASM and udder health in dairy cows.

AKNOWLEDGEMENTS

This study was funded by the Organic Agriculture Research and Extension Initiative

(OREI) from the National Institute of Food and Agriculture (grant number: 2018-51300-28563).

The first author, Felipe Peña-Mosca, was partially funded by Fulbright and Agencia Nacional de

Investigación e Innovación from Uruguay (scholarship number: POS_FUL_2019_1_1008441).

The authors thank the Laboratory for Udder Health at the University of Minnesota for

performing milk cultures and for their assistance in taxonomic identification of isolates. We also

want to express our deep gratitude to the numerous students, faculty, and staff for their assistance

with sample collection and laboratory work. Additionally, we extend our thanks to the owners

and managers of the herds who made this study possible. Felipe Peña-Mosca sampled in

Minnesota, performed laboratory work, cleaned, and wrangled the data, performed statistical

analysis, and prepared the initial and final draft of the manuscript. Tara Gaire performed

laboratory work and edited the manuscript. Chris Dean sampled in Minnesota and edited the

manuscript. Pablo Pinedo conceptualized the study, coordinated sampling in Colorado, and

edited the manuscript. Diego Manriquez sampled in Colorado and edited the manuscript. Noelle

Noyes conceptualized the study, acquired funding, coordinated sampling and laboratory work,

and edited the manuscript. Luciano Caixeta conceptualized the study, acquired funding,

coordinated sampling and laboratory work, and edited the manuscript. The authors have not

stated any conflicts of interest.

DATA AVAILABILITY

512   The raw sequence data generated in this study has been submitted to the Sequence Read

513   Archive on NCBI (Accession: PRJNA1070794).

REFERENCES

Andrews, S. 2010. FastQC: a quality control tool for high throughput sequence data.

Arkin, A.P., R.W. Cottingham, C.S. Henry, N.L. Harris, R.L. Stevens, S. Maslov, P.
Dehal, D. Ware, F. Perez, S. Canon, M.W. Sneddon, M.L. Henderson, W.J. Riehl, D. Murphy-
Olson, S.Y. Chan, R.T. Kamimura, S. Kumari, M.M. Drake, T.S. Brettin, E.M. Glass, D.
Chivian, D. Gunter, D.J. Weston, B.H. Allen, J. Baumohl, A.A. Best, B. Bowen, S.E. Brenner,
C.C. Bun, J.-M. Chandonia, J.-M. Chia, R. Colasanti, N. Conrad, J.J. Davis, B.H. Davison, M.
DeJongh, S. Devoid, E. Dietrich, I. Dubchak, J.N. Edirisinghe, G. Fang, J.P. Faria, P.M.
Frybarger, W. Gerlach, M. Gerstein, A. Greiner, J. Gurtowski, H.L. Haun, F. He, R. Jain, M.P.

Joachimiak, K.P. Keegan, S. Kondo, V. Kumar, M.L. Land, F. Meyer, M. Mills, P.S. Novichkov, T. Oh, G.J. Olsen, R. Olson, B. Parrello, S. Pasternak, E. Pearson, S.S. Poon, G.A. Price, S. Ramakrishnan, P. Ranjan, P.C. Ronald, M.C. Schatz, S.M.D. Seaver, M. Shukla, R.A. Sutormin, M.H. Syed, J. Thomason, N.L. Tintle, D. Wang, F. Xia, H. Yoo, S. Yoo, and D. Yu. 2018. KBase: The United States Department of Energy Systems Biology Knowledgebase. Nat. Biotechnol. 36:566–569. doi:10.1038/nbt.4163.

Bankevich, A., S. Nurk, D. Antipov, A.A. Gurevich, M. Dvorkin, A.S. Kulikov, V.M. Lesin, S.I. Nikolenko, S. Pham, A.D. Prjibelski, A.V. Pyshkin, A.V. Sirotkin, N. Vyahhi, G. Tesler, M.A. Alekseyev, and P.A. Pevzner. 2012. SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. J. Comput. Biol. 19:455–477. doi:10.1089/cmb.2012.0021.

Barkema, H.W., Y.H. Schukken, and R.N. Zadoks. 2006. Invited Review: The role of cow, pathogen, and treatment regimen in the therapeutic success of bovine Staphylococcus aureus mastitis. J. Dairy Sci. 89:1877–1895. doi:10.3168/jds.S0022-0302(06)72256-1.

Bolger, A.M., M. Lohse, and B. Usadel. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30:2114–2120. doi:10.1093/bioinformatics/btu170.

Braem, G., B. Stijlemans, W. Van Haken, S. De Vliegher, L. De Vuyst, and F. Leroy. 2014. Antibacterial activities of coagulase-negative staphylococci from bovine teat apex skin and their inhibitory effect on mastitis-related pathogens. J. Appl. Microbiol. 116:1084–1093. doi:10.1111/jam.12447.

Buelow, K.L., C.B. Thomas, W.J. Goodger, K.V. Nordlund, and M.T. Collins. 1996. Effect of milk sample collection strategy on the sensitivity and specificity of bacteriologic culture and somatic cell count for detection of *Staphylococcus aureus* intramammary infection in dairy cattle. Prev. Vet. Med. 26:1–8. doi:10.1016/0167-5877(95)00518-8.

Carson, D.A., H.W. Barkema, S. Naushad, and J. De Buck. 2017. Bacteriocins of Non-aureus Staphylococci Isolated from Bovine Milk. Appl. Environ. Microbiol. 83:e01015-17. doi:10.1128/AEM.01015-17.

Chaumeil, P.-A., A.J. Mussig, P. Hugenholtz, and D.H. Parks. 2020. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. Bioinformatics 36:1925–1927. doi:10.1093/bioinformatics/btz848.

Cicconi-Hogan, K.M., M. Gamroth, R. Richert, P.L. Ruegg, K.E. Stiglbauer, and Y.H. Schukken. 2013. Risk factors associated with bulk tank standard plate count, bulk tank coliform count, and the presence of Staphylococcus aureus on organic and conventional dairy farms in the United States. J. Dairy Sci. 96:7578–7590. doi:10.3168/jds.2012-6505.

Compton, C.W.R., C. Heuer, K. Parker, and S. McDougall. 2007. Risk factors for peripartum mastitis in pasture-grazed dairy heifers. J. Dairy Sci. 90:4171–4180. doi:10.3168/jds.2006-882.

De Buck, J., V. Ha, S. Naushad, D.B. Nobrega, C. Luby, J.R. Middleton, S. De Vliegher, and H.W. Barkema. 2021. Non-aureus Staphylococci and Bovine Udder Health: Current Understanding and Knowledge Gaps. Front. Vet. Sci. 8.

De Visscher, A., S. Piepers, F. Haesebrouck, and S. De Vliegher. 2016. Teat apex
colonization with coagulase-negative Staphylococcus species before parturition: Distribution and
species-specific risk factors. J. Dairy Sci. 99:1427–1439. doi:10.3168/jds.2015-10326.

De Vliegher, S., L.K. Fox, S. Piepers, S. McDougall, and H.W. Barkema. 2012. Invited
review: Mastitis in dairy heifers: nature of the disease, potential impact, prevention, and control.
J. Dairy Sci. 95:1025–1040. doi:10.3168/jds.2010-4074.

De Vliegher, S., H. Laevens, H.W. Barkema, I.R. Dohoo, H. Stryhn, G. Opsomer, and A.
de Kruif. 2004a. Management Practices and Heifer Characteristics Associated with Early
Lactation Somatic Cell Count of Belgian Dairy Heifers. J. Dairy Sci. 87:937–947.
doi:10.3168/jds.S0022-0302(04)73238-5.

De Vliegher, S., H. Laevens, L.A. Devriese, G. Opsomer, J.L.M. Leroy, H.W. Barkema,
and A. de Kruif. 2003. Prepartum teat apex colonization with Staphylococcus chromogenes in
dairy heifers is associated with low somatic cell count in early lactation. Vet. Microbiol. 92:245–
252. doi:10.1016/s0378-1135(02)00363-2.

De Vliegher, S., G. Opsomer, A. Vanrolleghem, L.A. Devriese, O.C. Sampimon, J. Sol,
H.W. Barkema, F. Haesebrouck, and A. de Kruif. 2004b. In vitro growth inhibition of major
mastitis pathogens by Staphylococcus chromogenes originating from teat apices of dairy heifers.
Vet. Microbiol. 101:215–221. doi:10.1016/j.vetmic.2004.03.020.

Dean, C.J., F. Peña-Mosca, T. Ray, B.J. Heins, V.S. Machado, P.J. Pinedo, L.S. Caixeta,
and N.R. Noyes. 2022. Evaluation of Contamination in Milk Samples Pooled From
Independently Collected Quarters Within a Laboratory Setting. Front. Vet. Sci. 9:818778.
doi:10.3389/fvets.2022.818778.

Dean, C.J., I.B. Slizovskiy, K.K. Crone, A.X. Pfennig, B.J. Heins, L.S. Caixeta, and N.R.
Noyes. 2021. Investigating the cow skin and teat canal microbiomes of the bovine udder using
different sampling and sequencing approaches. J. Dairy Sci. 104:644–661. doi:10.3168/jds.2020-
18277.

Ewels, P., M. Magnusson, S. Lundin, and M. Käller. 2016. MultiQC: summarize analysis
results for multiple tools and samples in a single report. Bioinforma. Oxf. Engl. 32:3047–3048.
doi:10.1093/bioinformatics/btw354.

Fernandes, L., M.L. Celestino, P.R. Menta, T.H. Silva, D. Paiva, T.L. Ribeiro, L.S.
Caixeta, N.R. Noyes, and V.S. Machado. 2022. Cow-related factors associated with
intramammary infections during the 1st week postpartum in primiparous dairy cows in certified
organic herds. Vet. J. 282:105822. doi:10.1016/j.tvjl.2022.105822.

Green, M.J., A.J. Bradley, G.F. Medley, and W.J. Browne. 2008. Cow, Farm, and Herd
Management Factors in the Dry Period Associated with Raised Somatic Cell Counts in Early
Lactation. J. Dairy Sci. 91:1403–1415. doi:10.3168/jds.2007-0621.

Gurevich, A., V. Saveliev, N. Vyahhi, and G. Tesler. 2013. QUAST: quality assessment
tool for genome assemblies. Bioinforma. Oxf. Engl. 29:1072–1075.
doi:10.1093/bioinformatics/btt086.

Jahan, N.A., S.M. Godden, E. Royster, T.C. Schoenfuss, C. Gebhart, J. Timmerman, and
R.C. Fink. 2021. Evaluation of the matrix-assisted laser desorption ionization time of flight mass

spectrometry (MALDI-TOF MS) system in the detection of mastitis pathogens from bovine milk samples. J. Microbiol. Methods 182:106168. doi:10.1016/j.mimet.2021.106168.

Lenth, R.V., B. Bolker, P. Buerkner, I. Giné-Vázquez, M. Herve, M. Jung, J. Love, F. Miguez, H. Riebl, and H. Singmann. 2024. emmeans: Estimated Marginal Means, aka Least-Squares Means.

Lüdecke, D., M. Ben-Shachar, I. Patil, P. Waggoner, and D. Makowski. 2021. performance: An R Package for Assessment, Comparison and Testing of Statistical Models. J. Open Source Softw. 6:3139. doi:10.21105/joss.03139.

Mahmmod, Y.S., I.C. Klaas, L. Svennesen, K. Pedersen, and H. Ingmer. 2018. Communications of Staphylococcus aureus and non-aureus Staphylococcus species from bovine intramammary infections and teat apex colonization. J. Dairy Sci. 101:7322–7333. doi:10.3168/jds.2017-14311.

McCubbin, K.D., E. de Jong, T.J.G.M. Lam, D.F. Kelton, J.R. Middleton, S. McDougall, S. De Vliegher, S. Godden, P.J. Rajala-Schultz, S. Rowe, D.C. Speksnijder, J.P. Kastelic, and H.W. Barkema. 2022. Invited review: Selective use of antimicrobials in dairy cattle at drying-off. J. Dairy Sci. 105:7161–7189. doi:10.3168/jds.2021-21455.

Nakatsuji, T., T.H. Chen, S. Narala, K.A. Chun, A.M. Two, T. Yun, F. Shafiq, P.F. Kotol, A. Bouslimani, A.V. Melnik, H. Latif, J.-N. Kim, A. Lockhart, K. Artis, G. David, P. Taylor, J. Streib, P.C. Dorrestein, A. Grier, S.R. Gill, K. Zengler, T.R. Hata, D.Y.M. Leung, and R.L. Gallo. 2017. Antimicrobials from human skin commensal bacteria protect against Staphylococcus aureus and are deficient in atopic dermatitis. Sci. Transl. Med. 9:eaah4680. doi:10.1126/scitranslmed.aah4680.

NMC. 2017. Laboratory Handbook on Bovine Mastitis. 3rd Edition. National Mastitis Council, Madison, Wisconsin. National Mastitis Council, Madison, Wisconsin.

NMC. 2019. Mastitis control on organic dairy farms in the United States. National Mastitis Council, Madison, Wisconsin.

Parks, D.H., M. Chuvochina, C. Rinke, A.J. Mussig, P.-A. Chaumeil, and P. Hugenholtz. 2022. GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank normalized and complete genome-based taxonomy. Nucleic Acids Res. 50:D785–D794. doi:10.1093/nar/gkab776.

Peña-Mosca, F., C. Dean, L. Fernandes, E. Doster, K. Sharpe, T. Ray, V. Feijoo, A. Antunes, C. Baumann, T. Wehri, B. Heins, P. Pinedo, V. Machado, N. Noyes, and L. Caixeta. 2023a. Associations between early lactation intramammary infections and udder health and performance during the first 180 days in milk in first-lactation organic dairy cows (in review). J. Dairy Sci.

Peña-Mosca, F., C. Dean, V. Machado, L. Fernandes, P. Pinedo, E. Doster, B. Heins, K. Sharpe, T. Ray, V. Feijoo, A. Antunes, C. Baumann, T. Wehri, N. Noyes, and L. Caixeta. 2023b. Investigation of intramammary infections in primiparous cows during early lactation on organic dairy farms. J. Dairy Sci. S0022-0302(23)00552–0. doi:10.3168/jds.2022-23036.

Piepers, S., G. Opsomer, H.W. Barkema, A. de Kruif, and S. De Vliegher. 2010. Heifers infected with coagulase-negative staphylococci in early lactation have fewer cases of clinical

mastitis and higher milk production in their first lactation than noninfected heifers. J. Dairy Sci.
93:2014–2024. doi:10.3168/jds.2009-2897.

Piepers, S., K. Peeters, G. Opsomer, H.W. Barkema, K. Frankena, and S. De Vliegher.
2011. Pathogen group specific risk factors at herd, heifer and quarter levels for intramammary
infections in early lactating dairy heifers. Prev. Vet. Med. 99:91–101.
doi:10.1016/j.prevetmed.2011.02.010.

Pol, M., and P.L. Ruegg. 2007. Relationship Between Antimicrobial Drug Usage and
Antimicrobial Susceptibility of Gram-Positive Mastitis Pathogens. J. Dairy Sci. 90:262–273.
doi:10.3168/jds.S0022-0302(07)72627-9.

Rajala-Schultz, P., A. Nødtvedt, T. Halasa, and K. Persson Waller. 2021. Prudent Use of
Antibiotics in Dairy Cows: The Nordic Approach to Udder Health. Front. Vet. Sci. 8.

Reyher, K.K., and I.R. Dohoo. 2011. Diagnosing intramammary infections: Evaluation of
composite milk samples to detect intramammary infections. J. Dairy Sci. 94:3387–3396.
doi:10.3168/jds.2010-3907.

Reyher, K.K., D. Haine, I.R. Dohoo, and C.W. Revie. 2012. Examining the effect of
intramammary infections with minor mastitis pathogens on the acquisition of new intramammary
infections with major mastitis pathogens—A systematic review and meta-analysis. J. Dairy Sci.
95:6483–6502. doi:10.3168/jds.2012-5594.

Rowe, S.M., S.M. Godden, E. Royster, J. Timmerman, B.A. Crooker, and M. Boyle.
2019. Cross-sectional study of the relationships among bedding materials, bedding bacteria
counts, and intramammary infection in late-lactation dairy cows. J. Dairy Sci. 102:11384–11400.
doi:10.3168/jds.2019-17074.

Ruegg, P.L. 2017. A 100-Year Review: Mastitis detection, management, and prevention.
J. Dairy Sci. 100:10381–10397. doi:10.3168/jds.2017-13023.

Ruegg, P.L. 2022. Realities, Challenges and Benefits of Antimicrobial Stewardship in
Dairy Practice in the United States. Microorganisms 10:1626.
doi:10.3390/microorganisms10081626.

Schreiner, D.A., and P.L. Ruegg. 2003. Relationship between udder and leg hygiene
scores and subclinical mastitis. J. Dairy Sci. 86:3460–3465. doi:10.3168/jds.S0022-
0302(03)73950-2.

Sordillo, L.M. 2018. Mammary Gland Immunobiology and Resistance to Mastitis. Vet.
Clin. North Am. Food Anim. Pract. 34:507–523. doi:10.1016/j.cvfa.2018.07.005.

Toft, N., T. Halasa, S.S. Nielsen, S. Hechinger, L.M. Zervens, D. Schwarz, and C.
Kirkeby. 2019. Composite or aseptic quarter milk samples: Sensitivity and specificity of PCR
and bacterial culture of *Staphylococcus aureus* based on Bayesian latent class evaluation. Prev.
Vet. Med. 171:104689. doi:10.1016/j.prevetmed.2019.05.002.

Toledo-Silva, B., L. Beuckelaere, A. De Visscher, C. Geeroms, E. Meyer, S. Piepers, D.
Thiry, F. Haesebrouck, and S. De Vliegher. 2022. Novel Quantitative Assay to Describe In Vitro
Bovine Mastitis Bacterial Pathogen Inhibition by Non-aureus Staphylococci. Pathog. Basel
Switz. 11:264. doi:10.3390/pathogens11020264.

USDA. 2007. Dairy 2007, Part I: Reference of Dairy Cattle Health and Management
Practices in the United States, 2007. #N480.1007. USDA-APHIS-VS, CEAH, Fort Collins, CO.

Wiegand, I., K. Hilpert, and R.E.W. Hancock. 2008. Agar and broth dilution methods to
determine the minimal inhibitory concentration (MIC) of antimicrobial substances. Nat. Protoc.
3:163–175. doi:10.1038/nprot.2007.521.

Woudstra, S., N. Wente, Y. Zhang, S. Leimbach, M.K. Gussmann, C. Kirkeby, and V.
Krömker. 2023. Strain diversity and infection durations of Staphylococcus spp. and
Streptococcus spp. causing intramammary infections in dairy cows. J. Dairy Sci. 106:4214–4231.
doi:10.3168/jds.2022-22942.

APPENDIX

 Tables

 Table 1. Descriptive characteristics of enrolled cases (n=34) and controls (n=34)

Item	Controls	Cases
Farm		
A	20.6% (7/34)	20.6% (7/34)
B	79.4% (27/34)	79.4% (27/34)
Parity		
First lactation	47.1% (16/34)	47.1% (16/34)
Second lactation	52.9% (18/34)	52.9% (18/34)
Case type		
Staph aureus only	-	44.1% (15/34)
SSLO only	-	44.1% (15/34)
SSLO and *Staph aureus*	-	11.8% (4/34)
Cows positive at calving	-	64.7% (22/34)
Days in milk new IMI	-	13.2 (12.0)
Days relative to calving teat apex sample †	-2.8 (15.2)	-1.2 (16.5)

 Cows positive at calving: Cows with an IMI by *Staphylococcus aureus* or *Streptococcus* spp. and

 Streptococcus-like organisms (SSLO) on the first postpartum sample. † Teat apex sample: gauze

 sample extracted on the week prior to the first time a cow tested positive for an IMI.

702 **Table 2.** Bacterial counts on teat-apex (log10 CFU/square inch of gauze) of controls (referent)
703 and cases (n=71 teat apex samples).

Outcome	Estimated marginal means (SE)		Estimate (95%CI)	*P-value*
	Controls	Cases		
Total bacterial count	1.81 (0.12)	1.91 (0.12)	0.10 (-0.19, 0.40)	0.50
NASM	1.12 (0.18)	1.02 (0.18)	-0.09 (-0.54, 0.35)	0.67
Staph chromogenes	0.26 (0.12)	0.20 (0.12)	-0.05 (-0.34, 0.23)	0.71
Staph haemolyticus	0.21 (0.13)	0.21 (0.13)	-0.005 (-0.33, 0.32)	0.98
Staph sciuri	0.19 (0.10)	0.21 (0.10)	0.02 (-0.23, 0.27)	0.86
Staph succinus [a]	0.17 (0.09)	0.17 (0.09)	-0.004 (-0.24, 0.24)	1.00
Staph xylosus/*pseudoxylosus*	0.46 (0.09)	0.53 (0.10)	0.06 (-0.17, 0.30)	0.60
Other	0.20 (0.07)	0.12 (0.07)	-0.08 (-0.26, 0.09)	0.35

704 [a] *Staph succinus*: Model did not account for farm, because it was not present in farm B. [b] Other:
705 *Staph cohnii, Staph devreseii* and *Staph equorum.* NASM: non-*aureus Staphylococci* and
706 *Mammaliicocci.*

708   **Table 3.** Presence of non-*aureus Staphylococci* and *Mammaliicocci* (NASM) isolates on teat-
709   apex of controls (referent) and cases (n=71 teat apex samples).

Outcome	Presence on teat apex		OR (95%CI)	*P-value*
	Controls	Cases		
NASM	66.7% (24/36)	62.9% (22/35)	0.85 (0.32, 2.25)	0.74
Staph chromogenes	25.0% (9/36)	22.9% (8/35)	0.89 (0.29, 2.73)	0.84
Staph haemolyticus	25.0% (9/36)	22.9% (8/35)	0.89 (0.29, 2.73)	0.84
Staph sciuri	8.3% (3/36)	5.7% (2/35)	0.66 (0.08, 4.28)	0.66
Staph succinus [a]	11.1% (4/36)	11.4% (4/35)	1.03 (0.23, 4.72)	0.97
Staph xylosus/*pseudoxylosus*	11.1% (4/36)	17.1% (6/35)	2.17 (0.39, 14.47)	0.38
Other [b]	8.3% (3/36)	2.9% (1/35)	0.31 (0.01, 2.63)	0.29

[a] *Staph succinus*: Model did not account for farm, because it was not present in farm B. [b] Other:
Staph cohnii, Staph devreseii and *Staph equorum.*

Table 4. Taxonomic identification of isolates harbored from teat apex samples and submitted for
taxonomic identification using Matrix-Assisted Laser Desorption/Ionization-Time of Flight mass
spectrometer (n=275 isolates) and confirmation for non-*aureus Staphylococci* and
Mammaliicocci (NASM) using whole genome sequencing (n=83 isolates).

Taxonomy classification	Controls	Cases
Staph aureus	0% (0/134)	1.4% (2/141)
NASM	32.8% (44/134)	27.7% (39/141)
Staph chromogenes	9.0% (12/134)	7.1% (10/141)
Staph cohnii	0% (0/134)	0.7% (1/141)
Staph devreseii	1.5% (2/134)	0% (0/141)
Staph equorum	0.7% (1/134)	0% (0/141)
Staph haemolyticus	9.0% (12/134)	6.4% (9/141)
Staph pseudoxylosus	1.5% (2/134)	2.1% (3/141)
Staph sciuri	2.1% (3/134)	2.1% (3/141)
Staph succinus	4.5% (6/134)	4.3% (6/141)
Staph xylosus	4.5% (6/134)	5.0% (7/141)
Aerococcus viridans	0% (0/134)	1.4% (2/141)
Aerococcus sp.	0.8% (1/134)	2.8% (4/141)
Bacillus sp.	61.9% (83/134)	61.7% (87/141)
Gram-positive rod	4.5% (6/134)	5.0% (7/141)

718 **Table 5.** Relationship between species identification and minimum inhibitory concentration
719 (MIC) of non-*aureus Staphylococci* and *Mammaliicocci* (NASM) against *Staphylococcus aureus*
720 and *Streptococcus uberis* (n=79 isolates).

NASM taxonomy	*Staphylococcus aureus*		*Streptococcus uberis*	
	MIC (SE)	*P-value*	MIC (SE)	*P-value*
Staph chromogenes	3.32 (0.13) [a]	0.001	3.06 (0.16)	0.23
Staph haemolyticus	3.99 (0.14) [b]		3.63 (0.26)	
Staph sciuri	3.32 (0.24) [a]		3.49 (0.30)	
Staph succinus	2.92 (0.17) [a]		2.92 (0.22)	
Staph xylosus/pseudoxylosus	3.05 (0.14) [a]		3.05 (0.17)	
Other	4.39 (0.32) [b]		3.29 (0.49)	

Other NASM: *Staph cohnii, Staph devreseii, Staph equorum.* Different super indexes within the
same column indicate statistical differences. Intraclass correlation coefficient cow-ID *Staph*
aureus: Not available, *Strep uberis*: Not available.Other: *Staph cohnii, Staph devreseii* and *Staph*
equorum. Only isolates in which MIC was determined were included for analysis; this included
a total of 75 isolates for *Staph aureus*: *Staph chromogenes:* 100% (19/19), *Staph haemolyticus*:
81.0% (17/21), *Staph sciuri*: 100% (6/6), *Staph succinus*: 100% (11/11), *Staph*
xylosus/pseudoxylosus: 100% (18/18), other NASM: 75.0% (3/4). For *Streptococcus uberis* MIC
was determined in 63 isolates: *Staph chromogenes:* 100% (19/19), *Staph haemolyticus*: 33.0%
(7/21), *Staph sciuri*: 100% (6/6), *Staph succinus*: 100.0% (11/11), *Staph xylosus/pseudoxylosus*:
100% (18/18), other NASM: 50.0% (2/4).

Table 5.6. Relationship between non-*aureus Staphylococci* and *Mammaliicocci* (NASM)
identified species and isolates classification as the top 10 with the lowest minimum inhibitory
concentration (MIC) against *Staphylococcus aureus* and *Streptococcus uberis* (n=79 isolates)

Outcome	NASM taxonomy	Presence of isolates with top 10 lowest MIC	*P-value*
Inhibition *Staph aureus*	*Staph chromogenes*	10.5% (2/19)	0.04
	Staph haemolyticus	4.8% (1/21)	
	Staph sciuri	16.9% (1/6)	
	Staph succinus	45.5% (5/11)	
	Staph xylosus/pseudoxylosus	5.6% (1/18)	
	Other	0.0% (0/4)	
Inhibition *Strep uberis*	*Staph chromogenes*	10.5% (2/19)	0.06
	Staph haemolyticus	4.8% (1/21)	
	Staph sciuri	0% (0/6)	
	Staph succinus	45.5% (5/11)	
	Staph xylosus/pseudoxylosus	11.1% (2/18)	
	Other	0.0% (0/4)	

735   Other: *Staph cohnii, Staph devreseii* and *Staph equorum.*

743   *Figures*

745   **Figure 1.** Outline of the study with number of observations lost at each step. NASM: non-*aureus*
746   *Staphylococci* and *Mammaliicocci*. AMA: antimicrobial activity.

Figure 2. Kaplan meier plots showing presence of growth of *Staphylococcus aureus* and *Streptococcus uberis* at different concentrations of non-*aureus Staphylococci* and *Mammaliicocci* (NASM) on cases and controls (Referent group) (n=79 isolates). *Staphylococcus aureus*: HR (95%CI): 0.92 (0.56, 1.49), *P* =0.72. *Streptococcus uberis*: HR (95%CI): 1.23 (0.74, 2.08), *P*=0.45.

Figure 3. Ranking of 79 non-*aureus Staphylococci* and *Mammaliicocci* (NASM) isolates by minimum inhibitory concentration (MIC) against *Staphylococcus aureus* and *Streptococcus uberis* harbored from teat apex stratified by case control status (panel A/B) and assigned species (panel C/D) (n=79 isolates). Undetermined MIC: NASM isolates that showed growth of *Staphylococcus aureus* or *Streptococcus uberis* at the maximum investigated concentration on media. SCH: *Staphylococcus chromogenes*, SHAEM: *Staphylococcus haemolyticus*, SSC: *Staphylococcus sciuri*, SSUC: *Staphylococcus succinus,* SXYL: *Staphylococcus xylosus/pseudoxylosus,* Other: *Staphylococcus cohnii, Staphylococcus devreseii* and *Staphylococcus equorum.*

Figure 4. Kaplan meier plots showing presence of growth of *Staphylococcus aureus* and *Streptococcus uberis* at different concentrations of non-*aureus Staphylococci* and *Mammaliicocci* (NASM) for the different species isolated in this study (n=79 isolates). SCH: *Staphylococcus chromogenes*, SHAEM: *Staphylococcus haemolyticus*, SSC: *Staph sciuri*, SSUC: *Staphylococcus succinus,* SXYL: *Staphylococcus xylosus/pseudoxylosus,* Other: *Staphylococcus cohnii, Staphylococcus devreseii* and *Staphylococcus equorum.* Type III *P-value Staphylococcus aureus* < 0.001. Type III *P-value Streptococcus uberis* < 0.001.

795