In vitro antimicrobial activity of non-*aureus Staphylococci* and *Mammaliicocci* against *Staphylococcus aureus* and *Streptococcus uberis* and their relationship with the presence of intramammary infections in organic dairy cows

F. Peña Mosca [1], T. N. Gaire [1], C. Dean [1], D. Manriquez [2], P. Pinedo [3], N. Noyes [1], L. Caixeta [*] [1]

[1] Department of Veterinary Population Medicine, University of Minnesota, St. Paul, 55108

[2] AgNext, Colorado State University, Fort Collins, 80523

[3] Department of Animal Science, Colorado State University, Fort Collins, 80523

* Corresponding author: lcaixeta@umn.edu

ABSTRACT

This study aimed to: (1) Assess the *in vitro* antimicrobial activity of non-aureus

Staphylococci and *Mammaliicocci* (**NASM**) isolates from organic cows teat apices with and

without intramammary infections (**IMI**) by *Staphylococcus aureus* (**SAU**) or *Streptococcus* spp.

and *Streptococcus*-like organisms (**SSLO**); (2) Investigate the relationship between IMI status

and NASM counts on teat apex; (3) Explore the relationship between NASM species taxonomy

and the inhibitory activity against SAU and *Streptococcus uberis* (**SUB**). A total of 114 cows

from 2 farms were sampled and followed from 8 weeks before to 5 weeks after calving. Weekly

milk samples were cultured and used to design a case control study. Cases were defined as cows

that had an IMI by SAU or SSLO, and controls were randomly chosen from cows without an IMI

by these microorganisms. One teat apex sample from the week prior to IMI diagnosis was

aerobically cultured. Taxonomy of recovered isolates was confirmed using mass spectrometry

and whole genome sequencing. The inhibitory activity against SAU and SUB was investigated

using agar dilution methods. Isolates were ranked based on in vitro inhibitory activity against

SAU and SUB and categorized as "top 10" with the lowest minimum inhibitory concentration or

not. The presence of NASM isolates classified as "top 10" against SAU was highest in controls

(19.4%) compared to cases (5.8%). No difference in NASM counts on teat apex was observed

between cases and controls. The inhibitory activity varied across NASM, with *Staphylococcus*

succinus showing the highest inhibitory activity against both SAU and SUB.

IMPORTANCE

Organic dairy farms need novel strategies to control mastitis. Non-*aureus Staphylococci* and *Mammaliicocci* (**NASM**) have been proposed as potential targets for the development of products to control mastitis, however, their protective activity against major mastitis pathogens requires further investigation. This study provides initial information about the inhibitory capabilities of naturally occurring NASM isolated against mastitis pathogens and its relationship with udder health in organic dairy cows.

INTRODUCTION

Mastitis is the leading cause of morbidity (USDA, 2007) and milk production loss in dairy farms (Heikkila et al., 2018). Although preventive measures are important for the control of mastitis, antimicrobials are still a major component of on-farm contagious mastitis control (Ruegg, 2017). However, the availability of antimicrobials for dairy farmers and veterinarians is at-risk due to concerns about antimicrobial resistance to medically important antibiotics and policy restrictions on antimicrobial use in livestock (Rajala-Schultz et al., 2021; McCubbin et al., 2022; Ruegg, 2022). This limitation is especially remarkable on United States organic-certified dairy farms, where regulations for production of organic milk include a strict prohibition against the use of antimicrobials and other synthetic substances (NMC, 2019).

Prior data from our group showed that *Staphylococcus aureus* (**SAU**) and *Streptococcus* spp. had a high prevalence and persistence in the mammary gland of dairy cows in organic farms in the first 35 days of lactation (Peña-Mosca et al., 2023b). Moreover, this study showed that SAU and *Streptococcus* spp. in early lactation were associated with a higher risk of high somatic cell count (**SCC**) during the first 180 days of lactation in organic dairy cows (Peña-Mosca et al., 2023a). Non-*aureus Staphylococci* and *Mammaliicocci* (**NASM**) are also highly prevalent in

milk samples (De Buck et al., 2021; Peña-Mosca et al., 2023b), but unlike SAU and

Streptococcus, NASM have been associated with increased milk production (Piepers et al.,

2010). Indeed, the presence of NASM has been associated with decreased risk for intramammary

infection (**IMI**) with SAU and *Streptococcus* spp., suggesting a potential protective effect of

NASM against these two pathogens (Reyher et al., 2012). *In vitro* studies using isolates obtained

from the mammary gland of dairy cows have shown that NASM are capable of inhibiting the

growth of SAU (De Vliegher et al., 2004a; Carson et al., 2017; Toledo-Silva et al., 2022). Using

the cross-streak method for inhibitory testing, these studies reported that between 9.1% and

14.3% of the isolates were able to inhibit the growth of SAU (Carson et al., 2017; Toledo-Silva

et al., 2022). Recently, agar dilution methods have also been utilized to investigate the inhibitory

capabilities of NASM species on mastitis pathogens (Toledo-Silva et al., 2022). Moreover,

studies in humans have also reported that specific NASM strains were associated with a

decreased risk of skin colonization by SAU (Nakatsuji et al., 2017).

These findings have raised the hypothesis that certain NASM species could be used for

the development of products to prevent or treat mastitis in dairy farms (De Buck et al., 2021).

However, despite the *in vitro* antimicrobial activity demonstrated by NASM, it remains uncertain

whether any species or strains within the NASM group would yield *in vivo* protective effects

against IMI by major mastitis pathogens. Lastly, the use of the agar dilution method has only

been investigated using a few isolates, and a more comprehensive investigation needs to be

performed to assess the inhibitory activity of different NASM species.

The first objective of this study was to investigate the phenotypic *in vitro* antimicrobial

activity of NASM isolated from the teat apex of organic cows with and without an IMI caused by

SAU or *Streptococcus* spp. and *Streptococcus*-like organisms (**SSLO**) on the week preceding

IMI diagnosis. We hypothesized that NASM isolates from cows without an IMI caused by SAU

or SSLO (i.e., controls) would exhibit higher *in vitro* inhibitory activity against SAU and

Streptococcus uberis (**SUB**) compared to NASM isolates from cows with SAU or SSLO IMI

(i.e., cases). Our second objective was to investigate the relationship between the presence of an

IMI caused by SAU or SSLO and NASM counts on teat-apex swab samples collected the week

prior to the diagnosis of SAU or SSLO IMI. We hypothesized that the NASM counts from the

teat apex samples would be higher in organic cows without an IMI compared to cows with an

IMI. Lastly, our third objective was to explore the relationship between the assigned taxonomy

of NASM species and the presence of inhibitory activity against SAU and SUB. We

hypothesized that there will be differences in the *in vitro* inhibitory across different NASM

species, with a few specific species showing high *in vitro* antimicrobial activity.

MATERIALS AND METHODS

The Strengthening the Reporting of Observational Studies in Epidemiology - Veterinary

(STROBE-Vet) Statement guidelines were followed for the preparation of this manuscript

(Sargeant et al., 2016). All study activities were approved by the University of Minnesota

Institutional Animal Care and Use Committee (**IACUC**) (Protocol Number: 1807: 36109A),

Colorado State University IACUC (Protocol number: 1442), and Texas Tech University IACUC

(Protocol number: 18068-10). Laboratory activities were approved by the Institutional Biosafety

Committee protocol: 2205-40001H.

Inclusion criteria

Only cows from USDA organic certified dairy farms were enrolled in this study. Farms,

from different regions of the United States and with different herd sizes, were selected for the

study. Enrollment was based on willingness to participate, and proximity to the Universities

involved in the study. This study was part of a larger research initiative to investigate potential

associations between the udder microbiome and udder health (Dean et al., 2021). For this

purpose, cows from 5 organic dairy farms were enrolled 8 weeks before calving and followed up

during the first 5 weeks of lactation (Peña-Mosca et al., 2023b). In this manuscript, data from

only two of these farms were included, as teat apex samples for bacteriological culture were only

collected in these two farms due to limited research labor availability. Specifically, this project

used samples collected from a subset of 114 cows (Farm A [n =21] and Farm B [n =93]) out of

the 596 cows enrolled in the main research project. Farm A samples were collected between

August 2019 and January of 2020 and Farm B samples were collected between February and

July of 2021.

Implementation of case control study

Cases were defined as cows that had an IMI caused by SAU or SSLO during the first 35

DIM. Controls, with 1 control chosen for each case, were randomly chosen from the list of

sampled cows that did not have an IMI caused by these microorganisms during the first 35 DIM.

It should be emphasized that milk samples collected from both cases and controls might have

contained an IMI caused by other microorganisms (e.g., gram-negative bacteria, NASM, etc.).

However, these infections were not considered during the process of selecting cases and controls

or in the subsequent analysis. Cases and controls were individually matched by farm, lactation

number, and DIM of when SAU or SSLO IMI were diagnosed (±7 DIM). From each selected

cow, one teat apex sample from the week prior to the diagnosis of a SAU or SSLO IMI was

processed using the protocol outlined below. Because 3 cows did not have a teat apex sample

available in the week prior to first diagnosis of SAU or SSLO IMI, 2 samples were processed for

these animals from the 2 closest available samples surrounding IMI diagnosis, resulting in a total

of 71 samples from 68 cows (34 cases and 34 controls).

Teat apex sample collection

All enrolled cows were sampled weekly from 8 weeks before calving to 5 weeks after

calving. First, teats were visually inspected, and any visible debris was removed using a paper

towel. Then, using a new pair of disposable gloves, a single sterile pre-moistened gauze was used

to thoroughly swab the distal third of all four teats. After sampling, gauze was placed in

individual tubes containing a pre-loaded solution of 50% glycerol. Gauze samples were

transported in a cooler with ice and stored in -80 °C after arrival to research facilities.

Postpartum samples were collected before pre-dipping disinfectant solutions were applied to the

teat and before milk samples were collected.

Teat apex culture and species identification

The methods for culture and species-level identification were adapted from protocols

utilized in the Laboratory for Udder Health at the University of Minnesota to process towel

samples (Rowe et al., 2019). Before sample processing and to reduce the possibility of batch

effects, sample IDs were randomly sorted using Excel (Microsoft Corporation). In accordance

with this order, isolates were selected for each sampling processing date in batches of 10 isolates.

Gauze samples were thawed at 4 °C overnight and then transferred into 50 mL round-bottom

falcon tubes (Corning Inc.). Afterwards, 5 mL of Phosphate-Buffered Saline (**PBS**; Gibco,

ThermoFisher Scientific) was added to reduce viscosity. Samples were homogenized carefully

by turning tubes upside down twice and vortexing for 10 seconds at low speed. Samples were

allowed to stand for 10 minutes, and the homogenization process was repeated once. Following

homogenization, a single 1 in 10 serial dilution was done using PBS. A 100 uL aliquot from each

sample (i.e. undiluted and 10^{-1}) was inoculated onto Mannitol salt agar (Oxoid). Plates were then

incubated in aerobic conditions for 24 hours at 37 °C. Following incubation, bacterial groups

were visually assessed and counted from the dilution plate with the optimal number of colonies

(25 to 250 per plate). Isolates were then picked from the plate using a sterile calibrated loop

(ThermoFisher Scientific), streaked onto blood agar (Hardy Diagnostics) and incubated for 24

hours at 37 °C.

Species identification of isolates

After reaching pure growth, the identity of representative colonies was further

investigated using a Matrix-Assisted Laser Desorption/Ionization-Time of Flight (**MALDI-**

TOF) mass spectrometer (**MS**) (Microflex; BrukerDaltonics Inc.) (Jahan et al., 2021). Mass

spectra profiles produced from each isolate were matched against the Biotyper reference library.

Confidence scores were used to assign genus- and species-level classifications ($\geq$1.8 and $\geq$ 2.0

for genus- and species- classification, respectively) (Jahan et al., 2021). Non-*aureus*

Staphylococci and *Mammaliicocci* isolates were then stored for further investigation in -80 °C

using commercial cryovials (Hardy Diagnostics). Isolates identified as NASM using MALDI-

TOF MS were further investigated using whole genome sequencing. Isolates were first grown on

brain heart infusion for 24 hours, centrifuged at 10,000 rpm for 5 minutes, and the pellet was

collected in 2 mL sterile polypropylene microcentrifuge tubes (Thermo Fisher Scientific).

Following this procedure, DNA extraction was performed utilizing commercial kits and

following manufacturer's protocol (Qiagen's DNeasy 96 Blood & Tissue Kit, Qiagen). The

amount and quality of extracted DNA was evaluated using PicoGreen fluorometer and

NanoDrop spectrophotometer (ThermoFisher Scientific). Subsequently, library preparation was

performed using the DNA Prep Sample Preparation Kit (Illumina Inc.), and libraries were pooled

for 2x300 paired-end sequencing on the Illumina MISEQ platform, with a targeted depth of 50x

coverage across the *Staphylococcus* spp. genome for each library. Quality control of raw

sequencing data was performed using FastQC (Andrews, 2010) and MultiQC (Ewels et al.,

2016). Trimmomatic was utilized to trim low quality bases and adapters from raw reads (Bolger

et al., 2014). Genomes were assembled utilizing SPAdes version 3.15.5 (Bankevich et al., 2012),

and assembly quality was evaluated using Quast version 5.2.0 (Gurevich et al., 2013). Species

identification of isolates was performed utilizing GTDB-Tk version 1.7.0 (Chaumeil et al., 2020)

and GTDB release 202 (Parks et al., 2022) as implemented in KBase (Arkin et al., 2018).

Milk sample collection

After calving, quarter milk samples were collected weekly during the first 5 postpartum

weeks following guidelines from National Mastitis Council (NMC, 2017). After teat apex

samples were collected, a pre-dipping disinfectant solution was applied. Upon removal of the

pre-dipping disinfectant, 3 to 4 streams of milk were discarded from each quarter, and the teat

apex of each quarter was scrubbed with a gauze soaked in 70% ethanol. Milk samples were

collected in a clockwise direction, beginning with the right rear quarter, and ending with the right

front quarter. Approximately 10 mL of milk from each quarter were collected into separate 60

mL plastic vials (Capitol Vials, Thermo Fisher Scientific). Samples were placed on ice

immediately after collection and stored in a freezer at -20 °C until further processing.

Milk pooling

Quarter milk samples were combined into composite samples using the following

protocol (Peña-Mosca et al., 2023b). First, milk samples were thawed at 4 °C overnight. Next,

quarter milk samples were homogenized by inverting them upside down twice. Then, 2 mL was

extracted from each vial and dispensed into a single plastic vial inside a laminar hood. Lab-

pooled composite samples were then submitted to the Laboratory for Udder Health at the

University of Minnesota (St. Paul, MN, USA) for milk culture.

Milk culture

Using a cotton swab, approximately 100 μL of milk were plated onto Columbia CNA

agar with 5% sheep blood and MacConkey agar. Agar plates were incubated in aerobic

conditions at 37 °C for a total of 42 to 48 h. Plates were examined by a trained technician to

evaluate the presence of growth of distinct isolates at 18 to 24 hours and 42 to 48 hours. Samples

were defined as contaminated if more than 3 different isolates were identified and omitted from

further analysis (Dean et al., 2022). Taxonomic assignments of cultured isolates were made using

a MALDI-TOF MS as previously described for teat-apex samples. An IMI was defined as a

composite sample containing one or more than one colony forming units (10 CFU/mL) of any

cultured isolates. Microorganisms were grouped according to their morphological and

epidemiological characteristics in different bacterial groups: SAU, NASM (members from

Staphylococcus genus that were not SAU), SSLO which include members from *Streptococcus*

genus, as well as *Streptococcus*-like organisms (e.g., *Enterococcus* spp., *Aerococcus* spp.,

Lactococcus spp. and *Micrococcus* spp.), gram negative microorganisms (e.g., *Escherichia* spp.,

Klebsiella spp., *Enterobacter* spp., *Pseudomonas* spp. and *Enterobacter* spp., *Pantoea* spp.) and

others (microorganisms that did not belong to any of the previous groups).

Determination of in vitro antimicrobial activity of NASM isolates using the agar dilution

method

The methods utilized in this study were based on procedures previously described by

others (Toledo-Silva et al., 2022). In each batch of samples, on day 1, isolates were randomly

sorted using Excel (Microsoft Corporation). Following this procedure, isolates that were

previously identified using MALDI-TOF MS as NASM were grown for 24 hours in blood agar.

If evidence of mixed growth (more than 1 colony type) was present, samples were replated. On

day 2, one colony was picked with a 10 uL disposable loop and was grown on brain heart

infusion (**BHI**; Research Products International Corporation) media for 18 to 20 hours. The next

day, 1 in 10 serial dilutions solutions were created using sterile PBS in a 96 well plate

(ThermoFisher Scientific). Bacterial counts (CFU/mL) were estimated by plating 100 uL of the

serial dilutions on blood agar plates that were then aerobically incubated for 18 to 20 h at 37 °C.

One hundred microliters of bacterial suspensions containing NASM isolates at different dilutions

(10^{-2}, 10^{-3}, 10^{-4}, 10^{-5}, 10^{-6}, 10^{-7}) were added to 10 mL of 2% Tryptic soy agar (**TSA**; Oxoid)

inside Falcon round-bottom polypropylene tubes (Corning Inc.). The temperature was examined

using a laboratory digital thermometer (Thermco) before the inclusion of the 2% TSA. The

media were not used until the temperature reached 50 °C (Toledo-Silva et al., 2022). Tubes were

then homogenized and poured into pre-heated (37 °C) TSA plates. Following aerobic incubation

for 18 to 20 h, 10 uL of the bacterial suspensions containing approximately 10,000 CFU of SAU

(American Type Culture Collection (**ATCC**) strain Seattle 1945; *Staphylococcus aureus* subsp.

aureus Rosenbach 25923 ™) and SUB (ATCC strain 01040 J; *Streptococcus uberis* Diernhofer

BAA-854 ™) were inoculated onto the top of plates containing different concentrations of NASM

isolates and incubated for 18 to 20 h. For each batch of samples, negative controls were included

by incubating a 2% TSA agar plate with the inclusion of 100 uL of BHI media only (instead of

NASM isolates). Additionally, a 10 μL droplet of PBS was added on top of the media on each

plate to serve as an additional negative control, ensuring the absence of PBS contamination.

Staphylococcus aureus and SUB were incorporated into the media as positive controls, and their

self-inhibition was tested to explore the determination of MIC driven by nutrient competition

rather than *in vitro* inhibitory activity. Growth of SAU and SUB (i.e., yes vs no) was evaluated

by visual examination of each of the plates and comparison with negative control plates within

each batch. Minimum inhibitory concentration (**MIC**) was defined as the minimum

concentration of NASM (CFU/mL) on the 2% TSA top layer able to inhibit the growth of SAU

or SUB.

Statistical analysis

All analyses were performed in R 4.3.2 (https://www.r-project.org/). Code and output can

be found online: https://fepenamosca.github.io/NASM_in_vitro_inhibitory_activity.io/ . Before

modeling, descriptive characteristics for numerical variables and categorical outcomes were

evaluated using the "table 1" package in R (Rich et al., 2023). Data distribution was evaluated

using density plots as implemented in the "car" package in R (Fox et al., 2022). The relationship

between IMI status (i.e., cases versus controls) and $\log_{10}$ -transformed bacterial counts from teat

apex samples was investigated using linear regression models. The relationship between IMI

status and growth (yes or no) of NASM from the teat apex samples was investigated using a

generalized linear model with a "binomial" family and "logit" link (logistic regression). For

analysis, NASM taxonomy was determined based on the assigned taxonomy using MALDI-TOF

MS and whole genome sequencing. It was represented as a six-level multi-level variable

(*Staphylococcus chromogenes*, *Staphylococcus haemolyticus*, *Staphylococcus sciuri*,

Staphylococcus *succinus*, *Staphylococcus xylosus/pseudoxylosus*, other rare NASM

[*Staphylococcus cohnii*, *Staphylococcus devrieseii*, *Staphylococcus equorum*]). To assess the

association between NASM taxonomy and concentration to NASM *in vitro* growth inhibition on

SUB and SAU, Cox proportional hazards regression was utilized. Observations were right

censored when inhibition was still not present at maximum NASM concentration analyzed (i.e.,

10^{-7} dilution). Proportional hazards assumption was tested by evaluating the Schoenfeld

residuals. Since multiple isolates were harbored from each cow, the non-independence of

observations within each enrolled animal was accounted for using the robust sandwich estimator.

The NASM MIC (log_{10}, CFU/mL) on SAU and SUB was further explored and compared across

cases and controls and different NASM species using mixed linear regression, incorporating

cow-ID as a random effect to address the non-independence of observations for isolates harbored

from the same cow. Subsequently, isolates were ranked according to their MIC (e.g., 1 to 79) and

classified as being in the top 10 isolates with lowest MIC or not ("top 10"). The relationship

between IMI status and presence of a "top 10" NASM on the teat apex was investigated using a

generalized linear model (logistic regression) with "binomial" family and "logit" link. When

performing this analysis by NASM species, we used the Fisher exact test due to the absence of

"top 10" isolates for some NASM species. The intraclass correlation coefficient was estimated

using the "performance" package (Lüdecke et al., 2021). Estimated marginal means were

estimated using the "emmeans" package in R (Lenth et al., 2022). Normality assumption was

investigated by evaluation of model residuals and the use of quantile-quantile plots. Multiple

comparisons were accounted for with Tukey adjustment as implemented in the "emmeans"

package (Lenth et al., 2022). Farm-ID was forced into all models that investigated the

relationship between IMI status and NASM inhibitory activity against SAU and SUB. Because

farm ID was strongly associated with the prevalence of different NASM species (i.e., each of the

NASM species were in most cases highly prevalent in one farm but rarely found in the other one)

, this variable was not included in the models comparing *in vitro* inhibitory activity across

different NASM species.

For all models, processing batch (i.e., the date at which gauze samples were processed or

antimicrobial activity of NASM was evaluated) and parity of the cow (i.e., first vs second

lactation) were offered as potential confounders. Presence of confounding was evaluated by

examining the change in the estimates after adjustment by each confounder ([Crude estimate –

Adjusted estimate] / Adjusted estimate). Confounders that changed the estimates by 10% or more

were kept in the models.

277   **RESULTS**

Descriptive characteristics

Farms enrolled in this study were located in Minnesota (farm A) and Colorado (farm B)

with herd sizes of 100 and 3,000 milking cows, respectively. Both farms utilized a free-stall

housing system and cows consumed at least 30% of dry matter intake from pasture, when

possible, following organic regulations. All cows in farm A (14/14) were starting their first

lactation, while for farm B 46.2% (43/93) of the cows were starting their first lactation and

53.8% (50/93) were transitioning to their second lactation. A total of 68 animals (34 cases and 34

controls) were included in this case control study (**Table 1**). Fifteen of these cases had a SAU-

IMI, 15 had a SSLO-IMI and 4 had an IMI caused by both microorganisms. In the first

postpartum sample, 64.7% (22/34) of the cases had the presence of a SAU or SSLO IMI. Teat

apex samples processed from cases and controls showed similar days relative to calving (mean

[SD]; Controls: -2.8 [15.2] vs Cases: -1.2 [16.5]). A flowchart describing the *in vitro* study as

well as the number of observations lost at each step is presented in **Figure 1**.

***In vitro inhibitory activity of** NASM isolated from case and control cows against*

Staphylococcus aureus and Streptococcus uberis

A total of 275 isolates were submitted for MALDI-TOF MS as shown in **Table 2**.

Surprisingly, *Bacillus* spp. were the most frequently identified genus in controls (61.9%

[83/134]) and cases (61.7% [87/141]). Non-*aureus Staphylococci Mammaliicocci* were identified

in 32.8% (44/134) and 27.7% (39/141) of the isolates obtained from the teat apex samples from

controls and cases, respectively. The most frequent species identified within this group included

Staphylococcus chromogenes (8.0% [22/275]) and *Staphylococcus haemolyticus* (7.6%

[21/275]). The overall inhibitory activity of NASM on SAU was not different between isolates

obtained from cases and controls (HR (95%CI): 0.92 (0.56, 1.49), $P = 0.72$; **Figure 2**). The

adjusted MIC ($\log_{10}$, CFU/mL) for SAU was 3.26 and 3.37 for isolates recovered from controls

and cases, respectively (Estimate [95%CI]: 0.11 [-0.21, 0.44]; $P = 0.49$). Similarly, we did not

find evidence of a difference in the inhibitory activity of NASM on SUB between isolates

obtained from cases and controls (HR (95%CI): 1.23 (0.74, 2.08), $P=0.45$; **Figure 2**). The

adjusted MIC ($\log_{10}$, CFU/ml) of NASM against SUB was 3.17 in isolates from controls and

3.08 in isolates from cases (Estimate [95%CI]: -0.10 [-0.45, 0.25]; $P = 0.58$).

During the study, we evaluated the inhibitory activity of 79 NASM isolates. We were not

able to determine the MIC against SAU and SUB for 5 and 14 NASM isolates, respectively

(**Figure 3**). Among these isolates, the average (SD) highest concentration with an undetermined

MIC was 3.98 (0.45) CFU/mL for SAU and 4.30 (0.52) CFU/mL for SUB.

When SAU was included in the media as a positive control, the MIC against SAU was

3.09. Among the studied NASM isolates, 34.2% (27/52) displayed an MIC below this value

(38.1% [16/42] and 30.0% [11/37] for isolates recovered from controls and cases, respectively).

Upon the inclusion of SUB in the media as a positive control, its MIC against SUB was 3.61.

Twenty percent of the isolates (16/79) displayed a MIC below this value (21.4% [9/42] for

isolates from controls and 18.9% [7/37] for those from cases).

317   To explore isolates with high inhibitory activity, we arranged the isolates based on their

MIC values against SAU and SUB (**Figure 3A and 3B**). The 10 isolates with lowest MICs

displayed a mean (SD) MIC ($\log_{10}$, CFU/ml) of 2.41 (0.26) and 2.16 (0.32) for SAU and SUB,

respectively. These isolates showed MICs that ranged between 1.92 to 2.66 $\log_{10}$ CFU/mL

against SAU growth and 1.65 to 2.56 $\log_{10}$ CFU/mL against SUB growth. Top 10 isolates with

lowest MIC against SAU were more prevalent in controls (19.4% (7/36)) than in cases (5.8%

(2/35)) (OR (95%CI): 0.25 (0.04, 1.13), $P = 0.07$). However, no evidence of an association was

found between IMI status and the likelihood of harboring a 10 top isolate for SUB (controls:

11.1% (4/35) vs cases: 14.7% (5/36); (OR (95%CI): 1.33 (0.32, 5.83), $P = 0.69$).

Relationship between case control status and bacterial counts on teat apex

327   The association between IMI status and bacterial count in the teat apex samples is

presented in **Table 3**. No evidence of a difference in NASM counts was found between cases and

controls (Estimate [95%CI]: -0.09 [-0.51, 0.36], $P = 0.72$). The likelihood of harboring NASM

on the teat apex was not different between controls (66.7% [22/36]) and cases (62.9% [24/35])

(OR (95%CI): 0.85 (0.32, 2.25), $P = 0.74$) (**Table 4**). The distribution of NASM species differed

by farm. *Staphylococcus chromogenes, Staphylococcus haemolyticus* and *Staphylococcus*

succinus showed a high prevalence in farm A (28.1% [16/57], 28.1% [16/57] and 14.1% [8/57],

respectively), but were rarely found on farm B (7.1% [1/14], 7.1% [1/14] and 0% [0/14],

respectively). On the other hand, *Staphylococcus xylosus* or *Staphylococcus pseudoxylosus* were

found in half of the samples from farm B (50.0% [7/14]) but showed a low prevalence in farm A

(1.8% [1/57]).

Relationship between NASM taxonomy and inhibitory activity against Staphylococcus aureus

and Streptococcus uberis

The inhibitory activity of NASM on SAU and SUB differed across different NASM

species ($P < 0.001$; **Figure 3, Figure 4**). *Staphylococcus haemolyticus* and other rare NASM

species (i.e., *Staphylococcus cohni*, *Staphylococcus devrieseii*, *Staphylococcus equorum*) showed

lower inhibitory activity against SAU compared to *Staphylococcus chromogenes* (HR [95%CI]:

0.28 [0.09, 0.90], $P=0.02$; and HR [95%CI]: 0.15 [0.03, 0.71], $P=0.006$, respectively),

Staphylococcus sciuri (HR [95%CI]: 0.21 [0.08, 0.58]; $P = 0.001$, and HR [95%CI]: 0.11 [0.02,

0.50], $P<0.001$, *respectively*), *Staphylococcus succinus* (HR [95%CI]: 0.15 [0.05, 0.42],

$P<0.001$; and HR [95%CI]: 0.08 [0.02, 0.36], $P<0.001$, respectively) and *Staphylococcus*

xylosus/*pseudoxylosus* (HR [95%CI]: 0.14 [0.05, 0.38], $P<0.001$; and HR [95%CI]: 0.07 [0.02,

0.28], $P<0.001$, respectively). The adjusted MIC ($\log_{10}$ CFU/ml) of NASM on SAU for different

NASM species ranged between 2.92 and 3.99 (**Table 5)**. Similarly, when inhibition of SUB was

the outcome of interest, *Staphylococcus haemolyticus* displayed a reduced ability to inhibit SUB

compared to *Staphylococcus chromogenes* (HR [95%CI]: 0.11 [0.02, 0.49]), *Staphylococcus*

sciuri (HR (95%CI): 0.17 (0.06, 0.48)), *Staphylococcus succinus* (HR [95%CI]: 0.13 [0.03,

0.50]) and *Staphylococcus xylosus*/*pseudoxylosus* (HR [95%CI]: 0.11 [0.03, 0.43]), all $P<0.001$).

However, the inhibitory activity of *Staphylococcus haemolyticus* against SUB was not different

from other rare NASM species (e.g., *Staphylococcus cohnii*, *Staphylococcus devriesei*,

Staphylococcus equorum) (HR [95%CI]: 0.62 [0.06, 6.31], $P=0.99$). The adjusted MIC ($\log_{10}$

CFU/ml) of NASM against SUB in different NASM species ranged between 2.92 and 3.63

(Table 5). The distribution of isolates ranked by their MIC on SAU and SUB growth is shown in

Figure 3C and 3D. The proportion of isolates classified as "top 10" with respect to SAU and

SUB was different between the NASM species ($P = 0.04$ and 0.06, respectively). Only 4.8%

(1/21) and 0% (0/5) of the isolates identified as *Staphylococcus haemolyticus* and other rare

NASM species, respectively, were classified as being "top 10" against both SAU and SUB. By

comparison, 45.5% (5/11) of *Staphylococcus succinus* isolates were classified as "top 10" against

both SAU and SUB (**Table 6**).

<div align="center">**DISCUSSION**</div>

This study aimed to explore the association of naturally occurring NASM isolates and

their ability to inhibit the growth of SAU and SUB, two major mastitis pathogens that are

difficult to control and eradicate on dairy farms (Barkema et al., 2006; Peña-Mosca et al., 2023b;

Woudstra et al., 2023). In our study, we observed an association between the presence of isolates

with high *in vitro* antimicrobial activity against SAU and absence of IMI by SAU or SSLO. In

addition, the inhibitory activity of NASM showed substantial variation across different NASM

species.

Relationship between in vitro antimicrobial activity of non-aureus Staphylococcus and

Mammaliicocci isolates and presence of intramammary infections

Our primary objective was to investigate the *in vitro* antimicrobial activity of NASM

isolates and their association with the presence of IMI. We discovered a wide variation in the

inhibitory activity across NASM isolates. A considerable number of the NASM isolates did not

show inhibitory activity at the maximum concentration studied suggesting the absence of

important inhibitory activity for those microorganisms. Moreover, we did not find evidence of a

difference in the average inhibitory activity of NASM on SAU or SUB across cases and controls.

One possible explanation for these findings is that average antimicrobial activity may not be an

important predictor of the risk of IMI, and only high antimicrobial activity against a specific

pathogen may be effective to prevent the risk of IMI (Nakatsuji et al., 2017). In this study, we

discovered that control cows had the majority of NASM isolates that were within the "top 10"

with the lowest MIC against SAU. In addition, cows in the control group had a higher chance of

having these highly inhibitory isolates on their teat apices than cows in the case group. Although

our results need to be interpreted with caution, given the small sample size and imprecision of

estimates, these results suggest the presence of a protective effect of NASM with high *in vitro*

antimicrobial activity against SAU on the presence of IMI by SAU or SSLO. These findings are

in line with studies performed in humans in which isolates with high phenotypic antimicrobial

activity showed a high prevalence on the skin on controls, but were rarely identified on people

with SAU skin infections (Nakatsuji et al., 2017). As whole, our results suggest that commensal

NASM may provide defense against mammary gland pathogen invasion, albeit these research

discoveries require more study.

Relationship between teat apex colonization by non-aureus Staphylococcus and

Mammaliicocci and presence of intramammary infections

The prevalence of NASM on teat apex (64.8%) was similar to that reported by other

studies in conventional dairy farms in which 70.1% (Mahmmod et al., 2018) and 71.8% (De

Visscher et al., 2016) of the teat-apex samples showed presence of NASM. Consistently, we

found a high prevalence of *Staphylococcus chromogenes* and differences in the prevalence of the

different NASM species across the enrolled farms, which concurs with these previous reports

(De Visscher et al., 2016; Mahmmod et al., 2018). These differences have been previously

attributed to differences in management across the enrolled farms (De Buck et al., 2021) and are

not surprising, given the differences in the prevalence and distribution of different

microorganisms that cause IMI in organic dairy farms (Peña-Mosca et al., 2023b).

Contrary to our hypothesis, the counts or the presence of NASM on teat apex were not

associated with the presence of SAU or SSLO IMI. These results are inconsistent to those

reported by prior studies, in which prepartum colonization of teat apices by NASM was related

to a lower risk of postpartum IMI (Piepers et al., 2011) and subclinical mastitis (De Vliegher et

al., 2003). These findings suggest that, in our study population, the mere presence of NASM on

the teat apex was not a significant factor driving the risk of IMI in early lactation cows. Other

factors, which could potentially account for NASM's ability to prevent major pathogens IMI, as

has been proposed in previous research (Reyher et al., 2012), including the presence of *in vitro*

antimicrobial activity (Braem et al., 2014; Carson et al., 2017; Toledo-Silva et al., 2022).

Relationship between non-aureus Staphylococci and Mammaliicocci taxonomy and presence

of in vitro antimicrobial activity

In our study, nine different NASM species isolated were identified from teat apices of

organic dairy cows. This findings align with previous research (Braem et al., 2014; Carson et al.,

2017), highlighting important differences among various NASM species. In these studies, the

assessment of NASM's antimicrobial activity relied on the cross-streak method. This

methodology revealed that between 9.1% and 13% of the isolates showed signs of antimicrobial

activity (Braem et al., 2014; Carson et al., 2017). In our study, agar dilution methods were used

(Wiegand et al., 2008). This methodology not only aligns with the cross-streak technique but

also provides a quantitative measurement of the antimicrobial activity of bacterial isolates

(Toledo-Silva et al., 2022).

To the best of our knowledge, the quantitative method has only been investigated in 3

different NASM species isolates from the udder of dairy cows, which included *Staphylococcus*

chromogenes, *Staphylococcus epidermidis* and *Staphylococcus simulans*. The MIC of these

species against *Staphylococcus aureus* ranged between 5.10 and 8.24 CFU/mL ($\log_{10}$) (Toledo-

Silva et al., 2022). In our study, variations were observed in the inhibitory activity of NASM on

SAU or SUB. Our results suggested that *Staphylococcus haemolyticus* possesses little to no

inhibitory activity on SAU or SUB. This finding is consistent with earlier research that showed

that SAU was not inhibited by *Staphylococcus haemolyticus* (Carson et al., 2017). In contrast,

Staphylococcus succinus exhibited the lowest MIC values, with almost fifty percent of these

isolates falling within the "top 10" isolates possessing the lowest MIC. This differs from other

species, where this proportion did not exceed 20%. In addition, this contradicts prior research

showing no inhibitory activity of *Staphylococcus succinus* against SAU (Carson et al., 2017).

These results may be explained by the observation that antibacterial activity in earlier research in

humans appeared to be strain-specific (Nakatsuji et al., 2017). Hence, it is plausible that

antimicrobial activity of NASM could be attributed more to specific strains rather than distinct

species. Investigating the antimicrobial activity across different strains within each of the NASM

species was beyond the scope of this project and should be further investigated in the future.

Study validity and limitations

This study investigated the quantitative antimicrobial activity of NASM isolates against

SAU and SUB. While this method represents a novel approach that has only recently been

described for its use on mastitis pathogens (Toledo-Silva et al., 2022), it also has certain

limitations. For instance, the novelty of these methods in mastitis research, coupled with the

absence of predetermined MIC cut-off points, complicates the interpretation of results. This

makes it challenging to discern whether the absence or presence of inhibition is due to

antimicrobial activity or merely a competition for nutrients that creates unfavorable conditions

for the growth of target microorganisms in the media (De Vliegher et al., 2004b; Toledo-Silva et

al., 2022). To overcome this limitation, we tested the self-inhibition of SAU and SUB, and we

found that a considerable number of NASM isolates were able to inhibit the growth of these

mastitis pathogens at lower concentrations than those necessary for self-inhibition of SAU and

SUB, suggesting the presence of inhibitory activity.

In this investigation, steps to control for potential technical and biological confounders

were taken. This included the randomization of samples for processing, matching by important

confounders, and assessing the magnitude of confounding by including these variables in the

models used in this study. Nonetheless, it is vital to understand that some of our results could be

still biased by other unmeasured confounders. For instance, cow-level factors that increased the

likelihood of presence of IMI such as abnormally high or low body condition score, milk leakage

(Fernandes et al., 2022), dysfunctional immune response (Sordillo, 2018), or dirty udders

(Schreiner and Ruegg, 2003; Compton et al., 2007) could have influenced our results. Moreover,

the small sample size and resulting imprecision of estimates could hinder our ability to make

inference from the results in this exploratory study.

Lastly, one major limitation of our study was the fact that almost two thirds of SAU and

SSLO IMI were detected in the first postpartum milk sample. This suggests that IMI may have

been acquired before calving. This finding is not surprising given the challenges to control

mastitis without antibiotics (NMC, 2019) and the increased prevalence of SAU on organic dairy

farms (Pol and Ruegg, 2007; Cicconi-Hogan et al., 2013). All the above limits our ability to

determine if colonization of NASM with antimicrobial activity preceded the acquisition of SAU

or SUB IMI. Therefore, it is conceivable that other factors not explored in this study, such as the

presence of a close-up pen with poor bedding management, high stocking density, high fly

presence, and presence of IMI carried across lactations for multiparous cows, may be influencing

the results and explaining why some of these animals calved with IMI (Green et al., 2008; De

Vliegher et al., 2012; Fernandes et al., 2022).

External validity

This study was conducted in two organic dairy farms located in Minnesota and Colorado.

Considering the substantial differences in the prevalence NASM species across different dairy

farms in both this and previous studies (De Visscher et al., 2016; Mahmmod et al., 2018), it is

prudent to restrict the generalization of results to dairy farms managed under similar conditions.

In addition, because antimicrobial activity differed across different NASM species and even

within the species, results should be confirmed including a larger number of isolates

representative of different species that can be found on the mammary gland of dairy cows.

Finally, to address the wide heterogeneity of NASM antimicrobial activity, strain-level

identification is required to identify clades of bacteria and antimicrobial peptides connected to

the presence of *in vitro* antimicrobial activity.

489   **CONCLUSIONS**

The presence of NASM isolates classified as "top 10" with lowest MIC against SAU was

highest in cows without SAU or SUB IMI (controls) compared to cows that had an IMI by these

microorganisms (cases) in early lactation. However, no evidence of an association was found

between presence of isolates with "top 10" lowest MIC against SUB and IMI status. In this

study, no evidence of an association was found between NASM counts on teat apex and presence

of IMI by SAU or SSLO. The inhibitory activity of NASM was different across different NASM

species. This study provides initial information about the inhibitory capabilities of naturally

occurring NASM isolated against SAU and SUB and contributes to our understanding of the

relationship between NASM and udder health in dairy cows.

AKNOWLEDGEMENTS

This study was funded by the Organic Agriculture Research and Extension Initiative

(OREI) from the National Institute of Food and Agriculture (grant number: 2018-51300-28563).

The first author, Felipe Peña-Mosca, was partially funded by Fulbright and Agencia Nacional de

Investigación e Innovación from Uruguay (scholarship number: POS_FUL_2019_1_1008441).

The authors thank the Laboratory for Udder Health at the University of Minnesota for

performing milk cultures and for their assistance in taxonomic identification of isolates. We also

want to express our deep gratitude to the numerous students, faculty, and staff for their assistance

with sample collection and laboratory work. Additionally, we extend our thanks to the owners

and managers of the herds who made this study possible. Felipe Peña-Mosca sampled in

Minnesota, performed laboratory work, cleaned, and wrangled the data, performed statistical

analysis, and prepared the initial and final draft of the manuscript. Tara Gaire performed

laboratory work and edited the manuscript. Chris Dean sampled in Minnesota and edited the

manuscript. Pablo Pinedo conceptualized the study, coordinated sampling in Colorado, and

edited the manuscript. Diego Manriquez sampled in Colorado and edited the manuscript. Noelle

Noyes conceptualized the study, acquired funding, coordinated sampling and laboratory work,

and edited the manuscript. Luciano Caixeta conceptualized the study, acquired funding,

coordinated sampling and laboratory work, and edited the manuscript. The authors have not

stated any conflicts of interest.

DATA AVAILABILITY

The raw sequence data generated in this study has been submitted to the Sequence Read

Archive on NCBI (Accession: PRJNA1070794).

REFERENCES

Andrews, S. 2010. FastQC: a quality control tool for high throughput sequence data.

Arkin, A.P., R.W. Cottingham, C.S. Henry, N.L. Harris, R.L. Stevens, S. Maslov, P. Dehal, D.
Ware, F. Perez, S. Canon, M.W. Sneddon, M.L. Henderson, W.J. Riehl, D. Murphy-Olson, S.Y.
Chan, R.T. Kamimura, S. Kumari, M.M. Drake, T.S. Brettin, E.M. Glass, D. Chivian, D. Gunter,
D.J. Weston, B.H. Allen, J. Baumohl, A.A. Best, B. Bowen, S.E. Brenner, C.C. Bun, J.-M.
Chandonia, J.-M. Chia, R. Colasanti, N. Conrad, J.J. Davis, B.H. Davison, M. DeJongh, S.
Devoid, E. Dietrich, I. Dubchak, J.N. Edirisinghe, G. Fang, J.P. Faria, P.M. Frybarger, W.
Gerlach, M. Gerstein, A. Greiner, J. Gurtowski, H.L. Haun, F. He, R. Jain, M.P. Joachimiak,
K.P. Keegan, S. Kondo, V. Kumar, M.L. Land, F. Meyer, M. Mills, P.S. Novichkov, T. Oh, G.J.
Olsen, R. Olson, B. Parrello, S. Pasternak, E. Pearson, S.S. Poon, G.A. Price, S. Ramakrishnan,
P. Ranjan, P.C. Ronald, M.C. Schatz, S.M.D. Seaver, M. Shukla, R.A. Sutormin, M.H. Syed, J.
Thomason, N.L. Tintle, D. Wang, F. Xia, H. Yoo, S. Yoo, and D. Yu. 2018. KBase: The United
States Department of Energy Systems Biology Knowledgebase. Nat. Biotechnol. 36:566–569.
doi:10.1038/nbt.4163.

Bankevich, A., S. Nurk, D. Antipov, A.A. Gurevich, M. Dvorkin, A.S. Kulikov, V.M. Lesin, S.I.
Nikolenko, S. Pham, A.D. Prjibelski, A.V. Pyshkin, A.V. Sirotkin, N. Vyahhi, G. Tesler, M.A.
Alekseyev, and P.A. Pevzner. 2012. SPAdes: A New Genome Assembly Algorithm and Its
Applications to Single-Cell Sequencing. J. Comput. Biol. 19:455–477.
doi:10.1089/cmb.2012.0021.

Barkema, H.W., Y.H. Schukken, and R.N. Zadoks. 2006. Invited Review: The role of cow,
pathogen, and treatment regimen in the therapeutic success of bovine Staphylococcus aureus
mastitis. J. Dairy Sci. 89:1877–1895. doi:10.3168/jds.S0022-0302(06)72256-1.

Bolger, A.M., M. Lohse, and B. Usadel. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30:2114–2120. doi:10.1093/bioinformatics/btu170.

Braem, G., B. Stijlemans, W. Van Haken, S. De Vliegher, L. De Vuyst, and F. Leroy. 2014.
Antibacterial activities of coagulase-negative staphylococci from bovine teat apex skin and their
inhibitory effect on mastitis-related pathogens. J. Appl. Microbiol. 116:1084–1093.
doi:10.1111/jam.12447.

Carson, D.A., H.W. Barkema, S. Naushad, and J. De Buck. 2017. Bacteriocins of Non-aureus
Staphylococci Isolated from Bovine Milk. Appl. Environ. Microbiol. 83:e01015-17.
doi:10.1128/AEM.01015-17.

Chaumeil, P.-A., A.J. Mussig, P. Hugenholtz, and D.H. Parks. 2020. GTDB-Tk: a toolkit to
classify genomes with the Genome Taxonomy Database. Bioinformatics 36:1925–1927.
doi:10.1093/bioinformatics/btz848.

Cicconi-Hogan, K.M., M. Gamroth, R. Richert, P.L. Ruegg, K.E. Stiglbauer, and Y.H.
Schukken. 2013. Risk factors associated with bulk tank standard plate count, bulk tank coliform
count, and the presence of Staphylococcus aureus on organic and conventional dairy farms in the
United States. J. Dairy Sci. 96:7578–7590. doi:10.3168/jds.2012-6505.

Compton, C.W.R., C. Heuer, K. Parker, and S. McDougall. 2007. Risk factors for peripartum
mastitis in pasture-grazed dairy heifers. J. Dairy Sci. 90:4171–4180. doi:10.3168/jds.2006-882.

De Buck, J., V. Ha, S. Naushad, D.B. Nobrega, C. Luby, J.R. Middleton, S. De Vliegher, and
H.W. Barkema. 2021. Non-aureus Staphylococci and Bovine Udder Health: Current
Understanding and Knowledge Gaps. Front. Vet. Sci. 8.

De Visscher, A., S. Piepers, F. Haesebrouck, and S. De Vliegher. 2016. Teat apex colonization
with coagulase-negative Staphylococcus species before parturition: Distribution and species-
specific risk factors. J. Dairy Sci. 99:1427–1439. doi:10.3168/jds.2015-10326.

De Vliegher, S., L.K. Fox, S. Piepers, S. McDougall, and H.W. Barkema. 2012. Invited review:
Mastitis in dairy heifers: nature of the disease, potential impact, prevention, and control. J. Dairy
Sci. 95:1025–1040. doi:10.3168/jds.2010-4074.

De Vliegher, S., H. Laevens, H.W. Barkema, I.R. Dohoo, H. Stryhn, G. Opsomer, and A. de
Kruif. 2004a. Management Practices and Heifer Characteristics Associated with Early Lactation
Somatic Cell Count of Belgian Dairy Heifers. J. Dairy Sci. 87:937–947. doi:10.3168/jds.S0022-
0302(04)73238-5.

De Vliegher, S., H. Laevens, L.A. Devriese, G. Opsomer, J.L.M. Leroy, H.W. Barkema, and A.
de Kruif. 2003. Prepartum teat apex colonization with Staphylococcus chromogenes in dairy
heifers is associated with low somatic cell count in early lactation. Vet. Microbiol. 92:245–252.
doi:10.1016/s0378-1135(02)00363-2.

De Vliegher, S., G. Opsomer, A. Vanrolleghem, L.A. Devriese, O.C. Sampimon, J. Sol, H.W.
Barkema, F. Haesebrouck, and A. de Kruif. 2004b. In vitro growth inhibition of major mastitis
pathogens by Staphylococcus chromogenes originating from teat apices of dairy heifers. Vet.
Microbiol. 101:215–221. doi:10.1016/j.vetmic.2004.03.020.

Dean, C.J., F. Peña-Mosca, T. Ray, B.J. Heins, V.S. Machado, P.J. Pinedo, L.S. Caixeta, and
N.R. Noyes. 2022. Evaluation of Contamination in Milk Samples Pooled From Independently
Collected Quarters Within a Laboratory Setting. Front. Vet. Sci. 9:818778.
doi:10.3389/fvets.2022.818778.

Dean, C.J., I.B. Slizovskiy, K.K. Crone, A.X. Pfennig, B.J. Heins, L.S. Caixeta, and N.R. Noyes.
2021. Investigating the cow skin and teat canal microbiomes of the bovine udder using different
sampling and sequencing approaches. J. Dairy Sci. 104:644–661. doi:10.3168/jds.2020-18277.

Ewels, P., M. Magnusson, S. Lundin, and M. Käller. 2016. MultiQC: summarize analysis results
for multiple tools and samples in a single report. Bioinforma. Oxf. Engl. 32:3047–3048.
doi:10.1093/bioinformatics/btw354.

Fernandes, L., M.L. Celestino, P.R. Menta, T.H. Silva, D. Paiva, T.L. Ribeiro, L.S. Caixeta, N.R.
Noyes, and V.S. Machado. 2022. Cow-related factors associated with intramammary infections
during the 1st week postpartum in primiparous dairy cows in certified organic herds. Vet. J.
282:105822. doi:10.1016/j.tvjl.2022.105822.

Green, M.J., A.J. Bradley, G.F. Medley, and W.J. Browne. 2008. Cow, Farm, and Herd
Management Factors in the Dry Period Associated with Raised Somatic Cell Counts in Early
Lactation. J. Dairy Sci. 91:1403–1415. doi:10.3168/jds.2007-0621.

Gurevich, A., V. Saveliev, N. Vyahhi, and G. Tesler. 2013. QUAST: quality assessment tool for
genome assemblies. Bioinforma. Oxf. Engl. 29:1072–1075. doi:10.1093/bioinformatics/btt086.

Jahan, N.A., S.M. Godden, E. Royster, T.C. Schoenfuss, C. Gebhart, J. Timmerman, and R.C.
Fink. 2021. Evaluation of the matrix-assisted laser desorption ionization time of flight mass
spectrometry (MALDI-TOF MS) system in the detection of mastitis pathogens from bovine milk
samples. J. Microbiol. Methods 182:106168. doi:10.1016/j.mimet.2021.106168.

Mahmmod, Y.S., I.C. Klaas, L. Svennesen, K. Pedersen, and H. Ingmer. 2018. Communications
of Staphylococcus aureus and non-aureus Staphylococcus species from bovine intramammary
infections and teat apex colonization. J. Dairy Sci. 101:7322–7333. doi:10.3168/jds.2017-14311.

McCubbin, K.D., E. de Jong, T.J.G.M. Lam, D.F. Kelton, J.R. Middleton, S. McDougall, S. De
Vliegher, S. Godden, P.J. Rajala-Schultz, S. Rowe, D.C. Speksnijder, J.P. Kastelic, and H.W.
Barkema. 2022. Invited review: Selective use of antimicrobials in dairy cattle at drying-off. J.
Dairy Sci. 105:7161–7189. doi:10.3168/jds.2021-21455.

Nakatsuji, T., T.H. Chen, S. Narala, K.A. Chun, A.M. Two, T. Yun, F. Shafiq, P.F. Kotol, A.
Bouslimani, A.V. Melnik, H. Latif, J.-N. Kim, A. Lockhart, K. Artis, G. David, P. Taylor, J.
Streib, P.C. Dorrestein, A. Grier, S.R. Gill, K. Zengler, T.R. Hata, D.Y.M. Leung, and R.L.
Gallo. 2017. Antimicrobials from human skin commensal bacteria protect against
Staphylococcus aureus and are deficient in atopic dermatitis. Sci. Transl. Med. 9:eaah4680.
doi:10.1126/scitranslmed.aah4680.

NMC. 2017. Laboratory Handbook on Bovine Mastitis. 3rd Edition. National Mastitis Council,
Madison, Wisconsin. National Mastitis Council, Madison, Wisconsin.

NMC. 2019. Mastitis control on organic dairy farms in the United States. National Mastitis
Council, Madison, Wisconsin.

Parks, D.H., M. Chuvochina, C. Rinke, A.J. Mussig, P.-A. Chaumeil, and P. Hugenholtz. 2022.
GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically
consistent, rank normalized and complete genome-based taxonomy. Nucleic Acids Res.
50:D785–D794. doi:10.1093/nar/gkab776.

Peña-Mosca, F., C. Dean, L. Fernandes, E. Doster, K. Sharpe, T. Ray, V. Feijoo, A. Antunes, C.
Baumann, T. Wehri, B. Heins, P. Pinedo, V. Machado, N. Noyes, and L. Caixeta. 2023a.
Associations between early lactation intramammary infections and udder health and performance
during the first 180 days in milk in first-lactation organic dairy cows (in review). J. Dairy Sci.

Peña-Mosca, F., C. Dean, V. Machado, L. Fernandes, P. Pinedo, E. Doster, B. Heins, K. Sharpe,
T. Ray, V. Feijoo, A. Antunes, C. Baumann, T. Wehri, N. Noyes, and L. Caixeta. 2023b.
Investigation of intramammary infections in primiparous cows during early lactation on organic
dairy farms. J. Dairy Sci. S0022-0302(23)00552–0. doi:10.3168/jds.2022-23036.

Piepers, S., G. Opsomer, H.W. Barkema, A. de Kruif, and S. De Vliegher. 2010. Heifers infected
with coagulase-negative staphylococci in early lactation have fewer cases of clinical mastitis and
higher milk production in their first lactation than noninfected heifers. J. Dairy Sci. 93:2014–
2024. doi:10.3168/jds.2009-2897.

Piepers, S., K. Peeters, G. Opsomer, H.W. Barkema, K. Frankena, and S. De Vliegher. 2011.
Pathogen group specific risk factors at herd, heifer and quarter levels for intramammary
infections in early lactating dairy heifers. Prev. Vet. Med. 99:91–101.
doi:10.1016/j.prevetmed.2011.02.010.

Pol, M., and P.L. Ruegg. 2007. Relationship Between Antimicrobial Drug Usage and
Antimicrobial Susceptibility of Gram-Positive Mastitis Pathogens. J. Dairy Sci. 90:262–273.
doi:10.3168/jds.S0022-0302(07)72627-9.

Rajala-Schultz, P., A. Nødtvedt, T. Halasa, and K. Persson Waller. 2021. Prudent Use of
Antibiotics in Dairy Cows: The Nordic Approach to Udder Health. Front. Vet. Sci. 8.

Reyher, K.K., D. Haine, I.R. Dohoo, and C.W. Revie. 2012. Examining the effect of
intramammary infections with minor mastitis pathogens on the acquisition of new intramammary
infections with major mastitis pathogens—A systematic review and meta-analysis. J. Dairy Sci.
95:6483–6502. doi:10.3168/jds.2012-5594.

Rowe, S.M., S.M. Godden, E. Royster, J. Timmerman, B.A. Crooker, and M. Boyle. 2019.
Cross-sectional study of the relationships among bedding materials, bedding bacteria counts, and
intramammary infection in late-lactation dairy cows. J. Dairy Sci. 102:11384–11400.
doi:10.3168/jds.2019-17074.

Ruegg, P.L. 2017. A 100-Year Review: Mastitis detection, management, and prevention. J. Dairy
Sci. 100:10381–10397. doi:10.3168/jds.2017-13023.

Ruegg, P.L. 2022. Realities, Challenges and Benefits of Antimicrobial Stewardship in Dairy
Practice in the United States. Microorganisms 10:1626. doi:10.3390/microorganisms10081626.

Schreiner, D.A., and P.L. Ruegg. 2003. Relationship between udder and leg hygiene scores and
subclinical mastitis. J. Dairy Sci. 86:3460–3465. doi:10.3168/jds.S0022-0302(03)73950-2.

Sordillo, L.M. 2018. Mammary Gland Immunobiology and Resistance to Mastitis. Vet. Clin.
North Am. Food Anim. Pract. 34:507–523. doi:10.1016/j.cvfa.2018.07.005.

Toledo-Silva, B., L. Beuckelaere, A. De Visscher, C. Geeroms, E. Meyer, S. Piepers, D. Thiry,
F. Haesebrouck, and S. De Vliegher. 2022. Novel Quantitative Assay to Describe In Vitro
Bovine Mastitis Bacterial Pathogen Inhibition by Non-aureus Staphylococci. Pathog. Basel
Switz. 11:264. doi:10.3390/pathogens11020264.

USDA. 2007. Dairy 2007, Part I: Reference of Dairy Cattle Health and Management Practices in
the United States, 2007. #N480.1007. USDA-APHIS-VS, CEAH, Fort Collins, CO.

Wiegand, I., K. Hilpert, and R.E.W. Hancock. 2008. Agar and broth dilution methods to
determine the minimal inhibitory concentration (MIC) of antimicrobial substances. Nat. Protoc.
3:163–175. doi:10.1038/nprot.2007.521.

Woudstra, S., N. Wente, Y. Zhang, S. Leimbach, M.K. Gussmann, C. Kirkeby, and V. Krömker.
2023. Strain diversity and infection durations of Staphylococcus spp. and Streptococcus spp.
causing intramammary infections in dairy cows. J. Dairy Sci. 106:4214–4231.
doi:10.3168/jds.2022-22942.

APPENDIX

Tables

Table 1. Descriptive characteristics of enrolled cases (n=34) and controls (n=34)

Item	Controls	Cases
Farm		
A	20.6% (7/34)	20.6% (7/34)
B	79.4% (27/34)	79.4% (27/34)
Parity		
First lactation	47.1% (16/34)	47.1% (16/34)
Second lactation	52.9% (18/34)	52.9% (18/34)
Case type		
Staph aureus only	-	44.1% (15/34)
SSLO only	-	44.1% (15/34)
SSLO and *Staph aureus*	-	11.8% (4/34)
Cows positive at calving	-	64.7% (22/34)
Days in milk new IMI	-	13.2 (12.0)
Days relative to calving teat apex sample †	-2.8 (15.2)	-1.2 (16.5)

681 Cows positive at calving: Cows with an IMI by *Staphylococcus aureus* or *Streptococcus* spp. and

682 *Streptococcus*-like organisms (SSLO) on the first postpartum sample. † Teat apex sample: gauze

683 sample extracted on the week prior to the first time a cow tested positive for an IMI.

685 **Table 2.** Bacterial counts on teat-apex (log10 CFU/square inch of gauze) of controls (referent)
686 and cases (n=71 teat apex samples).

Outcome	Estimated marginal means (SE)		Estimate (95%CI)	*P-value*
	Controls	Cases		
Total bacterial count	1.81 (0.12)	1.91 (0.12)	0.10 (-0.19, 0.40)	0.50
NASM	1.12 (0.18)	1.02 (0.18)	-0.09 (-0.54, 0.35)	0.67
Staph chromogenes	0.26 (0.12)	0.20 (0.12)	-0.05 (-0.34, 0.23)	0.71
Staph haemolyticus	0.21 (0.13)	0.21 (0.13)	-0.005 (-0.33, 0.32)	0.98
Staph sciuri	0.19 (0.10)	0.21 (0.10)	0.02 (-0.23, 0.27)	0.86
Staph succinus [a]	0.17 (0.09)	0.17 (0.09)	-0.004 (-0.24, 0.24)	1.00
Staph xylosus/*pseudoxylosus*	0.46 (0.09)	0.53 (0.10)	0.06 (-0.17, 0.30)	0.60
Other	0.20 (0.07)	0.12 (0.07)	-0.08 (-0.26, 0.09)	0.35

687 [a] *Staph succinus*: Model did not account for farm, because it was not present in farm B. [b] Other:

688 *Staph cohnii, Staph devreseii* and *Staph equorum.* NASM: non-*aureus Staphylococci* and

689 *Mammaliicocci.*

690 **Table 3.** Presence of non-*aureus Staphylococci* and *Mammaliicocci* (NASM) isolates on teat-
691 apex of controls (referent) and cases (n=71 teat apex samples).

Outcome	Presence on teat apex		OR (95%CI)	*P-value*
	Controls	Cases		
NASM	66.7% (24/36)	62.9% (22/35)	0.85 (0.32, 2.25)	0.74
Staph chromogenes	25.0% (9/36)	22.9% (8/35)	0.89 (0.29, 2.73)	0.84
Staph haemolyticus	25.0% (9/36)	22.9% (8/35)	0.89 (0.29, 2.73)	0.84
Staph sciuri	8.3% (3/36)	5.7% (2/35)	0.66 (0.08, 4.28)	0.66
Staph succinus [a]	11.1% (4/36)	11.4% (4/35)	1.03 (0.23, 4.72)	0.97
Staph xylosus/*pseudoxylosus*	11.1% (4/36)	17.1% (6/35)	2.17 (0.39, 14.47)	0.38
Other [b]	8.3% (3/36)	2.9% (1/35)	0.31 (0.01, 2.63)	0.29

[a] *Staph succinus*: Model did not account for farm, because it was not present in farm B. [b] Other:
Staph cohnii, Staph devreseii and *Staph equorum.*

Table 4. Taxonomic identification of isolates harbored from teat apex samples and submitted for
taxonomic identification using Matrix-Assisted Laser Desorption/Ionization-Time of Flight mass
spectrometer (n=275 isolates) and confirmation for non-*aureus Staphylococci* and
Mammaliicocci (NASM) using whole genome sequencing (n=83 isolates).

Taxonomy classification	Controls	Cases
Staph aureus	0% (0/134)	1.4% (2/141)
NASM	32.8% (44/134)	27.7% (39/141)
Staph chromogenes	9.0% (12/134)	7.1% (10/141)
Staph cohnii	0% (0/134)	0.7% (1/141)
Staph devreseii	1.5% (2/134)	0% (0/141)
Staph equorum	0.7% (1/134)	0% (0/141)
Staph haemolyticus	9.0% (12/134)	6.4% (9/141)
Staph pseudoxylosus	1.5% (2/134)	2.1% (3/141)
Staph sciuri	2.1% (3/134)	2.1% (3/141)
Staph succinus	4.5% (6/134)	4.3% (6/141)
Staph xylosus	4.5% (6/134)	5.0% (7/141)
Aerococcus viridans	0% (0/134)	1.4% (2/141)
Aerococcus sp.	0.8% (1/134)	2.8% (4/141)
Bacillus sp.	61.9% (83/134)	61.7% (87/141)
Gram-positive rod	4.5% (6/134)	5.0% (7/141)

699 **Table 5.** Relationship between species identification and minimum inhibitory concentration
700 (MIC) of non-*aureus Staphylococci* and *Mammaliicocci* (NASM) against *Staphylococcus aureus*
701 and *Streptococcus uberis* (n=79 isolates).

NASM taxonomy	*Staphylococcus aureus*		*Streptococcus uberis*	
	MIC (SE)	*P-value*	MIC (SE)	*P-value*
Staph chromogenes	3.32 (0.13) [a]	0.001	3.06 (0.16)	0.23
Staph haemolyticus	3.99 (0.14) [b]		3.63 (0.26)	
Staph sciuri	3.32 (0.24) [a]		3.49 (0.30)	
Staph succinus	2.92 (0.17) [a]		2.92 (0.22)	
Staph xylosus/pseudoxylosus	3.05 (0.14) [a]		3.05 (0.17)	
Other	4.39 (0.32) [b]		3.29 (0.49)	

Other NASM: *Staph cohnii, Staph devreseii, Staph equorum.* Different super indexes within the
same column indicate statistical differences. Intraclass correlation coefficient cow-ID *Staph*
aureus: Not available, *Strep uberis*: Not available.Other: *Staph cohnii, Staph devreseii* and *Staph*
equorum. Only isolates in which MIC was determined were included for analysis; this included
a total of 75 isolates for *Staph aureus*: *Staph chromogenes:* 100% (19/19), *Staph haemolyticus*:
81.0% (17/21), *Staph sciuri*: 100% (6/6), *Staph succinus*: 100% (11/11), *Staph*
xylosus/pseudoxylosus: 100% (18/18), other NASM: 75.0% (3/4). For *Streptococcus uberis* MIC
was determined in 63 isolates: *Staph chromogenes:* 100% (19/19), *Staph haemolyticus*: 33.0%
(7/21), *Staph sciuri*: 100% (6/6), *Staph succinus*: 100.0% (11/11), *Staph xylosus/pseudoxylosus*:
100% (18/18), other NASM: 50.0% (2/4).

Table 5.6. Relationship between non-*aureus Staphylococci* and *Mammaliicocci* (NASM)
identified species and isolates classification as the top 10 with the lowest minimum inhibitory
concentration (MIC) against *Staphylococcus aureus* and *Streptococcus uberis* (n=79 isolates)

Outcome	NASM taxonomy	Presence of isolates with top 10 lowest MIC	*P-value*
Inhibition *Staph aureus*	*Staph chromogenes*	10.5% (2/19)	0.04
	Staph haemolyticus	4.8% (1/21)	
	Staph sciuri	16.9% (1/6)	
	Staph succinus	45.5% (5/11)	
	Staph xylosus/pseudoxylosus	5.6% (1/18)	
	Other	0.0% (0/4)	
Inhibition *Strep uberis*	*Staph chromogenes*	10.5% (2/19)	0.06
	Staph haemolyticus	4.8% (1/21)	
	Staph sciuri	0% (0/6)	
	Staph succinus	45.5% (5/11)	
	Staph xylosus/pseudoxylosus	11.1% (2/18)	
	Other	0.0% (0/4)	

715   Other: *Staph cohnii, Staph devreseii* and *Staph equorum.*

716   *Figures*

718   **Figure 1.** Outline of the study with number of observations lost at each step. NAS: non-*aureus*
719   *Staphylococci* and *Mammaliicocci*. AMA: antimicrobial activity.

Figure 2. Kaplan meier plots showing presence of growth of *Staphylococcus aureus* and
Streptococcus uberis at different concentrations of non-*aureus Staphylococci* and
Mammaliicocci (NASM) on cases and controls (Referent group) (n=79 isolates). *Staphylococcus
aureus*: HR (95%CI): 0.92 (0.56, 1.49), *P* =0.72. *Streptococcus uberis*: HR (95%CI): 1.23 (0.74,
2.08), *P*=0.45.

Figure 3. Ranking of 79 non-*aureus Staphylococci* and *Mammaliicocci* (NASM) isolates by minimum inhibitory concentration (MIC) against *Staphylococcus aureus* and *Streptococcus uberis* harbored from teat apex stratified by case control status (panel A/B) and assigned species (panel C/D) (n=79 isolates). Undetermined MIC: NASM isolates that showed growth of *Staphylococcus aureus* or *Streptococcus uberis* at the maximum investigated concentration on media. SCH: *Staphylococcus chromogenes*, SHAEM: *Staphylococcus haemolyticus*, SSC: *Staphylococcus sciuri*, SSUC: *Staphylococcus succinus*, SXYL: *Staphylococcus xylosus/pseudoxylosus*, Other: *Staphylococcus cohnii, Staphylococcus devreseii* and *Staphylococcus equorum*.

Figure 4. Kaplan meier plots showing presence of growth of *Staphylococcus aureus* and
Streptococcus uberis at different concentrations of non-*aureus Staphylococci* and
Mammaliicocci (NASM) for the different species isolated in this study (n=79 isolates). SCH:
Staphylococcus chromogenes, SHAEM: *Staphylococcus haemolyticus*, SSC: *Staph sciuri*, SSUC:
Staphylococcus succinus, SXYL: *Staphylococcus xylosus/pseudoxylosus,* Other: *Staphylococcus*
cohnii, Staphylococcus devreseii and *Staphylococcus equorum.* Type III *P-value Staphylococcus*
aureus < 0.001. Type III *P-value Streptococcus uberis* < 0.001.

```
Teat apex culture

n = 71 samples
```

```
Isolates with missing
MALDI-TOF MS ID

n = 6 isolates
```

```
Isolates with MALDI-
TOF MS ID

n = 275 isolates
```

```
Isolates with other
taxonomy assigned

n = 192 isolates
```

```
Isolates identified as
NASM

n = 83 isolates
```

```
Isolates with missing
in-vitro AMA

n = 4 isolates
```

```
Isolates with quantified
in-vitro AMA

n = 79 isolates
```