Exploring the phylogenetic diversity and antimicrobial activity of non-*aureus staphylococci*

and *mammaliicocci* isolated from teat apices of organic dairy cows

F. Peña-Mosca [1,2], T. N. Gaire [1], C. Dean [1], P. Ferm [1], D. Manriquez [3], P. Pinedo [4], N. Noyes [1], L.

Caixeta [1]

[1] Department of Veterinary Population Medicine, University of Minnesota, St. Paul, 55108

[2] Department of Public and Ecosystem Health, Cornell University, Ithaca, 14853.

[3] AgNext, Colorado State University, Fort Collins, 80523

[4] Department of Animal Science, Colorado State University, Fort Collins, 80523

ABSTRACT

Prior studies have suggested that non-*aureus staphylococci* and *mammaliicoci* (**NASM**)

possess inhibitory activity against mastitis pathogens. However, their impact on udder health

outcomes and the mechanisms underlying this potential protective effect remain poorly

understood. Our first objective was to identify NASM species on teat apices of organic dairy

cows, assess their within-species phylogeny, and explore associations with presence of

intramammary infections (**IMI**) and genomic features, including antimicrobial peptides (**AMPs**),

virulence, and resistance genes. The second objective was to evaluate the *in vitro* antimicrobial

activity of NASM isolates against mastitis pathogens and examine its associations with

taxonomy, phylogeny, AMP genes, and IMI. Milk and teat apex swabs were collected weekly

from 114 cows on two organic farms. Milk was cultured to identify *Staphylococcus aureus*

(**SAU**) or *Streptococcus* spp. and *Streptococcus*-like organisms (**SSLO**) IMI. A case-control was

designed to include cows with and without SAU or SSLO IMI. For each selected cow, the teat

apex gauze swab collected during the week preceding IMI diagnosis (or corresponding time for

controls) was aerobically cultured, and the taxonomy of isolates was determined using mass

spectrometry. Isolates classified as NASM were subjected to whole genome sequencing using

Illumina MiSeq. The inhibitory activity of NASM isolates was tested against SAU and

Streptococcus uberis. Phylogenetic trees were constructed using Snippy and IQ-TREE. Genomes

were assembled and annotated to identify species, AMP genes, virulence, and antimicrobial

resistance markers. The *in vitro* antimicrobial activity of NASM varied across species and

between cows with and without an IMI. *Staphylococcus succinus* was the species most

frequently associated with highly inhibitory isolates, which were more prevalent in cows without

IMI (19.4% vs 5.8%). Organic dairy cow teat apices harbored multiple NASM species and

strains. All isolates had at least 1 AMP associated gene in their genome. *In vitro* antimicrobial

activity was generally unrelated to clade membership, except for isolates classified as

Staphylococcus succinus. *Staphylococcus aureus* had high virulence gene prevalence, while

NASM species showed lower, species-specific prevalence. This study advances understanding of

NASM antimicrobial activity and virulence potential.

Key words: Antimicrobial peptides; Whole genome sequencing; Phylogenetic Analysis; Udder

health; Mastitis.

39   **INTRODUCTION**

Mastitis is the leading cause of morbidity (USDA, 2007) and milk production loss in

dairy farms (Heikkila et al., 2018). Although preventive measures are important for the control

of mastitis, antimicrobials are still a major component of on-farm contagious mastitis control

(Ruegg, 2017). However, the availability of antimicrobials for dairy farmers and veterinarians is

at-risk due to concerns about antimicrobial resistance to medically important antibiotics and

policy restrictions on antimicrobial use in livestock (McCubbin et al., 2022; Rajala-Schultz et al.,

2021; Ruegg, 2022). This limitation is especially remarkable on United States organic-certified

dairy farms, where regulations for production of organic milk include a strict prohibition against

the use of antimicrobials and other synthetic substances (NMC, 2019).

Prior data from our group showed that *Staphylococcus aureus* (**SAU**) and *Streptococcus*

spp. had a high prevalence and persistence in the mammary gland of dairy cows in organic farms

in the first 35 days of lactation (Peña-Mosca et al., 2023). Moreover, this study showed that SAU

and *Streptococcus* spp. in early lactation were associated with a higher risk of high somatic cell

count (**SCC**) during the first 180 days of lactation in organic dairy cows (Peña-Mosca et al.,

2024). Non-*aureus staphylococci* and *mammaliicocci* (**NASM**) are also highly prevalent in milk

samples (De Buck et al., 2021; Peña-Mosca et al., 2023), but unlike SAU and *Streptococcus*, in

some studies, NASM IMI have been associated with increased milk production (Compton et al.,

2007a; Piepers et al., 2010); while others suggested no association between NASM IMI and milk

production (Nyman et al., 2018; Tomazi et al., 2015; Valckenier et al., 2020). The protective

effects of NASM, have been supported by results from experimental studies in which the

infusion into the mammary gland of NASM has been associated with decreased risk for IMI with

SAU and *Streptococcus* spp., suggesting a potential protective effect of NASM against these two

pathogens (Reyher et al., 2012). *In vitro* studies using isolates obtained from the mammary gland

of dairy cows (i.e., from milk or teat apex) have shown that NASM are capable of inhibiting the

growth of SAU (Carson et al., 2017; De Vliegher et al., 2004a; Toledo-Silva et al., 2022). Using

the cross-streak method for inhibitory testing, these studies reported that between 9.1% and

14.3% of the isolates were able to inhibit the growth of SAU (Carson et al., 2017; Toledo-Silva

et al., 2022). Recently, agar dilution methods have also been utilized to investigate the inhibitory

capabilities of NASM species on mastitis pathogens (Toledo-Silva et al., 2022). One potential

explanation for the *in vitro* antimicrobial activities is the ability of NASM to produce

antimicrobial peptides (**AMPs**) (Braem et al., 2014; Carson et al., 2017; Nascimento et al.,

2005), commonly referred to as bacteriocins or *Staphylococcins* (de Freire Bastos et al., 2020;

Newstead et al., 2020). While AMP-encoding genes on bovine NASM isolates have been

explored (Braem et al., 2014; Carson et al., 2017; Nascimento et al., 2005), further research is

needed to identify specific AMPs that could be targeted for development of products for mastitis

control. Lastly, although there has been significant expansion in our understanding of NASM

epidemiology in recent years, the number of studies that have comprehensively investigated the

entire genome of NASM are still limited (De Buck et al., 2021; Fergestad et al., 2021; Naushad

et al., 2019). This limits our ability to investigate associations between NASM genotypes,

phenotypes, and cow-level udder health outcomes. Therefore, more research is needed to

describe the genomic diversity of NASM isolates from dairy cows, including important

phenotypic drivers such as genes related to AMPs, virulence factors and antimicrobial resistance

genes. Understanding the presence and diversity of these genes could provide further insights

into the potential for NASM isolates to act as protective commensals or, conversely, as emerging

pathogens in the udder environment.

Our first objective was to identify the NASM species present on teat apices of organic

dairy cows and examine their within-species phylogeny, as well as explore associations with

intramammary infections (**IMI**) and genomic features, including genes related to AMPs,

virulence, and antimicrobial resistance. The second objective was to assess the *in vitro*

antimicrobial activity of NASM isolates cultured from teat apices of cows with and without IMI,

and explore its associations with NASM taxonomy, within-species phylogeny, the presence of

AMP genes, and IMI.

MATERIALS AND METHODS

This manuscript was prepared following the guidelines outlined in the Strengthening the

Reporting of Observational Studies in Epidemiology - Veterinary (STROBE-Vet) Statement

(Sargeant et al. in 2016). All study activities were approved by the University of Minnesota

Institutional Animal Care and Use Committee (IACUC) (Protocol Number: 1807: 36109A), and

Colorado State University IACUC (Protocol number: 1442). Laboratory activities were approved

by the University of Minnesota Institutional Biosafety Committee (IBC) protocol: 2205-40001H.

Inclusion criteria

Only cows from USDA organic certified dairy farms were enrolled in this study. Farms

from different regions of the United States and with different herd sizes were selected.

Enrollment was based on willingness to participate, and proximity to the Universities involved in

the study. This work was part of a larger research initiative to investigate potential associations

between the udder microbiome and udder health (Dean et al., 2021). Following the study design

to assess that objective, cows from 5 organic dairy farms were enrolled 8 weeks before calving

and followed up during the first 5 weeks of lactation (Peña-Mosca et al., 2023). In this

manuscript, data from only two of these farms were included, as teat apex samples for

bacteriological culture were only collected in these two farms due to limited research labor

availability. Specifically, this manuscript used samples collected from a subset of 114 cows

(Farm A [n =21] and Farm B [n =93]) out of the 596 cows enrolled in the main research project,

where both milk and teat apex samples were collected. Samples from Farm A were collected

between August 2019 and January 2020 and included all first-lactation cows that calved during

that period. Samples from Farm B were collected between February and July 2021 and included

both first and second lactation cows that calved throughout that interval.

Milk sample collection

After calving, quarter milk samples were collected weekly during the first 5 postpartum

weeks prior to the morning milking by members of the research team, following guidelines from

the National Mastitis Council (NMC, 2017). Upon removal of the pre-dipping disinfectant, 3 to 4

streams of milk were discarded from each quarter, and the teat apex of each quarter was

scrubbed with a gauze soaked in 70% ethanol. Milk samples were collected in a clockwise

direction, beginning with the right rear quarter, and ending with the right front quarter.

Approximately 10 mL of milk from each quarter were collected into separate 60 mL plastic vials

(Capitol Vials, Thermo Fisher Scientific). Samples were placed on ice immediately after

collection and stored in a freezer at -20 °C until further processing.

Milk pooling

Quarter milk samples were combined into composite samples using the following

protocol (Peña-Mosca et al., 2023). First, milk samples were thawed at 4 °C overnight. Next,

quarter milk samples were homogenized by inverting them upside down twice. Then, 2 mL was

extracted from each vial and dispensed into a single plastic vial inside a laminar hood. Lab-

pooled composite samples were then submitted to the Laboratory for Udder Health at the

University of Minnesota (St. Paul, MN, USA) for milk culture.

Milk culture

Using a cotton swab, approximately 100 μL of milk were plated onto Columbia CNA agar with 5% sheep blood and MacConkey agar. Agar plates were incubated in aerobic conditions at 37 °C for a total of 42 to 48 h. Plates were examined by a trained technician to evaluate the presence of growth of distinct isolates at 18 to 24 hours and 42 to 48 hours. Samples were defined as contaminated if more than 3 different isolates were identified and omitted from further analysis (Dean et al., 2022). Taxonomic assignments of cultured isolates were made using a Matrix-Assisted Laser Desorption/Ionization-Time of Flight (**MALDI-TOF**) mass spectrometer (**MS**) (Microflex; BrukerDaltonics Inc.) (Jahan et al., 2021). Mass spectra profiles produced from each isolate were matched against the Biotyper reference library. Confidence scores were used to assign genus- and species-level classifications (≥ 1.8 and ≥ 2.0 for genus- and species- classification, respectively) (Jahan et al., 2021). An IMI was defined as a composite sample containing one or more than one colony forming units (10 CFU/mL) of any cultured isolate.

Teat apex sample collection

Cows enrolled from two of the participating farms, where labor was available, were sampled weekly from 8 weeks before calving to 5 weeks after calving in the milking parlor prior to the morning milking. First, teats were visually inspected, and any visible debris was removed using a paper towel. Then, using a new pair of disposable gloves, a single sterile pre-moistened gauze was used to thoroughly swab the teat apices of all four teats. After sampling, each gauze was placed in individual tubes containing a pre-loaded solution of 50% glycerol. Gauze samples were transported in a cooler with ice and stored in -80 °C after arrival to research facilities.

Postpartum samples were collected before pre-dipping disinfectant solutions were applied to the

teat and before milk samples were collected.

Implementation of case control study

Milk culture results were used to select a subset of gauze samples based on the presence

or absence of an IMI caused by SAU or *Streptococcus* spp. and *Streptococcus*-like organisms

(**SSLO**) during the first 35 DIM. These mastitis pathogens were used to guide sample selection

because they are known to persist in the mammary gland and are strongly associated with

elevated SCC and impaired udder health outcomes (Peña-Mosca et al., 2024b). Cows with and

without an IMI were individually matched based on farm, lactation number, and the timing of

IMI detection (within ±7 DIM). Selected teat apex samples were processed for bacterial culture

and species identification as described below. For three cows without a sample available from

the week prior to IMI detection, two samples were processed from the closest available time

points. In total, 71 samples were collected from 68 cows (34 with an IMI and 34 without an

IMI).

Teat apex sample culture and species identification

The methods for culturing gauze samples collected from teat apex were adapted from

protocols utilized in the Laboratory for Udder Health at the University of Minnesota to process

towel samples (Rowe et al., 2019). Before sample processing and to reduce the possibility of

batch effects, sample IDs for each gauze sample were randomly sorted using Excel (Microsoft

Corporation). In accordance with this order, isolates were selected for each sampling processing

date in batches of 10 isolates. Gauze samples were thawed at 4 °C overnight and then transferred

into 50 mL round-bottom falcon tubes (Corning Inc.). Afterwards, 5 mL of Phosphate-Buffered

Saline (**PBS**; Gibco, ThermoFisher Scientific) was added to reduce viscosity. Samples were

homogenized carefully by turning tubes upside down twice and vortexing for 10 seconds at low

speed. Samples were allowed to stand for 10 minutes, and the homogenization process was

repeated once. Following homogenization, a single 1 in 10 serial dilution was done using PBS. A

100 uL aliquot from each sample (i.e. undiluted and 10^{-1}) was inoculated onto Mannitol salt agar

(Oxoid). Plates were then incubated in aerobic conditions for 24 hours at 37 °C. Following

incubation, bacterial groups were visually assessed and counted from the dilution plate with the

optimal number of colonies (25 to 250 per plate). Isolates of visually distinct colonies were then

picked from the plate using a sterile calibrated loop (Thermo Fisher Scientific), streaked onto

blood agar (Hardy Diagnostics) and incubated for 24 hours at 37 °C. After achieving pure

growth, the identity of representative colonies was determined using MALDI-TOF MS, using

similar methods to those described for isolates identified in milk samples. Non-*aureus*

staphylococci and *mammaliicocci* isolates were then preserved at -80 °C using commercial

cryovials (Hardy Diagnostics) until further analysis. NASM species-level taxonomy was further

investigated through whole genome sequencing, as reported below. Colony forming units (**CFU**)

per square inch of gauze were estimated for each distinct colony type based on the number of

colonies present on each plate type using a correction factor based on the used dilution.

Specifically, a 10x correction factor was applied for 10^{-1} plates, and this calculation also

accounted for inoculum size (10x), dilution with PBS (2x), and gauze size (1/4x).

Determination of in vitro inhibitory activity against Staphylococcus aureus and Streptococcus

uberis

The methods utilized in this study were based on procedures previously described by

others (Toledo-Silva et al., 2022). For each batch of isolates, on day 1, isolates previously

identified as NASM using MALDI-TOF MS were randomly sorted using Excel (Microsoft

Corporation). Following this procedure, isolates previously isolated from teat apex were grown

for 24 hours in blood agar. If evidence of mixed growth (more than 1 colony type) was present,

samples were replated. On day 2, one colony was picked with a 10 uL disposable loop and was

grown on brain heart infusion (**BHI**; Research Products International Corporation) media for 18

to 20 hours. The next day, 1 in 10 serial dilutions solutions were created using sterile PBS in a 96

well plate (Thermo Fisher Scientific). Bacterial counts (CFU/mL) were estimated by plating 100

uL of the serial dilutions on blood agar plates that were then aerobically incubated for 18 to 20 h

at 37 °C. One hundred microliters of bacterial suspensions containing NASM isolates at different

dilutions (10^{-2}, 10^{-3}, 10^{-4}, 10^{-5}, 10^{-6}, 10^{-7}) were added to 10 mL of 2% Tryptic soy agar (**TSA**;

Oxoid) inside Falcon round-bottom polypropylene tubes (Corning Inc.). The temperature was

examined using a laboratory digital thermometer (Thermco) before the inclusion of the 2% TSA.

The media were not used until the temperature reached 50 °C (Toledo-Silva et al., 2022). Tubes

were then homogenized and poured into pre-heated (37 °C) TSA plates. Following aerobic

incubation for 18 to 20 h, 10 uL of the bacterial suspensions containing approximately 10,000

CFU of SAU (American Type Culture Collection (**ATCC**) strain Seattle 1945; *Staphylococcus*

aureus subsp. aureus Rosenbach 25923 [TM]) and *Streptococcus uberis* (**SUB**; ATCC strain 01040

J; *Streptococcus uberis* Diernhofer BAA-854 [TM]) were inoculated onto the top of plates

containing different concentrations of NASM isolates and incubated for 18 to 20 h. For each

batch of samples, negative controls were included by incubating a 2% TSA agar plate with the

inclusion of 100 uL of BHI media only (instead of NASM isolates). Additionally, a 10 μL

droplet of PBS was added on top of the media on each plate to serve as an additional negative

control, ensuring the absence of PBS contamination. *Staphylococcus aureus* and SUB were

incorporated into the media as positive controls, and their self-inhibition was tested to explore

the determination of MIC driven by nutrient competition rather than *in vitro* inhibitory activity.

Growth of SAU and SUB (i.e., yes vs no) was evaluated by visual examination of each of the

plates and comparison with negative control plates within each batch. Minimum inhibitory

concentration (**MIC**) was defined as the minimum concentration of NASM (CFU/mL) on the 2%

TSA top layer able to inhibit the growth of SAU or SUB.

DNA extraction, library preparation and whole genome sequencing

Isolates were allowed to grow on BHI for 24 hours, centrifuged at 10,000 rpm for 5

minutes, and the pellet was collected in 2mL sterile polypropylene microcentrifuge tubes

(Thermo Fisher Scientific). Following this, DNA extraction was performed utilizing commercial

kits (Qiagen's DNeasy 96 Blood & Tissue Kit, Qiagen). Amount and quality of extracted DNA

was evaluated using PicoGreen fluorometer and NanoDrop spectrophotometer (Thermo Fisher

Scientific). Library preparation was performed using the DNA Prep Sample Preparation Kit

(Illumina Inc.). Resultant DNA was subjected to paired-end sequencing (2x300) utilizing the

Illumina MISEQ platform aiming for 50x coverage of *Staphylococcus* spp. genome.

Bioinformatics

Quality control of raw sequencing data was performed using FastQC (Andrews, 2010)

and MultiQC (Ewels et al., 2016). Trimmomatic was utilized to trim low quality base pairs and

Nextera XT adapter sequences from raw reads (Bolger et al., 2014). MultiQC was then utilized

to summarize the QC metrics during the filtering and trimming process. Trimmed sequence reads

were assembled into contigs using SPAdes version 3.15.5 (Bankevich et al., 2012). Assembly

quality was evaluated using Quast version 5.2.0 (Gurevich et al., 2013). Taxonomic

classification of contigs was performed using GTDB-Tk version 1.7.0 (Chaumeil et al., 2020)

and GTDB release 202 (Parks et al., 2022) as implemented in KBase (Arkin et al., 2018). The

ABRIcate pipeline (Seemann, 2023a). was used with default settings to identify antimicrobial

resistance genes (**ARGs**) from contigs using the MEGARes v2 reference database (Doster et al.,

2020). To identify virulence factors from DNA contigs, we used BLAST+ (Camacho et al.,

2009) and a custom comprehensive protein database (Naushad et al., 2019), previously shown to

be effective for identifying virulence genes in *Staphylococcus* genomes (Fergestad et al., 2021;

Naushad et al., 2019). Only BLAST+ results with an e-value of less than or equal to 10^{e-5}

(Choudhuri, 2014) and amino acid identity of greater than or equal to 35% (Rost, 1999) were

considered for analysis. The BAGEL4 online server with default settings was used to identify

gene clusters associated with the production of AMPs on contigs (van Heel et al., 2018). Snippy

was used with default settings (i.e., minimum coverage of 10x and a minimum variant fraction of

90%) to identify core single nucleotide polymorphisms (Seemann, 2023b) by aligning trimmed

reads from each of the isolates onto reference genomes obtained from Genbank (Clark et al.,

2016). The following reference genomes were used for each NASM species: *Staphylococcus*

chromogenes (**SCH**): GenBank accession number: GCA_002994305.1, *Staphylococcus*

haemolyticus (**SHAEM**): GenBank accession number: GCA_001611955.1, *Mammaliicocci*

sciuri (**MSCf**): GenBank accession number: GCA_002209165.2, *Staphylococcus succinus*

(**SSUC**): GenBank accession number: GCA_001902315.1, *Staphylococcus xylosus* (**SXYL**) and

Staphylococccus pseudoxylosus (**SPXYL**): GenBank accession number: GCA_000709415.1. As

a result of the limited availability of only one or two genomes for *Staphylococcus cohnii* (**SCO**) ,

Staphylococcus devrieseii (**SDEV**), and *Staphylococcus equorum* (**SEQ**)*,* SNP calling was not

performed on these species. The generated Snippy core files were used to build a maximum

likelihood tree using IQ-TREE, with 1,000 bootstrap replicates to assess the robustness of the

phylogenetic tree (Nguyen et al., 2015). Trees were visualized and annotated using iTOL

(Letunic and Bork, 2021).

Statistical analysis

All statistical analyses were performed in R version 4.3.2 (https://www.r-project.org/).

Code and output are available online: https://fepenamosca.github.io/nasm_antimicrobial_wgs/.

Association between IMI status and NASM colonization of teat apex

The relationship between IMI status and $\log_{10}$ -transformed bacterial counts from teat

apex samples was investigated using linear regression models. The relationship between IMI

status and presence of NASM on the teat apex samples (i.e., yes vs no) was investigated using a

generalized linear model with a "binomial" family and "logit" link (logistic regression). For

analysis, NASM taxonomy was determined based on the assigned taxonomy using MALDI-TOF

MS and whole genome sequencing. It was represented as a six-level multi-level variable (SCH,

SHAEM, MSC, SSUC, SXYL*,* and other rare NASM [SCO, SDEV, SEQ]).

Association between IMI status, NASM taxonomy and presence of NASM with in vitro

antimicrobial activity on teat apex

Cox proportional hazards regression was used to assess the association between presence

of IMI, NASM taxonomy and the concentration to NASM *in vitro* growth inhibition against

SAU and SUB. These models were utilized to estimate hazard ratios (**HR**) and their 95%

confidence intervals (**CI**), used to compare the inhibitory activity across groups, where higher

HR indicates higher NASM inhibitory activity and the presence of inhibition of mastitis

pathogens at lower NASM concentrations on the media. As some isolates exhibited no inhibitory

activity at the maximum concentration analyzed, observations were right-censored if inhibitory

activity remained absent at the highest NASM concentration examined (i.e., 10^{-2} dilution).

Proportional hazards assumption was tested by evaluating the Schoenfeld residuals. Since

multiple isolates were obtained from teat apices each cow, the non-independence of observations

within each enrolled animal was accounted for using the robust sandwich estimator. The NASM

MIC ($\log_{10}$, CFU/mL) against SAU and SUB was further explored and compared across cows

with and without an IMI and different NASM species using mixed linear regression. These

models incorporated cow-ID as a random effect to address the non-independence of observations

for isolates obtained from the same cow. Subsequently, isolates were ranked according to their

MIC (e.g., 1 to 77) and classified as being in the top 10 isolates with lowest MIC or not ("top

10"). The relationship between IMI status and presence of a "top 10" NASM on the teat apex

was investigated using a generalized linear model (logistic regression) with "binomial" family

and "logit" link. When performing this analysis by NASM species, we used the Fisher exact test

due to the absence of "top 10" isolates for some NASM species. The intraclass correlation

coefficient was estimated using the "performance" package (Lüdecke et al., 2021). Estimated

marginal means were estimated using the "emmeans" package in R (Lenth et al., 2024) .

Normality assumption was investigated by evaluation of model residuals and the use of quantile-

quantile plots. Multiple comparisons were accounted for with Tukey adjustment as implemented

in the "emmeans" package (Lenth et al., 2024). Farm-ID was forced into all models that

investigated the relationship between IMI status and NASM inhibitory activity against SAU and

SUB. Because farm ID was strongly associated with the prevalence of different NASM species

(i.e., each of the NASM species were in most cases highly prevalent in one farm but rarely found

in the other one), this variable was not included in the models comparing *in vitro* inhibitory

activity across different NASM species. Processing batch (i.e., the date at which gauze samples

were processed or antimicrobial activity of NASM was evaluated) and parity of the cow (i.e.,

first vs second lactation) were offered as potential confounders. Presence of confounding was

evaluated by examining the change in the estimates after adjustment by each confounder ([Crude

estimate – Adjusted estimate] / Adjusted estimate). Confounders that changed the estimates by

10% or more were kept in the models.

Association between the phylogeny, genotypic, and in vitro antimicrobial activity of NASM

isolated from the teat apex

Because NASM species were strongly associated with *in vitro* inhibitory activity and

presence of AMP gene clusters, all statistical analyses were performed separately for each of the

NASM species (SCH, SHAEM, MSC, SSUC, SXYL, SPXYL, Others [SCO, SDEV, SEQ]). A

clade, also known as a monophyletic group, was defined as a group of genomes on a

phylogenetic tree that included their most recent shared ancestor and their descendants (Kapli et

al., 2020). Only AMP gene clusters that were both present and absent in at least 2 isolates within

a given species were investigated using regression analysis. For modeling, the association

between IMI status, clade membership and the presence of AMP gene clusters was investigated

using logistic regression. The presence of AMP gene clusters across different NASM species,

and clades within species, was compared using the Fisher exact test. This approach was chosen

because, in several cases, gene clusters were found exclusively in a single clade or species. The

association between the presence of AMP gene clusters, clade membership, and the MIC of

NASM against SAU and SUB for each NASM species was investigated using linear regression.

 RESULTS

 Descriptive characteristics

  This study examined two USDA-organic certified farms, one in Minnesota (A) and one in

 Colorado (B), with varying herd sizes and housing systems. On Farm A, all cows included in this

 study (14/14) were in their first lactation, while in Farm B, roughly half (43/93) were in their

 first lactation, and half (50/93) were transitioning to their second lactation. Isolates from 68 cows

 (34 cows with an IMI and 34 cows without an IMI) from these two farms were included in this

 study. Fifteen of the cows with an IMI had a SAU-IMI, 15 had a SSLO-IMI, and 4 had an IMI

 caused by both microorganisms. In the first postpartum composite milk sample, 64.7% (22/34) of

 the cases were already positive for SAU or SSLO IMI. Teat apex samples from cows with and

 without an IMI were on average collected on similar days relative to calving (mean ±SD; no IMI:

 -2.8 ±15.2 vs IMI: -1.2 ±16.5).

 Association between IMI status and NASM colonization of teat apex

  NASM were isolated from the teat apex of dairy cows during the transition period in a

 similar proportion of those without (66.7% [22/36]) and with an IMI (62.9% [24/35]) (OR [95%

 CI]: 0.85 [0.32, 2.25], $P = 0.74$; **Table 1**). Likewise, NASM counts ($\log_{10}$ CFU/gauze inch^2) on

 the teat apex were similar between groups (Estimate [95% CI]: -0.09 [-0.51, 0.35], $P = 0.69$;

 Table 1). The distribution of NASM species differed by farm. *Staphylococcus chromogenes,*

 SHAEM, and SSUC showed a high prevalence in farm A (28.1% [16/57], 28.1% [16/57] and

 14.1% [8/57], respectively), but were rarely found on farm B (7.1% [1/14], 7.1% [1/14] and 0%

 [0/14], respectively). On the other hand, SXYL or SPXYL were found in more than half of the

 samples from farm B (57.1% [8/14]) but showed a low prevalence in farm A (3.5% [2/57]).

Association between IMI status, NASM taxonomy and presence of NASM with in vitro

antimicrobial activity on teat apex

A total of 273 isolates were obtained from the teat apices of the enrolled cows and submitted for identification using MALDI-TOF MS as shown in **Table 2**. Surprisingly, *Bacillus* spp. was the most frequently identified genus. Non-*aureus staphylococci mammaliicocci* were identified in 27.7% (39/141) and 31.8% (42/132) of the isolates obtained from the teat apex samples from cows with and without an IMI, respectively. The most frequent species identified within this group included SCH (8.1% [22/273]) and SHAEM (7.0% [19/273]). Results of MALDI-TOF MS and whole genome sequencing for NASM isolates are presented in supplementary materials (https://doi.org/10.6084/m9.figshare.25832938.v2). All isolates classified as SCH (n=20) or SHAEM (n=15) using MALDI-TOF MS showed similar taxonomy when analyzed using whole genome sequencing. Isolates initially classified as *Staphylococcus xylosus/saprophyticus* (n=12) using MALDI-TOF MS were subsequently reclassified *as* SXYL (n=8) or SPXYL (n=4) by whole genome sequencing. The remaining isolates (34/83; 41.0%) were not identified up to the species-level using MALDI-TOF MS. These isolates were identified using whole genome sequencing as SCH (n=2), SCO (n=1), SDEV (n=2), SEQ (n=1), SHAEM (n=4), SPXYL (n=1), MSC (n=6), SSUC (n=12), and SXYL (n=5).

We investigated the inhibitory activity of NASM included in a top layer inside the media at varying concentrations against SAU and SUB. In some cases, inhibitory activity was not observed even at the highest concentration tested. In fact, among the 77 NASM isolates evaluated, no inhibitory activity was observed for 5 isolates against SAU and 14 isolates against SUB (**Figure 1**). For these isolates with no inhibitory activity, the highest concentration tested averaged (mean±SD) 3.98 ± 0.45 $\log_{10}$ CFU/mL for SAU and 4.30 ± 0.52 $\log_{10}$ CFU/mL for SUB.

For isolates where inhibition was observed, MIC values did not differ between those from

cows with and without an IMI (**Table 3**). To account for the varying concentrations required to

reach inhibition and the presence of right-censored data (i.e., isolates without inhibition at the

maximum NASM concentration tested), we analyzed the data using survival analysis. The

overall presence of inhibitory activity was not different between isolates from cows with and

without an IMI (HR [95%CI] for SAU: 0.89 [0.56, 1.45]; for SUB: 1.17 [0.68, 1.97]; **Figure 2**).

To provide a reference for interpreting NASM's inhibitory activity, we tested SAU and

SUB against themselves by plating each pathogen on media containing varying concentrations of

the same microorganism. When SAU was tested as a control on the 2% TSA top layer, its MIC

was 3.42 $\log_{10}$ CFU/mL, and 51.9% (40/77) of NASM isolates had an MIC below this value,

including 52.5% (21/40) of isolates from cows without an IMI and 51.4% (19/37) from cows

with an IMI. Similarly, when SUB was evaluated, its MIC was 2.91 $\log_{10}$ CFU/mL, and 32.4%

(25/77) of NASM isolates had an MIC below this threshold. Among these isolates, 35.0%

(14/40) were collected from cows without an IMI, and 29.7% (11/37) were from cows with an

IMI

Building on these MIC distributions, we ranked the isolates based on their inhibitory

activity against SAU and SUB to identify those with the highest inhibitory activity (**Figure 1;**

Table 3). The "top 10" isolates with the lowest MIC ($\log_{10}$ CFU/mL) had a mean ± SD MIC of

2.41 ± 0.26 for SAU and 2.16 ± 0.32 for SUB, with MICs ranging from 1.92 to 2.66 for SAU

and 1.65 to 2.56 for SUB. These highly inhibitory isolates were more common in cows without

an IMI (19.4% [7/36]) compared to those with an IMI (5.8% [2/35]) (OR [95% CI]: 0.25 [0.04,

1.13], $P = 0.07$). However, an association with IMI status was not observed for cows harboring a

"top 10" isolate for SUB (controls: 11.1% [4/36] vs. cases: 14.3% [5/35]; OR [95% CI]: 1.33

[0.32, 5.83], $P = 0.69$).

The inhibitory activity of NASM against SAU and SUB varied significantly among

NASM species ($P < 0.001$; **Figure 2**). *Staphylococcus haemolyticus* and other less common

NASM species (e.g., SCO, SDEV, SEQ) exhibited lower inhibitory activity against SAU

compared to SCH (HR [95% CI]: 0.29 [0.12, 0.59], $P = 0.02$; and 0.15 [0.03, 0.71], $P < 0.01$,

respectively), MSC (0.21 [0.07, 0.58], $P < 0.001$; and 0.12 [0.03, 0.52], $P < 0.001$, respectively),

SSUC (0.15 [0.05, 0.42], $P < 0.001$; and 0.08 [0.02, 0.38], $P < 0.001$, respectively), and

SXYL/SPXYL (0.13 [0.05, 0.38], $P < 0.001$; and 0.08 [0.02, 0.29], $P < 0.001$, respectively). The

adjusted MIC ($\log_{10}$ CFU/mL) of NASM against SAU across species ranged from 2.92 to 4.39

(**Table 3**).

Similarly, when inhibition of SUB was assessed, SHAEM displayed reduced inhibitory

activity compared to SCH (HR [95% CI]: 0.10 [0.02, 0.50]), MSC (0.17 [0.06, 0.54]), SSUC

(HR [95% CI]: 0.13 [0.03, 0.55]), and SXYL/SPXYL (HR [95% CI]: 0.11 [0.03, 0.47], all $P <$

0.001). The inhibitory activity against SUB of rare NASM species was not different from that of

other NASM species identified in this study (**Figure 2**). The adjusted MIC ($\log_{10}$ CFU/mL) of

NASM species against SUB ranged between 2.92 and 3.51 (**Table 3**).

The distribution of isolates ranked by MIC against SAU and SUB is shown in **Figure 3**.

The proportion of "top 10" inhibitory isolates differed across NASM species ($P = 0.05$ for SAU

and $P = 0.07$ for SUB; **Table 3**). Among the species with lower inhibitory activity, only 5.3%

(1/19) of SHAEM isolates and 0% (0/5) of rare NASM species isolates were classified as "top

10" against SAU and/or SUB. In contrast, 45.5% (5/11) of SSUC isolates were "top 10"

inhibitory isolates against both SAU and SUB (**Figure 3** and **Table 3**)

Association between the phylogeny, genotypic, and in vitro antimicrobial activity of NASM

isolated from the teat apex

<u>*Assembly quality of NASM isolates submitted for WGS*</u>

All isolates classified as NASM were submitted for WGS. After quality filtering the

genomes were assembled using spades. Across the NASM assembled genomes, an average $\pm$ SD

of 48.4 $\pm$ 137 contigs were identified. The mean $\pm$ SD genome size was 2,610,000 $\pm$ 276,000,

with the N50 value averaging 585,000 $\pm$ 485,000 across all genomes. The longest contig

exhibited an average size of 883,000 $\pm$ 367,000 across all genomes.

<u>*Phylogeny and antimicrobial activity*</u>

Five phylogenetic trees were constructed representing six different NASM species (SCH,

SHAEM, MSC, SSUC, SXYL/SPXYL). The assigned NASM taxonomy showed a strong

association with the presence of various AMP gene clusters. For example, it was observed that

AIP class I gene clusters (NCBI accession number: WP_001093929.1) were present in all

isolates identified as SCH and SHAEM, as well as in half of those identified as MSC, but were

absent in SSUC, PXYL, and SPSXYL (*P*<0.001). Similarly, putative bacteriocin 193.2

(Genbank accession number: AJ002203.2) gene clusters were highly prevalent in SXYL or SEQ,

but were not detected in other NASM species (*P*<0.001).

Staphylococcus chromogenes

Among the 21 SCH isolates (**Figure 3**), 12 and 9 were from cows without and with IMI,

respectively. The majority (95.2% [20/21]) were from farm A. Genes for AIP class I and II

(NCBI accession: WP_001094921.1) were detected in 76.2% (16/21) and 23.8% (5/21) of SCH

isolates, respectively. Sactipeptides (InterPro: PF04055) and subtilosin A (NCBI: NP_391616.1)

gene clusters were found in 100% (21/21) and 61.9% (13/21) of SCH isolates. Lanthipeptides

class II (Interpro accession number: PF05147) gene clusters were identified in 1 out of the 21

isolates (4.8%). AIP class I and II genes (NCBI: WP_001094921.1) were detected in 76.2%

[16/21] and 23.8% [5/21] of isolates, respectively. AIP class I was more frequent in isolates from

cows without IMI (91.7% [11/12]) than with IMI (55.6% [5/9]) ($P = 0.08$), while an opposite

pattern was observed for AIP class II (no IMI: 8.3% [1/12] vs. IMI: 44.4% [4/9]; $P = 0.08$).

Sactipeptides (InterPro accession number: PF04055) were present in all isolates. Subtilosin A

(NCBI: NP_391616.1) was detected in isolates from cows without IMI (41.7% [5/12]) and with

IMI (33.3% [3/9]) ($P = 0.70$). Lanthipeptides class II (InterPro: PF05147) were rare (4.8%

[1/21]). None of these gene clusters were associated with MIC against SAU or SUB ($P > 0.05$ for

all linear regression models).

Phylogenetic analysis of the 21 SCH genomes revealed two major clades, with one clade

containing just two isolates, both from the same cow (SCH7 and SCH14 from cow 46, **Figure**

3). The largest subclade contained 9 isolates (SCH1, SCH3, SCH5, SCH8, SCH15, SCH16,

SCH17, SCH18, SCH20). The genomes within this clade were characterized by the consistent

presence of subtilosin A (100% [9/9]) and AIP class I (100% [9/9]) gene clusters, which were

less common among other SCH isolates (subtilosin A: 33% [4/12], $P < 0.001$; AIP class I: 58.3%

[7/12], $P = 0.045$). In contrast, AIP class II gene clusters were not identified in this clade but

identified in 41.7% (5/12) of others ($P = 0.045$). This clade also contained the only 2 SCH

isolates classified as "top 10" with highest antimicrobial activity. However, MIC (CFU/mL)

against SAU or SUB did not differ between this clade and others (SAU: estimate [95%CI]: 0.05

[-0.45, 0.56]), $P=0.82$; SUB: -0.35 [-0.93, 0.22], $P=0.51$). These 21 isolates were obtained from

16 cows, with 5 cows carrying two SCH isolates each. In some cases, both isolates from the

same cow showed minimal divergence and clustered within the same clade (e.g., cow 461: SCH7

and SCH14; cow 863: SCH3 and SCH18; cow 1086: SCH2 and SCH21). In others, isolates from

different clades were recovered from the same cow on the same day (e.g., cow 87835: SCH5 and

SCH11; cow 89386: SCH5 and SCH15), suggesting the presence of distinct strains within the

SCH population colonizing the teat apex.

Staphylococcus haemolyticus

A total of 19 isolates were identified as SHAEM (**Figure 4**). Out of these isolates, 10 and

9, were from cows without and with an IMI, respectively. Most, 94.7% [18/19]) were from farm

A. Genes encoding AIP class I and sactipeptides were found in all isolates (100% [19/19]), while

lanthipeptides class IV were detected in 5.3% [1/19]

The 19 SHAEM isolates were grouped into 2 distinct clades, with one clade comprising

just two isolates, as shown in **Figure 4.** The largest subclade contained 10 isolates: SHAEM2,

SHAEM3, SHAEM4, SHAEM7, SHAEM9, SHAEM13, SHAEM14, SHAEM16, SHAEM18,

SHAEM19. Half of these isolates were obtained from cows without IMI (5/10) and half from

cows with an IMI (5/10). Like other SHAEM isolates, this clade showed limited inhibitory

activity, with many unable to inhibit SAU (30.0% [3/10]) or SUB (90.0% [9/10]). Interestingly, a

small subclade containing 2 isolates (SHAEM 8 and SHAEM10), encompassed isolates obtained

from cows in different farms and exhibiting different inhibitory activity against SAU or SUB.

We did not investigate potential associations with genes associated with AMP production

because they either had low prevalence or were present in all SHAEM isolates.

Mammaliicoccus sciuri

Six isolates were identified as MSC (**Figure 5**)*.* Half of these isolates were cultured from

controls (3/6) and half from cases (3/6). In addition, 4 of these isolates were obtained from cows

in farm A and 2 from cows in farm B. The genomes of these isolates contained genes that

encoded AIPs class I (no IMI: 66.7% [2/3], IMI: 33.3% [1/3]) (*P*=0.42), sactipeptides (present in

all isolates) and putative bacteriocin 193.2 (Genbank accession number: AJ002203.2) (no IMI:

33.3% [1/3], IMI: 66.7% [2/3], P=1.00). No associations were found between presence of these

genes, and the MIC against SAU or SUB (*P*>0.05 for all linear models).

Phylogenetic analysis showed that isolates were grouped in 2 distinct clades, one

including isolates from farm A and the other one isolates from farm B (**Figure 5**). Isolates from

farm A formed two different subclades, one including two isolates obtained from cows with IMI

(MSC1 and MSC2) and the other from cows without an IMI (MSCI5 and MSCI6). Putative

bacteriocin 193.2 was identified in all isolates isolated from cows in farm A (4/4) but not

identified in those from farm B (0/2) (*P*=0.08). No differences were found in the MIC against

SAU or SUB across these clades (*P*=0.92 and *P*=0.21, respectively).

Staphylococcus succinus

A total of 12 isolates were identified as SSUC (**Figure 6**). Half (6/12) were isolated from

cows without IMI, and the remaining half (6/12) from cows with IMI. Furthermore, all SSUC

isolates were found in cows from farm A. Genes encoding AIPs class II and Sactipeptides were

identified in 100% (19/19) of the isolates. Carnolysins were identified in 1 out of the 12 isolates

(8.3%). *Staphylococcus succinus* isolates were grouped in multiple clades (**Figure 4**). One of

these clades contained isolates SSUC1, SSUC2, SSUC3, SSUC4, SSUC5, SSUC10 and

SSUC12, and the MICs of these isolates were lower than SSUC isolates in other clades (estimate

[95%CI]: MIC SAU: -1.06 [-1.66, -0.46], *P*=0.003; MIC SUB: -1.06 [-2.14, 0.02], *P*=0.05). This

clade also contained all of the SSUC isolates classified as "top 10" with lowest MIC against both

SAU and SUB. Lastly, an interesting finding was that there were 2 cows from which multiple

isolates were recovered (cow 94706: SSUC1, SSUC7 and SSUC12; cow 1333: SSUC6 and

SSUC10), and these isolates were distributed across distinct subclades, indicating the presence of

distinct strains within the teat apex of a given cow. Associations between the presence of genes

related to the production of AMPs were not investigated because they showed a low prevalence

or were present in all SSUC isolates.

Staphylococcus xylosus & Staphylococcus pseudoxylosus

We identified 13 isolates as SXYL and 5 as SPXYL (**Figure 7**). Among SXYL isolates,

53.8% [7/13] were collected from cows without IMI and 46.2% [6/13] from cows with IMI. For

SPXYL, 60.0% [3/5] were harbored from cows without IMI and 40.0% [2/5] with IMI. Most

SXYL (92.3% [12/13]) and SPXYL (80.0% [4/5]) isolates were from farm B. The phylogenetic

tree showed that isolates could be divided into 3 distinct clades, two of which contained isolates

from SXYL (SXYL clade A: SXYL1, SXYL2, SXYL4, SXYL10, SXYL11, SXYL13, SXYL14;

SXYL clade B: other SXYL isolates) and the other of which contained isolates classified as

PSXYL. The MIC ($\log_{10}$ CFU/mL) against SAU was (mean±SE) 3.14±0.18, 2.98±0.17, and

3.06±0.20 for isolates in SXYL clade A and B and for those identified as SPXYL, respectively

(Type III *P-value* = 0.81). In addition, for isolates belonging to SXYL clades A and B and those

identified as SPXYL, the MIC against SUB was 3.12±0.26, 2.97±0.28, and 3.06±0.30,

respectively (Type III *P-value* = 0.92). Sactipeptides and subtilosin A were present in all SXYL

and SPXYL genomes. On the other hand, AIPs class II were present in 53.8% (7/13) of the

SXYL isolates (clade A: 33.3% [2/6]; clade B: 71.4% [5/7] and 100% [5/5] of the PSXYL

isolates) (*P*=0.07). In addition, AIPs class III (NCBI accession number: WP_000735197.1) were

highly prevalent in SXYL clade A (66.7% [4/6]); showed a low prevalence in SXYL clade B

(14.3% [1/7]); and were not identified in SPXYL genomes (P=0.03). Putative bacteriocin 193.2

genes were identified in all isolates from SXYL clade B (100%, 6/6), had a lower prevalence in

SXYL clade A (14.3%, 1/7), and were not found in SPXYL isolates (P<0.001). None of these

genes were associated with MIC against SAU or SUB (P>0.05 for all linear regression models).

Genetic determinants of virulence

We investigated the prevalence of virulence genes associated with adherence,

exoenzymes, immune evasion, iron uptake, and toxin production across NASM and SAU, with

SAU carrying a higher number of these genes. Among adherence-related genes (n=37), 81.1%

were detected, with most highly prevalent in SAU, while NASM species showed lower

prevalence but retained some key adhesion and biofilm formation genes (**Figure 8**). Of the 22

exoenzyme genes examined, SAU carried 77.3%, whereas NASM species harbored fewer,

though nucleases and aureolysins were widely present (**Figure 8**). Immune evasion genes (n=35)

were nearly ubiquitous in SAU (97.1%) but varied across NASM, with capsular genes

widespread and immune evasion factors such as staphylococcal binding immunoglobulin protein

and staphylococcal complement inhibitor mainly found in SAU and SCH (**Figure 9**). For iron

uptake and metabolism (n=29), all genes were present in SAU, while NASM species harbored

fewer, with prevalence varying distribution (**Figure 9**). Among 87 enterotoxin and exotoxin

genes, 59 were identified, with enterotoxins exclusive to SAU and exotoxins detected in SAU

and SCH but absent in other NASM (**Figure 10**). Of the 36 toxin-related genes, SAU carried

75.0%, while NASM species had fewer, except for some cases such as exfoliative toxin type C,

which was present in all NASM isolates (**Figure 10**), beta phenol-soluble modulins, which were

widespread except in SCH and MSC, and type VII secretion system proteins, primarily detected

in SCH (**Figure 10**).

Genetic determinants of antimicrobial resistance

In this research, we identified the presence of 28 different ARGs (**Figure 11**).

Staphylococcus aureus had 15 of these genes in its genome, whereas each NASM species had

between 1 and 6 ARGs in at least one isolate. Ninety-seven percent of the isolates (81/83)

showed the presence of at least 1 ARG. The most prevalent ARGs were 23S rRNA

methyltransferases (**RLMH**) (53.0% [44/83]), involved in the synthesis of an rRNA

methyltransferase related to resistance against lincosamides and macrolides. Another gene that

exhibited high prevalence was aminoglycoside O-phosphotransferases (**APH3-PRIME**) (50.6%

[42/83]), which encodes an aminoglycoside-modifying enzyme. Additionally, macrolide

phosphotransferases (**MPHC**), a gene associated with macrolide resistance, was present in 16.9%

(14/83) of the isolates. Other genes, including those linked to penicillin resistance (e.g., class A

betalactamases [**blaZ gene**]) or methicillin resistance (e.g., penicillin binding proteins [**mecA**,

mecB, **mecI**]), demonstrated low prevalence (<10%) among SAU and NASM isolates.

DISCUSSION

This study improved our understanding of NASM genomes isolated from teat apices of

organic dairy cows around parturition, with a focus on their species distribution, phylogeny,

antimicrobial activity, virulence, and antimicrobial resistance. Multiple NASM species, and

occasionally multiple strains within the same species, were identified on teat apices. NASM

isolates ranked among the "top 10" with lowest MIC against SAU were more prevalent in cows

without IMI compared to those with IMI. Although all NASM genomes contained at least one

AMP, no association was found between presence of AMPs in NASM genomes and *in vitro*

inhibitory activity against SAU or SUB. One clade within the SSUC group comprised 7 NASM

isolates, with the majority demonstrating high *in vitro* antimicrobial activity. However, for other

species, no association was found between clade membership and *in vitro* antimicrobial activity.

Lastly, our results showed that although SAU genomes showed a higher prevalence of virulence

genes and ARGs, all NASM contained at least some of these genes, with a distribution that

appeared to be in most cases species-dependent.

Relationship between teat apex colonization by* NASM, *in vitro inhibitory activity and

presence of intramammary infections

The prevalence of NASM on teat apex was similar to that reported by other studies in

conventional dairy farms (De Visscher et al., 2016; Mahmmod et al., 2018). Consistent with

previous studies, we found a high prevalence of Staphylococcus chromogenes and observed

differences in NASM species distribution across farms (De Visscher et al., 2016; Mahmmod et

al., 2018). These differences have been previously attributed to differences in management

practices across farms (De Buck et al., 2021) and are not surprising, given the differences in the

prevalence and distribution of different microorganisms that cause IMI on organic dairy farms

(Peña-Mosca et al., 2023).

The counts or the presence of NASM on teat apex were not associated with the presence

of SAU or SSLO IMI. These results are inconsistent to those reported by prior studies, in which

prepartum colonization of teat apices by NASM was related to a lower risk of postpartum IMI

(Piepers et al., 2011) and subclinical mastitis (De Vliegher et al., 2003). These findings suggest

that, in our study population, the mere presence of NASM on the teat apex did not appear to be a

major factor influencing IMI risk. Other characteristics, such as the antimicrobial activity of

NASM isolates, may be more relevant, as proposed by previous research (Reyher et al., 2012),

and supported by studies reporting the in vitro antimicrobial activity of NASM isolates (Braem et

al., 2014; Carson et al., 2017; Toledo-Silva et al., 2022).

We observed wide variation in the in vitro antimicrobial activity among NASM isolates.

A considerable number of isolates showed no inhibitory activity at the maximum concentration

tested, suggesting the absence of effective antimicrobial activity in these isolates. Moreover, we

did not find evidence of a difference in the average inhibitory activity of NASM isolates against

SAU or SUB between cows with and without an IMI. One possible explanation is that average

antimicrobial activity may not be a major determinant of IMI risk, and that strong antimicrobial

activity against specific pathogens may play a more important role in preventing infection

(Nakatsuji et al., 2017). Supporting this idea, we found that transitioning dairy cows without an

IMI were more likely to carry NASM isolates with the lowest MIC values against SAU

compared to cows with an IMI. Although these findings should be interpreted with caution given

the small sample size and wide confidence intervals, they suggest the potential for a protective

effect of NASM isolates exhibiting strong *in vitro* antimicrobial activity against SAU on IMI risk

by SAU or SSLO. Our findings are in line with human studies, where isolates with high

phenotypic antimicrobial activity were commonly found on the skin of healthy individuals but

were rarely identified in individuals with SAU skin infections (Nakatsuji et al., 2017). As a

whole, our results suggest that commensal NASM may contribute to defense against mammary

gland pathogen invasion, although further research is needed to confirm these observations.

Relationship between* NASM taxonomy *and presence of in vitro antimicrobial activity

Nine NASM species were identified from teat apices of organic dairy cows. Consistent

with previous studies, we observed important differences in antimicrobial activity across NASM

species (Braem et al., 2014; Carson et al., 2017). Earlier studies using the cross-streak method

reported antimicrobial activity in 9.1% to 13% of NASM isolates (Braem et al., 2014; Carson et

al., 2017). In contrast, we used agar dilution methods (Wiegand et al., 2008), which aligns with

the cross-streak method but provides quantitative measurements (Toledo-Silva et al., 2022).

To our knowledge, quantitative MIC data for NASM against mastitis pathogens, have

only been reported for SCH, *Staphylococcus epidermidis*, and *Staphylococcus simulans*, with

MIC values against SAU ranging from 5.10 to 8.24 log10 CFU/mL (Toledo-Silva et al., 2022).

In our study, we observed variation in NASM inhibitory activity against SAU and SUB.

Staphylococcus haemolyticus exhibited little to no inhibition, consistent with prior findings

(Carson et al., 2017). In contrast, SSUC showed the strongest inhibitory activity, with nearly

50% of its isolates ranked among the "top 10" lowest MIC values. This proportion was notably

higher than that of other species, where it did not exceed 20%, and differs from prior reports

(Carson et al., 2017). These discrepancies may be explained by the observation that antimicrobial

activity in previous human research appeared to be strain-specific rather than species-specific

(Nakatsuji et al., 2017). Hence, it is plausible that the antimicrobial activity of NASM isolates is

influenced more by strain-level variation rather than by species species.

Association between the phylogeny, genotypic, and in vitro antimicrobial activity of NASM

isolated from the teat apex

To investigate strain-level variation, we constructed five phylogenetic trees to examine

the evolutionary relationships among the most common NASM species in this study. For most

species, clade membership was not associated with IMI status or *in vitro* antimicrobial activity.

In SSUC, however, we observed that isolates with the strongest *in vitro* inhibitory activity were

concentrated within a single clade. In particular, one clade containing 7 genomes showed a lower

MIC against both SAU and SUB compared to isolates in other clades. Additionally, all 5 isolates

classified as part of the "top 10" group for highest inhibitory activity within SSUC were found in

this clade, suggesting that specific genome features of these SSUC isolates may confer enhanced

protection against SAU and SUB.

Our phylogenetic analysis revealed that a single cow could harbor multiple strains of a

NASM species within a single sample (**Figures 2-5**; e.g., cow 171: SHAEM1 and SXYL1; cow

87835: SCH5 and SCH11). This finding aligns with previous research showing that the teat apex

can be colonized by multiple strains of the same NASM species (Woudstra et al., 2023). Multi-

strain infections have been documented in various human and animal pathogens, including

Staphylococcus spp. (Balmer and Tanner, 2011) and similar patterns have been observed among

mastitis-causing pathogens colonizing the mammary gland. Specifically multiple strains of SAU

(Smith et al., 2005) or SUB (Zadoks et al., 2003) have been isolated from the mammary gland.

However, evidence suggests that when multiple strains colonize the mammary gland, a single

strain often dominates, as demonstrated with SUB (Pryor et al., 2009). The implications of multi-

strain infections are still not well understood, but it has been hypothesized it could have an

impact on disease dynamics (Balmer and Tanner, 2011). This could have important

consequences for interpreting mastitis pathogen epidemiology and for decision-making on dairy

farms (Exel et al., 2022). Previous studies have documented inter-strain differences in

epidemiology (Zadoks et al., 2011), virulence potential (França et al., 2021; Monistero et al.,

2020), and antimicrobial activity (Nakatsuji et al., 2017). These findings emphasize the

importance of considering strain-level variation in study design. Methodological approaches that

do not differentiate between strains and species may yield ambiguous or inaccurate results and

could partly explain contradictory findings regarding NASM epidemiology (De Buck et al.,

2021).

Genotypic and in vitro antimicrobial activity

One intriguing observation was the ubiquitous presence of AMP-associated gene clusters

in all examined genomes. This finding contradicts a prior study conducted with isolates from

conventional dairies, reporting that 21.5% of isolates contained gene clusters associated with the

production of AMPs (Carson et al., 2017). Several factors could explain this discrepancy,

including updates to the databases used for gene annotation in recent years. Differences in

methodology, such as the inclusion of additional steps to assess AMP gene cluster completeness

(Carson et al., 2017), may also have contributed. Lastly, caution is warranted, as the presence of

AMP-associated genes does not guarantee their expression or the production of functional

peptides (Sánchez-Romero and Casadesús, 2020).

None of the detected AMP gene clusters were associated with *in vitro* antimicrobial

activity. The prevalence of AIP class I was higher in cows without an IMI, whereas AIP class II

was more prevalent in cows with an IMI, specifically among isolates classified as SCH. Auto-

induced peptides are important for SAU pathogenesis, since they play an important role for the

induction of expression of virulence factors through the accessory regulation system(Le and

Otto, 2015; Wang and Muir, 2016). In NASM, the production of AIPs can repress the expression

of this system in SAU, thus reducing the virulence potential of this pathogen (Canovas et al.,

2016; Toledo-Silva et al., 2021). Extensive research has explored the use of AIP-mediated

quorum sensing interference as a strategy for controlling SAU infections (Gray et al., 2013; Tan

et al., 2018). Therefore, a higher prevalence of AIPs might be expected in cows without an IMI,

consistent with the observed pattern for AIP class I in SCH, but not for AIP class II. Additional

research is required to gain a deeper understanding of the potential applications of AIPs in the

management of mastitis.

Genetic determinants of virulence

We investigated 207 virulence genes involved in adherence, immune evasion, iron

uptake, and toxin production. These genes were highly prevalent in SAU but less common

among NASM. Adherence-related genes (i.e., aap and bap) were more prevalent than in previous

studies (Fergestad et al., 2021; Naushad et al., 2019; Srednik et al., 2017; Tremblay et al., 2013).

The IcaA gene, essential for biofilm formation, was detected in SCH and SSUC, aligning with

earlier reports (Tremblay et al., 2013). Fibronectin binding factors (fnbA and fnbB), critical for

SAU invasion (Campos et al., 2022), were nearly ubiquitous in SAU and also present in SCH,

SXYL, and SPSXYL.

Short-chain dehydrogenase/reductase genes, key for *Staphylococcus* pathogenesis

(França et al., 2021), were identified in all isolates. Genes encoding exoenzymes that facilitate

host invasion were commonly detected (Tam and Torres, 2019). Nucleases and aur genes were

found in all NASM isolates, similar to previous research (Fergestad et al., 2021; Naushad et al.,

2019). Other genes exhibited variable prevalence. For example, adenosine synthase A was

exclusive to SCH, SHAEM, and SDEV. The triacyglycerol lipase and glycerol ester hydrolase

genes were highly prevalent in SCH, SHAEM, and SXYL but less common in SSUC and absent

in MSC, consistent with previous findings (Naushad et al., 2019).

Capsular genes, which help bacteria resist phagocytosis (Cheung et al., 2021), were

present in all isolates, as reported in previous studies (Fergestad et al., 2021; Naushad et al.,

2019). Genes encoding staphylococcal binding immunoglobulin protein and staphylococcal

complement inhibitor, both important for SAU immune evasion (Rooijakkers et al., 2005; Smith

et al., 2011), were detected in SAU and SCH but not in other NASM species, aligning with prior

research (Naushad et al., 2019).

Enterotoxins were identified only in SAU, contrasting with previous studies that reported

them in NASM (Fergestad et al., 2021; Naushad et al., 2019). Staphylococcal exotoxins were

only identified in SAU and SCH. This finding is interesting considering that SCH is the most

prevalent NASM isolated from milk samples (De Buck et al., 2021). Some studies suggest SCH

negatively impacts udder health (Fry et al., 2014; Supré et al., 2011; Valckenier et al., 2020),

while others indicate a protective effect (De Vliegher et al., 2003). However, the presence of

exotoxin genes in all SCH isolates raises questions about the extent of strain-level differences in

virulence. Exfoliative toxin type C was prevalent in almost all NASM isolates, consistent with

one prior study (Fergestad et al., 2021), but not with another investigation (Naushad et al., 2019).

Beta phenol-soluble modulins (**PSMs**), important for pathogenesis, were widespread in most

NASM except SCH and MSC. Intriguingly, alpha PSMs, which are considered highly potent,

were absent in both SAU and NASM. Type VII secretion system proteins are important for

virulence of SAU and other gram-positive pathogens (Spencer and Doran, 2022), and were

primarily found in SCH, consistent with previous studies (Fergestad et al., 2021; Naushad et al.,

2019).

Genetic determinants of antimicrobial resistance

In this investigation, we identified that nearly all NASM isolates (97.7%) contained at

least 1 ARG in their genome. In most cases, distribution of ARGs was species dependent; and if

a given ARG was present within a species, it was typically present across all isolates in that

species.

The most prevalent ARG was *rlmH*, a gene that halts action of macrolides and

lincosamides, whose identification was not reported in NASM in a prior study (Nobrega et al.,

2018). In addition, the prevalence of *mphC*, another gene involved in resistance to macrolides,

showed a 16.9% prevalence in our study, which is comparable to that of a prior study (Nobrega

et al., 2018).

Half of the isolates carried the *aph(3′)-III* gene, encoding an aminoglycoside-inactivating

enzyme, a much higher prevalence than reported in a previous study where these ARGs were

nearly absent (Nobrega et al., 2018).

Penicillin resistance in SAU is typically mediated by beta-lactamase production (Nobrega

et al., 2018). Previous studies have shown various degrees of penicillin resistance in both SAU

and NASM isolates (Fergestad et al., 2021; Nobrega et al., 2018; Oliver and Murinda, 2012).

Methicillin resistance in SAU strains, mediated by acquisition of the staphylococcal

chromosomal cassette containing *mec* genes, remains a major concern for both human and

animal health(Holmes and Zadoks, 2011; McCarthy et al., 2012; Patel et al., 2021; Paterson et

al., 2014). In our study, the prevalence of mec genes was low, with *mecA* detected only in SSC

isolates. Previous studies also reported a low prevalence of *mecA* in MSC, although it has

occasionally been identified in other NASM species (Fergestad et al., 2021; Nobrega et al.,

2018).

Internal validity

This study investigated the quantitative antimicrobial activity of NASM isolates against

SAU and SUB. While this method represents a novel approach that has only recently been

described for its use on mastitis pathogens, it allows for the investigation of more subtle

variations in the inhibitory activity of bacteria (Toledo-Silva et al., 2022); however, it also has

certain limitations.

For instance, the novelty of these methods in mastitis research, coupled with the absence

of predetermined MIC cut-off points, complicates the interpretation of results. This makes it

challenging to discern whether the absence or presence of inhibition is due to antimicrobial

activity or merely a competition for nutrients that creates unfavorable conditions for the growth

of target microorganisms in the media (De Vliegher et al., 2004b; Toledo-Silva et al., 2022). To

overcome this limitation, we tested the self-inhibition of SAU and SUB, and we found that a

considerable number of NASM isolates were able to inhibit the growth of these mastitis

pathogens at lower concentrations than those necessary for self-inhibition of SAU and SUB,

suggesting the presence of inhibitory activity. In addition, it is important to highlight that the

utilized culture-based methods captured data from a few morphologically distinct isolates from

each teat apex sample (De Visscher et al., 2016). These data were used to draw conclusions

about the counts and prevalence of inhibitory NASM on the teat apex, which may not accurately

represent the diversity or relative abundance of inhibitory NASM on the teat apex (Dean et al.,

2024). This limitation is particularly relevant given that multiple species and strains can

simultaneously colonize the teat apex, as was observed in our study.

In this investigation, steps to control for potential technical and biological confounders

were taken. This included the randomization of samples for processing, matching by important

confounders, and assessing the magnitude of confounding by including these variables in the

models used in this study.

Nonetheless, it is vital to understand that some of our results could be still biased by other

unmeasured confounders. For instance, cow-level factors that increased the likelihood of

presence of IMI such as abnormally high or low body condition score, milk leakage (Fernandes

et al., 2022), dysfunctional immune response (Sordillo, 2018), or dirty udders (Compton et al.,

2007b; Schreiner and Ruegg, 2003) could have influenced our results. Moreover, the small

sample size and resulting imprecision of estimates could hinder our ability to make inference

from the results in this exploratory study. The use of composite milk samples represents a

potential limitation of this observational study, considering the lower sensitivity of milk culture

to detect IMI (Reyher and Dohoo, 2011; Toft et al., 2019). However, by culturing composite

milk samples from each cow multiple times and defining IMI positivity as the presence of an IMI

in at least one sample, we addressed this limitation, improving milk culture sensitivity for IMI

detection (Buelow et al., 1996). Nevertheless, evaluating exposure and outcome at the cow-level

poses a challenge, as it is possible that NASM with *in vitro* inhibitory activity could be present

on the teat apex of one quarter while mastitis pathogens lead to an IMI in a different quarter,

compromising the validity of this investigation (Bravata and Olkin, 2001).

Lastly, one major limitation of our study was the fact that almost two thirds of SAU and

SSLO IMI were detected in the first postpartum milk sample. This suggests that IMI may have

been acquired before calving. This finding is not surprising given the challenges to control

mastitis without antibiotics (NMC, 2019) and the increased prevalence of SAU on organic dairy

farms (Cicconi-Hogan et al., 2013; Pol and Ruegg, 2007). All the above limits our ability to

determine if colonization of NASM with antimicrobial activity preceded the acquisition of SAU

or SSLO IMI. Therefore, it is conceivable that other factors not explored in this study, such as

the presence of a close-up pen with poor bedding management, high stocking density, high fly

presence, and presence of IMI carried across lactations for multiparous cows, may be influencing

the results and explaining why some of these animals calved with IMI (De Vliegher et al., 2012;

Fernandes et al., 2022; Green et al., 2008).

External validity

This study was conducted in two organic dairy farms located in Minnesota and Colorado.

Considering the substantial differences in the prevalence NASM species across different dairy

farms in both this and previous studies (De Visscher et al., 2016; Mahmmod et al., 2018), it is

prudent to restrict the generalization of results to dairy farms managed under similar conditions.

In addition, because antimicrobial activity differed across different NASM species and

even within the species, results should be confirmed including a larger number of isolates

representative of different species that can be found on the mammary gland of dairy cows.

Finally, to address the wide heterogeneity of NASM antimicrobial activity, strain-level

identification might be warranted in future studies to further investigate bacterial clades and

antimicrobial peptides associated with the presence of *in vitro* antimicrobial activity.

One important limitation of our approach is that we focused our investigation on DNA

sequences, while it is known that the presence of a gene does not necessarily guarantee its

expression or the production of a protein (Sánchez-Romero and Casadesús, 2020). Hence, it is

plausible that this could explain some of the inconsistencies between the genotypes (e.g.,

presence of AMP gene clusters) and phenotypes (e.g., *in vitro* inhibitory activity). In addition, it

is conceivable that the AMPs responsible for the observed *in vitro* antimicrobial activity against

SAU or SUB in our study's isolates have not been characterized, are not present in the utilized

database (i.e., Bagel 4), or could not be identified due to bioinformatic challenges (e.g., the lack

of homology in the genes encoding precursor peptides for AMPs) (Morton et al., 2015).

827   Another limitation of our study was the fact that in the majority of the dairy cows the IMI

was already present on the first postpartum sample after calving. This limits our ability to assess

the temporality between the presence of AMP gene clusters or *in vitro* inhibitory activity and the

onset of new IMI by SAU or SSLO. Lastly, the low sample size, and small number of farms

limits our ability to make inferences about the different NASM species, especially those with

low prevalence (n<5). This is especially important, considering the substantial variation in the

distribution of NASM species and strains across different farms. Therefore, results from this

study should only be generalized to the NASM species identified in the current study and

extrapolation to other species should be done with caution.

CONCLUSIONS

837   Teat apices of organic dairy cows around parturition harbored multiple NASM species,

and in some cases, multiple strains within the same species. NASM isolates classified among the

"top 10" with the lowest MIC against SAU were more frequently found in cows without SAU or

SSLO IMI compared to those with an IMI in early lactation. At least one AMP-associated gene

was identified in all NASM isolates, indicating a high prevalence across species; however, their

presence was not associated with *in vitro* (i.e., phenotypic) antimicrobial activity. The

reconstructed phylogenetic tree for each NASM species contained multiple clades, indicating a

non-clonal population that may reflect the population structure of the cows both within and

across the two farms, and/or the evolutionary history of these bacteria within the two farms.

NASM in vitro antimicrobial activity differed across species but, apart from SSUC, was not

associated with clade membership. Virulence genes were highly prevalent in SAU, but in most

cases showed a low prevalence in NASM species, with a distribution that appeared to be species

dependent.

ACKNOWLEDGEMENTS

This study was funded by the Organic Agriculture Research and Extension Initiative

(OREI) from the National Institute of Food and Agriculture (grant number: 2018-51300-28563).

The first author, Felipe Peña-Mosca, was partially funded by Fulbright and Agencia Nacional de

Investigación e Innovación from Uruguay (scholarship number: POS_FUL_2019_1_1008441).

The authors thank the Laboratory for Udder Health at the University of Minnesota for

performing milk cultures and for their assistance in taxonomic identification of isolates. We

convey our appreciation to the University of Minnesota Genomics Core (UMGC) for their

assistance in DNA extraction, library preparation, and sequencing support, as well as to the

Minnesota Supercomputing Institute (MSI) for supplying the necessary resources for data

storage and computation. We also want to express our deep gratitude to the numerous students,

faculty, and staff for their assistance with sample collection and laboratory work. Additionally,

we extend our thanks to the owners and managers of the herds who made this study possible.

Felipe Peña-Mosca sampled in Minnesota, conducted laboratory work, cleaned and wrangled the

data, conducted bioinformatics and statistical analysis, and prepared the initial and final draft of

the manuscript. Tara Gaire conducted laboratory work and edited the manuscript. Chris Dean

sampled in Minnesota, collaborated on bioinformatics, and edited the manuscript. Peter Ferm

collaborated on bioinformatics and edited the manuscript. Pablo Pinedo conceptualized the

study, coordinated sampling in Colorado, and edited the manuscript. Diego Manriquez sampled

in Colorado and edited the manuscript. Noelle Noyes conceptualized the study, acquired funding,

collaborated on bioinformatics, coordinated sampling and laboratory work, and edited the

manuscript. Luciano Caixeta conceptualized the study, acquired funding, coordinated sampling

and laboratory work, and edited the manuscript. The authors have not stated any conflicts of

interest.

DATA AVAILABITY

875 The raw sequence data generated in this study has been submitted to the Sequence Read Archive

876 on NCBI (Accession: PRJNA1070794).

REFERENCES

Andrews, S., 2010. FastQC A Quality Control tool for High Throughput Sequence Data
[WWW Document]. URL https://www.bioinformatics.babraham.ac.uk/projects/fastqc/
(accessed 2.23.25).

Arkin, A.P., Cottingham, R.W., Henry, C.S., Harris, N.L., Stevens, R.L., Maslov, S., Dehal,
P., Ware, D., Perez, F., Canon, S., Sneddon, M.W., Henderson, M.L., Riehl, W.J., Murphy-Olson,
D., Chan, S.Y., Kamimura, R.T., Kumari, S., Drake, M.M., Brettin, T.S., Glass, E.M., Chivian, D.,
Gunter, D., Weston, D.J., Allen, B.H., Baumohl, J., Best, A.A., Bowen, B., Brenner, S.E., Bun, C.C.,
Chandonia, J.-M., Chia, J.-M., Colasanti, R., Conrad, N., Davis, J.J., Davison, B.H., DeJongh, M.,
Devoid, S., Dietrich, E., Dubchak, I., Edirisinghe, J.N., Fang, G., Faria, J.P., Frybarger, P.M.,
Gerlach, W., Gerstein, M., Greiner, A., Gurtowski, J., Haun, H.L., He, F., Jain, R., Joachimiak, M.P.,
Keegan, K.P., Kondo, S., Kumar, V., Land, M.L., Meyer, F., Mills, M., Novichkov, P.S., Oh, T.,
Olsen, G.J., Olson, R., Parrello, B., Pasternak, S., Pearson, E., Poon, S.S., Price, G.A.,
Ramakrishnan, S., Ranjan, P., Ronald, P.C., Schatz, M.C., Seaver, S.M.D., Shukla, M., Sutormin,
R.A., Syed, M.H., Thomason, J., Tintle, N.L., Wang, D., Xia, F., Yoo, H., Yoo, S., Yu, D., 2018. KBase:
The United States Department of Energy Systems Biology Knowledgebase. Nat Biotechnol 36,
566–569. https://doi.org/10.1038/nbt.4163

Balmer, O., Tanner, M., 2011. Prevalence and implications of multiple-strain infections.
The Lancet Infectious Diseases 11, 868–878. https://doi.org/10.1016/S1473-3099(11)70241-9

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin,
V.M., Nikolenko, S.I., Pham, S., Prjibelski, A.D., Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N., Tesler, G.,
Alekseyev, M.A., Pevzner, P.A., 2012. SPAdes: A New Genome Assembly Algorithm and Its
Applications to Single-Cell Sequencing. J Comput Biol 19, 455–477.
https://doi.org/10.1089/cmb.2012.0021

Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30, 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Braem, G., Stijlemans, B., Van Haken, W., De Vliegher, S., De Vuyst, L., Leroy, F., 2014. Antibacterial activities of coagulase-negative staphylococci from bovine teat apex skin and their inhibitory effect on mastitis-related pathogens. J Appl Microbiol 116, 1084–1093. https://doi.org/10.1111/jam.12447

Bravata, D.M., Olkin, I., 2001. Simple Pooling versus Combining in Meta-Analysis. Eval Health Prof 24, 218–230. https://doi.org/10.1177/01632780122034885

Buelow, K.L., Thomas, C.B., Goodger, W.J., Nordlund, K.V., Collins, M.T., 1996. Effect of milk sample collection strategy on the sensitivity and specificity of bacteriologic culture and somatic cell count for detection of *Staphylococcus aureus* intramammary infection in dairy cattle. Preventive Veterinary Medicine 26, 1–8. https://doi.org/10.1016/0167-5877(95)00518-8

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T.L., 2009. BLAST+: architecture and applications. BMC Bioinformatics 10, 421. https://doi.org/10.1186/1471-2105-10-421

Campos, B., Pickering, A.C., Rocha, L.S., Aguilar, A.P., Fabres-Klein, M.H., de Oliveira Mendes, T.A., Fitzgerald, J.R., de Oliveira Barros Ribon, A., 2022. Diversity and pathogenesis of Staphylococcus aureus from bovine mastitis: current understanding and future perspectives. BMC Vet Res 18, 115. https://doi.org/10.1186/s12917-022-03197-5

Canovas, J., Baldry, M., Bojer, M.S., Andersen, P.S., Gless, B.H., Grzeskowiak, P.K., Stegger, M., Damborg, P., Olsen, C.A., Ingmer, H., 2016. Cross-Talk between Staphylococcus aureus and Other Staphylococcal Species via the agr Quorum Sensing System. Frontiers in Microbiology 7.

Carson, D.A., Barkema, H.W., Naushad, S., De Buck, J., 2017. Bacteriocins of Non-aureus Staphylococci Isolated from Bovine Milk. Appl Environ Microbiol 83, e01015-17. https://doi.org/10.1128/AEM.01015-17

Chaumeil, P.-A., Mussig, A.J., Hugenholtz, P., Parks, D.H., 2020. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. Bioinformatics 36, 1925–1927. https://doi.org/10.1093/bioinformatics/btz848

Cheung, G.Y.C., Bae, J.S., Otto, M., 2021. Pathogenicity and virulence of Staphylococcus aureus. Virulence 12, 547–569. https://doi.org/10.1080/21505594.2021.1878688

Choudhuri, S., 2014. Chapter 6 - Sequence Alignment and Similarity Searching in Genomic Databases: BLAST and FASTA**The opinions expressed in this chapter are the author's own and they do not necessarily reflect the opinions of the FDA, the DHHS, or the Federal Government., in: Choudhuri, S. (Ed.), Bioinformatics for Beginners. Academic Press, Oxford, pp. 133–155. https://doi.org/10.1016/B978-0-12-410471-6.00006-2

Cicconi-Hogan, K.M., Gamroth, M., Richert, R., Ruegg, P.L., Stiglbauer, K.E., Schukken, Y.H., 2013. Risk factors associated with bulk tank standard plate count, bulk tank coliform count, and the presence of Staphylococcus aureus on organic and conventional dairy farms in the United States. Journal of Dairy Science 96, 7578–7590. https://doi.org/10.3168/jds.2012-6505

Clark, K., Karsch-Mizrachi, I., Lipman, D.J., Ostell, J., Sayers, E.W., 2016. GenBank. Nucleic Acids Res 44, D67–D72. https://doi.org/10.1093/nar/gkv1276

Compton, C.W.R., Heuer, C., Parker, K., McDougall, S., 2007a. Epidemiology of mastitis
in pasture-grazed peripartum dairy heifers and its effects on productivity. J Dairy Sci 90, 4157–
4170. https://doi.org/10.3168/jds.2006-880

Compton, C.W.R., Heuer, C., Parker, K., McDougall, S., 2007b. Risk factors for peripartum
mastitis in pasture-grazed dairy heifers. J Dairy Sci 90, 4171–4180.
https://doi.org/10.3168/jds.2006-882

De Buck, J., Ha, V., Naushad, S., Nobrega, D.B., Luby, C., Middleton, J.R., De Vliegher, S.,
Barkema, H.W., 2021. Non-aureus Staphylococci and Bovine Udder Health: Current
Understanding and Knowledge Gaps. Frontiers in Veterinary Science 8.

de Freire Bastos, M. do C., Miceli de Farias, F., Carlin Fagundes, P., Varella Coelho, M.L.,
2020. Staphylococcins: an update on antimicrobial peptides produced by staphylococci and their
diverse potential applications. Appl Microbiol Biotechnol 104, 10339–10368.
https://doi.org/10.1007/s00253-020-10946-9

De Visscher, A., Piepers, S., Haesebrouck, F., De Vliegher, S., 2016. Teat apex
colonization with coagulase-negative Staphylococcus species before parturition: Distribution
and species-specific risk factors. Journal of Dairy Science 99, 1427–1439.
https://doi.org/10.3168/jds.2015-10326

De Vliegher, S., Fox, L.K., Piepers, S., McDougall, S., Barkema, H.W., 2012. Invited review:
Mastitis in dairy heifers: nature of the disease, potential impact, prevention, and control. J Dairy
Sci 95, 1025–1040. https://doi.org/10.3168/jds.2010-4074

De Vliegher, S., Laevens, H., Barkema, H.W., Dohoo, I.R., Stryhn, H., Opsomer, G., de
Kruif, A., 2004a. Management Practices and Heifer Characteristics Associated with Early
Lactation Somatic Cell Count of Belgian Dairy Heifers. Journal of Dairy Science 87, 937–947.
https://doi.org/10.3168/jds.S0022-0302(04)73238-5

De Vliegher, S., Laevens, H., Devriese, L.A., Opsomer, G., Leroy, J.L.M., Barkema, H.W.,
de Kruif, A., 2003. Prepartum teat apex colonization with Staphylococcus chromogenes in dairy
heifers is associated with low somatic cell count in early lactation. Vet Microbiol 92, 245–252.
https://doi.org/10.1016/s0378-1135(02)00363-2

De Vliegher, S., Opsomer, G., Vanrolleghem, A., Devriese, L.A., Sampimon, O.C., Sol, J.,
Barkema, H.W., Haesebrouck, F., de Kruif, A., 2004b. In vitro growth inhibition of major mastitis
pathogens by Staphylococcus chromogenes originating from teat apices of dairy heifers. Vet
Microbiol 101, 215–221. https://doi.org/10.1016/j.vetmic.2004.03.020

Dean, C.J., Peña-Mosca, F., Ray, T., Heins, B.J., Machado, V.S., Pinedo, P.J., Caixeta, L.S.,
Noyes, N.R., 2022. Evaluation of Contamination in Milk Samples Pooled From Independently
Collected Quarters Within a Laboratory Setting. Front Vet Sci 9, 818778.
https://doi.org/10.3389/fvets.2022.818778

Dean, C.J., Peña-Mosca, F., Ray, T., Wehri, T.J., Sharpe, K., Antunes, Jr., A.M., Doster, E.,
Fernandes, L., Calles, V.F., Bauman, C., Godden, S., Heins, B., Pinedo, P., Machado, V.S., Caixeta,
L.S., Noyes, N.R., 2024. Exploring associations between the teat apex metagenome and
Staphylococcus aureus intramammary infections in primiparous cows under organic directives.
Applied and Environmental Microbiology 90, e02234-23. https://doi.org/10.1128/aem.02234-23

Dean, C.J., Slizovskiy, I.B., Crone, K.K., Pfennig, A.X., Heins, B.J., Caixeta, L.S., Noyes, N.R.,
2021. Investigating the cow skin and teat canal microbiomes of the bovine udder using different

sampling and sequencing approaches. Journal of Dairy Science 104, 644–661.
https://doi.org/10.3168/jds.2020-18277

Doster, E., Lakin, S.M., Dean, C.J., Wolfe, C., Young, J.G., Boucher, C., Belk, K.E., Noyes,
N.R., Morley, P.S., 2020. MEGARes 2.0: a database for classification of antimicrobial drug,
biocide and metal resistance determinants in metagenomic sequence data. Nucleic Acids Res 48,
D561–D569. https://doi.org/10.1093/nar/gkz1010

Ewels, P., Magnusson, M., Lundin, S., Käller, M., 2016. MultiQC: summarize analysis
results for multiple tools and samples in a single report. Bioinformatics 32, 3047–3048.
https://doi.org/10.1093/bioinformatics/btw354

Exel, C.E., Halasa, T., Koop, G., Steeneveld, W., Lam, T.J.G.M., Benedictus, L., Gussmann,
M., 2022. A stochastic modelling approach to determine the effect of diverse Staphylococcus
aureus strains on the economic and epidemiological outcomes of mastitis intervention
strategies in dairy cattle. Preventive Veterinary Medicine 199, 105566.
https://doi.org/10.1016/j.prevetmed.2021.105566

Fergestad, M.E., Touzain, F., De Vliegher, S., De Visscher, A., Thiry, D., Ngassam
Tchamba, C., Mainil, J.G., L'Abee-Lund, T., Blanchard, Y., Wasteson, Y., 2021. Whole Genome
Sequencing of Staphylococci Isolated From Bovine Milk Samples. Front Microbiol 12, 715851.
https://doi.org/10.3389/fmicb.2021.715851

Fernandes, L., Celestino, M.L., Menta, P.R., Silva, T.H., Paiva, D., Ribeiro, T.L., Caixeta,
L.S., Noyes, N.R., Machado, V.S., 2022. Cow-related factors associated with intramammary
infections during the 1st week postpartum in primiparous dairy cows in certified organic herds.
The Veterinary Journal 282, 105822. https://doi.org/10.1016/j.tvjl.2022.105822

França, A., Gaio, V., Lopes, N., Melo, L.D.R., 2021. Virulence Factors in Coagulase-
Negative Staphylococci. Pathogens 10, 170. https://doi.org/10.3390/pathogens10020170

Fry, P.R., Middleton, J.R., Dufour, S., Perry, J., Scholl, D., Dohoo, I., 2014. Association of
coagulase-negative staphylococcal species, mammary quarter milk somatic cell count, and
persistence of intramammary infection in dairy cattle. Journal of Dairy Science 97, 4876–4885.
https://doi.org/10.3168/jds.2013-7657

Gray, B., Hall, P., Gresham, H., 2013. Targeting agr- and agr-Like Quorum Sensing
Systems for Development of Common Therapeutics to Treat Multiple Gram-Positive Bacterial
Infections. Sensors 13, 5130–5166. https://doi.org/10.3390/s130405130

Green, M.J., Bradley, A.J., Medley, G.F., Browne, W.J., 2008. Cow, Farm, and Herd
Management Factors in the Dry Period Associated with Raised Somatic Cell Counts in Early
Lactation. Journal of Dairy Science 91, 1403–1415. https://doi.org/10.3168/jds.2007-0621

Gurevich, A., Saveliev, V., Vyahhi, N., Tesler, G., 2013. QUAST: quality assessment tool
for genome assemblies. Bioinformatics 29, 1072–1075.
https://doi.org/10.1093/bioinformatics/btt086

Holmes, M.A., Zadoks, R.N., 2011. Methicillin Resistant S. aureus in Human and Bovine
Mastitis. J Mammary Gland Biol Neoplasia 16, 373–382. https://doi.org/10.1007/s10911-011-
9237-x

Jahan, N.A., Godden, S.M., Royster, E., Schoenfuss, T.C., Gebhart, C., Timmerman, J.,
Fink, R.C., 2021. Evaluation of the matrix-assisted laser desorption ionization time of flight mass

spectrometry (MALDI-TOF MS) system in the detection of mastitis pathogens from bovine milk
samples. Journal of Microbiological Methods 182, 106168.
https://doi.org/10.1016/j.mimet.2021.106168

Kapli, P., Yang, Z., Telford, M.J., 2020. Phylogenetic tree building in the genomic age. Nat
Rev Genet 21, 428–444. https://doi.org/10.1038/s41576-020-0233-0

Le, K.Y., Otto, M., 2015. Quorum-sensing regulation in staphylococci—an overview.
Front Microbiol 6, 1174. https://doi.org/10.3389/fmicb.2015.01174

Lenth, R.V., Bolker, B., Buerkner, P., Giné-Vázquez, I., Herve, M., Jung, M., Love, J.,
Miguez, F., Riebl, H., Singmann, H., 2024. emmeans: Estimated Marginal Means, aka Least-
Squares Means. [WWW Document]. URL https://cran.r-
project.org/web/packages/emmeans/index.html (accessed 2.17.25).

Letunic, I., Bork, P., 2021. Interactive Tree Of Life (iTOL) v5: an online tool for
phylogenetic tree display and annotation. Nucleic Acids Research 49, W293–W296.
https://doi.org/10.1093/nar/gkab301

Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P., Makowski, D., 2021. performance:
An R Package for Assessment, Comparison and Testing of Statistical Models. JOSS 6, 3139.
https://doi.org/10.21105/joss.03139

Mahmmod, Y.S., Klaas, I.C., Svennesen, L., Pedersen, K., Ingmer, H., 2018.
Communications of Staphylococcus aureus and non-aureus Staphylococcus species from bovine
intramammary infections and teat apex colonization. J Dairy Sci 101, 7322–7333.
https://doi.org/10.3168/jds.2017-14311

McCarthy, A.J., Lindsay, J.A., Loeffler, A., 2012. Are all meticillin-resistant Staphylococcus
aureus (MRSA) equal in all hosts? Epidemiological and genetic comparison between animal and
human MRSA. Veterinary Dermatology 23, 267-e54. https://doi.org/10.1111/j.1365-
3164.2012.01072.x

McCubbin, K.D., de Jong, E., Lam, T.J.G.M., Kelton, D.F., Middleton, J.R., McDougall, S.,
De Vliegher, S., Godden, S., Rajala-Schultz, P.J., Rowe, S., Speksnijder, D.C., Kastelic, J.P.,
Barkema, H.W., 2022. Invited review: Selective use of antimicrobials in dairy cattle at drying-off.
Journal of Dairy Science 105, 7161–7189. https://doi.org/10.3168/jds.2021-21455

Monistero, V., Barberio, A., Biscarini, F., Cremonesi, P., Castiglioni, B., Graber, H.U.,
Bottini, E., Ceballos-Marquez, A., Kroemker, V., Petzer, I.M., Pollera, C., Santisteban, C., Veiga
Dos Santos, M., Bronzo, V., Piccinini, R., Re, G., Cocchi, M., Moroni, P., 2020. Different
distribution of antimicrobial resistance genes and virulence profiles of Staphylococcus aureus
strains isolated from clinical mastitis in six countries. Journal of Dairy Science 103, 3431–3446.
https://doi.org/10.3168/jds.2019-17141

Morton, J.T., Freed, S.D., Lee, S.W., Friedberg, I., 2015. A large scale prediction of
bacteriocin gene blocks suggests a wide functional spectrum for bacteriocins. BMC
Bioinformatics 16, 381. https://doi.org/10.1186/s12859-015-0792-9

Nakatsuji, T., Chen, T.H., Narala, S., Chun, K.A., Two, A.M., Yun, T., Shafiq, F., Kotol, P.F.,
Bouslimani, A., Melnik, A.V., Latif, H., Kim, J.-N., Lockhart, A., Artis, K., David, G., Taylor, P.,
Streib, J., Dorrestein, P.C., Grier, A., Gill, S.R., Zengler, K., Hata, T.R., Leung, D.Y.M., Gallo, R.L.,
2017. Antimicrobials from human skin commensal bacteria protect against Staphylococcus

aureus and are deficient in atopic dermatitis. Sci Transl Med 9, eaah4680.
https://doi.org/10.1126/scitranslmed.aah4680

Nascimento, J. dos S., Fagundes, P.C., Brito, M.A.V. de P., Santos, K.R.N. dos, Bastos, M.
do C. de F., 2005. Production of bacteriocins by coagulase-negative staphylococci involved in
bovine mastitis. Veterinary Microbiology 106, 61–71.
https://doi.org/10.1016/j.vetmic.2004.10.014

Naushad, S., Naqvi, S.A., Nobrega, D., Luby, C., Kastelic, J.P., Barkema, H.W., De Buck, J.,
2019. Comprehensive Virulence Gene Profiling of Bovine Non-aureus Staphylococci Based on
Whole-Genome Sequencing Data. mSystems 4, e00098-18.
https://doi.org/10.1128/mSystems.00098-18

Newstead, L.L., Varjonen, K., Nuttall, T., Paterson, G.K., 2020. Staphylococcal-Produced
Bacteriocins and Antimicrobial Peptides: Their Potential as Alternative Treatments for
Staphylococcus aureus Infections. Antibiotics (Basel) 9, 40.
https://doi.org/10.3390/antibiotics9020040

Nguyen, L.-T., Schmidt, H.A., von Haeseler, A., Minh, B.Q., 2015. IQ-TREE: A Fast and
Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol 32,
268–274. https://doi.org/10.1093/molbev/msu300

NMC, 2019. Mastitis control on organic dairy farms in the United States. National
Mastitis Council, Madison, Wisconsin.

NMC, 2017. Laboratory Handbook on Bovine Mastitis. 3rd edition. National Mastitis
Council, Madison, Wisconsin. National Mastitis Council, Madison, Wisconsin.

Nobrega, D.B., Naushad, S., Naqvi, S.A., Condas, L.A.Z., Saini, V., Kastelic, J.P., Luby, C.,
De Buck, J., Barkema, H.W., 2018. Prevalence and Genetic Basis of Antimicrobial Resistance in
Non-aureus Staphylococci Isolated from Canadian Dairy Herds. Front Microbiol 9, 256.
https://doi.org/10.3389/fmicb.2018.00256

Nyman, A.-K., Fasth, C., Waller, K.P., 2018. Intramammary infections with different non-
aureus staphylococci in dairy cows. J Dairy Sci 101, 1403–1418.
https://doi.org/10.3168/jds.2017-13467

Oliver, S.P., Murinda, S.E., 2012. Antimicrobial Resistance of Mastitis Pathogens.
Veterinary Clinics of North America: Food Animal Practice, Mastitis in Dairy Cows 28, 165–185.
https://doi.org/10.1016/j.cvfa.2012.03.005

Parks, D.H., Chuvochina, M., Rinke, C., Mussig, A.J., Chaumeil, P.-A., Hugenholtz, P.,
2022. GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically
consistent, rank normalized and complete genome-based taxonomy. Nucleic Acids Research 50,
D785–D794. https://doi.org/10.1093/nar/gkab776

Patel, K., Godden, S.M., Royster, E.E., Crooker, B.A., Johnson, T.J., Smith, E.A.,
Sreevatsan, S., 2021. Prevalence, antibiotic resistance, virulence and genetic diversity of
Staphylococcus aureus isolated from bulk tank milk samples of U.S. dairy herds. BMC Genomics
22, 367. https://doi.org/10.1186/s12864-021-07603-4

Paterson, G.K., Harrison, E.M., Holmes, M.A., 2014. The emergence of mecC methicillin-
resistant Staphylococcus aureus. Trends Microbiol 22, 42–47.
https://doi.org/10.1016/j.tim.2013.11.003

Peña-Mosca, F., Dean, C., Fernandes, L., Doster, E., Sharpe, K., Ray, T., Feijoo, V., Antunes, A., Baumann, C., Wehri, T., Heins, B., Pinedo, P., Machado, V., Noyes, N., Caixeta, L., 2024. Associations between early lactation intramammary infections and udder health and performance during the first 180 days in milk in first-lactation organic dairy cows. J Dairy Sci 107, 2426–2443. https://doi.org/10.3168/jds.2023-23924

Peña-Mosca, F., Dean, C., Machado, V., Fernandes, L., Pinedo, P., Doster, E., Heins, B., Sharpe, K., Ray, T., Feijoo, V., Antunes, A., Baumann, C., Wehri, T., Noyes, N., Caixeta, L., 2023. Investigation of intramammary infections in primiparous cows during early lactation on organic dairy farms. J Dairy Sci S0022-0302(23)00552–0. https://doi.org/10.3168/jds.2022-23036

Piepers, S., Opsomer, G., Barkema, H.W., de Kruif, A., De Vliegher, S., 2010. Heifers infected with coagulase-negative staphylococci in early lactation have fewer cases of clinical mastitis and higher milk production in their first lactation than noninfected heifers. J Dairy Sci 93, 2014–2024. https://doi.org/10.3168/jds.2009-2897

Piepers, S., Peeters, K., Opsomer, G., Barkema, H.W., Frankena, K., De Vliegher, S., 2011. Pathogen group specific risk factors at herd, heifer and quarter levels for intramammary infections in early lactating dairy heifers. Preventive Veterinary Medicine 99, 91–101. https://doi.org/10.1016/j.prevetmed.2011.02.010

Pol, M., Ruegg, P.L., 2007. Relationship Between Antimicrobial Drug Usage and Antimicrobial Susceptibility of Gram-Positive Mastitis Pathogens. Journal of Dairy Science 90, 262–273. https://doi.org/10.3168/jds.S0022-0302(07)72627-9

Pryor, S.M., Cursons, R.T., Williamson, J.H., Lacy-Hulbert, S.J., 2009. Experimentally induced intramammary infection with multiple strains of Streptococcus uberis. J Dairy Sci 92, 5467–5475. https://doi.org/10.3168/jds.2009-2223

Rajala-Schultz, P., Nødtvedt, A., Halasa, T., Persson Waller, K., 2021. Prudent Use of Antibiotics in Dairy Cows: The Nordic Approach to Udder Health. Frontiers in Veterinary Science 8.

Reyher, K.K., Dohoo, I.R., 2011. Diagnosing intramammary infections: Evaluation of composite milk samples to detect intramammary infections. Journal of Dairy Science 94, 3387–3396. https://doi.org/10.3168/jds.2010-3907

Reyher, K.K., Haine, D., Dohoo, I.R., Revie, C.W., 2012. Examining the effect of intramammary infections with minor mastitis pathogens on the acquisition of new intramammary infections with major mastitis pathogens—A systematic review and meta-analysis. Journal of Dairy Science 95, 6483–6502. https://doi.org/10.3168/jds.2012-5594

Rooijakkers, S.H.M., Ruyken, M., Roos, A., Daha, M.R., Presanis, J.S., Sim, R.B., van Wamel, W.J.B., van Kessel, K.P.M., van Strijp, J.A.G., 2005. Immune evasion by a staphylococcal complement inhibitor that acts on C3 convertases. Nat Immunol 6, 920–927. https://doi.org/10.1038/ni1235

Rost, B., 1999. Twilight zone of protein sequence alignments. Protein Engineering, Design and Selection 12, 85–94. https://doi.org/10.1093/protein/12.2.85

Rowe, S.M., Godden, S.M., Royster, E., Timmerman, J., Crooker, B.A., Boyle, M., 2019. Cross-sectional study of the relationships among bedding materials, bedding bacteria counts, and intramammary infection in late-lactation dairy cows. Journal of Dairy Science 102, 11384–11400. https://doi.org/10.3168/jds.2019-17074

Ruegg, P.L., 2022. Realities, Challenges and Benefits of Antimicrobial Stewardship in Dairy Practice in the United States. Microorganisms 10, 1626. https://doi.org/10.3390/microorganisms10081626

Ruegg, P.L., 2017. A 100-Year Review: Mastitis detection, management, and prevention. J Dairy Sci 100, 10381–10397. https://doi.org/10.3168/jds.2017-13023

Sánchez-Romero, M.A., Casadesús, J., 2020. The bacterial epigenome. Nat Rev Microbiol 18, 7–20. https://doi.org/10.1038/s41579-019-0286-2

Schreiner, D.A., Ruegg, P.L., 2003. Relationship between udder and leg hygiene scores and subclinical mastitis. J Dairy Sci 86, 3460–3465. https://doi.org/10.3168/jds.S0022-0302(03)73950-2

Seemann, T., 2023a. Abricate: Mass screening of contigs for antimicrobial and virulence gen. Accesed: 2/6/2024.

Seemann, T., 2023b. Snippy: Rapid haploid variant calling and core genome alignment. Accessed: 2/6/2024.

Smith, E.J., Visai, L., Kerrigan, S.W., Speziale, P., Foster, T.J., 2011. The Sbi Protein Is a Multifunctional Immune Evasion Factor of Staphylococcus aureus. Infection and Immunity 79, 3801–3809. https://doi.org/10.1128/iai.05075-11

Smith, E.M., Green, L.E., Medley, G.F., Bird, H.E., Dowson, C.G., 2005. Multilocus Sequence Typing of Staphylococcus aureus Isolated from High-Somatic-Cell-Count Cows and the Environment of an Organic Dairy Farm in the United Kingdom. Journal of Clinical Microbiology 43, 4731–4736. https://doi.org/10.1128/jcm.43.9.4731-4736.2005

Sordillo, L.M., 2018. Mammary Gland Immunobiology and Resistance to Mastitis. Veterinary Clinics of North America: Food Animal Practice, Mastitis 34, 507–523. https://doi.org/10.1016/j.cvfa.2018.07.005

Spencer, B.L., Doran, K.S., 2022. Evolving understanding of the type VII secretion system in Gram-positive bacteria. PLOS Pathogens 18, e1010680. https://doi.org/10.1371/journal.ppat.1010680

Srednik, M.E., Archambault, M., Jacques, M., Gentilini, E.R., 2017. Detection of a mecC-positive Staphylococcus saprophyticus from bovine mastitis in Argentina. Journal of Global Antimicrobial Resistance 10, 261–263. https://doi.org/10.1016/j.jgar.2017.05.016

Supré, K., Haesebrouck, F., Zadoks, R.N., Vaneechoutte, M., Piepers, S., De Vliegher, S., 2011. Some coagulase-negative Staphylococcus species affect udder health more than others. J Dairy Sci 94, 2329–2340. https://doi.org/10.3168/jds.2010-3741

Tam, K., Torres, V.J., 2019. Staphylococcus aureus Secreted Toxins and Extracellular Enzymes. Microbiology Spectrum 7, 10.1128/microbiolspec.gpp3-0039–2018. https://doi.org/10.1128/microbiolspec.gpp3-0039-2018

Tan, L., Li, S.R., Jiang, B., Hu, X.M., Li, S., 2018. Therapeutic Targeting of the Staphylococcus aureus Accessory Gene Regulator (agr) System. Front Microbiol 9, 55. https://doi.org/10.3389/fmicb.2018.00055

Toft, N., Halasa, T., Nielsen, S.S., Hechinger, S., Zervens, L.M., Schwarz, D., Kirkeby, C., 2019. Composite or aseptic quarter milk samples: Sensitivity and specificity of PCR and bacterial

culture of *Staphylococcus aureus* based on Bayesian latent class evaluation. Preventive Veterinary Medicine 171, 104689. https://doi.org/10.1016/j.prevetmed.2019.05.002

Toledo-Silva, B., Beuckelaere, L., De Visscher, A., Geeroms, C., Meyer, E., Piepers, S., Thiry, D., Haesebrouck, F., De Vliegher, S., 2022. Novel Quantitative Assay to Describe In Vitro Bovine Mastitis Bacterial Pathogen Inhibition by Non-aureus Staphylococci. Pathogens 11, 264. https://doi.org/10.3390/pathogens11020264

Toledo-Silva, B., de Souza, F.N., Mertens, K., Piepers, S., Haesebrouck, F., De Vliegher, S., 2021. Bovine-associated non-aureus staphylococci suppress Staphylococcus aureus biofilm dispersal in vitro yet not through agr regulation. Vet Res 52, 114. https://doi.org/10.1186/s13567-021-00985-z

Tomazi, T., Gonçalves, J.L., Barreiro, J.R., Arcari, M.A., dos Santos, M.V., 2015. Bovine subclinical intramammary infection caused by coagulase-negative staphylococci increases somatic cell count but has no effect on milk yield or composition. J Dairy Sci 98, 3071–3078. https://doi.org/10.3168/jds.2014-8466

Tremblay, Y.D.N., Lamarche, D., Chever, P., Haine, D., Messier, S., Jacques, M., 2013. Characterization of the ability of coagulase-negative staphylococci isolated from the milk of Canadian farms to form biofilms. J Dairy Sci 96, 234–246. https://doi.org/10.3168/jds.2012-5795

USDA, 2007. Dairy 2007, Part I: Reference of Dairy Cattle Health and Management Practices in the United States, 2007. #N480.1007. USDA-APHIS-VS, CEAH, Fort Collins, CO.

Valckenier, D., Piepers, S., De Visscher, A., De Vliegher, S., 2020. The effect of intramammary infection in early lactation with non-aureus staphylococci in general and Staphylococcus chromogenes specifically on quarter milk somatic cell count and quarter milk yield. J Dairy Sci 103, 768–782. https://doi.org/10.3168/jds.2019-16818

van Heel, A.J., de Jong, A., Song, C., Viel, J.H., Kok, J., Kuipers, O.P., 2018. BAGEL4: a user-friendly web server to thoroughly mine RiPPs and bacteriocins. Nucleic Acids Research 46, W278–W281. https://doi.org/10.1093/nar/gky383

Wang, B., Muir, T.W., 2016. Regulation of virulence in Staphylococcus aureus: molecular mechanisms and remaining puzzles. Cell Chem Biol 23, 214–224. https://doi.org/10.1016/j.chembiol.2016.01.004

Wiegand, I., Hilpert, K., Hancock, R.E.W., 2008. Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. Nat Protoc 3, 163–175. https://doi.org/10.1038/nprot.2007.521

Woudstra, S., Wente, N., Zhang, Y., Leimbach, S., Gussmann, M.K., Kirkeby, C., Krömker, V., 2023. Strain diversity and infection durations of Staphylococcus spp. and Streptococcus spp. causing intramammary infections in dairy cows. Journal of Dairy Science 106, 4214–4231. https://doi.org/10.3168/jds.2022-22942

Zadoks, R.N., Gillespie, B.E., Barkema, H.W., Sampimon, O.C., Oliver, S.P., Schukken, Y.H., 2003. Clinical, epidemiological and molecular characteristics of Streptococcus uberis infections in dairy herds. Epidemiology & Infection 130, 335–349. https://doi.org/10.1017/S0950268802008221

Zadoks, R.N., Middleton, J.R., McDougall, S., Katholm, J., Schukken, Y.H., 2011. Molecular epidemiology of mastitis pathogens of dairy cattle and comparative relevance to

1238   humans. J Mammary Gland Biol Neoplasia 16, 357–372. https://doi.org/10.1007/s10911-011-
1239   9236-y

1240

1242 **Table 1.** Non-aureus *staphylococci* and *mammaliicocci* counts ($\log_{10}$ CFU/square inch of gauze)
1243 and prevalence in cows without and with a *Staphylococcus aureus* or *Streptococcus* spp. and
1244 *Streptococcus*-like organisms (n=71 teat apex samples).

Outcome	SAU SSLO IMI		Estimate (95%CI)	*P-value*
	No	Yes		
Counts				
Overall	1.10 ± 0.17	1.02 ± 0.17	-0.09 (-0.54, 0.35)	0.69
Staph chromogenes	0.26 ± 0.11	0.20 ± 0.12	-0.06 (-0.34, 0.23)	0.71
Staph haemolyticus	0.20 ± 0.13	0.21 ± 0.13	0.01 (-0.31, 0.32)	0.96
Mammaliicoccus sciuri	0.19 ± 0.10	0.20 ± 0.10	0.01 (-0.23, 0.26)	0.91
Staph succinus [a]	0.17 ± 0.09	0.13 ± 0.08	-0.04 (-0.27, 0.19)	0.75
Staph xylosus/*pseudoxylosus*	0.46 ± 0.09	0.53 ± 0.10	0.07 (-0.17, 0.31)	0.56
Other	0.20 ± 0.08	0.15 ± 0.08	-0.05 (-0.24, 0.14)	0.60
Prevalence				
Overall	66.7% (24/36)	62.9% (22/35)	0.85 (0.32, 2.25)	0.74
Staph chromogenes	25.0% (9/36)	22.9% (8/35)	0.89 (0.29, 2.73)	0.84
Staph haemolyticus	25.0% (9/36)	22.9% (8/35)	0.89 (0.29, 2.73)	0.84
Mammaliicoccus sciuri	8.3% (3/36)	5.7% (2/35)	0.66 (0.08, 4.28)	0.66
Staph succinus [a]	11.1% (4/36)	11.4% (4/35)	1.03 (0.23, 4.72)	0.97
Staph xylosus/*pseudoxylosus*	11.1% (4/36)	17.1% (6/35)	1.66 (0.43, 7.02)	0.46
Other [b]	8.3% (3/36)	2.9% (1/35)	0.31 (0.01, 2.63)	0.29

1245 [a] *Staph succinus*: Model did not account for farm, because it was not present in farm B. [b] Other:
1246 *Staph cohnii, Staph devriesei* and *Staph equorum.* NASM: non-*aureus staphylococci* and
1247 *mammaliicocci*.

1248

1249 **Table 2.** Taxonomic identification of isolates isolated from teat apex samples and submitted for
1250 taxonomic identification using Matrix-Assisted Laser Desorption/Ionization-Time of Flight mass
1251 spectrometer (n=273 isolates) and confirmation of species-level taxonomy for non-*aureus*
1252 *staphylococci* and *mammaliicocci* (**NASM**) using whole genome sequencing.

Taxonomy classification	SAU SSLO IMI	
	No	Yes
Staph aureus	0% (0/132)	1.4% (2/141)
NASM	32.8% (44/132)	27.7% (39/141)
Staph chromogenes	9.1% (12/132)	7.1% (10/141)
Staph cohnii	0% (0/132)	0.7% (1/141)
Staph devriesei	1.5% (2/132)	0% (0/141)
Staph equorum	0.8% (1/132)	0% (0/141)
Staph haemolyticus	7.6% (10/132)	6.4% (9/141)
Staph pseudoxylosus	1.5% (2/132)	2.1% (3/141)
Mammaliicoccus sciuri	2.3% (3/132)	2.1% (3/141)
Staph succinus	4.5% (6/132)	4.3% (6/141)
Staph xylosus	4.5% (6/132)	5.0% (7/141)
Aerococcus viridans	0% (0/132)	1.4% (2/141)
Aerococcus sp.	0.8% (1/132)	2.8% (4/141)
Bacillus sp.	62.9% (83/132)	61.7% (87/141)
Gram-positive rod	4.5% (6/132)	5.0% (7/141)

1253

1254

Table 3. Relationship between species identification and minimum inhibitory concentration (MIC; log CFU/mL) of non-*aureus*
staphylococci and *mammaliicocci* (NASM) isolated from teat apex against *Staphylococcus aureus* and *Streptococcus uberis* (n=77
isolates).

NASM taxonomy	MIC SAU *		Top 10 lowest MIC SAU [¥]		MIC SUB *		Top 10 lowest MIC SUB [¥]	
	Mean (SE)	*P-value*	Mean (SE)	*P-value*	Mean (SE)	*P-value*	*% (n/N)*	*P-value*
Counts								
SCH	3.32 (0.13) [a]	0.001	10.5% (2/19)	0.05	3.06 (0.16)	0.41	10.5% (2/19)	0.07
SHAEM	3.97 (0.14) [b]		5.2% (1/19)		3.51 (0.28)		4.8% (1/19)	
MSC	3.32 (0.24) [a]		16.9% (1/6)		3.49 (0.30)		0% (0/6)	
SSUC	2.92 (0.17) [a]		45.5% (5/11)		2.92 (0.22)		45.5% (5/11)	
SXYL	3.05 (0.14) [a]		5.6% (1/18)		3.05 (0.17)		11.1% (2/18)	
Other NASM	4.39 (0.32) [b]		0.0% (0/4)		3.29 (0.49)		0.0% (0/4)	

Other NASM: *Staph cohnii, Staph devriesei, Staph equorum.* [¥] Top 10 lowest MIC: top 10 isolates with lowest MIC against
Staphylococcus aureus (SAU) or *Streptococcus uberis* (SUB). * For MIC comparison across species, only isolates with a determined
MIC were included, as some did not exhibit inhibitory activity at the maximum concentration evaluat

1261

Figure 1. Ranking of 77 non-*aureus staphylococci* and *mammaliicocci* (NASM) isolates by

minimum inhibitory concentration (MIC) against *Staphylococcus aureus* and *Streptococcus*

uberis isolated from teat apex stratified by presence of intramammary infections by

Staphyloccocus aureus and *Streptococcus uberis* (SAU SSLO IMI) (panel A/B) and NASM

taxonomy, as determined using Matrix-Assisted Laser Desorption/Ionization-Time of Flight and

whole genome sequencing (panel C/D) (n=79 isolates). No inhibitory activity: NASM isolates

1268   that showed growth of *Staphylococcus aureus* or *Streptococcus uberis* at the maximum

1269   investigated concentration on media. SCH: *Staphylococcus chromogenes*, SHAEM:

1270   *Staphylococcus haemolyticus*, MSC: *Mammaliicoccus sciuri*, SSUC: *Staphylococcus succinus,*

1271   SXYL: *Staphylococcus xylosus/pseudoxylosus,* Other: *Staphylococcus cohnii, Staphylococcus*

1272   *devriesei* and *Staphylococcus equorum.*

1273

Figure 2. Kaplan-Meier plots showing the presence of *Staphylococcus aureus* and *Streptococcus*

uberis growth on top of the media at different concentrations of non-*aureus staphylococci* and

mammaliicocci (NASM) in the top layer, stratified by the presence of intramammary infections

caused by *Staphylococcus aureus* and *Streptococcus uberis* and NASM taxonomy, as determined

using Matrix-Assisted Laser Desorption/Ionization-Time of Flight and whole genome

sequencing (n=77 isolates). SCH: *Staphylococcus chromogenes*, SHAEM: *Staphylococcus*

1280   *haemolyticus*, MSC: *Mammaliicoccus sciuri*, SSUC: *Staphylococcus succinus,* SXYL:

1281   *Staphylococcus xylosus/pseudoxylosus,* Other: *Staphylococcus cohnii, Staphylococcus devriesei*

1282   and *Staphylococcus equorum.*

1283

Figure 3. Maximum likelihood phylogenetic tree including 21 isolates identified as

Staphylococcus chromogenes (SCH). This tree represents a total of 64,794 single nucleotide

polymorphisms. Reference: GenBank accession number: GCA_002994305.1. MIC SAU:

Minimum inhibitory concentration against *Staphylococcus aureus*. MIC SUB: Minimum

inhibitory concentration against *Streptococcus uberis*. Ranking inhibitory activity on SAU:

Ranking of isolates according to MIC on SAU. Ranking inhibitory activity on SUB: Ranking of

isolates according to MIC on SUB. Auto induced peptide class I (aip_1): NCBI accession

number: WP_001093929.1. Auto induced peptide class II (aip_2). NCBI accession number

WP_001094921.1. Auto induced peptide class III (aip_3): CBI accession number

WP_000735197.1. Auto induced peptide class IV (aip_4): NCBI accession number:

WP_001094303.1). Sactipeptides: Interpro accession number PF04055. Subtilosin A NCBI

accession number: NP_391616.1. Lanthipeptides class II Interpro accession number: PF05147.

Lanthipeptides class IV (Uniprot accession number: O88037). putative bacteriocins 193.2 and

194.2 (putative_bacteriocin_193/194): Genbank accession number AJ002203.2. lacticin Z

(lacticin): Interpro accession number PF11758. Carnolysins: Interpo accession number PF00082.

1299

Figure 4. Maximum likelihood phylogenetic tree including 19 isolates identified as

Staphylococcus haemolyticus (SHAEM). This tree represents a total of 52,188 single nucleotide

polymorphisms. Reference: GenBank accession number: GCA_001611955.1. MIC SAU:

Minimum inhibitory concentration against *Staphylococcus aureus*. MIC SUB: Minimum

inhibitory concentration against *Streptococcus uberis*. Ranking inhibitory activity on SAU:

Ranking of isolates according to MIC on SAU. Ranking inhibitory activity on SUB: Ranking of

isolates according to MIC on SUB. Auto induced peptide class I (aip_1): NCBI accession

number: WP_001093929.1. Auto induced peptide class II (aip_2). NCBI accession number

WP_001094921.1. Auto induced peptide class III (aip_3): CBI accession number

WP_000735197.1. Auto induced peptide class IV (aip_4): NCBI accession number:

WP_001094303.1). Sactipeptides: Interpro accession number PF04055. Subtilosin A NCBI

accession number: NP_391616.1. Lanthipeptides class II Interpro accession number: PF05147.

Lanthipeptides class IV (Uniprot accession number: O88037). putative bacteriocins 193.2 and

194.2 (putative_bacteriocin_193/194): Genbank accession number AJ002203.2. lacticin Z

(lacticin): Interpro accession number PF11758. Carnolysins: Interpo accession number PF00082.

1315

Figure 5. Maximum likelihood phylogenetic tree including 6 isolates identified as

Mammaliicoccus sciuri (MSC). This tree represents a total of 93,312 single nucleotide

polymorphisms. Reference: GenBank accession number: GCA_002209165.2. MIC SAU:

Minimum inhibitory concentration against *Staphylococcus aureus*. MIC SUB: Minimum

inhibitory concentration against *Streptococcus uberis*. Ranking inhibitory activity on SAU:

Ranking of isolates according to MIC on SAU. Ranking inhibitory activity on SUB: Ranking of

isolates according to MIC on SUB. Auto induced peptide class I (aip_1): NCBI accession

number: WP_001093929.1. Auto induced peptide class II (aip_2). NCBI accession number

WP_001094921.1. Auto induced peptide class III (aip_3): CBI accession number

WP_000735197.1. Auto induced peptide class IV (aip_4): NCBI accession number:

WP_001094303.1). Sactipeptides: Interpro accession number PF04055. Subtilosin A NCBI

accession number: NP_391616.1. Lanthipeptides class II Interpro accession number: PF05147.

Lanthipeptides class IV (Uniprot accession number: O88037). putative bacteriocins 193.2 and

194.2 (putative_bacteriocin_193/194): Genbank accession number AJ002203.2. lacticin Z

(lacticin): Interpro accession number PF11758. Carnolysins: Interpo accession number PF00082.

1331

Figure 6. Maximum likelihood phylogenetic tree including 12 isolates identified as

Staphylococcus succinus (SSUC). This tree represents a total of 69,716 single nucleotide

polymorphisms. Reference: GenBank accession number: GCA_001902315.1. MIC SAU:

Minimum inhibitory concentration against *Staphylococcus aureus*. MIC SUB: Minimum

inhibitory concentration against *Streptococcus uberis*. Ranking inhibitory activity on SAU:

Ranking of isolates according to MIC on SAU. Ranking inhibitory activity on SUB: Ranking of

isolates according to MIC on SUB. Auto induced peptide class I (aip_1): NCBI accession

number: WP_001093929.1. Auto induced peptide class II (aip_2). NCBI accession number

WP_001094921.1. Auto induced peptide class III (aip_3): CBI accession number

WP_000735197.1. Auto induced peptide class IV (aip_4): NCBI accession number:

WP_001094303.1). Sactipeptides: Interpro accession number PF04055. Subtilosin A NCBI

accession number: NP_391616.1. Lanthipeptides class II Interpro accession number: PF05147.

Lanthipeptides class IV (Uniprot accession number: O88037). putative bacteriocins 193.2 and

194.2 (putative_bacteriocin_193/194): Genbank accession number AJ002203.2. lacticin Z

(lacticin): Interpro accession number PF11758. Carnolysins: Interpo accession number PF00082.

1347

Figure 7. Maximum likelihood phylogenetic tree including 18 isolates identified as

Staphylococcus xylosus (XYL) and *Staphylococcus pseudoxylosus* (PXYL). This tree represents

a total of 209,336 single nucleotide polymorphisms. Reference: GenBank accession number:

GCA_000709415.1. MIC SAU: Minimum inhibitory concentration against *Staphylococcus*

aureus. MIC SUB: Minimum inhibitory concentration against *Streptococcus uberis*. Ranking

inhibitory activity on SAU: Ranking of isolates according to MIC on SAU. Ranking inhibitory

activity on SUB: Ranking of isolates according to MIC on SUB. Auto induced peptide class I

(aip_1): NCBI accession number: WP_001093929.1. Auto induced peptide class II (aip_2).

NCBI accession number WP_001094921.1. Auto induced peptide class III (aip_3): CBI

accession number WP_000735197.1. Auto induced peptide class IV (aip_4): NCBI accession

number: WP_001094303.1). Sactipeptides: Interpro accession number PF04055. Subtilosin A

NCBI accession number: NP_391616.1. Lanthipeptides class II Interpro accession number:

PF05147. Lanthipeptides class IV (Uniprot accession number: O88037). putative bacteriocins

193.2 and 194.2 (putative_bacteriocin_193/194): Genbank accession number AJ002203.2.

1362   lacticin Z (lacticin): Interpro accession number PF11758. Carnolysins: Interpo accession number

1363   PF00082.

1364

Figure 8. Distribution of virulence genes involved in adherence and exoenzymes in

Staphylococcus aureus and non-*aureus staphylococci* and *mammaliicocci* (NASM).

Staphylococcus aureus strain 1 (SAU 1) represents a reference strain: American Type Culture

Collection strain Seattle 1945; *Staphylococcus aureus* subsp. *aureus* Rosenbach 25923 ™.

Rosenbach 25923™. aap: accumulation associated protein. bap: biofilm associated surface

protein. atl: autolysin. atlC: fibronectin binding autolysin. clfA-B: clumping factor class A-B.

cna: collagen adhesin precursor. ebh: cell wall associated fibronectin-binding protein. ebhA: cell

wall associated fibronectin-binding protein class A. efb: fibrinogen-binding protein. fnbA-B:

fibronectin-binding protein class A-B. eap/map: extracellular adherence protein/MHC analogous

protein. sasC-G: cell wall surface anchor family protein type C-G. sraP: serine rich adhesion for

platelets. icaA-R: intercellular adhesion proteins class A-R. sdrC-I: Ser-Asp rich fibrinogen-

binding bone sialoprotein-binding protein class C-I. uafA: cell wall anchored protein class A.

adsA: adenosine synthase A. aur: aureolysin. coa: staphylocoagulase. geh: glycerol ester

hydrolase. hysA: hyaluronate lyase. lip: triacylglycerol lipase. sak: staphylokinase. nuc:

thermonuclease. spLA-F: serine proteinase class A-F. sspA-C: serine protease class A-C. vWbp:

von Willebrand factor-binding protein.

Figure 9. Distribution of virulence genes involved in immune evasion and iron uptake and

metabolism in *Staphylococcus aureus* and non-*aureus staphylococci* and *mammaliicoci*

(NASM). *Staphylococcus aureus* strain 1 (SAU 1) represents a reference strain: American Type

Culture Collection strain Seattle 1945; *Staphylococcus aureus* subsp. *aureus* Rosenbach

25923 ™. cap8A-O: capsular polysaccharide-synthesis enzyme class 8A-O. cap5A-P: capsular

polysaccharide-biosynthesis protein class 5A-P. chp: chemotaxis-inhibiting protein. sbi:

1388   staphylococcal binding immunoglobulin protein. scn: staphylococcal complement inhibitor. spa:

1389   staphylococcal protein A. isdA-I: iron-regulated surface determinant protein class A-I. htsA-C:

1390   FecCD iron compound ABC transporter permease family protein class A-C. sbnA-I: siderophore

1391   biosynthesis class A-I. srtB: sortase B. sirA-C: staphylococcal iron regulated class A-C. sfaA-D:

1392   staphyloferrin A biosynthesis protein class A-D.

1393

Figure 10. Distribution of toxin related genes in *Staphylococcus aureus* and non-*aureus*

staphylococci and *mammaliicocci* (NASM). *Staphylococcus aureus* strain 1 (SAU 1) represents a

reference strain: American Type Culture Collection strain Seattle 1945; *Staphylococcus aureus*

subsp. *aureus* Rosenbach 25923 ™. se/sel a-v: Staphylococcal enterotoxin A-V. selu2:

Staphylococcal enterotoxin U2. sec3: Staphylococcal enterotoxin C1 precursor. set1-40:

staphylococcal exotoxin 1-40. yent1-2: enterotoxin yent1-2. hly/hla: alpha-hemolysin. hlb: beta-

hemolysin. hld: delta-hemolysin. hlgA-C: gamma-hemolysin component A-C. lukD-M:

leukocidin D-M. lukF-like: Panton-Valentine leukocidin LukF-PV chain precursor. lukF-PV:

Panton-Valentine leukocidin chain F precursor. lukS-PV: Panton-Valentine leukocidin chain S

1403   precursor. tsst: toxic shock syndrome toxin-1. eta-d: exfoliative toxin class A-D. esaA-C: type

1404   VII secretion system protein class A-C. essA-C: type VII secretion system protein class A-C.

1405   esxA-B: type VII secretion system secreted protein class A-B. PSMβ1-6: phenol-soluble

1406   modulin beta class 1-6.

1407

Figure 11. Distribution of antimicrobial resistance genes in *Staphylococcus aureus* and non-

aureus staphylococci and *mammaliicoci* (NASM). *Staphylococcus aureus* strain 1 (SAU 1)

represents a reference strain: American Type Culture Collection strain Seattle 1945;

Staphylococcus aureus subsp. *aureus* Rosenbach 25923 ™. AAC3: Aminoglycoside N-

acetyltransferase. ANT9: Aminoglycoside nucleotidyltransferase. APH3-PRIME:

Aminoglycoside O-phosphotransferases. ArlR: Arl Response Regulator. ArlS: Arl sensor kinase.

BLAZ: Class A betalactamases. DHAP: Phenicol resistance MFS efflux pumps. ERM: 23S

rRNA methyltransferases. FOSB: Fosfomycin thiol transferases class B. FOSD: Fosfomycin

thiol transferases class D. FUSF: Fusidic acid esterases. LMRS: Drug and biocide MFS efflux

pumps. LNUA: Lincosamide nucleotidyltransferases. MECA: Penicillin binding protein class A.

MECC: Penicillin binding protein class C. MECI: Penicillin binding protein class I. MEPA:

Drug and biocide MATE efflux pumps class A. MEPB: Drug and biocide MATE efflux pumps

class B. MEPR: Drug and biocide MATE efflux pumps class R. MGRA: MDR regulator.

MPHC: Macrolide phosphotransferases. MSRA: MLS resistance ABC efflux pumps. NORA:

Drug and biocide MFS efflux pumps class A. NORB: Drug and biocide MFS efflux pumps class

B. RLMH: 23S rRNA methyltransferases. SALA: Multi-drug ABC efflux pumps. TET38:

Tetracycline resistance MFS efflux pumps. TETK: Tetracycline resistance MFS efflux pump.

Tree scale: 0.01

Tree scale: 0.1

Tree scale: 0.1

	IMI	Cow ID	DIM	Farm	MIC SAU (log10; CFU/mL)	Ranking inhibitory activity on SAU	MIC SUB (log10; CFU/mL)	Ranking inhibitory activity on SUB
SXYL2	No	1276	3	B	2.78	11 to 20	3.78	51 to 60
SXYL10	No	1283	3	B	2.90	21 to 30	2.90	21 to 30
Reference								
SXYL4	Yes	171	-5	A	3.17	31 to 40	3.17	31 to 40
SXYL11	Yes	1294	-7	B	2.71	11 to 20	1.71	Top 10
SXYL13	Yes	1294	-7	B	3.17	31 to 40	2.17	Top 10
SXYL1	No	1276	3	B	4.09	51 to 60	4.09	61 to 70
SPXYL3	Yes	41010	-7	A	2.70	11 to 20	2.70	11 to 20
SPXYL5	Yes	41010	-7	A	3.06	21 to 30	3.06	21 to 30
SPXYL1	No	1288	12	B	3.93	51 to 60	3.93	51 to 60
SPXYL2	Yes	1294	-7	B	3.01	21 to 30	3.01	21 to 30
SPXYL4	No	1283	3	B	2.61	Top 10	2.61	11 to 20
SXYL6	Yes	1298	4	B	3.16	31 to 40	3.16	31 to 40
SXYL12	No	1276	3	B	2.80	11 to 20	3.80	51 to 60
SXYL7	Yes	1289	6	B	2.74	11 to 20	2.74	21 to 30
SXYL3	No	1286	-7	B	2.85	11 to 20	3.85	51 to 60
SXYL9	Yes	1285	20	B	2.93	21 to 30	2.93	21 to 30
SXYL5	Yes	1294	-7	B	2.81	11 to 20	2.81	21 to 30
SXYL8	No	1286	-7	B	3.56	41 to 50	2.56	11 to 20

Tree scale: 0.1