Dextran sodium sulfate-induced colitis alters the proportion and composition of replicating gut bacteria

Eve T. Beauchemin[1], Claire Hunter[2], & Corinne F. Maurice[1,3]

[1]Department of Microbiology & Immunology, Faculty of Medicine and Health Sciences, McGill University, Montreal, Quebec, Canada

[2]Department of Public Health and Primary Care, School of Clinical Medicine, University of Cambridge, Cambridge, England

[3]McGill Centre for Microbiome Research, Montreal, Quebec, Canada

Corresponding author: corinne.maurice@mcgill.ca

Abstract

The bacteria living in the human gut are essential for host health. Though the composition and metabolism of these bacteria is well described in both healthy hosts and those with intestinal disease, less is known about the activity of the gut bacteria prior to, and during, disease development – especially regarding gut bacterial replication. Here, we use a recently developed single-cell technique alongside existing metagenomics-based tools to identify, track, and quantify the replicating gut bacteria and their replication dynamics in the dextran sodium sulfate mouse model of colitis. We show that the proportion of replicating gut bacteria decreases when mice have the highest levels of inflammation and returns to baseline levels as mice begin recovering. We additionally report significant alterations in the composition of the total replicating gut bacterial community during colitis development. On the taxa level, we observe significant changes in the abundance of taxa such as the mucus-degrading *Akkermansia muciniphila* and the poorly described *Erysipelatoclostridium* genus. We further demonstrate that many taxa exhibit variable replication rates during colitis, including *A. muciniphila*. Lastly, we show that colitis

development is positively correlated with increases in the presence and abundance of
bacteria predicted to be fast replicators, suggesting that taxa with the potential to replicate
quickly may have an advantage during intestinal inflammation. These data support the
need for additional research using activity-based approaches to further characterize the
gut bacterial response to intestinal inflammation and its consequences for both the host
and the gut microbial community at large.

Importance

It is well known that the bacteria living inside the gut are important for human health.
Indeed, the type of bacteria which are present and their metabolism is different in healthy
people versus those with intestinal disease. However, less is known about how these gut
bacteria are replicating, especially as someone begins to develop intestinal disease. This
is especially important as it is thought that the active gut bacteria may be more relevant
to health. Here, we begin addressing this gap by using several complementary
approaches to characterize the replicating gut bacteria in a mouse model of intestinal
inflammation. We reveal which gut bacteria are replicating, and how quickly, as mice
develop and recover from inflammation. This work can serve as a model for future
research to identify how the active gut bacteria may be impacting health, or why these
particular bacteria tend to thrive during intestinal inflammation.

Introduction

Inflammatory bowel disease (IBD) is an umbrella term for several chronic inflammatory
intestinal diseases, including Crohn's disease (CD) and ulcerative colitis (UC).[1] The
incidence of IBD has been increasing in both high-income and low- and middle-income
countries over the past 10 years and shows no signs of slowing down.[2] Though the exact
etiology of IBD is unknown, evidence suggests that IBD results from a combination of
genetic defects in the immune system and environmental triggers, which results in an
aberrant host immune response to the gut microbiota.[3,4] Indeed, this disease cannot
occur without the presence of the gut bacteria.[5] As such, these bacteria may play a crucial
role in the manifestation and/or perpetuation of IBD.

Several studies have characterized the composition of the colonic or fecal bacteria in patients with IBD versus those without. Patients with IBD generally exhibit decreased bacterial diversity, increased abundances of aerotolerant and facultative anaerobic bacteria (such as members of the *Enterobacteriaceae* family), and decreased abundances of beneficial obligate anaerobes.[6] It remains unclear what roles the gut bacteria play in IBD, and whether the observed diversity changes are a cause or consequence of disease. One emerging hypothesis is that the immune dysregulation apparent in IBD patients could initially alter the gut environment, favouring the survival and growth of bacterial taxa with pro-inflammatory properties, which further contribute to inflammation.[1,7,8] This self-perpetuating cycle is broken primarily through the use of medication, such as corticosteroid drugs and 5-aminosalicylic acid, also known as mesalamine.[7,9]

Studies which have evaluated changes in gut bacterial transcripts, proteins, and metabolites have shown more consistent alterations than changes in gut bacterial composition in IBD patients, supporting the notion that gut bacterial metabolism plays a larger role in perpetuating inflammation than changes in community structure.[3,10] For example, in IBD there is an increased abundance of bacterial genes encoding for, and proteins involved in, oxidative stress and DNA replication;[11,12] increased levels of bacterial proteins related to sulfur metabolism and hydrogen sulfide production;[12] increased transcriptional activity of specific bacteria, including *Clostridia* species and the mucin-degrader *Ruminococcus gnavus*;[6] and significant alterations in bacterial metabolites, such as decreased levels of anti-inflammatory short-chain fatty acids like butyrate, and increased levels of primary bile acids.[6]

One fundamental aspect of bacterial activity which remains understudied in the gut is bacterial replication,[13] specifically in the context of inflammation. Pioneering work from Olm et al. (2019) showed that bacterial replication rates were one of the best predictors of the development of necrotizing enterocolitis (NEC), an inflammatory intestinal disease in pre-term infants. Indeed, the authors reported a significant increase in gut bacterial

replication rates, especially members of the *Enterobacteriaceae* family, preceding the
development of NEC.[14] In the same year, Riglar et al. (2019) tested their newly developed
a synthetic gene oscillator, the repressilator 2.0, which reported on the growth dynamics
of a transformed *E. coli* strain in a mouse model of dextran sodium sulfate (DSS)-induced
colitis. The authors found that during active inflammation, there was increased variability
in the growth rates of the transformed *E. coli*.[15]

In a more recent publication, Joseph et al. (2022) used an updated version of their
replication rate measurement tool, termed Co-PTR, to identify taxa with altered replication
rates in healthy individuals versus those with IBD.[16] They found that gut bacterial
replication rates exhibited large interindividual variability but seemed relatively stable
within healthy individuals. They also reported that replication rates of specific taxa were
associated with CD (*Subdoligranulum* species) or UC (*Roseburia intestinalis*,
Ruminiclostridium species, and *Subdoligranum* species), whereas the relative
abundances of these same taxa were not associated with their respective disease.
Notably, they highlight that at least one other study also reported *R. intestinalis* as having
increased replication rates in individuals with CD and UC, emphasizing the relevance of
this taxon in IBD.

Collectively, these studies demonstrate how gut bacterial replication provides information
about gut bacterial activity that complements relative abundance measurements,
contributing to a better understanding of which taxa will thrive during intestinal
inflammation. This could allow us to characterize how gut bacteria contribute to
inflammatory conditions and shape the composition and function of the post-inflammatory
gut microbiota.

Commonly used tools to study bacterial replication in community contexts rely on whole
genome shotgun sequencing (WGS) data. Though these tools are informative and
quantitative, they can be prohibitively expensive to implement and are data-analysis
intensive.[17] As such, there is a need for more physiologically-based techniques,[18] which

are more affordable and easily implementable, to complement existing WGS-focused
techniques.

Most studies to date which evaluated gut bacterial activity or replication in IBD and other
inflammatory conditions were cross-sectional and did not focus on disease development
and progression. This is partly because it is challenging to identify and track individuals
who will later develop IBD, despite new efforts in this direction.[19-21] We hypothesize that
tracking gut bacterial replication dynamics during the development and recovery from
colitis will better inform us about the bacteria most relevant to intestinal inflammation.

Here, we use a combination of bacterial physiology- and bioinformatics-based techniques
to capture changes in the composition and replication dynamics of gut bacteria during the
development and recovery from chemically induced colitis in mice. The dextran sodium
sulfate (DSS) colitis model is one of the most widely used rodent models of intestinal
inflammation, reproducibly recapitulating the colonic damage and mucosal recovery of
UC,[22] despite its limitations.[22-27] Importantly, this model allows the precise timing of colitis
onset and development. We thus tracked the proportion and identity of actively replicating
gut bacterial cells, quantified their replication rates when possible, and estimated the
replication potential of the entire bacterial community over time.

As mice develop colitis, there are decreases in the proportions of replicating gut bacteria,
which revert to baseline values as the mice recover. We also report significant changes
in the composition of the replicating community of gut bacteria, as well as significant
changes in the abundance of specific taxa, such as the mucus-degrader *Akkermansia*
muciniphila and the poorly described *Erysipelatoclostridium* genus. Whole genome
shotgun sequencing approaches reveal dynamic replication rates for many taxa
throughout colitis development, including *A. muciniphila*. Lastly, we observed that the
development of colitis coincided with the preferential retention in the gut of bacterial
genomes with faster predicted doubling times. Our data support the need for additional
functional approaches to better understand the gut bacterial dynamics underlying the
progression of intestinal inflammation and the resulting altered microbial community.

Results

DSS reproducibly causes colitis and changes in fecal bacterial composition

We first confirmed that administration of 2% DSS for 5 days in the drinking water of the mice in our facility led to reproducible colitis. To do so, we monitored weight loss, blood in stool, and levels of fecal lipocalin-2 (LCN-2), a commonly used and sensitive marker of intestinal inflammation,[23] before, during, and after administration of DSS (**Figure 1A,** Error! Reference source not found.). We incorporated these three parameters into a modified scoring metric of colitis as described by Kim et al. (2012) based on the disease activity index (DAI) (**Error! Reference source not found.**, **Error! Reference source not found.**).[28] In this way, we defined the different health states of the mice during colitis development: baseline, pre-symptomatic, symptomatic, and recovery,[29] as detailed in the Methods.

Figure 1. *Experimental timeline and changes in gut bacterial community during DSS-colitis.*

(A) Timeline for each mouse experiment. After a 2-week acclimatization period, the feces of four cages of mice (two cages per sex, with 3 mice per cage) were collected over 7 days. Then, dextran sodium sulfate (DSS) was administered in the drinking water of the mice at a concentration of 2% (w/v) for 5 days, after which normal water was returned. Mouse feces continued to be collected throughout. Once DSS administration began, and thereafter, mouse health was monitored via measurements of blood in stool and lipocalin-2. Based on mouse health parameters, the phases of this study were split into four health states: baseline, pre-symptomatic, symptomatic, and recovery. (B) Taxonomic bar plots at the genus level of the fecal gut bacteria from each mouse during each health state (C) Beta diversity (Jaccard index) of the fecal gut bacterial communities from all cages, grouped per health state. D) Heat maps of standardized relative abundances of taxa from all cages of mice which were differentially abundant as compared to the baseline state. BA = Baseline; PS = Pre-symptomatic; SY = Symptomatic; RE = Recovery.

Amplicon sequencing of the 16S V4-5 region was conducted on bacterial DNA extracted
from mouse feces collected throughout the experiment, as described in the Methods.
Following this, QIIME2 was used to taxonomically identify the bacteria present in each
sample (**Figure 1B**).[30] The bacterial communities of these mice included the genera
Muribaculaceae (F1: 25.37% ± 9.36; F2: 34.20% ± 12.42; M1: 17.62% ± 9.30; M2: 17.93%
± 12.80) and *Bacteroides* (F1: 17.34% ± 7.50; F2: 14.08% ± 8.37; M1: 15.68% ± 11.89;
M2: 12.04% ± 7.91) of the Bacteroidota phylum; along with *Dubosiella* (F1: 0.64% ± 1.11;
F2: 10.15% ± 10.96; M1: 21.55% ± 15.63; M2: 26.41% ± 12.61), *Lachnospiraceae*
NK4136 group (F1: 7.58% ± 3.88; F2: 11.72% ± 10.09; M1: 5.01% ± 2.92; M2: 8.05% ±
5.10), and *Faecalibaculum* (F1: 24.01% ± 18.33; F2: 6.66 * 10^-3 % ± 2.40 * 10^-2; M1:
7.85% ± 1.72; M2: 7.09% ± 1.17) of the Bacillota phylum (**Figure 1B**).
190
When combining the data from all cages, we report no significant differences in the alpha
diversity between the different health states for any of the metrics we assessed (Error!
Reference source not found.**A**). Nevertheless, we saw a trend towards significance for
the Shannon index, supported by a significant Friedman's test (p = 0.044) but non-
significant post-hoc test (Wilcoxon signed-rank test; p = 0.375 for all comparisons) (**Error!**
Reference source not found.).
197
At the beta diversity level, we report significant differences in the bacterial communities
between baseline and both the symptomatic and recovery states based on the Jaccard
index (PERMANOVA, q = 0.039; **Figure 1C,** Error! Reference source not found.). Despite
no significant differences between the health states for the other calculated beta diversity
metrics (**Error! Reference source not found.**), we observed the same pattern of
deviation from baseline during the symptomatic state and partial return to near-baseline
values during recovery (Error! Reference source not found.**B-D**).
205
206   When grouping the bacterial diversity data by sex, significant differences were observed
207   in the beta-diversity values between health states (Error! Reference source not found.). F
208   or female mice (cages F1 and F2), this included significant differences in the unweighted
209   Unifrac distance between the baseline and symptomatic health states (PERMANOVA, q

= 0.027), and between the recovery and both the pre-symptomatic (PERMANOVA, q = 0.042) and the symptomatic (PERMANOVA, q = 0.027) health states (**Error! Reference source not found.iv**). There were no significant differences for the other beta diversity metrics (Error! Reference source not found.**i-iii**). For male mice (cages M1 and M2), there were significant differences in the unweighted Unifrac distance (UU), Jaccard index (JC), and Bray-Curtis dissimilarity (BC) between the baseline and symptomatic (PERMANOVA, UU: q = 0.02 ; JC: q = 0.036; BC: q = 0.039) and recovery (PERMANOVA , UU: q = 0.018; JC: q = 0.012; BC: q = 0.033) health states, and between the recovery and pre-symptomatic (PERMANOVA , UU: q = 0.0315; JC: q = 0.036; BC: q = 0.039) and symptomatic (PERMANOVA , UU: q = 0.02; JC: q = 0.015; BC: q = 0.033) health states (Error! Reference source not found.**i, ii, iv**). There were no significant differences for the weighted Unifrac distance (Error! Reference source not found.**iii**). These results suggest possible sex-specific differences in gut bacterial communities during the development of colitis.

We next sought to determine whether there were any changes in the abundance of specific taxa during the progression of colitis, using ANCOM-BC2 to conduct differential abundance analysis.[31] We compared the abundances of each taxon at a genus level between baseline and the other health states (see Methods). We identified 3, 9, and 17 taxa as differentially abundant in the pre-symptomatic, symptomatic, and recovery states, respectively (Error! Reference source not found.).

We noted a consistent increase in the abundance of *Erysipelatoclostridium* and a *Eubacterium fissicatena* group member in all health states as compared to the baseline, as well as the increased abundance of *Parasutterella* and *Faecalibaculum* during the symptomatic and recovery states (**Figure 1D;** Error! Reference source not found.). A *Lactobacillus* member consistently appeared at significantly higher abundances during baseline relative to the pre-symptomatic and symptomatic health states; whereas UCG-005 (from the *Oscillospiraceace* family), a *Ruminococcaceae* family member, a *Eubacterium coprostanoligenes* group member, and RF39 (from the *Bacilli* class) all had

240 higher relative abundances during baseline when compared to the symptomatic and
241 recovery health states (**Figure 1D;** Error! Reference source not found.).

Colitis alters the replicating and whole communities of gut bacteria

Next, we wanted to assess the proportions and identity of the replicating gut bacteria
during the development of, and recovery from, DSS colitis. As such, a second
independent mouse experiment with the same design was conducted, wherein the
recently optimized EdU-click technique was used to label newly synthesized bacterial
DNA.[13] EdU-click was combined with flow cytometry, as well as fluorescence-activated
cell sorting and 16S rRNA amplicon sequencing (FACSeq), to quantify the proportions of
actively replicating cells in fresh mouse feces and to taxonomically identify these bacteria
(see Methods and **Figure 2A**), respectively. The health status of the mice was monitored
and health states were determined as described earlier (**Error! Reference source not**
found.).

Figure 2. *Experimental workflow and changes in the proportion and composition of replication gut bacteria during DSS colitis.*

(A) Workflow for each mouse experiment. Whenever fecal pellets were collected, half were immediately used for EdU-click and FACSeq; the other half were frozen for later DNA extraction, whole genome shotgun sequencing, and subsequent analyses. (B-C) Median proportion of replicating (%EdU⁺) cells, represented as a line graph showing per-cage dynamics over time (B) and as boxplots, grouping all cages per health state (C). (D-E) Beta diversity (Jaccard indices) of the replicating (EdU⁺) (D) and whole community (Whole) (E) of bacterial cells for all cages, grouped by health state

The proportion of replicating fecal bacterial cells, denoted as %EdU$^+$ cells, changed as
the mice developed and recovered from DSS-induced colitis (**Figure 2B**). Specifically,
there was a notable but non-significant decrease in the median proportion of replicating
cells from 4.75% ± 3.73 during baseline to 2.33% ± 1.35 during the pre-symptomatic state
and 1.79% ± 0.94 during the symptomatic state, with a return to near-baseline values
during the recovery period (4.40% ± 1.89) (**Figure 2C**). Notably, the variability in the
median proportion of replicating cells between cages decreased during the pre-
symptomatic (1.35) and symptomatic (0.94) time periods as compared to the baseline
(3.73) and recovery periods (1.89) (Error! Reference source not found.**A,B**),
demonstrating that the proportion of replicating cells was most similar between cages
during active colitis.

We next wanted to determine if certain taxa preferentially replicate as mice develop colitis
and recover. To accomplish this, we conducted 16S rRNA amplicon sequencing on the
V4-5 region from DNA extracted from the replicating cells (EdU$^+$) and the whole
community (Whole) of bacteria originating from the same fecal sample (**Figure 2A**).

When considering the alpha diversity metrics, there were no significant differences in any
metric between the different health states for either the EdU$^+$ fraction or the Whole
community (Error! Reference source not found.**;** Error! Reference source not found.). The
lack of statistically significant differences is likely due to both inter-cage variability and low
numbers of cages (n = 4 cages).

We then compared the fecal bacterial communities of these mice across the different
health states using beta diversity metrics, for both the EdU$^+$ and Whole community of
sorted bacteria. We saw that there was grouping and separation between the different
health states in both fractions (**Figure 2D, E**). However, these differences were only
statistically significant with the Jaccard index for the EdU$^+$ fraction, between the
symptomatic state and both the baseline (PERMANOVA, q = 0.012) and the pre-
symptomatic (PERMANOVA, q = 0.012) health states (Error! Reference source not
found.**i**). As with the alpha diversity measurements, the lack of other statistically

significant differences (Error! Reference source not found.; Error! Reference source not

found.**ii-iv**) is likely due to both inter-cage variability as well as low numbers of cages (n

= 4 cages). When the two sexes were considered separately, only the Whole community

of the male mice (cages M1 and M2) showed a statistical difference in any alpha diversity

metric, being the weighted Unifrac distance, between the pre-symptomatic and recovery

states (PERMANOVA, q = 0.042) (Error! Reference source not found.**P;** Error! Reference

source not found.**iii**). No other beta diversity metrics were significant (Error! Reference

source not found.**A-O;** Error! Reference source not found.**,** Error! Reference source not

found.**,** Error! Reference source not found.**,** Error! Reference source not found.**i, ii, iv**)

We then sought to identify taxa which were differentially abundant in either the EdU$^+$ or

the Whole fraction in the different health states relative to the baseline. As before, we first

taxonomically identified the bacteria present in our samples using QIIME2 (**Figure 3A,**

B).

In the EdU$^+$ fraction, the most abundant genera included *Erysipelatoclostridium* (mean

28.76% ± 26.65), *Muribaculaceae* (mean 15.12% ± 11.19), *Lactobacillus* (mean 8.33% ±

13.05), *Lachnospiraceae* NK4A136 group (mean 6.38% ± 11.73), and *Dubosiella* (mean

9.23% ± 11.53) (**Figure 3A, B**).

C.

EdU+ Cells

Whole community

D.

In the Whole fraction, the most abundant genera included *Muribaculaceae* (mean 20.22%

± 14.59), *Dubosiella* (mean 36.09% ± 20.71), *Akkermansia* (mean 10.69% ± 11.58),

Turicibacter (mean 5.50 % ± 7.40), and *Lachnospiraceae* NK4A136 group (4.39% ± 6.38)

(**Figure 3A, B**).

We then used ANCOM-BC2 to conduct differential abundance analysis at the genus level.

In the replicating (EdU⁺) fraction, *Akkermansia* significantly increased its abundance

during the pre-symptomatic and symptomatic time points compared to the baseline state,

whereas in the Whole fraction, this taxon only increased during the symptomatic state

(Error! Reference source not found.). In contrast, *Erysipelatoclostridium* was significantly

increased in both the EdU⁺ fraction and the Whole fraction in all health states compared

to the baseline state (Error! Reference source not found.). In the Whole fraction during

the symptomatic state, there were also transient increases in a *Eubacterium fissicatena*

group member and *Turicibacter*, and decreases in *Lachnoclostridum*, *Blautia*, and

Roseburia, with no corresponding changes in the EdU⁺ fraction (Error! Reference source

not found.**A-D**).

The changes in the proportion of replicating cells (EdU⁺) of both *Akkermansia* and

Erysipelatoclostridium species mirrored their changes in total abundance (Whole),

suggesting the active and successful replication of these taxa, leading to increased total

biomass (**Figure 3C, D;** Error! Reference source not found.**G**). This increase was

transient for *Akkermansia* but seemed to persist for the *Erysipelatoclostridium* taxon.

Altered replication dynamics of fecal bacteria during colitis development

As a complementary method to EdU-click, we also used whole genome shotgun (WGS)

sequencing to quantify the replication dynamics of fecal bacteria on a subset of samples.

Bacterial genomes were assembled from WGS data into metagenome-assembled

genomes (MAGs) as described in the Methods. WGS sequencing was performed on fecal

samples from mice in cages F2 and M2 (see **Error! Reference source not found.** for

WGS statistics), as these mice had more severe signs of inflammation compared to their
biological replicate cages (Error! Reference source not found.).

The taxonomic composition and differentially abundant taxa of the MAGs, as well as their
beta diversity, are summarized in **Figure 4** and Error! Reference source not found.,
respectively. None of the beta diversity measurements showed significant differences in
the fecal bacterial composition (Error! Reference source not found.), likely due to the
limited number of genomes for which taxonomy could be determined (37 MAGs for cage
F2; 43 MAGs for cage M2). Nevertheless, differential abundance analysis showed a
significant increase in the relative abundance of an *Akkermansia muciniphila* MAG, and
a significant decrease in the relative abundance of two MAGs from the *Lachnospiraceae*
family, a 14-2 species and a COE1 species, during all health states as compared to the
baseline state (**Figure 4C-E**). There was also a significant increase in the relative
abundance of a *Duncaniella* MAG during the symptomatic state (**Figure 4D**).

370
Figure 4. *Relative abundance and differentially abundant taxa from whole genome shotgun sequencing*
(WGS) data.(A-B) Taxonomic bar plots at the genus level for the metagenome-assembled genomes. (C) Volcano
plots of metagenome-assembled genomes determined to be differentially abundant from the baseline. Note that
genomes 14-2 & COE1 are both from the Lachnospiraceae *family*

Next, we wanted to quantify the replication dynamics of these MAGs using two distinct

methods: the peak-to-trough (PTR) method, which estimates instantaneous replication

rates,[32] and the codon usage bias (CUB) method, which estimates the maximal replication

potential or minimal doubling time (DT).[33-35]

To calculate the replication rates during the development of DSS colitis for as many

quality genomes as possible, we used the PTR-based tool GRiD[36], which works well with

medium quality MAGs at low coverage (**Error! Reference source not found.**). GRiD is

specialized in using genomes with a coverage of as little as 0.2x and as many as 90

scaffolds/Mbp, and for genomes with at least 1x coverage, as little as 50% completion

and as much as 200 scaffolds/Mbp.[36] The other main PTR-based tools, iRep,[37] DEMIC,[38]

and CoPTR,[16] all have requirements or assumptions which are not appropriate for the

quality, coverage, or temporal nature of our data. The differences between these tools

have been thoroughly evaluated elsewhere and will not be covered here.[16,35,39]

From cage F2, we were able to calculate PTR values using GRiD across all time points

for 32 MAGs (corresponding to 53% of all MAGs from cage F2) (**Error! Reference source**

not found., **Error! Reference source not found.**); from cage M2, we calculated PTR

values across all time points for 43 MAGs (corresponding to 65% of all MAGs from cage

M2) (**Error! Reference source not found.**, **Error! Reference source not found.**). Of

these MAGs, we highlighted the replication rate dynamics of the following four:

Akkermansia muciniphila (**Figure 5A**), a *Duncaniella* species (**Figure 5B**),

Bifidobacterium globosum (**Figure 5C**), and a *Muribaculaceae* species (**Figure 5D**). We

selected these taxa based on whether they met all or most of the following criteria: they

were differentially abundant between the health states, had the highest median replication

rates during the symptomatic period, had replication rate measurements for all

experimental days, were one of the most abundant taxa overall or during the symptomatic

period, and had replication rate measurements estimated in both mouse cages.

A. *Akkermansia muciniphila*

B. *Duncaniella* sp910589485

C. *Bifidobacterium globosum*

D. *Muribaculaceae* CAG-485 sp002362485

409
Figure 5. *Replication dynamics of gut bacteria during DSS colitis. (A-D) Replication rates of representative*
taxa as calculated by GRiD, coloured by health state. (E) Correlation between health states and minimal doubling
times (DTs) for both cages, as calculated by gRodon.

These taxa appear to have cage-specific dynamics. For instance, in cage F2, the *A.*

muciniphila MAG had no correlation with the health status of the mice; whereas, in cage

M2, the *A. muciniphila* MAG had consistently higher replication rates once the mice

developed colitis relative to the baseline state (**Figure 5A**).

For the *Duncaniella* species, this MAG had its highest peak of replication rates during the

symptomatic period in cage F2; while the same MAG had the highest replication rates

during the baseline period relative to other periods in cage M2 (**Figure 5B**).

For *B. globosum*, this MAG had relatively steady replication rates in cage F2, except for

a rapid and transient decrease on the first day of the pre-symptomatic period. This same

MAG had decreased replication rates throughout the experiment starting after the first

baseline day in cage M2 (**Figure 5C**).

Finally, for the *Muribaculaceae* species, this MAG had its lowest replication rates during

the pre-symptomatic state in cage F2, whereas in cage M2 its replication rates were

highest during the baseline state and were consistently lower afterwards (**Figure 5D**).

We further determined no correlation between bacterial replication rates and the relative

abundance of their corresponding taxa in the whole community (Pearson r = -0.14756;

r^2 = 0.02177; Error! Reference source not found.).[16]

To glean more information about the replication abilities of the taxa we identified, we next

used gRodon, a CUB-based tool, to calculate the minimal doubling times (DTs) for each

MAG (Error! Reference source not found.).[34] This method provides information on the

replication potential of bacteria, as opposed to their instantaneous replication rates.

Previous studies have shown that measurements of replication potential using CUB-

based tools can more accurately predict the replication behaviour of certain taxa,

especially slower growing ones in complex communities.[39]

The median minimal DTs for all phyla in both cages was between 1.39 and 3.46 hours

(Error! Reference source not found.**A;** Error! Reference source not found.**,** Error!

Reference source not found.), supporting the notion that gut bacteria are fast replicators.[33]

Within the Bacillota phylum, there was a larger range (F2: 0.89-7.09 hrs; M2: 0.63-7.79

hrs) and more variability (F2: median absolute deviation (MAD) 1.67 hrs; M2: MAD 1.48

hrs) in the minimal DTs as compared to the other phyla with more than one genome

present (F2: Actinomycetota range: 0.98-2.33 hrs, MAD: 0.74 hrs; Bacteroidota range:

1.44-2.59 hrs, MAD: 0.66 hrs; M2: Actinomycetota range: 1.04-2.72 hrs, MAD: 0.64 hrs;

Bacteroidota range: 1.50-2.53 hrs, MAD: 0.55 hrs) (**Error! Reference source not found.**,

Error! Reference source not found.). This range and variability specifically occurred

within the Clostridia class (F2: range: 0.89 -7.09 hrs; MAD: 1.69 hrs; M2: range: 1.15-7.79

hrs; MAD: 1.40 hrs), which were comprised mostly of genomes from the *Lachnospiraceae*

family (F2: 15 genomes: range: 1.53-7.00 hrs; MAD: 0.57 hrs; M2: 30 genomes: range:

1.64-7.79 hrs; MAD: 1.28 hrs) (Error! Reference source not found.**B,C; Error! Reference**

source not found., **Error! Reference source not found.**).

To see whether there were any changes in the replication potential of all bacteria detected

over time, rather than individual taxa, we next used gRodon's metagenome mode to

calculate the minimal DTs of the whole community of gut bacteria during the development

of colitis.[35] There was a negative but non-significant ($p > 0.05$) slope when plotting the

minimal DTs of both cages of mice as a function of the health state, while accounting for

cage and day as random effects (see Methods) (**Figure 5E,** Error! Reference source not

found.**;** Error! Reference source not found.). However, this trend was close to significant

($p = 0.0743$) when the minimal DTs were plotted as a function of fecal LCN-2 levels, a

sensitive marker of colitis (Error! Reference source not found.**;** Error! Reference source

not found.)[40]. Both data indicate that the colitic environment could select for bacteria with

the capacity to replicate quickly.

As fast replication could be a strategy for bacterial taxa to persist in a perturbed

environment,[41] we also sought to describe the link between MAGs with the potential to

replicate quickly (i.e., with low minimal DTs estimated by gRodon) and those with

increased replication rates and/or increased relative abundances during the symptomatic
period.

When combining information from all of the MAGs and both cages together, we report no
correlation between the minimal DT of a MAG and its median replication rate during the
symptomatic period (r = 0.049; Error! Reference source not found.). In contrast, there
seemed to be a weak negative correlation between the minimal DT of a MAG and its
median relative abundance during the symptomatic period (r = -0.329), suggesting that
bacteria with the potential to replicate quickly tend to be at higher abundances during
active colitis (Error! Reference source not found.). However, our analyses are limited by
the rare high-abundance bacterial taxa and the prevalent low-abundance taxa.

Discussion

This study sought to determine the replication dynamics of gut bacteria during the
development of, and recovery from, intestinal inflammation in a mouse model of
chemically induced colitis. We found that the development of colitis resulted in decreased
proportions of replicating gut bacterial cells, as well as significant changes in the
composition of the replicating gut bacterial community. At the genus level, we report
significantly increased abundances of replicating *Akkermansia* and *Erysipelatoclostridium*
taxa, which preceded increases in their relative abundances in the community.
Replication rates of *A. muciniphilia, Bifidobacterium globosum,* a *Muribaculaceae*
species*,* and a *Duncaniella* species also changed during colitis. Lastly, we saw that the
progression of colitis coincided with an overall decrease in the predicted minimal doubling
time of the whole bacterial community. Collectively, our data suggest that the composition
of the mouse fecal bacterial community shifts towards taxa with the potential to replicate
faster as intestinal inflammation progresses.

DSS colitis results in broad-level and taxon-specific changes in
gut bacterial communities

While the role of gut microbial communities in intestinal inflammation is increasingly
studied, most studies remain cross-sectional. Thus, the dynamic alterations of the gut

microbiota during colitis progression remain less characterized.[19,20,29,42-50] By monitoring

murine fecal microbial communities in DSS colitis, we aimed to determine some of these

underlying microbial dynamics of colitis. While we do not report significant decreases in

alpha-diversity, unlike other DSS studies[48,50-52], our beta-diversity analyses suggest that

these gut bacterial communities may have reached an alternative stable state distinct

from baseline.[29,43,48,49,53] These dynamics were especially marked in male mice, possibly

explaining the more reproducible and severe phenotype typically observed for males in

this colitis model.

Our longitudinal approach allowed us to determine that differentially abundant gut

bacterial taxa could be categorized into 4 broad response groups, according to their

dynamics during colitis progression (Error! Reference source not found.). Taxa in Group

1 – which include RF39, *Oscillibacter,* a *Ruminococcaceae* member, and a *Clostridium*

vadin BB60 group member – had decreased abundances during the recovery period,

suggesting that they likely cannot survive a colitic environment. In contrast, Group 2 taxa,

including *Erysipelatoclostridium* and a *Eubacterium fissicatena* group member, had

increased abundances during colitis development and onwards, suggesting that these

taxa expanded when the host environment was perturbed, and were able to maintain their

increased abundances even as the host recovered. Other response groups include taxa

which decreased temporarily below the limit of detection during colitis but which returned

during the recovery period (Group 3; *Lactobacillus*); and those which only expanded

during recovery, possibly taking over a niche made available by the loss of other taxa

(Group 4; *Lachnospiraceace* UCG-008).

The limited existing literature on the above highlighted taxa suggests some possible

explanations for their dynamics. For instance, RF39 of Group 1, which has been shown

to decrease in abundance in mouse models of colitis,[54-65] has also been described as

having a reduced genome and as being metabolically dependent on other bacteria for

some amino acids and vitamins.[66,67] Our data support the hypothesis that auxotrophic

bacteria such as RF39 are less likely to survive harsh colitic environments, as has been

suggested to occur in IBD patients.[68,69]

In contrast, Group 2 bacterial taxa *Erysipelatoclostridium* and *E. fissicatena* persistently

increase after colitis development, as reported elsewhere.[70-87] While the mechanisms

behind this increase are unclear, our data align with the literature while also contributing

a finer time scale regarding their changes in abundance during colitis.

At least one other DSS colitis study has seen the growth patterns of Group 3 member

Lactobacillus reported here.[88] One putative explanation for such behaviour is that, since

Lactobacillus species colonize the forestomach of mice, these taxa can be washed into

the cecum and colon – especially during the diarrhea commonly seen in DSS colitis.[22-

24,49,89] If a favourable niche in these distal intestinal locations becomes available during a

perturbation, *Lactobacillus* may be able to establish there; whereas under homeostatic

conditions, this would be less likely to occur.[90] Furthermore, similarly to the earlier

discussed taxon RF39, the metabolic dependence of most *Lactobacillus* species likely

precludes their ability to thrive in an inflamed environment.[90] After the inflammation,

however, it may be possible that *Lactobacillus* from the forestomach establish in the

colon.

For the Group 4 member *Lachnospiraceae* UCG-008, little is known about its metabolism;

or indeed, the metabolism of most *Lachnospiraceae* species,[91] despite including many

SCFA-producers and metabolizers of complex plant-derived carbohydrates.[92]

Lachnospiraceae UCG-008 is likely a butyrate producer, since the *Lachnospiraceae*

family is known to contain many butyrate-producing members.[91-94] There is discrepancy

between different studies as to whether *Lachnospiraceae* members are increased or

decreased in various intestinal perturbations, and whether this is beneficial or detrimental

to the host.[91,94-99] It has been suggested that the impact of *Lachnospiraceae* members on

the host is likely species or strain specific.[91,94,96] As such, the overall role of

Lachnospiraceae UCG-008 in colitis remains unclear. One speculation as to how this

taxon thrives during colitis recovery is that it could create endospores.[100] Upon the return

to a more favourable environment, such as when the mice begin recovering from colitis,

these endospores may allow *Lachnospiraceae* UCG-008 to rapidly take advantage of
newly opened niches in the gut.

Decreased proportions of replicating gut bacteria during colitis

To determine if the aforementioned patterns of altered abundances reflected changes in
gut bacterial replication, we quantified and taxonomically identified the replicating gut
bacteria using EdU-click in an independent study. We observed a decreased proportion
of replicating bacteria during the pre-symptomatic and symptomatic stages, suggesting
that the inflammatory environment is detrimental for the replication of most taxa.

Importantly, we noted that the replicating gut bacterial community during the symptomatic
state was markedly different from other health states. Narrowing down to a per-taxon
basis, we report significant differences in the abundance of *Akkermansia* and
Erysipelatoclostridium in both the replicating fraction and in the whole bacterial
community.

Since *Akkermansia* began increasing its number of replicating cells during the pre-
symptomatic state, and then subsequently increased its abundance in the whole fraction
during the symptomatic state, this suggests that its increased number of replicating cells
successfully led to an increase in its biomass. Once the mice began to recover, however,
the abundance of this taxon quickly decreased back to baseline values.

The impact of *Akkermansia* in DSS colitis is unclear, as it is associated with health in
humans and is depleted in individuals with IBD,[101] whereas it is commonly seen to
increase during DSS colitis in mice.[29,102-104] In addition, while some studies have found
that mice with higher levels of *Akkermansia* exhibit less severe colitis,[105] supporting a
protective role for this taxon, other studies have shown the reverse.[103] More recent studies
highlight that the impact of *Akkermansia* on host health could be strain specific[106] or
depend on the microbial and immunological status of the host.[107] As a mucus degrader,
Akkermansia could worsen inflammation,[108] rendering the host intestinal epithelium more
accessible to microbial translocation. Alternatively, its mucus metabolism could stimulate

increased mucus secretion by the host,[109] as some *Bacteroides* species have been

shown to do.[110] An emerging hypothesis for the transient increase in *Akkermansia* during

DSS colitis is that a DSS-dependent increase in mucus permeability[89] allows this taxon

greater access to its preferred carbon source.[111] *Akkermansia* is also likely to be

somewhat resistant to the increased oxidative environment of colitis, given that there are

higher levels of oxygen present at the mucus layer where this taxon resides.[112] The

combination of its resistance to oxidative stress and an increased access to its preferred

energy source creates a perfect environment for *Akkermansia* to thrive, at least until host

mucosal recovery begins.[113]

The *Erysipelatoclostridium* taxon was increased at all time points relative to the baseline

state, in both the replicating and the whole bacterial community. Despite a limited number

of studies noting the contrary[99,114-116], these results are in accordance with the behaviour

of *Erysipelatoclostridium* reported in prior studies during intestinal disturbance. Indeed,

this genus and one of its species, namely *Clostridium ramosum* (also known as *E.*

ramosum or *Thomasclavelia ramosum*),[117-119] have been associated with intestinal

inflammation or injury in humans,[46,48,73,120-127] increasing in abundance in mouse models

of colitis[71-82] and infection[128], and being positively correlated with fecal water content (i.e.,

diarrhea), a common colitis symptom[2,22-24,129,130]. The high replication activity of

Erysipelatoclostridium we report demonstrates that it is a particularly active taxon during

colitis, resulting in a possibly outsized impact on the microbial environment and the host.

What allows *Erysipelatoclostridium* to benefit from perturbations can only be speculated,

since its metabolism is not well described,[70] despite being a dominant gut Clostridial

taxon.[131] The best described member of this genus is *C. ramosum*, which is reported to

produce low amounts of acetate and possibly butyrate, and can significantly increase the

number and function of anti-inflammatory Treg cells.[132,133] These capabilities point

towards the potential anti-inflammatory role of this taxon, without explaining its continual

increase in abundance in perturbed intestinal conditions.

As more thoroughly discussed by Beauchemin et al.,[13] not all bacterial taxa can be
labelled with EdU, and labeled bacteria likely do not equally incorporate EdU into their
DNA. As such, this study can only report on replicating taxa amenable to EdU-click, and
likely does not capture all replicating taxa. As one way to address this limitation, we also
used metagenomics-based bioinformatic tools to calculate bacterial replication dynamics
on a subset of our data, as discussed below.

Taxon-specific changes in gut bacterial replication rates and increased abundances of fast replicating gut bacteria during colitis

Stool samples from ten non-consecutive sampling times between days 1-18 in one cage
of female mice (F2) and one cage of male mice (M2) underwent shotgun sequencing.
This allowed us to quantify the replication dynamics of several metagenome-assembled
genomes (MAGs) during colitis development, using both PTR-based and CUB-based
approaches. For the most part, the oscillatory replication rates for each taxon in each
cage went back to near-baseline values, with no clear link to the host health states – with
the exception of *A. muciniphila*. In agreement with our EdU-click and differential
abundance analyses, the *A. muciniphila* genome seen in both cages replicated most
rapidly once DSS was administered, after which its replication rates decreased back to
near baseline values as colitis developed.

We also observed cage-specific replication rate dynamics of the same gut taxa. For
instance, the *Duncaniella*, *Bifidobacterium globosum,* and *Muribaculaceae* CAG-485
genomes in cage M2 had decreased replication rates as colitis developed, whereas in
cage F2 these same taxa had variable replication rates. These differences suggest
individual-specific gut bacterial replication dynamics, as reported in previous human
cohorts[16], in addition to possible sex- and cage-specific responses to inflammation. Since
Duncaniella, Muribaculaceae CAG-485, and *B. globosum* are not as well described as *A.*
muciniphila, the possible explanations underlying their replication rate patterns remain to
be determined.

We additionally report a lack of correlation between gut bacterial replication rates and

their associated relative abundances, as noted by other studies[16,32,37,134]. Indeed, relative

abundance measurements account for the culmination of bacterial growth and death,

whereas replication rates only account for the former, emphasizing that both approaches

measure distinct aspects of the gut bacterial community.[16,32,37]

We next used an independent metric based on genome-specific CUB bias, using gRodon,

to determine possible links between gut bacterial replication and the response of specific

taxa during inflammation. When calculating the replication potential of the whole gut

bacterial community over time, we noted slightly lower minimal doubling times (i.e.,

increased maximal replication rates) as colitis developed or as levels of LCN-2 increased.

This suggests that faster replicating gut bacteria may be preferentially selected for in an

inflammatory environment.

These findings were further corroborated by the negative correlation between the minimal

doubling time (DT) of a MAG and its median relative abundance during the symptomatic

period, suggesting that taxa with the ability to replicate quickly are present at higher

abundances during colitis. This is in line with previous research which suggests that faster

replicating bacteria can adapt more quickly to changing environments,[33,135,136] that gut

bacteria during IBD flares may be replicating more than during remission,[137] and that fast

replication is one strategy gut bacteria can use to maintain or increase their biomass when

cell turnover and intestinal sloughing are high [41] as in DSS-induced colitis.[22,23,138,139]

We further report a lack of correlation between gut bacterial minimal doubling times and

replication rates during the symptomatic period. This suggests that replication potential

and replication rates are also measurements of fundamentally different processes,

although this remains to be validated.

It should be noted that our WGS analysis has limitations. We could only assemble

approximately 35-36% of the sequences obtained into MAGs, of which only 53-65% could

be both taxonomically characterized and have their replication rates estimated using

GRiD (**Error! Reference source not found.**). This can be partly attributed to the relatively

shallow sequencing (median 12,775,255 reads; range 8,448,754 – 19,756,090 reads),

resulting in the low coverage of many genomes for most time points. Despite this, we

believe that this study provides a foundational representation of the replication dynamics

of the most abundant gut taxa during DSS colitis development, and argue that bacterial

replication rates should be measured in ongoing microbiome workflows where WGS data

is acquired. We have created and shared on GitHub (see Data Accessibility) our workflow

for genome assembly and binning, as well as subsequent MAG classification and

bacterial replication measurements, to help researchers proceed with such analyses.

694 Conclusions and Future Perspectives

Overall, this work demonstrates the utility of tracking various facets of gut bacterial

replication alongside measurements of cell abundance and taxonomy during the

development of intestinal inflammation. We provide further evidence for the relevance of

some taxa, such as *Akkermansia*, and describe new gut bacterial replication dynamics,

such as those for *Erysipelatoclostridium*, in a DSS mouse model of colitis.

These taxa should be further studied for their relevance to host health.[64] For instance,

determining the shared metabolic characteristics of gut bacteria which persist, and/or

thrive, in the inflamed gut could reveal how these taxa or their metabolism may be linked

to the development of an inflammatory environment. Such taxa or metabolic activities

could act as biomarkers, wherein changes in their presence could indicate the onset of a

flare before overt symptoms manifest. This information could further help predict how a

particular gut bacterial community will respond to inflammation, depending on the

composition and activity of the community members.[140]

Simplified *in vitro or in vivo* systems (e.g., artificial guts, gut-on-a-chip, or bacterial

consortia like the OMM12 community[141]) will be particularly useful, since the complexity

of the native gut microbiota and the host environment severely limit our ability to

disentangle the myriad microbe-microbe and host-microbe interactions occurring during

colitis development. Such approaches could, for example, allow for the direct comparison

of transcripts or metabolites produced by thriving and non-thriving gut taxa in an
inflammatory intestinal environment.

Evaluating gut bacterial replication dynamics provides insight into bacterial activity which
cannot be captured from abundance measurements alone. For instance, quantifying
bacterial replication rates could be beneficial when following colonization dynamics in
early life, to improve our understanding of how microbial activity is linked to the known
succession dynamics observed therein.[33,37] Characterizing bacterial replication dynamics
could also be useful when determining whether members of a probiotic cocktail are
actively replicating in the gut after administration,[142] or how the introduction of a pathogen
affects bacterial community replication patterns. On a conceptual level, describing various
aspects of gut bacterial replication could help uncover the role of r- and K-selection in the
gut. Such research will contribute to increasing our knowledge in a variety of ecosystems
on this fundamental, yet poorly described, characteristic of complex microbial
communities.

730 Methods

731 Animals
Equal numbers of 6–8-week-old specific-pathogen free (SPF) male and female wild-type
C57BL/6 mice (Jackson Laboratories) were kept at the Goodman Cancer Research
Center Animal Facility at McGill University, in accordance with the McGill Ethics Research
Board (animal ethics protocol MCGL-7999). For each experiment, there were six mice of
each sex, and mice were housed three per cage according to sex, with a total of four
cages per experiment.

Before administration of dextran sodium sulfate (DSS), mice were allowed to acclimate
to the animal facility for two weeks. On the third week, fecal pellets were collected from
the mice on three non-consecutive days. Fecal pellets were collected per cage by putting
all mice from a single cage into a new, sterile cage, empty of corncob bedding and nesting
material, with no access to food or water. Feces were collected soon after defecation

using ethanol-cleaned tweezers reserved for this purpose. Feces from mice of the same

cage were put into the same 1.5 mL microcentrifuge tube. This procedure lasted no longer

than one hour, after which mice were placed back into their respective cages. Immediately

after collection, the fecal pellets were either processed in an anaerobic chamber or frozen

at -80°C for later DNA extraction, as detailed below.

The week following acclimatization, mice were administered 2% DSS (molecular weight:

36-50 kDA; MP Biomedicals) in the drinking water for 5 days, and fecal pellets were

collected daily except on weekends. Afterwards, the mice were returned to the facility

water line, and fecal pellets continued to be collected daily, except on weekends, for

another 6 days, after which fecal pellets were collected twice weekly for an additional two

to three weeks.

Mouse health parameters, including body weight, blood in stool, and levels of fecal

lipocalin-2, were monitored throughout the experiment. Body weight was measured per

mouse daily except on weekends. Blood in stool was detected using the Hemoccult

Sensa kit (Beckman Coulter). Levels of fecal lipocalin-2 (LCN-2) were quantified using

the mouse lipocalin 2 DuoSet enzyme-linked immunosorbent assay (ELISA) (R&D). Fecal

samples were prepared for the ELISA using the method of Chassaing et al. (2012)[40] and

diluted 10-fold to 10,000-fold depending on the concentration of LCN-2 in the sample.

The ELISA was performed according to the manufacturer's instructions.

The health states of the mice were determined based on body weight, blood in stool, and

fecal lipocalin-2 levels, as detailed in **Error! Reference source not found.**. Baseline

days were defined as those before DSS administration, during which the mice had no

clinical symptoms of colitis. Pre-symptomatic days were defined as those during DSS

administration where the mice had minimal symptoms of colitis, including minor weight

loss and increased levels of fecal LCN-2 up to 100 ng per gram of feces (ng/g).

Symptomatic days were defined as those during or after DSS administration where the

mice had weight loss of 1-10%, increased levels of fecal LCN-2 exceeding 100 ng/g,

and/or the presence of blood in their stool. Recovery days were defined as the days after

DSS administration cessation and during which the mice began to gain weight and no
longer had blood in their stool, though high levels (≥100 ng/g) of fecal LCN-2 still remained
(**Figure 1**).
778
We randomly selected fecal samples collected from each cage of mice, rather than
tracking each individual mouse, due to our need for large quantities of fecal samples to
conduct our experiments. The microbial composition of the mice in a single cage is
thought to be very similar due to mouse coprophagic behaviour,[143-145] and as such, fecal
samples taken on a per-cage basis are expected to be broadly representative of all the
mice in a single cage.
785
786   This experimental setup was conducted independently two times, as detailed in the main
787   text.

Isolation of bacteria from mouse feces, EdU-click, and flow cytometry

Freshly collected fecal samples from mice were transferred to an anaerobic chamber
(Coy Laboratory Products; 5% H_2, 20% CO_2, 75% N_2) within one hour of collection.
Isolation of bacteria from mouse feces, EdU labeling, EdU-click, flow cytometry, and cell
sorting were performed as described previously.[13] EdU-positive (EdU⁺) and EdU-negative
(EdU⁻) cells and the whole community were sorted based on their level of fluorescence in
the appropriate channel and quantified as described previously.[13]

DNA extraction and 16S rRNA amplicon sequencing and analysis

Bacterial DNA was extracted from sorted cells and mouse fecal pellets using the AllPrep
PowerFecal DNA/RNA kit (Qiagen) as previously described.[13] Extracted DNA underwent
library preparation and 16S rRNA gene sequencing of the V4-V5 hypervariable region
with 515F/926R primers using the Illumina MiSeq PE250 system at the Université de
Québec à Montréal (UQAM) Center of Excellence in Research on Orphan Diseases –
Foundation Courtois (CERMO-FC) genomics platform.
803

Sequenced DNA was analyzed using QIIME2[30] version 2022.2 as previously described.[13]
Sequences were rarified to the depth of the sample with the fewest reads before
conducting alpha and beta diversity analyses. Taxonomic classification and relative
abundance measurements were conducted on the quality-controlled, filtered sequences
using QIIME2's machine-learning program trained on the 16S V4-V5 regions from our
data set, using the Silva 132 database on 99% operational taxonomic units. Reads
present in the sheath fluid but absent in the Whole community samples were identified as
contaminants and removed.

Further analysis was performed using phyloseq (version 1.44.0)[146] in R (version 4.3.0)
running in RStudio (2023.03.1). Beta diversity measurements were calculated on
weighted or unweighted UniFrac distances, Jaccard indices, or Bray-Curtis dissimilarities
using rbiom (version 1.0.3).[147] Repeated measures, multiple comparisons
PERMANOVAs were calculated with 999 permutations to test for significance using the
pairwiseAdonis package (version 0.4.1),[148] with multiple comparisons being corrected
using the Benjamini-Hochberg (BH) false discovery rate (FDR) correction. Differential
abundance analyses were performed on the combined data from all cages, using
ANCOM-BC2 (version 2.1.4)[31], and correcting for cage origin and the repeated measures
nature of the data, as described in *Statistical tests* and shown in the code available on
GitHub (see *Data availability*).

824   Whole genome shotgun sequencing and analysis

For fecal samples from two cages (F2 and M2, corresponding to a cage of female mice
and a cage of male mice, respectively) from the second experiment, DNA extracted from
the feces collected from these cages underwent library preparation and whole genome
shotgun (WGS) sequencing using the Illumina Miseq PE150 system at SeqCenter
(Pittsburgh, PA). The resulting bacterial genomes were assessed for their quality using
FastQC (version 0.11.9)[149] and the resulting FastQC files were visualized using MultiQC
(version 1.12).[150] Sequences were then co-assembled per cage across all sampling days
using Megahit (version 1.2.9).[151] Bowtie2 (version 2.4.4)[152] was used to remove reads of
host origin and to assess the extent to which assembled contigs represented the original,

quality-controlled reads. Binning of bacterial genomes from the co-assemblies was
performed using CONCOCT (version 1.1.0),[153] MaxBin2 (version 2.2.7),[154] and MetaBAT
2 (version 2.14),[155] followed by DASTool (version 1.1.4),[156] the latter of which identifies
the best bins from the three aforementioned binning tools. Bin quality was assessed using
CheckM (version 1.2.0).[157] Only bins meeting the minimum MIMAGS standards of
medium quality genomes[158] went on to be taxonomically identified using GTDB-Tk
(version 2.1.0).[159] The relative abundance of each bin was calculated based on the length
of each genome and its coverage (see code under *Data availability*).

842 Quantification of replication rate dynamics

Bins meeting the minimum MIMAGS standards of medium quality (≥50% complete, <10%
contamination)[158] were assessed for their potential maximal replication rates using
gRodon2 (version 2.3.0).[35] The maximal potential replication rate for the entire bacterial
community each day was calculated from the prokka (version 1.14.6)[160] protein
annotation output of per-cage, per-day assemblies as specified in the gRodon2 vignette
(http://microbialgamut.com/gRodon-vignette), using metagenome mode V2 and with n_le
= 1000 and all other parameters as default.

Bins meeting the minimum standards for GRiD (>=50% completeness, <=90
scaffolds/Mbp, >=x0.2 coverage) were assessed with this tool using default parameters.
An index was created for each bin using Bowtie2, and each sample (i.e., each day) per
cage was mapped to each appropriate index (i.e., each bin) that came from that cage.
The resulting SAM files from mapping were used as input for GRiD, using the "grid single"
mode.

857 Correlations

To correlate gut bacterial replication rates with their relative abundances, we used the
approach of Joseph TA et al. (2020).[16] Briefly, we first log2-transformed all relative
abundance and replication rate (GRiD) values. Then, we standardized the log2-
transformed relative abundance and GRiD values per taxon using the scale() function in
R. The standardized, log2-transformed values for abundance and GRiD were then

correlated using Pearson correlation. We proceeded with the same approach to correlate

gut bacterial replication potentials with their corresponding median replication rates during

the symptomatic period, and to correlate gut bacterial replication potential with their

corresponding median relative abundances during the symptomatic period.

Statistical tests

Statistical testing was performed according to the non-parametric, repeated measures

(i.e., longitudinal), and multiple comparisons nature of the study design as necessary, as

described in the results and/or figure legends, where appropriate.

Comparisons of the proportion of EdU^+ cells (%EdU^+ cells) between health states were

performed by first calculating the median %EdU^+ cells per cage per health state, as the

median is more robust to outliers than the mean. Then, the median %EdU^+ cells per health

state, considering all four cages in each health state, were compared to each other using

the Friedman test, accounting for cage as a random intercept effect.

Comparisons of alpha diversity metric values (Shannon index, Simpson evenness, and

observed features) between health states was performed by first calculating the median

alpha diversity value per cage per health state. Then, the median alpha diversity values

per health state, considering all four cages in each health state, were compared to each

other using the Friedman test, accounting for cage as a random variable. Where

appropriate, the Wilcoxon sign-rank post-hoc test was conducted, wherein the baseline

group was compared to all of the other health states, with the p-value adjusted for multiple

comparisons using the Bonferroni correction.

PERMANOVA comparisons of the beta diversity between different health states of mice

were conducted using pairwiseAdonis (version 0.4.1).[148] These comparisons accounted

for the repeated measures design of the study as described in the available code and as

first described here: https://thebiobucket.blogspot.com/2011/04/repeat-measure-adonis-

lately-i-had-to.html. Multiple comparisons were conducted using the pairwiseAdonis

package in R. When data from the various cages were compiled together, cage origin
was considered as a random intercept effect.
894
895   ANCOM-BC2 (version 2.1.4)[31] was used to detect differentially abundant bacteria
896   between health states as described in the main text and by the creators of the ANCOM-
897   BC2 package. All health states were compared to the baseline state, using Dunnett's test.
898   When data from all cages of mice were compiled together, cage origin was considered
899   as a random intercept effect.
900
901   To determine to what extent the health state of the host could explain the minimal doubling
902   time of the whole community of bacteria, we fit a linear mixed-effects model to our data
903   using the lme4 (version 1.1-34)[161] and lmerTest (version 3.1-3)[162] packages in R, with
904   health state and minimal doubling time as fixed effects, and cage and sampling day as
905   random intercept effects.

Data availability
Bacterial 16S rRNA gene sequencing data from all experimental arms are available in the
NCBI SRA database under project number PRJNA1040262 and PRJNA1040873,
respectively. Whole genome shotgun sequencing data from the second experiment is
available in the NCBI SRA database under project number PRJNA1040965. Code related
to the analysis of these data have been deposited in GitHub under the account at
https://github.com/evetb, repository DSS_Manuscript. Raw flow cytometry data have
been deposited in FigShare (https://figshare.com/projects/DSS_Manuscript/193004).

Declarations

Disclosure of interest
916   The authors report there are no competing interests to declare.

Authors' contributions

ETB designed, performed, analyzed, and interpreted experiments. CH aided with extracting DNA and supernatant for lipocalin-2 assays for the first mouse experiment. CFM helped develop the project, obtained funding, and helped design experiments and interpret the data. ETB and CFM edited the manuscript and approved the final draft.

Funding

This work was supported by a Frederick Banting and Charles Best Canada Graduate Scholarship-Master's (CGS M) from the Canadian Institutes of Health Research (CIHR), a Ferrings Pharmaceuticals Fellowship from the McGill Faculty of Medicine and Health Sciences, and a doctoral research grant from the Fonds de Recherche du Québec – Nature et Technologies (FRQNT) to ETB. This work was also supported by the Canada Research Chair program (grant 950-230748 X-242502), a CIHR transition grant, the Kenneth Rainin Foundation, and an Owens Catchpaugh IBD Graduate Research Award from the MUHC Foundation to CFM.

References

931 1   Caruso, R., Lo, B. C. & Núñez, G. Host-microbiota interactions in inflammatory
932 bowel disease. *Nat Rev Immunol* **20**, 411-426 (2020). https://doi.org:10.1038/s41577-
933 019-0268-7
934
2   Wang, R., Li, Z., Liu, S. & Zhang, D. Global, regional and national burden of
inflammatory bowel disease in 204 countries and territories from 1990 to 2019: a
systematic analysis based on the Global Burden of Disease Study 2019. *BMJ Open* **13**,
e065186 (2023). https://doi.org:10.1136/bmjopen-2022-065186
3   Li, J., Butcher, J., Mack, D. & Stintzi, A. Functional impacts of the intestinal
microbiome in the pathogenesis of inflammatory bowel disease. *Inflamm Bowel Dis* **21**,
139-153 (2015). https://doi.org:10.1097/mib.0000000000000215
4   Schirmer, M., Garner, A., Vlamakis, H. & Xavier, R. J. Microbial genes and
pathways in inflammatory bowel disease. *Nat Rev Microbiol* **17**, 497-511 (2019).
https://doi.org:10.1038/s41579-019-0213-6
5   Lane, E. R., Zisman, T. L. & Suskind, D. L. The microbiota in inflammatory bowel
disease: current and therapeutic insights. *J Inflamm Res* **10**, 63-73 (2017).
https://doi.org:10.2147/jir.S116088
6   Lloyd-Price, J. *et al.* Multi-omics of the gut microbial ecosystem in inflammatory
bowel diseases. *Nature* **569**, 655-662 (2019). https://doi.org:10.1038/s41586-019-1237-
9
7   Lee, J. Y., Tsolis, R. M. & Bäumler, A. J. The microbiome and gut homeostasis.
Science **377**, eabp9960 (2022). https://doi.org:10.1126/science.abp9960
8   Pisani, A. *et al.* Dysbiosis in the Gut Microbiota in Patients with Inflammatory
Bowel Disease during Remission. *Microbiol Spectr* **10**, e0061622 (2022).
https://doi.org:10.1128/spectrum.00616-22
9   Ho, G. T., Cartwright, J. A., Thompson, E. J., Bain, C. C. & Rossi, A. G.
Resolution of Inflammation and Gut Repair in IBD: Translational Steps Towards
Complete Mucosal Healing. *Inflamm Bowel Dis* **26**, 1131-1143 (2020).
https://doi.org:10.1093/ibd/izaa045
10   Schirmer, M. *et al.* Dynamics of metatranscription in the inflammatory bowel
disease gut microbiome. *Nat Microbiol* **3**, 337-346 (2018).
https://doi.org:10.1038/s41564-017-0089-z
11   Morgan, X. C. *et al.* Dysfunction of the intestinal microbiome in inflammatory
bowel disease and treatment. *Genome Biol* **13**, R79 (2012). https://doi.org:10.1186/gb-
2012-13-9-r79
12   Zhang, X. *et al.* Metaproteomics reveals associations between microbiome and
intestinal extracellular vesicle proteins in pediatric inflammatory bowel disease. *Nat
Commun* **9**, 2873 (2018). https://doi.org:10.1038/s41467-018-05357-4
13   Beauchemin, E. T., Hunter, C. & Maurice, C. F. Actively replicating gut bacteria
identified by 5-ethynyl-2'-deoxyuridine (EdU) click chemistry and cell sorting. *Gut
Microbes* **15**, 2180317 (2023). https://doi.org:10.1080/19490976.2023.2180317
14   Olm, M. R. *et al.* Necrotizing enterocolitis is preceded by increased gut bacterial
replication, Klebsiella, and fimbriae-encoding bacteria. *Science Advances* **5**, eaax5727
https://doi.org:10.1126/sciadv.aax5727

15   Riglar, D. T. *et al.* Bacterial variability in the mammalian gut captured by a single-
cell synthetic oscillator. *Nat Commun* **10**, 4665 (2019). https://doi.org:10.1038/s41467-
019-12638-z
16   Joseph, T. A., Chlenski, P., Litman, A., Korem, T. & Pe'er, I. Accurate and robust
inference of microbial growth dynamics from metagenomic sequencing reveals
personalized growth rates. *Genome Res* **32**, 558-568 (2022).
https://doi.org:10.1101/gr.275533.121
17   Quince, C., Walker, A. W., Simpson, J. T., Loman, N. J. & Segata, N. Shotgun
metagenomics, from sampling to analysis. *Nat Biotechnol* **35**, 833-844 (2017).
https://doi.org:10.1038/nbt.3935
18   Koch, B. J. *et al.* Estimating taxon-specific population dynamics in diverse
microbial communities. *Ecosphere* **9**, e02090 (2018).
https://doi.org:https://doi.org/10.1002/ecs2.2090
19   Galipeau, H. J. *et al.* Novel Fecal Biomarkers That Precede Clinical Diagnosis of
Ulcerative Colitis. *Gastroenterology* **160**, 1532-1545 (2021).
https://doi.org:10.1053/j.gastro.2020.12.004
20   Raygoza Garay, J. A. *et al.* Gut Microbiome Composition is associated with
future onset of Crohn's Disease in Healthy First-Degree Relatives. *Gastroenterology*
(2023). https://doi.org:10.1053/j.gastro.2023.05.032
21   Sasson, G. *et al.* A223 THE GEM PROJECT: ASSOCIATIONS OF DIETARY
PATTERNS WITH MICROBIOME AND FECAL CALPROTECTIN IN HEALTHY FIRST-
DEGREE RELATIVES OF CROHN'S DISEASE PATIENTS. *Journal of the Canadian*
Association of Gastroenterology **3**, 98-99 (2020).
https://doi.org:10.1093/jcag/gwz047.222
22   Eichele, D. D. & Kharbanda, K. K. Dextran sodium sulfate colitis murine model:
An indispensable tool for advancing our understanding of inflammatory bowel diseases
pathogenesis. *World J Gastroenterol* **23**, 6016-6029 (2017).
https://doi.org:10.3748/wjg.v23.i33.6016
23   Chassaing, B., Aitken, J. D., Malleshappa, M. & Vijay-Kumar, M. Dextran sulfate
sodium (DSS)-induced colitis in mice. *Curr Protoc Immunol* **104**, 15.25.11-15.25.14
(2014). https://doi.org:10.1002/0471142735.im1525s104
24   Perše, M. & Cerar, A. Dextran sodium sulphate colitis mouse model: traps and
tricks. *J Biomed Biotechnol* **2012**, 718617 (2012). https://doi.org:10.1155/2012/718617
25   Rogler, G. *et al.* Diet and Inflammatory Bowel Disease: What Quality Standards
Should Be Applied in Clinical and Laboratory Studies? *Molecular Nutrition & Food*
Research **65**, 2000514 (2021). https://doi.org:https://doi.org/10.1002/mnfr.202000514
26   Solomon, L. *et al.* The dextran sulphate sodium (DSS) model of colitis: an
overview. *Comparative Clinical Pathology* **19**, 235-239 (2010).
https://doi.org:10.1007/s00580-010-0979-4
27   Wirtz, S. *et al.* Chemically induced mouse models of acute and chronic intestinal
inflammation. *Nature Protocols* **12**, 1295-1309 (2017).
https://doi.org:10.1038/nprot.2017.044
28   Kim, J. J., Shajib, M. S., Manocha, M. M. & Khan, W. I. Investigating intestinal
inflammation in DSS-induced model of IBD. *J Vis Exp* (2012).
https://doi.org:10.3791/3678

29   Taguer, M. *et al.* Changes in Gut Bacterial Translation Occur before Symptom
Onset and Dysbiosis in Dextran Sodium Sulfate-Induced Murine Colitis. *mSystems* **6**,
e0050721 (2021). https://doi.org:10.1128/mSystems.00507-21
30   Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome
data science using QIIME 2. *Nat Biotechnol* **37**, 852-857 (2019).
https://doi.org:10.1038/s41587-019-0209-9
31   Lin, H. & Peddada, S. D. Analysis of compositions of microbiomes with bias
correction. *Nat Commun* **11**, 3514 (2020). https://doi.org:10.1038/s41467-020-17041-7
32   Korem, T. *et al.* Growth dynamics of gut microbiota in health and disease inferred
from single metagenomic samples. *Science* **349**, 1101-1106 (2015).
https://doi.org:10.1126/science.aac4812
33   Vieira-Silva, S. & Rocha, E. P. The systemic imprint of growth and its uses in
ecological (meta)genomics. *PLoS Genet* **6**, e1000808 (2010).
https://doi.org:10.1371/journal.pgen.1000808
34   Weissman, J. L., Hou, S. & Fuhrman, J. A. Estimating maximal microbial growth
rates from cultures, metagenomes, and single cells via codon usage patterns. *Proc Natl*
Acad Sci U S A **118** (2021). https://doi.org:10.1073/pnas.2016810118
35   Weissman, J. L., Peras, M., Barnum, T. P. & Fuhrman, J. A. Benchmarking
Community-Wide Estimates of Growth Potential from Metagenomes Using Codon
Usage Statistics. *mSystems* **7**, e0074522 (2022).
https://doi.org:10.1128/msystems.00745-22
36   Emiola, A. & Oh, J. High throughput in situ metagenomic measurement of
bacterial replication at ultra-low sequencing coverage. *Nat Commun* **9**, 4956 (2018).
https://doi.org:10.1038/s41467-018-07240-8
37   Brown, C. T., Olm, M. R., Thomas, B. C. & Banfield, J. F. Measurement of
bacterial replication rates in microbial communities. *Nat Biotechnol* **34**, 1256-1263
(2016). https://doi.org:10.1038/nbt.3704
38   Gao, Y. & Li, H. Quantifying and comparing bacterial growth dynamics in multiple
metagenomic samples. *Nat Methods* **15**, 1041-1044 (2018).
https://doi.org:10.1038/s41592-018-0182-0
39   Long, A. M., Hou, S., Ignacio-Espinoza, J. C. & Fuhrman, J. A. Benchmarking
microbial growth rate predictions from metagenomes. *ISME J* **15**, 183-195 (2021).
https://doi.org:10.1038/s41396-020-00773-1
40   Chassaing, B. *et al.* Fecal lipocalin 2, a sensitive and broadly dynamic non-
invasive biomarker for intestinal inflammation. *PLoS One* **7**, e44328 (2012).
https://doi.org:10.1371/journal.pone.0044328
41   Vandeputte, D. *et al.* Stool consistency is strongly associated with gut microbiota
richness and composition, enterotypes and bacterial growth rates. *Gut* **65**, 57-62 (2016).
https://doi.org:10.1136/gutjnl-2015-309618
42   Park, H., Yeo, S., Kang, S. & Huh, C. S. Longitudinal Microbiome Analysis in a
Dextran Sulfate Sodium-Induced Colitis Mouse Model. *Microorganisms* **9** (2021).
https://doi.org:10.3390/microorganisms9020370
43   Van de Guchte, M. *et al.* Alternative stable states in the intestinal ecosystem:
proof of concept in a rat model and a perspective of therapeutic implications.
Microbiome **8**, 153 (2020). https://doi.org:10.1186/s40168-020-00933-7

44   Berry, D. *et al.* Intestinal Microbiota Signatures Associated with Inflammation
History in Mice Experiencing Recurring Colitis. *Front Microbiol* **6**, 1408 (2015).
https://doi.org:10.3389/fmicb.2015.01408
45   Maharshak, N. *et al.* Altered enteric microbiota ecology in interleukin 10-deficient
mice during development and progression of intestinal inflammation. *Gut Microbes* **4**,
316-324 (2013). https://doi.org:10.4161/gmic.25486
46   Osaka, T. *et al.* Meta-Analysis of Fecal Microbiota and Metabolites in
Experimental Colitic Mice during the Inflammatory and Healing Phases. *Nutrients* **9**
(2017). https://doi.org:10.3390/nu9121329
47   Glymenaki, M. *et al.* Compositional Changes in the Gut Mucus Microbiota
Precede the Onset of Colitis-Induced Inflammation. *Inflammatory Bowel Diseases* **23**,
912-922 (2017). https://doi.org:10.1097/MIB.0000000000001118
48   Schwab, C. *et al.* Longitudinal study of murine microbiota activity and interactions
with the host during acute inflammation and recovery. *Isme j* **8**, 1101-1114 (2014).
https://doi.org:10.1038/ismej.2013.223
49   De Fazio, L. *et al.* Longitudinal analysis of inflammation and microbiota dynamics
in a model of mild chronic dextran sulfate sodium-induced colitis in mice. *World J*
Gastroenterol **20**, 2051-2061 (2014). https://doi.org:10.3748/wjg.v20.i8.2051
50   Smith, P. *et al.* Host Genetics and Environmental Factors Regulate Ecological
Succession of the Mouse Colon Tissue-Associated Microbiota. *PLOS ONE* **7**, e30273
(2012). https://doi.org:10.1371/journal.pone.0030273
51   Haange, S. B. *et al.* Disease Development Is Accompanied by Changes in
Bacterial Protein Abundance and Functions in a Refined Model of Dextran Sulfate
Sodium (DSS)-Induced Colitis. *J Proteome Res* **18**, 1774-1786 (2019).
https://doi.org:10.1021/acs.jproteome.8b00974
52   Munyaka, P. M., Rabbi, M. F., Khafipour, E. & Ghia, J. E. Acute dextran sulfate
sodium (DSS)-induced colitis promotes gut microbial dysbiosis in mice. *J Basic*
Microbiol **56**, 986-998 (2016). https://doi.org:10.1002/jobm.201500726
53   Sommer, F., Anderson, J. M., Bharti, R., Raes, J. & Rosenstiel, P. The resilience
of the intestinal microbiota influences health and disease. *Nature Reviews Microbiology*
15, 630-638 (2017). https://doi.org:10.1038/nrmicro.2017.58
54   Bao, C. *et al.* Acupuncture improves the symptoms, intestinal microbiota, and
inflammation of patients with mild to moderate Crohn's disease: A randomized
controlled trial. *eClinicalMedicine* **45** (2022).
https://doi.org:10.1016/j.eclinm.2022.101300
55   Eri, I., Masaya, Y. & Shigetada, K. Gut microbiota-mediated alleviation of dextran
sulfate sodium-induced colitis in mice. *bioRxiv*, 2023.2008.2007.552384 (2023).
https://doi.org:10.1101/2023.08.07.552384
56   Gao, Z. *et al.* Microbiota of Inflammatory Bowel Disease Models. *Annu Int Conf*
IEEE Eng Med Biol Soc **2018**, 2374-2377 (2018).
https://doi.org:10.1109/embc.2018.8512848
57   Hu, S. *et al.* Ameliorating Effects of Vitamin K2 on Dextran Sulfate Sodium-
Induced Ulcerative Colitis in Mice. *International Journal of Molecular Sciences* **24**
(2023).
58   Liu, Y. *et al.* β-patchoulene simultaneously ameliorated dextran sulfate sodium-
induced colitis and secondary liver injury in mice via suppressing colonic leakage and

flora imbalance. *Biochemical Pharmacology* **182**, 114260 (2020).
https://doi.org:https://doi.org/10.1016/j.bcp.2020.114260
59	Luo, S. *et al.* Emodin targeting the colonic metabolism via PPARγ alleviates UC
by inhibiting facultative anaerobe. *Phytomedicine* **104**, 154106 (2022).
https://doi.org:https://doi.org/10.1016/j.phymed.2022.154106
60	Pérez-Brocal, V. *et al.* Study of the viral and microbial communities associated
with Crohn's disease: a metagenomic approach. *Clin Transl Gastroenterol* **4**, e36
(2013). https://doi.org:10.1038/ctg.2013.9
61	Robinson, A. M. *et al.* Fecal Microbiota and Metabolome in a Mouse Model of
Spontaneous Chronic Colitis: Relevance to Human Inflammatory Bowel Disease.
Inflammatory Bowel Diseases **22**, 2767-2787 (2016).
https://doi.org:10.1097/MIB.0000000000000970
62	Sankarasubramanian, J., Ahmad, R., Avuthu, N., Singh, A. B. & Guda, C. Gut
Microbiota and Metabolic Specificity in Ulcerative Colitis and Crohn's Disease. *Front
Med (Lausanne)* **7**, 606298 (2020). https://doi.org:10.3389/fmed.2020.606298
63	Souza, E. L. S. *et al.* Beneficial effects resulting from oral administration of
Escherichia coli Nissle 1917 on a chronic colitis model. *Benef Microbes* **11**, 779-790
(2020). https://doi.org:10.3920/bm2020.0045
64	Wiredu Ocansey, D. K. *et al.* The diagnostic and prognostic potential of gut
bacteria in inflammatory bowel disease. *Gut Microbes* **15**, 2176118 (2023).
https://doi.org:10.1080/19490976.2023.2176118
65	Zuo, W. *et al.* 16S rRNA and metagenomic shotgun sequencing data revealed
consistent patterns of gut microbiome signature in pediatric ulcerative colitis. *Scientific
Reports* **12**, 6421 (2022). https://doi.org:10.1038/s41598-022-07995-7
66	Nayfach, S., Shi, Z. J., Seshadri, R., Pollard, K. S. & Kyrpides, N. C. New
insights from uncultivated genomes of the global human gut microbiome. *Nature* **568**,
505-510 (2019). https://doi.org:10.1038/s41586-019-1058-x
67	Wang, Y. *et al.* Phylogenomics of expanding uncultured environmental
Tenericutes provides insights into their pathogenicity and evolutionary relationship with
Bacilli. *BMC Genomics* **21**, 408 (2020). https://doi.org:10.1186/s12864-020-06807-4
68	Veseli, I. *et al.* Microbes with higher metabolic independence are enriched in
human gut microbiomes under stress. *bioRxiv* (2023).
https://doi.org:10.1101/2023.05.10.540289
69	Watson, A. R. *et al.* Metabolic independence drives gut microbial colonization
and resilience in health and disease. *Genome Biology* **24**, 78 (2023).
https://doi.org:10.1186/s13059-023-02924-x
70	Silamiķele, L. *et al.* Spatial variation of the gut microbiome in response to long-
term metformin treatment in high-fat diet-induced type 2 diabetes mouse model of both
sexes. *Gut Microbes* **15**, 2188663 (2023).
https://doi.org:10.1080/19490976.2023.2188663
71	Liu, Z., Peng, Y., Zhao, L. & Li, X. MFE40—the active fraction of Mume Fructus
alcohol extract—alleviates Crohn's disease and its complications. *Journal of
Ethnopharmacology* **296**, 115465 (2022).
https://doi.org:https://doi.org/10.1016/j.jep.2022.115465

72 Xu, H.-M. *et al.* Selection strategy of dextran sulfate sodium-induced acute or
chronic colitis mouse models based on gut microbial profile. *BMC Microbiology* **21**, 279
(2021). https://doi.org:10.1186/s12866-021-02342-8
73 Zhao, X. *et al.* Intervention with the crude polysaccharides of Physalis pubescens
L. mitigates colitis by preventing oxidative damage, aberrant immune responses, and
dysbacteriosis. *Journal of Food Science* **85**, 2596-2607 (2020).
https://doi.org:https://doi.org/10.1111/1750-3841.15330
74 Rui Zhang, Q. Z., Yanni Chen et al. Combined Rg1 and adipose-derived stem
cells alleviate DSS-induced colitis in a mouse model. *Research Square* (2021).
75 Zha, Z. *et al.* An orally administered butyrate-releasing xylan derivative reduces
inflammation in dextran sulphate sodium-induced murine colitis. *International Journal of*
Biological Macromolecules **156**, 1217-1233 (2020).
https://doi.org:https://doi.org/10.1016/j.ijbiomac.2019.11.159
76 Tian, X., Wang, G., Jin, K., Ding, Y. & Cheng, D. Rice hull insoluble dietary fiber
alleviated experimental colitis induced by low dose of dextran sulfate sodium in
cadmium-exposed mice. *Food Funct* **13**, 7215-7225 (2022).
https://doi.org:10.1039/d2fo00891b
77 Guo, X. *et al.* The protective effect of Schisandra chinensis (Turcz.) Baill.
polysaccharide on DSS-induced ulcerative colitis in mice via the modulation of gut
microbiota and inhibition of NF-κB activation. *Journal of the Science of Food and*
Agriculture **n/a** (2023). https://doi.org:https://doi.org/10.1002/jsfa.12905
78 Cheng, H. *et al.* Atractylodes macrocephala Koidz. volatile oil relieves acute
ulcerative colitis via regulating gut microbiota and gut microbiota metabolism. *Frontiers*
in Immunology **14** (2023).
79 Tian, M. *et al.* Barley Leaf Insoluble Dietary Fiber Alleviated Dextran Sulfate
Sodium-Induced Mice Colitis by Modulating Gut Microbiota. *Nutrients* **13** (2021).
https://doi.org:10.3390/nu13030846
80 Bai, J., Wang, B., Tan, X., Huang, L. & Xiong, S. Regulatory effect of lactulose on
intestinal flora and serum metabolites in colitis mice: In vitro and in vivo evaluation.
Food Chemistry: X **19**, 100821 (2023).
https://doi.org:https://doi.org/10.1016/j.fochx.2023.100821
81 Li, H. *et al.* Dealcoholized muscadine wine was partially effective in preventing
and treating dextran sulfate sodium-induced colitis and restoring gut dysbiosis in mice.
Food Funct **14**, 5994-6011 (2023). https://doi.org:10.1039/d3fo00047h
82 Wang, Y., Guo, Y., Chen, H., Wei, H. & Wan, C. Potential of Lactobacillus
plantarum ZDY2013 and Bifidobacterium bifidum WBIN03 in relieving colitis by gut
microbiota, immune, and anti-oxidative stress. *Canadian Journal of Microbiology* **64**,
327-337 (2018). https://doi.org:10.1139/cjm-2017-0716
83 Kansal, S. *et al.* The Microbiome in Paediatric Crohn's Disease—A Longitudinal,
Prospective, Single-Centre Study. *Journal of Crohn's and Colitis* **13**, 1044-1054
(2019). https://doi.org:10.1093/ecco-jcc/jjz016
84 Mao, B. *et al.* Blautia producta displays potential probiotic properties against
dextran sulfate sodium-induced colitis in mice. *Food Science and Human Wellness* **13**,
709-720 (2024). https://doi.org:10.26599/FSHW.2022.9250060
85 Liu, Y. *et al.* Sophora japonica flowers and their main phytochemical, rutin,
regulate chemically induced murine colitis in association with targeting the NF-κB

signaling pathway and gut microbiota. *Food Chemistry* **393**, 133395 (2022).
https://doi.org:https://doi.org/10.1016/j.foodchem.2022.133395
86 Min, Z., Xiaona, H., Aziz, T., Jian, Z. & Zhennai, Y. Exopolysaccharides from
Lactobacillus plantarum YW11 improve immune response and ameliorate inflammatory
bowel disease symptoms. *Acta Biochim Pol* **67**, 485-493 (2020).
https://doi.org:10.18388/abp.2020_5371
87 Li, Y. *et al.* Gegen Qinlian Decoction Alleviates Experimental Colitis and
Concurrent Lung Inflammation by Inhibiting the Recruitment of Inflammatory Myeloid
Cells and Restoring Microbial Balance. *J Inflamm Res* **15**, 1273-1291 (2022).
https://doi.org:10.2147/jir.S352706
88 Gu, W., Zhang, L., Han, T., Huang, H. & Chen, J. Dynamic Changes in Gut
Microbiome of Ulcerative Colitis: Initial Study from Animal Model. *Journal of*
Inflammation Research **15**, 2631-2647 (2022). https://doi.org:10.2147/JIR.S358807
89 Johansson, M. E. V. *et al.* Bacteria Penetrate the Inner Mucus Layer before
Inflammation in the Dextran Sulfate Colitis Model. *PLOS ONE* **5**, e12238 (2010).
https://doi.org:10.1371/journal.pone.0012238
90 Walter, J. Ecological Role of Lactobacilli in the Gastrointestinal Tract:
Implications for Fundamental and Biomedical Research. *Applied and Environmental*
Microbiology **74**, 4985-4996 (2008). https://doi.org:10.1128/AEM.00753-08
91 Abdugheni, R. *et al.* Metabolite profiling of human-originated Lachnospiraceae at
the strain level. *iMeta* **1**, e58 (2022). https://doi.org:https://doi.org/10.1002/imt2.58
92 Biddle, A., Stewart, L., Blanchard, J. & Leschine, S. Untangling the Genetic Basis
of Fibrolytic Specialization by Lachnospiraceae and Ruminococcaceae in Diverse Gut
Communities. *Diversity* **5**, 627-640 (2013).
93 Wei, H. *et al.* Diverse gut microbiota pattern between mild and severe cancer-
related fatigue in lung cancer patients treated with first-line chemotherapy: A pilot study.
Thorac Cancer **14**, 309-319 (2023). https://doi.org:10.1111/1759-7714.14765
94 Vacca, M. *et al.* The Controversial Role of Human Gut Lachnospiraceae.
Microorganisms **8** (2020). https://doi.org:10.3390/microorganisms8040573
95 Wang, T. *et al.* Astilbin from Smilax glabra Roxb. alleviates high-fat diet-induced
metabolic dysfunction. *Food Funct* **13**, 5023-5036 (2022).
https://doi.org:10.1039/d2fo00060a
96 Zhu, Y., Li, Y., Liu, M., Hu, X. & Zhu, H. Guizhi Fuling Wan, Chinese Herbal
Medicine, Ameliorates Insulin Sensitivity in PCOS Model Rats With Insulin Resistance
via Remodeling Intestinal Homeostasis. *Frontiers in Endocrinology* **11** (2020).
97 Qi, Y. *et al.* Effects of Schisandra chinensis polysaccharides on rats with
antibiotic-associated diarrhea. *International Journal of Biological Macromolecules* **124**,
627-634 (2019). https://doi.org:https://doi.org/10.1016/j.ijbiomac.2018.11.250
98 Su, L. *et al.* The Anti-colitis Effect of Schisandra chinensis Polysaccharide Is
Associated With the Regulation of the Composition and Metabolism of Gut Microbiota.
Frontiers in Cellular and Infection Microbiology **10** (2020).
99 Liu, C., Sun, C. & Cheng, Y. β-Glucan alleviates mice with ulcerative colitis
through interactions between gut microbes and amino acids metabolism. *Journal of the*
Science of Food and Agriculture **103**, 4006-4016 (2023).
https://doi.org:https://doi.org/10.1002/jsfa.12357

100   Meehan, C. J. & Beiko, R. G. A Phylogenomic View of Ecological Specialization
in the Lachnospiraceae, a Family of Digestive Tract-Associated Bacteria. *Genome*
Biology and Evolution **6**, 703-713 (2014). https://doi.org:10.1093/gbe/evu050
101   Belzer, C. & de Vos, W. M. Microbes inside—from diversity to function: the case
of Akkermansia. *The ISME Journal* **6**, 1449-1458 (2012).
https://doi.org:10.1038/ismej.2012.6
102   Berry, D. *et al.* Phylotype-level 16S rRNA analysis reveals new bacterial
indicators of health state in acute murine colitis. *Isme j* **6**, 2091-2106 (2012).
https://doi.org:10.1038/ismej.2012.39
103   Li, M., Wu, Y., Hu, Y., Zhao, L. & Zhang, C. Initial gut microbiota structure affects
sensitivity to DSS-induced colitis in a mouse model. *Science China Life Sciences* **61**,
762-769 (2018). https://doi.org:10.1007/s11427-017-9097-0
104   Nagalingam, N. A., Kao, J. Y. & Young, V. B. Microbial ecology of the murine gut
associated with the development of dextran sodium sulfate-induced colitis. *Inflamm*
Bowel Dis **17**, 917-926 (2011). https://doi.org:10.1002/ibd.21462
105   Xue, L. *et al.* The effects of live and pasteurized Akkermansia muciniphila on
DSS-induced ulcerative colitis, gut microbiota, and metabolomics in mice. *Food &*
Function **14**, 4632-4646 (2023). https://doi.org:10.1039/D2FO03493J
106   Liu, Q. *et al.* Akkermansia muciniphila Exerts Strain-Specific Effects on DSS-
Induced Ulcerative Colitis in Mice. *Frontiers in Cellular and Infection Microbiology* **11**
(2021).
107   Ansaldo, E. *et al.* Akkermansia muciniphila induces intestinal adaptive immune
responses during homeostasis. *Science* **364**, 1179-1184 (2019).
https://doi.org:10.1126/science.aaw7479
108   Sugihara, K. *et al.* Mucolytic bacteria license pathobionts to acquire host-derived
nutrients during dietary nutrient restriction. *Cell Reports* **40** (2022).
https://doi.org:10.1016/j.celrep.2022.111093
109   Ganesh, B. P., Klopfleisch, R., Loh, G. & Blaut, M. Commensal Akkermansia
muciniphila Exacerbates Gut Inflammation in Salmonella Typhimurium-Infected
Gnotobiotic Mice. *PLOS ONE* **8**, e74963 (2013).
https://doi.org:10.1371/journal.pone.0074963
110   Falk, P. G., Hooper, L. V., Midtvedt, T. & Gordon, J. I. Creating and maintaining
the gastrointestinal ecosystem: what we know and need to know from gnotobiology.
Microbiol Mol Biol Rev **62**, 1157-1170 (1998). https://doi.org:10.1128/mmbr.62.4.1157-
1170.1998
111   van Passel, M. W. J. *et al.* The Genome of Akkermansia muciniphila, a
Dedicated Intestinal Mucin Degrader, and Its Use in Exploring Intestinal Metagenomes.
PLOS ONE **6**, e16876 (2011). https://doi.org:10.1371/journal.pone.0016876
112   Ouwerkerk Janneke, P. *et al.* Adaptation of Akkermansia muciniphila to the Oxic-
Anoxic Interface of the Mucus Layer. *Applied and Environmental Microbiology* **82**, 6983-
6993 (2016). https://doi.org:10.1128/AEM.01641-16
113   Vidal-Lletjós, S. *et al.* Mucosal healing progression after acute colitis in mice.
World J Gastroenterol **25**, 3572-3589 (2019). https://doi.org:10.3748/wjg.v25.i27.3572
114   Chang, Z. Y., Liu, H. M., Leu, Y. L., Hsu, C. H. & Lee, T. Y. Modulation of Gut
Microbiota Combined with Upregulation of Intestinal Tight Junction Explains Anti-

Inflammatory Effect of Corylin on Colitis-Associated Cancer in Mice. *Int J Mol Sci* **23** (2022). https://doi.org:10.3390/ijms23052667

Li, R. *et al.* Integrated Analysis Reveals the Targets and Mechanisms in Immunosuppressive Effect of Mesalazine on Ulcerative Colitis. *Frontiers in Nutrition* **9** (2022).

Wang, W. *et al.* Swimming Impedes Intestinal Microbiota and Lipid Metabolites of Tumorigenesis in Colitis-Associated Cancer. *Frontiers in Oncology* **12** (2022).

Lawson, P. A., Saavedra Perez, L. & Sankaranarayanan, K. Reclassification of Clostridium cocleatum, Clostridium ramosum, Clostridium spiroforme and Clostridium saccharogumia as Thomasclavelia cocleata gen. nov., comb. nov., Thomasclavelia ramosa comb. nov., gen. nov., Thomasclavelia spiroformis comb. nov. and Thomasclavelia saccharogumia comb. nov. *International Journal of Systematic and Evolutionary Microbiology* **73** (2023). https://doi.org:https://doi.org/10.1099/ijsem.0.005694

Milosavljevic, M. N. *et al.* Antimicrobial treatment of Erysipelatoclostridium ramosum invasive infections: a systematic review. *Rev Inst Med Trop Sao Paulo* **63**, e30 (2021). https://doi.org:10.1590/S1678-9946202163030

Zakham, F. *et al.* Molecular diagnosis and enrichment culture identified a septic pseudoarthrosis due to an infection with Erysipelatoclostridium ramosum. *Int J Infect Dis* **81**, 167-169 (2019). https://doi.org:10.1016/j.ijid.2019.02.001

Qiu, Z. *et al.* Targeted Metagenome Based Analyses Show Gut Microbial Diversity of Inflammatory Bowel Disease patients. *Indian Journal of Microbiology* **57**, 307-315 (2017). https://doi.org:10.1007/s12088-017-0652-6

Nagayama, M. *et al.* TH1 cell-inducing Escherichia coli strain identified from the small intestinal mucosa of patients with Crohn's disease. *Gut Microbes* **12**, 1788898 (2020). https://doi.org:10.1080/19490976.2020.1788898

Sun, M. *et al.* Combined Signature of the Fecal Microbiome and Plasma Metabolome in Patients with Ulcerative Colitis. *Med Sci Monit* **25**, 3303-3315 (2019). https://doi.org:10.12659/msm.916009

Mancabelli, L. *et al.* Identification of universal gut microbial biomarkers of common human intestinal diseases by meta-analysis. *FEMS Microbiology Ecology* **93**, fix153 (2017). https://doi.org:10.1093/femsec/fix153

Cao, R.-R., He, P. & Lei, S.-F. Novel microbiota-related gene set enrichment analysis identified osteoporosis associated gut microbiota from autoimmune diseases. *Journal of Bone and Mineral Metabolism* **39**, 984-996 (2021). https://doi.org:10.1007/s00774-021-01247-w

Cai, S. *et al.* Gut Bacteria Erysipelatoclostridium and Its Related Metabolite Ptilosteroid A Could Predict Radiation-Induced Intestinal Injury. *Frontiers in Public Health* **10** (2022).

Farooq, P. D., Urrunaga, N. H., Tang, D. M. & von Rosenvinge, E. C. Pseudomembranous colitis. *Disease-a-Month* **61**, 181-206 (2015). https://doi.org:https://doi.org/10.1016/j.disamonth.2015.01.006

Liangen, L., Man, C., Hongcheng, Z., Bangzhou, Z. & Chuanxing, X. IDDF2023-ABS-0092 Multi-cohort retrospective microbiota analysis on fecal microbiota transplantation in ulcerative colitis. *Gut* **72**, A192 (2023). https://doi.org:10.1136/gutjnl-2023-IDDF.181

128 Feng, Y. *et al.* Barley Leaf Ameliorates Citrobacter rodentium-Induced Colitis
through Preventive Effects. *Nutrients* **14** (2022). https://doi.org:10.3390/nu14183833
129 Chai, M. *et al.* Different Bifidobacterium bifidum strains change the intestinal flora
composition of mice via different mechanisms to alleviate loperamide-induced
constipation. *Food & Function* **12**, 6058-6069 (2021).
https://doi.org:10.1039/D1FO00559F
130 Sullivan, S. *et al.* Downregulation of sodium transporters and NHERF proteins in
IBD patients and mouse colitis models: potential contributors to IBD-associated
diarrhea. *Inflamm Bowel Dis* **15**, 261-274 (2009). https://doi.org:10.1002/ibd.20743
131 Brook, I. & Long, S. S. in *Principles and Practice of Pediatric Infectious Diseases*
(Fifth Edition) (eds Sarah S. Long, Charles G. Prober, & Marc Fischer) 987-995.e982
(Elsevier, 2018).
132 Atarashi, K. *et al.* Treg induction by a rationally selected mixture of Clostridia
strains from the human microbiota. *Nature* **500**, 232-236 (2013).
https://doi.org:10.1038/nature12331
133 Narushima, S. *et al.* Characterization of the 17 strains of regulatory T cell-
inducing human-derived Clostridia. *Gut Microbes* **5**, 333-339 (2014).
https://doi.org:10.4161/gmic.28572
134 Vitalii, S., Musfiqur, S., Camilo, V., Kalai, M. & Giri, N. Novel Approach for
Microbiome Analysis Using Bacterial Replication Rates and Causal Inference with
Applications. *bioRxiv*, 2020.2005.2021.108514 (2020).
https://doi.org:10.1101/2020.05.21.108514
135 Li, S. *et al.* Frequency of change determines effectiveness of microbial response
strategies. *The ISME Journal* **17**, 2047-2057 (2023). https://doi.org:10.1038/s41396-
023-01515-9
136 Beller, Z. W. *et al.* Inducible CRISPR-targeted "knockdown" of human gut
Bacteroides in gnotobiotic mice discloses glycan utilization strategies. *Proceedings of*
the National Academy of Sciences **120**, e2311422120 (2023).
https://doi.org:10.1073/pnas.2311422120
137 Lloyd-Price, J. *et al.* Multi-omics of the gut microbial ecosystem in inflammatory
bowel diseases. *Nature* **569**, 655-662 (2019). https://doi.org:10.1038/s41586-019-1237-
9
138 Araki, Y., Mukaisyo, K., Sugihara, H., Fujiyama, Y. & Hattori, T. Increased
apoptosis and decreased proliferation of colonic epithelium in dextran sulfate sodium-
induced colitis in mice. *Oncol Rep* **24**, 869-874 (2010).
https://doi.org:10.3892/or.2010.869
139 Vetuschi, A., Latella, G., Sferra, R., Caprilli, R. & Gaudio, E. Increased
Proliferation and Apoptosis of Colonic Epithelial Cells in Dextran Sulfate Sodium-
Induced Colitis in Rats. *Digestive Diseases and Sciences* **47**, 1447-1457 (2002).
https://doi.org:10.1023/A:1015931128583
140 Raymond, F. *et al.* The initial state of the human gut microbiome determines its
reshaping by antibiotics. *Isme j* **10**, 707-720 (2016).
https://doi.org:10.1038/ismej.2015.148
141 Brugiroux, S. *et al.* Genome-guided design of a defined mouse microbiota that
confers colonization resistance against Salmonella enterica serovar Typhimurium.
Nature Microbiology **2**, 16215 (2016). https://doi.org:10.1038/nmicrobiol.2016.215

142 FitzGerald, J. *et al.* Improved gut microbiome recovery following drug therapy is
linked to abundance and replication of probiotic strains. *Gut Microbes* **14**, 2094664
(2022). https://doi.org:10.1080/19490976.2022.2094664
143 Robertson, S. J. *et al.* Comparison of Co-housing and Littermate Methods for
Microbiota Standardization in Mouse Models. *Cell Rep* **27**, 1910-1919 e1912 (2019).
https://doi.org:10.1016/j.celrep.2019.04.023
144 Russell, A. *et al.* Reduced housing density improves statistical power of murine
gut microbiota studies. *Cell Rep* **39**, 110783 (2022).
https://doi.org:10.1016/j.celrep.2022.110783
145 Singh, G., Brass, A., Cruickshank, S. M. & Knight, C. G. Cage and maternal
effects on the bacterial communities of the murine gut. *Sci Rep* **11**, 9841 (2021).
https://doi.org:10.1038/s41598-021-89185-5
146 McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible
interactive analysis and graphics of microbiome census data. *PLoS One* **8**, e61217
(2013). https://doi.org:10.1371/journal.pone.0061217
147 rbiom: Read/Write, Analyze, and Visualize 'BIOM' Data v. 2.0.0.9003 (2023).
148 pairwiseAdonis: Pairwise multilevel comparison using adonis v. 0.4 (2020).
149 FastQC: A Quality Control Tool for High Throughput Sequence Data (2010).
150 Ewels, P., Magnusson, M., Lundin, S. & Kaller, M. MultiQC: summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047-3048
(2016). https://doi.org:10.1093/bioinformatics/btw354
151 Li, D., Liu, C. M., Luo, R., Sadakane, K. & Lam, T. W. MEGAHIT: an ultra-fast
single-node solution for large and complex metagenomics assembly via succinct de
Bruijn graph. *Bioinformatics* **31**, 1674-1676 (2015).
https://doi.org:10.1093/bioinformatics/btv033
152 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat*
Methods **9**, 357-359 (2012). https://doi.org:10.1038/nmeth.1923
153 Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition.
Nat Methods **11**, 1144-1146 (2014). https://doi.org:10.1038/nmeth.3103
154 Wu, Y. W., Simmons, B. A. & Singer, S. W. MaxBin 2.0: an automated binning
algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**,
605-607 (2016). https://doi.org:10.1093/bioinformatics/btv638
155 Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and
efficient genome reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).
https://doi.org:10.7717/peerj.7359
156 Sieber, C. M. K. *et al.* Recovery of genomes from metagenomes via a
dereplication, aggregation and scoring strategy. *Nat Microbiol* **3**, 836-843 (2018).
https://doi.org:10.1038/s41564-018-0171-1
157 Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W.
CheckM: assessing the quality of microbial genomes recovered from isolates, single
cells, and metagenomes. *Genome Res* **25**, 1043-1055 (2015).
https://doi.org:10.1101/gr.186072.114
158 Bowers, R. M. *et al.* Minimum information about a single amplified genome
(MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and archaea.
Nat Biotechnol **35**, 725-731 (2017). https://doi.org:10.1038/nbt.3893

159   GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database
(2020).
160   Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**,
2068-2069 (2014). https://doi.org:10.1093/bioinformatics/btu153
161   Fitting Linear Mixed-Effects Models Using lme4 (2015).
162   Kuznetsova A., B. P. B., Christensen R.H.B. lmerTest Package: Tests in Linear
Mixed Effects Models. *Journal of Statistical Software* **82** (2017).
https://doi.org:https://doi.org/10.18637/jss.v082.i1310.18637/jss.v082.i13