Dextran sodium sulfate-induced colitis alters the proportion and composition of replicating gut bacteria

Eve T. Beauchemin[1], Claire Hunter[2], & Corinne F. Maurice[1,3]

[1]Department of Microbiology & Immunology, Faculty of Medicine and Health Sciences, McGill University, Montreal, Quebec, Canada

[2]Department of Public Health and Primary Care, School of Clinical Medicine, University of Cambridge, Cambridge, England

[3]McGill Centre for Microbiome Research, Montreal, Quebec, Canada

Corresponding author: corinne.maurice@mcgill.ca

Abstract

The bacteria living in the human gut are essential for host health. Though the composition and metabolism of these bacteria is well described in both healthy hosts and those with intestinal disease, less is known about the activity of the gut bacteria prior to, and during, disease development – especially regarding gut bacterial replication. Here, we use a recently developed single-cell technique alongside existing metagenomics-based tools to identify, track, and quantify the replicating gut bacteria and their replication dynamics in the dextran sodium sulfate mouse model of colitis. We show that the proportion of replicating gut bacteria decreases when mice have the highest levels of inflammation and returns to baseline levels as mice begin recovering. We additionally report significant alterations in the composition of the total replicating gut bacterial community during colitis development. On the taxa level, we observe significant changes in the abundance of taxa such as the mucus-degrading *Akkermansia muciniphila* and the poorly described *Erysipelatoclostridium* genus. We further demonstrate that many taxa exhibit variable replication rates during colitis,

including *A. muciniphila*. Lastly, we show that colitis development is positively correlated

with increases in the presence and abundance of bacteria predicted to be fast

replicators, suggesting that taxa with the potential to replicate quickly may have an

advantage during intestinal inflammation. These data support the need for additional

research using activity-based approaches to further characterize the gut bacterial

response to intestinal inflammation and its consequences for both the host and the gut

microbial community at large.

Importance

It is well known that the bacteria living inside the gut are important for human health.

Indeed, the type of bacteria which are present and their metabolism is different in

healthy people versus those with intestinal disease. However, less is known about how

these gut bacteria are replicating, especially as someone begins to develop intestinal

disease. This is especially important as it is thought that the active gut bacteria may be

more relevant to health. Here, we begin addressing this gap by using several

complementary approaches to characterize the replicating gut bacteria in a mouse

model of intestinal inflammation. We reveal which gut bacteria are replicating, and how

quickly, as mice develop and recover from inflammation. This work can serve as a

model for future research to identify how the active gut bacteria may be impacting

health, or why these particular bacteria tend to thrive during intestinal inflammation.

Introduction

Inflammatory bowel disease (IBD) is an umbrella term for several chronic inflammatory

intestinal diseases, including Crohn's disease (CD) and ulcerative colitis (UC).[1] The

incidence of IBD has been increasing in both high-income and low- and middle-income

countries over the past 10 years and shows no signs of slowing down.[2] Though the

exact etiology of IBD is unknown, evidence suggests that IBD results from a

combination of genetic defects in the immune system and environmental triggers, which

results in an aberrant host immune response to the gut microbiota.[3,4] Indeed, this

disease cannot occur without the presence of the gut bacteria.[5] As such, these bacteria

may play a crucial role in the manifestation and/or perpetuation of IBD.

Several studies have characterized the composition of the colonic or fecal bacteria in

patients with IBD versus those without. Patients with IBD generally exhibit decreased

bacterial diversity, increased abundances of aerotolerant and facultative anaerobic

bacteria (such as members of the *Enterobacteriaceae* family), and decreased

abundances of beneficial obligate anaerobes.[6] It remains unclear what roles the gut

bacteria play in IBD, and whether the observed diversity changes are a cause or

consequence of disease. One emerging hypothesis is that the immune dysregulation

apparent in IBD patients could initially alter the gut environment, favouring the survival

and growth of bacterial taxa with pro-inflammatory properties, which further contribute to

inflammation.[1,7,8] This self-perpetuating cycle is broken primarily through the use of

medication, such as corticosteroid drugs and 5-aminosalicylic acid, also known as

mesalamine.[7,9]

Studies which have evaluated changes in gut bacterial transcripts, proteins, and

metabolites have shown more consistent alterations than changes in gut bacterial

composition in IBD patients, supporting the notion that gut bacterial metabolism plays a

larger role in perpetuating inflammation than changes in community structure.[3,10] For

example, in IBD there is an increased abundance of bacterial genes encoding for, and

proteins involved in, oxidative stress and DNA replication;[11,12] increased levels of

bacterial proteins related to sulfur metabolism and hydrogen sulfide production;[12]

increased transcriptional activity of specific bacteria, including *Clostridia* species and the

mucin-degrader *Ruminococcus gnavus*;[6] and significant alterations in bacterial

metabolites, such as decreased levels of anti-inflammatory short-chain fatty acids like

butyrate, and increased levels of primary bile acids.[6]

One fundamental aspect of bacterial activity which remains understudied in the gut is

bacterial replication,[13] specifically in the context of inflammation. Pioneering work from

Olm et al. (2019) showed that bacterial replication rates were one of the best predictors

of the development of necrotizing enterocolitis (NEC), an inflammatory intestinal
disease in pre-term infants. Indeed, the authors reported a significant increase in gut
bacterial replication rates, especially members of the *Enterobacteriaceae* family,
preceding the development of NEC.[14] In the same year, Riglar et al. (2019) tested their
newly developed a synthetic gene oscillator, the repressilator 2.0, which reported on the
growth dynamics of a transformed *E. coli* strain in a mouse model of dextran sodium
sulfate (DSS)-induced colitis. The authors found that during active inflammation, there
was increased variability in the growth rates of the transformed *E. coli*.[15]

In a more recent publication, Joseph et al. (2022) used an updated version of their
replication rate measurement tool, termed Co-PTR, to identify taxa with altered
replication rates in healthy individuals versus those with IBD.[16] They found that gut
bacterial replication rates exhibited large interindividual variability but seemed relatively
stable within healthy individuals. They also reported that replication rates of specific
taxa were associated with CD (*Subdoligranulum* species) or UC (*Roseburia intestinalis*,
Ruminiclostridium species, and *Subdoligranum* species), whereas the relative
abundances of these same taxa were not associated with their respective disease.
Notably, they highlight that at least one other study also reported *R. intestinalis* as
having increased replication rates in individuals with CD and UC, emphasizing the
relevance of this taxon in IBD.

Collectively, these studies demonstrate how gut bacterial replication provides
information about gut bacterial activity that complements relative abundance
measurements, contributing to a better understanding of which taxa will thrive during
intestinal inflammation. This could allow us to characterize how gut bacteria contribute
to inflammatory conditions and shape the composition and function of the post-
inflammatory gut microbiota.

Commonly used tools to study bacterial replication in community contexts rely on whole
genome shotgun sequencing (WGS) data. Though these tools are informative and
quantitative, they can be prohibitively expensive to implement and are data-analysis

intensive.[17] As such, there is a need for more physiologically-based techniques,[18] which

are more affordable and easily implementable, to complement existing WGS-focused

techniques.

Most studies to date which evaluated gut bacterial activity or replication in IBD and other

inflammatory conditions were cross-sectional and did not focus on disease development

and progression. This is partly because it is challenging to identify and track individuals

who will later develop IBD, despite new efforts in this direction.[19-21] We hypothesize that

tracking gut bacterial replication dynamics during the development and recovery from

colitis will better inform us about the bacteria most relevant to intestinal inflammation.

Here, we use a combination of bacterial physiology- and bioinformatics-based

techniques to capture changes in the composition and replication dynamics of gut

bacteria during the development and recovery from chemically induced colitis in mice.

The dextran sodium sulfate (DSS) colitis model is one of the most widely used rodent

models of intestinal inflammation, reproducibly recapitulating the colonic damage and

mucosal recovery of UC,[22] despite its limitations.[22-27] Importantly, this model allows the

precise timing of colitis onset and development. We thus tracked the proportion and

identity of actively replicating gut bacterial cells, quantified their replication rates when

possible, and estimated the replication potential of the entire bacterial community over

time.

As mice develop colitis, there are decreases in the proportions of replicating gut

bacteria, which revert to baseline values as the mice recover. We also report significant

changes in the composition of the replicating community of gut bacteria, as well as

significant changes in the abundance of specific taxa, such as the mucus-degrader

Akkermansia muciniphila and the poorly described *Erysipelatoclostridium* genus. Whole

genome shotgun sequencing approaches reveal dynamic replication rates for many

taxa throughout colitis development, including *A. muciniphila*. Lastly, we observed that

the development of colitis coincided with the preferential retention in the gut of bacterial

genomes with faster predicted doubling times. Our data support the need for additional

functional approaches to better understand the gut bacterial dynamics underlying the
progression of intestinal inflammation and the resulting altered microbial community.

151 Results

152 DSS reproducibly causes colitis and changes in fecal bacterial
153 composition

We first confirmed that administration of 2% DSS for 5 days in the drinking water of the
mice in our facility led to reproducible colitis. To do so, we monitored weight loss, blood
in stool, and levels of fecal lipocalin-2 (LCN-2), a commonly used and sensitive marker
of intestinal inflammation,[23] before, during, and after administration of DSS (**Figure 1A;**
Supplementary Figure 1). We incorporated these three parameters into a modified
scoring metric of colitis as described by Kim et al. (2012) based on the disease activity
index (DAI) (**Supplementary Table 1, Supplementary Table 2**).[28] In this way, we
defined the different health states of the mice during colitis development: baseline, pre-
symptomatic, symptomatic, and recovery,[29] as detailed in the Methods.
163

A.

B.

167
168 **Figure 1.** *Experimental timeline and changes in gut bacterial community during DSS-colitis.*

169 *(A) Timeline for each mouse experiment. After a 2-week acclimatization period, the feces of four cages of mice (two*
170 *cages per sex, with 3 mice per cage) were collected over 7 days. Then, dextran sodium sulfate (DSS) was*

administered in the drinking water of the mice at a concentration of 2% (w/v) for 5 days, after which normal water was returned. Mouse feces continued to be collected throughout. Once DSS administration began, and thereafter, mouse health was monitored via measurements of blood in stool and lipocalin-2. Based on mouse health parameters, the phases of this study were split into four health states: baseline, pre-symptomatic, symptomatic, and recovery. (B) Taxonomic bar plots at the genus level of the fecal gut bacteria from each mouse during each health state (C) Beta diversity (Jaccard index) of the fecal gut bacterial communities from all cages, grouped per health state. D) Heat maps of standardized relative abundances of taxa from all cages of mice which were differentially abundant as compared to the baseline state. BA = Baseline; PS = Pre-symptomatic; SY = Symptomatic; RE = Recovery.

Amplicon sequencing of the 16S V4-5 region was conducted on bacterial DNA extracted
from mouse feces collected throughout the experiment, as described in the Methods.
Following this, QIIME2 was used to taxonomically identify the bacteria present in each
sample (**Figure 1B**).[30] The bacterial communities of these mice included the genera
Muribaculaceae (F1: 25.37% ± 9.36; F2: 34.20% ± 12.42; M1: 17.62% ± 9.30; M2:
17.93% ± 12.80) and *Bacteroides* (F1: 17.34% ± 7.50; F2: 14.08% ± 8.37; M1: 15.68%
± 11.89; M2: 12.04% ± 7.91) of the Bacteroidota phylum; along with *Dubosiella* (F1:
0.64% ± 1.11; F2: 10.15% ± 10.96; M1: 21.55% ± 15.63; M2: 26.41% ± 12.61),
Lachnospiraceae NK4136 group (F1: 7.58% ± 3.88; F2: 11.72% ± 10.09; M1: 5.01% ±
2.92; M2: 8.05% ± 5.10), and *Faecalibaculum* (F1: 24.01% ± 18.33; F2: 6.66 * 10^-3 %
± 2.40 * 10^-2; M1: 7.85% ± 1.72; M2: 7.09% ± 1.17) of the Bacillota phylum (**Figure
1B**).

When combining the data from all cages, we report no significant differences in the
alpha diversity between the different health states for any of the metrics we assessed
(**Supplementary Figure 2A**). Nevertheless, we saw a trend towards significance for the
Shannon index, supported by a significant Friedman's test ($p = 0.044$) but non-
significant post-hoc test (Wilcoxon signed-rank test; $p = 0.375$ for all comparisons)
(**Supplementary Table 3**).

At the beta diversity level, we report significant differences in the bacterial communities
between baseline and both the symptomatic and recovery states based on the Jaccard
index (PERMANOVA, $q = 0.039$; **Figure 1C; Supplementary Table 4**). Despite no
significant differences between the health states for the other calculated beta diversity
metrics (**Supplementary Table 4**), we observed the same pattern of deviation from
baseline during the symptomatic state and partial return to near-baseline values during
recovery (**Supplementary Figure 2B-D**).

When grouping the bacterial diversity data by sex, significant differences were observed
in the beta-diversity values between health states (**Supplementary Figure 3**). For
female mice (cages F1 and F2), this included significant differences in the unweighted

Unifrac distance between the baseline and symptomatic health states (PERMANOVA, q
= 0.027), and between the recovery and both the pre-symptomatic (PERMANOVA, q =
0.042) and the symptomatic (PERMANOVA, q = 0.027) health states (**Supplementary**
Table 5iv). There were no significant differences for the other beta diversity metrics
(**Supplementary Table 5i-iii**). For male mice (cages M1 and M2), there were significant
differences in the unweighted Unifrac distance (UU), Jaccard index (JC), and Bray-
Curtis dissimilarity (BC) between the baseline and symptomatic (PERMANOVA, UU: q =
0.02 ; JC: q = 0.036; BC: q = 0.039) and recovery (PERMANOVA , UU: q = 0.018; JC: q
= 0.012; BC: q = 0.033) health states, and between the recovery and pre-symptomatic
(PERMANOVA , UU: q = 0.0315; JC: q = 0.036; BC: q = 0.039) and symptomatic
(PERMANOVA , UU: q = 0.02; JC: q = 0.015; BC: q = 0.033) health states
(**Supplementary Table 6i, ii, iv**). There were no significant differences for the weighted
Unifrac distance (**Supplementary Table 6iii**). These results suggest possible sex-
specific differences in gut bacterial communities during the development of colitis.

We next sought to determine whether there were any changes in the abundance of
specific taxa during the progression of colitis, using ANCOM-BC2 to conduct differential
abundance analysis.[31] We compared the abundances of each taxon at a genus level
between baseline and the other health states (see Methods). We identified 3, 9, and 17
taxa as differentially abundant in the pre-symptomatic, symptomatic, and recovery
states, respectively (**Supplementary Figure 4**).

We noted a consistent increase in the abundance of *Erysipelatoclostridium* and a
Eubacterium fissicatena group member in all health states as compared to the baseline,
as well as the increased abundance of *Parasutterella* and *Faecalibaculum* during the
symptomatic and recovery states (**Figure 1D; Supplementary Figure 4**). A
Lactobacillus member consistently appeared at significantly higher abundances during
baseline relative to the pre-symptomatic and symptomatic health states; whereas UCG-
005 (from the *Oscillospiraceace* family), a *Ruminococcaceae* family member, a
Eubacterium coprostanoligenes group member, and RF39 (from the *Bacilli* class) all

had higher relative abundances during baseline when compared to the symptomatic
and recovery health states (**Figure 1D; Supplementary Figure 4**).

243 Colitis alters the replicating and whole communities of gut
244 bacteria

Next, we wanted to assess the proportions and identity of the replicating gut bacteria

during the development of, and recovery from, DSS colitis. As such, a second

independent mouse experiment with the same design was conducted, wherein the

recently optimized EdU-click technique was used to label newly synthesized bacterial

DNA.[13] EdU-click was combined with flow cytometry, as well as fluorescence-activated

cell sorting and 16S rRNA amplicon sequencing (FACSeq), to quantify the proportions

of actively replicating cells in fresh mouse feces and to taxonomically identify these

bacteria (see Methods and **Figure 2A**), respectively. The health status of the mice was

monitored and health states were determined as described earlier (**Supplementary**

Figure 5).

Figure 2. *Experimental workflow and changes in the proportion and composition of replication gut bacteria during DSS colitis.*

(A) Workflow for each mouse experiment. Whenever fecal pellets were collected, half were immediately used for EdU-click and FACSeq; the other half were frozen for later DNA extraction, whole genome shotgun sequencing, and subsequent analyses. (B-C) Median proportion of replicating (%EdU⁺) cells, represented as a line graph showing per-cage dynamics over time (B) and as boxplots, grouping all cages per health state (C). (D-E) Beta diversity (Jaccard indices) of the replicating (EdU⁺) (D) and whole community (Whole) (E) of bacterial cells for all cages, grouped by health state

The proportion of replicating fecal bacterial cells, denoted as %EdU$^+$ cells, changed as
the mice developed and recovered from DSS-induced colitis (**Figure 2B**). Specifically,
there was a notable but non-significant decrease in the median proportion of replicating
cells from 4.75% ± 3.73 during baseline to 2.33% ± 1.35 during the pre-symptomatic
state and 1.79% ± 0.94 during the symptomatic state, with a return to near-baseline
values during the recovery period (4.40% ± 1.89) (**Figure 2C**). Notably, the variability in
the median proportion of replicating cells between cages decreased during the pre-
symptomatic (1.35) and symptomatic (0.94) time periods as compared to the baseline
(3.73) and recovery periods (1.89) (**Supplementary Figure 6A,B**), demonstrating that
the proportion of replicating cells was most similar between cages during active colitis.
279
280 We next wanted to determine if certain taxa preferentially replicate as mice develop
281 colitis and recover. To accomplish this, we conducted 16S rRNA amplicon sequencing
282 on the V4-5 region from DNA extracted from the replicating cells (EdU$^+$) and the whole
283 community (Whole) of bacteria originating from the same fecal sample (**Figure 2A**).
284
285 When considering the alpha diversity metrics, there were no significant differences in
286 any metric between the different health states for either the EdU$^+$ fraction or the Whole
287 community (**Supplementary Figure 7; Supplementary Table 8**). The lack of
288 statistically significant differences is likely due to both inter-cage variability and low
289 numbers of cages (n = 4 cages).
290
We then compared the fecal bacterial communities of these mice across the different
health states using beta diversity metrics, for both the EdU$^+$ and Whole community of
sorted bacteria. We saw that there was grouping and separation between the different
health states in both fractions (**Figure 2D, E**). However, these differences were only
statistically significant with the Jaccard index for the EdU$^+$ fraction, between the
symptomatic state and both the baseline (PERMANOVA, q = 0.012) and the pre-
symptomatic (PERMANOVA, q = 0.012) health states (**Supplementary Table 9i**). As
with the alpha diversity measurements, the lack of other statistically significant
differences (**Supplementary Figure 8; Supplementary Table 9ii-iv**) is likely due to

both inter-cage variability as well as low numbers of cages (n = 4 cages). When the two

sexes were considered separately, only the Whole community of the male mice (cages

M1 and M2) showed a statistical difference in any alpha diversity metric, being the

weighted Unifrac distance, between the pre-symptomatic and recovery states

(PERMANOVA, q = 0.042) (**Supplementary Figure 9P; Supplementary Table 14iii**).

No other beta diversity metrics were significant (**Supplementary Figure 9A-O;**

Supplementary Table 11, Supplementary 12, Supplementary 13, Supplementary

14i, ii, iv)

We then sought to identify taxa which were differentially abundant in either the EdU$^+$ or

the Whole fraction in the different health states relative to the baseline. As before, we

first taxonomically identified the bacteria present in our samples using QIIME2 (**Figure**

3A, B).

In the EdU$^+$ fraction, the most abundant genera included *Erysipelatoclostridium* (mean

28.76% ± 26.65), *Muribaculaceae* (mean 15.12% ± 11.19), *Lactobacillus* (mean 8.33%

± 13.05), *Lachnospiraceae* NK4A136 group (mean 6.38% ± 11.73), and *Dubosiella*

(mean 9.23% ± 11.53) (**Figure 3A, B**).

A.

B.

DSS #2 - Whole

320
Figure 3. Taxonomy and differentially abundant taxa during DSS colitis.(A-B) Taxonomic bar plots of the
fecal gut bacteria for each cage during each health state for the replicating (EdU⁺) (A) and whole community (Whole)
(B) of bacterial cells. (C-D) Heat maps of the standardized abundance of taxa determined as differentially abundant
as compared to the baseline state for the replicating (EdU⁺) cells (C) and the whole community (Whole) (D). BA =
baseline; PS = pre-symptomatic; SY = symptomatic; RE = recovery.

In the Whole fraction, the most abundant genera included *Muribaculaceae* (mean
20.22% ± 14.59), *Dubosiella* (mean 36.09% ± 20.71), *Akkermansia* (mean 10.69% ±
11.58), *Turicibacter* (mean 5.50 % ± 7.40), and *Lachnospiraceae* NK4A136 group
(4.39% ± 6.38) (**Figure 3A, B**).

We then used ANCOM-BC2 to conduct differential abundance analysis at the genus
level. In the replicating (EdU$^+$) fraction, *Akkermansia* significantly increased its
abundance during the pre-symptomatic and symptomatic time points compared to the
baseline state, whereas in the Whole fraction, this taxon only increased during the
symptomatic state (**Supplementary Figure 10**). In contrast, *Erysipelatoclostridium* was
significantly increased in both the EdU$^+$ fraction and the Whole fraction in all health
states compared to the baseline state (**Supplementary Figure 10**). In the Whole
fraction during the symptomatic state, there were also transient increases in a
Eubacterium fissicatena group member and *Turicibacter*, and decreases in
Lachnoclostridum, *Blautia*, and *Roseburia*, with no corresponding changes in the EdU$^+$
fraction (**Supplementary Figure 10A-D**).

The changes in the proportion of replicating cells (EdU$^+$) of both *Akkermansia* and
Erysipelatoclostridium species mirrored their changes in total abundance (Whole),
suggesting the active and successful replication of these taxa, leading to increased total
biomass (**Figure 3C, D; Supplementary 10G**). This increase was transient for
Akkermansia but seemed to persist for the *Erysipelatoclostridium* taxon.

Altered replication dynamics of fecal bacteria during colitis development

As a complementary method to EdU-click, we also used whole genome shotgun (WGS)
sequencing to quantify the replication dynamics of fecal bacteria on a subset of
samples. Bacterial genomes were assembled from WGS data into metagenome-
assembled genomes (MAGs) as described in the Methods. WGS sequencing was
performed on fecal samples from mice in cages F2 and M2 (see **Supplementary Table**

15 for WGS statistics), as these mice had more severe signs of inflammation compared

to their biological replicate cages (**Supplementary Figure 5**).

The taxonomic composition and differentially abundant taxa of the MAGs, as well as

their beta diversity, are summarized in **Figure 4** and **Supplementary Figure 11**,

respectively. None of the beta diversity measurements showed significant differences in

the fecal bacterial composition (**Supplementary Figure 11**), likely due to the limited

number of genomes for which taxonomy could be determined (37 MAGs for cage F2; 43

MAGs for cage M2). Nevertheless, differential abundance analysis showed a significant

increase in the relative abundance of an *Akkermansia muciniphila* MAG, and a

significant decrease in the relative abundance of two MAGs from the *Lachnospiraceae*

family, a 14-2 species and a COE1 species, during all health states as compared to the

baseline state (**Figure 4C-E**). There was also a significant increase in the relative

abundance of a *Duncaniella* MAG during the symptomatic state (**Figure 4D**).

E.

375
Figure 4. *Relative abundance and differentially abundant taxa from whole genome shotgun sequencing*
(WGS) data.(A-B) Taxonomic bar plots at the genus level for the metagenome-assembled genomes. (C) Volcano
plots of metagenome-assembled genomes determined to be differentially abundant from the baseline. Note that
genomes 14-2 & COE1 are both from the Lachnospiraceae *family. BA = baseline; PS = pre-symptomatic; SY =*
symptomatic; RE = recovery.

Next, we wanted to quantify the replication dynamics of these MAGs using two distinct
methods: the peak-to-trough (PTR) method, which estimates instantaneous replication
rates,[32] and the codon usage bias (CUB) method, which estimates the maximal
replication potential or minimal doubling time (DT).[33-35]

To calculate the replication rates during the development of DSS colitis for as many
quality genomes as possible, we used the PTR-based tool GRiD[36], which works well
with medium quality MAGs at low coverage (**Supplementary Table 15**). GRiD is
specialized in using genomes with a coverage of as little as 0.2x and as many as 90
scaffolds/Mbp, and for genomes with at least 1x coverage, as little as 50% completion
and as much as 200 scaffolds/Mbp.[36] The other main PTR-based tools, iRep,[37]
DEMIC,[38] and CoPTR,[16] all have requirements or assumptions which are not
appropriate for the quality, coverage, or temporal nature of our data. The differences
between these tools have been thoroughly evaluated elsewhere and will not be covered
here.[16,35,39]

From cage F2, we were able to calculate PTR values using GRiD across all time points
for 32 MAGs (corresponding to 53% of all MAGs from cage F2) (**Supplementary Table
15, Supplementary Table 16**); from cage M2, we calculated PTR values across all time
points for 43 MAGs (corresponding to 65% of all MAGs from cage M2) (**Supplementary
Table 15, Supplementary Table 17**). Of these MAGs, we highlighted the replication
rate dynamics of the following four: *Akkermansia muciniphila* (**Figure 5A**), a *Duncaniella*
species (**Figure 5B**), *Bifidobacterium globosum* (**Figure 5C**), and a *Muribaculaceae*
species (**Figure 5D**). We selected these taxa based on whether they met all or most of
the following criteria: they were differentially abundant between the health states, had
the highest median replication rates during the symptomatic period, had replication rate
measurements for all experimental days, were one of the most abundant taxa overall or
during the symptomatic period, and had replication rate measurements estimated in
both mouse cages.

A. *Akkermansia muciniphila*

B. *Duncaniella* sp910589485

C. *Bifidobacterium globosum*

D.

Figure 5. *Replication dynamics of gut bacteria during DSS colitis. (A-D) Replication rates of representative taxa as calculated by GRiD, coloured by health state. (E) Correlation between health states and minimal doubling times (DTs) for both cages, as calculated by gRodon.*

431   These taxa appear to have cage-specific dynamics. For instance, in cage F2, the *A.*
432   *muciniphila* MAG had no correlation with the health status of the mice; whereas, in cage
433   M2, the *A. muciniphila* MAG had consistently higher replication rates once the mice
434   developed colitis relative to the baseline state (**Figure 5A**).
435
436   For the *Duncaniella* species, this MAG had its highest peak of replication rates during
437   the symptomatic period in cage F2; while the same MAG had the highest replication
438   rates during the baseline period relative to other periods in cage M2 (**Figure 5B**).
439
440   For *B. globosum*, this MAG had relatively steady replication rates in cage F2, except for
441   a rapid and transient decrease on the first day of the pre-symptomatic period. This
442   same MAG had decreased replication rates throughout the experiment starting after the
443   first baseline day in cage M2 (**Figure 5C**).
444
445   Finally, for the *Muribaculaceae* species, this MAG had its lowest replication rates during
446   the pre-symptomatic state in cage F2, whereas in cage M2 its replication rates were
447   highest during the baseline state and were consistently lower afterwards (**Figure 5D**).
448
449   We further determined no correlation between bacterial replication rates and the relative
450   abundance of their corresponding taxa in the whole community (Pearson r = -0.14756;
451   r^2 = 0.02177; **Supplementary Figure 12**).[16]
452
To glean more information about the replication abilities of the taxa we identified, we
next used gRodon, a CUB-based tool, to calculate the minimal doubling times (DTs) for
each MAG (**Supplementary Figure 13**).[34] This method provides information on the
replication potential of bacteria, as opposed to their instantaneous replication rates.
Previous studies have shown that measurements of replication potential using CUB-
based tools can more accurately predict the replication behaviour of certain taxa,
especially slower growing ones in complex communities.[39]
460

The median minimal DTs for all phyla in both cages was between 1.39 and 3.46 hours
(**Supplementary Figure 13A; Supplementary Table 16, Supplementary Table 17**),
supporting the notion that gut bacteria are fast replicators.[33] Within the Bacillota phylum,
there was a larger range (F2: 0.89-7.09 hrs; M2: 0.63-7.79 hrs) and more variability (F2:
median absolute deviation (MAD) 1.67 hrs; M2: MAD 1.48 hrs) in the minimal DTs as
compared to the other phyla with more than one genome present (F2: Actinomycetota
range: 0.98-2.33 hrs, MAD: 0.74 hrs; Bacteroidota range: 1.44-2.59 hrs, MAD: 0.66 hrs;
M2: Actinomycetota range: 1.04-2.72 hrs, MAD: 0.64 hrs; Bacteroidota range: 1.50-2.53
hrs, MAD: 0.55 hrs) (**Supplementary Table 18, Supplementary Table 19**). This range
and variability specifically occurred within the Clostridia class (F2: range: 0.89 -7.09 hrs;
MAD: 1.69 hrs; M2: range: 1.15-7.79 hrs; MAD: 1.40 hrs), which were comprised mostly
of genomes from the *Lachnospiraceae* family (F2: 15 genomes: range: 1.53-7.00 hrs;
MAD: 0.57 hrs; M2: 30 genomes: range: 1.64-7.79 hrs; MAD: 1.28 hrs)
(**Supplementary Figure 13B,C; Supplementary Table 18, Supplementary Table**
19).
476
To see whether there were any changes in the replication potential of all bacteria
detected over time, rather than individual taxa, we next used gRodon's metagenome
mode to calculate the minimal DTs of the whole community of gut bacteria during the
development of colitis.[35] There was a negative but non-significant (p > 0.05) slope when
plotting the minimal DTs of both cages of mice as a function of the health state, while
accounting for cage and day as random effects (see Methods) (**Figure 5E,**
Supplementary Figure 14; Supplementary Table 20). However, this trend was close
to significant (p = 0.0743) when the minimal DTs were plotted as a function of fecal
LCN-2 levels, a sensitive marker of colitis (**Supplementary Figure 15; Supplementary**
Table 21)[40]. Both data indicate that the colitic environment could select for bacteria with
the capacity to replicate quickly.
488
As fast replication could be a strategy for bacterial taxa to persist in a perturbed
environment,[41] we also sought to describe the link between MAGs with the potential to
replicate quickly (i.e., with low minimal DTs estimated by gRodon) and those with

increased replication rates and/or increased relative abundances during the
symptomatic period.

When combining information from all of the MAGs and both cages together, we report
no correlation between the minimal DT of a MAG and its median replication rate during
the symptomatic period (r = 0.049; **Supplementary Figure 16**). In contrast, there
seemed to be a weak negative correlation between the minimal DT of a MAG and its
median relative abundance during the symptomatic period (r = -0.329), suggesting that
bacteria with the potential to replicate quickly tend to be at higher abundances during
active colitis (**Supplementary Figure 17**). However, our analyses are limited by the rare
high-abundance bacterial taxa and the prevalent low-abundance taxa.

503 Discussion

This study sought to determine the replication dynamics of gut bacteria during the
development of, and recovery from, intestinal inflammation in a mouse model of
chemically induced colitis. We found that the development of colitis resulted in
decreased proportions of replicating gut bacterial cells, as well as significant changes in
the composition of the replicating gut bacterial community. At the genus level, we report
significantly increased abundances of replicating *Akkermansia* and
Erysipelatoclostridium taxa, which preceded increases in their relative abundances in
the community. Replication rates of *A. muciniphilia, Bifidobacterium globosum,* a
Muribaculaceae species*,* and a *Duncaniella* species also changed during colitis. Lastly,
we saw that the progression of colitis coincided with an overall decrease in the
predicted minimal doubling time of the whole bacterial community. Collectively, our data
suggest that the composition of the mouse fecal bacterial community shifts towards taxa
with the potential to replicate faster as intestinal inflammation progresses.

517 DSS colitis results in broad-level and taxon-specific changes in
gut bacterial communities

While the role of gut microbial communities in intestinal inflammation is increasingly
studied, most studies remain cross-sectional. Thus, the dynamic alterations of the gut

microbiota during colitis progression remain less characterized.[19,20,29,42-50] By monitoring

murine fecal microbial communities in DSS colitis, we aimed to determine some of

these underlying microbial dynamics of colitis. While we do not report significant

decreases in alpha-diversity, unlike other DSS studies[48,50-52], our beta-diversity analyses

suggest that these gut bacterial communities may have reached an alternative stable

state distinct from baseline.[29,43,48,49,53] These dynamics were especially marked in male

mice, possibly explaining the more reproducible and severe phenotype typically

observed for males in this colitis model.

Our longitudinal approach allowed us to determine that differentially abundant gut

bacterial taxa could be categorized into 4 broad response groups, according to their

dynamics during colitis progression (**Supplementary Figure 4**). Taxa in Group 1 –

which include RF39, *Oscillibacter,* a *Ruminococcaceae* member, and a *Clostridium*

vadin BB60 group member – had decreased abundances during the recovery period,

suggesting that they likely cannot survive a colitic environment. In contrast, Group 2

taxa, including *Erysipelatoclostridium* and a *Eubacterium fissicatena* group member,

had increased abundances during colitis development and onwards, suggesting that

these taxa expanded when the host environment was perturbed, and were able to

maintain their increased abundances even as the host recovered. Other response

groups include taxa which decreased temporarily below the limit of detection during

colitis but which returned during the recovery period (Group 3; *Lactobacillus*); and those

which only expanded during recovery, possibly taking over a niche made available by

the loss of other taxa (Group 4; *Lachnospiraceace* UCG-008).

The limited existing literature on the above highlighted taxa suggests some possible

explanations for their dynamics. For instance, RF39 of Group 1, which has been shown

to decrease in abundance in mouse models of colitis,[54-65] has also been described as

having a reduced genome and as being metabolically dependent on other bacteria for

some amino acids and vitamins.[66,67] Our data support the hypothesis that auxotrophic

bacteria such as RF39 are less likely to survive harsh colitic environments, as has been

suggested to occur in IBD patients.[68,69]

In contrast, Group 2 bacterial taxa *Erysipelatoclostridium* and *E. fissicatena* persistently

increase after colitis development, as reported elsewhere.[70-87] While the mechanisms

behind this increase are unclear, our data align with the literature while also contributing

a finer time scale regarding their changes in abundance during colitis.

At least one other DSS colitis study has seen the growth patterns of Group 3 member

Lactobacillus reported here.[88] One putative explanation for such behaviour is that, since

Lactobacillus species colonize the forestomach of mice, these taxa can be washed into

the cecum and colon – especially during the diarrhea commonly seen in DSS colitis.[22-

24,49,89] If a favourable niche in these distal intestinal locations becomes available during

a perturbation, *Lactobacillus* may be able to establish there; whereas under homeostatic

conditions, this would be less likely to occur.[90] Furthermore, similarly to the earlier

discussed taxon RF39, the metabolic dependence of most *Lactobacillus* species likely

precludes their ability to thrive in an inflamed environment.[90] After the inflammation,

however, it may be possible that *Lactobacillus* from the forestomach establish in the

colon.

For the Group 4 member *Lachnospiraceae* UCG-008, little is known about its

metabolism; or indeed, the metabolism of most *Lachnospiraceae* species,[91] despite

including many SCFA-producers and metabolizers of complex plant-derived

carbohydrates.[92] *Lachnospiraceae* UCG-008 is likely a butyrate producer, since the

Lachnospiraceae family is known to contain many butyrate-producing members.[91-94]

There is discrepancy between different studies as to whether *Lachnospiraceae*

members are increased or decreased in various intestinal perturbations, and whether

this is beneficial or detrimental to the host.[91,94-99] It has been suggested that the impact

of *Lachnospiraceae* members on the host is likely species or strain specific.[91,94,96] As

such, the overall role of *Lachnospiraceae* UCG-008 in colitis remains unclear. One

speculation as to how this taxon thrives during colitis recovery is that it could create

endospores.[100] Upon the return to a more favourable environment, such as when the

mice begin recovering from colitis, these endospores may allow *Lachnospiraceae* UCG-
008 to rapidly take advantage of newly opened niches in the gut.

Decreased proportions of replicating gut bacteria during colitis

To determine if the aforementioned patterns of altered abundances reflected changes in
gut bacterial replication, we quantified and taxonomically identified the replicating gut
bacteria using EdU-click in an independent study. We observed a decreased proportion
of replicating bacteria during the pre-symptomatic and symptomatic stages, suggesting
that the inflammatory environment is detrimental for the replication of most taxa.

Importantly, we noted that the replicating gut bacterial community during the
symptomatic state was markedly different from other health states. Narrowing down to a
per-taxon basis, we report significant differences in the abundance of *Akkermansia* and
Erysipelatoclostridium in both the replicating fraction and in the whole bacterial
community.

Since *Akkermansia* began increasing its number of replicating cells during the pre-
symptomatic state, and then subsequently increased its abundance in the whole fraction
during the symptomatic state, this suggests that its increased number of replicating cells
successfully led to an increase in its biomass. Once the mice began to recover,
however, the abundance of this taxon quickly decreased back to baseline values.

The impact of *Akkermansia* in DSS colitis is unclear, as it is associated with health in
humans and is depleted in individuals with IBD,[101] whereas it is commonly seen to
increase during DSS colitis in mice.[29,102-104] In addition, while some studies have found
that mice with higher levels of *Akkermansia* exhibit less severe colitis,[105] supporting a
protective role for this taxon, other studies have shown the reverse.[103] More recent
studies highlight that the impact of *Akkermansia* on host health could be strain
specific[106] or depend on the microbial and immunological status of the host.[107] As a
mucus degrader, *Akkermansia* could worsen inflammation,[108] rendering the host
intestinal epithelium more accessible to microbial translocation. Alternatively, its mucus

metabolism could stimulate increased mucus secretion by the host,[109] as some *Bacteroides* species have been shown to do.[110] An emerging hypothesis for the transient increase in *Akkermansia* during DSS colitis is that a DSS-dependent increase in mucus permeability[89] allows this taxon greater access to its preferred carbon source.[111] *Akkermansia* is also likely to be somewhat resistant to the increased oxidative environment of colitis, given that there are higher levels of oxygen present at the mucus layer where this taxon resides.[112] The combination of its resistance to oxidative stress and an increased access to its preferred energy source creates a perfect environment for *Akkermansia* to thrive, at least until host mucosal recovery begins.[113]

The *Erysipelatoclostridium* taxon was increased at all time points relative to the baseline state, in both the replicating and the whole bacterial community. Despite a limited number of studies noting the contrary[99,114-116], these results are in accordance with the behaviour of *Erysipelatoclostridium* reported in prior studies during intestinal disturbance. Indeed, this genus and one of its species, namely *Clostridium ramosum* (also known as *E. ramosum* or *Thomasclavelia ramosum*),[117-119] have been associated with intestinal inflammation or injury in humans,[46,48,73,120-127] increasing in abundance in mouse models of colitis[71-82] and infection[128], and being positively correlated with fecal water content (i.e., diarrhea), a common colitis symptom[2,22-24,129,130]. The high replication activity of *Erysipelatoclostridium* we report demonstrates that it is a particularly active taxon during colitis, resulting in a possibly outsized impact on the microbial environment and the host.

What allows *Erysipelatoclostridium* to benefit from perturbations can only be speculated, since its metabolism is not well described,[70] despite being a dominant gut Clostridial taxon.[131] The best described member of this genus is *C. ramosum*, which is reported to produce low amounts of acetate and possibly butyrate, and can significantly increase the number and function of anti-inflammatory Treg cells.[132,133] These capabilities point towards the potential anti-inflammatory role of this taxon, without explaining its continual increase in abundance in perturbed intestinal conditions.

As more thoroughly discussed by Beauchemin et al.,[13] not all bacterial taxa can be

labelled with EdU, and labeled bacteria likely do not equally incorporate EdU into their

DNA. As such, this study can only report on replicating taxa amenable to EdU-click, and

likely does not capture all replicating taxa. As one way to address this limitation, we also

used metagenomics-based bioinformatic tools to calculate bacterial replication

dynamics on a subset of our data, as discussed below.

650 Taxon-specific changes in gut bacterial replication rates and
651 increased abundances of fast replicating gut bacteria during colitis

Stool samples from ten non-consecutive sampling times between days 1-18 in one cage

of female mice (F2) and one cage of male mice (M2) underwent shotgun sequencing.

This allowed us to quantify the replication dynamics of several metagenome-assembled

genomes (MAGs) during colitis development, using both PTR-based and CUB-based

approaches. For the most part, the oscillatory replication rates for each taxon in each

cage went back to near-baseline values, with no clear link to the host health states –

with the exception of *A. muciniphila*. In agreement with our EdU-click and differential

abundance analyses, the *A. muciniphila* genome seen in both cages replicated most

rapidly once DSS was administered, after which its replication rates decreased back to

near baseline values as colitis developed.

We also observed cage-specific replication rate dynamics of the same gut taxa. For

instance, the *Duncaniella*, *Bifidobacterium globosum,* and *Muribaculaceae* CAG-485

genomes in cage M2 had decreased replication rates as colitis developed, whereas in

cage F2 these same taxa had variable replication rates. These differences suggest

individual-specific gut bacterial replication dynamics, as reported in previous human

cohorts[16], in addition to possible sex- and cage-specific responses to inflammation.

Since *Duncaniella, Muribaculaceae* CAG-485, and *B. globosum* are not as well

described as *A. muciniphila*, the possible explanations underlying their replication rate

patterns remain to be determined.

We additionally report a lack of correlation between gut bacterial replication rates and
their associated relative abundances, as noted by other studies[16,32,37,134]. Indeed,
relative abundance measurements account for the culmination of bacterial growth and
death, whereas replication rates only account for the former, emphasizing that both
approaches measure distinct aspects of the gut bacterial community.[16,32,37]

We next used an independent metric based on genome-specific CUB bias, using
gRodon, to determine possible links between gut bacterial replication and the response
of specific taxa during inflammation. When calculating the replication potential of the
whole gut bacterial community over time, we noted slightly lower minimal doubling times
(i.e., increased maximal replication rates) as colitis developed or as levels of LCN-2
increased. This suggests that faster replicating gut bacteria may be preferentially
selected for in an inflammatory environment.

These findings were further corroborated by the negative correlation between the
minimal doubling time (DT) of a MAG and its median relative abundance during the
symptomatic period, suggesting that taxa with the ability to replicate quickly are present
at higher abundances during colitis. This is in line with previous research which
suggests that faster replicating bacteria can adapt more quickly to changing
environments,[33,135,136] that gut bacteria during IBD flares may be replicating more than
during remission,[137] and that fast replication is one strategy gut bacteria can use to
maintain or increase their biomass when cell turnover and intestinal sloughing are high
[41] as in DSS-induced colitis.[22,23,138,139]

We further report a lack of correlation between gut bacterial minimal doubling times and
replication rates during the symptomatic period. This suggests that replication potential
and replication rates are also measurements of fundamentally different processes,
although this remains to be validated.

It should be noted that our WGS analysis has limitations. We could only assemble
approximately 35-36% of the sequences obtained into MAGs, of which only 53-65%

could be both taxonomically characterized and have their replication rates estimated
using GRiD (**Supplementary Table 15**). This can be partly attributed to the relatively
shallow sequencing (median 12,775,255 reads; range 8,448,754 – 19,756,090 reads),
resulting in the low coverage of many genomes for most time points. Despite this, we
believe that this study provides a foundational representation of the replication
dynamics of the most abundant gut taxa during DSS colitis development, and argue that
bacterial replication rates should be measured in ongoing microbiome workflows where
WGS data is acquired. We have created and shared on GitHub (see Data Accessibility)
our workflow for genome assembly and binning, as well as subsequent MAG
classification and bacterial replication measurements, to help researchers proceed with
such analyses.

Conclusions and Future Perspectives

Overall, this work demonstrates the utility of tracking various facets of gut bacterial
replication alongside measurements of cell abundance and taxonomy during the
development of intestinal inflammation. We provide further evidence for the relevance of
some taxa, such as *Akkermansia*, and describe new gut bacterial replication dynamics,
such as those for *Erysipelatoclostridium*, in a DSS mouse model of colitis.

These taxa should be further studied for their relevance to host health.[64] For instance,
determining the shared metabolic characteristics of gut bacteria which persist, and/or
thrive, in the inflamed gut could reveal how these taxa or their metabolism may be
linked to the development of an inflammatory environment. Such taxa or metabolic
activities could act as biomarkers, wherein changes in their presence could indicate the
onset of a flare before overt symptoms manifest. This information could further help
predict how a particular gut bacterial community will respond to inflammation,
depending on the composition and activity of the community members.[140]

Simplified *in vitro or in vivo* systems (e.g., artificial guts, gut-on-a-chip, or bacterial
consortia like the OMM12 community[141]) will be particularly useful, since the complexity
of the native gut microbiota and the host environment severely limit our ability to

disentangle the myriad microbe-microbe and host-microbe interactions occurring during
colitis development. Such approaches could, for example, allow for the direct
comparison of transcripts or metabolites produced by thriving and non-thriving gut taxa
in an inflammatory intestinal environment.

Evaluating gut bacterial replication dynamics provides insight into bacterial activity
which cannot be captured from abundance measurements alone. For instance,
quantifying bacterial replication rates could be beneficial when following colonization
dynamics in early life, to improve our understanding of how microbial activity is linked to
the known succession dynamics observed therein.[33,37] Characterizing bacterial
replication dynamics could also be useful when determining whether members of a
probiotic cocktail are actively replicating in the gut after administration,[142] or how the
introduction of a pathogen affects bacterial community replication patterns. On a
conceptual level, describing various aspects of gut bacterial replication could help
uncover the role of r- and K-selection in the gut. Such research will contribute to
increasing our knowledge in a variety of ecosystems on this fundamental, yet poorly
described, characteristic of complex microbial communities.

Methods

Animals

Equal numbers of 6–8-week-old specific-pathogen free (SPF) male and female wild-
type C57BL/6 mice (Jackson Laboratories) were kept at the Goodman Cancer
Research Center Animal Facility at McGill University, in accordance with the McGill
Ethics Research Board (animal ethics protocol MCGL-7999). For each experiment,
there were six mice of each sex, and mice were housed three per cage according to
sex, with a total of four cages per experiment.

Before administration of dextran sodium sulfate (DSS), mice were allowed to acclimate
to the animal facility for two weeks. On the third week, fecal pellets were collected from
the mice on three non-consecutive days. Fecal pellets were collected per cage by

putting all mice from a single cage into a new, sterile cage, empty of corncob bedding
and nesting material, with no access to food or water. Feces were collected soon after
defecation using ethanol-cleaned tweezers reserved for this purpose. Feces from mice
of the same cage were put into the same 1.5 mL microcentrifuge tube. This procedure
lasted no longer than one hour, after which mice were placed back into their respective
cages. Immediately after collection, the fecal pellets were either processed in an
anaerobic chamber or frozen at -80°C for later DNA extraction, as detailed below.
770
The week following acclimatization, mice were administered 2% DSS (molecular weight:
36-50 kDA; MP Biomedicals) in the drinking water for 5 days, and fecal pellets were
collected daily except on weekends. Afterwards, the mice were returned to the facility
water line, and fecal pellets continued to be collected daily, except on weekends, for
another 6 days, after which fecal pellets were collected twice weekly for an additional
two to three weeks.
777
Mouse health parameters, including body weight, blood in stool, and levels of fecal
lipocalin-2, were monitored throughout the experiment. Body weight was measured per
mouse daily except on weekends. Blood in stool was detected using the Hemoccult
Sensa kit (Beckman Coulter). Levels of fecal lipocalin-2 (LCN-2) were quantified using
the mouse lipocalin 2 DuoSet enzyme-linked immunosorbent assay (ELISA) (R&D).
Fecal samples were prepared for the ELISA using the method of Chassaing et al.
(2012)[40] and diluted 10-fold to 10,000-fold depending on the concentration of LCN-2 in
the sample. The ELISA was performed according to the manufacturer's instructions.
786
The health states of the mice were determined based on body weight, blood in stool,
and fecal lipocalin-2 levels, as detailed in **Supplementary Table 1**. Baseline days were
defined as those before DSS administration, during which the mice had no clinical
symptoms of colitis. Pre-symptomatic days were defined as those during DSS
administration where the mice had minimal symptoms of colitis, including minor weight
loss and increased levels of fecal LCN-2 up to 100 ng per gram of feces (ng/g).
Symptomatic days were defined as those during or after DSS administration where the

mice had weight loss of 1-10%, increased levels of fecal LCN-2 exceeding 100 ng/g,
and/or the presence of blood in their stool. Recovery days were defined as the days
after DSS administration cessation and during which the mice began to gain weight and
no longer had blood in their stool, though high levels (≥100 ng/g) of fecal LCN-2 still
remained (**Figure 1**).
799
We randomly selected fecal samples collected from each cage of mice, rather than
tracking each individual mouse, due to our need for large quantities of fecal samples to
conduct our experiments. The microbial composition of the mice in a single cage is
thought to be very similar due to mouse coprophagic behaviour,[143-145] and as such, fecal
samples taken on a per-cage basis are expected to be broadly representative of all the
mice in a single cage.
806
807 This experimental setup was conducted independently two times, as detailed in the
808 main text.

Isolation of bacteria from mouse feces, EdU-click, and flow cytometry

Freshly collected fecal samples from mice were transferred to an anaerobic chamber
(Coy Laboratory Products; 5% H_2, 20% CO_2, 75% N_2) within one hour of collection.
Isolation of bacteria from mouse feces, EdU labeling, EdU-click, flow cytometry, and cell
sorting were performed as described previously.[13] EdU-positive (EdU$^+$) and EdU-
negative (EdU$^-$) cells and the whole community were sorted based on their level of
fluorescence in the appropriate channel and quantified as described previously.[13]

DNA extraction and 16S rRNA amplicon sequencing and analysis

Bacterial DNA was extracted from sorted cells and mouse fecal pellets using the AllPrep
PowerFecal DNA/RNA kit (Qiagen) as previously described.[13] Extracted DNA
underwent library preparation and 16S rRNA gene sequencing of the V4-V5
hypervariable region with 515F/926R primers using the Illumina MiSeq PE250 system at

the Université de Québec à Montréal (UQAM) Center of Excellence in Research on

Orphan Diseases – Foundation Courtois (CERMO-FC) genomics platform.

Sequenced DNA was analyzed using QIIME2[30] version 2022.2 as previously

described.[13] Sequences were rarified to the depth of the sample with the fewest reads

before conducting alpha and beta diversity analyses. Taxonomic classification and

relative abundance measurements were conducted on the quality-controlled, filtered

sequences using QIIME2's machine-learning program trained on the 16S V4-V5 regions

from our data set, using the Silva 132 database on 99% operational taxonomic units.

Reads present in the sheath fluid but absent in the Whole community samples were

identified as contaminants and removed.

Further analysis was performed using phyloseq (version 1.44.0)[146] in R (version 4.3.0)

running in RStudio (2023.03.1). Beta diversity measurements were calculated on

weighted or unweighted UniFrac distances, Jaccard indices, or Bray-Curtis

dissimilarities using rbiom (version 1.0.3).[147] Repeated measures, multiple comparisons

PERMANOVAs were calculated with 999 permutations to test for significance using the

pairwiseAdonis package (version 0.4.1),[148] with multiple comparisons being corrected

using the Benjamini-Hochberg (BH) false discovery rate (FDR) correction. Differential

abundance analyses were performed on the combined data from all cages, using

ANCOM-BC2 (version 2.1.4)[31], and correcting for cage origin and the repeated

measures nature of the data, as described in *Statistical tests* and shown in the code

available on GitHub (see *Data availability*).

Whole genome shotgun sequencing and analysis

For fecal samples from two cages (F2 and M2, corresponding to a cage of female mice

and a cage of male mice, respectively) from the second experiment, DNA extracted

from the feces collected from these cages underwent library preparation and whole

genome shotgun (WGS) sequencing using the Illumina Miseq PE150 system at

SeqCenter (Pittsburgh, PA). The resulting bacterial genomes were assessed for their

quality using FastQC (version 0.11.9)[149] and the resulting FastQC files were visualized

using MultiQC (version 1.12).[150] Sequences were then co-assembled per cage across

all sampling days using Megahit (version 1.2.9).[151] Bowtie2 (version 2.4.4)[152] was used

to remove reads of host origin and to assess the extent to which assembled contigs

represented the original, quality-controlled reads. Binning of bacterial genomes from the

co-assemblies was performed using CONCOCT (version 1.1.0),[153] MaxBin2 (version

2.2.7),[154] and MetaBAT 2 (version 2.14),[155] followed by DASTool (version 1.1.4),[156] the

latter of which identifies the best bins from the three aforementioned binning tools. Bin

quality was assessed using CheckM (version 1.2.0).[157] Only bins meeting the minimum

MIMAGS standards of medium quality genomes[158] went on to be taxonomically

identified using GTDB-Tk (version 2.1.0).[159] The relative abundance of each bin was

calculated based on the length of each genome and its coverage (see code under *Data*

availability).

Quantification of replication rate dynamics

Bins meeting the minimum MIMAGS standards of medium quality (≥50% complete,

<10% contamination)[158] were assessed for their potential maximal replication rates

using gRodon2 (version 2.3.0).[35] The maximal potential replication rate for the entire

bacterial community each day was calculated from the prokka (version 1.14.6)[160] protein

annotation output of per-cage, per-day assemblies as specified in the gRodon2 vignette

(http://microbialgamut.com/gRodon-vignette), using metagenome mode V2 and with

n_le = 1000 and all other parameters as default.

Bins meeting the minimum standards for GRiD (>=50% completeness, <=90

scaffolds/Mbp, >=x0.2 coverage) were assessed with this tool using default parameters.

An index was created for each bin using Bowtie2, and each sample (i.e., each day) per

cage was mapped to each appropriate index (i.e., each bin) that came from that cage.

The resulting SAM files from mapping were used as input for GRiD, using the "grid

single" mode.

Correlations

To correlate gut bacterial replication rates with their relative abundances, we used the approach of Joseph TA et al. (2020).[16] Briefly, we first log2-transformed all relative abundance and replication rate (GRiD) values. Then, we standardized the log2-transformed relative abundance and GRiD values per taxon using the scale() function in R. The standardized, log2-transformed values for abundance and GRiD were then correlated using Pearson correlation. We proceeded with the same approach to correlate gut bacterial replication potentials with their corresponding median replication rates during the symptomatic period, and to correlate gut bacterial replication potential with their corresponding median relative abundances during the symptomatic period.

Statistical tests

Statistical testing was performed according to the non-parametric, repeated measures (i.e., longitudinal), and multiple comparisons nature of the study design as necessary, as described in the results and/or figure legends, where appropriate.

Comparisons of the proportion of EdU$^+$ cells (%EdU$^+$ cells) between health states were performed by first calculating the median %EdU$^+$ cells per cage per health state, as the median is more robust to outliers than the mean. Then, the median %EdU$^+$ cells per health state, considering all four cages in each health state, were compared to each other using the Friedman test, accounting for cage as a random intercept effect.

Comparisons of alpha diversity metric values (Shannon index, Simpson evenness, and observed features) between health states was performed by first calculating the median alpha diversity value per cage per health state. Then, the median alpha diversity values per health state, considering all four cages in each health state, were compared to each other using the Friedman test, accounting for cage as a random variable. Where appropriate, the Wilcoxon sign-rank post-hoc test was conducted, wherein the baseline group was compared to all of the other health states, with the p-value adjusted for multiple comparisons using the Bonferroni correction.

PERMANOVA comparisons of the beta diversity between different health states of mice
were conducted using pairwiseAdonis (version 0.4.1).[148] These comparisons accounted
for the repeated measures design of the study as described in the available code and
as first described here: https://thebiobucket.blogspot.com/2011/04/repeat-measure-
adonis-lately-i-had-to.html. Multiple comparisons were conducted using the
pairwiseAdonis package in R. When data from the various cages were compiled
together, cage origin was considered as a random intercept effect.
916
917   ANCOM-BC2 (version 2.1.4)[31] was used to detect differentially abundant bacteria
918   between health states as described in the main text and by the creators of the ANCOM-
919   BC2 package. All health states were compared to the baseline state, using Dunnett's
920   test. When data from all cages of mice were compiled together, cage origin was
921   considered as a random intercept effect.
922
To determine to what extent the health state of the host could explain the minimal
doubling time of the whole community of bacteria, we fit a linear mixed-effects model to
our data using the lme4 (version 1.1-34)[161] and lmerTest (version 3.1-3)[162] packages in
R, with health state and minimal doubling time as fixed effects, and cage and sampling
day as random intercept effects.

Data availability
Bacterial 16S rRNA gene sequencing data from all experimental arms are available in
the NCBI SRA database under project number PRJNA1040262 and PRJNA1040873,
respectively. Whole genome shotgun sequencing data from the second experiment is
available in the NCBI SRA database under project number PRJNA1040965. Code
related to the analysis of these data have been deposited in GitHub under the account
at https://github.com/evetb, repository DSS_Manuscript. Raw flow cytometry data have
been deposited in FigShare (https://figshare.com/projects/DSS_Manuscript/193004).

Declarations

Disclosure of interest

The authors report there are no competing interests to declare.

Authors' contributions

ETB designed, performed, analyzed, and interpreted experiments. CH aided with extracting DNA and supernatant for lipocalin-2 assays for the first mouse experiment. CFM helped develop the project, obtained funding, and helped design experiments and interpret the data. ETB and CFM edited the manuscript and approved the final draft.

Funding

This work was supported by a Frederick Banting and Charles Best Canada Graduate Scholarship-Master's (CGS M) from the Canadian Institutes of Health Research (CIHR), a Ferrings Pharmaceuticals Fellowship from the McGill Faculty of Medicine and Health Sciences, and a doctoral research grant from the Fonds de Recherche du Québec – Nature et Technologies (FRQNT) to ETB. This work was also supported by the Canada Research Chair program (grant 950-230748 X-242502), a CIHR transition grant, the Kenneth Rainin Foundation, and an Owens Catchpaugh IBD Graduate Research Award from the MUHC Foundation to CFM.

References

Caruso, R., Lo, B. C. & Núñez, G. Host-microbiota interactions in inflammatory bowel disease. *Nat Rev Immunol* **20**, 411-426 (2020). https://doi.org:10.1038/s41577-019-0268-7

Wang, R., Li, Z., Liu, S. & Zhang, D. Global, regional and national burden of inflammatory bowel disease in 204 countries and territories from 1990 to 2019: a systematic analysis based on the Global Burden of Disease Study 2019. *BMJ Open* **13**, e065186 (2023). https://doi.org:10.1136/bmjopen-2022-065186

Li, J., Butcher, J., Mack, D. & Stintzi, A. Functional impacts of the intestinal microbiome in the pathogenesis of inflammatory bowel disease. *Inflamm Bowel Dis* **21**, 139-153 (2015). https://doi.org:10.1097/mib.0000000000000215

Schirmer, M., Garner, A., Vlamakis, H. & Xavier, R. J. Microbial genes and pathways in inflammatory bowel disease. *Nat Rev Microbiol* **17**, 497-511 (2019). https://doi.org:10.1038/s41579-019-0213-6

Lane, E. R., Zisman, T. L. & Suskind, D. L. The microbiota in inflammatory bowel disease: current and therapeutic insights. *J Inflamm Res* **10**, 63-73 (2017). https://doi.org:10.2147/jir.S116088

Lloyd-Price, J. *et al.* Multi-omics of the gut microbial ecosystem in inflammatory bowel diseases. *Nature* **569**, 655-662 (2019). https://doi.org:10.1038/s41586-019-1237-9

Lee, J. Y., Tsolis, R. M. & Bäumler, A. J. The microbiome and gut homeostasis. *Science* **377**, eabp9960 (2022). https://doi.org:10.1126/science.abp9960

Pisani, A. *et al.* Dysbiosis in the Gut Microbiota in Patients with Inflammatory Bowel Disease during Remission. *Microbiol Spectr* **10**, e0061622 (2022). https://doi.org:10.1128/spectrum.00616-22

Ho, G. T., Cartwright, J. A., Thompson, E. J., Bain, C. C. & Rossi, A. G. Resolution of Inflammation and Gut Repair in IBD: Translational Steps Towards Complete Mucosal Healing. *Inflamm Bowel Dis* **26**, 1131-1143 (2020). https://doi.org:10.1093/ibd/izaa045

Schirmer, M. *et al.* Dynamics of metatranscription in the inflammatory bowel disease gut microbiome. *Nat Microbiol* **3**, 337-346 (2018). https://doi.org:10.1038/s41564-017-0089-z

Morgan, X. C. *et al.* Dysfunction of the intestinal microbiome in inflammatory bowel disease and treatment. *Genome Biol* **13**, R79 (2012). https://doi.org:10.1186/gb-2012-13-9-r79

Zhang, X. *et al.* Metaproteomics reveals associations between microbiome and intestinal extracellular vesicle proteins in pediatric inflammatory bowel disease. *Nat Commun* **9**, 2873 (2018). https://doi.org:10.1038/s41467-018-05357-4

Beauchemin, E. T., Hunter, C. & Maurice, C. F. Actively replicating gut bacteria identified by 5-ethynyl-2'-deoxyuridine (EdU) click chemistry and cell sorting. *Gut Microbes* **15**, 2180317 (2023). https://doi.org:10.1080/19490976.2023.2180317

Olm, M. R. *et al.* Necrotizing enterocolitis is preceded by increased gut bacterial replication, Klebsiella, and fimbriae-encoding bacteria. *Science Advances* **5**, eaax5727 https://doi.org:10.1126/sciadv.aax5727

15   Riglar, D. T. *et al.* Bacterial variability in the mammalian gut captured by a single-
cell synthetic oscillator. *Nat Commun* **10**, 4665 (2019). https://doi.org:10.1038/s41467-
019-12638-z
16   Joseph, T. A., Chlenski, P., Litman, A., Korem, T. & Pe'er, I. Accurate and robust
inference of microbial growth dynamics from metagenomic sequencing reveals
personalized growth rates. *Genome Res* **32**, 558-568 (2022).
https://doi.org:10.1101/gr.275533.121
17   Quince, C., Walker, A. W., Simpson, J. T., Loman, N. J. & Segata, N. Shotgun
metagenomics, from sampling to analysis. *Nat Biotechnol* **35**, 833-844 (2017).
https://doi.org:10.1038/nbt.3935
18   Koch, B. J. *et al.* Estimating taxon-specific population dynamics in diverse
microbial communities. *Ecosphere* **9**, e02090 (2018).
https://doi.org:https://doi.org/10.1002/ecs2.2090
19   Galipeau, H. J. *et al.* Novel Fecal Biomarkers That Precede Clinical Diagnosis of
Ulcerative Colitis. *Gastroenterology* **160**, 1532-1545 (2021).
https://doi.org:10.1053/j.gastro.2020.12.004
20   Raygoza Garay, J. A. *et al.* Gut Microbiome Composition is associated with
future onset of Crohn's Disease in Healthy First-Degree Relatives. *Gastroenterology*
(2023). https://doi.org:10.1053/j.gastro.2023.05.032
21   Sasson, G. *et al.* A223 THE GEM PROJECT: ASSOCIATIONS OF DIETARY
PATTERNS WITH MICROBIOME AND FECAL CALPROTECTIN IN HEALTHY FIRST-
DEGREE RELATIVES OF CROHN'S DISEASE PATIENTS. *Journal of the Canadian
Association of Gastroenterology* **3**, 98-99 (2020).
https://doi.org:10.1093/jcag/gwz047.222
22   Eichele, D. D. & Kharbanda, K. K. Dextran sodium sulfate colitis murine model:
An indispensable tool for advancing our understanding of inflammatory bowel diseases
pathogenesis. *World J Gastroenterol* **23**, 6016-6029 (2017).
https://doi.org:10.3748/wjg.v23.i33.6016
23   Chassaing, B., Aitken, J. D., Malleshappa, M. & Vijay-Kumar, M. Dextran sulfate
sodium (DSS)-induced colitis in mice. *Curr Protoc Immunol* **104**, 15.25.11-15.25.14
(2014). https://doi.org:10.1002/0471142735.im1525s104
24   Perše, M. & Cerar, A. Dextran sodium sulphate colitis mouse model: traps and
tricks. *J Biomed Biotechnol* **2012**, 718617 (2012). https://doi.org:10.1155/2012/718617
25   Rogler, G. *et al.* Diet and Inflammatory Bowel Disease: What Quality Standards
Should Be Applied in Clinical and Laboratory Studies? *Molecular Nutrition & Food
Research* **65**, 2000514 (2021). https://doi.org:https://doi.org/10.1002/mnfr.202000514
26   Solomon, L. *et al.* The dextran sulphate sodium (DSS) model of colitis: an
overview. *Comparative Clinical Pathology* **19**, 235-239 (2010).
https://doi.org:10.1007/s00580-010-0979-4
27   Wirtz, S. *et al.* Chemically induced mouse models of acute and chronic intestinal
inflammation. *Nature Protocols* **12**, 1295-1309 (2017).
https://doi.org:10.1038/nprot.2017.044
28   Kim, J. J., Shajib, M. S., Manocha, M. M. & Khan, W. I. Investigating intestinal
inflammation in DSS-induced model of IBD. *J Vis Exp* (2012).
https://doi.org:10.3791/3678

29   Taguer, M. *et al.* Changes in Gut Bacterial Translation Occur before Symptom
Onset and Dysbiosis in Dextran Sodium Sulfate-Induced Murine Colitis. *mSystems* **6**,
e0050721 (2021). https://doi.org:10.1128/mSystems.00507-21
30   Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome
data science using QIIME 2. *Nat Biotechnol* **37**, 852-857 (2019).
https://doi.org:10.1038/s41587-019-0209-9
31   Lin, H. & Peddada, S. D. Analysis of compositions of microbiomes with bias
correction. *Nat Commun* **11**, 3514 (2020). https://doi.org:10.1038/s41467-020-17041-7
32   Korem, T. *et al.* Growth dynamics of gut microbiota in health and disease inferred
from single metagenomic samples. *Science* **349**, 1101-1106 (2015).
https://doi.org:10.1126/science.aac4812
33   Vieira-Silva, S. & Rocha, E. P. The systemic imprint of growth and its uses in
ecological (meta)genomics. *PLoS Genet* **6**, e1000808 (2010).
https://doi.org:10.1371/journal.pgen.1000808
34   Weissman, J. L., Hou, S. & Fuhrman, J. A. Estimating maximal microbial growth
rates from cultures, metagenomes, and single cells via codon usage patterns. *Proc Natl*
Acad Sci U S A **118** (2021). https://doi.org:10.1073/pnas.2016810118
35   Weissman, J. L., Peras, M., Barnum, T. P. & Fuhrman, J. A. Benchmarking
Community-Wide Estimates of Growth Potential from Metagenomes Using Codon
Usage Statistics. *mSystems* **7**, e0074522 (2022).
https://doi.org:10.1128/msystems.00745-22
36   Emiola, A. & Oh, J. High throughput in situ metagenomic measurement of
bacterial replication at ultra-low sequencing coverage. *Nat Commun* **9**, 4956 (2018).
https://doi.org:10.1038/s41467-018-07240-8
37   Brown, C. T., Olm, M. R., Thomas, B. C. & Banfield, J. F. Measurement of
bacterial replication rates in microbial communities. *Nat Biotechnol* **34**, 1256-1263
(2016). https://doi.org:10.1038/nbt.3704
38   Gao, Y. & Li, H. Quantifying and comparing bacterial growth dynamics in multiple
metagenomic samples. *Nat Methods* **15**, 1041-1044 (2018).
https://doi.org:10.1038/s41592-018-0182-0
39   Long, A. M., Hou, S., Ignacio-Espinoza, J. C. & Fuhrman, J. A. Benchmarking
microbial growth rate predictions from metagenomes. *ISME J* **15**, 183-195 (2021).
https://doi.org:10.1038/s41396-020-00773-1
40   Chassaing, B. *et al.* Fecal lipocalin 2, a sensitive and broadly dynamic non-
invasive biomarker for intestinal inflammation. *PLoS One* **7**, e44328 (2012).
https://doi.org:10.1371/journal.pone.0044328
41   Vandeputte, D. *et al.* Stool consistency is strongly associated with gut microbiota
richness and composition, enterotypes and bacterial growth rates. *Gut* **65**, 57-62 (2016).
https://doi.org:10.1136/gutjnl-2015-309618
42   Park, H., Yeo, S., Kang, S. & Huh, C. S. Longitudinal Microbiome Analysis in a
Dextran Sulfate Sodium-Induced Colitis Mouse Model. *Microorganisms* **9** (2021).
https://doi.org:10.3390/microorganisms9020370
43   Van de Guchte, M. *et al.* Alternative stable states in the intestinal ecosystem:
proof of concept in a rat model and a perspective of therapeutic implications.
Microbiome **8**, 153 (2020). https://doi.org:10.1186/s40168-020-00933-7

44   Berry, D. *et al.* Intestinal Microbiota Signatures Associated with Inflammation
History in Mice Experiencing Recurring Colitis. *Front Microbiol* **6**, 1408 (2015).
https://doi.org:10.3389/fmicb.2015.01408
45   Maharshak, N. *et al.* Altered enteric microbiota ecology in interleukin 10-deficient
mice during development and progression of intestinal inflammation. *Gut Microbes* **4**,
316-324 (2013). https://doi.org:10.4161/gmic.25486
46   Osaka, T. *et al.* Meta-Analysis of Fecal Microbiota and Metabolites in
Experimental Colitic Mice during the Inflammatory and Healing Phases. *Nutrients* **9**
(2017). https://doi.org:10.3390/nu9121329
47   Glymenaki, M. *et al.* Compositional Changes in the Gut Mucus Microbiota
Precede the Onset of Colitis-Induced Inflammation. *Inflammatory Bowel Diseases* **23**,
912-922 (2017). https://doi.org:10.1097/MIB.0000000000001118
48   Schwab, C. *et al.* Longitudinal study of murine microbiota activity and interactions
with the host during acute inflammation and recovery. *Isme j* **8**, 1101-1114 (2014).
https://doi.org:10.1038/ismej.2013.223
49   De Fazio, L. *et al.* Longitudinal analysis of inflammation and microbiota dynamics
in a model of mild chronic dextran sulfate sodium-induced colitis in mice. *World J*
Gastroenterol **20**, 2051-2061 (2014). https://doi.org:10.3748/wjg.v20.i8.2051
50   Smith, P. *et al.* Host Genetics and Environmental Factors Regulate Ecological
Succession of the Mouse Colon Tissue-Associated Microbiota. *PLOS ONE* **7**, e30273
(2012). https://doi.org:10.1371/journal.pone.0030273
51   Haange, S. B. *et al.* Disease Development Is Accompanied by Changes in
Bacterial Protein Abundance and Functions in a Refined Model of Dextran Sulfate
Sodium (DSS)-Induced Colitis. *J Proteome Res* **18**, 1774-1786 (2019).
https://doi.org:10.1021/acs.jproteome.8b00974
52   Munyaka, P. M., Rabbi, M. F., Khafipour, E. & Ghia, J. E. Acute dextran sulfate
sodium (DSS)-induced colitis promotes gut microbial dysbiosis in mice. *J Basic*
Microbiol **56**, 986-998 (2016). https://doi.org:10.1002/jobm.201500726
53   Sommer, F., Anderson, J. M., Bharti, R., Raes, J. & Rosenstiel, P. The resilience
of the intestinal microbiota influences health and disease. *Nature Reviews Microbiology*
15, 630-638 (2017). https://doi.org:10.1038/nrmicro.2017.58
54   Bao, C. *et al.* Acupuncture improves the symptoms, intestinal microbiota, and
inflammation of patients with mild to moderate Crohn's disease: A randomized
controlled trial. *eClinicalMedicine* **45** (2022).
https://doi.org:10.1016/j.eclinm.2022.101300
55   Eri, I., Masaya, Y. & Shigetada, K. Gut microbiota-mediated alleviation of dextran
sulfate sodium-induced colitis in mice. *bioRxiv*, 2023.2008.2007.552384 (2023).
https://doi.org:10.1101/2023.08.07.552384
56   Gao, Z. *et al.* Microbiota of Inflammatory Bowel Disease Models. *Annu Int Conf*
IEEE Eng Med Biol Soc **2018**, 2374-2377 (2018).
https://doi.org:10.1109/embc.2018.8512848
57   Hu, S. *et al.* Ameliorating Effects of Vitamin K2 on Dextran Sulfate Sodium-
Induced Ulcerative Colitis in Mice. *International Journal of Molecular Sciences* **24**
(2023).
58   Liu, Y. *et al.* β-patchoulene simultaneously ameliorated dextran sulfate sodium-
induced colitis and secondary liver injury in mice via suppressing colonic leakage and

flora imbalance. *Biochemical Pharmacology* **182**, 114260 (2020).
https://doi.org:https://doi.org/10.1016/j.bcp.2020.114260
59   Luo, S. *et al.* Emodin targeting the colonic metabolism via PPARγ alleviates UC
by inhibiting facultative anaerobe. *Phytomedicine* **104**, 154106 (2022).
https://doi.org:https://doi.org/10.1016/j.phymed.2022.154106
60   Pérez-Brocal, V. *et al.* Study of the viral and microbial communities associated
with Crohn's disease: a metagenomic approach. *Clin Transl Gastroenterol* **4**, e36
(2013). https://doi.org:10.1038/ctg.2013.9
61   Robinson, A. M. *et al.* Fecal Microbiota and Metabolome in a Mouse Model of
Spontaneous Chronic Colitis: Relevance to Human Inflammatory Bowel Disease.
Inflammatory Bowel Diseases **22**, 2767-2787 (2016).
https://doi.org:10.1097/MIB.0000000000000970
62   Sankarasubramanian, J., Ahmad, R., Avuthu, N., Singh, A. B. & Guda, C. Gut
Microbiota and Metabolic Specificity in Ulcerative Colitis and Crohn's Disease. *Front
Med (Lausanne)* **7**, 606298 (2020). https://doi.org:10.3389/fmed.2020.606298
63   Souza, E. L. S. *et al.* Beneficial effects resulting from oral administration of
Escherichia coli Nissle 1917 on a chronic colitis model. *Benef Microbes* **11**, 779-790
(2020). https://doi.org:10.3920/bm2020.0045
64   Wiredu Ocansey, D. K. *et al.* The diagnostic and prognostic potential of gut
bacteria in inflammatory bowel disease. *Gut Microbes* **15**, 2176118 (2023).
https://doi.org:10.1080/19490976.2023.2176118
65   Zuo, W. *et al.* 16S rRNA and metagenomic shotgun sequencing data revealed
consistent patterns of gut microbiome signature in pediatric ulcerative colitis. *Scientific
Reports* **12**, 6421 (2022). https://doi.org:10.1038/s41598-022-07995-7
66   Nayfach, S., Shi, Z. J., Seshadri, R., Pollard, K. S. & Kyrpides, N. C. New
insights from uncultivated genomes of the global human gut microbiome. *Nature* **568**,
505-510 (2019). https://doi.org:10.1038/s41586-019-1058-x
67   Wang, Y. *et al.* Phylogenomics of expanding uncultured environmental
Tenericutes provides insights into their pathogenicity and evolutionary relationship with
Bacilli. *BMC Genomics* **21**, 408 (2020). https://doi.org:10.1186/s12864-020-06807-4
68   Veseli, I. *et al.* Microbes with higher metabolic independence are enriched in
human gut microbiomes under stress. *bioRxiv* (2023).
https://doi.org:10.1101/2023.05.10.540289
69   Watson, A. R. *et al.* Metabolic independence drives gut microbial colonization
and resilience in health and disease. *Genome Biology* **24**, 78 (2023).
https://doi.org:10.1186/s13059-023-02924-x
70   Silamiķele, L. *et al.* Spatial variation of the gut microbiome in response to long-
term metformin treatment in high-fat diet-induced type 2 diabetes mouse model of both
sexes. *Gut Microbes* **15**, 2188663 (2023).
https://doi.org:10.1080/19490976.2023.2188663
71   Liu, Z., Peng, Y., Zhao, L. & Li, X. MFE40—the active fraction of Mume Fructus
alcohol extract—alleviates Crohn's disease and its complications. *Journal of
Ethnopharmacology* **296**, 115465 (2022).
https://doi.org:https://doi.org/10.1016/j.jep.2022.115465

72   Xu, H.-M. *et al.* Selection strategy of dextran sulfate sodium-induced acute or
chronic colitis mouse models based on gut microbial profile. *BMC Microbiology* **21**, 279
(2021). https://doi.org:10.1186/s12866-021-02342-8
73   Zhao, X. *et al.* Intervention with the crude polysaccharides of Physalis pubescens
L. mitigates colitis by preventing oxidative damage, aberrant immune responses, and
dysbacteriosis. *Journal of Food Science* **85**, 2596-2607 (2020).
https://doi.org:https://doi.org/10.1111/1750-3841.15330
74   Rui Zhang, Q. Z., Yanni Chen et al. Combined Rg1 and adipose-derived stem
cells alleviate DSS-induced colitis in a mouse model. *Research Square* (2021).
75   Zha, Z. *et al.* An orally administered butyrate-releasing xylan derivative reduces
inflammation in dextran sulphate sodium-induced murine colitis. *International Journal of*
Biological Macromolecules **156**, 1217-1233 (2020).
https://doi.org:https://doi.org/10.1016/j.ijbiomac.2019.11.159
76   Tian, X., Wang, G., Jin, K., Ding, Y. & Cheng, D. Rice hull insoluble dietary fiber
alleviated experimental colitis induced by low dose of dextran sulfate sodium in
cadmium-exposed mice. *Food Funct* **13**, 7215-7225 (2022).
https://doi.org:10.1039/d2fo00891b
77   Guo, X. *et al.* The protective effect of Schisandra chinensis (Turcz.) Baill.
polysaccharide on DSS-induced ulcerative colitis in mice via the modulation of gut
microbiota and inhibition of NF-κB activation. *Journal of the Science of Food and*
Agriculture **n/a** (2023). https://doi.org:https://doi.org/10.1002/jsfa.12905
78   Cheng, H. *et al.* Atractylodes macrocephala Koidz. volatile oil relieves acute
ulcerative colitis via regulating gut microbiota and gut microbiota metabolism. *Frontiers*
in Immunology **14** (2023).
79   Tian, M. *et al.* Barley Leaf Insoluble Dietary Fiber Alleviated Dextran Sulfate
Sodium-Induced Mice Colitis by Modulating Gut Microbiota. *Nutrients* **13** (2021).
https://doi.org:10.3390/nu13030846
80   Bai, J., Wang, B., Tan, X., Huang, L. & Xiong, S. Regulatory effect of lactulose on
intestinal flora and serum metabolites in colitis mice: In vitro and in vivo evaluation.
Food Chemistry: X **19**, 100821 (2023).
https://doi.org:https://doi.org/10.1016/j.fochx.2023.100821
81   Li, H. *et al.* Dealcoholized muscadine wine was partially effective in preventing
and treating dextran sulfate sodium-induced colitis and restoring gut dysbiosis in mice.
Food Funct **14**, 5994-6011 (2023). https://doi.org:10.1039/d3fo00047h
82   Wang, Y., Guo, Y., Chen, H., Wei, H. & Wan, C. Potential of Lactobacillus
plantarum ZDY2013 and Bifidobacterium bifidum WBIN03 in relieving colitis by gut
microbiota, immune, and anti-oxidative stress. *Canadian Journal of Microbiology* **64**,
327-337 (2018). https://doi.org:10.1139/cjm-2017-0716
83   Kansal, S. *et al.* The Microbiome in Paediatric Crohn's Disease—A Longitudinal,
Prospective, ☐Single-Centre Study. *Journal of Crohn's and Colitis* **13**, 1044-1054
(2019). https://doi.org:10.1093/ecco-jcc/jjz016
84   Mao, B. *et al.* Blautia producta displays potential probiotic properties against
dextran sulfate sodium-induced colitis in mice. *Food Science and Human Wellness* **13**,
709-720 (2024). https://doi.org:10.26599/FSHW.2022.9250060
85   Liu, Y. *et al.* Sophora japonica flowers and their main phytochemical, rutin,
regulate chemically induced murine colitis in association with targeting the NF-κB

signaling pathway and gut microbiota. *Food Chemistry* **393**, 133395 (2022). https://doi.org:https://doi.org/10.1016/j.foodchem.2022.133395

Min, Z., Xiaona, H., Aziz, T., Jian, Z. & Zhennai, Y. Exopolysaccharides from Lactobacillus plantarum YW11 improve immune response and ameliorate inflammatory bowel disease symptoms. *Acta Biochim Pol* **67**, 485-493 (2020). https://doi.org:10.18388/abp.2020_5371

Li, Y. *et al.* Gegen Qinlian Decoction Alleviates Experimental Colitis and Concurrent Lung Inflammation by Inhibiting the Recruitment of Inflammatory Myeloid Cells and Restoring Microbial Balance. *J Inflamm Res* **15**, 1273-1291 (2022). https://doi.org:10.2147/jir.S352706

Gu, W., Zhang, L., Han, T., Huang, H. & Chen, J. Dynamic Changes in Gut Microbiome of Ulcerative Colitis: Initial Study from Animal Model. *Journal of Inflammation Research* **15**, 2631-2647 (2022). https://doi.org:10.2147/JIR.S358807

Johansson, M. E. V. *et al.* Bacteria Penetrate the Inner Mucus Layer before Inflammation in the Dextran Sulfate Colitis Model. *PLOS ONE* **5**, e12238 (2010). https://doi.org:10.1371/journal.pone.0012238

Walter, J. Ecological Role of Lactobacilli in the Gastrointestinal Tract: Implications for Fundamental and Biomedical Research. *Applied and Environmental Microbiology* **74**, 4985-4996 (2008). https://doi.org:10.1128/AEM.00753-08

Abdugheni, R. *et al.* Metabolite profiling of human-originated Lachnospiraceae at the strain level. *iMeta* **1**, e58 (2022). https://doi.org:https://doi.org/10.1002/imt2.58

Biddle, A., Stewart, L., Blanchard, J. & Leschine, S. Untangling the Genetic Basis of Fibrolytic Specialization by Lachnospiraceae and Ruminococcaceae in Diverse Gut Communities. *Diversity* **5**, 627-640 (2013).

Wei, H. *et al.* Diverse gut microbiota pattern between mild and severe cancer-related fatigue in lung cancer patients treated with first-line chemotherapy: A pilot study. *Thorac Cancer* **14**, 309-319 (2023). https://doi.org:10.1111/1759-7714.14765

Vacca, M. *et al.* The Controversial Role of Human Gut Lachnospiraceae. *Microorganisms* **8** (2020). https://doi.org:10.3390/microorganisms8040573

Wang, T. *et al.* Astilbin from Smilax glabra Roxb. alleviates high-fat diet-induced metabolic dysfunction. *Food Funct* **13**, 5023-5036 (2022). https://doi.org:10.1039/d2fo00060a

Zhu, Y., Li, Y., Liu, M., Hu, X. & Zhu, H. Guizhi Fuling Wan, Chinese Herbal Medicine, Ameliorates Insulin Sensitivity in PCOS Model Rats With Insulin Resistance via Remodeling Intestinal Homeostasis. *Frontiers in Endocrinology* **11** (2020).

Qi, Y. *et al.* Effects of Schisandra chinensis polysaccharides on rats with antibiotic-associated diarrhea. *International Journal of Biological Macromolecules* **124**, 627-634 (2019). https://doi.org:https://doi.org/10.1016/j.ijbiomac.2018.11.250

Su, L. *et al.* The Anti-colitis Effect of Schisandra chinensis Polysaccharide Is Associated With the Regulation of the Composition and Metabolism of Gut Microbiota. *Frontiers in Cellular and Infection Microbiology* **10** (2020).

Liu, C., Sun, C. & Cheng, Y. β-Glucan alleviates mice with ulcerative colitis through interactions between gut microbes and amino acids metabolism. *Journal of the Science of Food and Agriculture* **103**, 4006-4016 (2023). https://doi.org:https://doi.org/10.1002/jsfa.12357

100 Meehan, C. J. & Beiko, R. G. A Phylogenomic View of Ecological Specialization
in the Lachnospiraceae, a Family of Digestive Tract-Associated Bacteria. *Genome*
Biology and Evolution **6**, 703-713 (2014). https://doi.org:10.1093/gbe/evu050
101 Belzer, C. & de Vos, W. M. Microbes inside—from diversity to function: the case
of Akkermansia. *The ISME Journal* **6**, 1449-1458 (2012).
https://doi.org:10.1038/ismej.2012.6
102 Berry, D. *et al.* Phylotype-level 16S rRNA analysis reveals new bacterial
indicators of health state in acute murine colitis. *Isme j* **6**, 2091-2106 (2012).
https://doi.org:10.1038/ismej.2012.39
103 Li, M., Wu, Y., Hu, Y., Zhao, L. & Zhang, C. Initial gut microbiota structure affects
sensitivity to DSS-induced colitis in a mouse model. *Science China Life Sciences* **61**,
762-769 (2018). https://doi.org:10.1007/s11427-017-9097-0
104 Nagalingam, N. A., Kao, J. Y. & Young, V. B. Microbial ecology of the murine gut
associated with the development of dextran sodium sulfate-induced colitis. *Inflamm*
Bowel Dis **17**, 917-926 (2011). https://doi.org:10.1002/ibd.21462
105 Xue, L. *et al.* The effects of live and pasteurized Akkermansia muciniphila on
DSS-induced ulcerative colitis, gut microbiota, and metabolomics in mice. *Food &*
Function **14**, 4632-4646 (2023). https://doi.org:10.1039/D2FO03493J
106 Liu, Q. *et al.* Akkermansia muciniphila Exerts Strain-Specific Effects on DSS-
Induced Ulcerative Colitis in Mice. *Frontiers in Cellular and Infection Microbiology* **11**
(2021).
107 Ansaldo, E. *et al.* Akkermansia muciniphila induces intestinal adaptive immune
responses during homeostasis. *Science* **364**, 1179-1184 (2019).
https://doi.org:10.1126/science.aaw7479
108 Sugihara, K. *et al.* Mucolytic bacteria license pathobionts to acquire host-derived
nutrients during dietary nutrient restriction. *Cell Reports* **40** (2022).
https://doi.org:10.1016/j.celrep.2022.111093
109 Ganesh, B. P., Klopfleisch, R., Loh, G. & Blaut, M. Commensal Akkermansia
muciniphila Exacerbates Gut Inflammation in Salmonella Typhimurium-Infected
Gnotobiotic Mice. *PLOS ONE* **8**, e74963 (2013).
https://doi.org:10.1371/journal.pone.0074963
110 Falk, P. G., Hooper, L. V., Midtvedt, T. & Gordon, J. I. Creating and maintaining
the gastrointestinal ecosystem: what we know and need to know from gnotobiology.
Microbiol Mol Biol Rev **62**, 1157-1170 (1998). https://doi.org:10.1128/mmbr.62.4.1157-
1170.1998
111 van Passel, M. W. J. *et al.* The Genome of Akkermansia muciniphila, a
Dedicated Intestinal Mucin Degrader, and Its Use in Exploring Intestinal Metagenomes.
PLOS ONE **6**, e16876 (2011). https://doi.org:10.1371/journal.pone.0016876
112 Ouwerkerk Janneke, P. *et al.* Adaptation of Akkermansia muciniphila to the Oxic-
Anoxic Interface of the Mucus Layer. *Applied and Environmental Microbiology* **82**, 6983-
6993 (2016). https://doi.org:10.1128/AEM.01641-16
113 Vidal-Lletjós, S. *et al.* Mucosal healing progression after acute colitis in mice.
World J Gastroenterol **25**, 3572-3589 (2019). https://doi.org:10.3748/wjg.v25.i27.3572
114 Chang, Z. Y., Liu, H. M., Leu, Y. L., Hsu, C. H. & Lee, T. Y. Modulation of Gut
Microbiota Combined with Upregulation of Intestinal Tight Junction Explains Anti-

Inflammatory Effect of Corylin on Colitis-Associated Cancer in Mice. *Int J Mol Sci* **23** (2022). https://doi.org/10.3390/ijms23052667

Li, R. *et al.* Integrated Analysis Reveals the Targets and Mechanisms in Immunosuppressive Effect of Mesalazine on Ulcerative Colitis. *Frontiers in Nutrition* **9** (2022).

Wang, W. *et al.* Swimming Impedes Intestinal Microbiota and Lipid Metabolites of Tumorigenesis in Colitis-Associated Cancer. *Frontiers in Oncology* **12** (2022).

Lawson, P. A., Saavedra Perez, L. & Sankaranarayanan, K. Reclassification of Clostridium cocleatum, Clostridium ramosum, Clostridium spiroforme and Clostridium saccharogumia as Thomasclavelia cocleata gen. nov., comb. nov., Thomasclavelia ramosa comb. nov., gen. nov., Thomasclavelia spiroformis comb. nov. and Thomasclavelia saccharogumia comb. nov. *International Journal of Systematic and Evolutionary Microbiology* **73** (2023). https://doi.org:https://doi.org/10.1099/ijsem.0.005694

Milosavljevic, M. N. *et al.* Antimicrobial treatment of Erysipelatoclostridium ramosum invasive infections: a systematic review. *Rev Inst Med Trop Sao Paulo* **63**, e30 (2021). https://doi.org/10.1590/S1678-9946202163030

Zakham, F. *et al.* Molecular diagnosis and enrichment culture identified a septic pseudoarthrosis due to an infection with Erysipelatoclostridium ramosum. *Int J Infect Dis* **81**, 167-169 (2019). https://doi.org/10.1016/j.ijid.2019.02.001

Qiu, Z. *et al.* Targeted Metagenome Based Analyses Show Gut Microbial Diversity of Inflammatory Bowel Disease patients. *Indian Journal of Microbiology* **57**, 307-315 (2017). https://doi.org/10.1007/s12088-017-0652-6

Nagayama, M. *et al.* TH1 cell-inducing Escherichia coli strain identified from the small intestinal mucosa of patients with Crohn's disease. *Gut Microbes* **12**, 1788898 (2020). https://doi.org:10.1080/19490976.2020.1788898

Sun, M. *et al.* Combined Signature of the Fecal Microbiome and Plasma Metabolome in Patients with Ulcerative Colitis. *Med Sci Monit* **25**, 3303-3315 (2019). https://doi.org/10.12659/msm.916009

Mancabelli, L. *et al.* Identification of universal gut microbial biomarkers of common human intestinal diseases by meta-analysis. *FEMS Microbiology Ecology* **93**, fix153 (2017). https://doi.org:10.1093/femsec/fix153

Cao, R.-R., He, P. & Lei, S.-F. Novel microbiota-related gene set enrichment analysis identified osteoporosis associated gut microbiota from autoimmune diseases. *Journal of Bone and Mineral Metabolism* **39**, 984-996 (2021). https://doi.org/10.1007/s00774-021-01247-w

Cai, S. *et al.* Gut Bacteria Erysipelatoclostridium and Its Related Metabolite Ptilosteroid A Could Predict Radiation-Induced Intestinal Injury. *Frontiers in Public Health* **10** (2022).

Farooq, P. D., Urrunaga, N. H., Tang, D. M. & von Rosenvinge, E. C. Pseudomembranous colitis. *Disease-a-Month* **61**, 181-206 (2015). https://doi.org:https://doi.org/10.1016/j.disamonth.2015.01.006

Liangen, L., Man, C., Hongcheng, Z., Bangzhou, Z. & Chuanxing, X. IDDF2023-ABS-0092 Multi-cohort retrospective microbiota analysis on fecal microbiota transplantation in ulcerative colitis. *Gut* **72**, A192 (2023). https://doi.org/10.1136/gutjnl-2023-IDDF.181

Feng, Y. *et al.* Barley Leaf Ameliorates Citrobacter rodentium-Induced Colitis through Preventive Effects. *Nutrients* **14** (2022). https://doi.org:10.3390/nu14183833

Chai, M. *et al.* Different Bifidobacterium bifidum strains change the intestinal flora composition of mice via different mechanisms to alleviate loperamide-induced constipation. *Food & Function* **12**, 6058-6069 (2021). https://doi.org:10.1039/D1FO00559F

Sullivan, S. *et al.* Downregulation of sodium transporters and NHERF proteins in IBD patients and mouse colitis models: potential contributors to IBD-associated diarrhea. *Inflamm Bowel Dis* **15**, 261-274 (2009). https://doi.org:10.1002/ibd.20743

Brook, I. & Long, S. S. in *Principles and Practice of Pediatric Infectious Diseases (Fifth Edition)*   (eds Sarah S. Long, Charles G. Prober, & Marc Fischer) 987-995.e982 (Elsevier, 2018).

Atarashi, K. *et al.* Treg induction by a rationally selected mixture of Clostridia strains from the human microbiota. *Nature* **500**, 232-236 (2013). https://doi.org:10.1038/nature12331

Narushima, S. *et al.* Characterization of the 17 strains of regulatory T cell-inducing human-derived Clostridia. *Gut Microbes* **5**, 333-339 (2014). https://doi.org:10.4161/gmic.28572

Vitalii, S., Musfiqur, S., Camilo, V., Kalai, M. & Giri, N. Novel Approach for Microbiome Analysis Using Bacterial Replication Rates and Causal Inference with Applications. *bioRxiv*, 2020.2005.2021.108514 (2020). https://doi.org:10.1101/2020.05.21.108514

Li, S. *et al.* Frequency of change determines effectiveness of microbial response strategies. *The ISME Journal* **17**, 2047-2057 (2023). https://doi.org:10.1038/s41396-023-01515-9

Beller, Z. W. *et al.* Inducible CRISPR-targeted "knockdown" of human gut Bacteroides in gnotobiotic mice discloses glycan utilization strategies. *Proceedings of the National Academy of Sciences* **120**, e2311422120 (2023). https://doi.org:10.1073/pnas.2311422120

Lloyd-Price, J. *et al.* Multi-omics of the gut microbial ecosystem in inflammatory bowel diseases. *Nature* **569**, 655-662 (2019). https://doi.org:10.1038/s41586-019-1237-9

Araki, Y., Mukaisyo, K., Sugihara, H., Fujiyama, Y. & Hattori, T. Increased apoptosis and decreased proliferation of colonic epithelium in dextran sulfate sodium-induced colitis in mice. *Oncol Rep* **24**, 869-874 (2010). https://doi.org:10.3892/or.2010.869

Vetuschi, A., Latella, G., Sferra, R., Caprilli, R. & Gaudio, E. Increased Proliferation and Apoptosis of Colonic Epithelial Cells in Dextran Sulfate Sodium-Induced Colitis in Rats. *Digestive Diseases and Sciences* **47**, 1447-1457 (2002). https://doi.org:10.1023/A:1015931128583

Raymond, F. *et al.* The initial state of the human gut microbiome determines its reshaping by antibiotics. *Isme j* **10**, 707-720 (2016). https://doi.org:10.1038/ismej.2015.148

Brugiroux, S. *et al.* Genome-guided design of a defined mouse microbiota that confers colonization resistance against Salmonella enterica serovar Typhimurium. *Nature Microbiology* **2**, 16215 (2016). https://doi.org:10.1038/nmicrobiol.2016.215

142   FitzGerald, J. *et al.* Improved gut microbiome recovery following drug therapy is
linked to abundance and replication of probiotic strains. *Gut Microbes* **14**, 2094664
(2022). https://doi.org:10.1080/19490976.2022.2094664
143   Robertson, S. J. *et al.* Comparison of Co-housing and Littermate Methods for
Microbiota Standardization in Mouse Models. *Cell Rep* **27**, 1910-1919 e1912 (2019).
https://doi.org:10.1016/j.celrep.2019.04.023
144   Russell, A. *et al.* Reduced housing density improves statistical power of murine
gut microbiota studies. *Cell Rep* **39**, 110783 (2022).
https://doi.org:10.1016/j.celrep.2022.110783
145   Singh, G., Brass, A., Cruickshank, S. M. & Knight, C. G. Cage and maternal
effects on the bacterial communities of the murine gut. *Sci Rep* **11**, 9841 (2021).
https://doi.org:10.1038/s41598-021-89185-5
146   McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible
interactive analysis and graphics of microbiome census data. *PLoS One* **8**, e61217
(2013). https://doi.org:10.1371/journal.pone.0061217
147   rbiom: Read/Write, Analyze, and Visualize 'BIOM' Data v. 2.0.0.9003 (2023).
148   pairwiseAdonis: Pairwise multilevel comparison using adonis v. 0.4 (2020).
149   FastQC: A Quality Control Tool for High Throughput Sequence Data (2010).
150   Ewels, P., Magnusson, M., Lundin, S. & Kaller, M. MultiQC: summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047-3048
(2016). https://doi.org:10.1093/bioinformatics/btw354
151   Li, D., Liu, C. M., Luo, R., Sadakane, K. & Lam, T. W. MEGAHIT: an ultra-fast
single-node solution for large and complex metagenomics assembly via succinct de
Bruijn graph. *Bioinformatics* **31**, 1674-1676 (2015).
https://doi.org:10.1093/bioinformatics/btv033
152   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat*
Methods **9**, 357-359 (2012). https://doi.org:10.1038/nmeth.1923
153   Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition.
Nat Methods **11**, 1144-1146 (2014). https://doi.org:10.1038/nmeth.3103
154   Wu, Y. W., Simmons, B. A. & Singer, S. W. MaxBin 2.0: an automated binning
algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**,
605-607 (2016). https://doi.org:10.1093/bioinformatics/btv638
155   Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and
efficient genome reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).
https://doi.org:10.7717/peerj.7359
156   Sieber, C. M. K. *et al.* Recovery of genomes from metagenomes via a
dereplication, aggregation and scoring strategy. *Nat Microbiol* **3**, 836-843 (2018).
https://doi.org:10.1038/s41564-018-0171-1
157   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W.
CheckM: assessing the quality of microbial genomes recovered from isolates, single
cells, and metagenomes. *Genome Res* **25**, 1043-1055 (2015).
https://doi.org:10.1101/gr.186072.114
158   Bowers, R. M. *et al.* Minimum information about a single amplified genome
(MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and archaea.
Nat Biotechnol **35**, 725-731 (2017). https://doi.org:10.1038/nbt.3893

159   GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database
(2020).
160   Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**,
2068-2069 (2014). https://doi.org:10.1093/bioinformatics/btu153
161   Fitting Linear Mixed-Effects Models Using lme4 (2015).
162   Kuznetsova A., B. P. B., Christensen R.H.B. lmerTest Package: Tests in Linear
Mixed Effects Models. *Journal of Statistical Software* **82** (2017).
https://doi.org:https://doi.org/10.18637/jss.v082.i1310.18637/jss.v082.i13