Synaptic plasticity in human thalamocortical assembloids

Authors

Mary H. Patton[1,†], Kristen T. Thomas[1,†], Ildar T. Bayazitov[1], Kyle D. Newman[1], Nathaniel B. Kurtz[2], Camenzind G. Robinson[2], Cody A. Ramirez[1], Alexandra J. Trevisan[1], Jay B. Bikoff[1], Samuel T. Peters[3], Shondra M. Pruett-Miller[3,4], Yanbo Jiang[1], Andrew B. Schild[1], Anjana Nityanandam[1], Stanislav S. Zakharenko[1,*]

Affiliations

[1]Department of Developmental Neurobiology, St. Jude Children's Research Hospital; Memphis, TN 38105, USA

[2]Cell and Tissue Imaging Center, St. Jude Children's Research Hospital; Memphis, TN 38105, USA

[3]Center for Advanced Genome Engineering, St. Jude Children's Research Hospital; Memphis, TN 38105, USA

[4]Department of Cell & Molecular Biology, St. Jude Children's Research Hospital; Memphis, TN 38105, USA

[*]Corresponding author: Stanislav S. Zakharenko, stanislav.zakharenko@stjude.org

[†]These authors contributed equally to this work.

Highlights

- Human thalamic organoids consist of mostly glutamatergic projection neurons.

- Thalamocortical assembloids form reciprocal glutamatergic synapses.

- Synapses are functional and undergo short-term plasticity resembling animal models.

- Long-term potentiation and depression reveal mechanisms distinct from rodents.

eTOC

Human organoids are often used to model diseases with synaptic pathology; however, few studies have examined synaptic function via single-cell or single-synapse recordings. Patton et al. fused human thalamic and cortical organoids into assembloids to examine synaptic transmission and short- and long-term synaptic plasticity in human thalamocortical and corticothalamic circuits.

35 **GRAPHICAL ABSTRACT**

SUMMARY

Synaptic plasticities, such as long-term potentiation (LTP) and depression (LTD), tune synaptic

efficacy and are essential for learning and memory. Current studies of synaptic plasticity in

humans are limited by a lack of adequate human models. Here, we modeled the thalamocortical

system by fusing human induced pluripotent stem cell–derived thalamic and cortical organoids.

Single-nucleus RNA-sequencing revealed that most cells in mature thalamic organoids were

glutamatergic neurons. When fused to form thalamocortical assembloids, thalamic and cortical

organoids formed reciprocal long-range axonal projections and reciprocal synapses detectable

by light and electron microscopy, respectively. Using whole-cell patch-clamp electrophysiology

and two-photon imaging, we characterized glutamatergic synaptic transmission. Thalamocortical

and corticothalamic synapses displayed short-term plasticity analogous to that in animal

models. LTP and LTD were reliably induced at both synapses; however, their mechanisms

differed from those previously described in rodents. Thus, thalamocortical assembloids provide

a model system for exploring synaptic plasticity in human circuits.

INTRODUCTION

Synaptic transmission through the release of neurotransmitter and the subsequent activation of postsynaptic receptors is the fundamental mode of communication between neurons. Synaptic plasticity enhances or diminishes synaptic transmission in an activity-dependent manner in response to environmental cues. Although the chemical synapse is evolutionarily conserved,[1] the molecular mechanisms underlying synaptic plasticity differ across species.[2–6] Aberrant synaptic plasticity is well documented in animal models of autism, schizophrenia, and other psychiatric disorders.[7–10] Ultimately, these human disorders would benefit from a human model system.

Ex vivo human model systems are currently limited to postmortem samples, which are often in a state of uncertain tissue quality, and biopsy samples, which are almost exclusively from individuals with epilepsy or brain tumors.[11] The recent development of human induced pluripotent stem cell (hiPSC)-derived organoids provides a promising *in vitro* model system that is accessible to experimental manipulation and recapitulates human-specific features of neural development and function.[12–18] Organoids are widely used to model synaptopathies (i.e., neurologic and psychiatric disorders associated with synaptic dysfunction).[19–25] Specific properties of synaptic transmission and synaptic plasticity differ across brain regions and neuronal subtypes;[2,26–28] however, few studies have examined synaptic transmission between defined pre- and postsynaptic cell populations in organoids.[29–31] Moreover, synaptic plasticity within organoids has never been reported using whole-cell patch-clamp electrophysiology, the gold standard approach for studying synaptic physiology that has been validated over decades of research in animal models.

To build a functional neural circuit, human cortical organoids (hCOs) can be fused with human thalamic organoids (hThOs) to produce thalamocortical (TC) assembloids.[32] The thalamus is a primary relay center for incoming sensory information that sends widespread, yet highly organized, projections to various cortical regions based on the sensory modality.[33–36] The cortex, in turn, sends projections back to thalamic nuclei to integrate and update sensory, motor, and

cognitive information.[37,38] Collectively, synaptic transmission within the thalamo–cortico–thalamic circuit creates cognitive representations of the outside world based on diverse incoming sensory inputs and provides the foundation for dynamic executive functioning.[39–42] Experience-dependent synaptic plasticity within this system is critical to learning and underlies the expression of sensorimotor behaviors, attention, and perceptual and working memory.[42–47]

Here we developed a human assembloid system containing functional glutamatergic TC and corticothalamic (CT) synaptic connections that undergo short- and long-term synaptic plasticity. We then used this system to explore the molecular mechanisms underlying synaptic plasticity in human neurons.

RESULTS

hThOs contain functional glutamatergic thalamic neurons

To build optimal hThOs, we generated a reporter line from an hiPSC line derived from a neurotypical male subject with normal karyotype (See **Figure S1** for reporter line validation). Specifically, the *tdTomato*-coding sequence was inserted into the endogenous *TCF7L2* locus (**Figure 1A**). We then modified a previously reported protocol[32] to increase the efficiency of thalamic neuron generation. After differentiation into hThOs, we performed bulk RNA-sequencing and then VoxHunt deconvolution analysis of the RNA-seq data. High *tdTomato* RNA levels identified hThOs with high representation of diencephalon, the developmental structure that gives rise to the thalamus, and low representation from contaminating structures, e.g., the pallium and midbrain (**Figure 1A**). Notably, in hThOs generated from five independent hiPSC lines, *TCF7L2* positively predicted the expression of the thalamic neural precursor genes *OLIG3* and *OTX2* and the thalamic neuron genes *GBX2* and *LHX9* (**Figure S2**). Visual assessment was sufficient to identify hThOs with high *tdTomato*, as differential expression analysis comparing hThOs with high TCF7L2-tdTomato fluorescence to hCOs revealed significant enrichment of thalamic markers in

the hThOs (**Figure S3A** and **S3B**). All subsequent experiments were performed with TCF7L2-tdTomato[+] hThOs.

At 60 days after the start of differentiation (D60), hThOs contained OTX2[+] and SOX2[+] neural progenitor domains surrounded by TUBB3[+] neurons (**Figure 1B**). By D92, most cells expressed markers consistent with glutamatergic neurons of the developing thalamus, specifically LHX2, FOXP2, and GBX2; GABA immunoreactivity was observed in only a small subset of cells (**Figure S3C**). At D70–D90 in culture, synaptic gene expression significantly increased, and precursor and mitotic gene expression decreased in hThOs (**Figure S3D and S3E**).

Single-nucleus RNA-sequencing (snRNA-seq) analysis of D90 hThOs revealed that the majority (~85%) of cells were glutamatergic neurons found primarily in Excitatory Neuron 1-4 (ExN1-4) clusters (**Figure 1C-1F**). Applying VoxHunt mapping to *BrainSpan* and *Allen Brain Atlas* references, we found that clusters ExN1-4 mapped to the human mediodorsal thalamus (**Figure 1D**) and embryonic day (E) 15 mouse caudal thalamus (**Figure 1G**), the diencephalon structure that produces the glutamatergic neurons of the mature thalamus. Consistent with thalamic neurons, ExN1-4 neurons primarily expressed *SLC17A6* (or *VGLUT2*); *SLC17A7* (or *VGLUT1*) was sparsely expressed (**Figure 1F**). Neurons in clusters ExN1-4 expressed additional thalamic markers of interest, including *GBX2*, *SHOX2*, *FOXP2*, *CADM1*, and *NTNG1* (**Figure S3F**). In line with reports from the mouse thalamus,[48] a subset of cells also expressed *SOX2*, a marker typically associated with precursors rather than mature neurons (**Figure S3F**). We also identified one cluster that mapped to E15 mouse pretectum (PT) (**Figure 1H**), as well as cells expressing markers of the thalamic organizer zona limitans intrathalamica (ZLI) and rostral thalamus (rTh) (**Figure 1I**). Like the caudal thalamus, these structures arise from the diencephalon during early development. A small subset of *SLC17A6*[+] glutamatergic neurons and all *GAD1*[+] GABAergic neurons within the hThOs were found in this PT/ZLI/rTh cluster.

To verify that most cells in hThOs were functional neurons, we used whole-cell patch-clamp electrophysiology to investigate the membrane properties of individual cells. Action

potentials (APs) were evoked in response to depolarizing current injections (**Figure 1J**). The resting membrane potential, membrane capacitance, input resistance, and AP properties measured in hThO cells were consistent with functional neurons (**Figure S4A-S4H**). Transmission electron microscopy (TEM) also revealed numerous asymmetric and symmetric synapses in hThOs (**Figure S4Q** and **S4R**). A subset of presynaptic terminals contained dense core vesicles (**Figure S4S**).

Finally, we examined the non-neuronal cell populations in our hThOs. Pseudotime analysis (**Figure 1K**) and cell cycle analysis (**Figure 1L-1N**) revealed a small cluster of precursor cells undergoing mitosis (i.e., Cycling Progenitors cluster). *TNC*⁺ progenitors not undergoing mitosis were labeled radial glia. Most of the remaining cells were *GFAP*⁺ astrocytes within the Glia cluster (**Figure 1O**). Pseudotime analysis also revealed differences in glutamatergic neuron maturity, with the ExN4 cluster containing the least mature thalamic neurons, and the ExN2 cluster containing the most mature thalamic neurons (**Figure 1K**). We conclude that our hThO protocol generated functional thalamic neurons with high efficiency, and that those neurons formed synaptic connections.

TC assembloids form functional glutamatergic TC and CT synapses

To form TC assembloids, we also required optimal hCOs. We generated an isogenic hiPSC reporter line by inserting the *tdTomato*-coding sequence into the endogenous *SLC17A7* (*VGLUT1*) locus (**Figure 2A**) (See **Figure S1** for reporter line validation). Using this reporter line, we generated hCOs via a previously reported protocol.[49] Voxhunt deconvolution analysis of bulk RNA-seq data demonstrated that VGLUT1-tdTomato⁺ hCOs exhibited high representation of the pallium, the developmental structure that gives rise to the neocortex, and low representation of contaminating structures, e.g., the subpallium and diencephalon (**Figure 2A**). High expression of *tdTomato* RNA also predicted high expression of cortical markers relative to hThOs (**Figure S3A**

and **S3B**) and VGLUT1-tdTomato⁻ hCOs (**Figure S5A**). All subsequent experiments were performed with VGLUT1-tdTomato⁺ hCOs.

The snRNA-seq analysis of D90 hCOs revealed that most cells fell within a neuronal developmental trajectory, beginning with neural precursors and ending with differentiated neurons that expressed markers of upper-layer (UL) or deep-layer (DL) excitatory cortical neurons (UL ExNs and DL ExNs, respectively) (**Figures 2B** and **S5**). Using VoxHunt mapping to *BrainSpan* and *Allen Brain Atlas* references, we found that clusters within this trajectory mapped to human neocortical structures (**Figure S5B**) and E15 mouse neocortex (**Figure 2C**). During early development, thalamic neurons first form synapses with the cortical subplate before transitioning to cortical Layer IV.[50] We identified a cluster in hCOs (Subplate/DL ExN) that was enriched for markers of the cortical subplate[51] (**Figure 2B**) and contained the most mature neurons based on pseudotime analysis (**Figure 2D**), which was in line with this cluster containing subplate-like neurons.[52] Consistent with previous reports in rodent models,[53] *SLC17A6* was expressed in intermediate progenitors, but expression declined with neuronal maturation (**Figure 2E**). Mature neurons in hCOs exclusively expressed *SLC17A7* (**Figure 2F**). Remaining cells included unidentified glutamatergic neurons (found in the Un.ExN1 and Un.ExN2 clusters), glia, and cells resembling those in choroid plexus (**Figure S5I and S5J**).

We investigated the firing properties of hCO cells by using whole-cell patch-clamp electrophysiological recordings. Delivering depolarizing current injections to hCO cells evoked AP firing (**Figures 2F** and **S4L-S4P**). These cells also displayed typical neuronal properties (**Figure S4I-S4P**). TEM revealed numerous asymmetric and symmetric synapses in hCOs (**Figure S4Q and S4R**). A subset of presynaptic terminals contained dense core vesicles (**Figure S4S**).

We then fused hThOs with hCOs to form TC assembloids. The hCOs were transduced with hSyn-GFP lentivirus prior to fusion, so each organoid could be identified within the assembloid (**Figure 2G**). GFP⁺ axons from the hCO were detectable within the hThO within 5

days postfusion (dpf). Furthermore, 2-dimensional fusion assays confirmed that hThOs and hCOs

sent reciprocal axonal projections (**Figure 2H**).

We then sought to identify TC and CT synapses formed between the organoids after

fusion. To that end, we transduced either the hThO or hCO with hSyn-V5-Mito-APEX2 lentivirus,

which localized the V5-tagged peroxidase APEX2 to the mitochondrial matrix in neurons, enabling

the identification of the hThO or hCO origin of the presynaptic terminal.[54] APEX2$^+$ hThOs were

fused with APEX2$^-$ hCOs (or vice versa) to form TC assembloids. Light microscopy and

immunolabeling identified V5$^+$ puncta that co-localized with neurons expressing hSyn-GFP

(**Figure 2I**). Reaction with DAB produced strong contrast in the matrix of APEX2$^+$ mitochondria in

TEM images. TEM images revealed APEX2$^+$ mitochondria in presynaptic terminals from the hThO

that formed TC synapses within the hCO after fusion (**Figure 2J**). Conversely, we observed

APEX2$^+$ mitochondria in presynaptic terminals from the hCOs that formed CT synapses within the

hThOs after fusion (**Figure 2K**). This analysis confirmed that assembloids contained both TC and

CT synapses.

Whole-cell patch-clamp electrophysiology recordings confirmed that these synapses were

functional (**Figure 2L-2N**). Electrical synaptic stimulation of the hThO (**Figure 2L**) or hCO (**Figure

2M**) evoked excitatory postsynaptic currents (EPSCs) in cells recorded in the hCO or hThO,

respectively. The likelihood of evoking a synaptic response varied among assembloids (**Figure

2L** and **2M**); on average, the chance of cells responding to electrical stimulation of the opposite

organoid was 61% for the TC synapses and 58% for the CT synapses (**Figure 2M**).

To begin characterizing this synaptic response, we calculated the paired-pulse ratio (PPR)

of EPSCs, a classic measure of presynaptic short-term plasticity, at TC and CT synapses.[51–57] In

response to a pair of stimuli applied to a presynaptic neuron, TC synapses elicited paired-pulse

depression, wherein the second postsynaptic response was weaker than the first. In contrast, CT

synapses were more prone to paired-pulse facilitation, wherein the second response was stronger

than the first (**Figure 2N**). Both results resemble previous observations of these synapses in animal models.[55–61]

Next, we investigated the potential cellular origin of the functional connections between hThOs and hCOs. We used NeuronChat[62] to determine the likelihood of neuronal communication between cell clusters in the hThOs and hCOs based on snRNA-seq data. We found that the hThO ExN clusters, which contain glutamatergic thalamic neurons, exhibited the highest probability of TC communication with cells in the Cycling Progenitor and Subplate/DL ExN clusters of the hCOs (**Figure 3A**). Conversely, Subplate/DL ExNs exhibited a higher probability of CT communication with hThO ExN clusters than exhibited by other hCO clusters (**Figure 3B**). NRXN signaling mediates synaptogenesis, whereas glutamate signaling mediates excitatory synaptic transmission. Further analysis revealed that hThO ExNs exhibited a high probability of TC communication with hCO Cycling Progenitors and Subplate DL/ExNs clusters by NRXN signaling (**Figure 3C**). The hCO Subplate/DL ExNs and hThO ExNs also exhibited a higher probability of TC and CT communication using glutamate than other clusters (**Figure 3D**). Our snRNA-seq data confirmed that these clusters expressed the necessary ligands and receptors for NRXN and glutamatergic signal transduction (**Figure S6**). Overall, NeuronChat predicted that TC glutamatergic synaptic transmission most likely occurs between the hThO ExN2 cluster, which contains the most mature thalamic neurons, and the hCO Subplate/DL ExNs. Conversely, CT glutamatergic synaptic transmission most likely occurs between the hCO Subplate/DL ExNs and the hThO ExN1 cluster, the largest cluster containing glutamatergic thalamic neurons.

Next, we used whole-cell patch-clamp electrophysiology to further characterize TC and CT synaptic transmission. Evoked TC and CT EPSCs showed typical glutamatergic ionotropic properties comprising a fast α-amino-3-hydroxy-5-methyl-4-isoxazoleproprionic acid receptor (AMPAR)-mediated component blocked by the AMPAR inhibitor NBQX (3 mM) [86.7% ± 3.3% AMPAR current reduction for TC synapses (**Figure 3E** and **3G**); 82.2% ± 3.8% AMPAR current reduction for CT synapses, (**Figure 3H** and **3J**)] and a slow *N*-methyl-D-aspartate receptor

(NMDAR)-mediated component blocked by the NMDAR inhibitor AP5 (50 µM) [84.0% ± 4.6% NMDAR current reduction for TC synapses (**Figure 3F** and **3G**); 78.4% ± 2.9% NMDA current reduction for CT synapses (**Figure 3I** and **3J**)].

Using whole-cell patch-clamp electrophysiology and two-photon calcium imaging, we identified the sites of synaptic transmission. We detected synaptically evoked calcium transients in hCO cells upon stimulation of hThOs (**Figure 3K-3N**). These postsynaptic sites in the hCO cells resembled dendritic spines described in cortical neurons of animal models (**Figure 3L**).[63] Stimulation of the hThO evoked stronger calcium transients in dendritic spines compared to that in parent dendritic shafts (**Figure 3M** and **3N**), which suggested that the dendritic spines were synaptically connected to hThO axons. Moreover, calcium transients in dendritic spines were blocked by AP5 [82.4% ± 5.2% reduction (**Figure 3O**)], indicative of glutamatergic synaptic transmission.

TC and CT synapses undergo LTP in assembloids

Having established the existence of functional synaptic connections between hThOs and hCOs, we tested whether the TC and CT pathways undergo long-term synaptic plasticities, specifically LTP and LTD. To examine whether TC synapses undergo LTP, we tested several LTP-induction protocols: spike-timing–dependent plasticity (STDP) induction protocols and high-frequency tetanization. TC LTP was reliably induced by high-frequency (40-Hz) tetanization of thalamic inputs (**Figure 4A**), which increased EPSC amplitudes by 168.2% ± 19.3% compared to baseline (**Figure 4B** and **4C**). STDP is induced by stimulating presynaptic inputs and directly depolarizing the postsynaptic cell (**Figure 4D** and **4G**); STDP is based on the precise order and timing of pre- and postsynaptic activity.[64–66] Following a short (×1) STDP-induction protocol, the amplitude of TC EPSCs increased by 144.4% ± 17.8%, compared to baseline (**Figure 4D-4F**). Following a long (×3) STDP-induction protocol, EPSC amplitudes increased by 223.9% ± 24.4%, compared to baseline (**Figure 4G-4L**). This TC LTP was observed in all tested (9/9) cells, from six separate

assembloids, across three batches of differentiation (**Figure 4H**). TC LTP was not caused by changes in series resistance (**Figure 4I**); thus, it represents a true activity-dependent potentiation of synaptic strength.

Next, we investigated the mechanisms underlying TC LTP in assembloids. Bath application of the metabotropic glutamate 5 (mGluR5) antagonist MPEP (10 µM) blocked TC LTP, but the NMDAR antagonist AP5 (50 µM) did not (**Figure 4J** and **4K**). TC LTP also required postsynaptic Ca^{2+}. When we included the Ca^{2+} chelator BAPTA (20 mM) in the internal pipette solution (iBAPTA), the long STDP protocol not only failed to induce LTP but also moderately reduced the TC EPSC amplitude (**Figure 4J** and **4K**). We then tested if TC LTP was expressed presynaptically by measuring changes in PPR after LTP induction. PPR decreased (suggesting an increase in the probability of glutamate release from presynaptic terminals) compared to baseline in control and AP5 conditions, and this change was blocked in the presence of MPEP (**Figure S7A-S7D**). Together, these findings demonstrate that the TC pathway in assembloids undergoes LTP via multiple induction protocols, with the long (×3) STDP-evoked LTP induced postsynaptically and expressed, at least partially, presynaptically through mGluR5-dependent mechanisms.

CT synapses also underwent LTP after the long STDP-induction protocol (**Figure 5**). LTP was observed in 12/14 cells recorded from nine separate assembloids across two differentiation batches (**Figure 5B**). On average, EPSC amplitudes increased by 158.3% ± 15.2%, compared to baseline after LTP induction (**Figure 5C** and **E**). CT LTP was not caused by changes in series resistance (**Figure 5D**), representing a true potentiation of synaptic strength. Inclusion of iBAPTA in the internal pipette solution blocked CT LTP (**Figure 5C** and **E**), and unlike TC LTP, CT LTP was also blocked by separate application of MPEP and AP5 (**Figure 5C** and **E**). There were no changes in PPR in any of the conditions (**Figure S7E-S7H**), suggesting that CT LTP does not involve changes in presynaptic-release probabilities. These data suggest that the long (×3) STDP

induction protocol generates LTP in the CT pathway in assembloids, and this LTP is induced and expressed postsynaptically through both mGluR5- and NMDAR-dependent mechanisms.

TC and CT synapses undergo LTD in assembloids

The activity-dependent weakening of synaptic transmission between brain regions through LTD is a key component of learning.[67] Therefore, we tested whether TC and CT synapses in assembloids undergo LTD. Delivering low-frequency (1-Hz) stimulation to the hThO while recording from hCO cells depressed EPSCs in 8/9 cells recorded in nine assembloids from three differentiation batches (**Figure 6A** and **B**). On average, TC EPSC amplitudes decreased by 59.4% ± 8.1% of baseline after low-frequency stimulation (**Figure 6C-6F**). TC LTD was blocked by iBAPTA or bath-applied MPEP or AP5 (**Figure 6C** and **6E**). Low-frequency stimulation of the hThO did not change PPR across any of the conditions (**Figure S7I-S7L**). These findings provide evidence for a postsynaptically induced and expressed LTD in the TC pathway that depends on both mGluR5s and NMDARs.

Low-frequency stimulation of hCO inputs to hThO cells also reliably induced LTD, as it was observed in 10/10 cells recorded in nine assembloids across two batches of differentiation (**Figure 7A** and **B**). On average, the expression of CT LTD was reflected in a 65.8% ± 5.2% reduction of baseline EPSC amplitudes (**Figure 7C-7F**). CT LTD was blocked by iBAPTA or bath application of MPEP or AP5 (**Figure 7C** and **7E**). PPR at CT synapses was unchanged after 1-Hz stimulation across all conditions (**Figure S7M-S7P**). Neither TC LTD nor CT LTD occurred due to changes in series resistance (**Figures 6D** and **7D**, respectively). These data suggest that CT LTD is induced and expressed postsynaptically and requires both mGluR5s and NMDARs Notably, reversing the order of presynaptic stimulation and postsynaptic depolarization, in a reverse long STDP protocol, did not induce LTD in either the TC or CT pathways (data not shown).

DISCUSSION

Here we describe a novel, hiPSC-derived TC assembloid system for exploring synaptic transmission and synaptic plasticity in human neural circuits. Within assembloids, hThOs and hCOs form reciprocal glutamatergic synapses capable of short- and long-term synaptic plasticity. Notably, the vast majority (92.9%) of synaptically connected cells underwent LTP or LTD during the respective induction protocols. However, these synapses displayed a degree of specificity, as one established induction protocol (reverse long STDP protocol) failed to induce LTD at either synapse. Together, our findings suggest that synaptic plasticity is robust, highly replicable, and selective in the TC assembloid system.

Previous work describing synaptic plasticity in organoids has exclusively relied on multielectrode array (MEA) recordings.[68] Although this approach is high throughput and convenient, MEAs measure extracellular spike and local field activities that arise from a large group of cells. The specific mechanisms underlying synaptic transmission and synaptic plasticity depend on pre- and postsynaptic cell identity. Neither pre- nor postsynaptic cell identities in organoids are easily determined in MEA recordings because organoids contain heterogeneous cell populations that lack defined electrophysiological profiles. Reflecting this heterogeneity, MEA recordings in organoids detected LTP, LTD, and bidirectional short-term synaptic plasticity in response to identical induction protocols.[68] Few studies have examined synaptic transmission within a specific circuit in human brain organoids.[29–31] More commonly, the properties of spontaneously released neurotransmitters onto single cells in organoids have been characterized,[69–71] but the presynaptic sources of that transmission were not defined.

Our study used whole-cell patch-clamp electrophysiology, a well-validated method that has enabled decades of synaptic physiology discoveries in multiple species. Here we accurately measured electrically evoked synaptic currents in single postsynaptic cells. Furthermore, visually differentiating between GFP$^+$ hCOs and GFP$^-$ hThOs within the assembloid enabled us to evoke responses from a defined neuronal population and measure changes in synaptic transmission in

distinct postsynaptic cells. To our knowledge, our results are the first to characterize functional

synaptic connections and the underlying mechanisms of synaptic plasticity in a human-derived

assembloid system by using these techniques.

Although there are some similarities between our LTP/LTD findings from human

assembloids and those from rodents, specifically the high prevalence for NMDAR-mediated LTP

and LTD,[72] in general the mechanisms underlying LTP/LTD in the rodent TC and CT pathways

are distinct from what we report here.[73–75] In mouse auditory and somatosensory TC pathways,

LTP depends on postsynaptically expressed group 1 mGluRs, whereas LTP in the barrel cortex

requires postsynaptic NMDAR activation and subsequent Ca^{2+} entry into the postsynaptic cell.[76–

80] CT LTP in rodents is expressed presynaptically, requires a rise in presynaptic Ca^{2+} and protein

kinase A activation,[81] but does not require the activation of NMDARs or mGluRs[81]. In mice, LTD

within the somatosensory cortex is mediated by NMDARs,[82] and LTD in the barrel cortex requires

presynaptic type 1 cannabinoid receptors.[78] In contrast, we found that three of the four types of

long-term synaptic plasticity we measured required both mGluR5 and NMDARs. A functional link

between group 1 mGluRs and NMDAR activity is seen in various brain regions.[83,84] In mouse

cortical neurons, the activation of mGluR1 potentiates NMDAR-mediated currents through

downstream signaling.[85] Given our findings, a similar mGluR5-dependent mechanism may exist

in human-derived TC assembloids.

Our results may reflect species-specific differences between rodents and humans in the

expression and maintenance of long-term synaptic plasticity. However, organoids resemble fetal

human brain more closely than they do postnatal structures,[86] and most rodent studies of synaptic

physiology are conducted in postnatal animals. Unlike the adult counterparts, the immature

thalamus does not contain well-differentiated nuclei, and the fetal cortical plate lacks well-defined

layers. Organoids in their current forms display similar limitations in their structural organization.

Therefore, the mechanisms we define in TC assembloid synaptic plasticity may differ from those

observed in rodents due to species differences, developmental differences, or a combination thereof.

The organoid field is constantly improving. A recent report described hThOs that more closely resemble the thalamic reticular nucleus.[87] Future studies will derive organoids that better model specific projection nuclei of the thalamus. In the developing brain, thalamic inputs mold the laminar, columnar, and functional organization of the cortex.[88,89] Similarly, more mature thalamic projections might promote organizational maturation in hCOs, and more organized TC assembloids might better model the diversity of synaptic plasticity mechanisms observed across TC sensory pathways. Assembloids modeling other brain structures[90–92] (or synaptic targets outside the brain)[93] might also elucidate mechanisms that differ between neural circuits.

We anticipate that hiPSC-derived organoids and assembloids will provide a particularly useful model system for exploring synaptic pathology in human neurologic and psychiatric disorders. To date, most organoid studies have focused on disease-associated changes in gene expression, cellular composition, or neural network activity. Our data suggest that assembloids derived from patient hiPSCs or from hiPSCs carrying disease-associated mutations can be used to model disease-associated deficits in synaptic transmission and synaptic plasticity. We expect that TC assembloids will be particularly useful in this respect, as functional abnormalities in many thalamic nuclei are linked to psychiatric disorders, including schizophrenia.[94] The findings we present here provide a foundation for those future studies.

REFERENCES AND NOTES

1. Ryan, T.J., and Grant, S.G.N. (2009). The origin and evolution of synapses. Preprint, 10.1038/nrn2717 10.1038/nrn2717.

2. Hensch, T.K. (2004). Critical period regulation. Annu Rev Neurosci *27*, 549–579. 10.1146/annurev.neuro.27.070203.144327.

3. Zha, C., and Sossin, W.S. (2022). The molecular diversity of plasticity mechanisms underlying memory: An evolutionary perspective. J Neurochem *163*, 444–460. 10.1111/JNC.15717.

4. Szegedi, V., Paizs, M., Csakvari, E., Molnar, G., Barzo, P., Tamas, G., and Lamsa, K. (2016). Plasticity in single axon glutamatergic connection to GABAergic interneurons regulates complex events in the human neocortex. PLoS Biol *14*. 10.1371/JOURNAL.PBIO.2000237.

5. Ataman, B., Boulting, G.L., Harmin, D.A., Yang, M.G., Baker-Salisbury, M., Yap, E.L., Malik, A.N., Mei, K., Rubin, A.A., Spiegel, I., et al. (2016). Evolution of Osteocrin as an activity-regulated factor in the primate brain. Nature *539*, 242–247. 10.1038/NATURE20111.

6. Qiu, J., McQueen, J., Bilican, B., Dando, O., Magnani, D., Punovuori, K., Selvaraj, B.T., Livesey, M., Haghi, G., Heron, S., et al. (2016). Evidence for evolutionary divergence of activity-dependent gene expression in developing neurons. Elife *5*. 10.7554/ELIFE.20337.

7. Mould, A.W., Hall, N.A., Milosevic, I., and Tunbridge, E.M. (2021). Targeting synaptic plasticity in schizophrenia: insights from genomic studies. Preprint at Elsevier Ltd, 10.1016/j.molmed.2021.07.014 10.1016/j.molmed.2021.07.014.

8. Appelbaum, L.G., Shenasa, M.A., Stolz, L., and Daskalakis, Z. (2023). Synaptic plasticity and mental health: methods, challenges and opportunities. Preprint at Springer Nature, 10.1038/s41386-022-01370-w 10.1038/s41386-022-01370-w.

9. Bourgeron, T. (2015). From the genetic architecture to synaptic plasticity in autism spectrum disorder. Preprint at Nature Publishing Group, 10.1038/nrn3992 10.1038/nrn3992.

10. Goto, Y., Yang, C.R., and Otani, S. (2010). Functional and Dysfunctional Synaptic Plasticity in Prefrontal Cortex: Roles in Psychiatric Disorders. Preprint, 10.1016/j.biopsych.2009.08.026 10.1016/j.biopsych.2009.08.026.

11. Lee, K., Park, T.I.-H., Heppner, P., Schweder, P., Mee, E.W., Dragunow, M., and Montgomery, J.M. (2020). Human in vitro systems for examining synaptic function and plasticity in the brain. J Neurophysiol *123*, 945–965. 10.1152/jn.00411.2019.

12. Kanton, S., Boyle, M.J., He, Z., Santel, M., Weigert, A., Sanchís-Calleja, F., Guijarro, P., Sidow, L., Fleck, J.S., Han, D., et al. (2019). Organoid single-cell genomic atlas uncovers human-specific features of brain development. Nature *574*, 418–422. 10.1038/S41586-019-1654-9.

13. Pollen, A.A., Bhaduri, A., Andrews, M.G., Nowakowski, T.J., Meyerson, O.S., Mostajo-Radji, M.A., Di Lullo, E., Alvarado, B., Bedolli, M., Dougherty, M.L., et al. (2019). Establishing cerebral organoids as models of human-specific brain evolution. Cell *176*, 743-756.e17. 10.1016/J.CELL.2019.01.017.

14. Agoglia, R.M., Sun, D., Birey, F., Yoon, S.J., Miura, Y., Sabatini, K., Paşca, S.P., and Fraser, H.B. (2021). Primate cell fusion disentangles gene regulatory divergence in neurodevelopment. Nature *592*, 421–427. 10.1038/S41586-021-03343-3.

15. Velasco, S., Kedaigle, A.J., Simmons, S.K., Nash, A., Rocha, M., Quadrato, G., Paulsen, B., Nguyen, L., Adiconis, X., Regev, A., et al. (2019). Individual brain organoids reproducibly form cell diversity of the human cerebral cortex. Nature *570*, 523–527. 10.1038/S41586-019-1289-X.

16. Lancaster, M.A., Renner, M., Martin, C.A., Wenzel, D., Bicknell, L.S., Hurles, M.E., Homfray, T., Penninger, J.M., Jackson, A.P., and Knoblich, J.A. (2013). Cerebral organoids model human brain development and microcephaly. Nature *501*, 373–379. 10.1038/NATURE12517.

17. Mora-Bermúdez, F., Badsha, F., Kanton, S., Camp, J.G., Vernot, B., Köhler, K., Voigt, B., Okita, K., Maricic, T., He, Z., et al. (2016). Differences and similarities between human and chimpanzee neural progenitors during cerebral cortex development. Elife *5*. 10.7554/ELIFE.18683.

18. Li, Y., Muffat, J., Omer, A., Bosch, I., Lancaster, M.A., Sur, M., Gehrke, L., Knoblich, J.A., and Jaenisch, R. (2017). Induction of expansion and folding in human cerebral organoids. Cell Stem Cell *20*, 385-396.e3. 10.1016/J.STEM.2016.11.017.

19. Li, C., Fleck, J.S., Martins-Costa, C., Burkard, T.R., Themann, J., Stuempflen, M., Peer, A.M., Vertesy, Á., Littleboy, J.B., Esk, C., et al. (2023). Single-cell brain organoid screening identifies developmental defects in autism. Nature *621*, 373–380. 10.1038/s41586-023-06473-y.

20. Wulansari, N., Darsono, W.H.W., Woo, H.-J., Chang, M.-Y., Kim, J., Bae, E.-J., Sun, W., Lee, J.-H., Cho, I.-J., Shin, H., et al. (2021). Neurodevelopmental defects and neurodegenerative phenotypes in human brain organoids carrying Parkinson's disease-linked DNAJC6 mutations. Sci Adv *7*. 10.1126/sciadv.abb1540.

21. Paulsen, B., Velasco, S., Kedaigle, A.J., Pigoni, M., Quadrato, G., Deo, A.J., Adiconis, X., Uzquiano, A., Sartore, R., Yang, S.M., et al. (2022). Autism genes converge on asynchronous development of shared neuron classes. Nature *602*, 268–273. 10.1038/s41586-021-04358-6.

22. Sebastian, R., Jin, K., Pavon, N., Bansal, R., Potter, A., Song, Y., Babu, J., Gabriel, R., Sun, Y., Aronow, B., et al. (2023). Schizophrenia-associated NRXN1 deletions induce developmental-timing- and cell-type-specific vulnerabilities in human brain organoids. Nat Commun *14*. 10.1038/s41467-023-39420-6.

23. Notaras, M., Lodhi, A., Dündar, F., Collier, P., Sayles, N.M., Tilgner, H., Greening, D., and Colak, D. (2022). Schizophrenia is defined by cell-specific neuropathology and multiple neurodevelopmental mechanisms in patient-derived cerebral organoids. Mol Psychiatry *27*, 1416–1434. 10.1038/s41380-021-01316-6.

24. Liu, C., Fu, Z., Wu, S., Wang, X., Zhang, S., Chu, C., Hong, Y., Wu, W., Chen, S., Jiang, Y., et al. (2022). Mitochondrial HSF1 triggers mitochondrial dysfunction and neurodegeneration in Huntington's disease. EMBO Mol Med *14*. 10.15252/emmm.202215851.

25. Kathuria, A., Lopez-Lengowski, K., Jagtap, S.S., McPhie, D., Perlis, R.H., Cohen, B.M., and Karmacharya, R. (2020). Transcriptomic Landscape and Functional Characterization of Induced Pluripotent Stem Cell-Derived Cerebral Organoids in Schizophrenia. JAMA Psychiatry *77*, 745–754. 10.1001/jamapsychiatry.2020.0196.

26. Magee, J.C., and Grienberger, C. (2020). Synaptic plasticity forms and functions. Annu Rev Neurosci *43*, 95–117. 10.1146/ANNUREV-NEURO-090919-022842.

27. Reha, R.K., Dias, B.G., Nelson, C.A., Kaufer, D., Werker, J.F., Kolbh, B., Levine, J.D., and Hensch, T.K. (2020). Critical period regulation across multiple timescales. Proc Natl Acad Sci U S A *117*, 23242–23251. 10.1073/PNAS.1820836117.

28. Citri, A., and Malenka, R.C. (2008). Synaptic plasticity: multiple forms, functions, and mechanisms. Neuropsychopharmacology *33*, 18–41. 10.1038/SJ.NPP.1301559.

29. Miura, Y., Li, M.-Y., Birey, F., Ikeda, K., Revah, O., Thete, M.V., Park, J.-Y., Puno, A., Lee, S.H., Porteus, M.H., et al. (2020). Generation of human striatal organoids and cortico-striatal assembloids from human pluripotent stem cells. Nat Biotechnol *38*, 1421–1430. 10.1038/s41587-020-00763-w.

30. Jo, J., Xiao, Y., Sun, A.X., Cukuroglu, E., Tran, H.D., Göke, J., Tan, Z.Y., Saw, T.Y., Tan, C.P., Lokman, H., et al. (2016). Midbrain-like Organoids from Human Pluripotent Stem

478 Cells Contain Functional Dopaminergic and Neuromelanin-Producing Neurons. Cell Stem
479 Cell *19*, 248–257. 10.1016/j.stem.2016.07.005.

31. Pasca, A.M., Sloan, S.A., Clarke, L.E., Tian, Y., Makinson, C.D., Huber, N., Kim, C.H.,
     Park, J.Y., O'Rourke, N.A., Nguyen, K.D., et al. (2015). Functional cortical neurons and
     astrocytes from human pluripotent stem cells in 3D culture. Nat Methods *12*, 671–678.
     10.1038/nmeth.3415.

32. Xiang, Y., Tanaka, Y., Cakir, B., Patterson, B., Kim, K.-Y., Sun, P., Kang, Y.-J., Zhong,
     M., Liu, X., Patra, P., et al. (2019). hESC-derived thalamic organoids form reciprocal
     projections when fused with cortical organoids. Cell Stem Cell *24*, 487-497.e7.
     10.1016/j.stem.2018.12.015.

33. Abramson, B.P., and Chalupa, L.M. (1985). The laminar distribution of cortical
     connections with the tecto- and cortico-recipient zones in the cat's lateral posterior
     nucleus. Neuroscience *15*, 81–95. 10.1016/0306-4522(85)90125-3.

34. Rockland, K.S., Andresen, J., Cowie, R.J., and Robinson, D.L. (1999). Single axon
     analysis of pulvinocortical connections to several visual areas in the Macaque. J Comp
     Neurol *406*, 221–250. 10.1002/(SICI)1096-9861(19990405)406:2<221::AID-
     CNE7>3.0.CO;2-K.

35. Halassa, M.M., and Sherman, S.M. (2019). Thalamocortical circuit motifs: a general
     framework. Neuron *103*, 762–770. 10.1016/j.neuron.2019.06.005.

36. Nakajima, M., and Halassa, M.M. (2017). Thalamic control of functional cortical
     connectivity. Curr Opin Neurobiol *44*, 127–131. 10.1016/j.conb.2017.04.001.

37. Briggs, F., and Usrey, W.M. (2008). Emerging views of corticothalamic function. Curr
     Opin Neurobiol *18*, 403–407. 10.1016/J.CONB.2008.09.002.

38. Sherman, S.M. (2016). Thalamus plays a central role in ongoing cortical functioning. Nat
     Neurosci *19*, 533–541. 10.1038/NN.4269.

39. Takahashi, N., Moberg, S., Zolnik, T.A., Catanese, J., Sachdev, R.N.S., Larkum, M.E.,
     and Jaeger, D. (2021). Thalamic input to motor cortex facilitates goal-directed action
     initiation. Curr Biol *31*, 4148-4155.e4. 10.1016/J.CUB.2021.06.089.

40. Hwang, K., Shine, J.M., Cole, M.W., and Sorenson, E. (2022). Thalamocortical
     contributions to cognitive task activity. Elife *11*. 10.7554/ELIFE.81282.

41. Saalmann, Y.B., Pinsk, M.A., Wang, L., Li, X., and Kastner, S. (2012). The pulvinar
     regulates information transmission between cortical areas based on attention demands.
     Science *337*, 753–756. 10.1126/SCIENCE.1223082.

42. Wimmer, R.D., Schmitt, L.I., Davidson, T.J., Nakajima, M., Deisseroth, K., and Halassa,
     M.M. (2015). Thalamic control of sensory selection in divided attention. Nature *526*, 705–
     709. 10.1038/NATURE15398.

43. Biane, J.S., Takashima, Y., Scanziani, M., Conner, J.M., and Tuszynski, M.H. (2016).
     Thalamocortical projections onto behaviorally relevant neurons exhibit plasticity during
     adult motor learning. Neuron *89*, 1173–1179. 10.1016/J.NEURON.2016.02.001.

44. Audette, N.J., Bernhard, S.M., Ray, A., Stewart, L.T., and Barth, A.L. (2019). Rapid
     plasticity of higher-order thalamocortical inputs during sensory learning. Neuron *103*, 277-
     291.e4. 10.1016/J.NEURON.2019.04.037.

45. Scott, G.A., Liu, M.C., Tahir, N.B., Zabder, N.K., Song, Y., Greba, Q., and Howland, J.G.
     (2020). Roles of the medial prefrontal cortex, mediodorsal thalamus, and their combined
     circuit for performance of the odor span task in rats: analysis of memory capacity and
     foraging behavior. Learn Mem *27*, 67–77. 10.1101/LM.050195.119.

46. Phillips, J.M., Kambi, N.A., Redinbaugh, M.J., Mohanta, S., and Saalmann, Y.B. (2021).
     Disentangling the influences of multiple thalamic nuclei on prefrontal cortex and cognitive
     control. Neurosci Biobehav Rev *128*, 487–510. 10.1016/J.NEUBIOREV.2021.06.042.

47. Blundon, J.A., Roy, N.C., Teubner, B.J.W., Yu, J., Eom, T.-Y., Sample, K.J.J., Pani, A.,
     Smeyne, R.J., Han, S.B., Kerekes, R.A., et al. (2017). Restoring auditory cortex plasticity

in adult mice by restricting thalamic adenosine signaling. Science *356*, 1352–1356. 10.1126/science.aaf4612.

48. Vue, T.Y., Aaker, J., Taniguchi, A., Kazemzadeh, C., Skidmore, J.M., Martin, D.M., Martin, J.F., Treier, M., and Nakagawa, Y. (2007). Characterization of progenitor domains in the developing mouse thalamus. J Comp Neurol *505*, 73–91. 10.1002/cne.21467.

49. Rai, M., Coleman, Z., Curley, M., Nityanandam, A., Platt, A., Robles-Murguia, M., Jiao, J., Finkelstein, D., Wang, Y.D., Xu, B., et al. (2021). Proteasome stress in skeletal muscle mounts a long-range protective response that delays retinal and brain aging. *33*, 1137-1154.e9. 10.1016/J.CMET.2021.03.005.

50. Herrmann, K., Antonini, A., and Shatz, C.J. (1994). Ultrastructural evidence for synaptic interactions between thalamocortical axons and subplate neurons. European Journal of Neuroscience *6*, 1729–1742. 10.1111/j.1460-9568.1994.tb00565.x.

51. Polioudakis, D., de la Torre-Ubieta, L., Langerman, J., Elkins, A.G., Shi, X., Stein, J.L., Vuong, C.K., Nichterwitz, S., Gevorgian, M., Opland, C.K., et al. (2019). A single-cell transcriptomic atlas of human neocortical development during mid-gestation. Neuron *103*, 785-801.e8. 10.1016/j.neuron.2019.06.011.

52. Kanold, P.O., and Luhmann, H.J. (2010). The subplate and early cortical circuits. Annu Rev Neurosci *33*, 23–48. 10.1146/annurev-neuro-060909-153244.

53. De Gois, S., Schäfer, M.K.-H., Defamie, N., Chen, C., Ricci, A., Weihe, E., Varoqui, H., and Erickson, J.D. (2005). Homeostatic scaling of vesicular glutamate and GABA transporter expression in rat neocortical circuits. The Journal of Neuroscience *25*, 7121–7133. 10.1523/JNEUROSCI.5221-04.2005.

54. Zhang, Q., Lee, W.-C.A., Paul, D.L., and Ginty, D.D. (2019). Multiplexed peroxidase-based electron microscopy labeling enables simultaneous visualization of multiple cell types. Nat Neurosci *22*, 828–839. 10.1038/s41593-019-0358-7.

55. Beierlein, M., and Connors, B.W. (2002). Short-term dynamics of thalamocortical and intracortical synapses onto layer 6 neurons in neocortex. J Neurophysiol *88*, 1924–1932.

56. Rose, H.J., and Metherate, R. (2005). Auditory thalamocortical transmission is reliable and temporally precise. J.Neurophysiol. *94*, 2019–2030.

57. Bayazitov, I.T., Westmoreland, J.J., and Zakharenko, S.S. (2013). Forward suppression in the auditory cortex is caused by the Ca3.1 calcium channel-mediated switch from bursting to tonic firing at thalamocortical projections. Journal of Neuroscience *33*, 18940–18950. 10.1523/JNEUROSCI.3335-13.2013.

58. Viaene, A.N., Petrof, I., and Sherman, S.M. (2011). Properties of the thalamic projection from the posterior medial nucleus to primary and secondary somatosensory cortices in the mouse. Proc Natl Acad Sci U S A *108*, 18156–18161. 10.1073/pnas.1114828108.

59. Gil, Z., Connors, B.W., and Amitai, Y. (1999). Efficacy of thalamocortical and intracortical synaptic connections: quanta, innervation, and reliability. Neuron *23*, 385–397.

60. Stratford, K.J., Tarczy-Hornoch, K., Martin, K.A., Bannister, N.J., and Jack, J.J. (1996). Excitatory synaptic inputs to spiny stellate cells in cat visual cortex. Nature *382*, 258–261.

61. Beierlein, M., Fall, C.P., Rinzel, J., and Yuste, R. (2002). Thalamocortical bursts trigger recurrent activity in neocortical networks: layer 4 as a frequency-dependent gate. J Neurosci *22*, 9885–9894.

62. Zhao, W., Johnston, K.G., Ren, H., Xu, X., and Nie, Q. (2023). Inferring neuron-neuron communications from single-cell transcriptomics through NeuronChat. Nat Commun *14*. 10.1038/s41467-023-36800-w.

63. Richardson, R.J., Blundon, J.A., Bayazitov, I.T., and Zakharenko, S.S. (2009). Connectivity patterns revealed by mapping of active inputs on dendrites of thalamorecipient neurons in the auditory cortex. Journal of Neuroscience *29*, 6406–6417. 10.1523/JNEUROSCI.0258-09.2009.

64. Debanne, D., and Inglebert, Y. (2023). Spike timing-dependent plasticity and memory. Curr Opin Neurobiol *80*, 102707. 10.1016/j.conb.2023.102707.

65. Feldman, D.E. (2012). The spike-timing dependence of plasticity. Neuron *75*, 556–571. 10.1016/j.neuron.2012.08.001.

66. Dan, Y., and Poo, M.M. (2004). Spike timing-dependent plasticity of neural circuits. Neuron *44*, 23–30.

67. Collingridge, G.L., Peineau, S., Howland, J.G., and Wang, Y.T. (2010). Long-term depression in the CNS. Nat.Rev.Neurosci *11*, 459–473. 10.1038/NRN2867.

68. Zafeiriou, M.P., Bao, G., Hudson, J., Halder, R., Blenkle, A., Schreiber, M.K., Fischer, A., Schild, D., and Zimmermann, W.H. (2020). Developmental GABA polarity switch and neuronal plasticity in Bioengineered Neuronal Organoids. Nat Commun *11*. 10.1038/s41467-020-17521-w.

69. Muguruma, K., Nishiyama, A., Kawakami, H., Hashimoto, K., and Sasai, Y. (2015). Self-organization of polarized cerebellar tissue in 3D culture of human pluripotent stem cells. Cell Rep *10*, 537–550. 10.1016/j.celrep.2014.12.051.

70. Sakaguchi, H., Kadoshima, T., Soen, M., Narii, N., Ishida, Y., Ohgushi, M., Takahashi, J., Eiraku, M., and Sasai, Y. (2015). Generation of functional hippocampal neurons from self-organizing human embryonic stem cell-derived dorsomedial telencephalic tissue. Nat Commun *6*. 10.1038/ncomms9896.

71. Mariani, J., Coppola, G., Zhang, P., Abyzov, A., Provini, L., Tomasini, L., Amenduni, M., Szekely, A., Palejev, D., Wilson, M., et al. (2015). FOXG1-Dependent Dysregulation of GABA/Glutamate Neuron Differentiation in Autism Spectrum Disorders. Cell *162*, 375–390. 10.1016/j.cell.2015.06.034.

72. Lüscher, C., and Malenka, R.C. (2012). NMDA receptor-dependent long-term potentiation and long-term depression (LTP/LTD). Cold Spring Harb Perspect Biol *4*. 10.1101/cshperspect.a005710.

73. Pisani, A., Gubellini, P., Bonsi, P., Conquet, F., Picconi, B., Centonze, D., Bernardi, G., and Calabresi, P. (2001). Metabotropic glutamate receptor 5 mediates the potentiation of N-methyl-D-aspartate responses in medium spiny striatal neurons. Neuroscience *106*, 579–587. 10.1016/s0306-4522(01)00297-4.

74. Benquet, P., Gee, C.E., and Gerber, U. (2002). Two distinct signaling pathways upregulate NMDA receptor responses via two distinct metabotropic glutamate receptor subtypes. J Neurosci *22*, 9679–9686. 10.1523/JNEUROSCI.22-22-09679.2002.

75. Heidinger, V., Manzerra, P., Wang, X.Q., Strasser, U., Yu, S.-P., Choi, D.W., and Behrens, M.M. (2002). Metabotropic glutamate receptor 1-induced upregulation of NMDA receptor current: mediation through the Pyk2/Src-family kinase pathway in cortical neurons. J Neurosci *22*, 5452–5461. 10.1523/JNEUROSCI.22-13-05452.2002.

76. Fox, K., Schlaggar, B.L., Glazewski, S., and O'Leary, D.D. (1996). Glutamate receptor blockade at cortical synapses disrupts development of thalamocortical and columnar organization in somatosensory cortex. Proc Natl Acad Sci U S A *93*, 5584–5589. 10.1073/pnas.93.11.5584.

77. Schlaggar, B.L., Fox, K., and O'Leary, D.D. (1993). Postsynaptic control of plasticity in developing somatosensory cortex. Nature *364*, 623–626. 10.1038/364623a0.

78. Itami, C., Huang, J.-Y., Yamasaki, M., Watanabe, M., Lu, H.-C., and Kimura, F. (2016). Developmental switch in spike timing-dependent plasticity and cannabinoid-dependent reorganization of the thalamocortical projection in the barrel Cortex. J Neurosci *36*, 7039–7054. 10.1523/JNEUROSCI.4280-15.2016.

79. Crair, M.C., and Malenka, R.C. (1995). A critical period for long-term potentiation at thalamocortical synapses. Nature *375*, 325–328. 10.1038/375325a0.

80. Chun, S., Bayazitov, I.T., Blundon, J.A., and Zakharenko, S.S. (2013). Thalamocortical long-term potentiation becomes gated after the early critical period in the auditory cortex. Journal of Neuroscience *33*, 7345–7357. 10.1523/JNEUROSCI.4500-12.2013.

81. Castro-Alamancos, M.A., and Calcagnotto, M.E. (1999). Presynaptic long-term potentiation in corticothalamic synapses. J Neurosci *19*, 9090–9097. 10.1523/JNEUROSCI.19-20-09090.1999.

82. Feldman, D.E., Nicoll, R.A., Malenka, R.C., and Isaac, J.T. (1998). Long-term depression at thalamocortical synapses in developing rat somatosensory cortex. Neuron *21*, 347–357. 10.1016/s0896-6273(00)80544-9.

83. Pisani, A., Gubellini, P., Bonsi, P., Conquet, F., Picconi, B., Centonze, D., Bernardi, G., and Calabresi, P. (2001). Metabotropic glutamate receptor 5 mediates the potentiation of N-methyl-D-aspartate responses in medium spiny striatal neurons. Neuroscience *106*, 579–587. 10.1016/s0306-4522(01)00297-4.

84. Benquet, P., Gee, C.E., and Gerber, U. (2002). Two distinct signaling pathways upregulate NMDA receptor responses via two distinct metabotropic glutamate receptor subtypes. J Neurosci *22*, 9679–9686. 10.1523/JNEUROSCI.22-22-09679.2002.

85. Heidinger, V., Manzerra, P., Wang, X.Q., Strasser, U., Yu, S.-P., Choi, D.W., and Behrens, M.M. (2002). Metabotropic glutamate receptor 1-induced upregulation of NMDA receptor current: mediation through the Pyk2/Src-family kinase pathway in cortical neurons. J Neurosci *22*, 5452–5461. 10.1523/JNEUROSCI.22-13-05452.2002.

86. Amiri, A., Coppola, G., Scuderi, S., Wu, F., Roychowdhury, T., Liu, F., Pochareddy, S., Shin, Y., Safi, A., Song, L., et al. (2018). Transcriptome and epigenome landscape of human cortical development modeled in organoids. Science *362*. 10.1126/science.aat6720.

87. Kiral, F.R., Cakir, B., Tanaka, Y., Kim, J., Yang, W.S., Wehbe, F., Kang, Y.-J., Zhong, M., Sancer, G., Lee, S.-H., et al. (2023). Generation of ventralized human thalamic organoids with thalamic reticular nucleus. Cell Stem Cell *30*, 677-688.e5. 10.1016/j.stem.2023.03.007.

88. Monko, T., Rebertus, J., Stolley, J., Salton, S.R., and Nakagawa, Y. (2022). Thalamocortical axons regulate neurogenesis and laminar fates in the early sensory cortex. Proc Natl Acad Sci U S A *119*, e2201355119. 10.1073/pnas.2201355119.

89. Antón-Bolaños, N., Sempere-Ferràndez, A., Guillamón-Vivancos, T., Martini, F.J., Pérez-Saiz, L., Gezelius, H., Filipchuk, A., Valdeolmillos, M., and López-Bendito, G. (2019). Prenatal activity from thalamic neurons governs the emergence of functional cortical maps in mice. Science *364*, 987–990. 10.1126/science.aav7617.

90. Miura, Y., Li, M.-Y., Birey, F., Ikeda, K., Revah, O., Thete, M.V., Park, J.-Y., Puno, A., Lee, S.H., Porteus, M.H., et al. (2020). Generation of human striatal organoids and cortico-striatal assembloids from human pluripotent stem cells. Nat Biotechnol *38*, 1421–1430. 10.1038/s41587-020-00763-w.

91. Birey, F., Andersen, J., Makinson, C.D., Islam, S., Wei, W., Huber, N., Fan, H.C., Metzler, K.R.C., Panagiotakos, G., Thom, N., et al. (2017). Assembly of functionally integrated human forebrain spheroids. Nature *545*, 54–59. 10.1038/nature22330.

92. Bagley, J.A., Reumann, D., Bian, S., Lévi-Strauss, J., and Knoblich, J.A. (2017). Fused cerebral organoids model interactions between brain regions. Nat Methods *14*, 743–751. 10.1038/nmeth.4304.

93. Andersen, J., Revah, O., Miura, Y., Thom, N., Amin, N.D., Kelley, K.W., Singh, M., Chen, X., Thete, M.V., Walczak, E.M., et al. (2020). Generation of functional human 3D cortico-motor assembloids. Cell *183*, 1913-1929.e26. 10.1016/j.cell.2020.11.017.

94. Jiang, Y., Patton, M.H., and Zakharenko, S.S. (2021). A case for thalamic mechanisms of schizophrenia: perspective from modeling 22q11.2 deletion syndrome. Front Neural Circuits *15*. 10.3389/fncir.2021.769969.

95. Sloan, S.A., Darmanis, S., Huber, N., Khan, T.A., Birey, F., Caneda, C., Reimer, R., Quake, S.R., Barres, B.A., and Paşca, S.P. (2017). Human astrocyte maturation captured in 3D cerebral cortical spheroids derived from pluripotent stem cells. Neuron *95*, 779-790.e6. 10.1016/j.neuron.2017.07.035.

96. Tsankov, A.M., Akopian, V., Pop, R., Chetty, S., Gifford, C.A., Daheron, L., Tsankova,
N.M., and Meissner, A. (2015). A qPCR ScoreCard quantifies the differentiation potential
of human pluripotent stem cells. Nat Biotechnol *33*, 1182–1192. 10.1038/nbt.3387.
97. Assou, S., Girault, N., Plinet, M., Bouckenheimer, J., Sansac, C., Combe, M., Mianné, J.,
Bourguignon, C., Fieldes, M., Ahmed, E., et al. (2020). Recurrent genetic abnormalities in
human pluripotent stem cells: definition and routine detection in culture supernatant by
targeted droplet digital PCR. Stem Cell Reports *14*, 1–8. 10.1016/j.stemcr.2019.12.004.
98. Baker, D., Hirst, A.J., Gokhale, P.J., Juarez, M.A., Williams, S., Wheeler, M., Bean, K.,
Allison, T.F., Moore, H.D., Andrews, P.W., et al. (2016). Detecting genetic mosaicism in
cultures of human pluripotent stem cells. Stem Cell Reports *7*, 998–1012.
10.1016/j.stemcr.2016.10.003.
99. Martins-Taylor, K., Nisler, B.S., Taapken, S.M., Compton, T., Crandall, L., Montgomery,
K.D., Lalande, M., and Xu, R.-H. (2011). Recurrent copy number variations in human
induced pluripotent stem cells. Nat Biotechnol *29*, 488–491. 10.1038/nbt.1890.
100. Nishino, K., Toyoda, M., Yamazaki-Inoue, M., Fukawatase, Y., Chikazawa, E.,
Sakaguchi, H., Akutsu, H., and Umezawa, A. (2011). DNA methylation dynamics in
human induced pluripotent stem cells over time. PLoS Genet *7*, e1002085.
10.1371/journal.pgen.1002085.
101. Johannesson, B., Sagi, I., Gore, A., Paull, D., Yamada, M., Golan-Lev, T., Li, Z., LeDuc,
C., Shen, Y., Stern, S., et al. (2014). Comparable frequencies of coding mutations and
loss of imprinting in human pluripotent cells derived by nuclear transfer and defined
factors. Cell Stem Cell *15*, 634–642. 10.1016/j.stem.2014.10.002.
102. Norrie, J.L., Nityanandam, A., Lai, K., Chen, X., Wilson, M., Stewart, E., Griffiths, L., Jin,
H., Wu, G., Orr, B., et al. (2021). Retinoblastoma from human stem cell-derived retinal
organoids. Nat Commun *12*, 4535. 10.1038/s41467-021-24781-7.
103. Sentmanat, M.F., Peters, S.T., Florian, C.P., Connelly, J.P., and Pruett-Miller, S.M.
(2018). A survey of validation strategies for CRISPR-Cas9 editing. Sci Rep *8*, 888.
10.1038/s41598-018-19441-8.
104. Connelly, J.P., and Pruett-Miller, S.M. (2019). CRIS.py: a versatile and high-throughput
analysis program for CRISPR-based genome editing. Sci Rep *9*, 4194.
doi.org/10.1038/s41598-019-40896-w.
105. Chen, Y., Tristan, C.A., Chen, L., Jovanovic, V.M., Malley, C., Chu, P.-H., Ryu, S., Deng,
T., Ormanoglu, P., Tao, D., et al. (2021). A versatile polypharmacology platform promotes
cytoprotection and viability of human pluripotent and differentiated cells. Nat Methods *18*,
528–541. 10.1038/s41592-021-01126-2.
106. Keaveney, M.K., Tseng, H.-A., Ta, T.L., Gritton, H.J., Man, H.-Y., and Han, X. (2018). A
microRNA-based gene-targeting tool for virally labeling interneurons in the rodent cortex.
Cell Rep *24*, 294–303. 10.1016/j.celrep.2018.06.049.
107. Miller, J.D., Ganat, Y.M., Kishinevsky, S., Bowman, R.L., Liu, B., Tu, E.Y., Mandal, P.K.,
Vera, E., Shim, J., Kriks, S., et al. (2013). Human iPSC-based modeling of late-onset
disease via progerin-induced aging. Cell Stem Cell *13*, 691–705.
10.1016/j.stem.2013.11.006.
108. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*, 15–21. 10.1093/bioinformatics/bts635.
109. Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq
data with or without a reference genome. BMC Bioinformatics *12*, 323. 10.1186/1471-
2105-12-323.
110. Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics *26*, 139–140. 10.1093/bioinformatics/btp616.

111. Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015).
limma powers differential expression analyses for RNA-sequencing and microarray
studies. Nucleic Acids Res *43*, e47. 10.1093/nar/gkv007.
112. Fleck, J.S., Sanchís-Calleja, F., He, Z., Santel, M., Boyle, M.J., Camp, J.G., and
Treutlein, B. (2021). Resolving organoid brain region identities by mapping single-cell
genomic data to reference atlases. Cell Stem Cell *28*, 1148-1159.e8.
10.1016/j.stem.2021.02.015.
113. Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., and Vilo, J.
(2019). g:Profiler: a web server for functional enrichment analysis and conversions of
gene lists (2019 update). Nucleic Acids Res *47*, W191–W198. 10.1093/NAR/GKZ369.
114. Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis (Springer-Verlag New
York).
115. Gu, Z., Eils, R., and Schlesner, M. (2016). Complex heatmaps reveal patterns and
correlations in multidimensional genomic data. Bioinformatics *32*, 2847–2849.
10.1093/bioinformatics/btw313.
116. Gu, Z. (2022). Complex heatmap visualization. iMeta *1*. 10.1002/imt2.43.
117. Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using
real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods *25*, 402–408.
10.1006/meth.2001.1262.
118. Matson, K.J.E., Sathyamurthy, A., Johnson, K.R., Kelly, M.C., Kelley, M.W., and Levine,
A.J. (2018). Isolation of adult spinal cord nuclei for massively parallel single-nucleus RNA
sequencing. J Vis Exp. 10.3791/58413.
119. Young, M.D., and Behjati, S. (2020). SoupX removes ambient RNA contamination from
droplet-based single-cell RNA sequencing data. Gigascience *9*.
10.1093/gigascience/giaa151.
120. Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J.,
Wilk, A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell
data. Cell *184*, 3573-3587.e29. 10.1016/j.cell.2021.04.048.
121. Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., Hao, Y.,
Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive integration of single-cell
data. Cell *177*, 1888-1902.e21. 10.1016/j.cell.2019.05.031.
122. Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-
cell transcriptomic data across different conditions, technologies, and species. Nat
Biotechnol *36*, 411–420. 10.1038/nbt.4096.
123. Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., and Regev, A. (2015). Spatial
reconstruction of single-cell gene expression data. Nat Biotechnol *33*, 495–502.
10.1038/nbt.3192.
124. Tirosh, I., Izar, B., Prakadan, S.M., Wadsworth, M.H., Treacy, D., Trombetta, J.J., Rotem,
A., Rodman, C., Lian, C., Murphy, G., et al. (2016). Dissecting the multicellular
ecosystem of metastatic melanoma by single-cell RNA-seq. Science *352*, 189–196.
10.1126/science.aad0501.
125. Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y.,
Brenner, M., Loh, P.-R., and Raychaudhuri, S. (2019). Fast, sensitive and accurate
integration of single-cell data with Harmony. Nat Methods *16*, 1289–1296.
10.1038/s41592-019-0619-0.
126. Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N.J.,
Livak, K.J., Mikkelsen, T.S., and Rinn, J.L. (2014). The dynamics and regulators of cell
fate decisions are revealed by pseudotemporal ordering of single cells. Nat Biotechnol
32, 381–386. 10.1038/nbt.2859.
127. Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H.A., and Trapnell, C. (2017).
Reversed graph embedding resolves complex single-cell trajectories. Nat Methods *14*,
979–982. 10.1038/nmeth.4402.

128. Qiu, X., Hill, A., Packer, J., Lin, D., Ma, Y.-A., and Trapnell, C. (2017). Single-cell mRNA
787   quantification and differential analysis with Census. Nat Methods *14*, 309–315.
788   10.1038/nmeth.4150.
789   129. Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D.M., Hill, A.J., Zhang, F., Mundlos,
790   S., Christiansen, L., Steemers, F.J., et al. (2019). The single-cell transcriptional
791   landscape of mammalian organogenesis. Nature *566*, 496–502. 10.1038/s41586-019-
792   0969-x.
793   130. Govek, K.W., Chen, S., Sgourdou, P., Yao, Y., Woodhouse, S., Chen, T., Fuccillo, M. V,
794   Epstein, D.J., and Camara, P.G. (2022). Developmental trajectories of thalamic
795   progenitors revealed by single-cell transcriptome profiling and Shh perturbation. Cell Rep
796   *41*, 111768. 10.1016/j.celrep.2022.111768.
797   131. Guo, Q., and Li, J.Y.H. (2019). Defining developmental diversification of diencephalon
798   neurons through single cell gene expression profiling. Development *146*.
799   10.1242/dev.174284.
800   132. Vue, T.Y., Aaker, J., Taniguchi, A., Kazemzadeh, C., Skidmore, J.M., Martin, D.M.,
801   Martin, J.F., Treier, M., and Nakagawa, Y. (2007). Characterization of progenitor domains
802   in the developing mouse thalamus. J Comp Neurol *505*, 73–91. 10.1002/cne.21467.
803   133. Martell, J.D., Deerinck, T.J., Lam, S.S., Ellisman, M.H., and Ting, A.Y. (2017). Electron
804   microscopy using the genetically encoded APEX2 tag in cultured mammalian cells. Nat
805   Protoc *12*, 1792–1816. 10.1038/nprot.2017.065.
806   134. Garad, M., Edelmann, E., and Leßmann, V. (2021). Impairment of spike-timing-
807   dependent plasticity at Schaffer collateral-CA1 synapses in adult APP/PS1 mice depends
808   on proximity of Aβ plaques. Int J Mol Sci *22*. 10.3390/ijms22031378.
809   135. Blundon, J.A., Bayazitov, I.T., and Zakharenko, S.S. (2011). Presynaptic gating of
810   postsynaptically expressed plasticity at mature thalamocortical synapses. Journal of
811   Neuroscience *31*, 16012–16025. 10.1523/JNEUROSCI.3281-11.2011.
812
813

Acknowledgments: We thank Yiping Fan, Dale Hedges, and Daniel Estevez Prado (St. Jude

Center for Applied Bioinformatics, Transcriptomics Group) for assistance with bulk RNA-seq data

analysis; Lawrence Reiter for providing the TP-190a and TP-189 dental pulp stem cells for

reprogramming; Sergiu Pasca for providing the 2242, 1205, and 8858 hiPSC lines; Angela

McArthur for manuscript editing; and Zakharenko lab members for constructive comments.

Funding: This work was funded, in part, by the National Institutes of Health grants R01

MH097742 and R01 DC012833 (to SSZ), K99 MH129617 (to MHP), the Stanford Maternal and

Child Health Research Institute Uytengsu-Hamilton 22q11 Neuropsychiatry Research Program

grants UH22QEXTFY21 and UH22QEXTFY23 (to SSZ), the National Cancer Institute grant P30

CA021765, and the American Lebanese Syrian Associated Charities (ALSAC). The content is

solely the responsibility of the authors and does not necessarily represent the official views of

the National Institutes of Health or other granting agencies.

Author contributions:

Conceptualization: MHP, KTT, SSZ

Electrophysiology experiments: MHP, ITB, YJ

Molecular biology experiments: KTT, AN, ABS

Two-photon imaging experiments: ITB

hiPSC line maintenance, organoid differentiation, and assembloid preparation: AN, KDN

Electron microscopy: NBK, CGR

snRNA-seq sample preparation: AJT, JBB

snRNA-seq data analysis: CAR, KTT

Design and production of hiPSC reporter lines: STP, SMP-M

Visualization: MHP, KTT

Funding acquisition: SSZ

Supervision: SSZ

Writing – original draft: MHP, KTT

Writing – review & editing: SSZ

Competing interests: The authors declare no competing interests.

Data and materials availability: The snRNA-seq data that support the findings of this study

have been deposited in SRA under the Bioproject ID PRJNA999219. Bulk RNA-seq data have

been deposited in SRA under Bioproject ID PRJNA1001283. Code used for the analysis of

snRNA-seq data is available at Github

(https://github.com/ZakharenkoLab/Thalamic_and_Cortical_Organoid_snRNASeq). Additional R

code is available upon request.

Supplemental Materials

Figures S1 to S7

Table S1

Figures

Figure 1. hThOs contain functional glutamatergic thalamic neurons.

(**A**) Reporter cell line validation for hThOs. Top: Schematic of *TCF7L2* exon 1 in the TP-190a-TCF7L2-tdTomato reporter line, which was used to generate all hThOs, except where indicated in Figure S2. Bottom: VoxHunt deconvolution analysis of bulk RNA-seq data from D69-D70 hThOs using E13 mouse brain data from the *Allen Brain Atlas* as a reference.

(**B**) Immunofluorescence images of TCF7L2, TUBB3, OTX2, and SOX2 labeling in D60 hThOs. Nuclei are indicated by DAPI (cyan). TCF7L2-tdTomato fluorescence is indicated in magenta.

Images were acquired from serial sections of the same organoid. Scale bars: 200 μm (whole
section), 100 μm (insets).
(**C**) UMAP plot with cluster annotations indicated by color.
(**D**) VoxHunt analysis by snRNA-seq cell cluster. Excitatory neuron (ExN) clusters exhibit the
highest correlations with *BrainSpan* samples from human mediodorsal nucleus of the thalamus
(MD), aged 13–24 pcw. Cluster annotations are indicated by color on the x-axis.
(**E**) Bar plot showing the number of nuclei per cell cluster, with clusters indicated by fill color.
(**F**) UMAP plots of glutamatergic markers *SLC17A6* (*VGLUT2*) and *SLC17A7* (*VGLUT1*). Color
indicates normalized transcript level.
(**G**) VoxHunt correlation analysis mapping clusters ExN1-4 onto the E15 mouse brain.
(**H**) VoxHunt correlation analysis mapping the PT/ZLI/rTh cluster onto the E15 mouse brain.
(**I**) UMAP plots of the PT/ZLI/rTh cluster, demonstrating the expression of markers associated
with the PT, ZLI, and rTh. Transcript information is indicated by color. The relative locations of
these structures within the developing diencephalon are shown in the schematic.
(**J**) Example traces showing voltage and AP responses to current injections in cells recorded
from an hThO.
(**K**) Pseudotime ordering of cells within the hThOs.
(**L**) UMAP plot of the neural progenitor marker *TNC*. Color indicates the normalized transcript
level.
(**M-N**) UMAP plots of cell cycle analysis results for the Cycling Progenitor, Radial Glia, and Glia
cell clusters. Color indicates S Score (**M**) or G2M Score (**N**).
(**O**) UMAP plot of the astrocyte marker *GFAP* in the Cycling Progenitor, Radial Glia, and Glia
cell clusters. Color indicates the normalized transcript level.
Data in (**C-I**) and (**K-O**) were produced by snRNA-seq analysis of 15,363 nuclei from D90
hThOs.
See also **Figures S1-S3** for additional data validating hiPSC lines and hThOs. See **Figure S4**
for additional data related to electrophysiological properties and synapses in hThOs.
893
894

895
896 **Figure 2. Fusing hThOs and hCOs produces assembloids with functional synapses.**
897 (**A**) Reporter line validation for hCOs. Top: Schematic of *SLC17A7* (*VGLUT1*) exon 12 in the
898 TP-190a-VGLUT1-tdTomato reporter line, which was used to generate all hCOs. Bottom:
899 VoxHunt deconvolution analysis of bulk RNA-seq data from D70 hCOs using E13 mouse brain
900 data from the *Allen Brain Atlas* as a reference. Organoids were visually categorized as positive

or negative for tdTomato fluorescence prior to sequencing. The *tdTomato* RNA level for each sample is indicated in TPM (transcripts per million). Each stacked bar indicates one bulk RNA-seq sample derived from 2-3 pooled organoids.

(**B**) The snRNA-seq analysis of hCOs. Left: UMAP plot with cluster annotations. ExN: excitatory neuron, DL: deep layer, UL: upper layer, Un.: unknown. Right: Dot plot showing subplate marker expression by cluster. Avg Exp: normalized average expression, % Cells: percentage of cells expressing a marker within a cluster.

(**C**) VoxHunt analysis mapping hCOs (all clusters) onto the E15 mouse brain.

(**D**) Pseudotime analysis of the neural cell trajectory (Cycling Progenitors to UL ExNs, DL ExNs, and Subplate/DL ExNs) from hCOs.

(**E**) UMAP plots of glutamatergic markers *SLC17A6* (*VGLUT2*) and *SLC17A7* (*VGLUT1*). Color indicates normalized transcript level.

(**F**) Traces showing the voltage and AP responses in cells recorded from an hCO.

(**G**) Fluorescence and bright field image of a TC assembloid at 5 days postfusion (dpf).

(**H**) Representative fluorescence images for 2-dimensional fusion assay. Thalamic neurons (magenta, right chamber) and cortical neurons (green, left chamber) extend processes from their respective chambers, across the barrier region (dashed yellow lines), and into the opposite chamber starting at D9. Elaborate processes extending from the opposite sides can be seen in both halves by D61.

(**I**) Fluorescence image of an hCO co-transduced with hSyn-GFP and hSyn-V5-Mito-APEX2 lentiviruses.

(**J**) Schematic and TEM image of an APEX2$^+$ mitochondrion (circled in magenta) in a TC synapse. Pre: presynaptic compartment, post: postsynaptic compartment.

(**K**) Schematic and TEM image of an APEX2$^+$ mitochondrion (circled in green) in a CT synapse. APEX2$^-$ mitochondria are indicated by asterisks (*).

(**L**) Left: Schematic of the recording configuration for the TC pathway. Right: Bar graph of the percentage of responsive (green) and unresponsive (gray) cells in 11 assembloids. The numbers of cells recorded per assembloid are shown in the bars.

(**M**) Left: Schematic of the recording configuration for the CT pathway. Middle: The percentages of hThO cells that responded (magenta) or did not respond (gray) to hCO stimulation across 10 assembloids. Right: Bar graph of the average percentage of responsive cells for TC and CT synapses, based on (**G**) and (**H**).

(**N**) Line graph of PPRs across five interstimulus intervals (ISIs) in CT (magenta) and TC (green) synapses [one-sample *t*-test: μ = 1, #p <0.05, ##p <0.01, n = 18-23 cells/9-13 assembloids (TC), n = 8-24/7-12 (CT)]. Differences between CT and TC synapses were evaluated by unpaired *t*-test (**p <0.01). Inset: Sample traces depicting PPRs in CT and TC synapses. Circles represent stimulus artifacts.

Grouped data are shown as mean ± SEM. Scale bars (**F**): 10 mV, 2.5 s. Scale bars (**N**): 20 pA, 200 ms.

See also **Figures S1 and S5** for additional data validating the TP-190a-VGLUT1-tdTomato reporter line and hCOs, respectively. See **Figure S4** for additional data related to electrophysiological properties and synapses in hCOs.

Figure 3. Assembloids contain glutamatergic TC and CT synapses.

(**A**) NeuronChat predictions showing communication between cell clusters containing glutamatergic neurons in hThOs and other cell clusters in hCOs and hThOs. Probability of communication (Prob) is indicated in the heatmap (bottom). Interactions with a Prob >0.6 are shown in the circle plot (top).

(**B**) NeuronChat predictions showing communication between cell clusters containing glutamatergic neurons in hCOs and other cell clusters in hCOs and hThOs. Probability is indicated in the heatmap (bottom). Interactions with a Prob >0.5 are shown in the circle plot (top).

(**C**) Heatmap showing the probability of NRXN-mediated communication between cell clusters in hCOs and hThOs. Predictions were made using NeuronChat.

(**D**) Heatmap showing the probability of glutamatergic communication between cell clusters in hCOs and hThOs. Predictions were made using NeuronChat.

(**E**) Average TC EPSC amplitude [holding potential (Vh) –70 mV] in the presence of NBQX (3 mM) is significantly decreased compared to control aCSF conditions (paired *t*-test: **p = 0.009, n = 5 cells/2 assembloids).

(**F**) The average TC EPSP amplitude (Vh +40 mV) in the presence of NBQX and AP5 (50 µM) is significantly lower than in control aCSF (paired *t*-test: *p = 0.038, n = 5 cells/3 assembloids).

(**G**) Traces of evoked TC AMPAR- and NMDAR-mediated currents in control aCSF and in the presence of NBQX or NBQX and AP5, respectively. Circle represents a stimulus artifact.

(**H**) Average CT EPSC amplitude (Vh –70 mV) in the presence of NBQX is significantly decreased compared to control aCSF conditions (paired *t*-test: *p = 0.012, n = 5 cells/3 assembloids).

(**I**) The average CT EPSC amplitudes (Vh +40 mV) are significantly reduced in the presence of NBQX and AP5 compared to control aCSF (paired *t*-test: ***p = 0.0006, n = 5 cells/3 assembloids).

(**J**) Example traces of evoked CT AMPAR- and NMDAR-mediated currents in aCSF and in the presence of NBQX or NBQX and AP5, respectively.

(**K**) Schematics of two-photon calcium imaging in postsynaptic dendritic spines of hCO cells upon hThO stimulation. Alexa Fluor 594: AF-594 (R), magenta; Fluo-5F (G), green.

(**L**) Image of a dendrite of an hCO cell. Line scans (white line) were performed across a dendritic spine (Sp) and parent dendritic shaft (Sh).

(**M**) Left: Representative changes in G/R of Sp and Sh responses over time to a single synaptic stimulation (arrowhead and black line). Right: Representative line scans of Sp (light gray) and Sh (dark gray).

(**N**) Average changes in synaptically evoked G/R (paired *t*-test: **p = 0.002, n = 9 cells/4 assembloids).

(**O**) Average changes in synaptically evoked Sp G/R in aCSF and in the presence of AP5 (paired *t*-test: *p = 0.018, n = 7 cells/5 assembloids).

Data in (**E**), (**F**), (**H**), (**I**), (**N**), and (**O**) are shown as the average values with individual responses overlaid. Scale bars (**G**), (**J**): 40 pA, 100 ms. Scale bar (**M**): 20% G/R.

See also **Figure S6** for snRNA-seq data supporting the NeuronChat predictions in (**A**) to (**D**).

990
Figure 4. TC synapses in assembloids undergo LTP via multiple protocols.
(**A**) Left: Schematic of the recording configuration. Right, top: 40-Hz electrical stimulation LTP-
induction protocol. Right, bottom: Representative trace of a response to 10-Hz stimulation.
(**B**) Left: Time course data demonstrating that 40-Hz stimulation repeated three times (arrows)
induces LTP in TC assembloids (n = 9 cells/9 assembloids). Right: Representative traces from
the first 5 min (1, dark) and final 5 min (2, light) of the experiment. Circles indicate electrical
stimuli.
(**C**) Bar graph of group data after 40-Hz induction from (**B**) shows EPSC amplitudes significantly
differ from baseline values (one-sample *t*-test, μ = 100, ##p = 0.0077).

(**D**) Top: Spike-timing–dependent plasticity (STDP) was induced by stimulating presynaptic hThO inputs (Pre) and then delivering four current injections (2-nA) to the postsynaptic cell (Post), repeated 50 times. Bottom: Representative trace of an hCO cell's response to stimulation and depolarization.

(**E**) Left: Time course data demonstrating that the short (×1) STDP protocol (arrow) in TC assembloids induces LTP (n = 7 cells /3 assembloids). Right: Representative traces from the first (dark) and final (light) 5 min of the experiment.

(**F**) Bar graph of group data following the ×1 STDP induction from (**E**) shows that EPSC amplitudes significantly differ from baseline values (one-sample t-test, μ = 100, #p = 0.04).

(**G**) Top: Long (×3) STDP-induction protocol, as in (**D**) but repeated three times every 5 min. Bottom: Representative trace of a response to stimulation and depolarization.

(**H**) Bar graph showing the average responses from nine cells from six assembloids after TC LTP induction. Shades of gray indicate different batches of assembloids; vertical lines denote separate assembloids.

(**I**) Time course of series resistance (Rs) normalized to the 5-min baseline period demonstrating TC LTP is not due to changes in Rs.

(**J**) Time course demonstrating the ×3 STDP protocol (arrows) induces LTP in TC synapses (black, n = 9 cells/6 assembloids). MPEP (blue, n = 5 cells/4 assembloids) or iBAPTA blocked LTP (orange, n = 5 cells/3 assembloids). AP5 did not block TC LTP (green, n = 5 cells/2 assembloids). Shaded area depicts the presence of bath-applied drugs.

(**K**) Bar graph of group data following ×3 STDP induction from (**J**). Differences from baseline were evaluated by one-sample t-test (μ = 100, ##p <0.01, ###p <0.0001). Differences between treatments and aCSF were evaluated by one-way ANOVA, p <0.0001. Dunnett's test: *p <0.05, *p <0.01.

(**L**) Example traces from the first (1) and final (2) 5 min of the experiment across conditions. Scale bars for (**A**): 20 mV, 200 ms. Scale bars for (**B**), (**E**): 50 pA, 200 ms. Scale bars for (**D**), (**G**): 40 mV, 100 ms. Scale bars for (**L**): 20 pA, 200 ms. Data shown are mean ± SEM (**B**), (**E**), (**I**), and (**J**), with individual data points overlaid as dots in (**C**), (**F**), and (**K**).

For (**B**), (**E**), (**J**), and (**L**) the first (1) and final (2) 5 min of the experiment are noted.

See also **Figure S7** for analysis of paired-pulse ratio (PPR) measures.

1032
Figure 5. CT synapses in assembloids undergo LTP.
(**A**) Left: Schematic of the recording configuration to induce CT LTP. Right: The long (×3)
STDP-induction protocol and example response.
(**B**) Bar graph showing the average responses in 14 cells from 9 assembloids after CT LTP
induction in aCSF. Shades of gray indicate different batches of assembloids; vertical lines
denote separate assembloids.
(**C**) Time course demonstrating that ×3 STDP delivery (arrows) induces LTP in CT synapses
(black, n = 14 cells/9 assembloids). MPEP (blue, n = 8 cells/6 assembloids), AP5 (green, n = 15
cells/7 assembloids), or iBAPTA (orange, n = 6 cells/2 assembloids) blocked LTP. Shaded area
depicts the presence of bath-applied drugs. The first (1) and final (2) 5 min of the experiment
are noted.
(**D**) Time course of Rs demonstrating that CT LTP is not due to changes in Rs.
(**E**) Bar graph of group data from (**C**). Differences from baseline were evaluated by one-sample
t-test ($\mu = 100$, ##p <0.01). Differences between treatments and aCSF were evaluated by one-
way ANOVA, p = 0.0083. Dunnett's test: *p <0.05.
(**F**) Example traces from the first (1) and final (2) 5 min of the experiment across conditions.
Circles indicate electrical stimulation.
Scale bars for (**A**): 40 mV, 100 ms. Scale bars for (**F**): 50 pA, 200 ms. Data shown are mean ±
SEM (**C**), (**D**) with individual data points overlaid in (**E**).
See also **Figure S7** for PPR analysis.
1053
1054

Figure 6. TC synapses in assembloids undergo LTD.
(**A**) Left: Schematic of the recording configuration to induce TC LTD. Right, top: LTD was induced with electrical stimulation delivered at 1 Hz for 900 pulses. Bottom: Example responses of a cell to a subset of the 900 pulses, dark-to-light traces depict the responses as the number of pulses progressed (from pulse (p) 1 to p900).
(**B**) Bar graph showing the average responses from nine individual cells from nine assembloids after TC LTD induction in aCSF. Shades of gray indicate different batches of assembloids; vertical lines denote separate assembloids.
(**C**) Time course data demonstrating that 1-Hz electrical stimulation (thick dashed line) induces LTD in TC synapses (black, n = 9 cells/9 assembloids). MPEP (blue, n = 6 cells/4 assembloids), AP5 (green, n = 6 cells/5 assembloids), or iBAPTA (orange, n = 5 cells/4 assembloids) blocked LTD. Shaded area depicts the presence of bath-applied drugs. The first (1) and final (2) 5 min of the experiment are noted.
(**D**) Time course of Rs normalized to the 5-min baseline period, demonstrating that TC LTD is not due to changes in Rs.
(**E**) Bar graph of group data after 1-Hz stimulation from (**C**). Differences from baseline were evaluated by one-sample *t*-test (μ = 100, ##p <0.01). Differences between treatments and aCSF were evaluated by one-way ANOVA, p = 0.0034. Dunnett's test: **p <0.01.
(**F**) Example traces from the first (1) and final (2) 5 min of the experiment across conditions. Circles indicate electrical stimulation.
Scale bars for (**A**): 5 mV, 100 ms. Scale bars for (**F**): 40 pA, 200 ms. Data shown are mean ± SEM (**C**), (**D**), and (**E**), with individual data points overlaid in (**E**).
See also **Figure S7** for PPR analysis.

Figure 7. CT synapses in assembloids undergo LTD.
(**A**) Schematic of the experimental condition for CT LTD induction and the 1-Hz LTD-induction protocol.
(**B**) Bar graph of the average responses of 10 individual cells from 9 assembloids after CT LTD induction in aCSF. Shades of gray indicate different batches of assembloids; vertical lines denote separate assembloids.
(**C**) Time course data show that 1-Hz stimulation (thick dashed line) induced LTD in CT synapses (black, n = 10 cells/9 assembloids). MPEP (blue, n = 5 cells/5 assembloids), AP5 (green, n = 4 cells/3 assembloids), or iBAPTA (orange, n = 4 cells/4 assembloids) blocked LTD induction. Shaded area depicts the presence of bath-applied drugs. The first (1) and final (2) 5 min of the experiment are noted.
(**D**) Time course of Rs normalized to the 5-min baseline period, demonstrating that CT LTD is not due to changes in Rs.
(**E**) Bar graph of group data after 1-Hz stimulation from (**C**). Differences from baseline were evaluated by one-sample *t*-test (μ = 100, #p <0.05). Differences between treatments and aCSF were evaluated by one-way ANOVA, p = 0.0004. Dunnett's test: ***p <0.001.
(**F**) Example traces from the first (1) and final (2) 5 min of the experiment across conditions. Circles indicate electrical stimulation.
Scale bars for (**A**): 5 mV, 100 ms. Scale bars for (**F**): 50 pA, 200 ms. Data shown are mean ± SEM (**C**), (**D**), and (**E**), with individual data points overlaid in (**E**).
See also **Figure S7** for PPR analysis.

STAR Methods

Human iPSC culture

The use of hiPSCs for the generation of organoids was approved by the St. Jude Institutional Review Board. TP-190a and TP-189 were derived from dental pulp cells from neurotypical male and female subjects, respectively, with normal karyotypes. Cells were reprogrammed using episomal plasmids (ALSTEM LLC). The 2242 (i.e., 2242-1), 1205 (i.e., 1205-4), and 8858 (i.e., 8858-3) lines/clones were previously published.[95]

All hiPSC lines were maintained in culture on human ES-qualified Matrigel (5264004, Corning) in complete mTeSR Plus (100-0276, STEMCELL Technologies) at 5% O_2, 37 °C, and 5% CO_2. The cultures were passaged with Versene (15040066, ThermoFisher). Genetically modified reporter hiPSC lines were validated before differentiation (**Figure S1**). Specifically, six assays were performed: (1) Colonies were immunostained for six pluripotency markers[96] (Stemlight Pluripotency Antibody Kit 9656S, Cell Signaling); (2) Expression of five additional pluripotency markers[96] was assayed using a Custom TaqMan RT-qPCR assay designed in-house (manufactured by ThermoFisher); (3) G-banding; (4) Copy number variation at the seven most frequently aberrant chromosomal locations[97–99] was assayed using a custom TaqMan RT-qPCR assay designed in-house (manufactured by ThermoFisher); (5) Global methylation analysis (Infinium MethylationEPICv1.0 850K Beadchip) was performed to identify methylation status at select epigenetic markers,[100,101] and results were then compared to established and previously published hiPSC lines;[102]; (6) Trilineage assay (STEMdiff Trilineage Differentiation Kit 05230, STEMCELL Technologies) was performed, and markers of interest[96] were analyzed using a Custom TaqMan RT-qPCR assay designed in-house (manufactured by ThermoFisher).

Generating reporter lines

Genome-edited TP-190a hiPSCs were generated using CRISPR-Cas9 technology. Briefly, hiPSCs were pretreated with StemFlex (Thermo-Fisher Scientific) supplemented with 1×

RevitaCell (ThermoFisher Scientific) for 1 h. Then, approximately 10^6 cells were transiently

transfected with precomplexed ribonuclear proteins consisting of 250 pmol chemically modified

single-guide RNA (sgRNA; Synthego), 165 pmol Cas9 protein (St. Jude Protein Production Core),

500 ng pMaxGFP (Lonza), and 3 µg ssODN donor (for deletion) or 1 µg dsDNA donor (for tagging)

via nucleofection (Lonza, 4D-Nucleofector™ X-unit) using solution P3 and program CA-137 in a

large (100-µL) cuvette, according to the manufacturer's recommended protocol. Five days

postnucleofection, cells were single-cell sorted by FACS for GFP⁺ (transfected) cells and plated

onto Vitronectin XF (Stem Cell Technologies)-coated plates into prewarmed (37°C) StemFlex

media supplemented with 1× CloneR (Stem Cell Technologies).

Clones were screened for the desired modification via targeted deep sequencing using

gene-specific primers with partial Illumina adapter overhangs on a Miseq Illumina sequencer, as

previously described,[103] or by junction PCR followed by sequencing. Briefly, cell pellets of

approximately 10,000 cells were lysed and used to generate gene-specific amplicons with partial

Illumina adapters in PCR#1. Amplicons were indexed in PCR#2 and pooled with targeted

amplicons from other loci to create sequence diversity. Additionally, 10% PhiX Sequencing

Control V3 (Illumina) was added to the pooled amplicon library prior to running the sample on a

Miseq Sequencer System (Illumina) to generate paired 2 × 250-bp reads. Samples were de-

multiplexed using the index sequences, fastq files were generated, and next-generation

sequencing (NGS) analysis of clones was performed using CRIS.py.[104] Correctly edited clones

were identified, expanded, and sequence confirmed. Final clones were authenticated using the

PowerPlex® Fusion System (Promega), which was performed at the St. Jude Hartwell Center

and tested for mycoplasma by using the MycoAlertTMPlus Mycoplasma Detection Kit (Lonza).

Editing-construct sequences and relevant primers are listed in **Table S1**.

1154

Thalamic organoid generation

The hThOs were generated from TCF7L2-tdTomato reporter hiPSC lines, except for indicated organoids in **Figure S2**. For differentiation, cryovials were plated and maintained in culture in mTeSR1 (85850, STEMCELL Technologies). At 80% confluency, they were dissociated into single cells with Accutase (AT-104, Innovative Cell Technologies) and plated into low-attachment 96-well V-bottom plates (MS-9096VZ, Sbio) at 10,000 cells/well, in gfCDM media (1:1 IMDM (12440053, Thermofisher): Ham's F12 (12-615F, Lonza), 1× lipid concentrate (11905031, Thermofisher), 1× antibiotic-antimycotic (15240062, Gibco), 450 µM monothioglycerol (M6145, Sigma), 15 µG/mL apotransferrin (T1428, Sigma), 5 mg/mL BSA (50-255-465, Fisher Scientific) supplemented with 5 µM SB-431542 (TGFβ inhibitor, 1614, Tocris), 1 µg/mL insulin (I9278, Sigma), 1% v/v growth factor–reduced (GFR) Matrigel (354230, Corning), 2 µM thiazovivin (72254, STEMCELL Technologies). On Day (D) 2, half the media was replaced with the same media supplemented with 4 µM dorsomorphin (3093, Tocris). On D4 and D6, half of the media was replaced with gfCDM supplemented with 5 µM SB-431542, 100 nM Smoothened agonist (73414, STEMCELL Technologies), and 20 ng/mL Fgf8b (100-25, PeproTech). In some differentiations, Matrigel was added on D2 instead of D0, but no difference was detected in the resulting organoids. On D8, D10, and D12, three-fourths of the media was replaced with the same but further supplemented with 30 ng/mL BMP7 (354BP010, R&D Systems) and 1 µM MEKi PD0325901 (S1036, R&D Systems). On D14, D16, and D18 half the media was replaced with the same, but gfCDM was substituted with thalamic N2 media (DMEM:F12, 10% ES-FBS (ES-009-C, SIGMA), 1× N2 supplement (17502-048, Gibco), 1× Glutamax (35050061, Gibco), and 1× antibiotic-antimycotic.

On D20, all organoids were transferred to a magnetic stir bioreactor (BWS-S03N0S-6, ABLE Corporation, Tokyo) in thalamic N2 media and agitated at 40 rpm. On D22 and D24, half the media was replaced with thalamic N2 supplemented with 1× B27 without vitamin A (12587-010, Gibco), 20 ng/mL heat-stable bFGF (PHG0367, ThermoFisher) and 20 ng/mL EGF (AF-100-

15-100UG, Peprotech). On D26 and D28, half the media was replaced with the same, but the concentrations of bFGF and EGF were reduced to 10 ng/mL. On D30 and D32, all the media was replaced with thalamic N2 supplemented with 1× B27 without vitamin A.

Starting D35, full media was replaced every 4 days with BrainPhys (05790, STEMCELL Technologies) supplemented with 1× N2, 1× B27 without vitamin A, 10% ES-FBS, 10 ng/mL BDNF (450-02, Peprotech), and 10 ng/mL GDNF (450-10, Peprotech). Starting D70, all the media was changed to BrainPhys supplemented with 1× N2, 1× B27 without vitamin A, 1× glutamax, 1× NEAA (11140050, Gibco), 1× antibiotic-antimycotic, 200 μM ascorbic acid (A4403, Sigma), 100 μM dibutyryl cAMP (D0627, Sigma), 1% ES-FBS, 10 μM DAPT (2634, Tocris), 20 ng/mL BDNF, and 20 ng/mL GDNF. Starting D82, the concentration of BDNF and GDNF was reduced to 10 ng/mL. In addition, after D30, large organoids were pinched into two halves by using a pair of ultra-fine clipper scissors (15300-00, Fine Science Tools) every 5–7 days to avoid large necrotic centers.

Cortical organoid generation

The hCOs were generated from VGLUT1-tdTomato reporter iPSC lines. At 80% confluency, cell cultures were dissociated into single cells with Accutase (AT-104, Innovative Cell Technologies), and plated into low-attachment 96-well V-bottom plates (MS-9096VZ, Sbio) at 9000 cells/well, in EB media (DMEM:F12, 20% knockout serum replacement (KSR) (10828, Life Technologies), 3% ES-FBS, 1× Glutamax, 1× β-mercaptoethanol (2020-07-30, Gibco), 1× antibiotic-antimycotic) supplemented with 5 μM SB-431542 (TGFβ inhibitor, 1614, Tocris), 2 μM dorsomorphin (3093, Tocris), 3 μM IWR1e (Wnt inhibitor, 681669, EMD Millipore), 1% v/v GFR-Matrigel, and 2 μM thiazovivin.

In some differentiations, 0% or 0.5% v/v GFR-Matrigel was added on D0 but no difference in the resulting organoids was detected. On D2, half the media was replaced with the same but without Matrigel. On D4 and D6, half the media was replaced with GMEM KSR media (GMEM,

20% KSR, 1× NEAA (Gibco), 1× sodium pyruvate (11360070, Gibco), 1× β-mercaptoethanol, 1× antibiotic-antimycotic) supplemented with 5 μM SB-431542, 3 μM IWR1e, 2.5 μM cyclopamine (72074, STEMCELL Technologies), and 2 μM thiazovivin. On D8, half the media was replaced with GMEM KSR media supplemented with 5 μM SB-431542, 3 μM IWR1e, and 2.5 μM cyclopamine. On D10, D12, D14, and D16, half the media was replaced with GMEM KSR media supplemented with 5 μM SB-431542 and 3 μM IWR1e. On D18 and D20, half the media was replaced with CBO N2 media (DMEM:F12, 1× chemically defined lipid concentrate (11905-031, Life Technologies), 1× N2 supplement (17502-048, Gibco) and 1× antibiotic-antimycotic) supplemented with 1× B27 supplement without vitamin A (12587-010, Gibco), 20 ng/mL heat-stable bFGF, and 20 ng/mL EGF. On D22, organoids were transferred to a magnetic stir bioreactor (BWS-S03N0S-6, ABLE Corporation) in CBO N2 media supplemented with 1× B27 supplement without vitamin A, 20 ng/mL heat-stable bFGF and 20 ng/mL EGF, and agitated at 40 rpm. Half of the media was replaced with the same on D24, D26, and D28. On D30, the media was changed to CBO FBS media (DMEM:F12, 1× chemically defined lipid concentrate (11905-031, Life Technologies), 1× N2 supplement, 10% ES-FBS, 5 μg/mL heparin and 1× antibiotic-antimycotic) supplemented with 1× B27 supplement without vitamin A. Full media was replaced every 4 days. On D42 and D46, the media was changed to CBO FBS media supplemented with 1× B27 supplement without vitamin A, 10 ng/mL BDNF (450-02, Peprotech), and 10 ng/mL GDNF (450-10, Peprotech). Starting D50, all the media was replaced every 4 days with BrainPhys (05790, STEMCELL Technologies) supplemented with 1× N2 supplement, 50× B27 supplement without vitamin A, 10% ES-FBS, 10 ng/mL BDNF and 10 ng/mL GDNF. Starting D70, media was changed to BrainPhys supplemented with 1× N2, 50× B27 without vitamin A, 1× glutamax, 1× NEAA, 1× antibiotic-antimycotic, 200 μM ascorbic acid, 100 μM cAMP, 1% ES-FBS, 10 μM DAPT, 20 ng/mL BDNF, and 20 ng/mL GDNF. Starting at D82, the concentration of BDNF and GDNF was reduced to 10 ng/mL. In addition, after D30, large organoids were pinched into 2 halves by using a pair of ultra-fine clipper scissors every 5–7 days to avoid large necrotic centers.

1233

Generation of thalamocortical assembloids

Between D90 and D120, TCF7L2-tdTomato⁺ hThOs were paired with VGLUT1-tdTomato⁺ hCOs

of similar age. Each pair was transferred to a well of a low-attachment, 24-well plate in 500 mL

Fusion media (BrainPhys supplemented with 1× N2, 50× B27 without vitamin A, 1× glutamax, 1×

NEAA, 1× antibiotic-antimycotic, 200 µM ascorbic acid, 100 µM cAMP, 1% ES-FBS, 10 µM DAPT,

20 ng/mL BDNF, 20 ng/mL GDNF, and the CEPT cocktail (50 nM Chroman 1 [HY-15392,

MedChem Express], 5 µM emricasan (S7775, Selleckchem), 0.7 µM trans-ISRIB (#5284, Tocris),

and polyamine supplement (#P8483, Sigma-Aldrich[105]) supplemented with 0.5% v/v GFR-

Matrigel. The plate was left tilted and undisturbed in the incubator. After 3 days, 60% of the media

in each well was replaced with fusion media. This was done slowly, while keeping the plate tilted

with minimal disturbance to the "fused" organoid pair in each well. The same was done on D6 and

D9. Subsequently, 80% of the media was replaced every 3 days. On D4, the plate was transferred

to an orbital shaker at 80 rpm. The shaker speed was increased to 90 rpm on D5, 100 rpm on D6,

and starting D7, the assembloids were kept at 110 rpm. Between 5 and 10 weeks postfusion,

assembloids were harvested for electrophysiological experiments.

1249

Plasmids and lentiviruses

Synapsin-EGFP (hSyn-GFP) lentiviruses with the VSV-G pseudo-type were generated using the

pHR-hSyn-eGFP plasmid[106] (Addgene: 114215, a gift from Xue Han) by the St. Jude Viral Vector

Core. For APEX2 experiments, pLenti-hSyn-V5-COX4-APEX2 plasmid was generated by cloning

the V5-COX4-APEX2 sequence from pAAV-COX4-dAPEX2[54] (Addgene: 117176, a gift from

David Genty) into the pLenti backbone containing the human *SYN* promoter. Briefly, pAAV-COX4-

APEX2 was digested with BspE1 and EcoR1, and pLenti-hSyn-nucGFP[107] (Addgene: 140190, a

gift from Lorenz Studer) was digested with EcoRI and AgeI to remove the nucGFP-coding

sequence. Insert containing the V5-COX4-APEX2 sequence was then ligated into the pLenti-hSyn

backbone. The resulting plasmid sequence was confirmed by Sanger sequencing. Lenti-hSyn-V5-COX4-APEX2 (hSyn-V5-Mito-APEX2) lentiviruses with the VSV-G pseudo-type were generated by the St. Jude Viral Vector Core.

Lentiviral vectors prepared at 1-3bn TU/mL were added to organoids in the bioreactor at 200×. For vectors at lower titer, 2 µg/mL Polybrene was also added to the media. After 18–20 h, organoids were washed twice with DMEM:F12 and fed with fresh media. Media changes were continued according to the protocol. Lentiviral expression was detected at 72 h posttransduction.

Immunofluorescence and light microscopy

Organoids were briefly rinsed in phosphate-buffered saline (PBS) and then fixed in 4% paraformaldehyde in PBS overnight at 4°C. Following rinses in PBS, organoids were cryoprotected by incubation overnight in 30% sucrose in PBS. Organoids were then mounted in Optimal Cutting Temperature (O.C.T.) Compound (Tissue-Tek). Samples were stored at –80°C until cryosectioning. Cryosectioning was performed on a Leica CM 3050 cryostat set to –20°C. Serial sections of 20-µm thickness were mounted onto FisherBrand Superfrost Plus microscope slides and stored at –20°C.

Slides were briefly rehydrated with PBS and then blocked for 1 h at room temperature in blocking buffer (PBS, 5% normal donkey serum, 0.2% Triton-X100, 0.02% sodium azide, filter sterilized). Slides were incubated overnight at 4°C in primary antibodies diluted in blocking buffer, washed with PBS-Tween (0.1%), and incubated 1 h at room temperature in secondary antibodies diluted 1:500 in blocking buffer. Slides were then washed with PBS-Tween (0.1%), and nuclei were labeled with DAPI. Excess DAPI was removed by washing with PBS, and slides were dried and mounted for imaging using Prolong Diamond (Thermo Fisher, P36961). Images were acquired on a Zeiss Axio Imager M2 equipped with a 20× Plan-Apochromat Objective (Zeiss, 0.8 NA), 40× EC Plan-NeoFluar Objective (Zeiss, 1.3 NA), and Apotome.2 (Zeiss). Images including GABA were acquired using the 40× objective. All other images were acquired using the 20×

objective. During imaging, exposure times were kept below saturation, and imaging conditions were constant within experiments. For images acquired with the Apotome.2, Z-series were acquired at software-recommended intervals and image stacks were then deconvolved using ZEN software and a constrained iterative algorithm. Images are shown as maximum intensity projections prepared in Zeiss ZEN 3.7 software.

The following primary antibodies and dilutions were used: TCF7L2 (Cell Signaling Technologies 2569, 1:1000), OTX2 (R&D Systems AF1979, 1:100), TUBB3 (Abcam Ab107216, 1µg/mL), SOX2 (R&D Systems MAB2018, 1:200), V5 (Invitrogen R960-25, 1:1000), FOXP2 (Abcam ab16046, 1:250), LHX2 (Sigma ABE1402,1:250), GBX2 (R&D Systems AF4638, 1:250), and GABA (Sigma A2052, 1:5000). The following f(ab')$_2$ secondary antibodies from Jackson Immunoresearch were used: donkey anti–rabbit Alexa Fluor (AF) 488 (711-546-152), donkey anti–goat AF 647 (705-606-147), donkey anti–mouse AF 488 (715-546-150), donkey anti–mouse AF 647 (715-606-150), and donkey anti–chicken AF 647 (703-606-155).

Bulk RNA-seq

Each sample consisted of 2-3 pooled organoids from the indicated condition. Total RNA was isolated using the Direct-zol RNA Microprep Kits (Zymo, R2061), and DNA contamination was removed using the DNA-free DNA Removal Kit (Thermo Fisher, AM1906). RNA was quantified using the Quant-iT RiboGreen RNA assay (ThermoFisher) and quality checked by the 2100 Bioanalyzer RNA 6000 Nano assay (Agilent) or 4200 TapeStation High Sensitivity RNA ScreenTape assay (Agilent) prior to library generation. Libraries were prepared from total RNA with the TruSeq Stranded mRNA Library Prep Kit, according to the manufacturer's instructions (Illumina PN 20020595). Libraries were analyzed for insert-size distribution using the 2100 BioAnalyzer High Sensitivity kit (Agilent), 4200 TapeStation D1000 ScreenTape assay (Agilent), or 5300 Fragment Analyzer NGS fragment kit (Agilent). Libraries were quantified using the Quant-iT PicoGreen ds DNA assay (ThermoFisher) or by low-pass sequencing with a MiSeq nano kit

(Illumina). Paired-end 100-cycle sequencing was performed on a NovaSeq 6000 (Illumina) in the St. Jude Hartwell Center Genome Sequencing Core.

For tdTomato-expression analysis, we built a custom reference genome by adding the tdTomato sequence to the human genome (hg38, gencode v31). The tdTomato sequence was also added to the gene-annotation gtf file (gencode v31). Read alignment to the custom genome was performed with STAR (version 2.7) software.[108] Gene-level read count was determined using RSEM (version 1.3.1).[109]

For differential gene expression analysis, only protein-coding genes validated at GENECODE confidence level 1 to 3 were considered. To remove genes that were lowly expressed, we first calculated the cutoff as 10 read counts per million library size, where the library size was defined as the median library size in the data set. We then kept genes with expression level (counts per million) equal to or above the cutoff in a minimum number of samples, where the number of samples was chosen according to the minimum group sample size. The data were then normalized by TMM function in edgeR package,[110] followed by the limma package with its voom method, linear modeling, and empirical Bayes moderation to assess differential expression.[111]

Markers of interest were identified by performing a differential-expression analysis using *BrainSpan* Developmental Transcriptome data. Thalamic structures (mediodorsal nucleus of the thalamus and dorsal thalamus) were compared with all cortical structures. The top 100 up- or down-regulated genes in thalamic vs cortical structures were identified as "thalamic" or "cortical" markers, respectively.

Deconvolution of bulk RNA-seq data was performed using the VoxHunt (v1.0.1)[112] R package using the default workflow (https://quadbio.github.io/VoxHunt/articles/deconvolution.html). *Allen Brain Atlas* data derived from E13 mouse brain were used as a reference. The "broad" gene set contained the top 50

markers for each region of interest. The top 15 markers were then used as input for the deconvolution tool.

GO term enrichment analysis was performed using g:Profiler.[113] For all analyses, a custom background was uploaded containing genes detected in the data set of interest. Driver terms containing fewer than 300 genes were selected for graphing. All graphs, except heatmaps, were prepared in R using ggplot2 (v3.4.0).[114] Heatmaps were prepared using the ComplexHeatmap (v2.10.0) R package.[115,116]

RT-qPCR

RNA was isolated and DNase-treated, as described above. Reverse transcription was performed using 100 ng RNA and the iScript cDNA Synthesis Kit (Bio-Rad, 1708891). A qPCR analysis was then performed using SYBR Green Master Mix (Thermo Fisher, 4309155) and a C1000 Touch Thermal Cycler (Bio-Rad). The following primers were used: *GAPDH* Forward 5′-AATCCCATCACCATCTTCCA-3′, *GAPDH* Reverse 5′-TGGACTCCACGACGTACTCA-3′, *TCF7L2* Forward 5′-GAATCGTCCCAGAGTGATGTCG-3′, *TCF7L2* Reverse 5′-TGCACTCAGCTACGACCTTTGC-3′, *OLIG3* Forward 5′-TGAGGCTGAAGATCAACGGACG-3′, *OLIG3* Reverse 5′-AGTTTCTGGCGAGCAGGAGTGT-3′, *GBX2* Forward 5′-GCGGAGGACGGCAAAGGCTTC-3′, *GBX2* Reverse 5′-GTCGTCTTCCACCTTTGACTCG-3′, *LHX9* Forward 5′-ACCTGCTTTGCCAAGGACGGTA-3′, *LHX9* Reverse 5′-TGACCATCTCCGAGGCGGAAAT-3′, *OTX2* Forward 5′-GGAAGCACTGTTTGCCAAGACC-3′, *OTX2* Reverse 5′-CTGTTGTTGGCGGCACTTAGCT-3′, *FOXG1* Forward 5′-GTATGTGGTCACTAACAGGTC-3′, and *FOXG1* Reverse 5′-ACCACAGTATCACAATCAAG-3′.

Data were analyzed using the $2^{-\Delta\Delta Cq}$ method [previously known as the $2^{-\Delta\Delta Ct}$ method, first described in the Applied Biosystems User Bulletin 2 (P/N 4303859)].[117] Transcripts of interest were normalized first to *GAPDH* (within sample), then to the mean ΔCq of the hThO samples with high tdTomato that were previously used for bulk RNA-seq. Regression analyses were performed

in R using normalized values and graphed using the ggscatter function from ggpubr (v0.5.0). For all transcripts, except *FOXG1*, r and p were calculated using the Pearson correlation method, where *r* represents the correlation coefficient and p represents the p-value, and lines were fit using linear regression. Due to the presence of extreme outliers, for *FOXG1* the r_s and p-valuewere calculated using the Spearman correlation method, in which r_s represents the correlation coefficient, p represents the p-value, and the nonlinear curve is fit using the Loess local polynomial-regression method.

Preparation and sequencing of the snRNA-seq library

Two independent biological replicates were performed per the differentiation protocol (either cortical or thalamic), each containing 36 organoids pooled together. The hThOs were D91 or D96; the hCOs were D91. All organoids were flash frozen in liquid nitrogen and stored at –80°C until dissociation. Nuclei dissociation was performed as previously described.[118] Briefly, frozen tissue was mechanically dissociated with a Dounce homogenizer in detergent lysis buffer containing 0.32 M sucrose, 10 mM HEPES (pH 8.0), 5 mM $CaCl_2$, 3 mM magnesium acetate, 0.1 mM EDTA, 1 mM DTT, and 0.1% Triton-X100. The resulting homogenate was filtered through a 40-µm strainer and washed with the same solution described, without the Triton-X100 added. Nuclei were then centrifuged at 3200 ×*g* for 10 min at 4°C, and the supernatant was decanted. A sucrose-dense solution containing 1 M sucrose, 10 mM HEPES (pH 8.0), 3 mM magnesium acetate, and 1 mM DTT was carefully layered underneath the remaining supernatant and then spun at 3200 ×*g* for 20 min at 4°C. The supernatant was discarded, and the final remaining nuclei were resuspended in 0.4 mg/mL BSA and 0.2 U/µL Lucigen RNAse inhibitor (catalog number 30281-1) diluted in PBS. Between 5000 and 10,000 nuclei were inspected and counted on a hemacytometer before loading onto the 10× Chromium Controller (10× Genomics, catalog number 1000171). The snRNA-seq libraries were prepared using the 10× Genomics Chromium

Next GEM Single Cell Kit, version 3.1 single index gene expression profiling assay, according to

the manufacturer's instructions.

Libraries were analyzed for insert-size distribution by using the 2100 BioAnalyzer High

Sensitivity kit (Agilent), 4200 TapeStation D1000 ScreenTape assay (Agilent), or 5300 Fragment

Analyzer NGS fragment kit (Agilent). Libraries were quantified using the Quant-iT PicoGreen ds

DNA assay (ThermoFisher) or by low-pass sequencing with a MiSeq nano kit (Illumina). Paired-

end 100-cycle sequencing was performed on a NovaSeq 6000 (Illumina) in the St. Jude Hartwell

Center Genome Sequencing Core.

1395

Analysis of snRNA-seq data

1396

Sequences from each Illumina-sequencing data set were de-multiplexed using bcl2fastq

v2.20.0.422 (Illumina). The sequencing data were aligned to the human reference genome

GRCh38 (10× Genomics, v2020-A) using the CellRanger "count" algorithm (10× Genomics,

v7.0.0); however, the "–force-cells" option was set to the estimated number of cells loaded for

each sample (snCBO1: 6,000; snCBO2: 8,000; snTha1: 8,000; snTha2: 10,000). From the gene

expression matrix, the downstream analysis was carried out in R (v4.2.1). First, the ambient RNA

signal was removed using the default SoupX (v1.6.2) workflow (autoEstCounts and adjustCounts;

github.com/constantAmateur/SoupX).[119]

Each data set was initially filtered so that genes that were expressed in at least three cells,

and cells that expressed at least 200 genes were included. Additionally, cells with fewer than 300

genes (presumed to be droplets or cellular debris), fewer than 500 UMIs, more than 1% unique

transcripts derived from mitochondrial genes, or more than 3 median absolute deviations (MADs)

from the median number of unique transcripts derived from mitochondrial genes were removed.

Afterwards, cells with more than 3 MADs from the median number of genes expressed were

removed.

Samples were then preprocessed using the standard Seurat (v4.3.0) workflow (NormalizeData, ScaleData, FindVariables, RunPCA, FindNeighbors, FindClusters, and RunUMAP; github.com/satijalab/Seurat).[120–123] Data sets were individually log-normalized using Seurat's NormalizeData with default parameters. Cell cycle scoring was conducted using the associated S- and G2M-phase gene list from Tirosh et al.[124] and the CellCycleScoring command in Seurat. We calculated 3000 features exhibiting high cell-to-cell variation in the data set by using Seurat's FindVariableFeature function. Next, we scaled the data by linear regression against the number of reads by using Seurat's ScaleData function with default parameters. The variable genes were projected onto a low-dimensional subspace by performing principal component analysis using Seurat's RunPCA function with default parameters. The number of principal components (n = 30) was selected based on inspection of the plot of variance explained.

Data sets were integrated using Harmony (v 0.1.1) with default parameters.[125] A shared-nearest-neighbor graph was constructed based on the Euclidean distance in the low-dimensional subspace using Seurat's FindNeighbors with dims = 1:30 and default parameters. Integrated data sets then underwent nonlinear dimensional reduction and visualization using UMAP. Clusters were identified using a resolution of 0.4 and the Leiden algorithm for the integrated data sets. Pseudotime analysis was conducted using Monocle3 (v1.3.1) with default parameters.[126–129] Trajectory starting points were manually selected based on the expression of mitotic markers (e.g., *MKI67*) and/or neural precursor markers (e.g., *TNC*). Mapping of snRNA-seq data sets onto *Allen Brain Atlas* and *BrainSpan* reference data sets was performed using the VoxHunt (1.0.1) R package with the suggested workflows (https://quadbio.github.io/VoxHunt/articles/getting_started.html; https://quadbio.github.io/VoxHunt/articles/other_references.html).[112] For *Allen Brain Atlas* comparisons, data derived from E15 mouse embryos were used. For *BrainSpan* comparisons, data derived from human fetal tissue 13–24 postconception weeks (pcw) were used. Neural communication patterns were predicted and visualized using the NeuronChat (v1.0.0) R package

with the suggested workflow (https://github.com/Wei-

BioMath/NeuronChat/blob/main/vignettes/NeuronChat-Tutorial.html).[62]

Cell types were assigned to each cell/cluster based on marker expression and cell cycle

analysis. For hThO annotation, markers of interest were identified based on a comparison to

previously published scRNA-seq or snRNA-seq studies in developing mouse thalamus or

diencephalon.[130,131] Additional markers were identified based on previously published *in situ*

studies examining embryonic rodent thalamus. For example, within the mouse thalamus[132] and

hThOs, *SOX2* is expressed in a subset of postmitotic neurons. For hCO cluster annotation,

markers of interest were identified using a previously published scRNA-seq study that examined

human neocortex at midgestation.[51] Additional markers were identified based on previously

published *in situ* studies examining embryonic rodent cortex.

1449

Electron microscopy for DAB-labeled samples

Prior to fusion, TCF7L2-tdTomato[+] thalamic and VGLUT1-tdTomato[+] hSyn-GFP[+] cortical

organoids were separately transduced in low-attachment 6-well plates with 10^7 TU/mL Lenti-

hSyn-V5-COX4-APEX2 lentiviral vector. After 18–20 h, organoids were washed twice with

DMEM:F12 and fed fresh media. After 3 days, APEX2-transduced thalamic organoids were fused

with cortical organoids, and APEX2-transduced cortical organoids were fused with thalamic

organoids to generate the assembloids described above. At 6–7 weeks postfusion, the

assembloids were prepared for electron microscopy analysis. Specifically, each assembloid was

embedded at the center of a UV-sterilized Nunc Thermanox plastic coverslip (Thermo Fisher,

174950) in 5 µL GFR-Matrigel for 1 h at 37°C. They were then transferred to the fusion media in

6-well plates and placed in the incubator overnight. The following day, a sterile blade was used

to cut a V-shaped notch out of the coverslip on the side containing the APEX2[+] half of the

assembloid. DAB labeling was then performed using an adapted protocol.[133]

Briefly, after 1 additional day at 37°C, assembloids were fixed for 1 h in 2% glutaraldehyde in 0.1 M sodium cacodylate at room temperature, after which the fixative was replaced, and samples were incubated 1 h at 4°C. The samples were then washed thrice for 5 min in ice-cold wash solution (0.1 M sodium cacodylate). Next, the samples were incubated 5 min in 20 mM glycine in 1× sodium cacodylate, then washed thrice for 5 min on ice. The samples were preincubated in 0.5 mg/mL DAB in 0.1 M sodium cacodylate for 30 min on ice. The samples then underwent DAB staining in 0.5 mg/mL DAB and 50 mM H_2O_2 in 0.1 M sodium cacodylate on ice. The reaction was terminated by washing the samples thrice for 5 min on ice in wash solution.

After the DAB labeling developed, samples were postfixed in 2% osmium tetroxide in 0.1 M cacodylate buffer on ice for 30 min. Samples were subsequently washed 5 times for 2 min in ice-cold water, and then they were contrasted with 2% uranyl acetate overnight at 4°C. Samples were washed five times for 2 min in ice-cold water. Samples were dehydrated on ice by an ascending series of ethanol to 100%, followed by 100% propylene oxide at room temperature. Samples were infiltrated with EmBed-812 and polymerized at 60°C. Embedded samples were sectioned at ~70 nm on a Leica ultramicrotome and examined in a ThermoFisher Scientific TF20 transmission electron microscope at 80 kV. Digital micrographs were captured with an Advanced Microscopy Techniques imaging system. Unless otherwise indicated, all reagents were from Electron Microscopy Sciences.

Identification of synapses by transmission electron microscopy

Individual organoids were harvested between D102 and D121 for electron microscopy imaging. Samples were fixed in 0.1 M cacodylate buffer containing 2.5% glutaraldehyde and 2% paraformaldehyde. Samples were postfixed in reduced osmium tetroxide and contrasted with aqueous uranyl acetate. Dehydration was by an ascending series of ethanol to 100%, followed by 100% propylene oxide. Samples were infiltrated with EmBed-812 and polymerized at 60°C. Embedded samples were sectioned at ~70 nm on a Leica ultramicrotome and examined in a

ThermoFisher Scientific TF20 transmission electron microscope at 80 kV. Digital micrographs were captured with an Advanced Microscopy Techniques imaging system. Unless otherwise indicated, all reagents were from Electron Microscopy Sciences.

Fusion of 2-dimensional organoids

TCF7L2-tdTomato$^+$ thalamic and VGLUT1-tdTomato$^+$ hSyn-GFP$^+$ cortical organoids were halved using a pair of ultra-fine clipper scissors and plated in a culture-insert 2-well in µ-dish 35 mm (81176, Ibidi). Specifically, each half chamber was first coated with human ES-qualified Matrigel diluted 1:200 in DMEM:F12, for 1 h at room temperature. The coating solution was aspirated and 100 µL fusion media was added to each half. One thalamic organoid half was placed in one chamber and 1-2 cortical organoid halves were placed in the other and allowed to attach and extend neural processes for 5 days. On D5, the barrier was removed using sterilized blunt forceps, and the organoids were maintained in culture with media changes every 7 days. The barrier region in each dish was imaged every 3–7 days, from D9 to D61, at the same exposure time on a Zeiss AxioObserver D1.

Whole-cell patch-clamp electrophysiology

Whole-cell recordings were made in individual organoids between D90 and D141 or in assembloids between D19 and D78 postfusion. Organoids were placed in a recording chamber mounted on a two-photon laser-scanning microscope (Bruker) and superfused (2-3 mL/min) with aCSF containing the following solution: 125 mM NaCl, 2.5 KCl, 2 mM $CaCl_2$, 2 mM $MgCl_2$, 1.25 mM NaH_2PO_4, 26 mM $NaHCO_3$, and 20 mM glucose at 300–310 mOsm, equilibrated with 95% O_2/5% CO_2 at 32°C.

Cells were visualized under two-photon guidance by using PrairieView v5.5 software. Whole-cell voltage- and current-clamp recordings were made from visually identified thalamic or cortical cells. Short-term synaptic plasticity, 1-Hz induction of LTD, and 40-Hz LTP were recorded

in voltage-clamp mode (V_{Hold} = –60 mV), with an internal pipette containing the following solution: 125 mM CsMeSO$_3$, 2 mM CsCl, 10 mM HEPES, 0.1 mM EGTA, 4 mM ATP-Mg$_2$, 0.3 mM GTP-Na, 10 mM creatine phosphate-Na$_2$, 5 mM QX-314 chloride, and 5 mM TEA-Cl at pH 7.4 and 290-295 mOsm. Borosilicate glass pipettes (Sutter, 3-6 MΩ open pipette resistance) were used.

For investigating membrane properties and LTP induced by the STDP protocol, the internal solution contained 115 mM potassium gluconate (KGlu), 20 mM KCl, 10 mM HEPES, 4 mM MgCl$_2$, 0.1 mM EGTA, 4 mM ATP-Mg$_2$, 0.4 mM GTP-Na, and 10 mM creatine phosphate-Na$_2$ at pH 7.4 and 290-295 mOsm. Recordings were obtained using a Multiclamp 700B amplifier (Axon Instruments). Signals were digitized with an Axon Digidata 1550B (Axon Instruments) at 20 kHz and filtered at 2 kHz using Clampex 10.7 software. The liquid-junction potential was calculated to be –10 mV and was corrected for in each recording.

In current-clamp experiments, the rheobase was measured by first injecting a hyperpolarizing current step (–20 pA), followed by a depolarizing ramp (from –20 pA to +200 pA) into cells. The current at which the first AP was generated was recorded and averaged across cells. A series of hyperpolarizing and depolarizing step currents were injected into cells in current-clamp mode (+10 pA steps from –50 pA to +240 pA for 1 s) to measure input resistance and evoked firing rates.

In voltage-clamp experiments, synaptic currents were evoked using a bipolar concentric stimulating electrode (World Precision Instruments) or a homemade 2-prong stimulating electrode connected to a stimulus-isolation unit (Iso-Flex, A.M.P.I.) positioned in either the thalamic or cortical side of assembloids. Stimulus intensities were adjusted prior to each experiment to elicit measurable EPSCs in cortical or thalamic neurons. Because of the variability between assembloids, the amplitudes of evoked EPSCs ranged from –20 pA to –300 pA. PPRs were measured by delivering 2 stimuli 50, 100, 200, 500, or 1000 ms apart for both TC and CT synapses.

TC LTP was induced by high-frequency stimulation: 10 trains of 40-Hz stimulation for 200 ms every 5 s, repeated 3× every 5 min.[80] Additional TC LTP and CT LTP were induced via an STDP protocol: a single presynaptic electrical stimulation preceded four postsynaptic APs by 10 s. Postsynaptic APs were induced by four somatic current injections of 2 nA (2-ms duration) at 40 Hz. This protocol was repeated 50 times (at 1 Hz) every 5 min, for a total of three times (long STDP-induction protocol).[134] In a subset of experiments, the STDP protocol was delivered only 1× (short STDP-induction protocol). LTD was induced at TC and CT synapses by delivering electrical low-frequency stimulation at 1 Hz for 900 pulses.[135] In all long-term synaptic plasticity induction protocols, cells were current-clamped at –60 mV. EPSC peak amplitudes were measured before and after synaptic plasticity induction in voltage-clamp mode (V_{hold} = –60 mV) using paired electrical stimulation (10 Hz) delivered every 20 s for a 5-min baseline period and a 30-min postinduction period.

All electrophysiological experiments were analyzed offline using Clampfit 10.7 software. For all long-term synaptic plasticity experiments, raw EPSC amplitudes were measured, averaged per minute, and expressed as a percent change from baseline. The amplitude of the first EPSC peak was measured if a polysynaptic response was elicited. To determine a change in synaptic strength after the plasticity-induction protocols, the full postinduction time periods of all the cells in the experiment were averaged and compared to a theoretical baseline of 100% by using a one-sample *t*-test (GraphPad Prism 8.4.2), unless noted. To compare between experimental drug conditions, a one-way ANOVA with Dunnett's multiple comparisons post-hoc test was used. PPR was calculated by measuring the peak amplitude of the evoked EPSC from both pulses and dividing the EPSC2 peak amplitude by the EPSC1 peak amplitude. The PPR for each ISI of each synapse was compared against 1.0 by using a one-sample *t*-test and between TC and CT synapses by using an unpaired two-tailed *t*-test.

Two-photon calcium imaging

Two-photon calcium imaging was performed as described previously.[63] Briefly, two-photon laser-scanning microscopy was performed using an Ultima imaging system (Bruker), a Ti:sapphire Chameleon Ultra femtosecond-pulsed laser (Coherent, 820 nm) and 60× [0.9 numerical aperture] water-immersion infrared objectives (Olympus). Fluo-5F (300 µM) and Alexa 594 (10-25 µM) were included in the internal solution containing 115 mM potassium gluconate, 20 mM KCl, 10 mM HEPES, 4 mM $MgCl_2$, 0.1 mM EGTA, 4 mM ATP-Mg_2, 0.4 mM GTP-Na, 5 mM QX-314 chloride, and 10 mM creatine phosphate-Na_2 at pH 7.4 and 290-295 mOsm. Synaptically evoked changes in fluorescence of both fluorophores were measured in line-scan mode in a dendritic spine and the parent dendritic shaft. Line scans were analyzed as a ratio of normalized green (G) (Fluo-5F) fluorescence to normalized red (R) (Alexa Fluor 594) fluorescence (G/R). A line-scan was performed through every visible dendritic spine on a targeted dendritic branch, in an orientation that was parallel to the dendritic spine neck and orthogonal to the dendritic shaft.

Statistical analyses

Statistical tests were performed using Prism (Graphpad) or Sigmaplot (Systat) software. Statistical comparisons are noted in the text or figure legends. Unless otherwise noted, distributions were tested for normality (Shapiro-Wilk test) and equal variance (Brown-Forsythe test). If the distribution passed, a paired or unpaired t-test was performed. If it failed, a rank-sum test or signed-rank test was performed. To compare more than two distributions, a one-way or repeated-measures ANOVAs was performed. Significance was designated as $P < 0.05$. All data are presented as the mean ± SEM, and the sample size (N) is presented as the number of cells per the number of assembloids.

Drugs

1589

All salts for aCSF were purchased from Sigma-Aldrich. QX-314 chloride was purchased from

Hello Bio. DL-AP5 and MPEP were purchased from Tocris Bioscience. To create stock

solutions, MPEP was dissolved in DMSO and DL-AP5 was dissolved in water; both were kept

frozen at –20°C until dilution in aCSF to the final concentration. For iBAPTA experiments,

BAPTA tetra-potassium salt and BAPTA tetra-cesium salt were included in KGlu- and Cs-based

intracellular solutions, respectively, at 20 mM.

1596

1597