Task-irrelevant phase but not contrast variability unlocks generalization in visual perceptual

learning

Beyza Akkoyunlu[1,2,3,4], Caspar M. Schwiedrzik[1,2,3,*]

[1]Neural Circuits and Cognition Lab, European Neuroscience Institute Göttingen – A Joint Initiative of the

University Medical Center Göttingen and the Max Planck Society, Grisebachstraße 5, 37077 Göttingen, Germany;

[2]Perception and Plasticity Group, German Primate Center – Leibniz Institute for Primate Research, Kellnerweg

4, 37077 Göttingen, Germany; [3]Leibniz ScienceCampus "Primate Cognition", Göttingen, Germany; [4]Systems

Neuroscience Program, Graduate School for Neurosciences, Biophysics and Molecular Biosciences (GGNB) at the

Georg August University Göttingen, 37077 Göttingen, Germany

*To whom correspondence should be addressed: c.schwiedrzik@eni-g.de

ORCID IDs: 0000-0002-4401-1444 (BA), 0000-0003-0661-8859 (CMS)

Abstract

Performance on visual tasks can be improved by practice, a process called visual perceptual learning.

However, learning-induced performance improvements are often limited to the specific stimuli and

visual field locations used during training. Recent research has shown that variability along task-

irrelevant stimulus dimensions during training can reduce this specificity. This has been related to

higher stages of visual processing that harbor neurons which are invariant to the task-irrelevant

dimension. Here, we test whether task-irrelevant trial-by-trial variability in two visual features for

which invariances arise at different stages of processing, contrast and spatial phase, results in different

degrees of generalization in space in an orientation discrimination task. We find that randomizing

spatial phase results in complete generalization of learning to a new spatial location, contrary to

randomizing contrast. Our results thus suggest that the neural population undergoing plasticity in

visual perceptual learning is determined by the training task, which, in turn, affects generalization.

This lends further support to the hypothesis that task-irrelevant variability is an independent factor in

determining the specificity of perceptual learning.

Keywords

perceptual learning, orientation discrimination, variability, generalization, contrast, spatial phase

Statements and declarations

The funders had no role in study design, data collection and interpretation, decision to publish, or preparation of the manuscript. The authors declare no competing financial or non-financial interests.

Author contributions

Beyza Akkoyunlu: Conceptualization, Methodology, Software, Formal analysis, Investigation, Writing - Original Draft, Visualization. *Caspar M. Schwiedrzik:* Conceptualization, Methodology, Writing - Original Draft, Supervision, Project administration, Funding acquisition.

Acknowledgements

We would like to thank Jannik Rotzinger, Josefine-Lena Berger, and Tina Zahrie for their help in data collection and Giorgio Manenti for valuable insights during study design. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant agreement No. 802482, to CMS). BA is supported by a scholarship from the Studienstiftung des Deutschen Volkes. CMS is supported by the German Research Foundation Emmy Noether Program (SCHW1683/2-1).

Introduction

Our visual environment is ever-changing. In turn, our brain adapts to this challenge by maintaining the

ability to learn throughout life. This capacity also extends to our visual sense: through experience, we

can mould and even expand our perceptual abilities (Dosher & Lu, 2020). Yet, how we learn from

diverse experiences remains a lingering question.

In the laboratory setting, we stimulate learning in the visual system by employing visual perceptual

learning (VPL) paradigms, e.g., training participants on orientation discrimination tasks. While VPL

results in profound and long-lasting performance increases, these learning-induced gains are usually

highly specific to the training parameters (Lu & Dosher, 2022). For instance, even if only the location

or spatial frequency of the stimuli are changed after training, learning effects disappear (Fiorentini &

Berardi, 1980; Schoups et al., 1995). This characteristic of VPL is scientifically intriguing, but limits the

practical use of VPL, e.g., in vision restoration therapies.

Because generalization is often considered the ultimate goal of learning, recent research has asked

the question whether it is possible to develop training regimes that overcome the characteristic

specificity of VPL. One factor that has been shown to lead to generalizable learning effects in many

other domains such as motor, language, or category learning is *variability* (Douvis, 2005; Perry et al.,

2010; Vukatana et al., 2015). Variability may promote learning at a more abstract level and thus

prevent "overfitting", i.e., learning effects that are extremely specific to the training material (Raviv et

al., 2022). Indeed, our own recent research has shown that systematically inducing variability in a task-

irrelevant dimension can lead to more generalized learning effects even in visual orientation

discrimination (Manenti et al., 2023), a domain that had so far been considered to yield only extremely

specific VPL effects (Schoups et al., 1995; Shiu & Pashler, 1992). Although the results from this new

training strategy answer one question, they raise another: does all variability lead to generalization?

Whether learning is specific or generalizes is thought to depend on the stage of processing at which

learning occurs in the brain. *Specificity* may arise at early stages of processing, such as the primary

visual cortex (V1), where neurons respond with high specificity to a narrow range of stimulus features,

e.g., a particular orientation or spatial location (Fahle, 2005). Hence, if the tuning properties of these

neurons or their readout changes with learning, highly specific learning effect should ensue. On the

other hand, *generalization* is thought to depend on higher-level brain areas, where neurons respond

to a broader range of features and wider regions of the visual field (Ahissar & Hochstein, 2004).

Variability is thought to tap into the so-called invariance properties of these higher-order neurons:

invariant representations consistently respond to task-relevant features irrespective of variability in

other, task-irrelevant features. Lower brain areas could only achieve this by distributing plasticity over

a large number of neurons co-tuned to the task-relevant and the task-irrelevant feature. Hence, relying on invariant neurons provides task-relevant information in face of variability while simultaneously limiting the number of neurons or readout weights that need to undergo plasticity (Manenti et al., 2023).

To test this hypothesis, we trained participants on an orientation discrimination task, which is known to lead to highly specific learning effects (Schoups et al., 1995; Shiu & Pashler, 1992). During training, we introduced trial-by-trial variability in one of two task-irrelevant dimensions, the spatial phase or the contrast of the stimuli, respectively. We opted for these dimensions because they constitute fundamental features for visual processing (Gladilin & Eils, 2015; Shapley, 1985), yet, above threshold, they are not essential for the orientation discrimination task. Furthermore, *invariance* to these two features arises at different levels of processing. In particular, phase invariance appears later in the visual processing hierarchy than contrast invariance: for instance, contrast invariance arises already in simple cells, whereas phase invariance is first seen in V1 complex cells (Hubel & Wiesel, 1962). As neurons at higher stages of processing typically respond to larger regions of the visual field (i.e., they have larger receptive fields), we hypothesized that plasticity involving such neurons would result in less spatial specificity. We thus tested whether participants trained with phase randomized stimuli would achieve more transfer to untrained spatial locations than participants trained with contrast-randomized stimuli.

To preview, we find that phase and contrast randomization lead to differential spatial transfer after orientation discrimination learning. In particular, participants trained with phase randomized stimuli show generalization of learning effects in space that extend further than the learning effects of subjects trained with contrast randomized stimuli. This supports the idea that variability induces plasticity of neurons higher in the visual processing hierarchy and suggests that the specific feature in which variability occurs can determine at which stage of processing this plasticity occurs.

Materials and methods

Participants

Forty healthy volunteers (18 female, 4 left-handed, mean age 25 years, SD 2.66 years) participated in the experiment. All participants had normal or corrected-to-normal vision, reported no neurological or psychiatric diseases, and gave written informed consent before participation. Subjects were assigned to one of four experimental groups: phase randomized upper visual field (UVF) training (n=9) or contrast randomized UVF training (n=9), phase randomized lower visual field (LVF) training (n=11)

or contrast randomized LVF training (n=12). Participants received 12€/hour compensation, and to
ensure proper motivation, they received a 2€ bonus each day if they improved their orientation
discrimination threshold by at least 10% compared to the previous day. All procedures were in
accordance with the principles put forward by the Declaration of Helsinki and approved by the Ethics
Committee of the University Medical Center Göttingen (protocol number 29/8/17).

Stimuli and procedure

Subjects were trained with a two-alternative forced choice (2AFC) orientation discrimination task
using Gabor gratings (Figure 1A). Each participant participated in a pre- and a post-training threshold
measurement, respectively, and four training sessions. Each session was performed on a separate day,
and participants had to complete all sessions of the experiment within ten days. On average, subjects
took 7.26 days (SD 1.2 days) to complete the study.

The stimuli were presented on a gamma-corrected LCD monitor (ViewPixx EEG, resolution 1920 × 1080
px, refresh rate 120 Hz) using Psychtoolbox (Brainard, 1997) running in Matlab (R2018B, The
Mathworks). A circular aperture was placed on top of the screen to cover the monitor's corners in
order to prevent participants from using it as a reference while determining the orientation of the
gratings. Participants sat at a distance of 65 cm from the screen and performed the task using a
response box (Millikey SR-5). During the experiment, eye movements were recorded using a video-
based eye tracker (SR Research Eyelink 1000+). Data were collected from the left eye at 1000 Hz.

Gabor gratings (size 2 dva, spatial frequency 2.5 cpd) were placed at 6° eccentricity in the upper or
lower left visual field, respectively, on a grey background (52.42 cd/m^2 luminance). For the phase
randomized group, the gratings were presented at a constant 45% Michelson contrast, and the spatial
phase of the stimulus was randomized between each trial between 0° and 180°. In the contrast
randomized group, spatial phase of the stimulus was kept constant at 0°, and contrast was randomized
between 33% and 66% Michelson contrast.

In the threshold measurement sessions, participants performed three blocks of the 2AFC orientation
discrimination task at each of three locations (Figure 1B). Participants started each trial with fixating
for 400 ms, then a reference Gabor grating was presented for 92 ms. After a 600 ms inter-stimulus
interval, the test grating was presented for 92 ms. The reference grating had an orientation of 45° and
the orientation of the test grating was determined with a 3 down – 1 up staircase procedure. Except
for the orientation, reference and test gratings had identical parameters. Participants were instructed
to report the orientation of the test grating in comparison to the reference grating (clockwise or

counterclockwise). They were further instructed to respond as accurately and quickly as possible (but

no limit on the responses time was imposed). If fixation was broken during a trial (2° tolerance

window), the trial was aborted and immediately repeated. No feedback was provided.

Every block consisted of two interleaved staircases, one starting from a 1° offset and the other starting

from a 3° offset from the 45° reference with 0.05 log step size. The blocks ended after 10 reversals or

when 200 trials were completed (in a subset of session, more reversals were collected because of a

coding issue). The thresholds were measured at three isoeccentric locations (Figure 1B), namely,

training, close transfer (C-Transfer) and far transfer (F-Transfer). The distance between training – C-

Transfer and C-Transfer – F-Transfer locations was 5°, whereas the distance between training – F-

Transfer location was 11.5°. The order of testing at the three locations during the threshold

measurement sessions was counterbalanced between participants and kept identical for pre and post

training measurements.

Figure 1. Task design. A) *Each trial started with a 400 ms fixation period. Participants were then presented with the 45° reference grating (92 ms), followed by a 600 ms inter-stimulus interval, and then the test grating (92 ms). Participants were instructed to respond after the test grating disappeared. If participants broke fixation during the trial, the trial was aborted and repeated immediately.* **B)** *Training and transfer locations. All participants' thresholds were collected at the respective training location and two transfer location, namely, close (C-Transfer) and far (F-Transfer) transfer locations. Training and F-Transfer locations were switched for the groups who trained in the upper visual field and lower visual field, respectively.*

Subjects were trained at one of two locations (upper or lower visual field) and with one of two types

of task-irrelevant variability (spatial phase or contrast). In each training session, participants

performed five blocks of 200 trials of the 2AFC task, i.e., 1000 trials per training day, for a total of 4000

training trials. The task design was identical to the threshold measurements, but participants now

received feedback: error feedback as a high-pitch sound through headphones (Sennheiser HDA 280)

after a mistake in a given trial; in addition, upon finishing the block, the percentage of correct trials

was displayed on the screen. Before each training session, participants were informed whether they

had earned the financial performance bonus for the preceding training session in order to ensure

continuous motivation.

Analyses

One participant had to be excluded because he/she fell asleep during the experiment. Eleven

participants were excluded because they did not learn (LI 20% or worse). There was no statistically

significant difference between number of participants excluded from each experimental group (Fisher

exact test, p = 0.490, odds ratio = 0.201). The final n was 29 (14 female, 2 left-handed, mean age 24.5,

SD 2.31).

All statistical analyses were conducted in R (version 4.2.2; R Core Team, 2022) on macOS Monterey

12.0.1.

Each subject performed three threshold blocks at each retinal location pre and post learning. To

calculate the threshold of each block, the first four reversals of the staircase were discarded, and the

mean of the following six reversals was taken. Blocks with an insufficient number of reversals were

discarded. To determine how learning affected orientation discrimination thresholds at the training

and transfer locations, we fitted a linear mixed effects model to the block-wise threshold data using

the lme4 package (version 1.1-34) (Bates et al., 2015). Data were aligned and rank transformed

(Higgins & Tashtoush, 1994) using the ARTool package (version 0.11.1) (Wobbrock et al., 2011), to

satisfy distributional assumptions. In the linear mixed effects model, we predicted thresholds with the

factors *time* (2 levels, pre/post), *randomization* (2 levels, phase/contrast), *location* (3 levels,

training/C-transfer/F-transfer) and *training location* (2 levels, upper/lower visual field) as fixed effects,

and participants as random intercept, as in Equation 1:

$$\textbf{Threshold} \sim \textbf{Time} * \textbf{Randomization} * \textbf{Location} * \textbf{Training Location} + (\textbf{1}|\textbf{Subject}) \qquad \textit{(1)}$$

To obtain p-values, we performed a repeated measures analyses of variance (rmANOVA) of the model

terms and their interactions using Wald's F-test and degrees of freedom determined following

Kenward and Roger (Kenward & Roger, 1997). Posthoc tests were performed on estimated marginal

means of the aligned and rank transformed data (Elkin et al., 2021) using the emmeans package

(version 1.8.8) (Lenth, 2023).

For quantifying learning, we also calculated the Learning Index (LI) (Xiao et al., 2008), as in Equation

2:

$$\text{Learning Index(LI)} = \frac{\text{Threshold}_{\text{Post}} - \text{Threshold}_{\text{Pre}}}{\text{Threshold}_{\text{Pre}}} \qquad (2)$$

LIs were used to test whether transfer to the C-Transfer location correlated with transfer to the F-
Transfer location across subjects; a one-sided Fisher's z-test was used to compare the correlation
coefficients between groups via the cocor package (version 1.1.4) (Diedenhofen & Musch, 2015).

To determine effect sizes, we calculated partial η^2 for rmANOVAs, Cohen's d for t-tests, and Cohen's
q for comparing correlation coefficients using the effectsize package (version 0.8.6) (Ben-Shachar et
al., 2020). Plots were prepared using the ggpubr (version 0.6.0) (Kassambara, 2023) and ggplot2
(version 3.4.4) (Wickham, 2016) packages.

Data availability

All data will be made available in a publicly accessible online repository upon acceptance of the
manuscript.

Results

We first examined whether there were any pre-existing differences between the contrast and phase
randomization groups in terms of their pretraining orientation discrimination thresholds. This was not
the case: at none of the three locations, there were statistically significant differences in pre-training
thresholds between the contrast and phase randomized groups (Welch's two-sample t-test, Training
location t(83.484) = 0.790, p = 0.4313, Cohen's d = 0.17; C-Transfer t(83.558) = 0.049, p = 0.9505,
Cohen's d = 0.01; F-Transfer t(82.089) = -1.512, p = 0.1344, Cohen's d = 0.33). Hence, any differences
in learning and/or transfer between the groups cannot be due to pre-existing differences in
orientation discrimination.

Upon training, both randomization groups significantly improved in discriminating orientations at the
training location: the group trained with phase randomization improved thresholds on average by 1.53
deg (paired t-test, t(38) = 8.025, p < 0.001, Cohen's d = 1.30), and the group trained with contrast
randomization improved on average by 1.40 deg (t(46) = 7.981, p < 0.001, Cohen's d = 1.18). We then
assessed whether learning and transfer differed between the groups considering all three locations
(Figure 2). Comparing learning (i.e., the pre-post difference in thresholds) at the training location, we

*Figure 2. Pre and post learning thresholds at the training and transfer locations. **A)** Pre and post learning thresholds at all locations for phase (orange) and contrast (green) randomization. Each dot represents a subject's mean threshold at the respective location and time point. Bars show the mean thresholds over subjects in each group and location. At the training location, there was no statistically significant difference between the phase randomized and the contrast randomized group (t(469) = 1.062, p = 0.2888, Cohen's d = 0.10). Transfer of learning to the C-Transfer location also did not reveal a statistically significant difference between the two groups (t(469) = 0.936, p = 0.3495, Cohen's d = 0.09). However, there was significantly more transfer to the F-Transfer location in the phase randomized group than the contrast randomized group (t(469) = 2.910, p = 0.0038, Cohen's d = 0.27). **B)** Change in the thresholds with learning. Change in the threshold calculated per subject, by subtracting mean post-learning threshold from mean pre-learning threshold. In both panels, error bars represent the standard error of the mean.*

found no statistically significant difference between the phase randomized and the contrast randomized group (t(469) = 1.062, p = 0.2888, Cohen's d = 0.10). This suggests that despite different kinds of variability, training was similarly effective at this location. Transfer of learning to the C-Transfer location (again in terms of the pre-post difference in threshold) also did not reveal a statistically significant difference between the two groups (t(469) = 0.936, p = 0.3495, Cohen's d = 0.09). This suggests that training transferred similarly to a close by location with both randomization manipulations. However, there was significantly more transfer to the F-Transfer location in the phase randomized group than the contrast randomized group (t(469) = 2.910, p = 0.0038, Cohen's d = 0.27): while thresholds improved on average only by 1.31 deg in the contrast randomized group, the average improvement was 1.99 deg in the phase randomized group. This suggests that training effects generalize more broadly in space with phase than with contrast variability during training. This differential effect was also supported by a significant time × location × randomization interaction (F(2,469.031) = 3.317, p = 0.037, η^2_p = 0.01, Supplementary Table 1 and Supplementary Figure S1).

Overall, these results support the hypothesis that training with phase variability engages neurons with

larger receptive fields than training with contrast variability.

Another prediction that can be derived from the hypothesis that phase randomization and contrast

randomization rely on differently sized receptive fields is that learning effects should correlate

between spatial locations if both locations fall into the same receptive fields, whereas they should not

correlate if neurons with small, non-overlapping receptive fields are involved (Kondat et al., 2023). We

thus investigated the correlation of LIs at the two transfer locations for both randomization groups

(Figure 3). We find that LIs at the C-Transfer and F-Transfer locations correlate highly in the phase

randomization group (Pearson's correlation, r = 0.71, t(11) = 3.39, p = 0.006). In contrast, there was

no statistically significant correlation of LIs between the two transfer locations in the contrast

randomization group (Pearson's correlation, r = 0.09, t(14) = 0.373, p=0.71). Furthermore, the

correlation in LIs was significantly larger in the phase than in the contrast randomized group (z =

1.8964, p = 0.0290, Cohen's q = 0.79). This provides further evidence that the two groups rely on

different neural populations for learning orientation discrimination.

Figure 3. Correlation of LIs between the two transfer locations. In the contrast randomized group, LIs at the C-Transfer and F-Transfer locations are uncorrelated (r = 0.09, p = 0.71). In contrast, LIs are correlated in the phase randomized group (r = 0.71, p = 0.006). The correlation in LIs was significantly larger in the phase than in the contrast randomized group (z = 1.8964, p = 0.0290, Cohen's q = 0.79). Shaded areas represent the 95% confidence intervals. Note that different scaling is used in each panel to ease comparison.

Discussion

We find that variability in spatial phase during training leads to more generalization of VPL in space

than variability in contrast. Moreover, participants trained with variable spatial phase transfer learning

similarly across two transfer locations, while subjects trained with variable contrast do not. Taken

together, these two findings support the hypothesis that the two forms of variability tap into neurons

with differently large receptive fields, and in turn, that the type of variability introduced during training
can determine the stage of processing at which VPL takes place.

When our visual system is confronted with variability in a task-irrelevant dimension, it can rely on
neurons that are invariant to this feature and thus subsume this variability (Manenti et al., 2023); such
neurons typically reside at a higher stage of processing and have large receptive fields. Alternatively,
the visual system can make use of neurons at earlier stages of processing that are conjunctively tuned
to the task-relevant and the task-irrelevant features of the stimuli. Because these neurons typically
have spatially restricted receptive fields, they cannot, by themselves, enable generalization across the
visual field. Importantly, different invariances arise at different levels of processing. Hence, creating
variability in different task-irrelevant features of a stimulus should result in different patterns of
generalization. This is what we find for the case of spatial phase and contrast variability.

Relevant to our experiments, it is known that contrast invariance arises earlier in the visual processing
hierarchy than phase invariance: contrast invariance gradually increase along the visual ventral stream
(Avidan et al., 2002), but is already seen at the level of simple cells in V1 (Hubel & Wiesel, 1962). Phase
invariance arises on the level of complex cells (Hubel & Wiesel, 1962). Both simple and complex cells
are orientation tuned. According to a classical model of orientation tuning in V1, simple cells feed into
complex cells (Hubel & Wiesel, 1962), and there is empirical evidence that supports this view (e.g.,
Alonso & Martinez, 1998). This principle also underlies hierarchical models of visual processing such
as HMAX (Serre & Riesenhuber, 2004). Furthermore, simple cells are predominantly found in the
granular input layer of V1, whereas complex cells are found in supra- and infragranular output layers
(Hubel & Wiesel, 1968). Hence, complex cells reside on a higher level of processing than simple cells.
Importantly, complex cells also have larger receptive fields than simple cells (Schiller et al., 1976) and
could thus support transfer that extends further in the visual field than simple cells. Of note, simple
and complex cells are not only found in V1, but also at higher stages of processing such as V2 (Levitt
et al., 1994) and V4 (Desimone & Schein, 1987), which could possibly support the large distances
between the different locations in our experiments with their larger receptive fields. Hence, the
principle that phase invariance arises at a higher stage of processing than contrast invariance holds
irrespective of where exactly plasticity occurs. Our results are thus in line with the hypothesis that the
dimension in which task-irrelevant variability occurs determines the stage of processing affected by
VPL, reflected by varying degrees of generalization.

Previous VPL studies have used phase or contrast randomization not as an experimental variable, but
to assure that subjects are not using spatial cues to solve orientation discrimination tasks (in the case
of spatial phase) and/or to reduce adaptation. Some of these studies have reported spatial

generalization effects (e.g., Zhang et al., 2010). In contrast, a study by Sowden and colleagues (2002)

reported training effects that were extremely specific to the spatial frequency, retinal location and

even eye of origin, consistent with learning effects in the granular input layer of V1 where neurons

which such tuning properties reside. Importantly, this study used a training procedure with no

variability in which only a single grating was presented for several thousand trials. Notably, a study

investigating the neural basis of orientation discrimination learning in macaque monkey V1 used phase

randomization and reported learning effects only outside the granular input layer (Schoups et al.,

2001), which would be compatible with an outsized involvement of complex cells. Overall, this

suggests that by using variability during training, previous studies may have coincidentally engaged

neurons at higher stages of processing and with larger receptive fields.

Apart from variability, spatial generalization of VPL also occurs when adaptation is reduced or

eliminated (Harris et al., 2012). While spatial transfer to the C-Transfer location we found in both

randomization groups may be explained by a reduction of adaptation with variability, the differential

transfer effects at the F-Transfer location and the differential correlation between learning effects at

the two transfer locations in the different groups are not easily accounted for by adaptation. Previous

studies have also shown that task difficulty affects generalization (Ahissar & Hochstein, 2004).

However, as pretraining thresholds did not differ significantly between contrast and phase

randomization, this factor also cannot easily explain our differential transfer results. Hence, variability

may contribute to generalization in VPL beyond the well-established factors task difficulty and

adaptation. On the other hand, studies using contrast randomization during orientation discrimination

learning in the context of double training (Xiao et al., 2008) and manipulation of attention (Donovan

& Carrasco, 2018) have reported sizable spatial transfer effects. Hence, variability is not the only factor

that enables spatial transfer after perceptual learning, but there are alternative paths of achieving

generalization.

Taken together, our results emphasize a role of targeted variability in achieving spatial generalization

and suggest higher stages of visual processing as a possible locus of VPL. Variability may thus be a

useful component for improving real life applications of VPL without burdening the learner with

additional training time and effort.

References

Ahissar, M., & Hochstein, S. (2004). The reverse hierarchy theory of visual perceptual learning. *Trends in cognitive sciences*, *8*(10), 457-464. https://doi.org/10.1016/j.tics.2004.08.011

Alonso, J. M., & Martinez, L. M. (1998). Functional connectivity between simple cells and complex cells in cat striate cortex. *Nature neuroscience*, *1*(5), 395-403. https://doi.org/10.1038/1609

Avidan, G., Harel, M., Hendler, T., Ben-Bashat, D., Zohary, E., & Malach, R. (2002). Contrast sensitivity in human visual areas and its relationship to object recognition. *Journal of neurophysiology*, *87*(6), 3102-3116. https://doi.org/10.1152/jn.2002.87.6.3102

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

Ben-Shachar, M., Lüdecke, D., & Makowski, D. (2020). effectsize: Estimation of Effect Size Indices and Standardized Parameters. *Journal of Open Source Software*, *5*(56), 2815. https://doi.org/10.21105/joss.02815

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial vision*, *10*(4), 433-436. https://www.ncbi.nlm.nih.gov/pubmed/9176952

Desimone, R., & Schein, S. J. (1987). Visual properties of neurons in area V4 of the macaque: sensitivity to stimulus form. *Journal of neurophysiology*, *57*(3), 835-868. https://doi.org/10.1152/jn.1987.57.3.835

Diedenhofen, B., & Musch, J. (2015). cocor: A Comprehensive Solution for the Statistical Comparison of Correlations. *PLoS One*, *10*(4), e0121945. https://doi.org/10.1371/journal.pone.0121945

Donovan, I., & Carrasco, M. (2018). Endogenous spatial attention during perceptual learning facilitates location transfer. *Journal of vision*, *18*(11), 7. https://doi.org/10.1167/18.11.7

Dosher, B. A., & Lu, Z. L. (2020). *Perceptual learning. How experience shapes visual perception*. MIT Press.

Douvis, S. J. (2005). Variable Practice in Learning the Forehand Drive in Tennis. *Perceptual and motor skills*, *101*(2), 531-545. https://doi.org/10.2466/pms.101.2.531-545

Elkin, L. A., Kay, M., Higgins, J. J., & Wobbrock, J. O. (2021). An Aligned Rank Transform Procedure for Multifactor Contrast Tests. UIST '21: The 34th Annual ACM Symposium on User Interface Software and Technology, Virtual Event.

Fahle, M. (2005). Perceptual learning: specificity versus generalization. *Current opinion in neurobiology*, *15*(2), 154-160. https://doi.org/10.1016/j.conb.2005.03.010

Fiorentini, A., & Berardi, N. (1980). Perceptual learning specific for orientation and spatial frequency. *Nature*, *287*(5777), 43-44. https://doi.org/10.1038/287043a0

Gladilin, E., & Eils, R. (2015). On the role of spatial phase and phase correlation in vision, illusion, and cognition. *Front Comput Neurosci*, *9*, 45. https://doi.org/10.3389/fncom.2015.00045

Harris, H., Gliksberg, M., & Sagi, D. (2012). Generalized perceptual learning in the absence of sensory adaptation. *Current biology*, *22*(19), 1813-1817. https://doi.org/10.1016/j.cub.2012.07.059

Higgins, J. J., & Tashtoush, S. (1994). An aligned rank transform test for interaction. *Nonlinear World*, *1*, 201-211.

Hubel, D. H., & Wiesel, T. N. (1962). Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. *Journal of physiology*, *160*(1), 106-154. https://doi.org/10.1113/jphysiol.1962.sp006837

Hubel, D. H., & Wiesel, T. N. (1968). Receptive fields and functional architecture of monkey striate cortex. *Journal of physiology*, *195*(1), 215-243. https://doi.org/10.1113/jphysiol.1968.sp008455

Kassambara, A. (2023). *ggpubr: 'ggplot2' Based Publication Ready Plots*. https://CRAN.R-project.org/package=ggpubr

Kenward, M. G., & Roger, J. H. (1997). Small Sample Inference for Fixed Effects from Restricted Maximum Likelihood. *Biometrics*, *53*(3). https://doi.org/10.2307/2533558

Kondat, T., Aderka, M., & Censor, N. (2023). Modulating temporal dynamics of performance across retinotopic locations enhances the generalization of perceptual learning. *iScience*, *26*(11), 108276. https://doi.org/10.1016/j.isci.2023.108276

Lenth, R. V. (2023). *emmeans: Estimated Marginal Means, aka Least-Squares Means*. https://CRAN.R-project.org/package=emmeans

Levitt, J. B., Kiper, D. C., & Movshon, J. A. (1994). Receptive fields and functional architecture of macaque V2. *Journal of neurophysiology*, *71*(6), 2517-2542. https://doi.org/10.1152/jn.1994.71.6.2517

Lu, Z.-L., & Dosher, B. A. (2022). Current directions in visual perceptual learning. *Nature Reviews Psychology*, *1*(11), 654-668. https://doi.org/10.1038/s44159-022-00107-2

Manenti, G. L., Dizaji, A. S., & Schwiedrzik, C. M. (2023). Variability in training unlocks generalization in visual perceptual learning through invariant representations. *Current biology*, *33*(5), 817-826 e813. https://doi.org/10.1016/j.cub.2023.01.011

Perry, L. K., Samuelson, L. K., Malloy, L. M., & Schiffer, R. N. (2010). Learn Locally, Think Globally: Exemplar Variability Supports Higher-Order Generalization and Word Learning. *Psychological science*, *21*(12), 1894-1902. https://doi.org/10.1177/0956797610389189

Raviv, L., Lupyan, G., & Green, S. C. (2022). How variability shapes learning and generalization. *Trends in cognitive sciences*, *26*(6), 462-483. https://doi.org/10.1016/j.tics.2022.03.007

Schiller, P. H., Finlay, B. L., & Volman, S. F. (1976). Quantitative studies of single-cell properties in
monkey striate cortex. I. Spatiotemporal organization of receptive fields. *Journal of*
neurophysiology, *39*(6), 1288-1319. https://doi.org/10.1152/jn.1976.39.6.1288

Schoups, A. A., Vogels, R., & Orban, G. A. (1995). Human perceptual learning in identifying the oblique
orientation: retinotopy, orientation specificity and monocularity. *Journal of physiology*, *483*
(Pt 3), 797-810. https://doi.org/10.1113/jphysiol.1995.sp020623

Schoups, A. A., Vogels, R., Qian, N., & Orban, G. A. (2001). Practising orientation identification
improves orientation coding in V1 neurons. *Nature*, *412*(6846), 549-553.
https://doi.org/10.1038/35087601

Serre, T., & Riesenhuber, M. (2004). Realistic Modeling of Simple and Complex Cell Tuning in the HMAX
Model, and Implications for Invariant Object Recognition in Cortex. *CSAIL Technical Reports*,
052. http://hdl.handle.net/1721.1/30491

Shapley, R. (1985). The importance of contrast in the perception of brightness and form. *Transactions*
of the American Philosophical Society, *75*(6), 20-29.

Shiu, L. P., & Pashler, H. (1992). Improvement in line orientation discrimination is retinally local but
dependent on cognitive set. *Perception & psychophysics*, *52*(5), 582-588.
https://doi.org/10.3758/bf03206720

Sowden, P. T., Rose, D., & Davies, I. R. (2002). Perceptual learning of luminance contrast detection:
specific for spatial frequency and retinal location but not orientation. *Vision research*, *42*(10),
1249-1258. https://doi.org/10.1016/s0042-6989(02)00019-6

Vukatana, E., Graham, S. A., Curtin, S., & Zepeda, M. S. (2015). One is Not Enough: Multiple Exemplars
Facilitate Infants' Generalizations of Novel Properties. *Infancy*, *20*(5), 548-575.
https://doi.org/10.1111/infa.12092

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer.
https://doi.org/10.1007/978-3-319-24277-4

Wobbrock, J. O., Findlater, L., Gergle, D., & Higgins, J. J. (2011). The Aligned Rank Transform for
Nonparametric Factorial Analyses Using Only Anova Procedures. *CHI '11: Proceedings of the*
SIGCHI Conference on Human Factors in Computing Systems, 143–146.
https://doi.org/10.1145/1978942.1978963

Xiao, L. Q., Zhang, J. Y., Wang, R., Klein, S. A., Levi, D. M., & Yu, C. (2008). Complete transfer of
perceptual learning across retinal locations enabled by double training. *Current biology*,
18(24), 1922-1926. https://doi.org/10.1016/j.cub.2008.10.030

Zhang, T., Xiao, L. Q., Klein, S. A., Levi, D. M., & Yu, C. (2010). Decoupling location specificity from

perceptual learning of orientation discrimination. *Vision research*, *50*(4), 368-374.

https://doi.org/10.1016/j.visres.2009.08.024

417