Task-irrelevant phase but not contrast variability unlocks generalization in visual perceptual
learning

Beyza Akkoyunlu[1,2,3,4], Caspar M. Schwiedrzik[1,2,3,5*]

[1]Neural Circuits and Cognition Lab, European Neuroscience Institute Göttingen – A Joint Initiative of the
University Medical Center Göttingen and the Max Planck Society, Grisebachstraße 5, 37077 Göttingen,
Germany; [2]Perception and Plasticity Group, German Primate Center – Leibniz Institute for Primate Research,
Kellnerweg 4, 37077 Göttingen, Germany; [3]Leibniz ScienceCampus "Primate Cognition", Göttingen, Germany;
[4]Systems Neuroscience Program, Graduate School for Neurosciences, Biophysics and Molecular Biosciences
(GGNB) at the Georg August University Göttingen, 37077 Göttingen, Germany; [5]Department of Cognitive
Neurobiology, Faculty of Biology and Biotechnology, Ruhr-University Bochum, 44801 Bochum, Germany

*To whom correspondence should be addressed: cschwiedrzik@dpz.eu

ORCID IDs: 0000-0002-4401-1444 (BA), 0000-0003-0661-8859 (CMS)

Abstract

Performance on visual tasks can be improved by practice, a process called visual perceptual learning.
However, learning-induced performance improvements are often limited to the specific stimuli and
visual field locations used during training. Recent research has shown that variability along task-
irrelevant stimulus dimensions during training can reduce this specificity. This has been related to
higher stages of visual processing that harbor neurons which are invariant to the task-irrelevant
dimension. Here, we test whether task-irrelevant trial-by-trial variability in two visual features for
which invariances arise at different stages of processing, contrast and spatial phase, results in
different degrees of generalization in space in an orientation discrimination task. We find that
randomizing spatial phase results in complete generalization of learning to a new spatial location,
contrary to randomizing contrast. Our results thus suggest that the neural population undergoing
plasticity in visual perceptual learning is determined by the training task, which, in turn, affects
generalization. This lends further support to the hypothesis that task-irrelevant variability is an
independent factor in determining the specificity of perceptual learning.

Keywords

31 perceptual learning, orientation discrimination, variability, generalization, contrast, spatial phase

Declarations

Funding

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant agreement No. 802482, to CMS). BA is supported by a scholarship from the Studienstiftung des Deutschen Volkes. CMS is supported by the German Research Foundation Emmy Noether Program (SCHW1683/2-1). The funders had no role in study design, data collection and interpretation, decision to publish, or preparation of the manuscript.

Conflict of interest

The authors have no conflict of interest to declare.

Ethical approval

All procedures in this study were in accordance with the principles put forward by the Declaration of Helsinki and approved by the Ethics Committee of the University Medical Center Göttingen (protocol number 29/8/17).

Informed consent

Informed consent was obtained from all individual participants included in the study.

Consent for publication

Consent for publication was obtained for every individual person's data included in the study.

Data availability

All data will be made available in a publicly accessible online repository upon acceptance of the manuscript.

Author contributions

Beyza Akkoyunlu: Conceptualization, Methodology, Software, Formal analysis, Investigation, Writing

- Original Draft, Visualization. *Caspar M. Schwiedrzik:* Conceptualization, Methodology, Writing -

Original Draft, Supervision, Project administration, Funding acquisition.

Acknowledgements

We would like to thank Jannik Rotzinger, Josefine-Lena Berger, and Tina Zahrie for their help in data

collection, Igor Kagan for helpful discussions, and Giorgio Manenti for valuable insights during study

design.

Introduction

Our visual environment is ever-changing. In turn, our brain adapts to this challenge by maintaining the ability to learn throughout life. This capacity also extends to our visual sense: through experience, we can mould and even expand our perceptual abilities (Dosher & Lu, 2020). Yet, how we learn from diverse experiences remains a lingering question.

In the laboratory setting, we stimulate learning in the visual system by employing visual perceptual learning (VPL) paradigms, e.g., training participants on orientation discrimination tasks. While VPL results in profound and long-lasting performance increases, these learning-induced gains are usually highly specific to the training parameters (Lu & Dosher, 2022). For instance, even if only the location or spatial frequency of the stimuli are changed after training, learning effects disappear (Fiorentini & Berardi, 1980; Schoups et al., 1995). This characteristic of VPL is scientifically intriguing, but limits the practical use of VPL, e.g., in vision restoration therapies.

Because generalization is often considered the ultimate goal of learning, recent research has asked the question whether it is possible to develop training regimes that overcome the characteristic specificity of VPL. One factor that has been shown to lead to generalizable learning effects in many other domains such as motor, language, or category learning is *variability* (Douvis, 2005; Perry et al., 2010; Vukatana et al., 2015). Variability may promote learning at a more abstract level and thus prevent "overfitting", i.e., learning effects that are extremely specific to the training material (Raviv et al., 2022). Indeed, our own recent research has shown that systematically inducing variability in a task-irrelevant dimension can lead to more generalized learning effects even in visual orientation discrimination (Manenti et al., 2023), a domain that had so far been considered to yield only extremely specific VPL effects (Schoups et al., 1995; Shiu & Pashler, 1992). Although the results from this new training strategy answer one question, they raise another: does all variability lead to generalization?

Whether learning is specific or generalizes is thought to depend on the stage of processing at which learning occurs in the brain. *Specificity* may arise at early stages of processing, such as the primary visual cortex (V1), where neurons respond with high specificity to a narrow range of stimulus features, e.g., a particular orientation or spatial location (Fahle, 2005). Hence, if the tuning properties of these neurons or their readout changes with learning, highly specific learning effect should ensue. On the other hand, *generalization* is thought to depend on higher-level brain areas, where neurons respond to a broader range of features and wider regions of the visual field (Ahissar & Hochstein, 2004). *Variability* is thought to tap into the so-called invariance properties of these higher-order neurons: invariant representations consistently respond to task-relevant features

irrespective of variability in other, task-irrelevant features. Lower brain areas could only achieve this
by distributing plasticity over a large number of neurons co-tuned to the task-relevant and the task-
irrelevant feature. Hence, relying on invariant neurons could provide task-relevant information in
face of variability while simultaneously limiting the number of neurons or readout weights that need
to undergo plasticity (Manenti et al., 2023). Alternatively, plasticity could always occur at the lowest
level of processing at which the relevant feature dimension is differentially encoded (Karni & Bertini,
1997).

To contrast these two hypotheses, we trained participants on an orientation discrimination task,
which is known to lead to highly specific learning effects (Schoups et al., 1995; Shiu & Pashler, 1992).
During training, we introduced trial-by-trial variability in one of two task-irrelevant dimensions, the
spatial phase or the contrast of the stimuli, respectively. We opted for these dimensions because
they constitute fundamental features for visual processing (Gladilin & Eils, 2015; Shapley, 1985), yet,
above threshold, they are not essential for the orientation discrimination task. Furthermore,
invariance to these two features arises at different levels of processing. In particular, phase
invariance appears later in the visual processing hierarchy than contrast invariance: for instance,
contrast invariance arises already in simple cells, whereas phase invariance is first seen in V1
complex cells (Hubel & Wiesel, 1962) and fully emerges in V2 (Cloherty & Ibbotson, 2015). As
neurons at higher stages of processing typically respond to larger regions of the visual field (i.e., they
have larger receptive fields), we hypothesized that plasticity involving such neurons would result in
less spatial specificity. Alternatively, given that both contrast and spatial phase are not essential to
orientation processing above threshold, both randomization types could result in a comparable
degree of spatial specificity if plasticity always occurs at the lowest stage of processing where the
task-relevant feature is encoded. We thus tested whether participants trained with phase
randomized stimuli would achieve more transfer to untrained spatial locations than participants
trained with contrast-randomized stimuli.

To preview, we find that phase and contrast randomization lead to differential spatial transfer after
orientation discrimination learning. In particular, participants trained with phase randomized stimuli
show generalization of learning effects in space that extend further than the learning effects of
subjects trained with contrast randomized stimuli. This supports the idea that variability induces
plasticity of neurons higher in the visual processing hierarchy and suggests that the specific feature
in which variability occurs can determine at which stage of processing this plasticity occurs.

Materials and methods

 Participants

Forty healthy volunteers (18 female, 4 left-handed, mean age 25 years, SD 2.66 years) participated in
the experiment. All participants had normal or corrected-to-normal vision, reported no neurological
or psychiatric diseases, and gave written informed consent before participation. Subjects were
assigned to one of four experimental groups: phase randomized upper visual field (UVF) training
(n=9) or contrast randomized UVF training (n=9), phase randomized lower visual field (LVF) training
(n=11) or contrast randomized LVF training (n=12). Participants received 12€/hour compensation,
and to ensure proper motivation, they received a 2€ bonus each day if they improved their
orientation discrimination threshold by at least 10% compared to the previous day. All procedures
were in accordance with the principles put forward by the Declaration of Helsinki and approved by
the Ethics Committee of the University Medical Center Göttingen (protocol number 29/8/17).

Stimuli and procedure

Subjects were trained with a two-alternative forced choice (2AFC) orientation discrimination task
using Gabor gratings (Figure 1A).   Each participant participated in a pre- and a post-training
threshold measurement, respectively, and four training sessions. Each session was performed on a
separate day, and participants had to complete all sessions of the experiment within ten days. On
average, subjects took 7.26 days (SD 1.2 days) to complete the study.

The stimuli were presented on a gamma-corrected LCD monitor (ViewPixx EEG, resolution 1920 ×
1080 px, refresh rate 120 Hz) using Psychtoolbox (Brainard, 1997) running in Matlab (R2018B, The
Mathworks). A circular aperture was placed on top of the screen to cover the monitor's corners in
order to prevent participants from using it as a reference while determining the orientation of the
gratings. Participants sat at a distance of 65 cm from the screen and performed the task using a
response box (Millikey SR-5). During the experiment, eye movements were recorded using a video-
based eye tracker (SR Research Eyelink 1000+). Data were collected from the left eye at 1000 Hz.

Our experimental stimuli and procedure parameters closely followed a study by Xiao and colleagues
(2008). Gabor gratings (size 2 dva, spatial frequency 2.5 cpd, standard deviation of the Gaussian 0.28
deg) were placed at 6° eccentricity in the upper or lower left visual field, respectively, on a grey
background (52.42 cd/m^2 luminance). For the phase randomized group, the gratings were presented
at a constant 45% Michelson contrast, and the spatial phase of the stimulus was randomized
between each trial between 0° and 180°. In the contrast randomized group, spatial phase of the

Figure 1. Task design. A) Each trial started with a 400 ms fixation period. Participants were then presented with the 45° reference grating (92 ms), followed by a 600 ms inter-stimulus interval, and then the test grating (92 ms). Participants were instructed to respond after the test grating disappeared. If participants broke fixation during the trial, the trial was aborted and repeated immediately. *B)* Training and transfer locations. All participants' thresholds were collected at the respective training location and two transfer location, namely, close (C-Transfer) and far (F-Transfer) transfer locations. Training and F-Transfer locations were switched for the groups who trained in the upper visual field and lower visual field, respectively.

stimulus was kept constant at 0°, and contrast was randomized between 33% and 66% Michelson

contrast. The fixation dot had a size of 0.23 dva.

In the threshold measurement sessions, participants performed three blocks of the 2AFC orientation

discrimination task at each of three locations (Figure 1B). Participants started each trial with fixating

for 400 ms, then a reference Gabor grating was presented for 92 ms. After a 600 ms inter-stimulus

interval, the test grating was presented for 92 ms. Our stimulus timing (92 ms), taken from Xiao and

colleagues (2008), falls squarely into the range of previously used stimulus durations, which vary

between 33 ms (Lu & Dosher, 2004) and 500 ms (Tan et al., 2019) in the VPL literature. The

reference grating had an orientation of 45° and the orientation of the test grating was determined

with a 3 down – 1 up staircase procedure. Except for the orientation, reference and test gratings had

identical parameters. Participants were instructed to report the orientation of the test grating in

comparison to the reference grating (clockwise or counterclockwise). They were further instructed

to respond as accurately and quickly as possible (but no limit on the responses time was imposed). If

fixation was broken during a trial (2° tolerance window), the trial was aborted and immediately

repeated. No feedback was provided.

Every block consisted of two interleaved staircases, one starting from a 1° offset and the other

starting from a 3° offset from the 45° reference with 0.05 log step size. The blocks ended after 10

reversals or when 200 trials were completed (in a subset of session, more reversals were collected

because of a coding issue). The thresholds were measured at three isoeccentric locations (Figure 1B),

namely, training, close transfer (C-Transfer) and far transfer (F-Transfer). The distance between

training – C-Transfer and C-Transfer – F-Transfer locations was 5°, whereas the distance between

training – F-Transfer location was 11.5°. The order of testing at the three locations during the
threshold measurement sessions was counterbalanced between participants and kept identical for
pre and post training measurements.

Subjects were trained at one of two locations (upper or lower visual field) and with one of two types
of task-irrelevant variability (spatial phase or contrast). In each training session, participants
performed five blocks of 200 trials of the 2AFC task, i.e., 1000 trials per training day, for a total of
4000 training trials. The task design was identical to the threshold measurements, but participants
now received feedback: error feedback as a high-pitch sound through headphones (Sennheiser HDA
280) after a mistake in a given trial; in addition, upon finishing the block, the percentage of correct
trials was displayed on the screen. Before each training session, participants were informed whether
they had earned the financial performance bonus for the preceding training session in order to
ensure continuous motivation.

Analyses

One participant had to be excluded because he/she fell asleep during the experiment. Eleven
participants were excluded because they did not learn (LI 20% or worse). There was no statistically
significant difference between number of participants excluded from each experimental group
(Fisher exact test, p = 0.490, odds ratio = 0.201). The final n was 29 (14 female, 2 left-handed, mean
age 24.5, SD 2.31).

All statistical analyses were conducted in R (version 4.2.2; R Core Team, 2022) on macOS Monterey
12.0.1.

Each subject performed three threshold blocks at each retinal location pre and post learning. To
calculate the threshold of each block, the first four reversals of the staircase were discarded, and the
mean of the following six reversals was taken. Blocks with an insufficient number of reversals were
discarded. To determine how learning affected orientation discrimination thresholds at the training
and transfer locations, we fitted a linear mixed effects model to the block-wise threshold data using
the lme4 package (version 1.1-34) (Bates et al., 2015). Data were aligned and rank transformed
(Higgins & Tashtoush, 1994) using the ARTool package (version 0.11.1) (Wobbrock et al., 2011), to
satisfy distributional assumptions. In the linear mixed effects model, we predicted thresholds with
the factors *time* (2 levels, pre/post), *randomization* (2 levels, phase/contrast), *location* (3 levels,
training/C-transfer/F-transfer) and *training location* (2 levels, upper/lower visual field) as fixed
effects, and participants as random intercept, as in Equation 1:

$$\textbf{Threshold} \sim \textbf{Time} * \textbf{Randomization} * \textbf{Location} * \textbf{Training Location} + (1|\textbf{Subject})$$ **(1)**

To obtain p-values, we performed a repeated measures analyses of variance (rmANOVA) of the model terms and their interactions using Wald's F-test and degrees of freedom determined following Kenward and Roger (Kenward & Roger, 1997). Posthoc tests were performed on estimated marginal means of the aligned and rank transformed data (Elkin et al., 2021) using the emmeans package (version 1.8.8) (Lenth, 2023).

For quantifying learning, we also calculated the Learning Index (LI) (Xiao et al., 2008), as in Equation 2:

$$\textbf{Learning Index(LI)} = \frac{\textbf{Threshold}_{\text{Post}} - \textbf{Threshold}_{\text{Pre}}}{\textbf{Threshold}_{\text{Pre}}}$$ **(2)**

LIs were used to test whether transfer to the C-Transfer location correlated with transfer to the F-Transfer location across subjects; a one-sided Fisher's z-test was used to compare the correlation coefficients between groups via the cocor package (version 1.1.4) (Diedenhofen & Musch, 2015).

To determine effect sizes, we calculated partial η^2 for rmANOVAs, Cohen's d for t-tests, and Cohen's q for comparing correlation coefficients using the effectsize package (version 0.8.6) (Ben-Shachar et al., 2020). Plots were prepared using the ggpubr (version 0.6.0) (Kassambara, 2023) and ggplot2 (version 3.4.4) (Wickham, 2016) packages.

Eye movements

Trials were aborted if subjects broke fixation during fixation phase of trial. To analyze residual miniature eye movements, we focused on microsaccades (MS), which have previously been implicated in VPL (Hung & Carrasco, 2023). To this end, we first identified blinks in the data using the Eyelink parsing algorithm and excluded these timepoints as well as 100 ms before and 150 ms after the blink from analyses. Raw gaze position was converted to dva. Data were segmented into trials (200 ms before the reference stimulus presentation until 350ms after the offset of the target stimulus). We then detected MS using a velocity-based algorithm (Engbert & Kliegl, 2003). This algorithm calculates 2D velocities per trial and identifies MS candidates when the velocity surpasses 6 SD above the average velocity of the trial for a minimum of 6 ms. From these, MS candidates with

less than 1° amplitude and a minimum 10 ms gap between MS were selected to prevent false

positives (Kapoula & Robinson, 1986).

To calculate the MS rate per s, we calculated the average number of MS per time point (1 ms) across

all trials in each session and multiplied these values by the sampling rate (1,000 Hz). To calculate MS

percentage following Hung & Carrasco (2023), we divided the number of MS made in a session by

the number of valid trials (trials which do not contain a blink) for each session and each participant.

This was done for the time window of the reference stimulus, the target stimulus, as well as the

whole trial.

Results

We first examined whether there were any pre-existing differences between the contrast and phase

randomization groups in terms of their pretraining orientation discrimination thresholds. This was

not the case: at none of the three locations, there were no statistically significant differences in pre-

training thresholds between the contrast and phase randomized groups (Welch's two-sample t-test,

Training location $t(83.484) = 0.790$, $p = 0.4313$, Cohen's d = 0.17; C-Transfer $t(83.558) = 0.049$, $p =$

0.9505, Cohen's d = 0.01; F-Transfer $t(82.089) = -1.512$, $p = 0.1344$, Cohen's d = 0.33). Hence, any

differences in learning and/or transfer between the groups cannot be due to pre-existing differences

in orientation discrimination.

Upon training, both randomization groups improved in discriminating orientations at the training

location: the group trained with phase randomization improved thresholds on average by 1.53 deg,

and the group trained with contrast randomization improved on average by 1.40 deg. Learning in all

groups was well characterized by a power law function (see Supplementary Figure S1), characteristic

of gradual VPL (B. A. Dosher & Z.-L. Lu, 2005; B. A. Dosher & Z. L. Lu, 2005).

*Figure 2. Pre and post learning thresholds at the training and transfer locations. **A)** Pre and post learning thresholds at all locations for phase (orange) and contrast (green) randomization. Each dot represents a subject's mean threshold at the respective location and time point. Bars show the mean thresholds over subjects in each group and location. At the training location, there was no statistically significant difference between the phase randomized and the contrast randomized group (t(469) = 1.062, p = 0.2888, Cohen's d = 0.10). Transfer of learning to the C-Transfer location also did not reveal a statistically significant difference between the two groups (t(469) = 0.936, p = 0.3495, Cohen's d = 0.09). However, there was significantly more transfer to the F-Transfer location in the phase randomized group than the contrast randomized group (t(469) = 2.910, p = 0.0038, Cohen's d = 0.27). **B)** Change in the thresholds with learning. Change in the threshold calculated per subject, by subtracting mean post-learning threshold from mean pre-learning threshold. In both panels, error bars represent the standard error of the mean.*

We then assessed whether learning and transfer differed between the groups considering all three
locations (Figure 2). Comparing learning (i.e., the pre-post difference in thresholds) at the training
location, we found no statistically significant difference between the phase randomized and the
contrast randomized group (t(469) = 1.062, p = 0.2888, Cohen's d = 0.10). This suggests that despite
different kinds of variability, training was similarly effective at this location. Transfer of learning to
the C-Transfer location (again in terms of the pre-post difference in threshold) also did not reveal a
statistically significant difference between the two groups (t(469) = 0.936, p = 0.3495, Cohen's d =
0.09). This suggests that training transferred similarly to a close by location with both randomization
manipulations. However, there was significantly more transfer to the F-Transfer location in the
phase randomized group than the contrast randomized group (t(469) = 2.910, p = 0.0038, Cohen's d
= 0.27): while thresholds improved on average only by 1.31 deg in the contrast randomized group,
the average improvement was 1.99 deg in the phase randomized group. This suggests that training
effects generalize more broadly in space with phase than with contrast variability during training.
This differential effect was also supported by a significant time × location × randomization

interaction (F(2,469.031) = 3.317, p = 0.037, η^2_p = 0.01, Supplementary Table 1 and Supplementary

Figure S2). Overall, these results support the hypothesis that training with phase variability engages

neurons with larger receptive fields than training with contrast variability.

Another prediction that can be derived from the hypothesis that phase randomization and contrast

randomization rely on differently sized receptive fields is that learning effects should correlate

between spatial locations if both locations fall into the same receptive fields, whereas they should

not correlate if neurons with small, non-overlapping receptive fields are involved (Kondat et al.,

2023). We thus investigated the correlation of LIs at the two transfer locations for both

randomization groups (Figure 3). We find that LIs at the C-Transfer and F-Transfer locations correlate

highly in the phase randomization group (Pearson's correlation, r = 0.71, t(11) = 3.39, p = 0.006). In

contrast, there was no statistically significant correlation of LIs between the two transfer locations in

the contrast randomization group (Pearson's correlation, r = 0.09, t(14) = 0.373, p=0.71).

Furthermore, the correlation in LIs was significantly larger in the phase than in the contrast

randomized group (z = 1.8964, p = 0.0290, Cohen's q = 0.79). This provides further evidence that the

two groups rely on different neural populations for learning orientation discrimination.

Figure 3. Correlation of LIs between the two transfer locations. In the contrast randomized group, LIs at the C-Transfer and F-Transfer locations are uncorrelated (r = 0.09, p = 0.71). In contrast, LIs are correlated in the phase randomized group (r = 0.71, p = 0.006). The correlation in LIs was significantly larger in the phase than in the contrast randomized group (z = 1.8964, p = 0.0290, Cohen's q = 0.79). Shaded areas represent the 95% confidence intervals. Note that different scaling is used in each panel to ease comparison.

Microsaccade results

MS velocities and amplitudes were highly correlated (Pearson's correlation, r=0.69, t(43618)=199.76,

p<2.2×10⁻¹⁶), i.e., they followed the so-called "main-sequence". After having established that we had

correctly detected MS, we examined how MS varied with our experimental factors. We first tested

whether MS percentage during target stimulus presentation changed with session, randomization
and/or training location. MS percentage during the target stimulus presentation did not change with
session (F(5,123.150)=0.63,p = 0.67), randomization (F(1,24.954)=0.45, p=0.50), or training location
(F(1,24.954)=1.12, p=0.29), and there were no statistically significant interactions (all p>0.05).
Similarly, MS percentage during reference stimulus presentation did not vary depending on session,
randomization, or training location (session F(5,123.047)=0.41, p=0.83; F(1,24.985)=0.11, p=0.73;
training location F(1,24.985)=0.24, p=0.62; all interaction effects p>0.05). When considering the
entire trial, MS percentages changed slightly with session (F(5,123.051)=3.07, p=0.01), but we did
not find any statistically significant effect of randomization (F(1,24984)=2.06, p=0.16) or training
location (F(1,24.984)=1.88, p=0.18), or interaction (all p>0.05). Hence, the different stimulation
parameters did not lead to statistically appreciable differences in the occurrence of MS.

We then assessed whether MS were related to the differential generalization effects we found in our
experimental groups. To this end, we correlated LIs at the two transfer locations, respectively, with
MS percentage. These analyses did not reveal any evidence for fixation variability leading to more
generalizable learning effects (Figure 4, all p>0.08).

Figure 4. MS Correlations between microsaccade percentage with LIs. The upper row shows correlations at C-Transfer and the lower row at F-Transfer during different periods: during the target stimulus, reference stimulus and the entire trial (which includes ISI and both stimuli). There were no significant correlations in both randomization groups in any of the trial phases tested (all p>0.08). Shaded areas represent 95% confidence intervals.

Discussion

We find that variability in spatial phase during training leads to more generalization of VPL in space than variability in contrast. Moreover, participants trained with variable spatial phase transfer learning similarly across two transfer locations, while subjects trained with variable contrast do not. Taken together, these two findings support the hypothesis that the two forms of variability tap into neurons with differently large receptive fields, and in turn, that the type of variability introduced during training can determine the stage of processing at which VPL takes place.

It has been suggested on the basis of behavioral data and simulations that when our visual system is confronted with variability in a task-irrelevant dimension, it can rely on neurons that are invariant to this feature and thus subsume this variability (Manenti et al., 2023); in the brain, such neurons typically reside at a higher stage of processing and have large receptive fields. Alternatively, the

visual system can make use of neurons at earlier stages of processing that are conjunctively tuned to

the task-relevant and the task-irrelevant features of the stimuli. Because these neurons typically

have spatially restricted receptive fields, they cannot, by themselves, enable generalization across

the visual field. Importantly, different invariances arise at different levels of processing. Hence,

creating variability in different task-irrelevant features of a stimulus should result in different

patterns of generalization. This is what we find for the case of spatial phase and contrast variability.

Relevant to our experiments, it is known that contrast invariance arises earlier in the visual

processing hierarchy than phase invariance: contrast invariance gradually increase along the visual

ventral stream (Avidan et al., 2002), but is already seen at the level of simple cells in V1 (Hubel &

Wiesel, 1962). The earliest stage at which phase invariance arises is on the level of complex cells in

V1 (Hubel & Wiesel, 1962). Both simple and complex cells are orientation tuned. According to a

classical model of orientation tuning in V1, simple cells feed into complex cells (Hubel & Wiesel,

1962), and there is empirical evidence that supports this view (e.g., Alonso & Martinez, 1998). This

principle also underlies hierarchical models of visual processing such as HMAX (Serre & Riesenhuber,

2004). Furthermore, simple cells are predominantly found in the granular input layer of V1, whereas

complex cells are found in supra- and infragranular output layers (Hubel & Wiesel, 1968). Hence,

complex cells reside on a higher level of processing than simple cells already in V1. Full phase

invariance only arises in complex cells in V2 (Cloherty & Ibbotson, 2015), and thus later than contrast

invariance.

Importantly, V1 complex cells have larger receptive fields than simple cells (Schiller et al., 1976),

especially at higher eccentricities (Wilson & Sherman, 1976), and fully phase-invariant complex cells

in V2 have larger receptive fields than contrast-invariant simple cells in V1 (Gattass et al., 1981).

Hence, the earliest stages of processing at which differential spatial transfer effects between

contrast- and phase-randomized training could hypothetically occur would be between simple and

complex cells in V1, and/or between V1 and V2. Of note, simple and complex cells are not only

found in V1, but also at higher stages of processing such as V2 (Levitt et al., 1994) and V4 (Desimone

& Schein, 1987). Hence, the principle that phase invariance arises at a higher stage of processing

than contrast invariance holds irrespective of where exactly plasticity occurs.

We note that the spatial generalization that we observe is substantial relative to the typical

receptive field size in early visual areas, including V4. Yet, neurophysiological recordings in macaque

monkeys have shown that orientation tuning can be found not only in early visual areas, where it is

typically studied, but that high-level visual cortical areas in the temporal lobe with large receptive

fields (Desimone & Gross, 1979) contain orientation-tuned neurons (Gross et al., 1972; Vogels &

Orban, 1994) that are causally relevant for orientation discrimination (Adab & Vogels, 2016). These
could possibly support the large distances between the different locations in our experiments with
their larger receptive fields. Unfortunately, at least to our knowledge, the difference in receptive
field size between simple and complex cells has not been mapped for higher visual areas and the
distinction between simple and complex cells has not been made for orientation tuned neurons that
have been found beyond area V4 in the macaque monkey. Neurophysiological and/or neuroimaging
experiments would be necessary to achieve this. Nevertheless, the distinction between simple and
complex cells as different stages of processing seems to hold irrespective of these unknowns, as
does the emergence of full phase invariance at a higher stage of processing than contrast invariance.
Our results are thus in line with the hypothesis that the dimension in which task-irrelevant variability
occurs determines the stage of processing affected by VPL, reflected by varying degrees of
generalization.

Previous VPL studies have used phase or contrast randomization not as an experimental variable,
but to assure that subjects are not using spatial cues to solve orientation discrimination tasks (in the
case of spatial phase) and/or to reduce adaptation. Some of these studies have reported spatial
generalization effects (e.g., Zhang et al., 2010). In contrast, a study by Sowden and colleagues (2002)
reported training effects that were extremely specific to the spatial frequency, retinal location and
even eye of origin, consistent with learning effects in the granular input layer of V1 where neurons
which such tuning properties reside. Importantly, this study used a training procedure with no
variability in which only a single grating was presented for several thousand trials. Notably, a study
investigating the neural basis of orientation discrimination learning in macaque monkey V1 used
phase randomization and reported learning effects only outside the granular input layer (Schoups et
al., 2001), which would be compatible with an outsized involvement of complex cells. Overall, this
suggests that by using variability during training, previous studies may have coincidentally engaged
neurons at higher stages of processing and with larger receptive fields.

Apart from variability, spatial generalization of VPL also occurs when adaptation is reduced or
eliminated (Harris et al., 2012). While spatial transfer to the C-Transfer location we found in both
randomization groups may be explained by a reduction of adaptation with variability, the differential
transfer effects at the F-Transfer location and the differential correlation between learning effects at
the two transfer locations in the different groups are not easily accounted for by adaptation.
Previous studies have also shown that task difficulty affects generalization (Ahissar & Hochstein,
2004). However, as pretraining thresholds did not differ significantly between contrast and phase
randomization, this factor also cannot easily explain our differential transfer results. Hence,
variability may contribute to generalization in VPL beyond the well-established factors task difficulty

and adaptation. On the other hand, studies using contrast randomization during orientation
discrimination learning in the context of double training (Xiao et al., 2008) and manipulation of
attention (Donovan & Carrasco, 2018) have reported sizable spatial transfer effects. These may rely
on more intricate neural mechanisms such as top-down influences beyond the effects discussed
here. Hence, variability is not the only factor that enables spatial transfer after perceptual learning,
but there are alternative paths and mechanisms of achieving generalization.

Small, involuntary eye movements, e.g., in the form of MS, could in principle constitute an additional
form of variability beyond the variability in stimulus features that we induced in our experiments:
even a small displacement in fixation can introduce phase variability in the receptive field of a given
neuron. Interestingly, it has previously been proposed that such miniature eye movements increase
stimulus discriminability (Hennig et al., 2002; Rucci et al., 2007). Furthermore, MS have been found
to change in frequency over the time course of perceptual learning (Hung & Carrasco, 2023).
However, we did not find any differences between the training regimes that would support a specific
role of MS in the differential spatial generalization patterns that we observed. We note that
recordings in macaque V1 have found no differential effect of eye movements on simple versus
complex cells (Kagan et al., 2002). It has previously been proposed that lateral connections in V1
implement a form of spatial invariance that counteracts the possibly detrimental effects of positional
uncertainty brought about by miniature eye movements during perceptual learning (Otto et al.,
2006; Zhaoping et al., 2003). Alternatively or additionally, physiological (Ringach, 2002; Victor &
Purpura, 1998) and behavioral (Field & Nachmias, 1984; Huang et al., 2006) data indicate that phase
tuning is relatively broad. This would suggest that substantial phase variability (such as the phase
variability in our experimental condition) is needed to tap into phase-invariant stages of processing.
Ultimately, future experiments using gaze-contingent displays will be needed to differentiate the
effect of stimulus- versus eye movement-induced variability on spatial generalization.

While our study exemplifies the differential effects of different kinds of task-irrelevant variability, it
is important to emphasize that our study only investigated a narrow range of stimulus parameters in
each group. E.g., as mentioned above, a narrower range of phase variability might result in higher
specificity. Predicting outcomes of VPL with a much wider range of variability or other features
becomes more challenging given multi-level nature of VPL (Maniglia & Seitz, 2018). E.g., extending
the range of contrast variability might cause interactions with other factors such as task difficulty
(Talluri et al., 2015), or perceived variability (Zaman et al., 2021) that could affect behavioral
outcomes. Hence, further systematic studies are needed to map out the stimulus and variability
parameters that promote generalization.

Taken together, our results emphasize a role of targeted variability in achieving spatial

generalization and suggest higher stages of visual processing as a possible locus of VPL. Variability

may thus be a useful component for improving real life applications of VPL without burdening the

learner with additional training time and effort.

References

Adab, H. Z., & Vogels, R. (2016). Perturbation of posterior inferior temporal cortical activity impairs
coarse orientation discrimination. *Cerebral cortex*, *26*(9), 3814-3827.
https://doi.org/10.1093/cercor/bhv178

Ahissar, M., & Hochstein, S. (2004). The reverse hierarchy theory of visual perceptual learning.
Trends in cognitive sciences, *8*(10), 457-464. https://doi.org/10.1016/j.tics.2004.08.011

Alonso, J. M., & Martinez, L. M. (1998). Functional connectivity between simple cells and complex
cells in cat striate cortex. *Nature neuroscience*, *1*(5), 395-403. https://doi.org/10.1038/1609

Avidan, G., Harel, M., Hendler, T., Ben-Bashat, D., Zohary, E., & Malach, R. (2002). Contrast sensitivity
in human visual areas and its relationship to object recognition. *Journal of neurophysiology*,
87(6), 3102-3116. https://doi.org/10.1152/jn.2002.87.6.3102

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

Ben-Shachar, M., Lüdecke, D., & Makowski, D. (2020). effectsize: Estimation of Effect Size Indices and
Standardized Parameters. *Journal of Open Source Software*, *5*(56), 2815.
https://doi.org/10.21105/joss.02815

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial vision*, *10*(4), 433-436.
https://www.ncbi.nlm.nih.gov/pubmed/9176952

Cloherty, S. L., & Ibbotson, M. R. (2015). Contrast-dependent phase sensitivity in V1 but not V2 of
macaque visual cortex. *Journal of neurophysiology*, *113*(2), 434-444.
https://doi.org/10.1152/jn.00539.2014

Desimone, R., & Gross, C. G. (1979). Visual areas in the temporal cortex of the macaque. *Brain*
Research, *178*(2-3), 363-380. https://doi.org/10.1016/0006-8993(79)90699-1

Desimone, R., & Schein, S. J. (1987). Visual properties of neurons in area V4 of the macaque:
sensitivity to stimulus form. *Journal of neurophysiology*, *57*(3), 835-868.
https://doi.org/10.1152/jn.1987.57.3.835

Diedenhofen, B., & Musch, J. (2015). cocor: A Comprehensive Solution for the Statistical Comparison
of Correlations. *PLoS One*, *10*(4), e0121945. https://doi.org/10.1371/journal.pone.0121945

Donovan, I., & Carrasco, M. (2018). Endogenous spatial attention during perceptual learning
facilitates location transfer. *Journal of vision*, *18*(11), 7. https://doi.org/10.1167/18.11.7

Dosher, B. A., & Lu, Z.-L. (2005). Perceptual learning in clear displays optimizes perceptual expertise:
Learning the limiting process. *Proceedings of the National Academy of Sciences*, *102*(14),
5286–5290.

Dosher, B. A., & Lu, Z. L. (2005). Perceptual learning in clear displays optimizes perceptual expertise:
learning the limiting process. *Proceedings of the national academy of sciences of the united*
states of america, 102(14), 5286-5290. https://doi.org/10.1073/pnas.0500492102

Dosher, B. A., & Lu, Z. L. (2020). *Perceptual learning. How experience shapes visual perception.* MIT
Press.

Douvis, S. J. (2005). Variable Practice in Learning the Forehand Drive in Tennis. *Perceptual and motor*
skills, 101(2), 531-545. https://doi.org/10.2466/pms.101.2.531-545

Elkin, L. A., Kay, M., Higgins, J. J., & Wobbrock, J. O. (2021). An Aligned Rank Transform Procedure for
Multifactor Contrast Tests. UIST '21: The 34th Annual ACM Symposium on User Interface
Software and Technology, Virtual Event.

Engbert, R., & Kliegl, R. (2003). Microsaccades uncover the orientation of covert attention. *Vision*
research, 43(9), 1035-1045. https://doi.org/10.1016/s0042-6989(03)00084-1

Fahle, M. (2005). Perceptual learning: specificity versus generalization. *Current opinion in*
neurobiology, 15(2), 154-160. https://doi.org/10.1016/j.conb.2005.03.010

Field, D. J., & Nachmias, J. (1984). Phase reversal discrimination. *Vision research, 24*(4), 333–340.
https://doi.org/10.1016/0042-6989(84)90058-0

Fiorentini, A., & Berardi, N. (1980). Perceptual learning specific for orientation and spatial frequency.
Nature, 287(5777), 43-44. https://doi.org/10.1038/287043a0

Gattass, R., Gross, C. G., & Sandell, J. H. (1981). Visual topography of V2 in the macaque. *Journal of*
Comparative Neurology, 201(4), 519-539. https://doi.org/10.1002/cne.902010405

Gladilin, E., & Eils, R. (2015). On the role of spatial phase and phase correlation in vision, illusion, and
cognition. *Frontiers in computational neuroscience, 9,* 45.
https://doi.org/10.3389/fncom.2015.00045

Gross, C. G., Rocha-Miranda, C. E., & Bender, D. B. (1972). Visual properties of neurons in
inferotemporal cortex of the Macaque. *Journal of neurophysiology, 35*(1), 96-111.
https://doi.org/10.1152/jn.1972.35.1.96

Harris, H., Gliksberg, M., & Sagi, D. (2012). Generalized perceptual learning in the absence of sensory
adaptation. *Current biology, 22*(19), 1813-1817. https://doi.org/10.1016/j.cub.2012.07.059

Hennig, M. H., Kerscher, N. J., Funke, K., & Wörgötter, F. (2002). Stochastic resonance in visual
cortical neurons: does the eye-tremor actually improve visual acuity? *Neurocomputing, 44-*
46, 115-120. https://doi.org/10.1016/s0925-2312(02)00371-5

Higgins, J. J., & Tashtoush, S. (1994). An aligned rank transform test for interaction. *Nonlinear World,*
1, 201-211.

Huang, P.-C., Kingdom, F., & Hess, R. (2006). Only two phase mechanisms,+-cosine, in human vision.

Vision research, *46*(13), 2069–2081. https://doi.org/10.1016/j.visres.2005.12.020

Hubel, D. H., & Wiesel, T. N. (1962). Receptive fields, binocular interaction and functional

architecture in the cat's visual cortex. *Journal of physiology*, *160*(1), 106-154.

https://doi.org/10.1113/jphysiol.1962.sp006837

Hubel, D. H., & Wiesel, T. N. (1968). Receptive fields and functional architecture of monkey striate

cortex. *Journal of physiology*, *195*(1), 215-243.

https://doi.org/10.1113/jphysiol.1968.sp008455

Hung, S. C., & Carrasco, M. (2023). Microsaccades as a long-term oculomotor correlate in visual

perceptual learning. *Psychonomic Bulletin & Review*, *30*(1), 235-249.

https://doi.org/10.3758/s13423-022-02151-8

Kagan, I., Gur, M., & Snodderly, D. M. (2002). Spatial Organization of Receptive Fields of V1 Neurons

of Alert Monkeys: Comparison With Responses to Gratings. *Journal of neurophysiology*,

88(5), 2557-2574. https://doi.org/10.1152/jn.00858.2001

Kapoula, Z., & Robinson, D. A. (1986). Saccadic undershoot is not inevitable: Saccades can be

accurate. *Vision research*, *26*(5), 735-743. https://doi.org/10.1016/0042-6989(86)90087-8

Karni, A., & Bertini, G. (1997). Learning perceptual skills: behavioral probes into adult cortical

plasticity. *Current opinion in neurobiology*, *7*(4), 530-535. https://doi.org/10.1016/s0959-

4388(97)80033-5

Kassambara, A. (2023). *ggpubr: 'ggplot2' Based Publication Ready Plots*. https://CRAN.R-

project.org/package=ggpubr

Kenward, M. G., & Roger, J. H. (1997). Small Sample Inference for Fixed Effects from Restricted

Maximum Likelihood. *Biometrics*, *53*(3). https://doi.org/10.2307/2533558

Kondat, T., Aderka, M., & Censor, N. (2023). Modulating temporal dynamics of performance across

retinotopic locations enhances the generalization of perceptual learning. *iScience*, *26*(11),

108276. https://doi.org/10.1016/j.isci.2023.108276

Lenth, R. V. (2023). *emmeans: Estimated Marginal Means, aka Least-Squares Means*.

https://CRAN.R-project.org/package=emmeans

Levitt, J. B., Kiper, D. C., & Movshon, J. A. (1994). Receptive fields and functional architecture of

macaque V2. *Journal of neurophysiology*, *71*(6), 2517-2542.

https://doi.org/10.1152/jn.1994.71.6.2517

Lu, Z.-L., & Dosher, B. A. (2004). Perceptual learning retunes the perceptual template in foveal

orientation identification. *Journal of vision*, *4*(1), 5–5. https://doi.org/10.1167/4.1.5

Lu, Z.-L., & Dosher, B. A. (2022). Current directions in visual perceptual learning. *Nature Reviews*
Psychology, *1*(11), 654-668. https://doi.org/10.1038/s44159-022-00107-2

Manenti, G. L., Dizaji, A. S., & Schwiedrzik, C. M. (2023). Variability in training unlocks generalization
in visual perceptual learning through invariant representations. *Current biology*, *33*(5), 817-
826 e813. https://doi.org/10.1016/j.cub.2023.01.011

Maniglia, M., & Seitz, A. R. (2018). Towards a whole brain model of Perceptual Learning. *Current*
Opinion in Behavioral Sciences, *20*, 47-55. https://doi.org/10.1016/j.cobeha.2017.10.004

Otto, T. U., Herzog, M. H., Fahle, M., & Zhaoping, L. (2006). Perceptual learning with spatial
uncertainties. *Vision research*, *46*(19), 3223-3233.
https://doi.org/10.1016/j.visres.2006.03.021

Perry, L. K., Samuelson, L. K., Malloy, L. M., & Schiffer, R. N. (2010). Learn Locally, Think Globally:
Exemplar Variability Supports Higher-Order Generalization and Word Learning. *Psychological*
science, *21*(12), 1894-1902. https://doi.org/10.1177/0956797610389189

Raviv, L., Lupyan, G., & Green, S. C. (2022). How variability shapes learning and generalization.
Trends in cognitive sciences, *26*(6), 462-483. https://doi.org/10.1016/j.tics.2022.03.007

Ringach, D. L. (2002). Spatial structure and symmetry of simple-cell receptive fields in macaque
primary visual cortex. *Journal of neurophysiology*, *88*(1), 455–463.
https://doi.org/10.1152/jn.2002.88.1.455

Rucci, M., Iovin, R., Poletti, M., & Santini, F. (2007). Miniature eye movements enhance fine spatial
detail. *Nature*, *447*(7146), 851-854. https://doi.org/10.1038/nature05866

Schiller, P. H., Finlay, B. L., & Volman, S. F. (1976). Quantitative studies of single-cell properties in
monkey striate cortex. I. Spatiotemporal organization of receptive fields. *Journal of*
neurophysiology, *39*(6), 1288-1319. https://doi.org/10.1152/jn.1976.39.6.1288

Schoups, A. A., Vogels, R., & Orban, G. A. (1995). Human perceptual learning in identifying the
oblique orientation: retinotopy, orientation specificity and monocularity. *Journal of*
physiology, *483 (Pt 3)*, 797-810. https://doi.org/10.1113/jphysiol.1995.sp020623

Schoups, A. A., Vogels, R., Qian, N., & Orban, G. A. (2001). Practising orientation identification
improves orientation coding in V1 neurons. *Nature*, *412*(6846), 549-553.
https://doi.org/10.1038/35087601

Serre, T., & Riesenhuber, M. (2004). Realistic Modeling of Simple and Complex Cell Tuning in the
HMAX Model, and Implications for Invariant Object Recognition in Cortex. *CSAIL Technical*
Reports, *052*. http://hdl.handle.net/1721.1/30491

Shapley, R. (1985). The importance of contrast in the perception of brightness and form.
Transactions of the American Philosophical Society, *75*(6), 20-29.

Shiu, L. P., & Pashler, H. (1992). Improvement in line orientation discrimination is retinally local but
dependent   on   cognitive   set.   *Perception   &   psychophysics*,   *52*(5),   582-588.
https://doi.org/10.3758/bf03206720

Sowden, P. T., Rose, D., & Davies, I. R. (2002). Perceptual learning of luminance contrast detection:
specific for spatial frequency and retinal location but not orientation. *Vision research*,
42(10), 1249-1258. https://doi.org/10.1016/s0042-6989(02)00019-6

Talluri, B. C., Hung, S.-C., Seitz, A. R., & Seriès, P. (2015). Confidence-based integrated reweighting
model of task-difficulty explains location-based specificity in perceptual learning. *Journal of*
vision, *15*(10), 17. https://doi.org/10.1167/15.10.17

Tan, Q., Wang, Z., Sasaki, Y., & Watanabe, T. (2019). Category-induced transfer of visual perceptual
learning. *Current biology*, *29*(8), 1374–1378. https://doi.org/10.1016/j.cub.2019.03.003

Victor, J. D., & Purpura, K. P. (1998). Spatial phase and the temporal structure of the response to
gratings   in   V1.   *Journal   of   neurophysiology*,   *80*(2),   554–571.
https://doi.org/10.1152/jn.1998.80.2.554

Vogels, R., & Orban, G. A. (1994). Activity of inferior temporal neurons during orientation
discrimination with successively presented gratings. *Journal of neurophysiology*, *71*(4), 1428-
1451. https://www.ncbi.nlm.nih.gov/pubmed/8035226

Vukatana, E., Graham, S. A., Curtin, S., & Zepeda, M. S. (2015). One is Not Enough: Multiple
Exemplars Facilitate Infants' Generalizations of Novel Properties. *Infancy*, *20*(5), 548-575.
https://doi.org/10.1111/infa.12092

Wickham,   H.   (2016).   *ggplot2:   Elegant   Graphics   for   Data   Analysis*.   Springer.
https://doi.org/10.1007/978-3-319-24277-4

Wilson, J. R., & Sherman, S. M. (1976). Receptive-field characteristics of neurons in cat striate cortex:
Changes with visual field eccentricity. *Journal of neurophysiology*, *39*(3), 512-533.
https://doi.org/10.1152/jn.1976.39.3.512

Wobbrock, J. O., Findlater, L., Gergle, D., & Higgins, J. J. (2011). The Aligned Rank Transform for
Nonparametric Factorial Analyses Using Only Anova Procedures. *CHI '11: Proceedings of the*
SIGCHI   Conference   on   Human   Factors   in   Computing   Systems,   143–146.
https://doi.org/10.1145/1978942.1978963

Xiao, L. Q., Zhang, J. Y., Wang, R., Klein, S. A., Levi, D. M., & Yu, C. (2008). Complete transfer of
perceptual learning across retinal locations enabled by double training. *Current biology*,
18(24), 1922-1926. https://doi.org/10.1016/j.cub.2008.10.030

Zaman, J., Chalkia, A., Zenses, A.-K., Bilgin, A. S., Beckers, T., Vervliet, B., & Boddez, Y. (2021).

Perceptual variability: Implications for learning and generalization. *Psychonomic Bulletin &*

Review, *28*(1), 1-19. https://doi.org/10.3758/s13423-020-01780-1

Zhang, T., Xiao, L. Q., Klein, S. A., Levi, D. M., & Yu, C. (2010). Decoupling location specificity from

perceptual learning of orientation discrimination. *Vision research*, *50*(4), 368-374.

https://doi.org/10.1016/j.visres.2009.08.024

Zhaoping, L., Herzog, M. H., & Dayan, P. (2003). Nonlinear ideal observation and recurrent

preprocessing in perceptual learning. *Network*, *14*(2), 233-247.

https://www.ncbi.nlm.nih.gov/pubmed/12790183