The *homie* insulator has sub-elements with different insulating and long-range pairing properties

Miki Fujioka[1], Wenfan Ke[2], Paul Schedl[2,*], and James B Jaynes[1,*]

1: Department of Biochemistry and Molecular Biology, Thomas Jefferson University,

   Philadelphia, PA 19107

2: Department of Molecular Biology, Princeton University, Princeton NJ 08544

*corresponding authors: james.jaynes@jefferson.edu, pschedl@princeton.edu

running title: *homie* elements have distinct, overlapping function

19 **GRAPHICAL ABSTRACT**

ABSTRACT

Chromatin insulators are major determinants of chromosome architecture. Specific architectures induced by insulators profoundly influence nuclear processes, including how enhancers and promoters interact over long distances and between homologous chromosomes. Insulators can pair with copies of themselves in *trans* to facilitate homolog pairing. They can also pair with other insulators, sometimes with great specificity, inducing long-range chromosomal loops. Contrary to their canonical function of enhancer blocking, these loops can bring distant enhancers and promoters together to activate gene expression, while at the same time blocking other interactions in *cis*. The details of these effects depend on the choice of pairing partner, and on the orientation specificity of pairing, implicating the 3-dimensional architecture as a major functional determinant. Here we dissect the *homie* insulator from the Drosophila *even skipped* (*eve*) locus, to understand its substructure. We test pairing function in *cis* based on *homie*-carrying transgenes interacting with endogenous *eve*. The assay is sensitive to both pairing strength and orientation. Using this assay, we found that a Su(Hw) binding site in *homie* is required for efficient long-range interaction, although some activity remains without it. This binding site also contributes to the canonical insulator activities of enhancer blocking and barrier function. Based on this and other results from our functional dissection, each of the canonical insulator activities, chromosomal loop formation, enhancer blocking, and barrier activity, are partially separable. Our results show the complexity inherent in insulator functions, which can be provided by an array of different proteins with both shared and distinct properties.

Introduction

Genomic elements called boundaries or insulators function to separate genes from each other, or, in some cases, to separate distinct regulatory domains within a gene (e.g., in the Hox gene complexes). However, they have properties that have proven to be difficult to reconcile mechanistically. For example, while these elements can block enhancer-promoter interactions when inserted between regulatory elements, they can also mediate long-distance regulatory interactions (Chetverina *et al.* 2017; Schwartz and Cavalli 2017; Özdemir and Gambetta 2019).

One of the first insulator elements discovered is located within the *Drosophila gypsy* retrotransposon (Modolell *et al.* 1983; Peifer and Bender 1988). The *gypsy* transposon was found to be responsible for many spontaneous mutations in flies. In many cases, the mutant phenotypes arose because the *gypsy* transposon inserted between a gene and its regulatory elements, blocking regulatory interactions. The insulator properties of *gypsy* have been attributed to the binding of a set of chromatin proteins, the sequence-specific component of which is the protein Suppressor of Hairy Wing (Su(Hw) (Parkhurst *et al.* 1988; Holdridge and Dorsett 1991; Geyer and Corces 1992). Several insulators found in the bithorax complex (BxC) of Drosophila have also been extensively studied, including *Mcp*, *Fab7*, and *Fab8* (Karch *et al.* 1994; Zhou *et al.* 1999; Barges *et al.* 2000). Some but not all of these bind Su(Hw), while an array of other proteins, including Cp190, are also known to contribute to the activities of these and other insulators (Zhao *et al.* 1995; Gaszner *et al.* 1999; Ghosh *et al.* 2001; Pai *et al.* 2004; Moon *et al.* 2005; Rath *et al.* 2006; Roy *et al.* 2007; Maksimenko *et al.* 2008; Jiang *et al.* 2009; Aoki *et al.* 2012; Li and Gilmour 2013; Cuartero *et al.* 2014; Maksimenko *et al.* 2015; Zolotarev *et al.* 2016; Fedotova *et al.* 2018; Ramírez *et al.* 2018; Bag *et al.* 2021).

The Drosophila *even skipped* (*eve*) gene is flanked by two insulators, *nhomie* (*neighbor of homie*) and *homie* (*homing insulator at eve*), at its 5'- and 3'-ends, respectively. These two elements define the *eve* TAD (topologically associated domain) (Fujioka *et al.* 2009; Fujioka *et al.* 2016; Bing *et al.* 2024) (see map in Figure 1A). The properties of the 3' insulator *homie* have been well-characterized. Like many other insulators, it has enhancer blocking activity and an ability to prevent the spread of Polycomb-dependent silencing (Fujioka *et al.* 2013; Fujioka *et al.* 2021). *homie* abuts the promoter of an essential housekeeping gene, *TER94* (Pinter *et al.* 1998; Leon and McKearin 1999; Ruden *et al.* 2000). Since the *eve* TAD is assembled into a Polycomb group (PcG)-silenced domain in most cells during all but the early stages of development (Schwartz *et al.* 2006; Tolhuis *et al.* 2006), this barrier activity is thought to be a critically important function, as insulator protein binding sites are located at roughly half of the borders of Polycomb domains in Drosophila (De *et al.* 2020). In *eve-TER94* "pseudo-locus" transgenes, *homie* is required to keep the *eve* Polycomb response elements (PREs) from shutting down *TER94* through the spreading of the repressive PcG-dependent histone modification H3K27me3 (trimethylation of histone H3 residue lysine-27) (Fujioka *et al.* 2013).

In addition to these activities, *homie* and *nhomie* share another characteristic property of fly insulators, namely, an ability to physically pair with themselves and with each other (Fujioka *et al.* 2009; Fujioka *et al.* 2016; Chen *et al.* 2018). These pairing interactions have two important consequences. First, pairing between heterologous boundaries in *cis* generates a loop, or TAD. Depending on the orientation of the pairing interaction, this loop can be a circle-loop (where pairing is head-to-head) or a stem-loop (where pairing is head-to-tail). Since *homie* and *nhomie*

pair with each other head-to-tail, the *eve* TAD has a stem-loop topology. As we have shown (KE
et al. 2024), this loop topology mediates the physical isolation of the *eve* TAD from its
neighbors. Second, homologous chromosomes in flies are paired in register throughout much
of the life cycle, and evidence suggests that self-pairing interactions between TAD boundary
elements facilitate this (VIETS *et al.* 2019). While *homie:nhomie* pairing is head-to-tail, *homie*
and *nhomie* self-pairing is head-to-head, like that of other fly boundaries that have been tested
(e.g., (KYRCHANOVA *et al.* 2008). This orientation preference has been shown to favor
transvection (regulatory cross-talk) between paired homologous chromosomes (FUJIOKA *et al.*
2016), a phenomenon that is widespread in *Drosophila*.

As has been observed for the *gypsy* insulator and for boundaries from the Bithorax complex
(SIGRIST AND PIRROTTA 1997; MULLER *et al.* 1999; VAZQUEZ *et al.* 2006; LI *et al.* 2011), both *homie*
and *nhomie* can also mediate long-distance (100 kb to Mbs) regulatory interactions (FUJIOKA *et*
al. 2009; FUJIOKA *et al.* 2016; CHEN *et al.* 2018; BING *et al.* 2024). As an example, when a *homie*-
containing reporter transgene is inserted at an attP site located in the 1st exon of the *hebe*
gene, 142 kb upstream of the *eve* gene, enhancers in the *eve* TAD can drive reporter expression
(FUJIOKA *et al.* 2009; FUJIOKA *et al.* 2016; CHEN *et al.* 2018). Reporter activation depends on the
relative orientation of the *homie* element in the transgene, but not on the orientation of the
transgene in the chromosome. This constraint arises because *homie* in the transgene pairs with
homie in the *eve* locus head-to-head, and it pairs with *eve*-locus *nhomie* head-to-tail (FUJIOKA *et*
al. 2016; BING *et al.* 2024).

Here, we have undertaken a functional dissection of *homie*, focusing on three centrally
important activities of this class of elements: enhancer blocking, PRE blocking, and long-range
pairing. Detailed slicing and dicing of *homie* reveals a general correlation between the three
activities. However, the correlation is not perfect, indicating that while the activities are
related, there are some mechanistic differences. For example, we show that a minimal <300 bp
sequence retains all three activities. Within this minimal insulator, there is a single consensus
recognition sequence for the *gypsy* insulator-binding protein Su(Hw). While mutation of this
sequence results in a near-complete loss of long-range pairing activity, it only partially
compromises enhancer blocking activity, and has a minimal effect on PRE blocking. An
extensive comparison of subelement combinations provides clear cases of divergence between
the requirements for each of the three insulator functions.

Results

Multiple insulator sub-elements contribute to long-range pairing

In previous studies on long-range pairing interactions, we used a "minimal" *homie*, CDEF, which consisted of four contiguous ~100 bp sub-elements, as it appeared to have nearly full activity (Fᴜᴊɪᴏᴋᴀ *et al.* 2016; Cʜᴇɴ *et al.* 2018). To better understand how these *homie* sequences contribute to long-range pairing, we examined the long-distance pairing activity of different combinations of sub-elements from a larger 800 bp fragment, ABCDEFGH, that was previously identified as the "*homie*" insulator (Fᴜᴊɪᴏᴋᴀ *et al.* 2009). As indicated in Figure 1B, this larger fragment extends from just downstream of the *eve* PRE to just beyond the first exon of the *TER94-RD* transcript (Gʀᴀᴍᴀᴛᴇs *et al.* 2022), and it spans a DNase hypersensitive region (Sᴀᴄᴋᴇʀsᴏɴ *et al.* 1999). To test for long-range (LR) pairing interactions, we inserted *homie*-containing reporter gene constructs into an attP site in the 1st exon of the *hebe* gene, 142 kb upstream of *eve* (Fᴜᴊɪᴏᴋᴀ *et al.* 2009; Fᴜᴊɪᴏᴋᴀ *et al.* 2016; Cʜᴇɴ *et al.* 2018). For the first set of experiments, we used the single reporter transgene shown in Figure 1C, while in experiments described below, we used the dual reporter transgene shown in Figure 1D. When *homie* or its derivatives have LR pairing activity (and when the insulator is in the "correct" orientation relative to the reporter in the transgene: see Figure 1C, D) the reporter gene is subject to regulation by enhancers in the *eve* TAD and is expressed in an *eve*-like pattern (Figure 2). By contrast, no reporter expression is seen when the same reporter transgene has an equal-length DNA segment from phage λ ("λ DNA" in Figure 2, Figure S2). Also, as described further below in the section on enhancer blocking activity, a *hebe* enhancer is located upstream of the attP site at −142 kb (Figure S1B). When *homie* is interposed between this enhancer and the transgene reporter, it blocks the enhancer from activating the reporter.

In the experiments shown in Figure 2, we compared the LR pairing activity of different constructs using two criteria: the relative number of the cells expressing the reporter gene in an *eve* pattern and the relative level of expression as judged by the intensity of staining. These two measures generally correlated perfectly with each other. We first extended the minimal CDEF *homie* endpoint by ~200 bp downstream into the beginning of the *TER94* transcription unit to give CDEFGH (see map, Figure 1B). The addition of the GH sub-elements appeared to increase the LR pairing frequency and/or stability (compare CDEF with CDEFGH in Figure 2). This is reflected in a greater number of cells expressing the reporter in early stripes, and in the stronger expression at later stages, most noticeably in the CNS and mesoderm of stage 13 embryos (Figure S1A). To further test the contribution of the GH region to LR pairing activity, we combined GH with EF. As shown below (see Figure 3B), EF on its own has almost no LR pairing activity. In contrast, an *eve*-like pattern of *lacZ* expression is observed for EFGH throughout embryogenesis (Figure 2). However, the intensity of staining is considerably weaker than CDEF, and fewer cells express *lacZ*. This shows that GH can partially substitute for the CD sub-elements. On the other hand, as shown in Figure 2, the AB sub-elements cannot replace EF, as the ABCD combination has little or no LR pairing activity.

Next, we assayed the LR pairing activity of the four possible combinations of three different CDEF sub-elements. Of these sub-element combinations, only DEF has LR pairing activity comparable to that of CDEF (Figure 3A). However, C does facilitate LR pairing by DEF, since CDEF shows slightly stronger expression in more cells than does DEF (Figure 3A). This can be

seen as weaker expression with DEF in the mesoderm and the CNS (Figure S1A). However, the
reduction is small and somewhat variable. F facilitates LR pairing more than does C, since
expression with DEF is considerably stronger at all stages, and seen in more cells, than is
expression with CDE (Figure 3A). Both D and E are required for LR pairing activity, since neither
CDF nor CEF show any *eve*-patterned expression (Figure 3A).

Since DEF has significantly more LR pairing activity than any of the other tripartite
combinations, we tested the relative contributions of its three sub-elements. As shown in
Figure 3B, DF and EF have minimal LR pairing activity, as with each, there are at most only a few
cells in the embryo in which the reporter responds to *eve* enhancers. By contrast, the DE sub-
element combination clearly retains LR pairing activity. However, as evident from the number
of cells in which the *eve* enhancers activate reporter expression (compare DEF and DE) and the
level of expression (particularly at later stages), the F sub-element clearly bolsters the LR pairing
activity of the DE combination.

While EF does not have LR pairing activity on its own (Figure 3B), the EFGH combination
shows weak activity (Figure 2). This fits with our finding that GH also increases the activity of
CDEF. In other experiments, we further tested the AB region by creating ABDE in the context of
the double reporter transgene. The AB region is unable to enhance the activity of DE, since *lacZ*
expression in an *eve* pattern is not increased over that with DE alone (Figure S3).

Insulator protein binding sites are required for long-range pairing

The experiments in the previous section show that D and E harbor most of *homie*'s LR
pairing activity. Analysis of the *homie* sequence identified a Su(Hw) consensus site (RAMOS *et al.*
2006; BAXLEY *et al.* 2017) in D, and a Cp190-associated sequence (CUARTERO *et al.* 2014) in E
(Figure S7). We tested whether these sites are involved in LR pairing. We mutated the Su(Hw)
site in the context of DEF. As shown in Figure 4, this results in a loss of most but not all of the
LR pairing activity. Mutating the Cp190-associated sequence in the context of DEF reduced LR
pairing, although the effect is not as drastic as that of the Su(Hw) site mutation (Figure 4).

To further analyze the effects of the Su(Hw) site mutation, we used smFISH to probe for LR
pairing in stage 15 DEF and DEFΔSH dual reporter transgene embryos. In the experiment of
Figure 5, DEF and DEFΔSH dual reporters were inserted into the attP site at −142 kb in two
different orientations. In the Z5 orientation, the *LacZ* reporter is closer to the *eve* TAD, while in
the 3Z orientation, the *GFP* reporter is closer to the *eve* TAD (see Figure 1D for diagrams). We
then probed for *LacZ* and *GFP* expression in the anal plate region of the embryo. As expected
from studies on larger *homie* sub-fragments, the relative orientation of the DEF element in the
transgene, but not in the chromosome, determines which reporter is activated: *eve*-like *LacZ*
expression is observed in the anal plate, while *GFP* expression is not detected (Figure 5).
Consistent with the digoxigenin staining, mutation of the Su(Hw) recognition sequence in the D
sub-element impairs LR pairing activity, resulting in a substantial drop in *LacZ* expression (Figure
5).

Our prior studies showed that the orientation of *homie* (CDEF) in the transgene determines
which of the two transgene reporters is activated in an *eve* pattern, and this is true for both
orientations of the transgene in the chromosome (FUJIOKA *et al.* 2009; FUJIOKA *et al.* 2016; BING
et al. 2024). Figure 5 shows that this is also true for DEF. In addition to this, our quantitation of

LacZ mRNA in the anal plate reveals that for DEF, the anal plate enhancer drives a higher level
of reporter expression when the dual reporter transgene is inserted in the 3Z orientation than
when it is in the opposite, Z5, orientation (Figure 5). In this configuration, the topology of the
loop linking *homie* in the transgene to *homie* in the *eve* TAD is a circle-loop.

Next, we tested whether the Su(Hw) site is fully responsible for the activity of D in the
context of our dual reporter transgene. Here, we added back the 3' half of D, which contains
the Su(Hw) site, to EF, which has no LR pairing activity on its own (Figure 3B), to make D3EF.
Figure 6 shows that the addition of the D3 fragment restores LR pairing activity, and now the
lacZ reporter is expressed in an *eve* pattern. However, since the activity of D3EF is noticeably
weaker than DEF, it would appear that sequences in the 5' half of D must interact with a
factor(s) that is able to contribute to LR pairing.

Finally, we tested whether the order of sequences within *homie* is important for its activity.
To do this, we reversed the order of EF adjacent to the Su(Hw) site-containing 3' half of D, to
make D3FE. As shown in Figure 6, with the D3FE transgene, the *eve* enhancers do not activate
lacZ expression. It seemed possible that the change in sub-element order might alter the
orientation dependence of D3FE. Since D3FE is inserted in the dual reporter, we were able to
test this possibility by assaying GFP expression. Figure S4 shows that GFP is not expressed in
the D3FE dual reporter, indicating that LR pairing activity has indeed been lost with the change
in sub-element order, not switched in its orientation preference. Consistent with the
maintenance of orientation specificity with these constructs whenever there is significant LR
pairing, we found that none of the other reporters tested expresses GFP in an *eve* pattern to
any significant degree (DEF, DEFΔSu(Hw): Figure S4).

Contribution of *TER94*-associated elements to LR pairing activity

In the experiments described above, GH showed LR pairing activity in the context of either
CDEF or EF *homie*. G contains the 5'-most *TER94* transcription start site (which is transcribed
away from the *eve* locus and produces transcript *TER94-RA*), while H contains a non-coding
lncRNA start site (CR45324, which is transcribed toward the *eve* locus) and the initiation site for
transcript *TER94-RD* (GRAMATES *et al.* 2022) (see Figure 1B diagram). Because D3EF showed
weak but clear-cut LR pairing activity (Figure 6), we tested whether the adjacent GH region
enhances the LR pairing activity of D3EF. Indeed, D3EFGH shows considerably stronger activity
than does D3EF (Figure S5). We then trimmed back the added region to exclude the start sites
for both the non-coding RNA and *TER94-RD*, which resulted in no apparent loss of LR pairing
activity (D3EFGH5, Figure S5). We further trimmed it to just downstream of the *TER94-RA* start
site, again with no apparent loss of activity (D3EFG5, Figure S5). This suggests that it is the ~50
bp region just downstream of the F region that harbors most, if not all, of the LR pairing activity
of GH. Since this facilitating region includes the *TER94-RA* start site (to +2 nt), it may contain a
basal promoter element from this housekeeping gene. This increased activity is comparable to,
but not quite as strong as, that of CDEF, as evidenced by the more complete *eve* pattern in the
late-stage mesoderm in the latter. It is also worth noting our previous finding that *TER94*
promoter activity is enhanced by sequences downstream of its start site, which are not
included in this insulator-facilitating fragment. So, the sequences facilitating *homie*'s insulating
activity and those responsible for *TER94*'s transcriptional activity seem to be mostly, but
perhaps not entirely, separable.

Enhancer blocking activity only roughly correlates with other insulator activities

We tested many of the same *homie* fragments described above for their ability to block the
hebe enhancer upstream of the –142 kb attP site from activating transgene reporters (Figure 7).
Enhancer blocking activity was tested at the two stages of embryogenesis when the *hebe*
enhancer is active in cells in the CNS midline (stages 12 and 13). One set of fragments was
tested in the context of our single reporter transgene (the eZ vector, Figure 7A), and another,
overlapping set was tested in the double reporter transgene (the eZ-eG vector, Figure 7B). The
two assays gave a consistent order of activities with all of the constructs common to the two
assays. However, the enhancer blocking activity of DE in the eZ-eG vector is reduced relative to
that seen with the eZ vector, suggesting that eZ-eG provides a better context for distinguishing
among stronger enhancer blocking activities, while eZ is better for distinguishing among weaker
activities. The double reporter context also allowed us to see the effectiveness of blocking
expression from one reporter, located on the opposite side of the *homie* element from the
hebe enhancer (Figure S1B), at the same time that this enhancer is seen to activate the
proximal reporter, providing an internal control (for example, GFP expression in the CNS in
Figure S4).

When CDEFGH is located between the reporter and the *hebe* enhancer, it blocks the
enhancer from activating reporter expression (Figure S1A, CDEFGH: only *eve*-like expression is
seen in the CNS). While similar results are observed for both CDEF and EFGH, the ABCD
combination, like λ DNA, has little or no blocking activity, and so the *hebe* enhancers are able to
activate *LacZ* expression in CNS midline cell clusters, mimicking *hebe* expression at this stage
(Figure 2, arrowheads, Figure S2). For the tripartite combinations CDE, DEF, and CDF, blocking
activity follows a pattern roughly similar to LR pairing activity (Figure 7A). DEF is the most
effective enhancer blocker, followed by CDE, and then CDF. Because we obtained a transgenic
line carrying only one of the two possible orientations of CEF (the 3Z orientation), we were not
able to determine its enhancer blocking activity. For the three 2-element combinations DE, DF,
and EF, DE appears to be almost as effective in stage 13 embryos as DEF, while both DF and EF
have only partial blocking activity (Figure 3B), roughly paralleling their LR pairing activities.

Since the Su(Hw) and Cp190 sites contribute to the LR pairing activity of DEF, we tested the
effects of mutations in these recognition sequences. As was the case for LR pairing activity,
mutations in the Cp190 site had a modest effect on DEF's blocking activity (Figure 7B). In
contrast, mutating the Su(Hw) site led to a substantial loss in blocking activity (Figure 7B).
Again, these effects on enhancer blocking activity parallel the effects of these alterations on LR-
pairing activity. However, the correlation is not perfect for all constructs (summarized in Figure
10), suggesting that while the two activities are closely related, there are some mechanistic
differences.

The clearest example is provided by a comparison of the two *homie* derivatives D3EF and
D3FE (in which the order of the E and F regions of DEF *homie* are reversed – neither is inverted
in the chromosome, their order is simply switched). While D3FE is unable to mediate LR pairing
with the *eve* TAD (Figure 6), it has more enhancer blocking activity (showing 45% and 40% *hebe*-
like midline expression at stages 12 and 13, respectively, Figure 7B) than D3EF (with 70% and
54% *hebe*-like midline expression, Figure 7B). This suggests that, at least in the context of the

hebe locus, LR pairing activity is much more sensitive to the order of binding sites for insulator
proteins than is enhancer blocking activity.

Another example is provided by the quantitative effect of the Su(Hw) binding site in DEF
homie. DEFΔSH has a very low level of enhancer blocking activity, but still a clearly measurable
amount of LR pairing activity, while the *gypsy* insulator shows no LR pairing activity, but has
strong enhancer blocking activity (Figure S3, Figure 7B). This suggests that in the context of DEF
homie, Su(Hw) is particularly important for enhancer blocking, while other *homie*-binding
factors that are sufficient for LR pairing contribute little to enhancer blocking. And while
Su(Hw) contributes a significant fraction of DEF *homie*'s LR pairing activity, it contributes even
more of its enhancer blocking activity. This again speaks to the specificity of our LR pairing
assay, which reflects the ability of *homie* to specifically pair with copies of itself when the two
are located at considerable chromosomal distances from each other. Enhancer blocking in this
assay is a more local phenomenon, one that may not depend on specific pairing with another
insulator.

We used the PRE blocking and enhancer blocking assays to further compare the effects of
the Su(Hw) binding site in fragment D. D alone, but not E or F alone, has measurable activity in
the enhancer blocking assay (Figure 7C). When the Su(Hw) site is mutated, it abolishes activity
in the context of D alone (Figure 7C). Since E alone has no clearly measurably activity in the
enhancer blocking assay, mutating the Cp190-associated site can show no effect in this context.

PRE blocking activity

We showed previously that *homie* has the ability to block the spread of repressive
chromatin (barrier activity) from an *eve* pseudo-locus transgene into the neighboring gene,
TER94. This spreading depends on the *homie*-adjacent *eve* PRE (Fujioka *et al.* 2013). *TER94*
promoter activity is repressed when *homie* is removed, and this correlates with the spreading
of the histone modification H3K27me3, characteristic of PcG-dependent repressive chromatin.
This loss of expression is not due to reduced *TER94* promoter activity caused by removing
homie, since removing the *eve* PRE in addition to *homie* fully restores expression (Fujioka *et al.*
2013). In this transgenic context, when *homie* is present, *TER94* promoter-driven GFP
expression is high in ovaries dissected from adult females.

When full-length *homie* (ABCDEF) is replaced by CDEF, or any of the tripartite combinations
(DEF, CEF, CDF, or DEF) *TER94*-driven GFP expression is undiminished (Figure S6A, B), indicating
that PRE blocking is still complete. This is illustrated for DEF and "wt" in Figure 8A.
Furthermore, DE alone appears to have full activity (Figure S6A, B), while AB has no barrier
activity (Figure 8A, B). As shown in Figure 8A and B, we also tested individual *homie* sub-
elements. Of these, C has no barrier activity, while E alone is sufficient to completely block the
PRE's repressive effect. D and F both have partial barrier activity: GFP expression is diminished,
but not as much as when *homie* is completely replaced by λ DNA (compare λ DNA with D and E
in Figure 8A). It is also worth noting that this barrier activity does not depend on the
orientation of *homie* (Figure S6A, B, CDEF vs. FEDC).

The single *homie* sub-elements that retain barrier activity show some interesting
distinctions in their PRE blocking activities. D has strong activity in the somatic follicle cells that
form the outer layer of the ovariole, but clearly has less activity in the inner, germline cells
(nurse cells and oocyte). In contrast, E seems to have full activity in both the germline cells and

the follicle cells. F has somewhat less activity than E overall, and appears to have more activity
in germline cells than in follicle cells (compare D and F to E in Figure 8A, B). These results show
that there can be tissue specificity in barrier activity, at least when somatic follicle cells are
compared to ovarian germline cells.

Above, we showed that the Su(Hw) binding site in D is required for most, but not all, of the
activity of the D region in the LR pairing assay. We tested the same mutation here, and found
that this site is required for the D region to show any barrier activity (Figure 9A, B). We also
tested the requirement for the Cp190-associated site in region E. However, it does not seem to
have any effect on the barrier activity of E (Figure 9A, B). This last set of experiments was
performed in a different context than the one shown in Figure 8. Instead of the full-length *eve*
locus in the transgene, we used only the 3' end of the locus, which contains the *eve* 3' PRE,
homie, and the *TER94* promoter driving GFP (diagrammed at the top of Figure 9A). Despite the
absence of most of the *eve* locus from the transgene, the results for regions D and E (Figure 9)
are very similar to those shown in Figure 8. However, the germline specificity seen for the F
region in Figure 8 is not seen here. Instead of having less activity in follicle cells, here it shows
the opposite specificity, with stronger activity in follicle cells than in germline cells. We note
that in addition to the transgene construct being different, the insertion site of the transgene is
different. It will require further study to determine why the barrier activity of region F is so
sensitive to the chromosomal context, while that of regions D and E do not show this
sensitivity. As expected, when the PRE is not present in the transgene, there is no repression of
TER94 promoter-driven GFP expression, even in the absence of any insulator sequences ("no
PRE, no *homie*", Figure 9; here, both the PRE and the starting *homie* fragment are replaced by
equal-length stretches of λ DNA).

Discussion

A primary function of boundary elements/insulators in *Drosophila* is the subdivision of the chromosome into a series of looped domains, or TADs. TAD formation is thought to depend upon the physical pairing of neighboring boundary elements. In most cases that have been examined in detail, these pairing interactions are orientation-dependent (KYRCHANOVA *et al.* 2008; FUJIOKA *et al.* 2016). Orientation dependence determines the topology of the TAD. When boundaries pair with their neighbors head-to-head, a circle-loop is generated, while head-to-tail pairing generates a stem-loop. Since the two boundaries that delimit the *eve* TAD, *homie* and *nhomie*, pair with each other head-to-tail, the *eve* TAD is a stem-loop (KE *et al.* 2024). Like other fly boundaries, *homie* and *nhomie* also engage in self-pairing interactions, and in these cases, pairing is head-to-head (KYRCHANOVA *et al.* 2008; FUJIOKA *et al.* 2016). Unlike heterologous pairing, which occurs in *cis*, self-pairing interactions take place between the same boundary element in *trans* and are thought to be important for mediating the juxtapositioning and precise alignment of homologs.

In their endogenous location, *homie* and *nhomie* are separated by only 16 kb; however, both elements can pair with themselves and with each other when separated by multiple TADs and TAD boundaries (FUJIOKA *et al.* 2009; FUJIOKA *et al.* 2016; CHEN *et al.* 2018; BING *et al.* 2024). In addition to this long-range (LR) pairing activity, *homie* can block enhancer-promoter interactions (FUJIOKA *et al.* 2009; FUJIOKA *et al.* 2016; CHEN *et al.* 2018; BING *et al.* 2024) and act as a barrier to the spread of PcG silencing (FUJIOKA *et al.* 2013). In the studies reported here, we have undertaken a functional dissection of the *homie* boundary and identified sequences/sub-elements that are important for these three activities.

homie was originally defined as an 800 bp fragment that extended from a site close to the *eve* PRE to just beyond the 1st exon of the neighboring *TER94* gene, a region associated with DNase hypersensitivity (SACKERSON *et al.* 1999). For the purpose of analysis, this 800 bp fragment was subdivided into eight ~100 bp sequences or sub-elements, ABCDEFGH. As shown in Figure 1, G extends from upstream of the *TER94* transcription start site into the first *TER94-RA* mRNA exon, while H includes sequences from the first exon and the beginning of the first intron, including the start site for a *lncRNA* and the *TER94-RD* mRNA.

Barrier activity: Of the three activities we assayed, the least demanding is blocking PcG silencing. While both LR pairing and enhancer blocking activity require a combination of several sub-elements, single sub-elements are able to block PcG silencing to a greater or lesser extent. Of the sub-elements tested, D, E, and F are able to inhibit the spread of PcG silencing completely on their own (Figures 8 and 9). In contrast, AB and C are unable to function as barriers against PcG silencing. Since the *TER94* reporter in the mini-locus that contains a deletion of ABCDEF is also silenced by the *eve* PRE, the GH elements are unable to prevent PcG silencing (Figure 9). While the E sub-element is as effective in protecting the *TER94* reporter as the larger DEF element or the *homie* element in the intact pseudo-locus (ABCDEFGH), this is not the case for either D of F. The barrier activity of F is about half that of E or DEF and appears to be position-dependent, while the barrier activity of D is tissue-specific (Figures 8 and 9): it is greatly reduced in germline nurse cells, while it is close to that of DEF in somatic follicle cells. Interestingly, in spite of the clear tissue-specific difference in barrier activity, the D sub-element

requires the Su(Hw) binding site in both cell types. We also tested whether the barrier activity
of the E sub-element requires the Cp190-associated sequence; it does not.

Enhancer blocking/LR pairing: The requirements for enhancer blocking and LR pairing activity
are more stringent than for PcG barrier function. Among the single sub-elements, only D has
significant enhancer blocking activity (Figure 7C), and none of them can mediate LR pairing.
While the DE combination has both activities, its LR pairing activity is very weak (Figure 3B) and
its blocking activity is incomplete (Figure 7B). Of the tripartite sub-element combinations,
robust enhancer blocking and LR pairing activity are only observed with the DEF combination.
Consistent with a link between enhancer blocking and LR pairing activity, we find that in most
but not all cases, the different combinations of sub-elements exhibit a similar trend with
respect to their blocking and LR pairing activity. For example, among the *homie* combinations
with four sub-elements that we tested (Figures 2 and 7), CDEF has robust blocking and LR
pairing activity, and while EFGH blocks the *hebe* enhancers, blocking is incomplete and LR
pairing activity is only modest. The ABCD combination has only very little blocking activity and
no LR pairing activity. The same is true for the four tripartite combinations. The blocking
activity of DEF is equivalent to that of CDEF, while LR pairing activity is at most only slightly
reduced. At the other end of the spectrum, CDF has only weak blocking activity and virtually no
LR pairing activity. CDE has somewhat weaker blocking activity , and modest LR pairing. This is
likely because the DE element has some blocking and LR pairing activity on its own. The other
combinations, DF and EF, retain a low level of blocking activity, but have at most a very low
level of LR pairing activity.

Su(Hw) and Cp190 motifs: D has a sequences motif for the Su(Hw) protein while E has a
Cp190-associated sequence. Both of these proteins have been implicated in the functioning of
other fly boundaries. Consistent with an important role in *homie* activity, mutations in the
Su(Hw) site in the DEF combination led to a substantial drop in both blocking and LR pairing
activity (Figures 4 and 7). In contrast, Cp190 appears to have a much less important role, as
disruption of the Cp190-associated sequence had only a minor effect on blocking activity and
caused just a modest reduction in LR pairing. This was also true for barrier activity. Mutations
in the Su(Hw) recognition motif eliminated the barrier activity of the D element, while
mutations in the Cp190-associated sequence had minimal if any effect (Figures 7 and 9).

A previous study of the effects of a mutation in the *Cp190* gene on insulator activities
concluded that it affects enhancer blocking but not LR pairing by *homie* (KAUSHAL *et al.* 2022).
While our results with the Cp190-associated sequence mutation also implicate Cp190 in
enhancer blocking, the effects are not as dramatic as the *Cp190* gene mutation. Likewise, while
Kaushal et al. (KAUSHAL *et al.* 2022) observed little if any effect on LR pairing activity, our results
suggest that Cp190 contributes to *homie*'s LR pairing activity, albeit modestly. It is worth noting
here that Cp190 may not bind directly to DNA, but may instead function through interactions
with DNA binding proteins, including one that binds to the consensus "Cp190-associated" site
(CUARTERO *et al.* 2014) in *homie*.

The D sub-element: Both the enhancer blocking and LR pairing activities of DEF are nearly
equivalent to those of CDEF. While mutations in the D sub-element Su(Hw) recognition
sequence largely eliminates both blocking and LR pairing activity (Figure 4, Figure 7C),
combining a 50 bp sequence from the 3' half of D (D3), which contains the Su(Hw) site, with EF
(to give D3EF) does not restore the blocking and LR pairing activities of DEF (Figure 6, Figure

7B). The blocking activity of the D3EF fragment is close to that of DEFΔSH, while the LR pairing
activity is only marginally better than EF by itself. These findings argue that in addition to the
Su(Hw) recognition sequence, sequences elsewhere in D are required for both its enhancer
blocking and LR pairing activities.

Order of sub-elements is important for LR activity: Our analysis of the D3EF combination
uncovered another important feature of *homie* boundary activity. We found that the order of
the E and F sub-elements in combination with D3 is important for LR pairing interactions, but
not for enhancer blocking (Figure 6, Figure 7B). Thus, while D3EF mediates LR interactions,
changing the order of the EF sub-elements to give D3FE completely disrupts LR pairing activity.

This loss of LR pairing activity makes sense, as it is thought that the order of binding sites for
chromosomal architectural proteins like Su(Hw), CTCF and Pita is likely to play an important role
in generating orientation-dependent boundary:boundary pairing interactions (MAKSIMENKO *et al.*
2015). Members of this particular class of polydactyl zinc finger proteins have self-interaction
domains that can generate multimeric complexes and potentially directly link boundary
elements that have binding sites in common (FEDOTOVA *et al.* 2017; BONCHUK *et al.* 2021).

However, it has also been suggested that the Su(Hw) consensus site should be extended
beyond the core, because Su(Hw) contacts sequences flanking the core consensus binding site
using a part of the protein that contributes to insulator activity (BAXLEY *et al.* 2017). Since the
Su(Hw) site is very near the D-E junction, swapping E and F separates this extended region from
the core site, which might cause a reduction in Su(Hw) binding or insulator activity, and thereby
contribute to the observed loss of LR pairing. On the other hand, the D region without this
extended portion of the Su(Hw) site retains both enhancer blocking (Figure 7) and PRE blocking
activities (Figure 9) that are totally dependent on the core Su(Hw) binding site (D vs. DΔSH),
suggesting that the site has strong activity without the extended region. Therefore, it seems
unlikely that the large difference in LR pairing activity between D3EF and D3FE is due solely, or
even primarily, to a loss of Su(Hw) binding to the core consensus site that is present in both.

Changing the order of E and F in D3EF (which already has rather modest LR pairing activity)
to give D3FE could interfere with LR pairing activity at several levels. In order for the *eve*
enhancers to activate reporter expression at a distance of almost 150 kb and across nearly a
dozen intervening TADs, the transgene *homie* must recognize potential partners (*homie* and/or
nhomie) in the *eve* TAD and "initiate" physical pairing (BING *et al.* 2024). Changing the order of
E and F may make it impossible for proteins associate with each sub-element to simultaneously
interact with their partners in the endogenous *homie* and *nhomie*, and the initiation step could
fail. It is also possible that the initial interaction might involve only a subset of proteins
associated with each boundary. In this case, factors associated with a single sub-element might
be sufficient to initiate pairing, but steric hinderance arising from the mis-ordered sub-elements
could prevent the formation of a sufficient number of physical links (a zippering up) between
the boundary elements to generate a stable pairing interaction. Live imaging experiments
indicate that after pairing interactions are established and stabilized, activation of the
transgene reporter by the *eve* enhancers requires a further interaction that appears to involve
forming contacts between the *eve* enhancers and the promoter of the reporter (CHEN *et al.*
2018). Though less likely, this step might also be perturbed by steric hinderance arising from
misaligned sub-element-mediated contacts.

While D3FE is defective in LR pairing, this rearranged boundary nevertheless retains
enhancer blocking activity (Figure 7B). Several mechanisms have been proposed to explain how
boundaries block enhancers. Boundaries could act as roadblocks or sinks (GEYER 1997;
BLACKWOOD AND KADONAGA 1998; BI AND BROACH 1999; BLANTON et al. 2003; GOHL et al. 2011), in
which case blocking activity is autonomous and will depend on functional properties of the
proteins bound to D3EF and D3FE. Alternatively, insulation could be achieved by organizing the
chromatin fiber into looped domains (BING et al. 2024). In this case, the blocking activity of
D3EF and D3FE would depend upon whether boundaries in the neighborhood can pair with the
two homie derivatives. A 2-dimensional roadblock model cannot explain the pattern of MicroC
interactions within TADs, nor would it explain how E-P interactions between sequences that are
142 kb apart result in one reporter being activated and the other nearby reporter not (BING et
al. 2024). So it seems likely that blocking activity must be envisioned as a process in 3
dimensions that builds on the TAD organization of the chromosome.

Figure 10 summarizes all our results comparing the gypsy insulator and the various homie
subelements and their derivatives in our assays for LR pairing and enhancer blocking. The
constructs are listed in the order of their relative activities in each assay, highest to lowest,
from top to bottom. The 3 lines between the two columns connect constructs which are most
clearly "in different places" in the two lists. This indicates that their activities do not follow the
same trend in the two assays, suggesting that these two insulator activities have some
mechanistic distinctions. In particular, gypsy does not show LR pairing with the eve locus, but is
capable of enhancer blocking in our assay, which measures enhancer blocking at short range in
cis. This distinction may arise from the strong requirement for pairing specificity in the LR
pairing assay. The ability to pair with other insulators in the vicinity of the insertion site may be
sufficient for enhancer blocking, but not for LR pairing. Alternatively, enhancer blocking may
not require insulator pairing at all, even though it is clear that pairing can affect which
interactions are blocked and which are facilitated by insulators. Similar considerations may
explain the disparate positions of D3FE in the two lists, as discussed above. The discrepancy in
the positions of D3EF between the two lists is harder to explain. This activity, harbored within
the 5' portion of region D, appears to be more important for enhancer blocking in the context
of DEF than it is for LR pairing. Further investigation will be required to understand the basis of
this activity, but its existence suggests that there are yet to be uncovered mechanisms
underlying these distinctive insulator activities.

Enhancer blocking/PRE blocking: We also compared the enhancer blocking and PRE
blocking activities of an extensive set of homie subelements. These comparisons provide a
clear indication of a mechanistic distinction between enhancer blocking and PRE blocking by
homie: as Figure 7C shows, D has clearly more enhancer blocking activity than E (which has
little to none), while the opposite is true in the PRE blocking assay, where E has strong activity,
and D's is considerably weaker (particularly in the germline; Figures 8 and 9). One possible
explanation is that PRE blocking only requires the introduction of an extended nucleosome-
depleted region in 1-dimensional space, along the chromosome (reviewed in (GASZNER AND
FELSENFELD 2006; BUSHEY et al. 2008), while enhancer blocking may require an additional ability
to suppress looping in 3 dimensions.

Materials and Methods

Plasmid construction and transgenic fly production

Construction of eZ (used for Figures 2, 3, 7A, 7C and Figure S1) and eZ-eG (used for Figures 4, 5, 6, 7B and Suppl. Figures 2 – 5) vectors used for the LR interaction and enhancer blocking assays was described previously (FUJIOKA *et al.* 2009; FUJIOKA *et al.* 2016). These vectors carrying modified *homie* sequences were inserted into the attP site at –142kb relative to the *eve* locus (FUJIOKA *et al.* 2009). Analysis of *hebe* enhancer activity (Figure S1B) was done using the eZ vector, inserted into the attP site at cytological location 74A2 (FUJIOKA *et al.* 2013).

The *eve* pseudo-locus construct used for analyzing PRE blocking activity was described previously (FUJIOKA *et al.* 2013). The attP sites at 74A2 (used for Figure 8) and 95E5 (for Figure S6) were used. The smaller construct used to analyze PRE blocking activity (used for Figure 9) is the same as the pseudo-locus, except that the sequence upstream of +8.4 kb (relative to the *eve* transcription start site) was removed. The constructs using this vector were inserted into the miMIC 24B1 attP site (VENKEN *et al.* 2011; FUJIOKA *et al.* 2021). In both constructs, the *homie* region from +9.2 to +9.8 kb (relative to *eve* +1) were replaced with modified *homie* and control sequences.

Analysis of transgenic lines

In situ hybridization was performed as described previously (FUJIOKA *et al.* 2013). RNA was visualized using alkaline phosphatase conjugated anti-DIG antibody (Roche), using CBIP and NBT as substrates (Roche). Each set of experiments was carried out with control and experimental samples in parallel to minimize experimental variation. Representative expression patterns are shown in each figure. The same embryo populations were also subjected to counting of CNS midline cell clusters. Visible clusters were counted in about 20 embryos for each construct.

Ovaries were dissected from adult female flies aged 18-22 hr. Live images of ovaries were obtained using the same camera settings throughout, except as noted in the figures. In order to quantify GFP expression, ovaries were subjected to RT-PCR. Total RNA was extracted from 5-8 pairs of ovaries using an RNA extraction kit (Roche). 50 ng of total RNA was used to make cDNA using the Transcriptor First Strand cDNA Synthesis kit (Roche) with random primer. One μl out of 50 μl cDNA was used for each quantitative PCR reaction, each time performed in triplicate. Data are shown as GFP expression normalized to endogenous *TER94* RNA expression, similarly quantified in each sample. Each set of lines was analyzed in three independent assays. smFISH (TRCEK *et al.* 2017; LITTLE AND GREGOR 2018) was performed as previously described (KE *et al.* 2024).

Binding site analysis

Binding site searches was done using MacVector software (MacVector Inc). The Su(Hw) consensus sequences YWGCATACTTTT (RAMOS *et al.* 2006) and NWWWWNYRTWGCATACTTTTKGSDB (BAXLEY *et al.* 2017), and the Cp190-associated consensus sequence GGTTBDWRWMYYNGCTD (CUARTERO *et al.* 2014), were used. The results are shown in Figure S7.

Data Availability Statement

All data underlying this publication are included in the text and figures.

Acknowledgements

This work was supported by NIH grants to Paul Schedl (5R35GM126975) and James B. Jaynes (1R01GM137062). We thank Qing Liu for excellent technical assistance.

References

Aoki T, Sarkeshik A, Yates J, Schedl P. 2012. Elba, a novel developmentally regulated chromatin boundary factor is a hetero-tripartite DNA binding complex. eLife. 1:e00171.

Bag I, Chen S, Rosin LF, Chen Y, Liu CY, Yu GY, Lei EP. 2021. M1bp cooperates with cp190 to activate transcription at tad borders and promote chromatin insulator activity. Nat Commun. 12(1):4170.

Barges S, Mihaly J, Galloni M, Hagstrom K, Muller M, Shanower G, Schedl P, Gyurkovics H, Karch F. 2000. The fab-8 boundary defines the distal limit of the bithorax complex iab-7 domain and insulates iab-7 from initiation elements and a pre in the adjacent iab-8 domain. Development. 127(4):779-790.

Bateman JR, Lee AM, Wu CT. 2006. Site-specific transformation of drosophila via phic31 integrase-mediated cassette exchange. Genetics. 173(2):769-777.

Baxley RM, Bullard JD, Klein MW, Fell AG, Morales-Rosado JA, Duan T, Geyer PK. 2017. Deciphering the DNA code for the function of the drosophila polydactyl zinc finger protein suppressor of hairy-wing. Nucleic Acids Res. 45(8):4463-4478.

Bi X, Broach JR. 1999. Uasrpg can function as a heterochromatin boundary element in yeast. Genes Dev. 13(9):1089-1101.

Bing X, Ke W, Fujioka M, Kurbidaeva A, Levitt S, Levine M, Schedl P, Jaynes JB. 2024. Chromosome structure i: Loop extrusion or boundary: Boundary pairing? eLife. in press; preprint: https://elifesciences.org/reviewed-preprints/94070.

Blackwood EM, Kadonaga JT. 1998. Going the distance: A current view of enhancer action. Science. 281(5373):60-63.

Blanton J, Gaszner M, Schedl P. 2003. Protein:Protein interactions and the pairing of boundary elements in vivo. Genes Dev. 17(5):664-675.

Bonchuk A, Boyko K, Fedotova A, Nikolaeva A, Lushchekina S, Khrustaleva A, Popov V, Georgiev P. 2021. Structural basis of diversity and homodimerization specificity of zinc-finger-associated domains in drosophila. Nucleic Acids Res. 49(4):2375-2389.

Bushey AM, Dorman ER, Corces VG. 2008. Chromatin insulators: Regulatory mechanisms and epigenetic inheritance. Mol Cell. 32(1):1-9.

Chen H, Levo M, Barinov L, Fujioka M, Jaynes JB, Gregor T. 2018. Dynamic interplay between enhancer–promoter topology and gene activity. Nat Genet. 50(9):1296.

Chetverina D, Fujioka M, Erokhin M, Georgiev P, Jaynes JB, Schedl P. 2017. Boundaries of loop domains (insulators): Determinants of chromosome form and function in multicellular eukaryotes. Bioessays. 39(3):1600233.

Cuartero S, Fresan U, Reina O, Planet E, Espinas ML. 2014. Ibf1 and ibf2 are novel cp190-interacting proteins required for insulator function. EMBO J. 33(6):637-647.

De S, Gehred ND, Fujioka M, Chan FW, Jaynes JB, Kassis JA. 2020. Defining the boundaries of polycomb domains in drosophila. Genetics.

Fedotova A, Aoki T, Rossier M, Mishra RK, Clendinen C, Kyrchanova O, Wolle D, Bonchuk A, Maeda RK, Mutero A. 2018. The ben domain protein insensitive binds to the fab-7 chromatin boundary to establish proper segmental identity in drosophila. Genetics. 210(2):573-585.

Fedotova AA, Bonchuk AN, Mogila VA, Georgiev PG. 2017. C2h2 zinc finger proteins: The largest
but poorly explored family of higher eukaryotic transcription factors. Acta Naturae.
9(2):47-58.
Fujioka M, Emi-Sarker Y, Yusibova GL, Goto T, Jaynes JB. 1999. Analysis of an *even-skipped*
rescue transgene reveals both composite and discrete neuronal and early blastoderm
enhancers, and multi-stripe positioning by gap gene repressor gradients. Development.
126(11):2527-2538.
Fujioka M, Mistry H, Schedl P, Jaynes JB. 2016. Determinants of chromosome architecture:
Insulator pairing in *cis* and in *trans*. PLoS Genet. 12(2):e1005889.
Fujioka M, Nezdyur A, Jaynes JB. 2021. An insulator blocks access to enhancers by an
illegitimate promoter, preventing repression by transcriptional interference. PLoS
Genet. 17(4):e1009536.
Fujioka M, Sun G, Jaynes JB. 2013. The drosophila *eve* insulator homie promotes *eve* expression
and protects the adjacent gene from repression by polycomb spreading. PLoS Genet.
9(10):e1003883.
Fujioka M, Wu X, Jaynes JB. 2009. A chromatin insulator mediates transgene homing and very
long-range enhancer-promoter communication. Development. 136(18):3077-3087.
Fujioka M, Yusibova GL, Zhou J, Jaynes JB. 2008. The DNA-binding polycomb-group protein
pleiohomeotic maintains both active and repressed transcriptional states through a
single site. Development. 135(24):4131-4139.
Gaszner M, Felsenfeld G. 2006. Insulators: Exploiting transcriptional and epigenetic
mechanisms. Nat Rev Genet. 7(9):703-713. Epub 2006 Aug 2015.
Gaszner M, Vazquez J, Schedl P. 1999. The zw5 protein, a component of the scs chromatin
domain boundary, is able to block enhancer-promoter interaction. Genes Dev.
13(16):2098-2107.
Geyer PK. 1997. The role of insulator elements in defining domains of gene expression. Curr
Opin Genet Dev. 7(2):242-248.
Geyer PK, Corces VG. 1992. DNA position-specific repression of transcription by a drosophila
zinc finger protein. Genes Dev. 6(10):1865-1873.
Ghosh D, Gerasimova TI, Corces VG. 2001. Interactions between the su(hw) and mod(mdg4)
proteins required for gypsy insulator function. Embo J. 20(10):2518-2527.
Gohl D, Aoki T, Blanton J, Shanower G, Kappes G, Schedl P. 2011. Mechanism of chromosomal
boundary action: Roadblock, sink, or loop? Genetics. 187(3):731-748.
Gramates LS, Agapite J, Attrill H, Calvi BR, Crosby MA, Dos Santos G, Goodman JL, Goutte-Gattat
D, Jenkins VK, Kaufman T et al. 2022. Flybase: A guided tour of highlighted features.
Genetics. 220(4).
Holdridge C, Dorsett D. 1991. Repression of hsp70 heat shock gene transcription by the
suppressor of hairy-wing protein of drosophila melanogaster. Mol Cell Biol. 11(4):1894-
1900.
Jiang N, Emberly E, Cuvier O, Hart CM. 2009. Genome-wide mapping of boundary element-
associated factor (beaf) binding sites in drosophila melanogaster links beaf to
transcription. Mol Cell Biol. 29(13):3556-3568.
Karch F, Galloni M, Sipos L, Gausz J, Gyurkovics H, Schedl P. 1994. Mcp and fab-7: Molecular
analysis of putative boundaries of cis-regulatory domains in the bithorax complex of
drosophila melanogaster. Nucleic Acids Res. 22(15):3138-3146.

658 Kaushal A, Dorier J, Wang B, Mohana G, Taschner M, Cousin P, Waridel P, Iseli C, Semenova A,
659   Restrepo S et al. 2022. Essential role of cp190 in physical and regulatory boundary
660   formation. Sci Adv. 8(19):eabl8834.
661 Ke W, Fujioka M, Schedl P, Jaynes JB. 2024. Chromosome structure ii: Stem-loops and circle-
662   loops. eLife. in press; preprint: https://elifesciences.org/reviewed-
663   preprints/94114
664 Kyrchanova O, Chetverina D, Maksimenko O, Kullyev A, Georgiev P. 2008. Orientation-
665   dependent interaction between drosophila insulators is a property of this class of
666   regulatory elements. Nucleic Acids Res. 36(22):7019-7028.
667 Leon A, McKearin D. 1999. Identification of ter94, an aaa atpase protein, as a bam-dependent
668   component of the drosophila fusome. Molecular Biology of the Cell. 10(11):3825-3834.
669 Li H-B, Muller M, Bahechar IA, Kyrchanova O, Ohno K, Georgiev P, Pirrotta V. 2011. Insulators,
670   not polycomb response elements, are required for long-range interactions between
671   polycomb targets in drosophila melanogaster. Mol Cell Biol. 31(4):616-625.
672 Li J, Gilmour DS. 2013. Distinct mechanisms of transcriptional pausing orchestrated by gaga
673   factor and m1bp, a novel transcription factor. EMBO J. 32(13):1829-1841.
674 Little SC, Gregor T. 2018. Single mrna molecule detection in drosophila. Methods Mol Biol.
675   1649:127-142.
676 Maksimenko O, Bartkuhn M, Stakhov V, Herold M, Zolotarev N, Jox T, Buxa MK, Kirsch R,
677   Bonchuk A, Fedotova A. 2015. Two new insulator proteins, pita and zipic, target cp190
678   to chromatin. Genome Res. 25(1):89-99.
679 Maksimenko O, Golovnin A, Georgiev P. 2008. Enhancer-promoter communication is regulated
680   by insulator pairing in a drosophila model bigenic locus. Mol Cell Biol. 28(17):5469-5477.
681 Modolell J, Bender W, Meselson M. 1983. Drosophila melanogaster mutations suppressible by
682   the suppressor of hairy-wing are insertions of a 7.3-kilobase mobile element. Proc Natl
683   Acad Sci USA. 80(6):1678-1682.
684 Moon H, Filippova G, Loukinov D, Pugacheva E, Chen Q, Smith ST, Munhall A, Grewe B,
685   Bartkuhn M, Arnold R et al. 2005. Ctcf is conserved from drosophila to humans and
686   confers enhancer blocking of the fab-8 insulator. EMBO Rep. 6(2):165-170.
687 Muller M, Hagstrom K, Gyurkovics H, Pirrotta V, Schedl P. 1999. The mcp element from the
688   drosophila melanogaster bithorax complex mediates long-distance regulatory
689   interactions. Genetics. 153(3):1333-1356.
690 Özdemir I, Gambetta MC. 2019. The role of insulation in patterning gene expression. Genes
691   (Basel). 10(10).
692 Pai CY, Lei EP, Ghosh D, Corces VG. 2004. The centrosomal protein cp190 is a component of the
693   gypsy chromatin insulator. Mol Cell. 16(5):737-748.
694 Parkhurst SM, Harrison DA, Remington MP, Spana C, Kelley RL, Coyne RS, Corces VG. 1988. The
695   drosophila su(hw) gene, which controls the phenotypic effect of the gypsy transposable
696   element, encodes a putative DNA-binding protein. Genes Dev. 2(10):1205-1215.
697 Peifer M, Bender W. 1988. Sequences of the gypsy transposon of drosophila necessary for its
698   effects on adjacent genes. Proc Natl Acad Sci USA. 85(24):9650-9654.
699 Pinter M, Jekely G, Szepesi RJ, Farkas A, Theopold U, Meyer HE, Lindholm D, Nassel DR,
700   Hultmark D, Friedrich P. 1998. Ter94, a drosophila homolog of the membrane fusion
701   protein cdc48/p97, is accumulated in nonproliferating cells: In the reproductive organs
702   and in the brain of the imago. Insect Biochemistry & Molecular Biology. 28(2):91-98.

Ramírez F, Bhardwaj V, Arrigoni L, Lam KC, Grüning BA, Villaveces J, Habermann B, Akhtar A,
Manke T. 2018. High-resolution tads reveal DNA sequences underlying genome
organization in flies. Nature communications. 9(1):189.
Ramos E, Ghosh D, Baxter E, Corces VG. 2006. Genomic organization of gypsy chromatin
insulators in drosophila melanogaster. Genetics. 172(4):2337-2349.
Rath U, Ding Y, Deng H, Qi H, Bao X, Zhang W, Girton J, Johansen J, Johansen KM. 2006. The
chromodomain protein, chromator, interacts with jil-1 kinase and regulates the
structure of drosophila polytene chromosomes. J Cell Sci. 119(Pt 11):2332-2341.
Roy S, Gilbert MK, Hart CM. 2007. Characterization of beaf mutations isolated by homologous
recombination in drosophila. Genetics. 176(2):801-813.
Ruden DM, Sollars V, Wang X, Mori D, Alterman M, Lu X. 2000. Membrane fusion proteins are
required for oskar mrna localization in the drosophila egg chamber. Dev Biol.
218(2):314-325.
Sackerson C, Fujioka M, Goto T. 1999. The *even-skipped* locus is contained in a 16-kb chromatin
domain. Dev Biol. 211(1):39-52.
Schwartz YB, Cavalli G. 2017. Three-dimensional genome organization and function in
drosophila. Genetics. 205(1):5-24.
Schwartz YB, Kahn TG, Nix DA, Li XY, Bourgon R, Biggin M, Pirrotta V. 2006. Genome-wide
analysis of polycomb targets in drosophila melanogaster. Nat Genet. 38(6):700-705.
Sigrist CJ, Pirrotta V. 1997. Chromatin insulator elements block the silencing of a target gene by
the drosophila polycomb response element (pre) but allow trans interactions between
pres on different chromosomes. Genetics. 147(1):209-221.
Small S, Blair A, Levine M. 1992. Regulation of even-skipped stripe 2 in the drosophila embryo.
The EMBO journal. 11(11):4047-4057.
Small S, Blair A, Levine M. 1996. Regulation of two pair-rule stripes by a single enhancer in the
drosophila embryo. Dev Biol. 175(May 1):314-324.
Tolhuis B, de Wit E, Muijrers I, Teunissen H, Talhout W, van Steensel B, van Lohuizen M. 2006.
Genome-wide profiling of prc1 and prc2 polycomb chromatin binding in drosophila
melanogaster.[erratum appears in nat genet. 2006 jul;38(7):850]. Nat Genet. 38(6):694-
699.
Trcek T, Lionnet T, Shroff H, Lehmann R. 2017. Mrna quantification using single-molecule fish in
drosophila embryos. Nat Protoc. 12(7):1326-1348.
Vazquez J, Muller M, Pirrotta V, Sedat JW. 2006. The mcp element mediates stable long-range
chromosome-chromosome interactions in drosophila. Molecular Biology of the Cell.
17(5):2158-2165.
Venken KJT, Schulze KL, Haelterman NA, Pan H, He Y, Evans-Holm M, Carlson JW, Levis RW,
Spradling AC, Hoskins RA et al. 2011. Mimic: A highly versatile transposon insertion
resource for engineering drosophila melanogaster genes. Nat Methods. 8(9):737-743.
Viets K, Sauria ME, Chernoff C, Viales RR, Echterling M, Anderson C, Tran S, Dove A, Goyal R,
Voortman L. 2019. Characterization of button loci that promote homologous
chromosome pairing and cell-type-specific interchromosomal gene regulation. Dev Cell.
51(3):341-356. e347.
Zhao K, Hart CM, Laemmli UK. 1995. Visualization of chromosomal domains with boundary
element-associated factor beaf-32. Cell. 81(6):879-889.

Zhou J, Ashe H, Burks C, Levine M. 1999. Characterization of the transvection mediating region
of the abdominal-b locus in drosophila. Development. 126(14):3057-3065.
Zolotarev N, Fedotova A, Kyrchanova O, Bonchuk A, Penin AA, Lando AS, Eliseeva IA,
Kulakovskiy IV, Maksimenko O, Georgiev P. 2016. Architectural proteins pita, zw5,and
zipic contain homodimerization domain and support specific long-range interactions in
drosophila. Nucleic Acids Res. 44(15):7228-7241.
753

Figure legends

Figure 1. Maps of the *eve* TAD and transgenes inserted at –142kb from the *eve* locus.
A: map of the *eve* locus. Blue: stripe enhancers (SMALL *et al.* 1992; SMALL *et al.* 1996; FUJIOKA *et al.* 1999; SACKERSON *et al.* 1999). Pink: neuronal enhancers (FUJIOKA *et al.* 1999; SACKERSON *et al.* 1999). Orange: tissue-specific enhancers (FUJIOKA *et al.* 1999; SACKERSON *et al.* 1999). Yellow: PRE (FUJIOKA *et al.* 2008). Red: insulators *homie* (H) and *nhomie* (N) (FUJIOKA *et al.* 2009; FUJIOKA *et al.* 2016). Green arrowheads: DNase I hypersensitive sites (SACKERSON *et al.* 1999). **B: blowup of the *homie* region, subelements ABCDEFGH.** The *TER94-RA* transcript starts in G. Both the *TER94-RD* transcript and the *lncRNA:CR45324* start in H (GRAMATES *et al.* 2022). The positions of the Su(Hw) binding site and the Cp190-associated site (RAMOS *et al.* 2006; CUARTERO *et al.* 2014; BAXLEY *et al.* 2017) are shown as red ovals. **C, D. the eZ-CDEF and eZ-CDEF-eG transgenes.** The –142kb attP site is located in the 5' UTR of the *hebe* transcription unit. Because the attP site was originally inserted via P-element transposition, the 5'- and 3'-P-element ends flank it in the genome. The 3'-P-end ("3'P") is located closer to the CNS midline enhancer of the *hebe* gene, while the 5'-P-end ("5'P") is closer to the *eve* locus. The *hebe* transcription unit (all major transcripts start at the same site, according to Flybase) (GRAMATES *et al.* 2022) is shown as a blue arrow. Insertion of a transgene via RMCE then results in two recombined versions of the original attP and attB sites, which become the two "attR" sites (BATEMAN *et al.* 2006). With RMCE, transgene insertion can occur in either orientation: "Z5", where *lacZ* is closer to 5'P, or "3Z", where *lacZ* is closer to 3'P. In the case of Z5, a modified *homie* is located between 3'P and the *eve-lacZ* reporter gene (direction of transcription is shown as a black arrow). In the case of 3Z, modified *homie* is located between *eve-lacZ* and 5'P, as illustrated. The orientation of *homie* is indicated by the order of its subelements: either "CDEF" for the same orientation in the chromosome as endogenous *homie*, or "FEDC", for the opposite orientation. The eZ-CDEF-eG vector is the same, except for the additional presence of an *eve-GFP* reporter (green arrow), transcribed in the opposite direction as *eve-lacZ*. Each reporter is driven by the *eve* basal promoter, which confers no *eve*-like expression on its own (FUJIOKA *et al.* 2016).

Figure 2. 400 bp of *homie* (CDEF) contains virtually all of its long-range pairing activity.
The previously described 800bp "full-length" *homie* (FUJIOKA *et al.* 2009) was further dissected, by dividing it into roughly 100bp segments (A – H) and testing them in our functional assays. Each tested region is shown as black lines on the map. "λ DNA" is 500bp of phage λ DNA, which serves as a negative control. *lacZ* expression from eZ-vector transgenes (in the Z5 orientation) is shown at embryonic stages 5, 7, 11, and 13 (from top to bottom in each column). In stage 13 embryos (bottom row) with both ABCD and λ DNA, midline cell clusters in the CNS are seen, which are easily distinguished from smaller *eve*-positive neurons, seen in the 1st two columns. Some of these are indicated with arrowheads for ABCD.

Figure 3. Functional dissection of the CDEF fragment of *homie*.
A. *lacZ* expression of eZ-vector transgenes is shown, in either the Z5 (top panel) or 3Z orientation (bottom panel), at embryonic stages 5, 7, 11, and 13 (from top to bottom in each column). In stage 13 CDF embryos, clusters of midline cells are indicated by arrowheads. Note

798 that DEF contains most of the activity of CDEF. **B.** Further dissection of the DEF subfragment.
799 Transgenes in the Z5 orientation are shown, arranged as in A. Note that only DEF and DE
800 contains significant elements of expression in an *eve* pattern, as well as the ability to block
801 expression driven by the *hebe* CNS midline enhancer, as indicated by the absence of large
802 midline clusters of expression (bottom row).
803
Figure 4. A consensus binding site for Su(Hw) and a Cp190-associated site each contribute to
long-range pairing of *homie*.
lacZ expression is shown from eZ-eG transgenes in the Z5 orientation at embryonic stages 5, 7,
11, and 13 (from top to bottom in each column). "ΔSu(Hw)" and "ΔCp190" indicate versions
with the single consensus sites for those proteins point-mutated. Note that mutating the
Su(Hw) site in the context of DEF *homie* causes a severe loss of its LR pairing activity, while
mutating the Cp190 site causes a clear but incomplete loss of LR pairing (see text).
811
Figure 5. The single consensus Su(Hw) binding site in *homie* contributes to its LR pairing
activity.
A, B. smFISH analysis was performed on stage 15 DEF and DEFΔSH dual reporter transgene
embryos. DEF and DEFΔSH dual reporters were inserted into the attP site at −142 kb in two
different orientations. In the Z5 orientation, the *LacZ* reporter is closer to the *eve* TAD, while in
the 3Z orientation, the *GFP* reporter is closer to the *eve* TAD. We then probed for *LacZ* and *GFP*
expression in the anal plate region of the embryo (A). Note that *eve*-like *LacZ* expression is
observed in the anal plate, while *GFP* expression is not detected, and that mutation of the
Su(Hw) recognition sequence in the D sub-element impairs LR pairing activity, resulting in a
substantial drop in *LacZ* expression, quantified in B (P < 0.001).
822
Figure 6. The order of insulator elements is important for LR pairing activity. *lacZ* expression
is shown from eZ-eG transgenes in the Z5 orientation, at embryonic stages 5, 7, 11, and 13
embryos (from top to bottom in each column). In D3EF, the 5' half of D was removed, while in
D3FE, the orientation of section EF was reversed relative to that in D3EF. The orientation of EF
is indicated as blue arrows in the diagram at the top. Note that removing the 5' half of section
D causes a partial loss of LR pairing, although not as much as mutating the Su(Hw) site in the 3'
half of D (Figure 3B), and that reversing the EF region abolishes detectable LR pairing.
830
Figure 7. Enhancer blocking activity of *homie* and its derivatives. Average numbers of midline
clusters visible at stages 12 and 13 in transgenic embryos are shown as box-and-whiskers plots.
Those with a higher number of visible clusters have less enhancer blocking activity. The *homie*
subelements present in the transgene are listed below each pair of plots: names are as in the
text; e.g., "ΔSH" has the Su(Hw) consensus binding site mutated and "ΔCp" has the Cp190-
associated site mutated. Constructs were ordered based on their relative levels of activity in
the enhancer blocking and LR pairing assays. **A:** eZ vector transgenes. **B: Strength of**
enhancer blocking does not always correlate with LR pairing activity: eZ-eG vector
transgenes. Images are reproduced from Figure 4 for comparison. **C: Strength of enhancer**
blocking does not always correlate with PRE blocking activity. Analysis of enhancer blocking
by individual *homie* subelements and their derivatives in eZ vector transgenes shows that only
D, and neither DΔSH, E, EΔCp, nor F has significant enhancer blocking activity (top graph; DE is

reproduced from part A for comparison), while E has stronger PRE blocking activity than does D
(bottom graph, reproduced from Figure 8B for comparison).

Figure 8. Multiple, non-overlapping *homie* subelements are sufficient for PRE blocking
activity. A. Top: map of the *eve* pseudo-locus (FUJIOKA *et al.* 2013). Below: live images of
TER94-GFP in ovaries from transgenic flies carrying the pseudo-locus or modified versions, as
follows: **wt**: *homie* intact; **DEF, D, E, F, AB, C**: ABCDEF *homie* replaced with these
subelements. In each case, λ DNA was used to make their spacing similar to that in intact
homie; **λ DNA**: the same sequence used in Figure 2 replaced ABCDEF. For λ DNA, AB, and C,
longer exposures of the same ovaries are shown at the right. **B.** *GFP* RNA expression in
dissected ovaries, normalized to endogenous *TER94* RNA in the same samples, was quantified
using RT-PCR. Top panel: 3 independent RT-PCR results are shown. Bottom panel: averages
and standard deviations of the same 3 results.

Figure 9. The Su(Hw) binding site, but not the Cp190-associated site, contributes to PRE
blocking. A. Top: map of the PRE blocking assay construct, which consists of the 3' end of the
eve pseudo-locus, starting just upstream of the PRE. Below: live images of *TER94-GFP* in
ovaries from transgenic flies carrying the blocking assay construct or modified versions, as
follows: **wt**, **D, E, F, λ DNA**: same as in Figure 7C. **D-ΔSH, E-ΔCP**: the same point-mutations of
the Su(Hw) and Cp190 binding sites in *homie* used in Figure 7C; **no PRE, no *homie***; the *eve* PRE
and ABCDEF *homie* together were replaced by 1.3 kb of phage λ DNA. For λ DNA and D-ΔSH,
longer exposures of the same ovaries are shown at the right. **B.** *GFP* RNA expression in
dissected ovaries, normalized to endogenous *TER94* RNA in the same samples, was quantified
using RT-PCR. Top panel: 3 independent RT-PCR results are shown. Bottom panel: averages
and standard deviations of the same 3 results.

Figure 10. Relative strengths of enhancer blocking and LR pairing activities do not always
correlate. Constructs were ordered based on their relative levels of activity in the enhancer
blocking and LR pairing assays. Lines connect those (in bold) that clearly differ in their order of
activity in the two assays. Note: those shown in green lettering were tested in the eZ vector,
while those in black lettering were tested in the eZ-eG vector; CDEF, DEF, and DE were tested
in both.

Supplemental Figure legends

Figure S1. A. Expression of *eZ-lacZ* transgenes inserted at −142 kb in *eve*(+) embryonic
mesoderm and CNS. *lacZ* RNA expression in transgenic embryos carrying the indicated *homie*
fragments. Left panel: dorsal view of stage 13 embryos. Arrowheads indicate mesodermal
cells. Right panel: ventral view of stage 13 embryos. Arrowheads indicate *lacZ* expression in EL
(black) and CQ (red) neurons. In ABCD and λ DNA, only *hebe*-like midline expression is seen. **B.**
Mapping of *hebe* enhancers located near the transgene insertion site at −142 kb from
endogenous *eve*. The 1st intron of *hebe* was analyzed for enhancer activity using the eZ vector.
lacZ expression from eZ-vector transgenes carrying fragments from the *hebe* gene: *hebe*-1 (2.3
kb), *hebe*-2 (2.5 kb), and *hebe*-3 (2.8 kb), each inserted at the 74A2 attP site (on chromosome 3,

where there is no interaction with the *eve* locus). Note that the *hebe* CNS midline enhancer is
located in *hebe*-2, and that *hebe*-1 contains an anal plate ring enhancer active only after stage
16.
891
Figure S2. Negative control for transgenes inserted at –142 kb from endogenous *eve*.
893 Reporter gene RNA expression from eZ-eG vector transgenes inserted in the indicated
894 orientations and carrying 500 bp of λ DNA, in embryos at stages 5, 7, 11, and 13 (from top to
895 bottom), used as a negative control for eZ-eG vector transgenes carrying *homie* derivatives.
896
Figure S3. LR pairing activities of *homie* sub-elements and a *gypsy* element. *lacZ* RNA
898 expression from eZ-eG transgenes in the Z5 orientation at embryonic stages 5, 7, 11, and 13
899 (from top to bottom in each column). **CDEF, DEF, DE, ABDE**: map at top shows the analyzed
900 regions of *homie* as black lines. ***gypsy***: a 451 bp *gypsy* transposable element was tested in the
901 same context.
902
Figure S4. *GFP* RNA expression from transgenes used in Figure 4 and 6. Embryonic stages 5, 7,
904 11, and 13 (from top to bottom in each column) are shown. Note that no *eve*-like expression is
905 discernible, and that expression is strong in *hebe* CNS midline cell clusters.
906
Figure S5. Sequences within *TER94* do not contribute to LR pairing by *homie*. Diagram (top):
the green and red arrows show the directions of transcription of *TER94-RA* and non-coding
CR45324 RNAs, respectively. Shown is *lacZ* expression from eZ-eG transgenes in the Z5
orientation at embryonic stages 5, 7, 11, and 13 (from top to bottom in each column). CDEF is
the same as in Figure S3, and D3EF is the same as in Figure 5. In D3EFG5, D3EF is extended to
just downstream of the *TER94* start site; in D3EFGH5, it is extended to just downstream of the
CR45324 start site; in D3EFGH, it is extended through the end of *homie* subregion H. Note that
each of the 3'-extended versions of D3EF have about the same activity, which is noticeably
stronger than that of D3EF, but not quite as strong as that of CDEF, especially in the stage 13
mesoderm.
917
**Figure S6. A. Live images of *TER94-GFP* expression in ovaries carrying pseudo-locus
transgenes.** The ABCDEF region of the pseudo-locus inserted at 95E5 (Fᴜᴊɪᴏᴋᴀ *et al.* 2013) was
replaced with *homie* elements, as follows. **wt**: intact pseudo-locus; **ΔIns**: ABCDEF *homie*
deleted; **CDEF, FEDC, DEF, CDE, CEF, CDF, DE**: these regions of *homie* replaced ABCDEF. In
DEF, CDE, CEF, CDF, and DE, DNA from phage λ was inserted to make the position of the
remaining region similar to that in the intact pseudo-locus. Note that all except ΔIns retain full
PRE blocking activity. **B. *GFP* RNA expression in transgenic ovaries measured using RT-PCR.**
GFP RNA expression was normalized to endogenous *TER94* RNA expression. **Top panel**: 3
independent RT-PCR results are graphed; the 4[th] data set is a re-PCR analysis of the cDNAs of
the 1[st] set. **Bottom panel**: averages and standard deviations of the same 4 data sets are
shown. Labeling is the same as in A, except that "Δhomie" is used in place of "ΔIns".
929
Figure S7. DNA and primer sequences. See the main text and figure legends for how these
931 sequences were used.
932

933 **Figure 1**

Figure 2

939 **Figure 3**

Figure 3 (cont.)

944
945

Figure 4

Figure 5

 Figure 6

955 **Figure 7**

957   **Figure 7 (cont.)**

D3FE enhancer blocking is a bit better,
while LR pairing by D3EF is clearly better

958
959

Figure 7 (cont. 2)

enhancer blocking:
D > E

PRE blocking:
E > D

961
962

963 **Figure 8**

 Figure 8 (cont.)

969 **Figure 9**

970
971

Figure 9 (cont.)

Figure 10

 Supplemental Figure 1

A

Pericardial cells and muscles

CNS

B *hebe – RD transcript*

hebe-1; 2R:9829859 – 9832120
hebe-2; 2R:9831531 – 9834080
hebe-3; 2R:9833624 – 9836462

979 **Supplemental Figure 2**

eZ-λ DNA-eG

982 **Supplemental Figure 3**

985 **Supplemental Figure 4**

vector: eZ- -eG

DEF	DEF-ΔSu(Hw)	D3EF	D3FE

Supplemental Figure 5

991 **Supplemental Figure 6**

 Supplemental Figure 6 (cont.)

Supplemental Figure 7

CDEF *homie* sequence

AATACTAAAAAGTTTTTACTAGCATAAGCTGCGATTGAGCAGTTATTGTAGGGACAGGTTGATGGCCGA
GTGTTCTAGGGAAATGAATGAATGAAG**ATTTTTTCTTACTACCAAAACTTCTACAACTAAGTGAAATAG**
ACGGATTGAGTTCTACATACTTCCTACATATATTTTAGATATAGACATAGCCGAAAAGTATGCTGCAC*T*
*TTTT**CAATAAAAACGTAAGCAGCTAAGCAGCGTAAGGCAGCTTCATAGCAGCGATCATCTGCCAGCGAG*
*CAT**TAGCCAAGTTAG*GCACC<u>TGCACACATTGCCACTCTTTCTTTTAGCGTTGCCACTTCAATTTCTTTTACAA</u>
<u>ACCATCGCAGCGTGTAATC</u>

C: plain text
D: **bold**
E: *italics*
F: <u>underlined</u>
Red: Su(Hw) binding site
Green: Cp190-associated site

<u>Su(Hw) consensus binding site (on complementary strand)</u>
```
Searched:            YWGCATACTTTT
Found:               CAGCATACTTTT  (red = matches)

Searched:    NWWWWNYRTWGCATACTTTTKGSDB
Found:       AAAAAGTGCAGCATACTTTTCGGCT  (red = matches)
Mutated:     AAAAAGAACAGGATCCGAACATCCT
```

<u>Cp190-associated site (complementary strand)</u>
```
Searched:    GGTTBDWRWMYYNGCTD
Found:       GGTGCCTAACTTGGCTA   (green = matches)
Mutated:     GTTGCAAAGCTTGTCGA
```

D3EFGH sequence

ATATAGACATAGCCGAAAAGTATGCTGCACTTTTTCAATAAAAACGTAAGCAGCTAAGCAGCGTAAGGCA
GCTTCATAGCAGCGATCATCTGCCAGCGAGCATAGCCAAGTTAGGCACCTGCACACATTGCCACTCTTTC
TTTTAGCGTTGCCACTTCAATTTCTTTTACAAACCATCGCAGCGTGTAATC<u>GATATTTTCTAAGCCC</u>
<u>AATCGATGAATGTCCGGTAACGCACATC</u>**ACCAATCTGTTTCTGCTGCCTCATCTTTGAA**
TTTGACTAAAAGTGCCGATTTCCAGCTCAACGAATAGTCCAAGTCGCGCCGAATTGTAC
AATTAAGTTAAAAAGTTCAACGTCAACTAGTTTATCATGGCAGATTCCAAGGGGTAAGT
GAACTGGACGCCGAAATCTAGAGGCAGCAGGAGAAAT

D3EF: smaller, green font
<u>D3EFG5</u>: D3EF was extended by the <u>underlined</u> sequence
D3EFGH5: D3EFG5 was extended by the **bold** sequence

1042   D3EFGH: D3EFGH5 was extended to the end of the sequence

1043

1044

1045   **Primers used for RT-PCR**

1046   *GFP*

1047   GGGCACAAGCTGGAGTACAACTACAA

1048   TGGCGGATCTTGAAGTTCACCTTG

1049

1050   *TER94*

1051   TGAAGCCACCGCGTGGTATTCTTA

1052   TTTGGACATGATCTCCGGTCCGTT