TITLE

Selective epithelial expression of *KRAS*[G12D] drives ductal proliferation and desmoplasia in the Oncopig pancreas

Carlos P Jara[1], Al-Murtadha Al-Gahmi[1], Audrey J. Lazenby[2], Michael A. Hollingsworth[3], and Mark A. Carlson[1,4]

[1]Department of Surgery, University of Nebraska Medical Center, Omaha, NE, USA

[2]Department of Pathology, University of Nebraska Medical Center, Omaha, Nebraska, USA

[3]Eppley Institute for Research in Cancer and Allied Diseases, Fred and Pamela Buffett Cancer Center, University of Nebraska Medical Center, Omaha, NE, USA

[4]Department of Genetics, Cell Biology and Anatomy, University of Nebraska Medical Center, Omaha, NE USA

Carlos P. Jara (ORCID: 0000-0001-5444-721X)

Al-Murtadha Al-Gahmi (ORCID: 0009-0000-5484-467X)

Audrey J. Lazenby (NA)

Michael A. Hollingsworth (ORCID: 0000-0002-5255-8888)

Mark A. Carlson (ORCID: 0000-0002-8113-9199)

Corresponding author:

Mark A. Carlson
E-mail: macarlso@unmc.edu

ABSTRACT

Background. Pancreatic ductal adenocarcinoma (PDAC) remains a formidable challenge in oncology, characterized by a high mortality rate, largely attributable to delayed diagnosis and the intricacies of its tumor microenvironment. Innovations in modeling pancreatic epithelial transformation provide valuable insights into the pathogenesis and potential therapeutic strategies for PDAC.

Methods. We employed a porcine (Oncopig) model, utilizing the Ad-K8-Cre adenoviral vector, to investigate the effects of variable doses (10^7 to 10^{10} pfu) on pancreatic epithelial cells. This vector, the expression from which being driven by a Keratin-8 promoter, will deliver Cre-recombinase specifically to epithelial cells. Intraductal pancreatic injections in transgenic Oncopigs (*LSL-KRAS*[G12D]-*TP53*[R167H]) were performed with histologically-based evaluation at 2 months post-injection.

Results. Specificity of the adenoviral vector was validated through Keratin-8 expression and Cre-recombinase activity. We confirmed that the Ad-K8-Cre adenoviral vector predominantly targets ductal epithelial cells lining both large and small pancreatic ducts, as evidenced by Keratin 8 and CAM5.2 staining. Higher doses resulted in significant tissue morphology changes, including atrophy and enlarged lymph nodes. Microscopic examination revealed concentration-dependent proliferation of the ductal epithelium, cellular atypia, metaplasia, and stromal alterations. Transgene expression was confirmed with immunohistochemistry. Desmoplastic responses were evident through vimentin, α-SMA, and Masson's trichrome staining, indicating progressive collagen deposition, particularly at the higher vector doses.

Conclusion. Our study suggests a distinct dose-response relationship of Ad-K8-Cre in inducing pancreatic epithelial proliferation and possible neoplasia in an Oncopig model. All doses of the vector induced epithelial proliferation; the higher doses also produced stromal alterations, metaplasia, and possible neoplastic transformation. These findings highlight the potential for site-specific activation of oncogenes in large animal models of epithelial tumors, with the ability to induce stromal alterations reminiscent of human PDAC.

Keywords: PDAC, Ad-K8-Cre, Oncopigs, neoplasia, desmoplastic reaction, KRAS[G12D].

INTRODUCTION

Pancreatic ductal adenocarcinoma (PDAC) remains a challenge in public health, responsible for 7% of all cancer-related deaths. Five-year survival rate is approximately 9%, reflecting the disease's aggressive nature and modestly effective treatment options [1-3]. The use of large animal models, particularly porcine models, may provide a supplemental and unique approach to PDAC research, especially with respect to porcine physiological and anatomical similarities to humans [4]. Porcine size enables the replication of human interventional procedures which are challenging to perform in rodent models. Furthermore, the porcine immune system, lifespan, and genetic sequence mirrors those of humans better than mice do [4, 5].

In a previous study [6], we employed the transgenic Oncopig model (somatic *LSL-KRAS*^{G12D}*-TP53*^{R167H}) [7] to describe and characterize a porcine model of pancreatic cancer (proof-of principle for such a model was published earlier) [8, 9]. The KRAS^{G12D} mutation is a near-ubiquitous initiator and driver in PDAC [10-15]. In the Oncopig we used pancreatic ductal injection of an adenovirus that expressed Cre-Recombinase under the control of CMV promoter, allowing high Cre expression in almost any cell type, resulting in expression of the two mutants (KRAS^{G12D} and p53^{R167H}) in infected cells. The outcome was fulminant tumor growth which led to subject demise in 2-3 weeks; tumor pathology was mixed with an epithelial component [6].

In order to have a less fulminant, more tractable porcine pancreatic cancer model which also would be fully epithelial in origin, we utilized in the present study an adenovirus which would express Cre-Recombinase under control of the Keratin 8 promoter, a vector whose Cre expression would be restricted to epithelium [16]. We believed that the degree of transformation observed in response to varied adenoviral vector doses might provide an improved porcine model of pancreatic cancer that could be used for device development, testing of antitumor drugs, or even study of early tumor pathogenesis and tissue response.

RESULTS

Keratin-8 And Cre-Recombinase Expression In Pancreatic Cells. First, we explored Keratin-8 expression in pancreatic epithelial cells by using single-cell RNA sequencing from whole human pancreas. We successfully identified higher levels of Keratin-8 RNA expression mainly in pancreatic ductal cells and lower in acinar cells, both in human healthy and PDAC samples (Fig. 1a). Next, we explored Keratin 8 protein expression in the porcine pancreas using immunohistochemistry (IHC). . We identified epithelial staining (CAM5.2 staining and Keratin 8) in accessory and main ducts in both wild type and Oncopig pancreas (Fig. 1b). Despite human scRNA sequencing results indicating some K8 expression in acinar cells, our analysis in porcine pancreas showed Keratin-8 staining was restricted primarily to ductal structures, with no detectable expression in acinar cells. . Finally, we performed an e*x vivo* assessment of the Ad-K8-Cre Adenovirus (a vector construct with Keratin 8 promoter driving the expression of the Cre-Recombinase) in wildtype porcine pancreas (Fig. 1c, Supplementary S1a). This *ex vivo* infection assay showed that the expression of Cre-recombinase under the control of the Keratin 8 promoter was limited to the epithelial cells, confirming the selective activity of the adenoviral construct in pancreatic epithelial cells (Fig. 1d).

Dose-Dependent Macroscopic Alteration After Vector Administration. After confirming targeted expression of Cre-recombinase by Ad-K8-Cre in pancreatic epithelial cells, we evaluated for targeted epithelial activity of the Ad-K8-Cre adenovirus *in vivo* (Supplementary S1b). We injected Ad-K8-Cre adenovirus into the duct of the connecting lobe of the pancreas, as previously described [6], at different concentrations (ranging from 10^7 to 10^{10} pfu) in eight transgenic Oncopigs (Supplementary S2a-c). For the purposes of this report, the injection region, or the portion of pancreas subjected to the vector treatment, is meant to be synonymous to with the region of the connecting lobe of the pancreas (dashed oval in Fig. 2a,b). Technically speaking, the injection site would be the open end of the primary duct to the connecting lobe where the cannula was inserted (Supplementary Fig. S2). After the treatment was injected, however, it was presumed to diffuse into the regional pancreas (i.e., proximal connecting lobe) via the branching ductal system. Therefore, it is probably more accurate to refer to the treated pancreas as the "injection region" as opposed to "injection site".

After 2 months post-injection, all Oncopigs survived (Table 1). Oncopigs injected with 10^9 to 10^{10} pfu of vector exhibited signs of pancreatic atrophy in the connecting lobe (i.e., the injection region) with small (<5 mm) clustered lumps (nodularity) and with enlarged (>1 cm), firm lymph nodes around the pancreas (Fig. 2a,c-da-d). On the other hand, Oncopigs injected with vector doses ranging from 10^7 to 10^8 pfu showed conserved gross morphology of the pancreas (Fig. 2b). At macroscopic level, the injection region in Oncopigs treated with 10^7 to 10^8 pfu showed neither tangible damage nor change. While there were lymph nodes present around the pancreas as well, there was an absence of discernible masses or marked discoloration to suggest atypical growth or inflammation with the lower dose Oncopigs (Fig. 2b,c). All subjects gained weight appropriately during the eight-week observation, and there were no differences in weight at any time point among the four dosage groups (Table 1). Blood testing (hemoglobin, hematocrit, amylase, lipase, carcinoembryonic antigen, CA 19-9, and alpha-fetoprotein) prior to induction vs. prior to necropsy revealed no differences in any group (Table 2).

Microscopic Morphology. Two months after Ad-K8-Cre delivery, histological analysis of pancreas from the proximal connecting lobe, i.e., the region of injection (Fig. 3), showed architectural disruption and cellular atypia for all vector doses tested, whereas pancreatic samples remote from the injection site (i.e., located more than 10 cm away from the connecting lobe) exhibited normal pancreatic architecture. In this study,

each Oncopig served as its own control. That is, the pancreas in these swine was large enough so that a biopsy of normal appearing pancreas more than 10 cm away from the vector injection region (i.e., the proximal connecting lobe) was possible in each subject. So, samples of treated *versus* untreated pancreas could be acquired from the same pig. The above histologic changes were accentuated in the 10^9 and 10^{10} vector dose (Fig. 3a). In addition, the following histologic characteristics were observed within the injection region:

i.   *Acinar Cell Loss and Ductal Proliferation.* While the Ad-K8-Cre specifically drove expression in ductal cells, we observed loss of acinar morphology within the injection region, along with a significant increase in ductal cell proliferation (Fig. 4a-c, Supplementary S3a-d).
ii. *Intestinal metaplasia in the large ducts*. In some regions of the larger pancreatic ducts, the flat ductal epithelium showed the appearance of goblet cells (Supplementary S4a)
iii. *Inflammatory response*. At the two-month time point, the inflammatory response was assessed by identifying CD45-positive leukocytes, CD3-positive T lymphocytes, and MPO-positive neutrophils. We identified increased T lymphocytes with lower neutrophils infiltration (Fig. 5a-d).
iv. *Healing vs. regeneration vs. neoplasia*. Based on the H&E morphology and the immune cell staining, it was not possible to determine whether the above histologic changes were representative of a response to injury that would generate a scar (i.e., wound healing), or if a regenerative response would ensue, or if there was microscopic neoplasia present which would produce a tumor.

Proliferation Within The Injection Region. Ductal structures within the injection region were populated by Ki67 and PCNA-positive cells, which was evident at all Ad-K8-Cre concentrations tested (Fig. 4a-d). While the 10^7 to 10^9 pfu concentrations displayed a ductal proliferation pattern, the 10^{10} pfu showed a loss of distinct circular or ovoid proliferation areas, with stronger, more dispersed, and homogenous staining pattern throughout the sample. Pancreas remote from the injection region maintained normal microscopic morphology, including uniform tissue structures and limited Ki67 staining (Fig. 4a-b).

Targeted KRAS^{G12D} Expression Within Epithelium. After 2 months post Ad-K8-Cre injection, we observed a targeted induction of the KRAS^{G12D} expression within the epithelium (primarily in the ductal cells) within the injection region (Fig. 4c). These staining patterns were particularly evident in the areas with disrupted architecture and complex ductal proliferation. This staining confirms that the Ad-K8-Cre delivery induced KRAS^{G12D} expression within the epithelial cells. Interestingly, regions that tested positive for the KRAS^{G12D} mutation also displayed a corresponding rise in Ki-67 expression. We confirmed that proliferation is associated with cells expressing KRAS^{G12D} by performing double staining for PCNA and KRASG12D, as shown in Figure 4c. This underscores the relative quiescence of healthy pancreas compared to the aggressive growth pattern observed in the injected region.

Cytokeratin expression In Transformed, Proliferative Regions Suggest Epithelial Origin. We utilized epithelial markers, (Pankeratin, CAM5.2 and Cytokeratin 7) to confirm the epithelial nature of the transformed pancreatic cells. Pankeratin (Fig. 4b), CAM5.2 (Supplementary S3b), and Cytokeratin 7 (Supplementary S3c) stained strongly in cells corresponding to those transformed by KRAS within the injection region (Supplementary S3a) . We observed the highest staining intensity in samples from Oncopigs injected within 10^9 and 10^{10} concentrations, highlighting the effectiveness at elevated doses and suggesting a dose-response relationship. The epithelial markers were reduced in pancreas remote from

the injection, emphasizing the targeted expression of Cre recombinase, and suggesting that the KRAS^{G12D}-expressing cells maintained epithelial lineage post-transformation.

Desmoplasia, Matrix Deposition, and Mucin Production. Histologic features of stromal reaction to Ad-K8-Cre transfection were characterized with vimentin, α-SMA, and Masson's trichrome staining (Fig. 6a-c). Increased collagen deposition consistent with desmoplasia was found within the injection region for all concentrations (Fig. 6a). In particular, there was a substantial fibrotic and desmoplastic stromal reaction at the 10^9 and 10^{10} concentrations (blue staining in Fig. 6a). α-SMA (smooth muscle actin) immunohistochemistry demonstrated a vector dose-dependent increase in staining (Fig. 6b), also consistent with a stromal reaction within the injection region. Conversely, α-SMA staining was minimal within pancreas remote from the injection region (Fig. 6b). In addition, there was increased vimentin staining within the stroma surrounding proliferative ductal cells within the injection region, also consistent with desmoplasia (Fig. 6c). The 10^9 to 10^{10} vector doses had the most intense and widespread vimentin staining; pancreas remote from the injection region had minimal vimentin signal. There also was an increase in acidic mucin staining in the ductal epithelium and stroma within the injection region (purple-colored staining in Fig. 7), with an apparent dose-dependency on vector dose. In contrast to the injection region, remote healthy pancreas displayed minimal staining.

Comparative Human Microscopy. Examination of four human pancreatic cancer samples (Supplementary Fig. S5) demonstrated considerable architectural disruption, indicative of advanced disease. Human pancreatic cancer tissue also was characterized by a desmoplastic reaction, with increased collagen and vimentin deposition. These changes were comparable to the those observed with vector induction in the Oncopig model, particularly at the higher vector doses.

DISCUSSION

Intraductal Injection For Vector Delivery

The unique, tri-lobed anatomy of the porcine pancreas allows for isolation and injection of a large pancreatic duct (within the connecting lobe) without major physiologic impact to the subject [6]. This anatomic arrangement, along with the large subject size, allows for unique opportunities to study pancreatic neoplasia in swine. For example, the swine's unique anatomy allows for precise delivery of the adenoviral vector, Ad-K8-Cre. Intraductal administration of the vector ensures that the adenovirus will bathe the pancreatic epithelium, maximizing the potential for targeted gene activation. These theoretical advantages were borne out by the results of this study.

Endoscopic Retrograde Cholangiopancreatography (ERCP) offers advantages for pancreatic intraductal injections compared to traditional surgical methods. Its minimally invasive nature reduces the need for large incisions, leading to shorter recovery times and fewer postoperative complications [17][18-20]. Additionally, ERCP offers real-time imaging allowing precise targeting of the pancreatic ducts [21, 22] . This endoscopic approach is less invasive, but often presents challenges in terms of investigator accessibility (i.e., equipment and operator expertise requirements). It is not clear whether the duct to the connecting lobe can be reliably cannulated via ERCP. With the open surgical approach, our success rate in cannulation of the duct to the connecting lobe was 100%. Our laparotomy-based approach, involving dissection and direct visualization of the pancreatic duct, requires no special equipment other than what typically can be found in the typical Comparative Medicine Department, and can be taught to individuals who has veterinary operative experience.

Our ex vivo experiments using wild-type (WT) porcine pancreas were conducted to confirm that the Keratin 8 promoter-driven adenovirus is capable of Cre-Recombinase expression which is restricted to epithelial cells. To our knowledge, this was the first time this adenovirus was used in swine and in the pancreas, so the experiment aimed to verify the targeted expression of Cre-Recombinase.

Dose Dependence

At a lower vector concentration (10^7 pfu of virus per injection), Ad-K8-Cre promoted epithelial proliferation while preserving ductal structural integrity. This suggests that the adenoviral vector can precisely target the epithelial cells without disrupting the ductal architecture with only a mild desmoplastic response. In contrast, the highest vector concentration tested (10^{10}) produced a marked increase in disorganized epithelial proliferation (Ki67 staining), and a robust desmoplastic reaction characterized by extensive collagen deposition and α-SMA expression. This suggests that at elevated concentrations, Ad-K8-Cre not only impacted epithelial cells but also triggered a desmoplastic/fibrotic response that could potentially modify the tumor microenvironment. This dose-dependent response ultimately may prove useful for diverse research efforts on pancreatic cancer.

Intestinal metaplasia and Goblet Cells

Goblet cells are specialized epithelial cells found primarily in the mucosal lining of the intestines and respiratory tract, where they play a crucial role in secreting mucus to protect and lubricate the mucous membranes [23]. Intestinal metaplasia occurs when epithelial cells in one area of the body transform into

intestinal-type epithelial cells, typically as a response to chronic irritation or inflammation. The prototypical example of intestinal metaplasia is Barrett's esophagus, in which the stratified squamous epithelium of the esophagus transdifferentiates into columnar-type intestinal epithelium, which is associated with an increased risk for esophageal carcinoma [24]. Pancreatic ducts do not naturally contain goblet cells; their appearance in our study is consistent with intestinal metaplasia. Similar to ADM, this type of metaplasia can be triggered by chronic inflammation or injury. However, as in Barrett's esophagus, the appearance of goblet cells in the pancreatic ducts could signify an increased risk for epithelial neoplasia.

Large animal modeling of pancreatic cancer

The targeted expression of the KRAS^{G12D} mutant to pancreatic ductal epithelium within the Oncopig model underscores the potential of large animal models to advance PDAC research. Porcine pancreatic cancer models could provide a realistic and clinically-relevant platform for preclinical assessment of novel therapeutics and diagnostics, and potentially may permit mechanistic studies of early tumor progression [6, 9]. In contrast to a recent investigation on a porcine model of pancreatic cancer which used a nonspecific adenoviral vector (Ad5CMVCre-eGFP,) in the Oncopig and reported fulminant tumor growth, short survival times, and mixed tumor pathology [6], the present study on the use of the Ad-K8-Cre induction vector demonstrated enhanced specificity for pancreatic epithelial cells, which likely produced the 100% 2-mo survival. The implication of this result is that neoplasia induced by the K8-driven vector will be a slower growing tumor model which could present opportunities for early diagnostic and interventional technologies, along with mechanistic studies of tumor progression.

The genetic and physiological similarities between pigs and humans could better enable the study of tumor progression and therapeutic responses in a setting that is far more comparable to human biology than murine models. These porcine models allow researchers to explore complex multimodal treatments, including surgery, chemotherapy, and localized therapies, which are more difficult to evaluate in smaller animals like mice [25]. Furthermore, the size of the pig allows for the use of human-sized medical equipment, facilitating translational research in diagnostic and therapeutic device development [4, 9]. Unlike murine models, which often fail to replicate the full scope of human immune responses and tumor microenvironments, porcine models provide a robust platform to study interactions between cancer cells and the immune system, a critical component in the development of immunotherapies [4, 25].

The question of whether true neoplasia/ carcinoma was produced in this study is still open. Careful and independent pathologic analysis was inconclusive on this question. There was obvious evidence of mutant KRAS expression along with increased proliferation and desmoplasia typical of PDAC. Perhaps an early *in situ* transformative process was being observed; however, we cannot be conclusive with the present data. A longer observation period than the two months used in the present report may be needed to resolve this question.

Another open question from this study is the role of the host immune response to injection of the vector into the pancreas. It was demonstrated in previous work [6] that the vector itself has minimal pathologic effect; it was only when the vector expressed Cre recombinase (driven by the CMV promoter) that there was strong evidence of transformation accompanied by an overwhelming immune response. The immune response to epithelial-restricted Cre recombinase expression in this report was qualitatively less than in the prior study of unrestricted Cre recombinase expression [6] but a T-cell response was still present in the

current report. Perhaps the Oncopig's T-cell response to nascent ductal transformation will provide an opportunity for mechanistic studies on host immune response to PDAC.

CONCLUSION

Our study demonstrated that intraductal administration of the adenoviral vector Ad-K8-Cre had a qualitative dose-dependent effect on epithelial cell proliferation and stromal reaction within the injection region; the latter effect was reminiscent of human PDAC. These histologic findings were related to expression of the KRAS^{G12D} mutation. While we have no incontrovertible evidence for neoplastic transformation in the present study, we believe that future work will demonstrate that the histologic events noted in the present study were indicative of early stage PDAC.

DECLARATION

The datasets used and/or analyzed during the current study available from the corresponding author on reasonable request.

MATERIALS AND METHODS

Animal Welfare. The animals utilized for this report were maintained and treated in accordance with the *Guide for the Care and Use of Laboratory Animals* (8th ed.) and in accordance with the Animal Welfare Act of the United States (U.S. Code 7, Sections 2131 – 2159). The animal protocol pertaining to this manuscript was approved by the Institutional Animal Care and Use Committee (IACUC) of the University of Nebraska Medical Center (ID number 19-053-FC). All procedures were performed in animal facilities approved by the Association for Assessment and Accreditation of Laboratory Animal Care International (AAALAC; www.aaalac.org) and by the Office of Laboratory Animal Welfare of the Public Health Service (grants.nih.gov/grants/olaw/olaw.htm). All surgical procedures were performed under isoflurane anesthesia, and all efforts were made to minimize suffering. Euthanasia was performed in accordance with the AVMA Guidelines. Design and conduct of animal experiments was guided by the Animal Research: Reporting of In Vivo Experiments (ARRIVE) standards.

Porcine Subjects. Transgenic Oncopigs (Oncopig Model, or OCM; LSL-*KRAS*^G12D-IRES-*TP53*^R167H, Supplementary S2c). were purchased from Sus Clinicals (susclinicals.com). The OCM subjects were a hybrid of Minnesota minipigs and domestic pigs. The genotype of each porcine subject was confirmed with PCR upon subject delivery. Swine were housed two littermates per pen, except for one week of individual housing post-laparotomy (but with contact through the pen grating), in order to prevent wound cannibalism. Swine were fed ad lib with standard hog feed (Purina Nature's Match® Sow and Pig Complete Feed; www.purinamills.com). The basic experimental design included a ≥1 week acclimatization period after subject delivery to the research facility. Each subject then underwent an induction procedure (laparotomy under general anesthesia; one major survival procedure), followed by observation for 2 months.

Overview. The technique of vector administration into the Oncopig pancreas is based on a previous description [6]. Briefly, our laparotomy procedure initiated with a ventral midline incision, starting just inferior to the xiphoid process, and extending to the umbilicus in females and approaching the urethral meatus in males. After retracting the small and large bowel caudally, exposing the duodenal segment of the pancreas. The connecting lobe of the pancreas was transected just below its intersection with the duodenal lobe. Using surgical loupes (3.5x magnification), the duct within the connecting lobe was cannulated using a 22g Angiocath. The specified concentration of Ad-K8-Cre (from 10^7 to 10^{10} pfu) in 100-150 µL was injected into this duct , which was immediately ligated with 3-0 silk after the injection in order to prevent back flow of the injectate out of the duct. The pancreas remote (>10 cm) from the injection site was used as a same-subject control, providing a baseline for comparison with the treated regions.

Operative Details. All Oncopigs underwent tumor induction in the same fashion. The Oncopigs were pre-anesthetized using an intramuscular injection of Telazol®-Ketamine-Xylazine (2/4/2 mg/kg, respectively) placed supine on the procedure table, and then intubated with 6-7 French endotracheal tube. Anesthesia was maintained with 1% Isoflurane inhalation. Vitals were monitored noninvasively and maintenance fluid, (Lactated ringers' solution) was given throughout the case (200-300 mL total).

The skin was shaved and prepped with chlorohexidine in alcohol, weight-based bupivacaine was injected subcutaneously around the planned upper abdominal midline, and abdomen was draped in sterile fashion. An incision was made below the xiphoid extending down past umbilicus in females and proximal to the urethral meatus in males with #10 blade. This was then carried down through subcutaneous fat,

facia and muscle to the pre-peritoneal space using electrocautery. The peritoneum was carefully incised to enter the greater sac and Bookwalter retractor was assembled and used to provide adequate abdominal exposure.

The small and large bowel were retracted caudally and the stomach was grasped at the pylorus for reflection antero-superiorly to expose the duodenal lobe of the pancreas. Pancreatic adhesions to the small/large bowels were cut and a window between the connecting lobe of the pancreas and pancreatic head (duodenal lobe) was created. After the proximal connecting lobe of the pancreas had been dissected circumferentially, the proximal end of the connecting lobe of pancreas was tied off with 3-0 silk flush with the duodenal lobe.

The connecting lobe then was transected 0.5-1 cm distally from the silk ligature. With the aid of surgical loupe magnification (3.5x) the pancreatic duct in the distal connecting lobe was identified and cannulated with 22 g Angiocath. The injectate with the desired pfu of Ad-K8-Cre was injected into the pancreatic duct, the catheter was withdrawn, and the distal connecting lobe of pancreas was immediately ligated with 3-0 silk suture.

The peritoneum was reapproximated with 3-0 Vicryl suture; the midline abdominal fascia was closed with 0-Maxon suture (5 mm tissue bite, 2-3 mm stitch interval); the deep dermal layer was closed with 4-0 Vicryl suture; and a subcuticular closure with 4-0 Vicryl suture on a cutting needle. Dermabond adhesive glue was then placed over the incision. The Oncopigs were extubated after confirmation of spontaneous breathing and continued to be monitored until oxygen saturations remained greater than > 90 on room air.

Postoperative Care and Observation Period. Oncopigs were housed in individual pens until post-op day 4 to prevent wound cannibalism, after which they were regrouped with their respective sex. All animals were monitored carefully for 3 days and were kept NPO until postoperative day 2 once, at which time feeding was reinitiated if the subject was clinically stable. The swine continued to be assessed daily for signs of deterioration until term. Criteria for early removal from the study and euthanasia were symptoms of failure to thrive (anorexia, lethargy, decreased movement, abnormal breathing, or other signs of distress) or sepsis (fever, wound disruption, or drainage)

Euthanasia and Necropsy. At 2 months post AdCre induction, the Oncopigs underwent necropsy. Anesthesia and access to the abdomen was obtained in a fashion similar to the above, taking care to lyse adhesions and avoid bowel injury. At the time of euthanasia, inhalational isoflurane was increased to 5% via the ventilator. The prior midline incision was reopened. Inferiorly this incision was extended in paramedian fashion to avoid midline structures (such as the urethra in males). The completed necropsy incision extended from xiphoid process to the pelvic inlet. The bilateral thoracic cavity then was entered by transversely incising the diaphragm just inferior to the xiphoid process. The intrathoracic portion of the inferior vena cava was easily identified as it emerged from the liver in the posterior mediastinum. Phlebotomy was performed from the cava, and then Fatal-Plus® (pentobarbital sodium, 390 mg/mL; 1 mL per 4.5 kg body weight) was administered by caval injection. Two minutes after administration of Fatal-Plus®, the inferior vena cava was transected just above the diaphragm to exsanguinate the subject. A gross necropsy involving the lungs, heart, liver, kidneys, pancreas, intestines, bladder, and associated peritoneal surfaces then was performed. The pancreas and surrounding tissue then underwent gross evaluation and circumferential excision with tissue specimens collected from the injection region (connecting lobe), non-injection region (tail of pancreas), and lymph nodes for histopathologic analysis.

Ex Vivo Pancreatic Duct Procedures. For the *Ex Vivo* tissue culture, pancreatic samples were processed as previously described with some modifications [26, 27]. Briefly, porcine pancreases were harvested from wild type domestic pigs. The main pancreatic duct was carefully isolated from the harvested pancreas. The isolated pancreatic main duct was meticulously chopped into small pieces. The dissected pancreatic duct pieces were placed in a 6-well plate. Each well was filled with DMEM supplemented with 10% FBS (tetracycline-free). A dose of 10^9 pfu of Ad-K8-Cre was added to each well. The plate was incubated overnight in a 5% CO2 incubator. Following incubation, the samples were fixed using 4% paraformaldehyde overnight. The fixed samples were then embedded in paraffin and sectioned at 5µm thickness. The sections were incubated overnight with an anti-recombinase antibody. Post incubation, the sections were stained using DAB to visualize the recombinase expression.

Histology and Immunohistochemistry. For the histological analyses, pancreatic samples were processed and stained with hematoxylin and eosin (Hematoxylin Select Reserve Eosin Multichrome) as previously described [28]. Briefly, Pancreatic tissues were harvested from porcine subjects. Two distinct samples were obtained from each subject: Injection region (Connective lobe where AdCre was administered) and Non-injection region (pancreas distant from the injection site, particularly the tail or splenic lobe of the pancreas). Harvested samples were immediately fixed in cold 4% paraformaldehyde overnight. Post-fixation, the tissues were embedded in paraffin and serially sectioned at 5µm thickness. Three serial sections from both regions (injection and non-injection) per animal were prepared for IHC and histological analysis. We incubated sections with hematoxylin for 30 s, rinsed in distilled water, incubated for 30 s with eosin, rinsed again in distilled water, and dehydrated. The tissue sections were incubated overnight with the following primary antibodies: α-SMA (1:1000, Abcam ab5694), Anti-cytokeratin 8 (Invitrogen MA1-19037, 1:500) , Anti-CAM5.2 (1:500, Roche REF#790-4555) Anti-KRAS G12D (Genetex GTX132407 1:200), Anti-Ki67 (ab16667, 1:200), and Anti-vimentin (Roche REF#790-2917, 1:500). After incubation, the sections were treated with DAB to detect protein expression. For Masson's Trichrome Staining (Masson Trichrome Stain Kit, #KTMTR EA), we incubated Weigert's Iron Hematoxylin for 5 min, Biebrich Scarlet Acid Fuchsin solution for 15 min, differentiate in the phosphomolybdic acid solution for 15 min, Aniline Blue solution for 10 min and acetic acid solution for 5 min. Finally, we mounted the slices on Entellan®, digitally captured the images under bright field microscopy [29]. For the mucins and polysaccharides analyses, pancreatic samples were processed and stained with Alcian Blue-Periodic Acid Schiff (PAS, Alcian Blue PAS Stain Kit, Liter #KTAPALT EA) as previously described [30]. Briefly, Alcian Blue solution of pH 2,5 was made by dissolving Alcian Blue in distilled water and adding glacial acetic acid. The neutral red stain was made by dissolving 1 g of neutral red in 100 mL of distilled water and adding 0,1 ml of glacial acetic acid. In short, slides were deparaffinised and rehydrated, stained for 15 min in the Alcian Blue solution, rinsed for 5 min in tap water, counterstained with the Neutral Red solution for 1 min and then dehydrated, cleared, and mounted. All staining procedures were carried out in accordance with manufacturer's recommendations.

Pancreatic Transcriptomic analysis

For the single cell RNA sequencing analysis, publicly available healthy single cell samples [31, 32] were processed as previously described [33, 34] with some modifications. Briefly, cells with library size < 1000 UMI or few expressed genes were excluded from analysis and processed into the Seurat R package (v4.0). The Seurat objects and pipeline were used for downstream analyses and visualization: NormalizeData, ScaleData were used to calculate the comparable expression values; FindVariableFeatures were used to include the variable genes that contribute to the overall similarity/variability of cellular transcriptomic profiles; RunPCA, FindNeighbors, FindClusters, and RunUMAP were used to calculate the dimension-

reduction coordinates for visualization and to perform unsupervised clustering. In the downstream analyses, we used Uniform Manifold Approximation and Projection (UMAP) coordinates to visualize the layout of the cells [31].

REFERENCES

1.   Yu Y, Yang G, Huang H, Fu Z, Cao Z, Zheng L, et al. Preclinical models of pancreatic ductal adenocarcinoma: challenges and opportunities in the era of precision medicine. Journal of Experimental & Clinical Cancer Research. 2021;40(1):8.

2.   Sarantis P, Koustas E, Papadimitropoulou A, Papavassiliou AG, Karamouzis MV. Pancreatic ductal adenocarcinoma: Treatment hurdles, tumor microenvironment and immunotherapy. World J Gastrointest Oncol. 2020;12(2):173-81.

3.   Orth M, Metzger P, Gerum S, Mayerle J, Schneider G, Belka C, et al. Pancreatic ductal adenocarcinoma: biological hallmarks, current status, and future perspectives of combined modality treatment approaches. Radiation Oncology. 2019;14(1):141.

4.   Bailey KL, Carlson MA. Porcine Models of Pancreatic Cancer. Front Oncol. 2019;9:144.

5.   Jagadesan S, Mondal P, Carlson MA, Guda C. Evaluation of Five Mammalian Models for Human Disease Research Using Genomic and Bioinformatic Approaches. Biomedicines. 2023;11(8).

6.   Mondal P, Patel NS, Bailey K, Aravind S, Cartwright SB, Hollingsworth MA, et al. Induction of pancreatic neoplasia in the KRAS/TP53 Oncopig. Disease models & mechanisms. 2023;16(1).

7.   Schook LB, Collares TV, Hu W, Liang Y, Rodrigues FM, Rund LA, et al. A Genetic Porcine Model of Cancer. PLoS One. 2015;10(7):e0128864.

8.   Principe DR, Overgaard NH, Park AJ, Diaz AM, Torres C, McKinney R, et al. KRAS(G12D) and TP53(R167H) Cooperate to Induce Pancreatic Ductal Adenocarcinoma in Sus scrofa Pigs. Sci Rep. 2018;8(1):12548.

9.   Boas FE, Nurili F, Bendet A, Cheleuitte-Nieves C, Basturk O, Askan G, et al. Induction and characterization of pancreatic cancer in a transgenic pig model. PLoS One. 2020;15(9):e0239391.

10.   Collins MA, Bednar F, Zhang Y, Brisset JC, Galbán S, Galbán CJ, et al. Oncogenic Kras is required for both the initiation and maintenance of pancreatic cancer in mice. The Journal of clinical investigation. 2012;122(2):639-53.

11.   Hingorani SR, Petricoin EF, Maitra A, Rajapakse V, King C, Jacobetz MA, et al. Preinvasive and invasive ductal pancreatic cancer and its early detection in the mouse. Cancer Cell. 2003;4(6):437-50.

12.   Hingorani SR, Wang L, Multani AS, Combs C, Deramaudt TB, Hruban RH, et al. Trp53R172H and KrasG12D cooperate to promote chromosomal instability and widely metastatic pancreatic ductal adenocarcinoma in mice. Cancer Cell. 2005;7(5):469-83.

13.   Waddell N, Pajic M, Patch AM, Chang DK, Kassahn KS, Bailey P, et al. Whole genomes redefine the mutational landscape of pancreatic cancer. Nature. 2015;518(7540):495-501.

14.   Biankin AV, Waddell N, Kassahn KS, Gingras MC, Muthuswamy LB, Johns AL, et al. Pancreatic cancer genomes reveal aberrations in axon guidance pathway genes. Nature. 2012;491(7424):399-405.

15.   Jones S, Zhang X, Parsons DW, Lin JC, Leary RJ, Angenendt P, et al. Core signaling pathways in human pancreatic cancers revealed by global genomic analyses. Science. 2008;321(5897):1801-6.

16.   Tao L, van Bragt MP, Laudadio E, Li Z. Lineage tracing of mammary epithelial cells using cell-type-specific cre-expressing adenoviruses. Stem cell reports. 2014;2(6):770-9.

17.   Huang Y, Kruse RL, Ding H, Itani MI, Morrison J, Wang ZZ, et al. Parameters of biliary hydrodynamic injection during endoscopic retrograde cholangio-pancreatography in pigs for applications in gene delivery. PLoS One. 2021;16(4):e0249931.

18.   Søreide JA, Karlsen LN, Sandblom G, Enochsson L. Endoscopic retrograde cholangiopancreatography (ERCP): lessons learned from population-based national registries: a systematic review. Surgical Endoscopy. 2019;33(6):1731-48.

19.   Malheiro R, de Monteiro-Soares M, Hassan C, Dinis-Ribeiro M. Methodological quality of guidelines in gastroenterology. Endoscopy. 2014;46(6):513-25.

20.   Enochsson L, Blohm M, Sandblom G, Jonas E, Hallerbäck B, Lundell L, et al. Inversed relationship between completeness of follow-up and coverage of postoperative complications in gallstone surgery and ERCP: a potential source of bias in patient registers. BMJ Open. 2018;8(1):e019551.

21.   Baron TH, Kozarek RA, Carr-Locke DL. ERCP E-Book: Elsevier Health Sciences; 2012.

22.   Lin E, Schueler B. 3 - Radiologic Issues and Radiation Safety During ERCP. In: Baron TH, Kozarek RA, Carr-Locke DL, editors. ERCP (Third Edition). Philadelphia: Elsevier; 2019. p. 14-29.e1.

23.   Gustafsson JK, Johansson MEV. The role of goblet cells and mucus in intestinal homeostasis. Nat Rev Gastroenterol Hepatol. 2022;19(12):785-803.

24.   Killcoyne S, Fitzgerald RC. Evolution and progression of Barrett's oesophagus to oesophageal cancer. Nat Rev Cancer. 2021;21(11):731-41.

25.   Saur D, Schnieke A. Porcine cancer models for clinical translation. Nat Rev Cancer. 2022;22(7):375-6.

26.   Misra S, Moro CF, Del Chiaro M, Pouso S, Sebestyén A, Löhr M, et al. Ex vivo organotypic culture system of precision-cut slices of human pancreatic ductal adenocarcinoma. Sci Rep. 2019;9(1):2133.

27.   Kokkinos J, Sharbeen G, Haghighi KS, Ignacio RMC, Kopecky C, Gonzales-Aloy E, et al. Ex vivo culture of intact human patient derived pancreatic tumour tissue. Sci Rep. 2021;11(1):1944.

28.   Poblete Jara C, Nogueira G, Morari J, do Prado TP, de Medeiros Bezerra R, Velloso LA, et al. An older diabetes-induced mice model for studying skin wound healing. PLoS One. 2023;18(2):e0281373.

29.   Jara CP, Wang O, Paulino do Prado T, Ismail A, Fabian FM, Li H, et al. Novel fibrin-fibronectin matrix accelerates mice skin wound healing. Bioactive Materials. 2020;5(4):949-62.

30.   Martens S, Coolens K, Olsen C, Lefesvre P, Rooman I. Heterotopia of salivary gland tissue in the pancreas. Diagn Pathol. 2023;18(1):98.

31.   Peng J, Sun BF, Chen CY, Zhou JY, Chen YS, Chen H, et al. Single-cell RNA-seq highlights intra-tumoral heterogeneity and malignant progression in pancreatic ductal adenocarcinoma. Cell Res. 2019;29(9):725-38.

32.   Baron M, Veres A, Wolock SL, Faust AL, Gaujoux R, Vetere A, et al. A Single-Cell Transcriptomic Map of the Human and Mouse Pancreas Reveals Inter- and Intra-cell Population Structure. Cell Syst. 2016;3(4):346-60.e4.

33.   Han L, Jara CP, Wang O, Shi Y, Wu X, Thibivilliers S, et al. Isolating and cryopreserving pig skin cells for single-cell RNA sequencing study. PLoS One. 2022;17(2):e0263869.

34.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, III, Zheng S, Butler A, et al. Integrated analysis of multimodal single-cell data. Cell. 2021;184(13):3573-87.e29.

FIGURE LEGENDS

Figure 1: Analysis of Keratin 8 Expression and Activity in Pancreatic Cells.

(a) Pancreatic KRT8 Transcript Levels: Single-cell RNA sequencing data depicting the expression of KRT8 transcript in various pancreatic cell types from healthy and pancreatic ductal adenocarcinoma (PDAC) samples. The heatmap on the left shows the mean expression levels across healthy cell types, while the dot plot on the right indicates the percentage of cells expressing KRT8 and the average expression level in each PDAC cell type. The analysis was conducted using a total of 26 pancreatic cancer samples and 13 control pancreas samples. (b) Immunohistochemistry for Keratin 8: Representative images of pancreatic sections stained for CAM5.2 and Keratin 8. Top panels: wild type (WT) pancreas; Bottom panels: Oncopig pancreas. The images highlight the expression of Keratin 8 in both the main and accessory ductal structures. (c) Ex Vivo Adenovirus Activity in Porcine Pancreatic Ducts: Images of porcine pancreatic main duct before and after tissue preparation. This ex vivo assay assesses the activity of the Cre-recombinase driven by the Keratin 8 promoter, indicating selective expression in pancreatic ductal cells. (d) Cre-Recombinase Expression in Pancreatic Cells: Immunohistochemical analysis of Cre-recombinase expression in pancreatic sections. Left panel: Control; Right panel: After administration of Ad-K8-Cre showing Cre-recombinase expression in epithelial cells, confirming the specificity of the Keratin 8 promoter.

Figure 2: *In vivo* injection of Ad-K8-Cre into the duct of the connecting lobe of the pancreas in the Oncopig.

(a) Exposure of the connecting lobe of the pancreas (CLP). Identification of the connecting lobe of the pancreas. The stomach, pylorus, and duodenum have been reflected superiorly. The surgeon's finger (asterisk) has dissected between the CLP and the duodenal lobe of the pancreas (DLP). Du = duodenum; Co = colon; SB = small bowel; thick arrow = cephalad. (b) Same as panel a, with continued dissection. The junction of the CLP with the DLP has been isolated, and a loop of silk suture (asterisk) has been placed around this region to aid with retraction. Py = pylorus. (c) The CLP was divided at its junction with the DLP, and the proximal stub of the CLP was ligated with a silk ligature (asterisk). The distal end of the transected CLP has an exposed duct which then was cannulated with a 22 g catheter, and the Ad-K8-Cre vector was injected into the CLP duct using a syringe.

Figure 3: Effect of Ad-K8-Cre injection on gross pancreatic morphology in the Oncopig.

(a) Freshly excised connecting lobe of the pancreas from an Oncopig injected with a high dose (10^9 pfu) of Ad-K8-Cre, showing atrophy and nodularity in the region of the injection (dashed circle). Thin arrow = site of ductal cannulation for vector injection; dashed oval = region of gross effect; thick arrow = cephalad; ruler = in/ cm. (b) Same as panel a, except with a low dose (10^7 pfu) of Ad-K8-Cre. (c) Regional *in vivo* appearance (necropsy) in Oncopig with a high dose injection. The midline incision has been re-opened and the superior margin of the pancreas has been exposed. SB = small bowel; thick arrow = cephalad. (d) Same as panel c, but the duodenal lobe of the pancreas (DLP) has been reflected cephalad to expose the connecting lobe of the pancreas (CLP, the region of injection). Co = colon; Du = duodenum; Py = pylorus; St = stomach.

Figure 4: Histological and Immunohistochemical Analysis of Pancreatic Tissues Post Ad-K8-Cre Administration.

(a) Hematoxylin and Eosin (H&E) Staining: Sequential panels (from left to right) show H&E stained pancreatic tissue from Oncopigs injected with increasing doses of Ad-K8-Cre (10^7 to 10^10 pfu). Top row images are from the injection regions. The bottom row shows tissue samples from regions remote (>10 cm) from the injection region, maintaining normal architectural features.

(b) Ki67 Immunostaining: Demonstrates proliferation within the pancreatic ductal structures at various vector doses. Top row: Injection regions with marked increase in Ki67 expression. Bottom row: Non-injection regions with minimal Ki67 staining, indicative of normal tissue proliferation rates.

Figure 5: KRAS^{G12D} Expression and Cytokeratin Staining Post Ad-K8-Cre Administration

(a) KRAS^{G12D} Staining: Panels display pancreatic tissue sections stained to reveal KRAS^{G12D} expression across different doses (10^7 to 10^10 pfu). Top row: Tissue from the injection regions showing increased KRAS^{G12D} expression, primarily in ductal cells, with disrupted architecture and potential neoplastic transformation. Bottom row: Corresponding non-injection regions maintaining a baseline expression with minimal disruptions. (b) CAM5.2 Immunostaining: Immunohistochemical analysis of CAM5.2 expression, a marker for epithelial cells, across the same dose range. Top row: Marked increase in CAM5.2 staining within injection regions correlates with areas of high KRAS^{G12D} expression, suggesting epithelial origin and proliferation. Bottom row: Minimal CAM5.2 expression within non-injection regions, indicating low epithelial activity and the specificity of vector targeting.

Figure 6: Evaluation of Stromal Reaction in Pancreatic Tissues Post Ad-K8-Cre Injection.

(a) Masson's trichrome staining of pancreatic tissues. Top row: Increased blue staining with the injection regions across increasing vector doses (10^7 to 10^10 pfu). Bottom row: Minimal staining in pancreas remote from the injection regions, showing normal collagen levels. (b) α-SMA immunohistochemistry used to detect stromal cells involved in the fibrotic response. Top row: Enhanced α-SMA staining within the injection regions with increasing vector doses. Bottom row: staining within non-injection regions.(c) Vimentin immunostaining. Top row: Marked increase in vimentin staining within the injection regions, particularly notable at doses 10^9 and 10^10 pfu. Bottom row: Sparse vimentin staining from the non-injection regions.

Figure 7: Alcian Blue-PAS Staining of Pancreatic Tissues Post Ad-K8-Cre Injection.

Top Row: progression of acidic mucin production with the injection regions of Oncopigs administered with increasing doses of Ad-K8-Cre (10^7 to 10^10 pfu). Non-injection Region (Bottom Row): limited Alcian Blue-PAS staining.

SUPPLEMENTARY FIGURE LEGENDS

Figure S1: Schematic Overview of Ad-K8-Cre Adenovirus Experimental Procedure.

(a) Ex Vivo Assessment in Wild Type Porcine Pancreas: This panel illustrates the workflow starting from a wild type domestic pig, extraction of the main pancreatic duct, and subsequent ex vivo infection with Ad-K8-Cre adenovirus. The adenovirus, driven by the Keratin 8 promoter to express Cre-recombinase, targets pancreatic epithelial cells cultured in a petri dish, followed by harvesting for analysis. (b) In Vivo Targeted Activity in Oncopigs: This diagram outlines the in vivo process where the Oncopig, harboring latent $KRAS^{G12D}$ and $TP53^{R167H}$ mutations, is injected with the adenovirus directly into the pancreatic duct. The illustration shows the mechanism by which Cre-recombinase activates these mutations, leading to observable changes in the pancreas after 2 months.

Figure S2: Histopathological Analysis and Genotyping in Oncopigs Post Ad-K8-Cre Injection

(a) Histological Comparison. Left column shows the normal architecture within non-injection regions. Right column images display the injection regions where notable morphological changes are evident, including the presence of goblet cells (yellow arrows), indicative of metaplastic transformation. (b) Cytokeratin 7 Staining. Top row shows increased CK7 staining intensity within the injection regions across increasing vector doses (10^7 to 10^{10} pfu). Bottom row corresponds to non-injection regions, where there is minimal staining, reflecting the epithelial stability and absence of transformation. (c) Genotyping of Oncopigs: gel image represents the genotyping results for Oncopigs, confirming the genetic alterations associated with the study's interventions. Each lane represents an individual animal, with specific band patterns confirming the presence of targeted genetic modifications.

Figure S3: Comparative Histopathological Analysis of Human Pancreatic Cancer Samples.

Vimentin Staining (Top Two Rows): Showcases vimentin immunohistochemistry across PDAC samples from four patients, highlighting the stromal reaction characteristic of pancreatic cancer. Vimentin is abundantly expressed, indicating a robust stromal response. Higher magnifications (20x) provide a closer view of the stromal involvement within the cancer architecture. Masson's Trichrome Staining (Middle Two Rows): Depicts collagen deposition within the tumor microenvironment, stained in blue. These sections reveal varying levels of fibrosis across different samples, with increased collagen indicative of a desmoplastic reaction, which is a hallmark of pancreatic cancer. CAM5.2 Staining (Bottom Two Rows): Highlights the expression of cytokeratin in epithelial cells, underscoring the epithelial nature of the cancerous cells.

Table 1. Oncopig weights.

Ad-K8-Cre dose (pfu)	Survival post-induction	Mean weight (Kg)				
		T_0	2 wk	4 wk	6 wk	Necropsy (8 wk)
10^7	8 wk	27.5	26.8	34.5	38.8	45.8
10^8	8 wk	25.5	25.3	28.3	35.5	40.3
10^9	8 wk	20.0	25.8	27.5	27.0	28.5
10^{10}	8 wk	24.3	28.8	34.3	38.0	48.1
	p value	0.5	0.8	0.6	0.6	0.3

*ANOVA across all four dose groups at the given time point.

Table 2. Oncopig serum testing.

Adcre	Sex	HG		HCT		AMYLASE		LIPASE		CEA		CA 19-9		AFP	
		T0	Necropsy	T0	Necropsy	T0	Necropsy	T0	Necropsy	T0	Necropsy	T0	Necropsy	T0	Necropsy
10^9	M	NA	8.5	NA	30	NA	669	NA	3	NA	0.1	NA	1	NA	0.5
	F	9.7	9.2	34.9	33	1520	1388	19	18	0.1	0.1	1	1	0.5	0.5
10^8	M	11	11.1	39.3	38.6	1015	1095	4	3	0.1	0.1	1	1	0.5	2.5
	F	11	9.4	40.3	33.6	1452	1749	27	3	0.1	0.1	1	1.4	0.8	1
10^7	M	9.3	10.6	32.9	37.3	573	739	8	21	0.1	0.1	2	1	0.5	1
	F	10.3	10.4	36.4	36.9	1551	1865	4	6	0.1	0.5	1	1	0.5	0.5
10^10	M	9.1	clotted	34.2	clotted	674	799	3	3	0.1	0.3	1	1	0.5	0.5
	F	10.4	10.2	38.2	36.9	1477	2556	3	3	0.1	0.1	1	1	0.5	0.5

Hg = hemoglobin; Hct = hematocrit; CEA = carcinoembryonic antigen; CA 19-9 = cancer antigen 19-9; AFP = alpha fetoprotein; T0 = immediately prior to induction.

Fig. 1

Fig. 3

Fig. 4

Fig. 5

Fig. 6

a

Fig. 7

Supplementary Material

a

Wild type
domestic pig

Main Pancreatic
duct

Harvest

K8 promoter
Cre-recombinase
Adenovirus → **Cre-recombinase**

b

K8 promoter
Cre-recombinase
Adenovirus

Cre
Recombinase

silent
mutation

active
mutation

2 months

Oncopigs
KRAS ^{G12D}
PT53 ^{R167H}

Fig. S1

Fig. S2

Fig. S3

a

Non-Injection site	**Injection site**

20x

40x

b

Genotyping Oncopigs

Fig. S4

Patient 1 **Patient 2** **Patient 3** **Patient 4**

10x / 20x — Vimentin

10x / 20x — Masson's

10x / 20x — CAM5.2

Fig. S5